Title: Developmental co-emergence of cardiac and gut tissues modeled by human iPSC-derived organoids

Authors: A.C. Silva[1], O.B. Matthys[1,2], D.A. Joy[1,2], M.A. Kauss[1,3], V. Natarajan[1], M.H. Lai[1], D. Turaga[1], A.P. Blair[1], M. Alexanian[1], B.G. Bruneau[1,4,5,6], T.C. McDevitt[1,7,*]

Affiliations:

[1]Gladstone Institutes, San Francisco, CA.

[2]UC Berkeley-UC San Francisco Graduate Program in Bioengineering, San Francisco, CA.

[3]UC San Francisco Graduate Program in Biomedical Sciences, San Francisco, CA.

[4]Roddenberry Center for Stem Cell Biology and Medicine at Gladstone, San Francisco, CA.

[5]Cardiovascular Research Institute, University of California, San Francisco, CA.

[6]Department of Pediatrics, University of California, San Francisco, CA.

[7]Department of Bioengineering and Therapeutic Sciences, University of California, San Francisco, CA.

* Correspondence to: todd.mcdevitt@gladstone.ucsf.edu

Abstract:

During embryogenesis, paracrine signaling between tissues in close proximity contributes to the determination of their respective cell fate(s) and development into functional organs. Organoids are *in vitro* models that mimic organ formation and cellular heterogeneity, but lack the paracrine input of surrounding tissues. Here, we describe a human multilineage iPSC-derived organoid that recapitulates cooperative cardiac and gut development and displays extensive cellular and structural complexity of both tissues. We demonstrate that the presence of endoderm tissue (gut/intestine) in multilineage organoids contributed to the development of the cardiac tissue, specifically cardiomyocyte expansion, compartmentalization, enrichment of atrial/nodal cells, myocardial compaction and functional fetal-like maturation. Overall, this study demonstrates the ability to generate specific cooperative tissues originating from different germ lineages within a single organoid model, an advance that will further the examination of multi-tissue interactions during development and disease.

Keywords:

Organoids, pluripotent stem cells, morphogenesis, co-emergence, cooperative development, endoderm, heart, intestine, atrial/nodal cardiomyocytes

Introduction

The formation of organs during embryogenesis results from a coordinated sequence of morphogenic events that regulate the co-emergence and self-organization of distinct cell populations (Zhang and Hiiragi, 2018). Organoids derived from stem and progenitor cells provide an experimentally tractable model of tissue morphogenesis *in vitro* due to the fact that they differentiate into multiple cell types in a spatially-organized manner and recapitulate functional properties of developing native tissues (Lancaster and Knoblich, 2014;Matthys *et al.*, 2020). However, most organoids to date give rise to single tissues in isolation, and lack the mechanical cues and paracrine signaling contributed by neighboring organs that synergistically benefit the trajectory of tissue development *in vivo* (Varner and Taber, 2012;Jung *et al.*, 1999;Ishii *et al.*, 2007). Previous attempts to generate multi-tissue organoids in a controllable fashion have mainly involved the fusion of pre-differentiated single-lineage organoids (Koike *et al.*, 2019;Bagley *et al.*, 2017).

Mesoderm and endoderm display reciprocal interactions during embryonic development that are fundamental to the successful formation of several organs, including the heart (Lough and Sugi, 2000;Schultheiss *et al.*, 1995;Sugi and Lough, 1994). *In vitro* co-culture systems of embryonic germ layer explants and pluripotent stem cells demonstrate that endoderm cells secret cardiac-inductive factors (e.g. Activin A, FGF2) that promote cardiomyocyte formation (Anderson *et al.*, 2016;Rudy-Reil and Lough, 2004;Brown *et al.*, 2010;Uosaki *et al.*, 2012;Sugi and Lough, 1995). However, it has been challenging to determine the effect and sequence of direct interactions between endoderm and mesoderm *in vivo* because interrogation at each stage of development disrupts the natural course of morphogenesis.

Here, we generated 3D spheroids of early committed mesendoderm progenitors in a

culture medium permissive of multi-lineage differentiation. We observed the co-emergence of

mesoderm and endoderm cell lineages, resulting in unique multilineage organoids able to

spontaneously organize the formation of complex structures similar to the developing heart and

gut. Multiple *in vitro* models have been developed to study various aspects of the heart,

including patterning constructs (Ma *et al.*, 2015), gastruloids/organoids (Lee *et al.*, 2020;Rossi *et*

al., 2020),(Drakhlis *et al.*, 2021) (development), and engineered microtissues (homeostasis and

maturation) (Ronaldson-Bouchard *et al.*, 2018;Hookway *et al.*, 2019;Mills *et al.*, 2017;Macqueen

et al., 2018). However, models that recapitulate heart morphogenic dynamics (i.e. gastruloid/

organoids) have been mostly obtained with mouse stem cells (Rossi *et al.*, 2020;Lee *et al.*, 2020),

and only recently using human induced pluripotent stem cells (iPSCs)(Drakhlis *et al.*, 2021). On

the other hand, intestinal tissue has been generated *in vitro* with high reproducibility using mouse

and human stem cells derived from tissue biopsies and iPSC (Sato *et al.*, 2009;Munera *et al.*,

2017). The cardiac and gut (multilineage) organoid presented here constitutes a first step toward

a new generation of organoids that recapitulate the co-development and differentiation of two

defined tissues for long periods of time (>1 year). Such defined multi-lineage/multi-organ

organoids hold the unique promise of modeling the crosstalk between specific tissues that occurs

during human embryogenesis, tissue maturation and disease— a complex phenomenon that is

difficult to dissect *in vivo*.

Results

Formation of heart-like compartments in multilineage spheroids

We began by generating spheroids from mesendoderm progenitors differentiated for 5 days with the GiWi protocol (Lian *et al.*, 2012). Spheroids were then cultured in conventional, cardiac-permissive medium supplemented with insulin (RPMI/B27+), or in an ascorbic acid (AA)-supplemented medium formulated for epicardial cell differentiation to mimic the signaling that occurs during cardiac differentiation to support the emergence of a variety of cell types (i.e. fibroblasts and smooth muscle cells (SMC))(Bao *et al.*, 2016;Lian *et al.*, 2013) **(Figure 1A,B)**. Noticeable differences in size and structural arrangement between spheroids grown in conventional versus AA medium (hereafter referred to as "conventional" and "multilineage" microtissues, respectively) became evident within 2 weeks of differentiation, with multilineage microtissues growing to approximately 3 times the size of conventional microtissues by 30 days **(Figure 1C, D, E)**. After 10 days, conventional microtissues displayed a spherical, uniformly translucent morphology with spontaneous contractility throughout **(Figure 1C, D; Movie S1)**, whereas the multilineage microtissues exhibited a dark core of vigorously contracting cells surrounded by translucent non-contracting cells. Calcium transients were limited to the central region of contracting cells in the multilineage tissues **(Figure 1C,D; Movie S2)**. At early stages of culture, light-sheet fluorescence microscopy revealed that conventional microtissues were largely composed of cardiomyocytes, as identified by cardiac troponin (cTnT) immunostaining, that co-expressed TBX18 (cTnT+/TBX18+ cells), and an outer layer of TBX18+ epicardial cells (cTnT-/TBX18+) **(Figure 1F; S1A, S2A)**. However, by day 40, conventional microtissues had only small patches of epicardial cells at the surface **(Figure 1F; S2A)**. In contrast, multilineage microtissues displayed a core of cTnT+ cardiomyocytes surrounded by a non-myocyte, stromal-

like cell population (phalloidin+, cTnT-), and an outer layer of TBX18+ epicardial cells (cTnT-

/TBX18+), which was preserved for more than two month **(Figure 1F; S1A, S2A)**, and better

resembled the radial organization of the embryonic heart wall **(Figure S1A)**.

Additional histological analysis revealed that both conventional and multilineage microtissues

contained SMCs (SMA+/MYH7-; **Figure S2B**). However, in multilineage microtissues,

cardiomyocytes positive for the ventricular marker MLC2v aggregated together in the cardiac

tissue core, indicating spatial compartmentalization reminiscent of the distinct emergence of

atrial and ventricular cardiomyocytes in the primitive heart **(Figure 1G)**; distinct spatial

compartmentalization of cardiomyocytes was not observed in conventional microtissues.

The dramatic growth of multilineage microtissues over the first 30 days in culture was

paralleled by an increase in the volume of the cardiac region **(Figure 1H; S1C, S2C; Movie S3-**

4). Cardiac growth appeared to result from cardiomyocyte proliferation, based upon

immunostaining for Ki67, a marker of active cell cycle—analogous to cardiac wall thickening

during development **(Figure 1H; S1C)**. In addition, at day 30, we observed the emergence of

small epithelial cystic structures within the previously translucent layers of cells enclosing the

cardiac core in multilineage microtissues **(Figure 1C)**. These cystic structures expressed the

mid-hindgut (MHG) endoderm markers FOXA2 and CDX2 **(Figure 1C,I; S2D,E)**. Since the

multilineage microtissues contained endoderm epithelial clusters, attained greater structural

complexity typical of the cardiac compartment and exhibited growth dynamics comparable to the

embryonic heart, they were hereafter referred to as multilineage organoids **(Figure 1J)**

(Lancaster and Knoblich, 2014).

Long-term culture of multilineage organoids recapitulates embryonic heart and gut

development

We next examined whether multilineage organoids acquired further complexity with

extended time in culture. The growth of multilineage organoids accelerated throughout culture

duration, with the average organoid reaching ~2mm in diameter after 100 days **(Figure 2A-C)**.

After 50 days, the stromal cells surrounding the cardiac core were replaced by a mesenchymal

tissue **(Figure 2D)** enriched in extracellular matrix (ECM) components, such as hyaluronic acid

(Figure S2F), and a fine fibrillar network of glycosaminoglycans **(Figure 2D, light green/blue)**

that was populated by a few *WT1*-expressing cells **(Figure S1A)**. The mesenchymal tissue

resembled the interstitial tissue underlying the epicardium (or sub-epicardium) and the

peritoneum (protective layer of the intestine) **(Figure 2D)**. In addition, smooth muscle actin

(SMA)-positive cells were detected at the surface of the organoid and in the sub-epicardium, and

a few endothelial cells (PECAM-1+) appeared interspersed among the cardiomyocytes **(Figure**

2F). By day 60, SMCs (αSMA+ MYH7-) emerged prominently between the cardiac and the

primitive gut-like tissue. The SMC tissue displayed slow and infrequent calcium transients (~5s

duration per transient; 1-3 transients/minute) that corresponded to large contractions (>100 μm)

that resembled peristalsis **(Figure 2G,H; Movie S5-6)**. More endothelial cells were detected

between the cardiomyocytes, forming branched-like structures resembling pre-vascular networks

(Figure 2F). Between 70 and 100 days of differentiation, the primitive gut tissues matured, as

shown by the development of Paneth cells (antimicrobial peptide-producing cells) and goblet

cells (which produce mucins) **(Figure 2D)**. Thus, multilineage organoids undergo a sequence of

morphogenic events that recapitulates the development of the human embryonic heart and gut,

including the initial emergence of cardiac tissue followed by gut tissue, with interstitial tissue

(sub-epicardium/sub-peritoneum-like tissue) separating the two distinct tissues.

To assess the cellular heterogeneity of the multilineage organoids, we performed single-

cell RNA-sequencing (scRNA-seq) on dissociated organoids after 100 days. Seurat analysis

revealed 6 cell clusters corresponding to cardiomyocytes, SMCs, proliferating cells, fibroblasts,

gut and enteroendocrine cells **(Figure 3A-C; S3A; Table S1)**. GO term analysis of the

enteroendocrine cell cluster yielded hormone transport and secretion **(**expressing CHGA; **Figure**

3B; S3A), suggesting the development of chemosensing properties in the gut

compartment(Gribble and Reimann, 2016). Sub-clustering of the gut cells confirmed the

differentiation of specialized intestinal cells, including: 1) enterocyte-like cells (absorptive cells

expressing VIL1, ANPEP, MUC13/17); 2) Paneth cells (expressing LYZ); 3) LGR5-expressing

intestinal stem cells; and 4) small intestine cells marked by high expression of APOB, CDX2 and

VIL1, along with enrichment of genes for glucose and cholesterol transporters (SLC2A2 and

NPC1L1, respectively), and digestion (FABP2, SI, SLC5A1) **(Figure 3C,D; S3A,B)**. The

presence of these gut cell types was further validated by electron microscopy of 100-day old

multilineage organoids through the identification of key structural features of each cell type

(Figure 2I i-v). Goblet cells were not detected by scRNA-seq, presumably due to their larger

size.

Analysis of cell populations within the cluster enriched for proliferative markers revealed

the presence of proliferating cardiomyocytes (TTN, MYH6, NPPA) and SMCs (ACTG2,

MYH11) **(Figure S3C,D)**. Ki67 immunostaining demonstrated a decline in the number of

actively cycling cells over time **(Figure 2J)**; nevertheless, some cardiomyocytes (slow/fast

TnI+), SMCs, and gut cells remained Ki67+ after 100 days in culture, corroborating the scRNA-

seq results **(Figure S3G)**. Sub-clustering analysis of the cardiomyocytes indicated that after 100

days in culture, cardiomyocytes were divided into 3 groups, separated mainly by differences in

maturation stage markers: immature cells that either expressed ECM related genes(Cui *et al.*,

2019) (Cluster 2) or smooth muscle myosin (MYH11, Cluster 0), and more mature-like

cardiomyocytes (Cluster 1) that expressed higher levels of TNNI3 **(Figure 3E,F)**. A minority of

cardiomyocytes (~30%) expressed the ventricular cardiomyocyte markers IRX4 or MYH7

(Figure 3G; 3SA), indicating that multilineage organoids favor the differentiation and

maturation of atrial cardiomyocytes, as confirmed by the high expression of atrial markers such

as TBX5, MYH6 and GJA5 **(Figure 3G)**. Although key atrial and ventricular genes were

expressed in the same cluster of cells (Cluster 1), these genes were not co-expressed in individual

cells **(Figure S3F)**.

To identify *in vivo* correlates, we used a cell type classifier, trained on scRNA-seq of

human fetal heart tissue collected at 6.5-7 weeks **(Figure S3H)**[Asp *et al.*, 2019], to characterize a

subset of cardiomyocyte, fibroblast and SMC clusters that were identified by default Seurat

clustering and cardiac cell compartmentalization genes. The classifier estimated that 98% of the

organoid cardiomyocytes were atrial cardiomyocytes (prediction probability = 0.91) **(Figure 3H,**

I; S3G), with classification driven by MYH6 expression, a classifier top feature for atrial cell

predictions **(Table S2)**. Ventricular cells only represented 1% of the cardiomyocytes (prediction

probability = 0.69) **(Figure 3H, I; S3G)**. The cell classifier also identified fibroblast and

SMC/Fibroblast-like cells, as expected, which represented 44% and 11% of the cells analyzed

with prediction probabilities of 0.85 and 0.75, respectively **(Figure 3H,I)**. In addition, the *in vivo*

cell classifier assigned cell type identities not detected using default Seurat clustering analysis,

such as sub-epicardial and capillary endothelium/pericytes/adventia cells, which together

constituted less than 4% of the total cells evaluated **(Figure 3H,I; S3G)**. Although the capillary

endothelium/pericytes/adventia cells corresponded to only 0.1% of the cells, their assignment

had a high prediction probability of 1.00 **(Figure 3H; S3G)**. In summary, the *in vivo* cell

classifier corroborated the cellular complexity of the multilineage organoids first observed by

histological analysis, and suggested a high transcriptomic similarity between cells from

multilineage organoids and cells from fetal human hearts **(Figure 3I)**.

Multilineage organoids remained viable for over 1 year in culture, and stably maintained

their millimeter size-scale and structural complexity **(Figure S4)**. Contractile behavior was

observed only when electrical stimulation was applied, a feature consistent with functionally

mature iPSC-derived cardiomyocytes (Yang *et al.*, 2014) **(Figure S4B)**. The cardiac and

intestine tissues remained localized at opposite ends of the organoids, separated by stromal cells

and SMCs **(Figure S4C-E,G)**. In contrast to previous timepoints, small clusters of SMCs were

detected amongst the cardiomyocytes in these aged organoids, and SMCs were more pronounced

at the boundary of cardiac tissue regions **(Figure S4F)**. In addition, proliferating non-myocytes

(Ki67+, cTnT-) were detected within the cardiac tissue, along with structures that resembled

mature blood vessels (expressing lectin, a glycoprotein of the basal lamina of endothelial cells)

embedded within or adjacent to the cardiac regions in some organoids **(Figure S4C, G)**.

Organoids cultured for 1 year proved challenging to dissociate to single cells, potentially due to

the structural complexity and ECM composition, thus limiting our ability to capture the

morphological features and the transcriptional signature of individual cells.

Overall, these observations demonstrate that multilineage organoids represent a new type

of compartmentalized organoid model that exhibits the co-development and maturation of tissues

derived from two distinct germ layers, heart (mesoderm) and gut (endoderm)(Koike *et al.*,

2019;Faustino Martins *et al.*, 2020;Bagley *et al.*, 2017).

Multilineage organoids promote atrial/nodal cardiomyocyte specification and maturation

To further assess the maturity and sub-type specification of the cardiomyocytes, we

performed electrophysiological and structural analysis of the intact spheroids and dissociated

cells. Over time, 100% of conventional microtissues exhibited spontaneous contractility, and

33±21% of day-80 conventional microtissues could be paced at 2Hz **(Figure 4A; S5A,B)**.

Furthermore, conventional microtissues exhibited no dramatic changes over time in kinetic

calcium handling properties (amplitude and stroke velocities) under 1-Hz of stimulation,

reflecting relatively immature function **(Figure 4B,C)**. In contrast, multilineage organoids

demonstrated a decrease in spontaneous calcium activity over time, from 100% at day 10 to

37±16% at day 100, and an ability to respond to higher frequencies of electrical stimulation, up

to 8Hz after 80 days of culture **(Figure 4A-D; S5A)**. In addition, after 30 days, multilineage

organoids demonstrated progressively increasing amplitude and maximum stroke velocities

under 1-Hz of stimulation, consistent with greater functional maturation than the conventional

microtissues **(Figure 4B,C).**

Morphological analysis of dissociated tissues revealed greater structural maturation of

individual cardiomyocytes from day-30 multilineage organoids than conventional microtissues,

as determined by greater surface area and perimeter, elongated morphology, and highly-aligned

and defined sarcomeres **(Figure 4E-G)**(Lundy *et al.*, 2013). More elongated cardiomyocytes

with highly-aligned and defined sarcomeres were isolated from day-80 multilineage organoids,

reflecting additional structural maturation over time **(Figure 4F,G)**. Electron microscopic

imaging of cardiomyocytes from day-100 multilineage organoids revealed a series of

ultrastructural features typical of the third-trimester human fetal heart, including: 1) myofibrils

closely packed with defined striations and no apparent M bands **(Figure 4H,i)**; 2) cylindrical-

shaped mitochondria aligned between the myofibrils **(Figure 4H,ii)**; and 3) large pools of

glycogen vesicles, the major energy source during fetal heart development **(Figure 4H,iii)** (Kim

et al., 1992;Racca *et al.*, 2016). This advanced degree of iPSC-derived cardiomyocyte structural

features is not commonly observed without external stimuli (mechanical, electrical or chemical),

thus highlighting the intrinsic ability of the multilineage organoid model to promote cellular

maturation of cardiac muscle cells.

We next evaluated cardiomyocyte function using patch-clamp analysis of dissociated

single cells, considering multiple action potential parameters (APD_{90}/APD_{50} ratio, APD_{30-}

$_{40}/APD_{70-80}$ ratio, MDP, peak, amplitude, frequency, and dV/dTMax). Electrophysiology analysis

revealed a higher proportion of atrial and nodal-like (atrial/nodal) cardiomyocytes (day 30: 73%)

than found in conventional microtissues (day 30: 57%) **(Figure 4 I,J)**. In addition, the number of

atrial/nodal cardiomyocytes increased over time (multilineage organoids day 30: 73%, day 60:

80%) **(Figure 4 I,J)**, consistent with the assignment of 98% of cardiomyocytes as atrial-like cells

by the *in vivo* cell classifier at day 100 **(Figure 3H; S3G)**. These results reflect the ability of

multilineage organoids to specify and mature atrial/nodal cardiomyocytes, which has been

challenging to achieve by alternative 3D differentiation methods.

The primary differences between the multilineage and conventional media are the

concentrations of ascorbic acid (AA) and calcium. Previous studies have demonstrated that these

two components can influence cardiomyocyte differentiation and maturation, respectively(Cao *et*

al., 2012;Wheelwright *et al.*, 2018). To assess the potential impact of different culture medium

components alone on cardiomyocyte specification and maturation, patch-clamp analyses were

performed in iPSC-cardiomyocytes differentiated in monolayers in conventional cardiac media

(Ca^{2+}= 0.42mM), multilineage media (AA=100μg/ml, Ca^{2+}=1.05mM) and conventional cardiac

media supplemented with AA and calcium (AA=100μg/ml, Ca^{2+}=0.63mM). The AA- and Ca^{2+}-

supplemented conventional cardiac media enhanced ventricular cardiomyocyte maturation by

promoting an increase in peak, amplitude and upstroke velocity (V/s) of the action potentials, but

did not promote the emergence of atrial/nodal cardiomyocyte subtypes (no atrial/nodal cells were

identified) **(Figure S5C-E)**. Cardiomyocytes differentiated in multilineage medium (in 2D

monolayers) exhibited comparable maturation to those in the supplemented conventional media,

but yielded mainly ventricular-like cells (97%) **(Figure S5C-E)**. Therefore, although the media

components promoted functional cardiomyocyte maturation, they alone were not sufficient to

direct atrial/nodal specification. Furthermore, these experiments suggest that the co-development

of the gut tissue, which is absent in conventional cardiac microtissues and 2D monolayers,

appears to be a critical factor responsible for enrichment of atrial/nodal cardiomyocytes (>90% at

day 100; **Figure 3H, S3J**) and the physiological maturation of cardiac tissue in multilineage

organoids to an equivalent level of third-trimester human fetal hearts.

Multilineage organoids express cardiac and gut master regulatory genes

Many iPSC differentiation protocols suffer from inherent variability in consistency

(Volpato and Webber, 2020;Matthys *et al.*, 2020). In our system, when multilineage organoids

failed to develop, spheroids instead yielded either solely cardiac microtissues or gut organoids

(Figure 5A,B; S6). Despite batch-to-batch differences in spheroid phenotype, all spheroids

(n>200) within a single batch derived from the same initial differentiation of progenitor cells

manifested identical morphologies and cellularity **(Figure 5A)**. Thus, we hypothesized that

subtle differences in the cellular composition of the initial 2D populations of differentiating cells

were responsible for the phenotypic outcomes of the spheroids.

To evaluate the variability between the starting progenitor populations that gave rise to

cardiac microtissues, multilineage organoids, or gut organoids, we performed retrospective

analysis of the initial progenitor cell population at day 5 of differentiation by bulk RNA-

sequencing, immediately prior to spheroid formation **(Figure 5B)**. Multidimensional scaling

(MDS) analysis of the progenitor transcriptomes revealed that different batches had distinct gene

expression profiles, and that the progenitors of multilineage organoids represented a unique

transcriptional state, roughly equidistant from both cardiac and gut progenitors **(Figure 5C)**.

Comparative analysis between cardiac microtissue and gut organoid progenitors indicated that

cardiac microtissue progenitors displayed a muscle development-restricted gene expression

program. In contrast, progenitors of gut organoids revealed activation of a broader embryonic

developmental program, including forebrain, heart, visual and sensory system development

(Figure S7B). To dissect the specific transcriptomic features enabling multilineage organoid

formation, differential expression analysis was performed to compare the multilineage

transcriptional profile to gut organoid and cardiac microtissue progenitors independently.

Compared to cardiac microtissue progenitors, multilineage organoid progenitors were enriched

in genes related to muscle tissue development and function (*MYH6*, *MYOM1*, *TTN*), as well as

transcription factors (TFs) expressed by cardiomyocytes (*MEF2C*, *NKX2-5*, *ISL1*) **(Figure 5D)**,

suggesting an increased cardiac signature in the multilineage progenitors. Compared to gut

organoid progenitors, multilineage progenitors exhibited a gene expression enrichment of the

muscle tissue development program (*ACTC1*, *MYH6*, *MYOM1*) **(Figure 5E)**. No gene ontology

term enrichment was observed for the gut organoid progenitors relative to the multilineage

progenitors, presumably due to the fact that: 1) the population of gut progenitors express markers

of early cardiac differentiation similar to the multilineage progenitors, 2) at early time points

(before 20 days), gut organoids phenotypically resemble multilineage spheroids, and 3) and gut

tissue development required at least 2 months of 3D culture to manifest clearly with pronounced

villi only evident after 100 days **(Figure 5A, S6, S7A; Movie S8)**. Histological analysis revealed

that primitive gut structures emerged earlier in the gut organoids **(Figure S7A, day 12)**, however

similar structures were not seen in multilineage organoids until later time points **(Figure 5A, day**

20-30; Movie S8). Thus, the accelerated development of primitive cystic structures in gut

organoids might have disrupted the cardiac morphogenesis originally present in multilineage

organoids.

In order to dissect specific gene expression differences between multilineage and cardiac

or gut progenitors, a targeted pairwise analysis was performed for heart patterning transcription

factors, cardiac progenitors and endoderm-related genes. Multilineage progenitors expressed

higher levels of T-box (TBX) transcription factors (*TBX2, TBX5, TBX18, TBX20*) that regulate

the developing heart(Stennard and Harvey, 2005) , in comparison with the gut progenitors,

revealing a broader activation of the cardiac patterning program. Notably, in contrast to the

cardiac progenitors, multilineage progenitors expressed higher levels of TBX genes related to

sinoatrial node formation (*TBX18*) and chamber formation and septation (*TBX5*), in addition to

HEY1, a gene whose expression is restricted to the atria in the context of the heart(Rutenberg *et*

al., 2006), suggesting a transcriptomic profile more consistent with atrial/nodal cardiomyocyte

specification. Furthermore, multilineage progenitors expressed several TFs expressed by cardiac

cells (*ISL1*, *NKX2-5*, *TBX5*) and epicardial-related genes (*TBX18*, *CDH1, ALDH1A2, SEMA3D*)

at higher levels than cardiac progenitors **(Figure 5F, top)**, consistent with the greater cardiac cell

diversity observed in the multilineage organoids as compared to the cardiac microtissues **(Figure**

5A). Progenitors yielding multilineage organoids also expressed genes involved in gut tube

development (*SOX17*, *ONECUT2*), although at lesser levels than gut organoid progenitors

(notably for *SOX17*), suggesting that the multilineage populations contained a smaller number or

proportion of gut progenitors **(Figure 5F, bottom)**. In total, multilineage organoid progenitors

displayed a mixed transcriptomic profile of cardiac and gut progenitors, which could account for

their ability to give rise to organoids consisting of both heart and gut cells.

Multilineage organoid progenitors express cardiac and gut relevant paracrine factor genes

Distinct endoderm subtypes are known to play a role in cardiomyocyte formation through

paracrine signaling mechanisms (Holtzinger *et al.*, 2010;Brown and Badylak, 2014;Anderson *et*

al., 2016). To examine potential intercellular paracrine signaling affecting the co-emergent

populations, we compared the expression of genes associated with secreted factors between the

different progenitor populations **(Figure 5G)**. We first generated a list of secreted factors that

were differentially enriched in the cardiac or gut progenitors **(Figure S7D)**. FGF10 and NPPB

were identified among the gut-enriched factors. FGF10 is a growth factor that enhances

cardiogenic specification, fetal cardiomyocyte proliferation, and has been identified as a second

heart field (SHF)-specific marker, indicating a role in heart tissue development (Al Alam *et al.*,

2015;Rochais *et al.*, 2014;Kelly *et al.*, 2001;Chan *et al.*, 2010). NPPB is highly expressed during

cardiogenesis, particularly in the differentiating chamber myocardium (Sergeeva and

Christoffels, 2013). However, FGF10 and NPPB also promote intestinal cell proliferation,

induction of goblet cell differentiation and homeostasis of mesenteric and intestinal tissue(Al

351 Alam *et al.*, 2015;Rochais *et al.*, 2014;Kelly *et al.*, 2001;Sergeeva and Christoffels, 2013). Thus,

352 FGF10 and NPPB could enhance aspects of both cardiac and gut tissue formation. Indeed,

353 multilineage progenitors expressed higher levels of FGF10 and NPPB than the cardiac tissue

354 progenitors **(Figure 5G, top; S7E, left)**. Multilineage progenitors also expressed higher levels of

355 cardiac-enriched factors than the gut progenitors, such as the growth factors IGF1 and ANGPT1,

356 and the ECM proteins DCN and LUM **(Figure 5G, bottom; S7E, right)**. IGF1 enhances

357 mesodermal cell differentiation and subsequent cardiac progenitor cell formation (Engels *et al.*,

358 2013), and ANGPT1 supports proper atrium chamber morphogenesis (Kim *et al.*, 2018). DCN

359 and LUM, matrix molecule genes enriched in the multilineage progenitors, contribute to the

360 structure and mechanics of the heart by binding to core matrix proteins like collagen I and III,

361 and facilitate thickening of collagen fibrils, respectively, thereby impacting collagen

362 fibrillogenesis (Chakravarti, 2002;Ferdous *et al.*, 2007). Hence, multilineage progenitors

363 expressed increased levels of secreted factor genes relevant to both cardiac and gut tissue

364 morphogenesis compared to either individual tissue alone.

365  In summary, this retrospective analysis revealed that, in addition to the emergence of

366 cardiac cells within the multilineage organoids, the co-development of gut alongside cardiac

367 cells potentially contributes to more structurally and functionally mature cardiac tissue via the

368 secretion of key cardio-inductive molecules.

Discussion

Although organoids can recapitulate many physiological properties of tissue development *in vitro*, they typically lack the cooperative paracrine interactions between adjacent tissues that characterize embryogenesis, as well as the diverse multilineage progenitors essential for the development of complex organs such as the heart. Current strategies to increase organoid complexity have focused on the fusion of differentiated organoids (Koike *et al.*, 2019;Bagley *et al.*, 2017), but these approaches miss the interactions between germ layers that occur during the early steps of embryogenesis to initiate the formation of different organs. Although a couple of recent reports demonstrate the ability to generate primitive cardiac tissues coincident with endoderm derivatives from pluripotent stem cells with greater fidelity than traditional embryoid body differentiation (Rossi *et al.*, 2020;Drakhlis *et al.*, 2021), such organoids consist of many cell types from different germ lineages. However, in contrast, here we describe a multilineage organoid model system in which only cardiac (mesoderm) and gut (endoderm) tissues emerge in a sequential manner. This multilineage organoid recapitulates morphogenic events that lead to the emergence of key structural features of the developing human heart and gut. Past studies of the interactions between heart and gut during development indicated that endoderm-derived cells generate paracrine signals and mechanical forces that support heart formation (Rudy-Reil and Lough, 2004;Brown *et al.*, 2010;Uosaki *et al.*, 2012;Sugi and Lough, 1995;Kidokoro *et al.*, 2018)·(Hosseini *et al.*, 2017;Kelly *et al.*, 2001;Drakhlis *et al.*, 2021;Rossi *et al.*, 2020). Consistent with those findings, our multilineage organoid also demonstrates that endoderm-derived cells can positively impact cardiomyocyte phenotype, suggesting a broad contribution of endoderm-derived cells to heart morphogenesis.

We generated multilineage organoids by aggregating early committed mesendoderm

progenitors derived from human iPSCs in permissive media conditions. The sequential

emergence of functional cardiac and gut-like tissues mimicked human embryonic

morphogenesis, in which the first heart beat (3 weeks of gestation) is followed one week later by

the extension of the gut tube into fore-, mid- and hind-gut structures (Hill, 2020b;Hill, 2020a). At

early stages of *in vitro* culture, multilineage organoids resemble human fetal heart formation (~5

weeks of gestation), evidenced by the formation of a surface layer of epicardial cells and a core

of cardiomyocytes. Similar features were also observed in conventional microtissues, derived

from progenitors spheroids cultured in cardiac-permissive medium. However, only multilineage

organoids exhibited significant growth and preserved the exterior epicardial layer over the long

term, as well as, a preferential differentiation into atrial/nodal cells, in stark contrast to other

cardiac organoid models that also contain endoderm-derived cells(Rossi *et al.*, 2020;Drakhlis *et

al.*, 2021). The epicardium is an important source of IGF2, a potent cardiac mitogen involved in

cardiac wall thickening (Shen *et al.*, 2015;Perez-Pomares and De La Pompa, 2011). It also

generates cardiac fibroblasts and smooth muscle cells (SMCs) that invade the heart following an

epithelial-to-mesenchymal transition (Smits *et al.*, 2018). Similarly, in our organoid model, we

observed cardiomyocyte expansion, as well as the later emergence of SMCs and endothelial cells

(which occurs at >5 weeks of gestation in human fetal heart)(Smits *et al.*, 2018;Hirakow, 1983).

We propose that the co-emergence of multiple key progenitor cells contributed to the cellular and

structural complexity of the cardiac compartment of the multilineage organoids.

The presence of gut tissue developing alongside the cardiac tissue improved the

phenotypic and functional maturation of the organoid cardiomyocytes, whose structural features

resembled those of native third-trimester human fetal heart. To date, more mature features of

cardiomyocytes have only been achieved *in vitro* by subjecting early contractile iPSC-derived

cardiomyocytes on a 3D collagen gel to an increasing regimen of electrical stimulation

(Ronaldson-Bouchard *et al.*, 2018). Additionally, the prevailing view of atrial and ventricular

cardiomyocyte specification is that the different cardiomyocyte subtypes derive from distinct

mesoderm populations (Lee *et al.*, 2017;Devine *et al.*, 2014). Our organoid model shows that the

emergence of cardiomyocytes with an atrial/nodal phenotype occurs in parallel with the

formation of primitive gut: whereas at day 20, multilineage organoids contained equivalent

amounts of ventricular (MLC2V+) and atrial/nodal (MLC2V-) cardiomyocytes, by day 30, when

primitive mid-hindgut (MHG) structures emerged, the proportion of atrial/nodal cardiomyocytes

was greater **(Figure 2 D,E)**. Therefore, we speculate that similar to other endoderm subtypes

MHG endoderm secretes factors that impact the development of the heart, and specifically

promotes atrial/nodal specification. Indeed, previous studies have shown that other endoderm

derivatives, such as anterior visceral endoderm and anterior foregut, promote cardiac mesoderm

specification and formation of new cardiomyocytes through the secretion of cardiac-inductive

factors, like Activin A and FGF2 (Sugi and Lough, 1995;Mummery *et al.*, 2003;Anderson *et al.*,

2016;Lough and Sugi, 2000). FGF2 is also expressed by the epithelial, stromal, and SMCs in the

intestine, but its effects on gut development have not been fully defined (Dignass and Sturm,

2001;El-Hariry *et al.*, 1997). Our multilineage organoid model of cardiac and gut co-emergence

offers a novel platform to investigate how specific neighboring tissues synergistically interact to

promote co-development and functional maturation.

The complexity of the intestinal tissue in multilineage organoids mimics that of the small

intestinal mucosa that forms adjacent to enteric SMCs. The formation of an SMC layer in close

proximity to the gut epithelium in our multilineage organoids is notable since SMCs have so far

only been observed in gut organoids after *in vivo* transplantation (Munera *et al.*, 2017).

Furthermore, the SMCs in multilineage organoids exhibited spontaneous peristaltic-like

contractility, which has only been observed in fully mature cells *in vivo*, or in *in vitro* cultures of

primary cells together with stromal cells and gut pacemaker cells (interstitial cells of Cajal)

seeded onto a scaffold (Gays *et al.*, 2017;Kobayashi *et al.*, 2018). Other gut organoid models

form an intestine-like mucosa only upon embedding in an exogenous extracellular matrix (i.e.

Matrigel) (Munera *et al.*, 2017;Sato *et al.*, 2009). Thus, our organoid model creates a

microenvironment permissive for gut tissue formation in the absence of exogenous extracellular

matrix components or morphogenic signaling cues, likely due, in part, to the enrichment of ECM

components (e.g. glycosaminoglycans and hyaluronic acid) detected in the sub-epicardial/

peritoneum-like region where the gut epithelium clusters arise.

While multilineage organoids arose robustly from all the aggregated spheroids within a

single experiment, not all differentiation experiments gave rise to multilineage organoids. In

those alternative instances, we obtained gut organoids or cardiac microtissues instead, at similar

frequencies, from the same initial aggregation and culture conditions. Gene expression analysis

of the starting populations that yielded the different phenotypic outcomes indicated that

multilineage organoids result from progenitors with enriched expression of heart-related

patterning TFs, cardiomyocyte contractile machinery related genes, and SOX17, a definitive

endoderm master regulator gene. By contrast, cardiac microtissue progenitors expressed a

transcriptome that was less enriched for heart cell related genes, and gut organoid progenitors

strongly expressed the endoderm marker, SOX17. The expression of endoderm-specific genes at

the progenitor stage suggests that not all cells undergo the intended cardiac mesoderm

commitment, and that the subsequent emergence of intestinal tissue in the multilineage and gut

organoids results from the permissive culture medium supporting the expansion and

differentiation of the endoderm progenitors. The analysis of genes encoding secreted molecules

in the gut organoid progenitors revealed high expression of FGF10, which functions as an

autocrine and paracrine signaling molecule in heart development as well as in the gut, where it

regulates the balance between goblet and Paneth cells (Al Alam *et al.*, 2015;Rochais *et al.*,

2014;Kelly *et al.*, 2001). Our multilineage organoid model can be useful for the future dissection

of paracrine interactions between cardiac and gut tissues, and identification of specific factors

that promote morphogenesis and maturation of adjacent tissues.

Future efforts can also focus on more robustly controlling the proportion of mesoderm

and endoderm cells in the initial progenitor pool. By modulating the expression of cardiac and

gut regulatory genes in distinct sub-populations of progenitors, it should be possible to reduce

batch-to-batch variability when generating multilineage organoids. Genetic engineering

approaches have been demonstrated to control cellular patterning and subsequent adoption of

divergent cell fates – fundamental first steps to regulate phenotypic co-emergence and

morphogenesis of human iPSC-derived organoids(Libby *et al.*, 2019;Libby *et al.*, 2018;Mackay

et al., 2014). In addition, we envisage that "mosaic" transcriptional modulation approaches can

be expanded for the development of multilineage organoids that support the co-emergence of

distinct tissues other than heart and gut.

Overall, our model represents a new class of organoids characterized by the co-

development and maturation of two distinct tissues arising from a mixed pool of mesendoderm

progenitors. This organoid model opens new avenues to study genetic diseases that affect both

the heart and the gut that have been challenging to investigate, such as chronic atrial and

intestinal dysrhythmias (Chetaille *et al.*, 2014). Organoids have been primarily developed with

the intent to model individual tissues in isolation (Munera *et al.*, 2017;Takasato *et al.*, 2016), and

to date only limited attempts have been made to generate more complex and heterotypic

organoids, mostly via fusion of pre-differentiated cells and/or organoid tissues (Xiang *et al.*,

2017;Koike *et al.*, 2019;Bagley *et al.*, 2017). In contrast, multilineage organoids demonstrate that

two specific tissues from distinct germ lineages can arise in parallel within a single spheroid to

synergistically support each other's development, and thus will be useful to advance the

understanding of cell or tissue interactions during early tissue specification, structural and

functional maturation, and/or disease development. Although they may not mimic all aspects of

human development, multilineage organoids do offer a more direct view into tissue cross-talk

mechanisms than alternative *in vitro* systems or animal models currently in use. Multilineage

organoids represent a logical next step in the generation of more physiological *in vitro* models of

human development*,* and therefore broaden the possibilities for modeling early embryogenesis or

complex multi-organ disorders *ex vivo*.

Acknowledgments:

The authors would like to thank the Stem Cell Core (Dr. Po-Lin So), Histology and Light

Microscopy Core (Dr. Meredith Calvert) and Genomics Core (Dr. Natasha Carli and Jim

McGuire) at Gladstone Institutes, and Electron Microscopy Lab (Dr. Danielle Jorgens) at UC

Berkeley for all the help and expertise shared. The authors thank Dr. Bruce Conklin's laboratory

for providing WTC11 GCaMP iPS cell line, Dr. Deepak Srivastava's laboratory for providing

antibodies against MYL2 and MYL3, and Dr. Katja Schenke-Layland for providing human fetal

heart paraffin blocks. A special thanks to Drs. Tracy Hookway, Irfan Kathiriya, Sanjeev Ranade

and Elphège Nora for critical discussions of this work. The authors also acknowledge the

Gladstone Scientific Editing Department (Drs. Francoise Chanut and Kathryn Claiborn) for

assistance with manuscript editing. This work was supported by the California Institute of

Regenerative Medicine Grant (LA1_C14-08015) and the Gladstone BioFulcrum Heart Failure

Research Program.

Author contributions: A.C.S. and T.C.M. conceived the study, interpreted the data, and wrote

the manuscript. A.C.S. performed iPSC differentiations, microtissues formation and

maintenance, histology studies, preparation of the cells for patch clamp analysis, calcium

imaging, bulk and single-cell RNA-seq experiments and scRNA-seq analysis. O.B.M. acquired

and analyzed calcium flux analysis, assisted on microtissues maintenance. M.H.L. performed

patch clamp data acquisition and analysis. D.A.J. developed the python script for calcium flux

and patch clamp analysis, performed the alignment of single-cell and bulk RNA-seq data and

assisted with the single-cell and bulk RNA-seq analysis. M.A.K. and V.N. prepared bulk RNA-

seq libraries and assisted in microtissue maintenance. D.T. performed light sheet microscopy and

analysis. A.P.B. performed the *in vivo* cell classifier analysis and with B.G.B. contributed to

interpretation of the results. M.A. and B.G.B. assisted with bulk RNA-seq data analysis and

B.G.B. contributed to experimental design and interpretation of results. O.B.M., D.A.J., M.A.K,

V.N., M.A., and B.G.B. contributed to manuscript editing and discussion.

Competing interests: T.C.M. is a consultant for Tenaya Therapeutics. B.G.B. is a co-founder of

and owns equity in Tenaya Therapeutics. The other authors declare no competing interests.

Figures:

Figure 1. Derivation of a multilineage organoid with complex cardiac structures. **A-B** Overall

schematic of conventional and multilineage microtissue formation. **C.** Brightfield images of

conventional and multilineage microtissues over the course of 30 days in cell culture; high

magnification views of the regions within white squares (right). Multilineage microtissues

develop a compact dark core of cells (arrowheads) and epithelial-like structures (dashed outlines)

at the surrounding translucent region. **D.** Calcium flux transients visualized by GCAMP are

limited to the center area of the multilineage microtissues, whereas they occur throughout the

conventional microtissues. Dashed line: microtissue boundaries. **E.** Multilineage microtissues

increase in surface area throughout a 30-day culture. Representative experiment. Conventional

vs. Multilineage: ****$p<0.0001$. **F-I.** Structural dynamics and cell composition of conventional

and multilineage organoids over the course of 30-40 days in culture. **F.** Representative light-

sheet imaging of microtissues stained for cardiac Troponin T (cTnT) and TBX18. Conventional

microtissues are composed of cardiomyocytes positive for cTnT covered by a continuous outer

layer of epicardial-like cells (cTnT-/TBX18+). However, the epicardial coverage shrinks over

time to small patches of cells. Multilineage microtissues contain a core of cardiomyocytes

(cTnT+/TBX18-) surrounded by stromal-like cells positive for phalloidin and a continuous outer

layer of epicardial-like cells (cTnT-/TBX18+). Like in conventional microtissues, the epicardial

layer of the multilineage microtissues suffers some disruption throughout the course of the

culture, but overall is better preserved over time. Arrowhead: epicardial layer. **G.** Multilineage

microtissues evidence a collagen IV network and MYL2+ cardiomyocytes that cluster at the

center of the microtissues. Conventional microtissues demonstrate a network of collagen IV at

day 20 (d20), but it was not evident at later stages. No MYL2+ positives cells were detected in

conventional microtissues. **H.** Multilineage microtissues exhibit more actively cycling (ki67+)

cardiomyocytes and non-cardiomyocytes than conventional microtissues. **I.** At day 30,

multilineage organoids develop epithelial-like structures positive for mid-hindgut markers

(*FOXA2*/CDX2). **J.** Schematic of the main contrasting features between conventional and

multilineage microtissues. Due the fact multilineage microtissues contained epithelial endoderm,

displayed greater cardiac structural complexity, and exhibited extensive growth dynamics, they

were hereafter referred to as multilineage organoids.

Figure 2. Long-term culture of multilineage organoids recapitulates early heart and gut

development. **A.** Brightfield microscopy images of conventional and multilineage tissues over

days 30 to 80 of culture. Conventional microtissues are viable until day 80 and beyond, whereas

conventional microtissues degenerate. **B.** Stereoscope macroscopy (top) and brightfield (bottom)

images of day-100 multilineage organoids. **C.** Multilineage organoids display an increase in

surface area until day 100 of culture. Representative experiment. Conventional vs. Multilineage:

p<0.005, ****p<0.0001. **D-F. Structural dynamics and cell composition of multilineage

organoids over the course of the culture as revealed by Russel-Movat stain (D) and fluorescent

immunostaining (E-F). **G-H.** Representative images (G) and traces (H) of spontaneous calcium

flux transients of the smooth muscle cells (ROI 1) and cardiomyocytes (ROI 2) of a multilineage

organoid imaged via light sheet microscopy (Movie S6). **I.** Transmission electron microscopy of

the gut structures (i-v), epicardium/subepicardium (vi-vii) and cardiac core (viii-ix). GAG,

glycosaminoglycans; Ep, epithelial-like structures; subEp, subepicardium-like tissue; Endo,

endothelial cells network; Arrowheads, goblet cell (iii); Dashed line, Paneth cell (iv). **J.** The

number of actively cycling cells (Ki67+) decreases over time in multilineage organoids.

Figure 3. Single-cell transcriptomic analysis at day 100 reveals complex cellular heterogeneity

of multilineage organoids. **A.** UMAP plot of all cellular populations identified. Light gray:

Seurat cell clusters analyzed using an *in vivo* heart cells classifier. **B.** Cluster tree analysis of the

cell clusters with top differentially expressed genes defining each cluster (left) and gene ontology

analysis of associated biological processes (right). **C-D.** A sub-clustering analysis of the gut cells

evidenced 3 main cell clusters (C) distinguished by the expression of crypt-related or villi-related

genes (D): cluster 0 composed mainly of crypt cells (OLFM4), such as gut stem cells (LGR5)

and Paneth cells (LYZ); and cluster 1 and 2, constituted by cells at the villi, the enterocytes,

identified by the expression of genes related to transmembrane mucins (MUC13, MUC17),

digestion (ANPEP, FABP2, SI) and cholesterol transporter genes (NPC1L1), as well as APOB, a

gene expressed in the small intestine. **E-F.** Subset analysis of the cardiomyocyte cluster (E) and

top-6 genes of the three sub-clusters (F) identify clusters 0 and 2, composed of more immature

cardiomyocytes, and cluster 1, composed of more mature cells. **G.** Violin plots of secondary

cardiomyocyte analysis depicting expression of key cardiomyocyte subtype genes for

ventricular- (top) and atrial-like cells (bottom). **H.** Identity of the cardiac-like cells of

multilineage organoids, as determined by a cell classifier trained on expression data from human

fetal heart cells. The classifier was applied to the multilineage organoid cardiomyocyte,

fibroblast and smooth muscle cells (SMC) clusters identified by Seurat analysis. Data displayed

include cell percentage and assignment probability of transcriptomic resemblance with *in vivo*

heart cells. **I.** UMAP plot of *in vivo* human developing heart cells and *in vitro* multilineage

organoid cells classified as atrial cardiomyocyte, subepicardial cells, fibroblast-like, smooth

muscle cells/fibroblast-like cells (probability greater or equal to 0.85). The UMAP projection

demonstrates that *in vitro* multilineage organoid cells align closely to the *in vivo* human heart

cells suggesting that the transcriptome of the multilineage organoid heart cells closely mimic that

of human developing heart cells, and favors atrial cardiomyocyte specification.

Figure 4. Cardiomyocytes from multilineage organoids display phenotypic and functional

maturation throughout culture, and preferentially specify into atrial/nodal-like cells. **A.**

Multilineage organoids demonstrated a reduction in spontaneous beating throughout culture. **B.**

Representative calcium flux traces at day 30 and 80 of culture. **C.** Cardiomyocytes in

multilineage organoids display mature calcium flux features by showing higher amplitude, Vmax

upstroke and downstroke (Conventional vs. Multilineage, **p<0.01, **p<0.005, ***p<0.001,

****$p<0.0001$). **D.** Cardiomyocytes' ability to respond to electrical stimuli throughout culture

time. Black dashed line: electric stimuli applied**. E.** Representative immunofluorescence images

of dissociated cardiomyocytes from conventional microtissues and multilineage organoids. **F.**

Cardiomyocyte structural feature metrics. **G.** Cardiomyocyte sarcomere classification (from class

1—immature, undefined sarcomeres—to class 4—mature, parallel, striped sarcomeres). **H.** Day-

100 multilineage organoid cardiomyocyte sarcomeres (i), intercalated disks (ii, arrow heads) and

glycogen reservoir ultrastructure (iii). **I.** Electrophysiological characteristics of cardiomyocyte

subtypes. **J.** Percentage of ventricular and atrial/nodal cardiomyocytes dissociated from

conventional microtissues and multilineage organoids. **K.** Representative electrophysiological

traces of ventricular and atrial/nodal cardiomyocytes in multilineage organoids at day 30 of

culture.

Figure 5. Multilineage organoid progenitor cells have a transcriptomic profile that supports the

co-development of cardiac and gut tissues. **A.** Mesendoderm progenitors cultured in AA medium

sometimes failed to develop into multilineage organoids, and instead yielded cardiac

microtissues or gut organoids. Brightfield and histological analysis revealed phenotypic diversity

between these outcomes. **B.** Experimental work flow schematic. **C.** Multidimensional scaling

(MDS) plot describes similarity between the progenitor cells that give rise to cardiac

microtissues and multilineage and gut organoids. **D.** Heat map of the pairwise comparison of

multilineage organoids and cardiac microtissues progenitors evidence a gene transcription profile

that favors muscle tissue formation. **E.** Heat map of the pairwise comparison of multilineage

organoid and gut organoid progenitors and respective GO terms and genes demonstrated that

multilineage progenitor gene expression supports cardiac tissue formation. **F-G.** Targeted

pairwise comparison for cell type-specific transcription factor (TF) genes (F) and secreted factor

genes (G) in multilineage vs. cardiac (left) and multilineage vs. gut (right) progenitor cells. The

multilineage organoid progenitors expressed key heart and endoderm-related TFs and secreted

factors highly expressed in gut organoid and cardiac microtissues progenitors.

640
641
642
643   Supplementary Materials for
644
645   Developmental co-emergence of cardiac and gut tissues modeled by human iPSC-derived
646   organoids
647

648   A.C. Silva, O.B. Matthys, D.A. Joy, M.A. Kauss, V. Natarajan, M.H. Lai, D. Turaga, A.P. Blair,
649   M. Alexanian, B.G. Bruneau, T.C. McDevitt
650
651   Correspondence to: todd.mcdevitt@gladstone.ucsf.edu

652
653
654   **This PDF file includes:**
655
656   Materials and Methods (MM)
657   Figure MM1
658   Figs. S1 to S7
659   Table S1 to S3
660   Captions for Movies S1 to S8
661
662   **Other Supplementary Materials for this manuscript include the following:**
663
664   Movies S1 to S8
665
666
667

Materials and Methods

Human induced pluripotent stem cell culture

WTC11 human induced pluripotent stem cells (iPSCs) genetically modified with fast green

fluorescent calcium indicator GCaMP6f (Huebsch *et al.*, 2015;Mandegar *et al.*, 2016) were

maintained on tissue culture plates coated with 80μg/mL growth factor-reduced (GFR) Matrigel

(Corning) and fed daily with mTeSR1 medium (StemCell Technologies). When the iPSCs

reached ~70% confluency, about every 3 days, they were dissociated to single cells using

Accutase (Innovative Cell Technologies) and re-plated at density of $1x10^4$ cells/cm^2 in mTeSR1

medium containing 10μM Y-27632 ROCK inhibitor (ROCKi; Selleckchem) for 24h.

Differentiation to mesendoderm-derived progenitors

GCaMP-WTC11 iPSCs were differentiated to mesendoderm-derived progenitor cells following a

chemically-defined, serum-free protocol modulating Wnt/β-catenin signaling (Lian *et al.*, 2012).

Briefly, iPSCs were seeded at a density of $3x10^4$ cells/cm^2 and fed with mTeSR1 medium until

100% confluency was reached (~3 days; termed differentiation day 0). At day 0, cells were fed

with RPMI1640 medium with B27 Supplement minus insulin (denoted as RPMI/B27-; Life

Technologies) supplemented with 12μM CHIR-99021 (Selleckchem). Exactly 24h later, the

medium was changed to fresh RPMI/B27-. On day 3, cells were fed with RPMI/B27-

supplemented with 5μM IWP2 (Sellekchem) for 48h. On day 5, cells were either dissociated for

tissue formation or fed with RPMI/B27- medium.

Mesendoderm progenitor cell spheroid formation and culture

Mesendoderm progenitor cells were dissociated at differentiation day 5 with Accutase and

aggregated as previous described (Hookway *et al.*, 2016). Briefly, dissociated cells in

RPMI/B27- medium with 10μM ROCKi were seeded into 800μm pyramidal agarose wells at a

density of 2.4×10^4 cells/well. After 24h, cell spheroids were removed from the agarose wells by

gently pipetting using wide bore tips. Spheroids were maintained in rotary suspension culture at

a density of 65 spheroids/mL in 6-well low-attachment tissue culture plates (Hookway *et al.*,

2016). On day 7 of the differentiation protocol, 2 days after tissue aggregation, medium was

changed to either conventional cardiac-permissive medium, RPMI/B27+ (RPMI1640 medium

with B27 supplement with insulin; Life Technologies)(Lian *et al.*, 2012) or epicardial-permissive

medium (Advanced DMEM/F12 (Gibco) with 100μM L-Ascorbic Acid (Sigma))(Bao *et al.*,

2016) and refreshed every 3 days thereafter.

Calcium imaging and functional analysis

Calcium flux was analyzed by recording fluorescence of GCaMP signal in the spheroids. At day

30, 40, 50, 60, 70, 80, 90*, 100*, and 365*, conventional and multilineage microtissues (n = 4-

17 per condition) were equilibrated in Tyrode's solution (137 mM NaCl, 2.7 mM KCl, 1 mM

$MgCl_2$, 0.2 mM Na_2HPO_4, 12 mM $NaHCO_3$, 5.5 mM d-glucose, and 1.8 mM $CaCl_2$; Sigma-

Aldrich) for at least 30min at 37°C and then imaged on a Zeiss Axio Observer Z1 inverted

microscope equipped with a Hamamatsu ORCA-Flash 4.0 sCMOS camera. Electrical field

stimulation between 0.5Hz and 8Hz was applied with the MyoPacer (IonOptix) to evaluate

functional properties and eliminate differences in intrinsic beat rate between individual

spheroids. Calcium transients were recorded using Zen Professional software (v.2.0.0.0) with

10ms exposure and 100 frames per second. Circular regions of interest (ROI) with a diameter of

59.52 pixels were selected in the beating core of the spheroids and the mean fluorescence

intensity of the calcium transients was plotted against time. Additional metrics of calcium

handling properties, such as amplitude, stroke velocities, and beat rate, were analyzed using a

custom Python-script. Source code is available at https://github.com/david-a-joy/multilineage-

organoid (made public upon acceptance). *Of note, conventional samples were not evaluated at

these timepoints, because they were only viable until day 80 of culture.

Calcium imaging with light sheet microscopy

In order to assess calcium activity at higher resolution, light sheet fluorescence microscopy

(LSFM) was used to optically section live multilineage organoids and conventional microtissues

and record calcium transients *in situ* (Turaga *et al.*, 2020). Organoids/microtissues were

suspended in 1.5% low-melt agarose made up in Tyrode's solution in size 2 glass capillaries

(~1mm inner diameter; Zeiss). The Zeiss z.1 light sheet microscope used for imaging was

equipped with two PCO.edge sCMOS cameras, 10x 0.2 NA illumination lens, 20x 1.0 NA

detection lens, and 488nm laser, and the live samples were maintained at 37°C. 3D calcium

imaging stacks were acquired at ~120ms per frame and each z-spacing was 0.86μm per frame.

Single optical sections (3-4μm light thickness) were excited at 488nm and imaged at ~20Hz for

52 seconds (1000 total frames).

Patch clamp analysis

Conventional microtissues (n = 10 per condition), and multilineage organoids (n = 5-10 per

condition), as well as conventional 2D differentiated cardiomyocytes, were washed in PBS and

dissociated by successive 20min incubations in 0.25% Trypsin at 37°C (1 incubation for

monolayers and maximum of 5 incubations for tissues) until complete dissociation to single

cells. Dissociated cells at a density of 0.03×10^6 cells/cm^2 were seeded onto GFR Matrigel-coated

glass coverslips for patch-clamp recordings in medium containing 10μM ROCKi. 24h after

seeding, cells were kept in culture media without ROCKi until the experiments began. To

conduct experiments, coverslips containing hiPSC-CMs were transferred to a heated RC-26

recording chamber (Warner Instruments), mounted to a Nikon inverted microscope and

continuously perfused with warmed Tyrode's solution. Experiments were performed at 35 ± 1°C.

Action potentials were recorded from spontaneously beating cardiomyocytes using the perforated

patch technique with borosillicate glass pipettes (Sutter Instruments) with typical resistance of 2-

4 MΩ. The Tyrode's solution consisted of (in mM): 140 NaCl, 5.4 KCl, 1.8 $CaCl_2$, 1 $MgCl_2$, 10

glucose, and 10 HEPES, with the pH adjusted to 7.4 with NaOH. The patch pipette solution

consisted of (in mM): 150 KCl, 5 NaCl, 5 MgATP, 10 HEPES, 5 EGTA, 2 $CaCl_2$, and 240

μg/mL amphotericin B, with the pH adjusted to 7.2 with KOH. Data were acquired in current

clamp mode using an Axopatch 200B amplifier and Digidata 1440A digitizer coupled to pClamp

10 software (Molecular Devices). Data were digitized at 10 kHz and filtered at 1 kHz. Action

potential parameters were determined using Clampfit 10 software within pClamp. hiPSC-derived

cardiomyocytes were classified as either ventricular or nodal/atrial subtype after considering

multiple action potential parameters, including the APD_{90}/APD_{50} ratio, APD_{30-40}/APD_{70-80} ratio,

MDP, peak, amplitude, frequency, and dV/dT_{Max} (Ma *et al.*, 2011;Churko *et al.*, 2018).

Cardiomyocyte structure analysis (area, perimeter, AR, circularity and sarcomere score)

Day-30 and 80 conventional* (n =10 per condition) and multilineage microtissues (n = 5-10 per

condition) were dissociated into single cells as described above (patch clamp analysis sections)

and seeded onto GFR Matrigel-coated glass coverslips. 24h post-seeding, the cells were fed with

fresh medium without ROCKi for at least 48h to allow the cells to acquire their natural

morphology. Coverslips were then fixed in 4% paraformaldehyde (EMS), stained with the

cardiomyocyte-specific marker cardiac Troponin T (cTnT), labeled with Hoechst to detect

nuclei, and mounted on glass slides using ProLong™ Gold Antifade Mountant (ThermoFisher

Scientific). Representative images were acquired for each sample condition using an inverted

microscope (Zeiss Axio Observer Z1). For each cell, area, perimeter, circularity, and aspect ratio

(AR) were quantified using Fiji software. In addition, each cardiomyocyte was assigned a score

for sarcomere development between 1 and 4, defined as: 1) unaligned sarcomeres and absence of

a striped pattern; 2) continuous sarcomeres, well ordered but not in parallel; 3) continuous

sarcomeres, organized mostly in parallel, but with less-defined striped pattern; 4) continuous

sarcomeres with striped pattern, organized mostly in parallel **(Figure M1)**(Nguyen *et al.*,

2014;Judge *et al.*, 2017). The percentage of cells in each class of sarcomere development was

calculated per sample and timepoint. * Of note, conventional samples were not evaluated at day

80, because only a few microtissues were available and viable at this later stage.

Figure M1. Representative images of the different classes of sarcomere development.

Histology, immunostaining, and in situ hybridization

Conventional microtissues (n=10 per condition), cardiac organoids (n=10 per condition),

multilineage organoids (n=5-10 per condition), and gut (n=5-10 per condition) organoids were

fixed in 10% neutral buffered formalin (VWR) with 0.03% eosin (Newcomer Supply) at room

temperature for 1h and washed with PBS. Samples were embedded in HistoGel Specimen

Processing Gel (ThermoFisher Scientific), paraffin processed and embedded, cut into 5μm

sections, and adhered to positively charged glass slides. Slides were then deparaffinized with

xylene, re-hydrated through a series of decreasing ethanol concentrations (100%, 100%, 95%,

80%, 70%), and rinsed in tap water. Hematoxylin and eosin (H&E), Russell-Movat

Pentachrome, and Safranin O stainings were performed according to manufacturer's instructions

and the slides were mounted with Cytoseal 60 (Richard-Allan Scientific) and glass coverslips.

For immunofluorescence staining, heat-induced epitope retrieval was performed by immersing

slides in Tris-EDTA buffer (Cellgro) pH 9.0 or citrate buffer (Vector Laboratories) pH 6.0 in a

95°C water bath for 35min. Slides were cooled for 20min at RT and washed 3x with PBS.

Samples were permeabilized by slide immersion in 0.2% or 0.5% Triton X-100 (Sigma) in PBS,

washed 3x in PBS, and blocked in 1.5% normal donkey serum (NDS; Jackson

ImmunoResearch). Primary and secondary antibody cocktails were diluted in blocking solution

(Table S3). 3 PBS washes were performed after primary (overnight, 4°C) and secondary

antibody (1h, RT) incubations. For in situ hybridization analysis, RNAscope Multiplex

Fluorescent Reagent Kit v2 was used on paraffin sections, following user manual #323100-USM.

Human RNAscope probes used in this study were: WT1-C2 (catalog #415581); FOXA2-C1

(catalog #832221); TBX18-C3 (catalog #832231). Nuclei were counter-labeled with Hoechst and

coverslips were mounted on slides using ProLong™ Gold Antifade Mountant. Samples were

imaged on the Zeiss Axio Observer Z1.

Whole-mount immunostaining

Conventional microtissues (n=5) and multilineage organoids (n=5) were fixed in 4%

paraformaldehyde at room temperature for 1h with agitation (40 rpm). Samples were then

washed 3x in stain buffer (2% NDS, 0.1% Tween-20 (Sigma-Aldrich) in PBS) and

permeabilized for 30min in 1.5% Triton X-100, followed by a 3h incubation in blocking buffer

(3.5% NDS, 0.1% Tween-20 in PBS). Microtissues/organoids were incubated with TBX18 and

cTnT primary antibodies overnight at 4°C and secondary antibodies plus phalloidin and Hoechst

stains for 4h at RT **(Table S3)**, followed by 3 final washes in stain buffer prior to imaging.

Whole-spheroid structural analysis by light sheet microscopy

Whole-mount immunostained conventional microtissues and multilineage organoids were

suspended in 1.5% low-melt agarose (made up in PBS; IBI Scientific) in size 2 glass capillaries

immediately prior to imaging. Microtissues/organoids were each imaged at three angles (120°

rotations between views), and then stitched with multi-view reconstruction to provide isotropic

resolution throughout the sample, as previously described (Turaga *et al.*, 2020). Volumetric

reconstruction and size analyses were performed using Imaris 8.3 software.

Single-cell RNA sequencing

Multilineage and gut organoids at day 100 of culture (n = 5-8) were washed in PBS and

dissociated by successive 20-min incubations in 0.25% trypsin at 37°C (maximum of 5

incubations) until complete dissociation to single cells. 10,000 cells in PBS supplemented with

0.08% FBS were prepared for sequencing by droplet encapsulation using the Chromium

Controller and library preparation with the 10x Genomics Single Cell 3' v2 Library and Gel

Bead Kit (10x Genomics). Libraries were sequenced using the NovaSeq S4 (Illumina) to a

minimum depth of 50,000 mean reads per cell. Sequences were demultiplexed and aligned to

human reference genome grch38 using *CellRanger* v3.0.2, and downstream analysis was

performed with Seurat v.3.1. Cells with fewer than 1,500 genes were excluded from analysis and

cell identity was assigned to each cluster by cross-referencing cluster marker genes with known

cardiac and gut markers. Cell-cycle did not introduce heterogeneity in scRNA-seq data. Raw

data is available at Geo under the accession number GSE153075 (access token:

qvslsuecdpofjup).

Classification of In Vivo Cardiac Cell Types

To evaluate the transcriptomic similarity between multilineage organoid heart cells and human

fetal heart cells, an *in vivo* heart cell classifier was applied. A multiclass logistic regression (lr)

model was trained on *in vivo* cardiac cell types(Asp *et al.*, 2019). To have consistent features, we

ran SCTransform (default, version 3.1.5.9) independently on each dataset and then took an

intersection of their top 3,000 highly variable genes. A total of 1,418 genes, transformed as

Pearson residuals, were used to train the model. The resulting expression matrix of 3,717 single

cells from a 6.5–7 post conception week (PCW) heart with established cardiac cell types was

used to train the model using a 'sag' solver and cross-entropy loss cost function with the Python

package sklearn (version 0.22.2.post1).

A two-dimensional UMAP (version 0.4.4) latent space was fit to *in vivo* subepicardial cells,

fibroblast-like, smooth muscle cells/fibroblast-like, and atrial cardiomyocytes. We used the lr

model's feature space for UMAP dimensionality reduction. A sklearn C-support vector classifier

was then trained on the four *in vivo* cell types using the UMAP latent space. *In vitro* cells that the

lr model classified as one of the four *in vivo* cell types, with probabilities greater than or equal to

eighty-five percent, were transformed into the UMAP latent space. The trained C-support vector

classifier was then applied to the *in vitro* cell's UMAP transform embedding. The Seaborn

(version 0.10.1) library was used to visualize UMAP transformations.

Bulk RNA sequencing

To reduce batch effects between RNA-sequencing analysis of the mesendoderm-derived

progenitors between experiments, bulk RNA-sequencing was applied. Day 5 progenitor cells that

gave rise to cardiac (n=2), multilineage (n=1), or gut organoids (n=2) were lysed with Trizol

(Ambion) and RNA was extracted using the Direct-zol Miniprep kit (ZymoResearch) and

quantified using the NanoDrop 2000c (ThermoFisher Scientific). RNA-seq libraries were created

using the SMARTer Stranded Total RNA Sample Prep Kit (Takara Bio) and sequenced using

NextSeq500/550 High Output v2.5 kit (spiked with 10% PhiX; Illumina) to a minimum depth of

29 million reads per sample. The sequences were aligned to grch38 using HiSat2 (Kim *et al.*,

2019), reads were quantified using featureCounts (Liao *et al.*, 2014), and differential expression

was determined using GFOLD(Feng *et al.*, 2012). Relative gene expression level was determined

using the "l" normalization algorithm (Zhang *et al.*, 2019). Raw data is available at Geo under

the accession number GSE153071 (access token: utwxqsawhjcpfgz).

Statistical analysis

All results were obtained in three independent experiments, unless otherwise stated, and shown

as mean ± standard error of the mean (SEM). The statistical analysis of the data was performed

using Prism 8 with statistical significance determined at $p < 0.05$. Tukey's multiple comparison

test was applied. Whenever normality was verified through D'Agostino-Pearson omnibus

normality test, the t-Student test was applied and the Independent Samples Mann-Whitney U test

was used for the remaining analyses. A blinded approach (labeling samples with an

alphanumeric code) was implemented for analyzing cardiomyocyte morphological features (area,

perimeter, AR, circularity, sarcomere scoring), patch clamp, histology, and immunostaining.

Figure S1. *Related to Figure 1 and 2.* Comparison of conventional and multilineage

microtissues with human heart development (**A-B**). **A.** RNAscope validation of TBX18 and WT1

transcripts in native human heart tissue, multilineage organoids, and conventional microtissues.

Dashed line denotes boundary of the cardiac core. **B.** Identification of cardiomyocytes

(MYH7+SMA-, MYH7+SMA+), smooth muscle cells (SMA+MYH7-), and endothelial cells

(Pecam-1+) in native human heart tissue, multilineage organoids, and conventional microtissues.

C. Proliferating cardiomyocytes in human ventricle wall tissue (MYL3+Ki67+), multilineage

organoids, and conventional microtissues (slow/fast TnI+Ki67+). **D.** Comparative images

exhibiting similarities between the cardiac compartment of native human fetal heart tissue and

that of the multilineage organoids.

Figure S2. *Related to Figure 1 and 2.* Multilineage organoid histological and volumetric

analysis. **A.** Detailed light-sheet images of the epicardial layer (TBX18+) of conventional and

multilineage microtissues. **B.** Conventional and multilineage microtissues demonstrate the

presence of cardiomyocytes (MYH7+), smooth muscle cells (SMA+) and MYH7+/SMA+

cardiomyocytes (d20, more immature). **C.** Total volume (blue), cardiac core volume (yellow),

and ratio of cardiac:total volume (black) of the multilineage organoids at days 10 and 30 of

culture. *$p<0.05$, **$p<0.005$. **D-E.** Epithelial-like structures in multilineage organoids

correspond to primitive midgut based on the detection of FOXA2 transcript by RNAscope (D)

(absent in conventional microtissues) and CDX2 protein by immunostaining (E). **F.** Hyaluronic

acid (HA) is expressed in the interstitial tissue (tissue surrounding the cardiac core and the gut)

in the multilineage organoids. White squares regions in high magnification at the bottom panels.

Dashed line boundary of the cardiac core and gut.

Figure S3. *Related to Figure 3.* Transcriptional characterization of day-100 multilineage organoids. **A.** Dotplot of key genes that define cell type identity of UMAP clusters as cardiomyocytes (CM), proliferating cells (Prol.), smooth muscle cells (SMC), fibroblasts (FB), enteroendocrine cells (EnteroE), and gut. **B.** Violin plot of VIL1 expression across different endoderm-derived tissues using GTEx tissue dataset (https://www.gtexportal.org/). **C.** Heatmap of secondary analysis of re-clustered proliferating cells identified SMCs (cluster 0) and CMs (cluster 1) as the main proliferative cell types at day 100. **D.** Violin plot of key karyokinesis-related genes in proliferating SMCs and CMs. **E.** Histological validation of SMC proliferation (yellow square; ki67+, SMA+) and gut cell proliferation (blue square; ki67+, Russel-Movat) in a day-100 multilineage organoid. Dashed line denotes SMC region. **F.** Feature plot showing co-expression of key atrial, ventricular and pacemaker related genes in day-100 organoids. **G.** Table showing the distribution of organoid heart cells classified to various cell identities by the *in vivo* cell classifier. **H.** Confusion matrix comparing test vs. predicted cell type labels for human developing fetal cardiac cells(Asp *et al.*, 2019).

Figure S4. *Related to figure 2.* 1 year-old multilineage organoids. **A.** Macroscopic field of view of multiple 1 year-old organoids. **B.** Calcium handling response to electrical stimulation (0.5-4Hz). **C-E.** Hematoxylin and eosin stain (H&E) reveals vasculature-like structures (**C**), smooth muscle cells (**D**), and intestine (**E**). Dashed line denotes cardiac core; solid arrowhead indicates gut; outlined arrowhead indicates smooth muscle cells. **F.** Immunostaining highlights the presence of cardiomyocytes (MYH7+) and smooth muscle cells (SMA+). **G.** Lectin-positive vasculature-like structures are found close to the cardiac compartment (inset) and non-specific lectin expression was also identified in gut structures (solid arrowheads). **H.** Identification of

actively proliferating cells by Ki67+ immunostaining. **I.** Safranin O staining reveals presence of

goblet cells (counterstained in red) in the gut structures.

Figure S5. *Related to Figure 4.* Cardiomyocyte functional analyses. **A.** Calcium handling

response of day 80 conventional and multilineage spheroids subjected to electrical field

stimulation at increasing frequencies (0.5-8Hz). **B**. Percentage of conventional and multilineage

spheroids capable of responding to the electrical field (0.5-8Hz) stimulation applied at day 80 of

culture. **C.** Table displaying electrophysiological properties of day-30 cardiomyocytes

differentiated in 2D in various media conditions: conventional media (RPMI/B27+,

Ca^{2+}=0.42mM), conventional media supplemented with ascorbic acid (AA) and calcium

(Conventional Suppl., AA=100 μg/ml, Ca^{2+}=0.63mM) and multilineage-permissive media

(AA=100 μg/ml, Ca^{2+}=1.05mM). **D.** Representative traces of ventricular cardiomyocytes in the

different media conditions. **E.** Percentage of ventricular and atrial/nodal cardiomyocytes across

the different media conditions.

Figure S6. *Related to Figure 5.* Analysis of day-100 gut organoids by single-cell RNA

sequencing, histology, and immunostaining. **A.** UMAP plot colored by cell identity. **B.** Feature

plot of key genes used to assign cell-type identity to UMAP clusters. **C.** Hierarchical clustering

of the different UMAP cell populations. **D.** Heatmap of top 10 differentially expressed genes that

define each UMAP cluster with associated biological process gene ontology analysis. **E-F.**

Histological analysis of gut organoids from 2 independent experiments. Hematoxylin & eosin

staining (left, E), immunostaining to identify the presence of endothelial cells (Pecam-1+),

smooth muscle cells (SMA+, solid arrowhead), and remnant cardiomyocytes (cTnT+) (right, E)

and Safranin O staining to identify the presence of goblet cells (red, F). Outlined arrowhead,

intestine tissue; *, pancreas-like tissue. **G-H.** Representative images of spontaneous calcium flux

transients (G) in gut organoids at day 12 (top, ROI1) and 60 (bottom, ROI2) of culture and

respective ROIs traces (H) (Movie S8).

Figure S7. *Related to Figure 5.* Transcriptional variability of the progenitor populations across

experiments. **A.** Spheroid phenotype diversity observed across experiments at early stages of

culture. **B.** Heat map and respective gene ontology (GO) terms analysis of cardiac microtissues (top) versus gut organoids progenitors (bottom). **C.** Heatmap of the differentially expressed genes analyzed through a target analysis of relevant cardiac- and gut-related genes in the progenitor cell populations between experiments. **D.** List of highly expressed genes encoding secreted factors in cardiac microtissues and gut organoids. **E.** Heatmap of the differentially expressed genes detailed in (**D**) in the progenitor cell populations for each experiment.

968   **Table S1.** List of genes that describe each cell cluster identity derived from Seurat analysis.

970   **Table S2.** *In vivo* cell classifier gene features and weights for each cell type prediction.

972   **Table S3.** Panel of dyes, primary and secondary antibodies used for whole mount

973   immunostaining, immunocytochemistry and paraffin sections.

Protein	Source	Catalog #	Dilution
Alexa Fluor 488 Donkey anti-mouse IgG	Invitrogen		1:400
Alexa Fluor 555 Donkey anti-rabbit IgG	Invitrogen		1:400
Alexa Fluor 647 Donkey anti-goat IgG	Invitrogen		1:400
Alexa Fluor 647 Phalloidin	Invitrogen	A22287	1:100
CDX2	R&D Systems	AF3665	1:100
Collagen IV	Millipore	AB769	1:100
Ki67	Invitrogen	Ab15580	1:500
Smooth muscle actin	DAKO	M0851	1:100
HABP	Sigma-Aldrich	H9910	1:65
Hoechst 33342	Thermo Scientific	62246	1:10000
MYL3	Abcam	Ab680	1:1200
MYH7	Sigma-Aldrich	HPA001239	1:100
MLC2-v	Abcam	Ab79935	1:200
PECAM-1	Santa Cruz	SC-1506	1:1000
TBX18	R&D Systems	MAB63371	1:100
To-Pro-3 Iodide	Thermo Fisher	T3605	1:5000
Troponin I Slow and Fast	Abcam	ab110132	1:100
Troponin T	Abcam	ab45932	1:400
Vimentin	Millipore	AB1620	1:100
WGA	Thermo Fisher	AF350	1:100

975
976

Movie S1. Spontaneous calcium transients, depicted by GCaMP fluorescence signal, of day-30

conventional microtissues.

Movie S2. Spontaneous calcium transients of day-30 multilineage microtissues.

Movie S3. Representative 3D reconstruction of day-12 multilineage microtissue by light sheet

microscopy. Total volume was quantified based on a nuclear label (blue, Hoechst) and volume of

the cardiac core was based on cardiomyocyte-specific marker, cardiac troponin T (yellow,

cTnT). Stromal-like cell population identified by phalloidin stain (red).

Movie S4. Representative 3D reconstruction of day-30 multilineage microtissue by light sheet

microscopy, with Hoechst labeling nuclei (blue) and cTnT marking cardiomyocytes (yellow).

Stromal-like cell population identified by phalloidin stain (red).

Movie S5. Lightsheet microscopy-based acquisition of the rhythmic calcium flux transients of

cardiomyocytes in day-30 multilineage organoids in the absence of electrical stimulation.

Movie S6. Lightsheet microscopy-based acquisition of the rhythmic and sporadic calcium flux

transients from cardiomyocytes and smooth muscle cells, respectively, in day-76 multilineage

organoids in the absence of electrical stimulation.

Movie S7. Calcium transients of 1-year-old multilineage organoids subjected to electrical

stimulation from 0.5-4Hz.

Movie S8. Calcium transients of day-12 and day-60 gut organoids.

References:

996   Al Alam, D., Danopoulos, S., Schall, K., Sala, F. G., Almohazey, D., Fernandez, G. E., Georgia, S., Frey, M. R., Ford, H. R., Grikscheit, T.*, et al.* 2015. Fibroblast growth factor 10 alters the balance between goblet and Paneth cells in the adult mouse small intestine. *Am J Physiol Gastrointest Liver Physiol,* 308**,** G678-90.

Anderson, C., Khan, M. a. F., Wong, F., Solovieva, T., Oliveira, N. M. M., Baldock, R. A., Tickle, C., Burt, D. W. andStern, C. D. 2016. A strategy to discover new organizers identifies a putative heart organizer. *Nat Commun,* 7**,** 12656.

Asp, M., Giacomello, S., Larsson, L., Wu, C., Furth, D., Qian, X., Wardell, E., Custodio, J., Reimegard, J., Salmen, F.*, et al.* 2019. A Spatiotemporal Organ-Wide Gene Expression and Cell Atlas of the Developing Human Heart. *Cell,* 179**,** 1647-1660 e19.

Bagley, J. A., Reumann, D., Bian, S., Levi-Strauss, J. andKnoblich, J. A. 2017. Fused cerebral organoids model interactions between brain regions. *Nat Methods,* 14**,** 743-751.

Bao, X., Lian, X., Hacker, T. A., Schmuck, E. G., Qian, T., Bhute, V. J., Han, T., Shi, M., Drowley, L., Plowright, A.*, et al.* 2016. Long-term self-renewing human epicardial cells generated from pluripotent stem cells under defined xeno-free conditions. *Nat Biomed Eng,* 1.

Brown, B. N. andBadylak, S. F. 2014. Extracellular matrix as an inductive scaffold for functional tissue reconstruction. *Transl Res,* 163**,** 268-85.

Brown, K., Doss, M. X., Legros, S., Artus, J., Hadjantonakis, A. K. andFoley, A. C. 2010. eXtraembryonic ENdoderm (XEN) stem cells produce factors that activate heart formation. *PLoS One,* 5**,** e13446.

Cao, N., Liu, Z. M., Chen, Z. Y., Wang, J., Chen, T. T., Zhao, X. Y., Ma, Y., Qin, L. J., Kang, J. H., Wei, B.*, et al.* 2012. Ascorbic acid enhances the cardiac differentiation of induced pluripotent stem cells through promoting the proliferation of cardiac progenitor cells. *Cell Research,* 22**,** 219-236.

Chakravarti, S. 2002. Functions of lumican and fibromodulin: lessons from knockout mice. *Glycoconj J,* 19**,** 287-93.

Chan, S. S., Li, H. J., Hsueh, Y. C., Lee, D. S., Chen, J. H., Hwang, S. M., Chen, C. Y., Shih, E. andHsieh, P. C. 2010. Fibroblast growth factor-10 promotes cardiomyocyte differentiation from embryonic and induced pluripotent stem cells. *PLoS One,* 5**,** e14414.

Chetaille, P., Preuss, C., Burkhard, S., Cote, J. M., Houde, C., Castilloux, J., Piche, J., Gosset, N., Leclerc, S., Wunnemann, F.*, et al.* 2014. Mutations in SGOL1 cause a novel cohesinopathy affecting heart and gut rhythm. *Nat Genet,* 46**,** 1245-9.

Churko, J. M., Garg, P., Treutlein, B., Venkatasubramanian, M., Wu, H., Lee, J., Wessells, Q. N., Chen, S. Y., Chen, W. Y., Chetal, K.*, et al.* 2018. Defining human cardiac transcription factor hierarchies using integrated single-cell heterogeneity analysis. *Nat Commun,* 9**,** 4906.

Cui, Y., Zheng, Y., Liu, X., Yan, L., Fan, X., Yong, J., Hu, Y., Dong, J., Li, Q., Wu, X.*, et al.* 2019. Single-Cell Transcriptome Analysis Maps the Developmental Track of the Human Heart. *Cell Rep,* 26**,** 1934-1950 e5.

Devine, W. P., Wythe, J. D., George, M., Koshiba-Takeuchi, K. andBruneau, B. G. 2014. Early patterning and specification of cardiac progenitors in gastrulating mesoderm. *Elife,* 3.

Dignass, A. U. andSturm, A. 2001. Peptide growth factors in the intestine. *Eur J Gastroenterol Hepatol,* 13**,** 763-70.

Drakhlis, L., Biswanath, S., Farr, C.-M., Lupanow, V., Teske, J., Ritzenhoff, K., Franke, A.,
Manstein, F., Bolesani, E., Kempf, H., *et al.* 2021. Human heart-forming organoids
recapitulate early heart and foregut development. *Nature Biotechnology*.
El-Hariry, I., Pignatelli, M. andLemoine, N. 1997. Fibroblast growth factor 1 and fibroblast
growth factor 2 immunoreactivity in gastrointestinal tumours. *J Pathol,* 181**,** 39-45.
Engels, M. C., Rajarajan, K., Feistritzer, R., Sharma, A., Nielsen, U. B., Schalij, M. J., De Vries,
A. a. F., Pijnappels, D. A. andWu, S. M. 2013. IGF promotes cardiac lineage induction
by selective expansion of cardiogenic mesoderm in vitro. *European Heart Journal,* 34.
Faustino Martins, J. M., Fischer, C., Urzi, A., Vidal, R., Kunz, S., Ruffault, P. L., Kabuss, L.,
Hube, I., Gazzerro, E., Birchmeier, C., *et al.* 2020. Self-Organizing 3D Human Trunk
Neuromuscular Organoids. *Cell Stem Cell,* 26**,** 172-186 e6.
Feng, J., Meyer, C. A., Wang, Q., Liu, J. S., Shirley Liu, X. andZhang, Y. 2012. GFOLD: a
generalized fold change for ranking differentially expressed genes from RNA-seq data.
Bioinformatics, 28**,** 2782-8.
Ferdous, Z., Wei, V. M., Iozzo, R., Hook, M. andGrande-Allen, K. J. 2007. Decorin-
transforming growth factor- interaction regulates matrix organization and mechanical
characteristics of three-dimensional collagen matrices. *J Biol Chem,* 282**,** 35887-98.
Gays, D., Hess, C., Camporeale, A., Ala, U., Provero, P., Mosimann, C. andSantoro, M. M.
2017. An exclusive cellular and molecular network governs intestinal smooth muscle cell
differentiation in vertebrates. *Development,* 144**,** 464-478.
Gribble, F. M. andReimann, F. 2016. Enteroendocrine Cells: Chemosensors in the Intestinal
Epithelium. *Annu Rev Physiol,* 78**,** 277-99.
Hill, M. A. 2020a. *Embryology Cardiovascular System - Heart Development* [Online]. UNSW
Embryology. Available:
https://embryology.med.unsw.edu.au/embryology/index.php/Cardiovascular_System_-
_Heart_Development [Accessed 2020].
Hill, M. A. 2020b. *Embryology Gastrointestinal Tract Development* [Online]. UNSW
Embryology. Available:
https://embryology.med.unsw.edu.au/embryology/index.php/Gastrointestinal_Tract_Dev
elopment#:~:text=The%20gastrointestinal%20tract%20(GIT)%20arises,all%20the%20ge
rm%20cell%20layers. [Accessed 2020].
Hirakow, R. 1983. Development of the cardiac blood vessels in staged human embryos. *Acta*
Anat (Basel), 115**,** 220-30.
Holtzinger, A., Rosenfeld, G. E. andEvans, T. 2010. Gata4 directs development of cardiac-
inducing endoderm from ES cells. *Dev Biol,* 337**,** 63-73.
Hookway, T. A., Butts, J. C., Lee, E., Tang, H. andMcdevitt, T. C. 2016. Aggregate formation
and suspension culture of human pluripotent stem cells and differentiated progeny.
Methods, 101**,** 11-20.
Hookway, T. A., Matthys, O. B., Mendoza-Camacho, F. N., Rains, S., Sepulveda, J. E., Joy, D.
A. andMcdevitt, T. C. 2019. Phenotypic Variation Between Stromal Cells Differentially
Impacts Engineered Cardiac Tissue Function. *Tissue Eng Part A,* 25**,** 773-785.
Hosseini, H. S., Garcia, K. E. andTaber, L. A. 2017. A new hypothesis for foregut and heart tube
formation based on differential growth and actomyosin contraction. *Development,* 144**,**
2381-2391.
Huebsch, N., Loskill, P., Mandegar, M. A., Marks, N. C., Sheehan, A. S., Ma, Z., Mathur, A.,
Nguyen, T. N., Yoo, J. C., Judge, L. M., *et al.* 2015. Automated Video-Based Analysis of

Contractility and Calcium Flux in Human-Induced Pluripotent Stem Cell-Derived
Cardiomyocytes Cultured over Different Spatial Scales. *Tissue Engineering Part C-*
Methods, 21**,** 467-479.
Ishii, Y., Langberg, J. D., Hurtado, R., Lee, S. andMikawa, T. 2007. Induction of proepicardial
marker gene expression by the liver bud. *Development,* 134, 3627-37.
Judge, L. M., Perez-Bermejo, J. A., Truong, A., Ribeiro, A. J., Yoo, J. C., Jensen, C. L.,
Mandegar, M. A., Huebsch, N., Kaake, R. M., So, P. L.*, et al.* 2017. A BAG3 chaperone
complex maintains cardiomyocyte function during proteotoxic stress. *JCI Insight,* 2.
Jung, J., Zheng, M., Goldfarb, M. andZaret, K. S. 1999. Initiation of mammalian liver
development from endoderm by fibroblast growth factors. *Science,* 284**,** 1998-2003.
Kelly, R. G., Brown, N. A. andBuckingham, M. E. 2001. The arterial pole of the mouse heart
forms from Fgf10-expressing cells in pharyngeal mesoderm. *Dev Cell,* 1**,** 435-40.
Kidokoro, H., Yonei-Tamura, S., Tamura, K., Schoenwolf, G. C. andSaijoh, Y. 2018. The heart
tube forms and elongates through dynamic cell rearrangement coordinated with foregut
extension. *Development,* 145**,** dev152488.
Kim, D., Paggi, J. M., Park, C., Bennett, C. andSalzberg, S. L. 2019. Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol,* 37**,** 907-
915.
Kim, H. D., Kim, D. J., Lee, I. J., Rah, B. J., Sawa, Y. andSchaper, J. 1992. Human fetal heart
development after mid-term: morphometry and ultrastructural study. *J Mol Cell Cardiol,*
24**,** 949-65.
Kim, K. H., Nakaoka, Y., Augustin, H. G. andKoh, G. Y. 2018. Myocardial Angiopoietin-1
Controls Atrial Chamber Morphogenesis by Spatiotemporal Degradation of Cardiac Jelly.
Cell Rep, 23**,** 2455-2466.
Kobayashi, M., Khalil, H. A., Lei, N. Y., Wang, Q., Wang, K., Wu, B. M. andDunn, J. C. Y.
2018. Bioengineering functional smooth muscle with spontaneous rhythmic contraction
in vitro. *Sci Rep,* 8**,** 13544.
Koike, H., Iwasawa, K., Ouchi, R., Maezawa, M., Giesbrecht, K., Saiki, N., Ferguson, A.,
Kimura, M., Thompson, W. L., Wells, J. M.*, et al.* 2019. Modelling human hepato-
biliary-pancreatic organogenesis from the foregut-midgut boundary. *Nature,* 574**,** 112-
116.
Lancaster, M. A. andKnoblich, J. A. 2014. Organogenesis in a dish: Modeling development and
disease using organoid technologies. *Science,* 345.
Lee, J., Sutani, A., Kaneko, R., Takeuchi, J., Sasano, T., Kohda, T., Ihara, K., Takahashi, K.,
Yamazoe, M., Morio, T.*, et al.* 2020. In vitro generation of functional murine heart
organoids via FGF4 and extracellular matrix. *Nat Commun,* 11**,** 4283.
Lee, J. H., Protze, S. I., Laksman, Z., Backx, P. H. andKeller, G. M. 2017. Human Pluripotent
Stem Cell-Derived Atrial and Ventricular Cardiomyocytes Develop from Distinct
Mesoderm Populations. *Cell Stem Cell,* 21**,** 179-194 e4.
Lian, X., Hsiao, C., Wilson, G., Zhu, K., Hazeltine, L. B., Azarin, S. M., Raval, K. K., Zhang, J.,
Kamp, T. J. andPalecek, S. P. 2012. Robust cardiomyocyte differentiation from human
pluripotent stem cells via temporal modulation of canonical Wnt signaling. *Proc Natl*
Acad Sci U S A, 109**,** E1848-57.
Lian, X., Zhang, J., Azarin, S. M., Zhu, K., Hazeltine, L. B., Bao, X., Hsiao, C., Kamp, T. J.
andPalecek, S. P. 2013. Directed cardiomyocyte differentiation from human pluripotent

stem cells by modulating Wnt/beta-catenin signaling under fully defined conditions. *Nat Protoc,* 8**,** 162-75.

Liao, Y., Smyth, G. K. andShi, W. 2014. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics,* 30**,** 923-30.

Libby, A. R., Joy, D. A., So, P. L., Mandegar, M. A., Muncie, J. M., Mendoza-Camacho, F. N., Weaver, V. M., Conklin, B. R. andMcdevitt, T. C. 2018. Spatiotemporal mosaic self-patterning of pluripotent stem cells using CRISPR interference. *Elife,* 7.

Libby, A. R. G., Briers, D., Haghighi, I., Joy, D. A., Conklin, B. R., Belta, C. andMcdevitt, T. C. 2019. Automated Design of Pluripotent Stem Cell Self-Organization. *Cell Syst,* 9**,** 483-495 e10.

Lough, J. andSugi, Y. 2000. Endoderm and heart development. *Dev Dyn,* 217**,** 327-42.

Lundy, S. D., Zhu, W. Z., Regnier, M. andLaflamme, M. A. 2013. Structural and functional maturation of cardiomyocytes derived from human pluripotent stem cells. *Stem Cells Dev,* 22**,** 1991-2002.

Ma, J., Guo, L., Fiene, S. J., Anson, B. D., Thomson, J. A., Kamp, T. J., Kolaja, K. L., Swanson, B. J. andJanuary, C. T. 2011. High purity human-induced pluripotent stem cell-derived cardiomyocytes: electrophysiological properties of action potentials and ionic currents. *Am J Physiol Heart Circ Physiol,* 301**,** H2006-17.

Ma, Z., Wang, J., Loskill, P., Huebsch, N., Koo, S., Svedlund, F. L., Marks, N. C., Hua, E. W., Grigoropoulos, C. P., Conklin, B. R.*, et al.* 2015. Self-organizing human cardiac microchambers mediated by geometric confinement. *Nat Commun,* 6**,** 7413.

Mackay, J. L., Sood, A. andKumar, S. 2014. Three-dimensional patterning of multiple cell populations through orthogonal genetic control of cell motility. *Soft matter,* 10**,** 2372-2380.

Macqueen, L. A., Sheehy, S. P., Chantre, C. O., Zimmerman, J. F., Pasqualini, F. S., Liu, X., Goss, J. A., Campbell, P. H., Gonzalez, G. M., Park, S. J.*, et al.* 2018. A tissue-engineered scale model of the heart ventricle. *Nat Biomed Eng,* 2**,** 930-941.

Mandegar, M. A., Huebsch, N., Frolov, E. B., Shin, E., Truong, A., Olvera, M. P., Chan, A. H., Miyaoka, Y., Holmes, K., Spencer, C. I.*, et al.* 2016. CRISPR Interference Efficiently Induces Specific and Reversible Gene Silencing in Human iPSCs. *Cell Stem Cell,* 18**,** 541-53.

Matthys, O. B., Silva, A. C. andMcdevitt, T. C. 2020. Engineering human organoid development ex vivo—challenges and opportunities. *Current Opinion in Biomedical Engineering,* 13**,** 160-167.

Mills, R. J., Titmarsh, D. M., Koenig, X., Parker, B. L., Ryall, J. G., Quaife-Ryan, G. A., Voges, H. K., Hodson, M. P., Ferguson, C., Drowley, L.*, et al.* 2017. Functional screening in human cardiac organoids reveals a metabolic mechanism for cardiomyocyte cell cycle arrest. *Proc Natl Acad Sci U S A,* 114**,** E8372-E8381.

Mummery, C., Ward-Van Oostwaard, D., Doevendans, P., Spijker, R., Van Den Brink, S., Hassink, R., Van Der Heyden, M., Opthof, T., Pera, M., De La Riviere, A. B.*, et al.* 2003. Differentiation of human embryonic stem cells to cardiomyocytes: role of coculture with visceral endoderm-like cells. *Circulation,* 107**,** 2733-40.

Munera, J. O., Sundaram, N., Rankin, S. A., Hill, D., Watson, C., Mahe, M., Vallance, J. E., Shroyer, N. F., Sinagoga, K. L., Zarzoso-Lacoste, A.*, et al.* 2017. Differentiation of Human Pluripotent Stem Cells into Colonic Organoids via Transient Activation of BMP Signaling. *Cell Stem Cell,* 21**,** 51-+.

Nguyen, D. C., Hookway, T. A., Wu, Q., Jha, R., Preininger, M. K., Chen, X., Easley, C. A.,
Spearman, P., Deshpande, S. R., Maher, K.*, et al.* 2014. Microscale generation of
cardiospheres promotes robust enrichment of cardiomyocytes derived from human
pluripotent stem cells. *Stem Cell Reports,* 3**,** 260-8.
Perez-Pomares, J. M. andDe La Pompa, J. L. 2011. Signaling during epicardium and coronary
vessel development. *Circ Res,* 109**,** 1429-42.
Racca, A. W., Klaiman, J. M., Pioner, J. M., Cheng, Y., Beck, A. E., Moussavi-Harami, F.,
Bamshad, M. J. andRegnier, M. 2016. Contractile properties of developing human fetal
cardiac muscle. *J Physiol,* 594**,** 437-52.
Rochais, F., Sturny, R., Chao, C. M., Mesbah, K., Bennett, M., Mohun, T. J., Bellusci, S.
andKelly, R. G. 2014. FGF10 promotes regional foetal cardiomyocyte proliferation and
adult cardiomyocyte cell-cycle re-entry. *Cardiovasc Res,* 104**,** 432-42.
Ronaldson-Bouchard, K., Ma, S. P., Yeager, K., Chen, T., Song, L., Sirabella, D., Morikawa, K.,
Teles, D., Yazawa, M. andVunjak-Novakovic, G. 2018. Advanced maturation of human
cardiac tissue grown from pluripotent stem cells. *Nature*.
Rossi, G., Broguiere, N., Miyamoto, M., Boni, A., Guiet, R., Girgin, M., Kelly, R. G., Kwon, C.
andLutolf, M. P. 2020. Capturing Cardiogenesis in Gastruloids. *Cell Stem Cell*.
Rudy-Reil, D. andLough, J. 2004. Avian precardiac endoderm/mesoderm induces cardiac
myocyte differentiation in murine embryonic stem cells. *Circ Res,* 94**,** e107-16.
Rutenberg, J. B., Fischer, A., Jia, H., Gessler, M., Zhong, T. P. andMercola, M. 2006.
Developmental patterning of the cardiac atrioventricular canal by Notch and Hairy-
related transcription factors. *Development,* 133**,** 4381-90.
Sato, T., Vries, R. G., Snippert, H. J., Van De Wetering, M., Barker, N., Stange, D. E., Van Es, J.
H., Abo, A., Kujala, P., Peters, P. J.*, et al.* 2009. Single Lgr5 stem cells build crypt-villus
structures in vitro without a mesenchymal niche. *Nature,* 459**,** 262-265.
Schultheiss, T. M., Xydas, S. andLassar, A. B. 1995. Induction of avian cardiac myogenesis by
anterior endoderm. *Development,* 121**,** 4203-14.
Sergeeva, I. A. andChristoffels, V. M. 2013. Regulation of expression of atrial and brain
natriuretic peptide, biomarkers for heart development and disease. *Biochim Biophys Acta,*
1832**,** 2403-13.
Shen, H., Cavallero, S., Estrada, K. D., Sandovici, I., Kumar, S. R., Makita, T., Lien, C. L.,
Constancia, M. andSucov, H. M. 2015. Extracardiac control of embryonic cardiomyocyte
proliferation and ventricular wall expansion. *Cardiovasc Res,* 105**,** 271-8.
Smits, A. M., Dronkers, E. andGoumans, M. J. 2018. The epicardium as a source of multipotent
adult cardiac progenitor cells: Their origin, role and fate. *Pharmacol Res,* 127**,** 129-140.
Stennard, F. A. andHarvey, R. P. 2005. T-box transcription factors and their roles in regulatory
hierarchies in the developing heart. *Development,* 132**,** 4897-4910.
Sugi, Y. andLough, J. 1994. Anterior Endoderm Is a Specific Effector of Terminal Cardiac
Myocyte Differentiation of Cells from the Embryonic Heart Forming Region.
Developmental Dynamics, 200**,** 155-162.
Sugi, Y. andLough, J. 1995. Activin-A and FGF-2 mimic the inductive effects of anterior
endoderm on terminal cardiac myogenesis in vitro. *Dev Biol,* 168**,** 567-74.
Takasato, M., Er, P. X., Chiu, H. S., Maier, B., Baillie, G. J., Ferguson, C., Parton, R. G.,
Wolvetang, E. J., Roost, M. S., Lopes, S. M.*, et al.* 2016. Kidney organoids from human
iPS cells contain multiple lineages and model human nephrogenesis. *Nature,* 536**,** 238.

Turaga, D., Matthys, O. B., Hookway, T. A., Joy, D. A., Calvert, M. andMcdevitt, T. C. 2020.
Single-Cell Determination of Cardiac Microtissue Structure and Function Using Light
Sheet Microscopy. *Tissue Eng Part C Methods,* 26**,** 207-215.
Uosaki, H., Andersen, P., Shenje, L. T., Fernandez, L., Christiansen, S. L. andKwon, C. 2012.
Direct contact with endoderm-like cells efficiently induces cardiac progenitors from
mouse and human pluripotent stem cells. *PLoS One,* 7**,** e46413.
Varner, V. D. andTaber, L. A. 2012. Not just inductive: a crucial mechanical role for the
endoderm during heart tube assembly. *Development,* 139**,** 1680-1690.
Volpato, V. andWebber, C. 2020. Addressing variability in iPSC-derived models of human
disease: guidelines to promote reproducibility. *Dis Model Mech,* 13.
Wheelwright, M., Win, Z., Mikkila, J. L., Amen, K. Y., Alford, P. W. andMetzger, J. M. 2018.
Investigation of human iPSC-derived cardiac myocyte functional maturation by single
cell traction force microscopy. *PLoS One,* 13**,** e0194909.
Xiang, Y., Tanaka, Y., Patterson, B., Kang, Y. J., Govindaiah, G., Roselaar, N., Cakir, B., Kim,
K. Y., Lombroso, A. P., Hwang, S. M.*, et al.* 2017. Fusion of Regionally Specified hPSC-
Derived Organoids Models Human Brain Development and Interneuron Migration. *Cell*
Stem Cell, 21**,** 383-398 e7.
Yang, H., Borg, T. K., Wang, Z., Ma, Z. andGao, B. Z. 2014. Role of the Basement Membrane
in Regulation of Cardiac Electrical Properties. *Annals of biomedical engineering,* 42**,**
1148-1157.
Zhang, H. T. andHiiragi, T. 2018. Symmetry Breaking in the Mammalian Embryo. *Annu Rev*
Cell Dev Biol, 34**,** 405-426.
Zhang, Z., Yu, D., Seo, M., Hersh, C. P., Weiss, S. T. andQiu, W. 2019. Novel Data
Transformations for RNA-seq Differential Expression Analysis. *Sci Rep,* 9**,** 4820.
1248