Bacillus thuringiensis bioinsecticides induce developmental defects

in non-target *Drosophila melanogaster* larvae

Marie-Paule Nawrot-Esposito[a], Aurélie Babin[a], Matthieu Pasco[a], Marylène Poirié[a], Jean-Luc

Gatti[a] and Armel Gallet[a#]

[a] Université Côte d'Azur, CNRS, INRAE, ISA, France

Running Head: *Bt* impacts on *Drosophila* midgut

Corresponding author: A. Gallet, ISA, UMR CNRS 7254/INRAE 1355/UCA, 400 route des

Chappes, BP 167, 06903 Sophia Antipolis Cedex, France. gallet@unice.fr

Summary statement

Bioinsecticides based on *Bacillus thuringiensis kurstaki* spores are applied to specifically kill pest lepidopteran larvae. Unexpectedly they display unintended effects on the intestinal homeostasis of *Drosophila* midgut, resulting in developmental delay and growth defects.

Abstract

Bioinsecticides made from the bacterium *Bacillus thuringiensis* (*Bt*) are the best-selling bioinsecticide worldwide. Among *Bt* bioinsecticides, those based on the strain *Bt var. kurstaki* (*Btk*) are widely used in farming to specifically control pest lepidopteran larvae. Although there is much evidence of the lack of acute lethality of *Btk* products for non-target animals, only scarce data are available on their potential non-lethal developmental adverse effects. Using doses that could be reached in the field upon sprayings, we have shown that *Btk* products impair growth and developmental time of the non-target dipteran *Drosophila melanogaster*. These effects are mediated by the synergy between *Btk* bacteria and *Btk* insecticidal toxins, which induces a significant apoptosis of larval enterocytes, resulting in a decreased intestinal capacity to digest proteins. The harmful effects can be mitigated by a protein-rich diet or by adding the probiotic bacterium *Lactobacillus plantarum* into the food. Finally, we showed that the larval midgut maintain its integrity upon *Btk* aggression thanks to both the flattening of surviving enterocytes and the generation of new immature cells arising from the adult midgut precursor cells.

Keywords

Bioinsecticides, *Bacillus thuringiensis*, *Drosophila melanogaster*, Developmental defects, Gut homeostasis, *Lactobacillus plantarum.*

INTRODUCTION

The use of conventional synthetic chemical pesticides is controversial mainly because of their harmful effects on human health and ecosystems. Public opinion along with governmental policies encourage to reduce their use (Casida and Bryant, 2017; Frisvold, 2018). However, due to the world's population expansion, the use of pesticides will still be necessary to produce enough food to feed the 9.5 billion humans expected in 2050 (https://population.un.org/wpp/). An emerging solution to reduce the use of conventional pesticides is their replacement with biopesticides.

Among bioinsecticides, *Bacillus thuringiensis* (*Bt*) products are increasingly sprayed in both organic farming and conventional agriculture (Fiuza et al., 2017). *Bt* products are the second most used insecticides in the world with 32 000 tons sold in 2017 (Casida and Bryant, 2017). *Bt* is a Gram-positive spore-forming bacterium belonging to the *Bacillus cereus* group (Vilas-Boas et al., 2007). It was first identified and characterized for its specific entomopathogenic properties due to the presence of specific Cry toxins, which are produced in a crystalline form during the sporulation of bacteria (Rabinovitch et al., 2017). Seventy-eight different strains of *Bt* are currently inventoried (http://www.bgsc.org/), producing a total of more than 300 distinct Cry toxins (http://www.lifesci.sussex.ac.uk/home/Neil_Crickmore/Bt/) having a spectrum of toxicity ranging from nematodes to human tumor cells (Adang et al., 2014; Frankenhuyzen, 2017). However, only four strains are used commercially as bioinsecticides owing to the specific acute toxicity of their Cry toxins to pest larvae: *Bt var. kurstaki (Btk)* and *Bt var. aizawai* to kill lepidopteran larvae, *Bt var. israelensis* to kill mosquito larvae and *Bt var. morrisoni* to kill coleopteran larvae (http://sitem.herts.ac.uk/aeru/bpdb/index.htm). Upon ingestion of commercial formulations made of spores, crystals and additives, Cry protoxins are first released from the crystal before being processed by gut proteases to give rise to the active Cry toxins. Then, these Cry toxins bind to receptors at the surface of the intestinal cells, inducing

their death and ultimately the destruction of the gut epithelium. The spores germinate and, thanks to the holes made by Cry toxins in the gut lining, vegetative bacteria invade the internal cavity of the body, causing sepsis which kills the targeted pest in 2 or 3 days. The specificity of the acute Cry toxicity relies on both the capacity of gut proteases to cleave and activate Cry protoxins and on the presence of host receptors specifically recognized by Cry toxins (Adang et al., 2014). Each variant of *Bt* produces between 1 and 6 different Cry toxins. In general, all the Cry toxins contained in a *Bt* variant target a specific phylogenetic order (*e.g.* lepidopteran, coleopteran…) (Palma et al., 2014). The most used *Bt* variant is *Btk,* which produces 5 different Cry toxins (Cry1Aa, Cry1Ab, Cry1Ac, Cry2Aa and Cry2Ab) (Caballero et al., 2020). *Btk* is widely used in forestry and farming to fight lepidopteran larvae.

The increasing environmental dispersion of *Btk* bioinsecticides raises the question of their long-term putative effects on both health and environment. Although there are many data showing that the Cry toxins produced by *Btk* do not display harmful effects against non-target organisms (Haller et al., 2017; Rubio-Infante and Moreno-Fierros, 2016), there are also studies suggesting that Cry toxins and/or *Bt* products may have unintended impacts on non-target animals. For example, the presence of Cry1Ab toxins produced by GM maize in water streams surrounding maize fields delays the growth and increases the mortality of trichopteran species whose larvae have an aquatic lifestyle (Rosi-Marshall et al., 2007; Tank et al., 2010). More recently, Amichot and colleagues have shown that *Btk* bioinsecticides are acutely toxic against *Trichogramma chilonis* females, a tiny egg endoparasitoid hymenopteran used as a biocontrol agent in integrated pest management (Amichot et al., 2016). On vertebrates, Grisolia and colleagues observed that the exposure of *Danio rerio* fry to different *Btk* Cry toxins induced lethality and developmental delay (Grisolia et al., 2009).

Based on these data, we decided to use the model organism *Drosophila melanogaster*, not targeted by *Btk*, to identify and characterize the putative impacts of *Btk* products on the development. While the acute toxicity of the agricultural doses of *Btk* bioinsecticides

efficiently kills lepidopteran larva, *Drosophila* larvae, being non-susceptible to these doses,

allow the study of non-lethal impacts. In addition, *D. melanogaster* has been used

successfully in many studies for deciphering the toxicology of insecticides and is a toolbox

widely used to identify disturbed cellular and molecular mechanisms (Scott and Buchon,

2019).

Here, using commercial *Btk* products at two realistic doses, a dose detected on

vegetables after one spraying and a dose only tenfold higher (but equivalent to the ten

cumulative treatments authorized by the European Union), we observed that *Btk*

bioinsecticides induce developmental delay and reduce growth of *D. melanogaster* larvae. We

further showed that *Btk* bioinsecticides trigger intestinal cell death and alter protein digestion.

Interestingly, we showed that a protein-rich diet or supplementing food with the probiotic

Lactobacillus plantarum (*L. plantarum*), known to enhance protein uptake (Erkosar et al.,

2015), mitigates the impacts of *Btk* bioinsecticide on growth and development. We also found

that *Btk* products do not interfere with the commensal flora of the larvae. Finally, we

unraveled new cellular mechanisms for maintaining intestinal integrity mounted by the larval

midgut epithelium in response to *Btk* aggression*,* particularly the flattening of enterocytes and

the production of new cells from the adult midgut precursors. Overall, our data show that

though not lethal for larvae of the non-target organism *Drosophila melanogaster*, agricultural

doses of *Btk* bioinsecticides damage gut epithelium and consequently impair larval growth

and developmental time.

RESULTS

Btk **bioinsecticide induces developmental delay and growth defect**

European and French Food Safety Agencies (Efsa and Anses, respectively) have estimated

that the amount of *Btk* spores on fruits and vegetables after treatment can reach $5x10^6$ Colony

Forming Units (CFU)/g (ANSES Saisine n° 2013-SA-0039, 2013; EFSA BIOHAZ Panel,

2016). This quantification may be underestimated since up to 10 successive treatments can be

applied by farmers (ANSES Saisine n° 2013-SA-0039, 2013) (see also the European

directives SANCO/1541/08 – rev. 4 and SANCO/1543/08 – rev. 4). Therefore, we decided to

assess the impacts of the ingestion of two *Btk* bioinsecticides (i.e. Delfin and Dipel) on the

growth and development of *Drosophila* larvae at two field-realistic doses: $5x10^6$ CFU/g

(corresponding to the dose observed after one spraying) and $5x10^7$ CFU/g (a dose which

could be reached upon repeated treatments), named hereafter "1X" and "10X", respectively.

We first compared the impacts of Delfin with water (the diluent for commercial products) by

contaminating 2g of standard fly medium before depositing 20 eggs aged of 0-4h. We then

measured the timing of larval development by performing a pupation curve. In the control

experiment (H_2O), 90% of the larvae pupated at 169h after eggs laying (this threshold will

serve as a point of comparison throughout our study, Table 1 summarizes all the data at 10%,

50% and 90% of pupation entry). The 1X and 10X doses of Delfin postponed pupation by

10.5h (+6.2%) and 16h (+9.5%) respectively (Fig. 1A and Table 1). We then assessed the

impact of the second bioinsecticide Dipel. Because developmental delay was marked with

Delfin at the 10X dose, we used this same dose as a benchmark for Dipel. Strikingly, the

pupation delay was increased to reach 25h (+14.8%). Commercial products are made of

spores, toxin crystals and additives (see Materials And Methods). Dipel contains 46% of

"trade secret" components, while Delfin contains 15% of naphthalene sulfonic acid. To assess

the potential involvement of the additives present in Dipel in the augmented delay observed,

we plated the commercial product on LB agar, isolated a colony and prepared our own

mixture of spores and toxins (the *Btk* strain used in Dipel is named *ABTS-351*). When larvae

were fed with the 10X dose of the homemade preparation, we obtained developmental delays

similar to those with the Delfin treatments (Fig. 1A and Table 1), suggesting that the

unknown additives present in the commercial preparation of Dipel might contribute to the

increased delay we observed. Moreover, these data also suggested that the 15% of

naphthalene sulfonic acid presents in Delfin had no impact on the developmental timing.

Indeed feeding developing larvae with an amount of naphthalene sulfonic acid equivalent to

the 10X dose of Delfin had no effect on the developmental timing (Fig. 1B and Table 1).

Then, we wondered whether the developmental delay was due to the spores alone, the crystals

of toxins alone, or both. First, we fed the developing larvae with the 1X and 10X doses of

spores of a *Btk* strain devoid of Cry toxins (*Btk$^{\Delta Cry}$*). Whatever the dose, no developmental

delay was observed (Fig. 1B and Table 1). Then, we fed the developing larvae with an

equivalent 10X dose of toxin crystals, considering that crystals represent 30% of the dry

weight of bioinsecticides (Agaisse and Lereclus, 1995; Monro, 1961; Murty et al., 1994) (Fig.

S2). Again, we did not observe any developmental delay (Fig. 1B and Table 1). Therefore, we

assumed that the developmental delay we observed was mainly due to the mixture of spores

and crystals (Fig. 1A).

Since growth defects have a well-documented impact on the developmental timing

(Hietakangas and Cohen, 2009), we next measured the pupal volume, a known readout to

estimate growth defects (Delanoue et al., 2010). Interestingly, we observed a marked growth

defect when the larvae were fed with Delfin and Dipel, as well as a weaker one when the

larvae were fed with *ABTS- 351* (Fig. 1C). We did not detect any growth defect in all the

other conditions. Together, our data show that ingestion of food contaminated with a mixture

of *Btk* spores and crystals induces developmental delay and growth defect on the model

organism *D. melanogaster*, at doses that can be found in treated areas.

Since i) the composition of Delfin is clearly provided by the manufacturer compared

to Dipel (see Materials And Methods), ii) Delfin is composed of 85% of spores and toxin

crystals while only 54% for Dipel, and iii) the naphthalene sulfonic acid has no effect on the

developmental timing and growth, we decided to use only Delfin in the rest of the study.

***Btk* bioinsecticide does not alter feeding behavior and food intake**

The *Btk* load in the midgut could explain the differences observed between $Btk^{\Delta Cry}$ and Delfin.

When we estimated the amount of *Btk* cells in the midgut of the L3 larvae (Fig. 2A), no

difference was observed for 1X or 10X doses of Delfin compared to the equivalent doses of

$Btk^{\Delta Cry}$. Therefore, the variation in the quantity of *Btk* bacterial cells in the midgut could not

explain the disturbed developmental timing and growth observed with Delfin.

Adult flies and larvae can sense the presence of harmful microbes, displaying an

aversion to the contaminated food (Babin et al., 2014; Keita et al., 2017; Stensmyr et al.,

2012; Surendran et al., 2017). Therefore, we tested whether the feeding behavior was affected

by the presence of Delfin. We added a blue dye in the food and counted the number of feeding

larvae containing blue food in their gut, either in L3 (Fig. 2B) or in L2 stage (Fig. 2C).

Although we observed a slight decrease in L2 larvae reared under Delfin 1X dose condition,

we did not detect any change in feeding behavior. We then quantified the food intake and,

once again, we did not detect any significant modification in the L3 larvae (Fig. 2D), unlike in

the L2 larvae, which were affected whatever the condition (excepted for the $Btk^{\Delta Cry}$ 10X

condition which was not significant) compared to the control (Fig. 2E). Therefore, although

food contaminated with *Btk* spores (with or without crystals of toxins) disturbed food intake

of L2 larvae but not of L3 larvae, these data do not explain the specific impact of Delfin on

the developmental time and growth of the larvae.

Btk bioinsecticide impairs intestinal protein metabolism

Proteins are a key nutrient to drive normal larval development and growth. Increasing protein

content in the diet accelerates the developmental time and/or stimulates growth, while an

increase in sugar content delays development and a change in lipid amounts has only a

marginal impact (Anagnostou et al., 2010; Colombani et al., 2003; Layalle et al., 2008; Reis,

2016; Rodrigues et al., 2015; Soultoukis and Partridge, 2016). Because we have previously

shown that the presence of opportunistic bacteria in the adult midgut impede the digestion of

proteins (Loudhaief et al., 2017), we wondered whether this could also be the case in larvae in the presence of *Btk* bioinsecticide. While the 10X dose of $Btk^{\Delta Cry}$ had no consequence, we observed a decreased capacity of the midgut of L3 larvae to digest proteins in a Delfin dose-dependent manner (Fig. 3A).

We next wondered whether increasing the protein content of the diet could rescue the developmental delay and the growth defects caused by the ingestion of Delfin. Our rearing medium contains 2% yeast as a protein source (see Materials and Methods). We therefore performed a pupation curve with larvae raised on a medium containing 10% yeast with the 1X or 10X doses of Delfin (Fig. 3B). Compared to the control, no difference in the developmental time was observed for the 1X dose of Delfin and only a slight delay for the 10X dose (Table 1). Noteworthy, the developmental time for all conditions on the protein-rich medium was shortened compared to the control on our conventional medium (light green curve in Fig. 3B and Table 1). Similarly, the pupal volume (Fig. 3C) was completely restored for the 1X dose of Delfin and partially rescued for the 10X dose of Delfin (5.9% volume loss instead of 11.9% in the poor-protein diet). Taken together, our data suggest that Delfin impairs protein digestion, and thus gut functions, which can be rescued by adding extra dietary protein.

The commensal bacterium *L. plantarum* helps to overcome *Btk* bioinsecticide impacts

It has been recently shown that the commensal bacterium *L. plantarum* promotes the juvenile systemic growth of *Drosophila* larvae and infant mice by stimulating digestion and absorption of proteins, and protects the child against sepsis upon undernutrition conditions (Erkosar et al., 2015; Panigrahi et al., 2017; Schwarzer et al., 2016; Storelli et al., 2011; Tefit and Leulier, 2017). Therefore, we wondered whether complementing our conventional medium with *L. plantarum* would compensate for the impacts of the 10X dose of Delfin. First, we observed that adding *L. plantarum* in the conventional medium shortened the developmental time of control larvae (Fig. 4A, purple curve compared to green curve and Table 1). Interestingly, the

presence of *L. plantarum* fully counteracted Delfin-dependent developmental delay (Fig. 4,

pink curve compared to red curve and Table 1). We next observed that the presence of *L*

plantarum in the conventional medium reduced the pupal size of the control larvae by 4.4%

compared to the control without *L. plantarum* (Fig. 4B, purple bar compared to green bar) but

fully rescued the growth defects induced by the 10X dose of Delfin (Fig. 4B pink bar). Since

ingestion of *L. plantarum* and its establishment in the midgut (Loudhaief et al., 2017) might

compete for the presence of *Btk* and not act on protein intake as expected, we also controlled

the number of *Btk* CFUs in the absence or presence of *L. plantarum* in the midgut of L3

larvae and we found no difference (Fig. 4C). Taken together, our data demonstrated that

stimulation digestion and/or absorption of proteins by the commensal bacterium *L. plantarum*

helps to overcome the adverse effects of the *Btk* bioinsecticides on larval development.

The impacts of *Btk* bioinsecticide do not rely on commensal bacteria disturbance

L. plantarum is a widespread commensal bacterium of the *Drosophila* midgut (Broderick and

Lemaitre, 2012; Chandler et al., 2011; Fast et al., 2018). Because complementing the larval

diet with *L. plantarum* rescued the developmental delay and growth defects induced by *Btk*,

we wondered whether the indigenous commensal flora of larvae reared on our conventional

medium (i.e. poor in protein) was inhabited by *L. plantarum*. Only few *L. plantarum* and/or *L.*

brevis (belonging to the family of *Lactobacillaceae*) were present in the commensal flora of

our larvae (Fig. 4D). The dominant bacteria family we identified belonged to

Acetobacteraceae known to be common in flies feeding on sugary, acidic and alcoholic food

(Chandler et al., 2011; Téfit et al., 2018). This observation was in agreement with the

composition of our conventional rearing medium poor in protein but quite rich in simple sugar

(i.e. sucrose, see Materials And Methods) and it suggested that the *Btk* effects we observed

were independent of a putative deleterious impact on the indigenous *L. plantarum* of our

larvae.

However, we could not rule out an impact of the *Btk* bioinsecticide on other members

of the commensal flora. To test this hypothesis, we generated an axenic fly strain (see

Materials And Methods). Axenic larvae (Fig. 4D) were further fed on our conventional

medium treated or not with the 10X dose of Delfin. The developmental time of the control

axenic larvae was slightly shorter than those of the conventionally reared (CR) larvae (i.e.

non-axenic) (Fig. 4E and Table 1). Nevertheless, axenic larvae infected with Delfin 10X

displayed a 19h developmental delay compared to the control axenic larvae (+11.7%) (Fig. 4E

and Table 1), which was similar to the delay observed with *Btk*-treated CR larvae (see Fig. 1).

Then, while axenic larvae gave rise to pupae with a similar volume to that of CR larvae (Fig.

4F), Delfin 10X still induced a reduction of pupal volume in axenic larvae compared to the

axenic control (Fig. 4F) in a magnitude similar to what we observed for Delfin 10X with CR

larvae (see Fig. 1). Taken together our data strongly suggest that Delfin ingestion affects

developmental time and growth independently of the commensal bacteria.

Btk bioinsecticide induces midgut perturbations

Because we had previously shown that the ingestion of *Bt*k vegetative cells by adult

Drosophila reduced the accumulation of lipids in enterocytes (Loudhaief et al., 2017), we

wondered whether the commercial *Btk* product did the same in larvae. In unchallenged L3

larvae, lipid droplets strongly accumulate in enterocytes in the most anterior part of the

midgut and weakly in the posterior midgut (Fig. 5A) (Palm et al., 2012). Instead of lipid

droplets depletion, feeding the larvae with Delfin 10X provoked an accumulation of lipid

droplets in the posterior larval midgut (Fig. 5E). Furthermore, lipid droplets accumulated

apically, facing the lumen (Fig. 5F and F' compared to 5B and B'). Such an accumulation was

not observed when developing larvae were fed with $Btk^{\Delta Cry}$ spores (Fig. 5C, D and D').

Interestingly, it has been recently demonstrated that the pore-forming toxins hemolysin and

monalysin produced respectively by the enteropathogenic bacteria *Serratia marcescens* (*Sm*)

and *Pseudomonas entomophila* (*Pe*) can induce an apical accumulation of lipid droplets in adult enterocytes of *Drosophila*, honeybees and mice as well as in epithelial Caco-2 cells in culture. This phenomenon precedes the expulsion of cytoplasm required to get rid of toxins from the enterocytes (Lee et al., 2016). Since the Cry toxins produced by *Btk* are also pore-forming toxins (Adang et al., 2014), we fed developing larvae with purified toxin crystals and checked for the accumulation of lipid droplets. Unexpectedly, we did not observe any accumulation of lipid droplets (Fig. 5G, H and H'). However, feeding our developing larvae with the naphthalene sulfonic acid (the additive present in the commercial products, see Supplemental Materials) provoked an accumulation of apical lipid droplets posterior to the acidic domain (Fig. 5I, J and J'). To further assess whether such a lipid accumulation in the midgut would impair lipid metabolism in the whole larva, we quantified the circulating lipids (the diacylglycerides - DAG) in the hemolymph and the stored lipids (triacylglycerides - TAG) (Tennessen et al., 2014) in mid-third instar larvae. We did not notice any significant perturbation in lipid metabolism (Fig. 5K) though a downward trend was visible for circulating DAG in larvae fed with Delfin 10X. Thereby, it is possible that such an apical accumulation of lipid droplets serves for the trap and elimination of the naphthalene sulfonic acid similarly to what happens for the elimination of the *Sm* and *Pe* pore-forming toxins (Lee et al., 2016).

In adult *Drosophila*, pathogenic and opportunistic bacteria can induce enterocyte apoptosis upon ingestion (Apidianakis et al., 2009; Loudhaief et al., 2017). Therefore, we verified the appearance of dying cells in the midgut using the Caspase3::GFP sensor to visualize apoptosis in living tissue (Schott et al., 2017) (see Materials And Methods). While unchallenged and *Btk*$^{\Delta Cry}$ fed larvae did not display any obvious apoptosis, the developing larvae fed with Delfin harbored apoptotic enterocytes mainly in the posterior part of the intestine (Fig. 5L-M and Fig. S3). Because in target lepidopteran larvae, Cry toxins make holes in the gut lining, breaking down the gut barrier and allowing *Btk* cells to penetrate

inside the internal milieu (Adang et al., 2014), we checked whether the sealing of the gut

epithelium was maintained even in presence of widespread apoptosis. We fed our developing

larvae with the small fluorescent molecule Dextran-FITC that is unable to cross the midgut

epithelium (Fig. 5O) unless the intestinal barrier is broken, allowing the fluorescence to

spread throughout the larva like in the case of *Pe* infection (Fig. 5P). Interestingly, *Btk^{ΔCry}*

spores as well as Delfin did not induce the loss of gut barrier function (Fig. 5Q and R). Thus,

our data show that *Btk* bioinsecticide hampers intestinal lipid homeostasis and induces

enterocyte apoptosis without causing the rupture of the intestinal barrier. Such physiological

perturbations may be responsible for the developmental delay and the growth defect observed.

Enterocyte flattening and incomplete differentiation of Adult Midgut Precursors help to

maintain intestinal integrity

Intestinal physiology can also be impeded by a disturbance of the apico-basal polarity of the

gut epithelium. In the control L3 larval midgut, Armadillo (Arm, β-Catenin), a component of

the adherens junction, marks the basolateral compartment of the enterocytes and Disc Large

(Dlg), a component of the septate junction, marks the sub-apical compartment. Phalloidin that

binds to F-Actin reveals both the apical brush border (facing the lumen) and the basal visceral

muscles that surround the midgut lining (Fig. 6A and A'). This organization is similar to the

adult midgut (Chen et al., 2018; Loudhaief et al., 2017). Moreover, Arm also delineates the

islets of the a̲dult m̲idgut p̲recursors (AMPs), which will give birth to the adult midgut during

metamorphosis (Fig. 6A) (Jiang and Edgar, 2009; Mathur et al., 2010; Micchelli et al., 2011).

When larvae were fed with the 10X dose of *Btk^{ΔCry}* spores, we did not observe any major

perturbations of the architecture of late L3 midguts (Fig. 6B and B'). However, we observed

an increased number of AMPs per islet (Fig. 6D) correlated with an increase in AMP division

(Fig. 6E). When larvae were fed with Delfin 10X, we also observed an increased number of

AMPs by islet (Fig. 6D) and a huge increase in AMP proliferation (Fig. 6E) though at the

expense of the number of islets (Fig. 6F). Moreover, we also observed an increase in the

enterocyte surface (Fig. 6C and G). Interestingly, such an increase in enterocyte surface did

not rely on a higher ploidy (Fig. 6J) but rather on a change in cell shape, i.e. enterocyte

flattening (Fig. 6C' and H). Finally, we noticed the appearance of uncharacterized cells having

an "intermediate" size. These cells which were intercalated between large enterocytes and the

AMP islets, were neither mature polyploid enterocytes nor diploid AMPs or enteroendocrine

cells (Fig. 6I and I'). This observation was confirmed by analyzing the ploidy of the nuclei. In

the control larvae, 89% of the polyploid cells underwent 4 endocycles of replication (32C

nuclei, C corresponding to chromatin copy number) and 10% reached 5 cycles of

endoreplication (64C) (Fig. 6J). When larvae were raised on medium contaminated with

$Btk^{\Delta Cry}$ spores, 74% corresponded to 32C nuclei and 25% to 64C nuclei (Fig. 6J). When larvae

were fed with Delfin 10X, a new population of cells with 16C, 8C and 4C nuclei appeared,

representing 37% of the population of polyploid cells (Fig. 6J). While the adult midgut

quickly regenerates upon an oral infection thanks to the proliferation of adult intestinal stem

cells and the differentiation of their daughter cells (the enteroblasts) into new enterocytes

which repair the damages (Bonfini et al., 2016), such a mechanism has never been observed

in *Drosophila* larvae. Indeed, the number of enterocytes composing the L3 intestine is defined

during embryogenesis (Miguel-Aliaga et al., 2018) and the larval gut grows without cell

division but by increasing the ploidy and consequently the size of enterocytes. However, our

data suggested that upon Delfin ingestion, larval gut homeostasis is perturbed and the

apparition of undefined cells is promoted. In the larval intestine, AMPs are the only cells

capable of division (Jiang and Edgar, 2009; Mathur et al., 2010; Micchelli et al., 2011).

Therefore, we hypothesized that the unknown cells having a nucleus ranging from 4C to 16C

(Fig. 6J) and adjacent to the AMP islets (Fig. 6I) could arise from the differentiation of those

AMPs. To test our hypothesis, we used the lineage tracing genetic tools developed in

Drosophila named ReDDM (Antonello et al., 2015). The principle of this lineage tracing is

based on the expression of both a labile GFP and a stable RFP within the AMPs thanks to the

use of a promotor specifically expressed in AMPs and inducible at will (see Materials And

Methods). Accordingly, AMPs express both GFP and RFP, while the progenies, which are not

AMPs, only express RFP. In control larvae, GFP and RFP were only present in the islet cells

(Fig. 6K) (Jiang and Edgar, 2009; Mathur et al., 2010; Micchelli et al., 2011). When larvae

were fed with $Btk^{\Delta Cry}$ spores, GFP and RFP were also only expressed in the same AMP cells

(Fig. 6L). In contrast, when larvae were fed with Delfin 10X, cells harboring only RFP were

now detectable. Furthermore, the newborn cells had nuclei of intermediate size ranging from

4C to 16C (Fig. 6M). This result demonstrated that upon aggression by *Btk* bioinsecticides,

the larval midgut is able to produce new cells, most probably immature enterocytes arising

from AMPs. Altogether, our data suggest that the larval intestine maintains its integrity upon

aggression through two mechanisms, i) enterocyte flattening which increases the cell surface

without new cycles of endoreplication and ii) producing new immature enterocytes. Both

mechanisms may allow the intestine to maintain the intestinal barrier, avoiding the bacteria to

penetrate into the internal milieu and therefore preventing septicemia.

Discussion

The long-term effects of chronic exposure to low doses of potential noxious products cannot

be easily revealed by conventional epidemiological studies (which are long lasting and

expensive) or conventional (eco)toxicological approaches based on acute toxicity. The use of

amenable and cheap laboratory models becomes of prime importance to pinpoint discrete

chronic adverse effects. *D. melanogaster* fulfills all the required criteria to be such a model.

Indeed, *Drosophila* is easily handling, cheap to rear and the availability of genetic, cellular

and molecular tools allows to detect tenuous effects and to decipher the underlying

physiological mechanisms impaired (Scott and Buchon, 2019). Moreover, the main

physiological functions and genetic networks governing the *Drosophila* gut physiology are

well conserved among the animal kingdom (Pasco et al., 2015), allowing to provide

informative outputs for both human health and wildlife.

Btk bioinsecticides have recently been shown to induce larval lethality and delay the

imago emergence of many different *Drosophila* species at doses ranging from 5×10^7 to 1×10^9

CFU of *Btk* spores/g of fly food medium (Babin et al., 2019; Cossentine et al., 2016). Larval

lethality is observed for the highest doses (Babin et al., 2019) and could be explained by the

cross-order toxicity of the Cry2A toxins (Cry2Aa and Cry2Ab are present in *Btk*

bioinsecticides) towards larvae of certain mosquito species (Frankenhuyzen, 2017) and

references herein). In this study, we investigated the physiological and cellular mechanisms

disturbed by *Btk* products which are responsible for developmental delay and growth defect.

To best mimic environmental conditions, we used doses potentially present in the field after

agricultural spraying. The lower dose we used corresponds to the amount deposited on the

vegetables after one treatment. The highest dose reflects the amount that may be encountered

after multiple treatments, even though data regarding *Bt* accumulation in the fields remain

scarce. Indeed, the half-life of the entomopathogenic Cry1 toxins being estimated at ten days

in the field (Hung et al., 2016), weekly sprays are recommended by the manufacturer and up

to 10 are authorized. Regarding the persistence of spores, it has been shown that 72 hours

after spraying of chards with *Btk* bioinsecticides, almost all of the *Btk* was still present at the

surface of the leaves (Bizzarri and Bishop, 2008). Ten days later, the amount of *Btk*

bioinsecticides corresponded to 50% of the initial dose sprayed on the leaves. The amount of

Bt found on the leaves approaches normal values only 28 days after treatment ($<10^1$ CFU/g)

(Raymond et al., 2010). Furthermore, once *Bt* has reached the soil, it persists more than a year

in the treated soil (Vettori et al., 2003). Our study highlights that such a persistence and

accumulation of *Btk* bioinsecticides may affect the development of non-target insects.

Nonetheless, the impacts are mainly due to the synergy between *Btk* spores and crystals since

either the spores ($Btk^{\Delta Cry}$) or the purified crystals alone do not promote detectable

developmental delay or growth defects in *Drosophila* larvae. These data are in agreement

with the study analyzing the safety of *Btk* purified Cry toxins on *Drosophila* larval

development (Haller et al., 2017).

Interestingly, the effects we have characterized here are reminiscent of those observed

on the target organisms *Helicoverpa armigera* and *H. zea*, *Spodoptera frugiperda* and

Choristoneura fumiferana. Indeed, when fed with sublethal doses of *Btk* spores, all these

target organisms display developmental delays and/or reduced pupal size (Bauce et al., 2006;

Polanczyk and Alves, 2005; Rabinovitch et al., 2017; Sedaratian et al., 2013). Moreover,

recent work has also shown that the potent pathogens *P. entomophila* and *P. aeruginosa* or

huge amounts of the opportunistic bacterium *Erwinia carotovora carotovora* induce

developmental delay and growth defects in *Drosophila* larvae (Houtz et al., 2019). These data

suggest that developmental delay and/or growth defects must be general consequences of

intestinal infections. The compromised gut digestive capacity of the infected larvae (whatever

the insect) is likely the main reason for such developmental defects. Indeed, we and others

have shown in the adult (Buchon et al., 2009; Chakrabarti et al., 2012; Erkosar et al., 2015;

Loudhaief et al., 2017; Wong et al., 2016) or in larva (this work) that allochthonous bacteria

induce a more or less widespread apoptosis of enterocytes leading to a decreased digestive

and/or absorptive capacities. The developmental delay and the growth defect likely reflect the

degree of damages suffered by the intestine. Houtz and colleagues observed a delay in

pupation of up to 4 days using strong pathogens (Houtz et al., 2019) while we observed here a

delay between 16 hours (Delfin) and 24 hours (Dipel) with *Btk* products. Therefore, while *Btk*

behaves as a strong pathogen on target organisms, ultimately leading to death (in fact the

expected effect), it has milder but significant impacts on the development of non-target insect.

Our work demonstrates that the composition of the diet is important to overcome *Btk*

intestinal infection. Increasing the protein content of the nutritive medium rescues both

growth defects and developmental delay. Our data are in agreement with previous

observations in mammals and insects infected with various pathogenic bacteria or viruses

showing that protein intake is important to withstand attacks by pathogens, to support an

efficient immune defense and to promote growth, while intake of carbohydrates is deleterious

for all these functions (Bing et al., 2018; Lee et al., 2006; Reis, 2016; Rodrigues et al., 2015;

Shikano and Cory, 2014; Soultoukis and Partridge, 2016). Moreover, our data show that the

commensal bacterium *L. plantarum* also participates in the protection of larvae from *Btk*

intestinal aggression in accordance with previous works involving commensal bacteria in gut

protection against pathogens (Ubeda et al., 2017; Van Arnam et al., 2018). In *Drosophila*, *L.*

plantarum has been shown to protect adult flies from *Sm-* or *P. aeruginosa*-induced mortality

and this protection is specific since the commensal bacteria *Enterococcus faecalis* or *B.*

subtilis do not operate as such (Blum et al., 2013). Our work strongly suggests that the benefit

brought by *L plantarum* to the larvae of enduring *Btk* aggression passes through an enhanced

protein digestion and/or absorption by the gut in line with previous observations (Erkosar et

al., 2015). Finally, we did not observe a decrease in food intake suggesting that spores and

commercial products do not induce avoidance behavior. Together, our data support the model

that the impact of the *Btk* products on the development of the non-target organism *D.*

melanogaster (and probably many other arthropods) depends on both the nutritional

environment and the performance of its commensal flora. Consequently, spraying *Btk*

products when environmental conditions are stressful would amplify the adverse effects of

Btk products. Our work also sheds light on the pathogenic opportunism of *Btk* on a broad

range of organisms owing to the conservation of the digestive tract architecture and functions

(Pasco et al., 2015). For instance, the impacts of *Btk* on animals or humans suffering from

dietary deficiencies (very often associated with dysbiosis) may be more deleterious than on

individuals with a well-balanced diet. Of note, bacteria of the *B. cereus* group are well-known

opportunistic pathogens responsible for foodborne poisoning (EFSA BIOHAZ, 2016; Wells-

Bennik et al., 2016). Therefore, a more accurate monitoring of the presence of *Btk* in food and

environment is needed to avoid any setbacks, especially in the current context where organic

farming is gaining market share.

Finally, our investigations have unraveled two concomitant mechanisms allowing

Drosophila larvae to overcome an intestinal infection. It was thought so far that the

Drosophila larval gut was unable to regenerate due to the short period covering the larval

stages (about 5-6 days) and the inability to give birth to news cells to replenish the damaged

ones (Miguel-Aliaga et al., 2018). Only two choices were thus offered to the larvae: *Live or*

let die. However, we have uncovered a third choice for the infected larvae: *Die another day*.

First, we demonstrate that the adult midgut precursors (AMPs) can be diverted from their

original fate. Indeed, upon aggression of the larval intestinal epithelium, the AMPs can

engage toward a process of differentiation to give birth to enterocyte-like cells, which are

intercalated in between AMP islets and enterocytes. We can assume that such newborn cells

could plug the holes caused by the death of enterocytes, thus avoiding the entry of luminal

bacteria into the internal milieu. Unfortunately, newborn cells do not fully differentiate and

probably do not participate in digestive functions as efficiently as well-differentiated

enterocytes. The addition of these misdifferentiated cells to dying enterocytes can explain the

reduced protease activity of the intestine of infected larvae and the subsequent developmental

delay and growth defect. The huge increase in the proliferation of AMPs that we observed at

the end of L3 upon *Btk* spore ingestion could be explained by the necessity to replenish the

right number of AMPs before pupation to achieve a complete adult gut metamorphosis and

therefore produce a viable adult fly. These AMPs replenishment could also participate to the

developmental delay we observed. The work very recently published by Houtz and colleagues

reaches to similar conclusions (Houtz et al., 2019).

We have also uncovered a second unexpected mechanism involved in the maintenance

of gut integrity: the change in enterocyte shape. So far, it has been shown in many species

(insects, mammals or plants) that the loss of cells in response to stressors can be compensated

by the rapid increase in the size of neighboring cells thanks to successive endocycles (Edgar

et al., 2014). Here, we evidence an increase in enterocyte area upon *Btk* spore ingestion but

concomitantly we have observed a thinning of the epithelium. Above all, we did not observe

an increase in the ploidy of the enterocyte nuclei. Interestingly, it has been shown that old

enterocytes in adult *Drosophila* midgut cannot re-enter the endocycle (Xiang et al., 2017).

Therefore, as for old adult enterocytes, L3 enterocytes (born during embryogenesis 7 days

earlier) are probably unable to re-enter a cycle of endoreplication to promote cell growth.

Hence, changing their shape through flattening allows to increase the area covered by one

enterocyte and to participate in maintaining epithelium sealing. Here again, the absence of

endoreplication likely participates to the reduced digestive capacities of the intestine and

therefore to the developmental delay and growth defects.

It has been proposed that lepidopteran larvae succumb to *Btk* spores and toxins

because they are unable to maintain their gut integrity. Conversely, one mean for a

lepidopteran larva to acquire resistance to *Btk* bioinsecticides may rely on its capacity to

maintain/regenerate its intestinal integrity (Castagnola and Jurat-Fuentes, 2016; Castro et al.,

2019; Dubovskiy et al., 2016). Non-target organisms are likely as such because they are able

to regenerate their gut lining faster than *Btk* bioinsecticides destroy it. However, increasing

the amount of *Btk* products provided to *Drosophila* larvae ultimately leads to their death

(Babin et al., 2019), suggesting that the harmfulness of *Btk* products relies on both the

amounts of *Btk* products ingested and the affinity of Cry toxins for more or less specific

receptors. Altogether, these data suggest that *Btk* spores used as bioinsecticides may be not as

safe as expected on non-target organisms. Moreover, the opportunistic virulence of *Btk* for

many organisms should not be overlooked, especially for those suffering from nutritional

stress, dysbiosis or with a compromised innate immunity. A situation that more and more

species will encounter with the climatic changes occurring worldwide. Many investigations

remain to be done to improve the use and efficiency of *Btk* products and to make their use as

safe as possible for both handlers and consumers (wildlife or humans).

MATERIALS AND METHODS

Fly Stocks and genetics

Canton S (WT) (Bloomington #64349); *w, dlg::GFP cc01936* (FlyTrap) was provided by A.

Spradling; *w; myo1A-Gal4* was a gift from N. Tapon; *w, UAS-GC3Ai^{G7S}* were kindly provided

by Magali Suzanne and described in (Schott et al., 2017). Apoptosis was monitored on F1

progeny *w/w; myo1A-Gal4/+; UAS-GC3Ai^{G7S}/+* L3 larvae. The ReDDM fly stock to trace

cell progenies in the midgut was kindly provided by Maria Dominguez (Antonello et al.,

2015). To perform lineage tracing, synchronized eggs (*w/+; esg-Gal4 UAS-GFP/+; tub-*

Gal80^{ts} UAS-H2B::RFP/+) were hatched on contaminated food (see below for the method)

and maintained for 24h at 25°C. Then, L1 larvae were transferred and maintained at 29°C to

alleviate the inhibitory effect of Gal80^{ts} on the Gal4 transcription factor. Intestine were

dissected and observed at mid L3 (between days 4 and 5).

Bacterial strains

L. plantarum^{WJL} that has been described in (Ryu et al., 2008) and sequenced in (Martino et al.,

2015) was kindly provided by Bernard Charroux (IDBM, Marseille, France) and by Renata

Matos (F. Leulier's Lab, IGFL, Lyon, France). *Pseudomonas entomophila* (Vodovar et al.,

2005) was kindly provided by Bruno Lemaitre's Lab (EPFL, Lausanne, Switzerland). *Pe* and

L. plantarum^{WJL} were grown as described in (Loudhaief et al., 2017). The *Btk^{ΔCry}* strain

(identified under the code 4D22) (Gonzalez et al., 1982) was obtained from the *Bacillus*

Genetic Stock Center (www.bgsc.org). Delfin and Dipel formulated bioinsecticides were

bought in an agricultural shop center. The strain *Btk* ABTS-351 was isolated from Dipel on

LB agar.

Spore preparation

Btk vegetative cells were let to grow and sporulate during two weeks at 30°C with shaking in

2L of PGSM medium. The culture was then heated 1h at 70°C to eliminate vegetative cells

and centrifuged for 15 min at 5000g. The pellet was first washed with 1 L of 0.15M NaCl and

twice with sterile water. The final pellet was suspended in 30 mL of sterile water, aliquoted

and lyophilized 24h-48h. The spore titration was determined by serial dilution of a known

weight of lyophilized spores and counting the number of "colony forming unit" (CFU) on LB

agar after incubation at 30°C for 18h.

Crystal purification

Two grams of the commercial product were suspended in 60 mL of sterile water, dissolved

for 5h at 4°C with agitation and then sonicated 4 times 15s with a frequency set at 50%

(Fisherbrand™ Model 505 Sonic Dismembrator). Ten milliliters of the homogenate were

deposited on a step density gradient of 67/72/79% sucrose and centrifuged at 100 000g for

18h at 4°C. Crystals were collected on the interfaces between the different sucrose phases and

washed twice with sterile water. The final pellet was suspended in 12 mL of sterile water,

aliquoted and lyophilized 24h-48h.

Diet composition

548   *Drosophila* were reared at 25°C on conventional medium (referred as the "poor-protein"

549   medium in this study): 8% cornmeal (AB Celnat), 2% yeast (Springaline® BA10/0–PW,

550   Amcan), 2.5% sucrose, 0.8% agar (ref 20768-361, VWR) and 0.6% methylparaben

551   (Tegosept,, ref 789063 Dutscher). The protein rich medium contains 10% yeast instead of 2%.

Intestinal CFU counting

553   Colony forming unit counting were described in (Loudhaief et al., 2017).

Commensal flora estimation

555   Ten flies or ten mid-third instar larvae were washed first with ethanol 70% for 30 seconds,

556   then washed with PBS and crushed in 500 µL of LB medium with a Tissue Lyser (Qiagen ref.

557   85600) at 50Hz for 5 minutes. The homogenate was plated on different selective media to

identify the common commensal bacteria. MRS agar (Sigma ref 69964 + 1 mL of Tween 20/L) at 37°C in anaerobic was used to identify *Lactobacillus plantarum* and *brevis*. BHI agar (Sigma ref 70138) at 37°C allowed the identification of *Enterococcus faecalis*. LB agar (Fisher ref BP1426-2) at 37°C allowed the identification of *Escherichia coli* and *Staphylococcus aureus*. Mannitol medium (Bacto peptone 3g/L, Yeast extract 5g/L, D-Mannitol 25g/L, agar 15g/L) at 30°C allowed the identification of *Commensalibacter intestini* and *Acetobacter pomorum*.

Axenic flies

Adult Canton S flies were let to oviposit eggs for two days at 25°C and 12h/12h day/night cycle on our conventional medium supplemented with a cocktail of four antibiotics: ampicillin, kanamycin, tetracycline and erythromycin (50 µg/mL each). On the next generation, newly hatched adults were immediately removed and transferred on fresh conventional medium containing the four antibiotics. This operation was repeated until flies with axenic gut were obtained. The presence of commensal flora was checked at each fly generation (Fig. S1). Axenic flies were obtained after 6 generations.

Oral infection of larvae

Adult flies were let to oviposit eggs for 4h at 25°C on laying nests containing grape juice medium (25% grape juice, 3% agar, 1.2% sucrose, 2% ethanol, 1% acetic acid). Twenty eggs were collected and transferred in a *Drosophila* vial containing 2g of the desired medium mixed with *Btk* spores, *Btk* crystals or the naphthalene additive. The conditions used for 2g of medium and 20 eggs were: 10^7 CFU of *Btk* spores for the 1X dose; 10^8 CFU of *Btk* spores for the 10X dose; 0.67mg of *Btk* purified crystals (equivalent to the 10X dose and considering that crystals represents 15-30% of spore weight (Agaisse and Lereclus, 1995; Monro, 1961; Murty et al., 1994)); 10^8 CFU of *L. plantarum* or *Pe*; 0.4mg of naphtalene-2-sulfonic acid sodium salt (ref A13066, Alfa Aesar) (equivalent to the 10X dose). Vials were incubated at 25°C with a 12h/12h day/night cycle until further processing.

Pupation curve

Immobilized larvae forming white pupae were counted. The number of pupae was expressed

as the percentage of the final number of pupae. While 20 eggs were deposited in each tube, in

some conditions, especially with Dipel and Delfin, we never got 20 pupae probably due to

previously described larval lethality induced by the *Btk* products (Babin et al., 2019). Table 1

summarizes the developmental time at which 10%, 50% and 90% of the larvae were

immobilized for all the conditions we tested. For each condition described above, the

experiment was performed at least in three experimental blocks of triplicates.

Pupal size

On day 7, 8 and 9 after egg laying, pupae were gently removed from the vial, washed in water

and slightly dried with tissue paper. Pupae were placed on a microscope slide and pictures

were taken with a numeric microscope VHX-2000 (Keyence). Pupal length and width were

measured with Image J software and pupal volume was calculated using the following

formula for ellipsoid: $V= [(4/3) \times 3,14 \times (L/2) \times (l/2)^2]$ where V is the volume, L the length

and l the width of the pupa.

Larval feeding behavior and Food intake

Quantification of larval food intake was performed according to (Bjordal et al., 2014) with the

following modifications: we deposited a mix of 50% yeast/0.75% of the food blue dye

Erioglaucin (ref 861146, Sigma) in the center of a Petri dish lid (diameter: 35mm) to feed 20

mid-third larvae (previously infected or not) *ad libitum*. After 2h at 25°C, larvae were

collected and washed with water. The number of white larvae (indicating no feeding) and blue

larvae was counted. Ten blue larvae were transferred into a microtube containing 500 μL of

PBS, crushed at 50Hz for 2 minutes with a TissueLyser (Qiagen, ref 85600) and centrifuged

at 10000g for 5 minutes at 4°C. The amount of Erioglaucin in the supernatant was quantified

at 629 nm with a spectrophotometer (SpectraMax® Plus384, Molecular device).

Midgut permeability

The same technique as described above was used except that the mix contained 50% yeast

and 1.25 mg/ml Dextran FITC (ref 46944, Sigma) in water. Twenty previously orally infected

mid-third instar larvae were placed in the mix. After 2h at 25°C, the larvae were collected,

washed with water and placed at least 30 minutes at 4°C. Pictures were taken with a

fluorescent stereoscopic AxioZoom V16 (Zeiss) microscope.

Metabolic assay

Intestinal protease activity assay was performed as described in (Nawrot-Esposito et al., 2017)

with 8 mid-third instar larval midguts per condition. Lipids were assayed as described in

(Tennessen et al., 2014). The tri- and diacylglycerol (TAG and DAG) were measured on 3µl

of hemolymph obtained from 5 days old larvae using the Triglyceride Reagent (ref T2449,

Sigma) and Free Glycerol Reagent (ref F6428, Sigma). The metabolite quantity was reported

per mg of protein or per µL of hemolymph. Protein assay was achieved with Biorad Protein

Assay Dye Reagent Concentrate (ref 5000006, Biorad). All experiments were performed at

least 3 times in triplicates.

Lipids labelling

Five days after infection (mid-third instar larvae), ten larval guts were dissected in PBS and

fixed 40 min in 4% formaldehyde in PBS. Guts were then washed twice in PBS and further

incubated for 30 min. either in 5µg/ml of BODIPY™ 493/503 (ref D3922, Invitrogen™

Molecular Probes™) or in a 0.06% Oil-red-O solution (ref 189400250, Acros Organics). Guts

were then washed twice with PBS. For BODIPY™ labelling guts were mounted in

Fluoroshield/DAPI (ref F6057, Sigma) and immediately observed under a fluorescent

microscope (Zeiss Axioplan Z1 with Apotome 2). For Oil-red-O labelling, guts were mounted

in 80% Glycerol/PBS and observed with a numeric microscope (VHX 2000, Keyence).

Immunohistochemistry, image capture and processing

Ten guts of late third instar larvae were dissected in PBS and fixed for 40 min. in 4%

formaldehyde in PBS. Guts were then washed twice with PBS-0.1% Triton X100, incubated

for 3h in Blotto (2.5% FBS, 0.1% Triton X-100, 0.02% sodium azide in PBS) and further

incubated with the primary antibody overnight at 4°C in Blotto. The following dilutions were

used: Mouse anti-Armadillo (DSHB #N2 7A1) at 1/50; Rabbit anti-Phospho-histone-3 (ref

9701S, Cell signaling) at 1/500. Guts were washed once with Blotto before incubation in the

secondary antibody and/or phalloidin (2h at room temperature in Blotto). The secondary

antibodies were the followings: AlexaFluor^{TM} 546 donkey anti-Rabbit IgG (ref A10040,

Invitrogen) used at 1:500; AlexaFluor^{TM} 647 goat anti-Mouse IgG (H+L) (ref A21235,

Invitrogen) used at 1/500; Alexa Fluor™ 555 Phalloidin (ref A34055, Invitrogen) used at

1/1000. Guts were finally washed several times in PBS, mounted in Fluoroshield/DAPI (ref

F6057, Sigma) and observed with a Zeiss Axioplan Z1 with Apotome 2 microscope. Images

were analyzed using ZEN (Zeiss) software or Photoshop. For cell death monitoring (in

myo1A-Gal4 UAS- UAS-GC3Ai^{G7S} larvae), guts were dissected in PBS, immediately mounted

in 80% Schneider medium/20% glycerol and pictures were rapidly taken with a fluorescent

stereoscopic microscope (AxioZoom V16, Zeiss). Image acquisition was performed at the

Microscopy platform of the Institut Sophia Agrobiotech (INRAE 1355-UCA-CNRS 7254-

Sophia Antipolis).

Polyploidy estimation

Ploidy was calculated by measuring the area of the nuclei of late L3 larvae. Only round nuclei

were analyzed. The average nucleus area of diploid adult midgut precursor cells was

calculated. This average area (8µm²) was assigned to the 2C ploidy (C corresponding to

chromatin copy number). The theoretical average area of 4C, 8C, 16C, 32C and 64C nuclei

was calculated on the basis of the measured 2C average area. 109 nuclei for the control larvae,

184 nuclei for *Btk$^{\Delta Cry}$* 10X larvae and 306 nuclei for Dipel 10X larvae were counted and

binned into ploidy groups.

Statistics

When "n" was equal or superior to 30, statistical analysis was performed using a parametric

T-test. An F-test was systematically done before applying the T-test to verify the homogeneity

of variances. When "n" was inferior to 30, we used the non-parametric pairwise comparisons

of the Wilcoxon-Mann-Whitney test. Significance symbols are the following: ***

($P<=0.001$); ** ($P<=0.01$), * ($P<=0.05$).

Acknowledgement

We would like to thank Nathalie Arquier and Rénald Delanoue for their help in food intake

measurement, lipid dosage and staining, and pupae volume measurement. We thank Julien

Colombani for the fruitful discussions, Pierre Barbero for discussions and correction of the

manuscript, Bernard Charroux, François Leulier and Renata Matos for providing *L.*

plantarum. We also thank Marine Chesneau for technical help during her internship. Thanks

to Olivier Pierre from the Microscopy platform (Institut Sophia Agrobiotech, INRAE 1355-

UCA-CNRS 7254-Sophia Antipolis) for help with the newly acquired Zeiss AxioZoom

Microscope and to Raphaël Rousset for help with statistical analyses and manuscript

correction. Finally, we warmly thank Rénald Delanoue for his help in designing the

experiments and correcting the manuscript.

Competing interests

The authors state that they have no competing interests.

Funding

This work was supported by the French Government (Agence Nationale pour la Recherche,

ANR) through the "Investments for the Future" LABEX SIGNALIFE ANR-11-LABX-0028-

01 and through the ANR-13-CESA-0003-01, by the Institut Olga Triballat (PR2016-19) to

AG and the ANSES PNR-EST & ECOPHYTO II (2018) to AG.

Author contributions

Marie-Paule Nawrot-Esposito: Methodology, Validation, Formal analysis, Investigation,

Resources, Data Curation, Writing - Original Draft **Aurélie Babin:** Methodology, Validation,

Formal analysis, Investigation, Writing - Review & Editing **Matthieu Pasco:** Investigation

Marylène Poirié: Writing - Review & Editing, Funding acquisition **Jean-Luc Gatti:**

Validation, Writing - Review & Editing, Funding acquisition **Armel Gallet:**

Conceptualization, Validation, Formal analysis, Data Curation, Writing - Original Draft,

Writing - Review & Editing, Visualization, Supervision, Project administration, Funding

acquisition

References

Adang, M. J., Crickmore, N. and Jurat-Fuentes, J. L. (2014). Chapter Two – Diversity of
Bacillus thuringiensis Crystal Toxins and Mechanism of Action. *Advances in Insect*
Physiology **47**, 39–87.
Agaisse, H. and Lereclus, D. (1995). How does Bacillus thuringiensis produce so much
insecticidal crystal protein? *J Bacteriol.* **177**, 6027-32. doi: 10.1128/jb.177.21.6027-
6032.1995.
Amichot, M., Curty, C., Benguettat-Magliano, O., Gallet, A. and Wajnberg, E. (2016).
Side effects of Bacillus thuringiensis var. kurstaki on the hymenopterous parasitic wasp
Trichogramma chilonis. *Environ Sci Pollut Res Int* **23**, 3097-103. doi: 10.1007/s11356-015-
5830-7. Epub 2015 Nov 21.
Anagnostou, C., Dorsch, M. and Rohlfs, M. (2010). Influence of dietary yeasts on
Drosophila melanogaster life-history traits. *Entomologia Experimentalis et Applicata* **136**, 1–
11.
ANSES Saisine n° 2013-SA-0039. (2013). Risques en termes de sécurité sanitaire liés à
l'utilisation de souches de Bacillus thuringiensis (Bt) en tant que substances actives dans des
insecticides biologiques. Saisine n°2013-SA-0039,
https://www.anses.fr/fr/system/files/DPR2013sa0039.pdf.
Antonello, Z. A., Reiff, T., Ballesta-Illan, E. and Dominguez, M. (2015). Robust intestinal
homeostasis relies on cellular plasticity in enteroblasts mediated by miR-8-Escargot switch.
EMBO **34**, 2025-41. doi: 10.15252/embj.201591517. Epub 2015 Jun 15.
Apidianakis, Y., Pitsouli, C., Perrimon, N. and Rahme, L. (2009). Synergy between
bacterial infection and genetic predisposition in intestinal dysplasia. *Proc Natl Acad Sci U S A*
106, 20883-8. Epub 2009 Nov 23.
Babin, A., Kolly, S., Schneider, F., Dolivo, V., Zini, M. and Kawecki, T. J. (2014). Fruit
flies learn to avoid odours associated with virulent infection. *Biol Lett.* **10**, 20140048. doi:
10.1098/rsbl.2014.0048. Print 2014 Mar.
Babin, A., Nawrot-Esposito, M.-P., Gallet, A., Gatti, J.-L. and Poirié, M. (2019). Adverse
effects of *Bacillus thuringiensis* bioinsecticide on non-target *Drosophila* species. *bioRxiv*,
541847.
Bauce, É., Carisey, N. and Dupont, A. (2006). Carry over effects of the entomopathogen
Bacillus thuringiensis ssp. Kurstaki on Choristoneura fumiferana (Lepidoptera: Tortricidae)

progeny under various stressful environmental conditions. *Agricultural and Forest*
Entomology **8**, 63-76.
Bing, X., Gerlach, J., Loeb, G. and Buchon, N. (2018). Nutrient-Dependent Impact of
Microbes on Drosophila suzukii Development. *MBio.* **9(2).** mBio.02199-17. doi:
10.1128/mBio.02199-17.
Bizzarri, M. F. and Bishop, A. H. (2008). The ecology of Bacillus thuringiensis on the
Phylloplane: colonization from soil, plasmid transfer, and interaction with larvae of Pieris
brassicae. *Microb Ecol* **56**, 133-9. Epub 2007 Nov 2.
Bjordal, M., Arquier, N., Kniazeff, J., Pin, J. P. and Leopold, P. (2014). Sensing of amino
acids in a dopaminergic circuitry promotes rejection of an incomplete diet in Drosophila. *Cell.*
156, 510-21. doi: 10.1016/j.cell.2013.12.024.
Blum, J. E., Fischer, C. N., Miles, J. and Handelsman, J. (2013). Frequent replenishment
sustains the beneficial microbiome of Drosophila melanogaster. *MBio.* **4**, e00860-13. doi:
10.1128/mBio.00860-13.
Bonfini, A., Liu, X. and Buchon, N. (2016). From pathogens to microbiota: How Drosophila
intestinal stem cells react to gut microbes. S0145-305X(16)30032-5. doi:
10.1016/j.dci.2016.02.008.
Broderick, N. A. and Lemaitre, B. (2012). Gut-associated microbes of Drosophila
melanogaster. *Gut Microbes.* **3**, 307-21. doi: 10.4161/gmic.19896. Epub 2012 May 10.
Buchon, N., Broderick, N. A., Poidevin, M., Pradervand, S. and Lemaitre, B. (2009).
Drosophila intestinal response to bacterial infection: activation of host defense and stem cell
proliferation. *Cell Host Microbe* **5**, 200-11. doi: 10.1016/j.chom.2009.01.003.
Caballero, J., Jiménez-Moreno, N., Orera, I., Williams, T., Fernández, A. B., Villanueva,
M., Ferré, J., Caballero, P. and Ancín-Azpilicueta, C. (2020). Unraveling the Composition
of Insecticidal Crystal Proteins in Bacillus thuringiensis: a Proteomics Approach. *Appl*
Environ Microbiol. **86**, e00476-20. doi: 10.1128/AEM.00476-20. Print 2020 Jun 2.
Casida, J. E. and Bryant, R. J. (2017). The ABCs of pesticide toxicology: amounts, biology,
and chemistry. *Toxicol Res (Camb).* **6**, 755-763. doi: 10.1039/c7tx00198c. eCollection 2017
Nov 1.
Castagnola, A. and Jurat-Fuentes, J. L. (2016). Intestinal regeneration as an insect
resistance mechanism to entomopathogenic bacteria. *Curr Opin Insect Sci.* **15:104-10.**,
10.1016/j.cois.2016.04.008. Epub 2016 Apr 20.
Castro, B., Martinez, L. C., Barbosa, S. G., Serrao, J. E., Wilcken, C. F., Soares, M. A.,
da Silva, A. A., de Carvalho, A. G. and Zanuncio, J. C. (2019). Toxicity and cytopathology
mediated by Bacillus thuringiensis in the midgut of Anticarsia gemmatalis (Lepidoptera:
Noctuidae). *Sci Rep* **9**, 6667.
Chakrabarti, S., Liehl, P., Buchon, N. and Lemaitre, B. (2012). Infection-induced host
translational blockage inhibits immune responses and epithelial renewal in the Drosophila gut.
Cell Host Microbe **12**, 60-70. doi: 10.1016/j.chom.2012.06.001.
Chandler, J. A., Lang, J. M., Bhatnagar, S., Eisen, J. A. and Kopp, A. (2011). Bacterial
communities of diverse Drosophila species: ecological context of a host-microbe model
system. *PLoS Genet.* **7**, e1002272. doi: 10.1371/journal.pgen.1002272. Epub 2011 Sep 22.
Chen, J., Sayadian, A. C., Lowe, N., Lovegrove, H. E. and St Johnston, D. (2018). An
alternative mode of epithelial polarity in the Drosophila midgut. *PLoS Biol.* **16**, e3000041.
doi: 10.1371/journal.pbio.3000041. eCollection 2018 Oct.
Colombani, J., Raisin, S., Pantalacci, S., Radimerski, T., Montagne, J. and Leopold, P.
(2003). A nutrient sensor mechanism controls Drosophila growth. *Cell.* **114**, 739-49.
Cossentine, J., Robertson, M. and Xu, D. (2016). Biological Activity of Bacillus
thuringiensis in Drosophila suzukii (Diptera: Drosophilidae). *J Econ Entomol* **22**.
Delanoue, R., Slaidina, M. and Leopold, P. (2010). The steroid hormone ecdysone controls
systemic growth by repressing dMyc function in Drosophila fat cells. *Dev Cell.* **18**, 1012-21.
doi: 10.1016/j.devcel.2010.05.007.

Dubovskiy, I. M., Grizanova, E. V., Whitten, M. M., Mukherjee, K., Greig, C., Alikina,
T., Kabilov, M., Vilcinskas, A., Glupov, V. V. and Butt, T. M. (2016). Immuno-
physiological adaptations confer wax moth Galleria mellonella resistance to Bacillus
thuringiensis. *Virulence* **7**, 860-870. Epub 2016 Mar 30.
Edgar, B. A., Zielke, N. and Gutierrez, C. (2014). Endocycles: a recurrent evolutionary
innovation for post-mitotic cell growth. *Nat Rev Mol Cell Biol.* **15**, 197-210. doi:
10.1038/nrm3756.
EFSA BIOHAZ. (2016). Risks for public health related to the presence of Bacillus cereus
and other Bacillus spp. including Bacillus thuringiensis in foodstuffs. *EFSA Journal* **14**,
e04524.
EFSA BIOHAZ Panel. (2016). Scientific opinion on the risks for public health related to the
presence of Bacillus cereus and other Bacillus spp. including Bacillus thuringiensis in
foodstuffs. EFSA Journal 2016;14(7):4524, 93 pp. doi:10.2903/j.efsa.2016.4524.
Erkosar, B., Storelli, G., Mitchell, M., Bozonnet, L., Bozonnet, N. and Leulier, F. (2015).
Pathogen Virulence Impedes Mutualist-Mediated Enhancement of Host Juvenile Growth via
Inhibition of Protein Digestion. *Cell Host Microbe* **18**, 445-55. doi:
10.1016/j.chom.2015.09.001. Epub 2015 Oct 1.
Fast, D., Duggal, A. and Foley, E. (2018). Monoassociation with Lactobacillus plantarum
Disrupts Intestinal Homeostasis in Adult Drosophila melanogaster. *MBio.* **9(4).** mBio.01114-
18. doi: 10.1128/mBio.01114-18.
Fiuza, L. M., Polanczyk, R. A. and Crickmore, N. (2017). Bacillus thuringiensis and
Lysinibacillus sphaericus: Springer International Publishing.
Frankenhuyzen, K. (2017). Specificity and Cross-order Activity of Bacillus thuringiensis
Pesticidal Proteins. In *Bacillus thuringiensis and Lysinibacillus sphaericus*, (ed. L. M. Fiuza
R. A. Polanczyk and N. Crickmore), pp. 127-172: Springer International Publishing.
Frisvold, G. B. (2018). How low can you go? Estimating impacts of reduced pesticide use.
Pest Manag Sci **8**.
Gonzalez, J. M., Jr., Brown, B. J. and Carlton, B. C. (1982). Transfer of Bacillus
thuringiensis plasmids coding for delta-endotoxin among strains of B. thuringiensis and B.
cereus. *Proc Natl Acad Sci U S A* **79**, 6951-5.
Grisolia, C. K., Oliveira, R., Domingues, I., Oliveira-Filho, E. C., Monerat, R. G. and
Soares, A. M. (2009). Genotoxic evaluation of different delta-endotoxins from Bacillus
thuringiensis on zebrafish adults and development in early life stages. *Mutat Res* **672**, 119-23.
Haller, S., Romeis, J. and Meissle, M. (2017). Effects of purified or plant-produced Cry
proteins on Drosophila melanogaster (Diptera: Drosophilidae) larvae. *Sci Rep.* **7**, 11172. doi:
10.1038/s41598-017-10801-4.
Hietakangas, V. and Cohen, S. M. (2009). Regulation of tissue growth through nutrient
sensing. *Annu Rev Genet* **43**, 389-410.
Houtz, P., Bonfini, A., Bing, X. and Buchon, N. (2019). Recruitment of Adult Precursor
Cells Underlies Limited Repair of the Infected Larval Midgut in Drosophila. *Cell Host*
Microbe **28**, 30409-3.
Hung, T. P., Truong, L. V., Binh, N. D., Frutos, R., Quiquampoix, H. and Staunton, S.
(2016). Fate of insecticidal Bacillus thuringiensis Cry protein in soil: differences between
purified toxin and biopesticide formulation. *Pest Manag Sci.* **72**, 2247-2253. doi:
10.1002/ps.4262. Epub 2016 Mar 24.
Jiang, H. and Edgar, B. A. (2009). EGFR signaling regulates the proliferation of Drosophila
adult midgut progenitors. *Development.* **136**, 483-93. doi: 10.1242/dev.026955.
Keita, S., Masuzzo, A., Royet, J. and Kurz, C. L. (2017). Drosophila larvae food intake
cessation following exposure to Erwinia contaminated media requires odor perception, Trpa1
channel and evf virulence factor. *J Insect Physiol.* **99:25-32.**, 10.1016/j.jinsphys.2017.02.004.
Epub 2017 Feb 21.

Layalle, S., Arquier, N. and Leopold, P. (2008). The TOR pathway couples nutrition and
developmental timing in Drosophila. *Dev Cell* **15**, 568-77. doi: 10.1016/j.devcel.2008.08.003.
Lee, K. P., Cory, J. S., Wilson, K., Raubenheimer, D. and Simpson, S. J. (2006). Flexible
diet choice offsets protein costs of pathogen resistance in a caterpillar. *Proc Biol Sci.* **273**,
823-9. doi: 10.1098/rspb.2005.3385.
Lee, K. Z., Lestradet, M., Socha, C., Schirmeier, S., Schmitz, A., Spenle, C., Lefebvre,
O., Keime, C., Yamba, W. M., Bou Aoun, R. et al. (2016). Enterocyte Purge and Rapid
Recovery Is a Resilience Reaction of the Gut Epithelium to Pore-Forming Toxin Attack. *Cell*
Host Microbe. **20**, 716-730. doi: 10.1016/j.chom.2016.10.010. Epub 2016 Nov 23.
Loudhaief, R., Brun-Barale, A., Benguettat, O., Nawrot-Esposito, M. P., Pauron, D.,
Amichot, M. and Gallet, A. (2017). Apoptosis restores cellular density by eliminating a
physiologically or genetically induced excess of enterocytes in the Drosophila midgut.
Development **144**, 808-819. doi: 10.1242/dev.142539.
Martino, M. E., Bayjanov, J. R., Joncour, P., Hughes, S., Gillet, B., Kleerebezem, M.,
Siezen, R., van Hijum, S. A. and Leulier, F. (2015). Resequencing of the Lactobacillus
plantarum Strain WJL Genome. *Genome Announc* **3(6).** e01382-15. doi:
10.1128/genomeA.01382-15.
Mathur, D., Bost, A., Driver, I. and Ohlstein, B. (2010). A transient niche regulates the
specification of Drosophila intestinal stem cells. *Science.* **327**, 210-3. doi:
10.1126/science.1181958.
Micchelli, C. A., Sudmeier, L., Perrimon, N., Tang, S. and Beehler-Evans, R. (2011).
Identification of adult midgut precursors in Drosophila. *Gene Expr Patterns.* **11**, 12-21. doi:
10.1016/j.gep.2010.08.005. Epub 2010 Sep 23.
Miguel-Aliaga, I., Jasper, H. and Lemaitre, B. (2018). Anatomy and Physiology of the
Digestive Tract of Drosophila melanogaster. *Genetics.* **210**, 357-396. doi:
10.1534/genetics.118.300224.
Monro, R. E. (1961). Protein turnover and the formation of protein inclusions during
sporulation of Bacillus thuringiensis. *Biochem J.* **81:225-32.**, 10.1042/bj0810225.
Murty, M. G., Srinivas, G., Bora, R. S. and Sekar, V. (1994). A simple method for
separation of the protein crystal from Bacillus thuringiensis using carboxymethyl cellulose
column chromatography. *Journal of Microbiological Methods* **19**, 103-110.
Nawrot-Esposito, M., Loudhaie, f. R. and Gallet, A. (2017). Protease activity assay in fly
intestines. *Bio-protocol* **7**, e2560.
Palm, W., Sampaio, J. L., Brankatschk, M., Carvalho, M., Mahmoud, A., Shevchenko,
A. and Eaton, S. (2012). Lipoproteins in Drosophila melanogaster--assembly, function, and
influence on tissue lipid composition. *PLoS Genet* **8**, e1002828. doi:
10.1371/journal.pgen.1002828. Epub 2012 Jul 26.
Palma, L., Munoz, D., Berry, C., Murillo, J. and Caballero, P. (2014). Bacillus
thuringiensis toxins: an overview of their biocidal activity. *Toxins (Basel).* **6**, 3296-325. doi:
10.3390/toxins6123296.
Panigrahi, P., Parida, S., Nanda, N. C., Satpathy, R., Pradhan, L., Chandel, D. S.,
Baccaglini, L., Mohapatra, A., Mohapatra, S. S., Misra, P. R. et al. (2017). A randomized
synbiotic trial to prevent sepsis among infants in rural India. *Nature.* **548**, 407-412. doi:
10.1038/nature23480. Epub 2017 Aug 16.
Pasco, M. Y., Loudhaief, R. and Gallet, A. (2015). The cellular homeostasis of the gut:
what the Drosophila model points out. *Histol Histopathol.* **Vol 30**, pp 277-292.
Polanczyk, R. A. and Alves, S. B. (2005). Biological parameters of Spodoptera frugiperda
(J.E. Smith) (Lepidoptera: Noctuidae) assayed with Bacillus thuringiensis berliner. *Scientia*
Agricola **62**, 464-468.
Rabinovitch, L., Vivoni, A. M., Machado, V., Knaak, N., Berlitz, D. L., Polanczyk, R. A.
and Fiuza, L. M. (2017). Bacillus thuringiensis Characterization: Morphology, Physiology,
Biochemistry, Pathotype, Cellular, and Molecular Aspects. In *Bacillus thuringiensis and*

Lysinibacillus sphaericus, (ed. L. M. Fiuza R. A. Polanczyk and N. Crickmore), pp. 1-18: Springer International Publishing.

Raymond, B., Wyres, K. L., Sheppard, S. K., Ellis, R. J. and Bonsall, M. B. (2010). Environmental factors determining the epidemiology and population genetic structure of the Bacillus cereus group in the field. *PLoS* **6**, e1000905.

Reis, T. (2016). Effects of Synthetic Diets Enriched in Specific Nutrients on Drosophila Development, Body Fat, and Lifespan. *PLoS One.* **11**, e0146758. doi: 10.1371/journal.pone.0146758. eCollection 2016.

Rodrigues, M. A., Martins, N. E., Balance, L. F., Broom, L. N., Dias, A. J., Fernandes, A. S., Rodrigues, F., Sucena, E. and Mirth, C. K. (2015). Drosophila melanogaster larvae make nutritional choices that minimize developmental time. *J Insect Physiol.* **81:69-80.**, 10.1016/j.jinsphys.2015.07.002. Epub 2015 Jul 4.

Rosi-Marshall, E. J., Tank, J. L., Royer, T. V., Whiles, M. R., Evans-White, M., Chambers, C., Griffiths, N. A., Pokelsek, J. and Stephen, M. L. (2007). Toxins in transgenic crop byproducts may affect headwater stream ecosystems. *Proc Natl Acad Sci U S A* **104**, 16204-8.

Rubio-Infante, N. and Moreno-Fierros, L. (2016). An overview of the safety and biological effects of Bacillus thuringiensis Cry toxins in mammals. *J Appl Toxicol* **36**, 630-48. doi: 10.1002/jat.3252. Epub 2015 Nov 4.

Ryu, J. H., Kim, S. H., Lee, H. Y., Bai, J. Y., Nam, Y. D., Bae, J. W., Lee, D. G., Shin, S. C., Ha, E. M. and Lee, W. J. (2008). Innate immune homeostasis by the homeobox gene caudal and commensal-gut mutualism in Drosophila. *Science* **319**, 777-82. doi: 10.1126/science.1149357. Epub 2008 Jan 24.

Schott, S., Ambrosini, A., Barbaste, A., Benassayag, C., Gracia, M., Proag, A., Rayer, M., Monier, B. and Suzanne, M. (2017). A fluorescent toolkit for spatiotemporal tracking of apoptotic cells in living Drosophila tissues. *Development.* **144**, 3840-3846. doi: 10.1242/dev.149807. Epub 2017 Sep 4.

Schwarzer, M., Makki, K., Storelli, G., Machuca-Gayet, I., Srutkova, D., Hermanova, P., Martino, M. E., Balmand, S., Hudcovic, T., Heddi, A. et al. (2016). Lactobacillus plantarum strain maintains growth of infant mice during chronic undernutrition. *Science* **351**, 854-7. doi: 10.1126/science.aad8588.

Scott, J. G. and Buchon, N. (2019). Drosophila melanogaster as a powerful tool for studying insect toxicology. *Pesticide Biochemistry and Physiology* **161**, 95-103.

Sedaratian, A., Fathipour, Y., Talaei-Hassanloui, R. and Jurat-Fuentes, J. L. (2013). Fitness costs of sublethal exposure to Bacillus thuringiensis in Helicoverpa armigera: a carryover study on offspring. *Journal of Applied Entomology* **137**, 540-549.

Shikano, I. and Cory, J. S. (2014). Dietary mechanism behind the costs associated with resistance to Bacillus thuringiensis in the cabbage looper, Trichoplusia ni. *PLoS One* **9**, e105864. doi: 10.1371/journal.pone.0105864. eCollection 2014.

Soultoukis, G. A. and Partridge, L. (2016). Dietary Protein, Metabolism, and Aging. *Annu Rev Biochem* **85**, 5-34.

Stensmyr, M. C., Dweck, H. K., Farhan, A., Ibba, I., Strutz, A., Mukunda, L., Linz, J., Grabe, V., Steck, K., Lavista-Llanos, S. et al. (2012). A conserved dedicated olfactory circuit for detecting harmful microbes in Drosophila. *Cell.* **151**, 1345-57. doi: 10.1016/j.cell.2012.09.046.

Storelli, G., Defaye, A., Erkosar, B., Hols, P., Royet, J. and Leulier, F. (2011). Lactobacillus plantarum promotes Drosophila systemic growth by modulating hormonal signals through TOR-dependent nutrient sensing. *Cell Metab* **14**, 403-14. doi: 10.1016/j.cmet.2011.07.012.

Surendran, S., Huckesfeld, S., Waschle, B. and Pankratz, M. J. (2017). Pathogen-induced food evasion behavior in Drosophila larvae. *J Exp Biol.* **220**, 1774-1780. doi: 10.1242/jeb.153395. Epub 2017 Mar 2.

Tank, J. L., Rosi-Marshall, E. J., Royer, T. V., Whiles, M. R., Griffiths, N. A.,
Frauendorf, T. C. and Treering, D. J. (2010). Occurrence of maize detritus and a transgenic
insecticidal protein (Cry1Ab) within the stream network of an agricultural landscape. *Proc*
Natl Acad Sci U S A **107**, 17645-50. Epub 2010 Sep 27.
Téfit, M. A., Gillet, B., Joncour, P., Hughes, S. and Leulier, F. (2018). Stable association
of a Drosophila-derived microbiota with its animal partner and the nutritional environment
throughout a fly population's life cycle. *Journal of Insect Physiology* **106**, 2-12.
Tefit, M. A. and Leulier, F. (2017). Lactobacillus plantarum favors the early emergence of
fit and fertile adult Drosophila upon chronic undernutrition. *J Exp Biol.* **220**, 900-907. doi:
10.1242/jeb.151522. Epub 2017 Jan 6.
Tennessen, J. M., Barry, W. E., Cox, J. and Thummel, C. S. (2014). Methods for studying
metabolism in Drosophila. *Methods* **68**, 105-15. doi: 10.1016/j.ymeth.2014.02.034. Epub
2014 Mar 12.
Ubeda, C., Djukovic, A. and Isaac, S. (2017). Roles of the intestinal microbiota in pathogen
protection. *Clin Transl Immunology.* **6**, e128. doi: 10.1038/cti.2017.2. eCollection 2017 Feb.
Van Arnam, E. B., Currie, C. R. and Clardy, J. (2018). Defense contracts: molecular
protection in insect-microbe symbioses. *Chem Soc Rev.* **47**, 1638-1651. doi:
10.1039/c7cs00340d.
Vettori, C., Paffettib, D., Saxenac, D., Stotzkyc, G. and Giannini, R. (2003). Persistence of
toxins and cells of Bacillus thuringiensis subsp kurstaki introduced in sprays to Sardinia soils.
Soil biology & biochemistry **35**, 1635-1642.
Vilas-Boas, G. T., Peruca, A. P. and Arantes, O. M. (2007). Biology and taxonomy of
Bacillus cereus, Bacillus anthracis, and Bacillus thuringiensis. *Can J Microbiol* **53**, 673-87.
Vodovar, N., Vinals, M., Liehl, P., Basset, A., Degrouard, J., Spellman, P., Boccard, F.
and Lemaitre, B. (2005). Drosophila host defense after oral infection by an
entomopathogenic Pseudomonas species. *Proc Natl Acad Sci U S A* **102**, 11414-9. Epub 2005
Aug 1.
Wells-Bennik, M. H., Eijlander, R. T., den Besten, H. M., Berendsen, E. M., Warda, A.
K., Krawczyk, A. O., Nierop Groot, M. N., Xiao, Y., Zwietering, M. H., Kuipers, O. P. et
al. (2016). Bacterial Spores in Food: Survival, Emergence, and Outgrowth. *Annu Rev Food*
Sci Technol **7:457-82.**, 10.1146/annurev-food-041715-033144.
Wong, A. C., Vanhove, A. S. and Watnick, P. I. (2016). The interplay between intestinal
bacteria and host metabolism in health and disease: lessons from Drosophila melanogaster.
Dis Model Mech. **9**, 271-81. doi: 10.1242/dmm.023408.
Xiang, J., Bandura, J., Zhang, P., Jin, Y., Reuter, H. and Edgar, B. A. (2017). EGFR-
dependent TOR-independent endocycles support Drosophila gut epithelial regeneration. *Nat*
Commun. **8:15125.**, 10.1038/ncomms15125.
971
972

973   **Table 1**

974   Time in hours when 10%, 50% and 90% larvae enter the pupation stage (immobilized)

975   expressed in hours after egg laying. "n" corresponds to the number of larvae counted in each

976   condition. Note that we compiled the controls per year of experiments (encompassing years

977   from 2017 to 2019).

% of pupation	10%	50%	90%	n
Control 2017	140.5	156.0	169.0	183
Control 2018	142.0	155.0	168.5	211
Control 2019	142.5	155.5	171.0	203
Delfin 1X	143.0	159.5	179.5	262
Delfin 10X	149.5	165.5	185.0	214
Dipel 10X	151.5	170.0	194.0	152
ABTS-351 10X	147.5	165.0	189.0	159
$Btk^{\Delta Cry}$ 1X	141.5	157.0	170.5	263
$Btk^{\Delta Cry}$ 10X	141.5	157.5	170.5	204
Naphthalene	142.5	155.0	168.0	212
Crystals	142.0	156.5	169.5	143
Control protein-rich medium	117.0	128.5	137.5	212
Delfin 1X protein-rich medium	115.0	127.0	138.0	195
Delfin 10X protein-rich medium	117.0	130.5	141.0	207
L. plantarum	128.0	146.0	162.5	144
Delfin 10X + *L plantarum*	127.0	143.0	158.0	153
Control Axenic	134.0	151.5	162.5	171
Delfin 10X Axenic	142.0	159.0	182.0	172

Figure legends

Figure 1: *Btk* bioinsecticides impacts on developmental delay and growth defects.

(**A**) Pupation curve of larvae fed with commercial products (Delfin and Dipel) or with a

homemade preparation of spores of the *Btk* strain ABTS-351 (Dipel). In this figure and in all

the following figures, the cumulative percentage of pupation is shown over time. (**B**) Pupation

curve of larvae fed with a *Btk* strain devoid of Cry toxins ($Btk^{\Delta Cry}$) or with Cry toxin

containing crystals purified from Dipel or with Naphthalene sulfonic acid at a dose equivalent

to the amount found in Delfin 10X. In A and B, red dashed lines mark 90% of the population

reaching pupation. (**C**) Pupae volume in the different conditions listed on the abscissa. n=

number of individual counted. For the controls, n correspond to the total number of pupae

measured when one compiled all the internal controls. Errors bars represent the Standard

Deviation of the Mean (SEM). *=P<0.05; ***=P<0.001 compared with controls.

Figure 2: *Btk* bioinsecticides and larval feeding behavior.

(**A**) L3 midgut load of larvae fed with either spores of $Btk^{\Delta Cry}$ or Delfin. (**B** and **C**) Percentage

of L3 (B) and L2 (C) larvae presenting blue dyed food in the intestine. (**D** and **E**) Amount of

dye ingested by L3 (D) and L2 (E) larvae. Green bars: control larvae (fed with water); light

blue bars: 1X dose of $Btk^{\Delta Cry}$ spores; dark blue bars: 10X dose of $Btk^{\Delta Cry}$ spores; yellow bars:

1X dose of Delfin; Red bars: 10X dose of Delfin. n= number of individuals analyzed. Errors

bars represent the SEM. *=P<0.05, **=P<0.01 compared with controls.

1001

Figure 3: Influence of a protein rich-diet on the Delfin impacts

(**A**) Measurement of the protease activity of the intestine of L3 larvae raised on control

medium contaminated with H2O (control) or with 10X doses of spores of $Btk^{\Delta Cry}$ or Delfin or

1X dose of spores of Delfin. (**B**) Pupation curve of larvae raised on a protein rich medium

(10% yeast instead of 2%) contaminated with 1X or 10X doses of Delfin. Note the shortening

of the developmental time of the control larvae raised on the protein-rich medium (10% yeast)

compared to larvae raised on the protein poor-medium (2% yeast). (**C**) Pupae volume derived

from larvae raised on the protein-rich medium and in the different conditions listed on the

abscissa.

1011

Figure 4: *Btk* bioinsecticides interaction with commensal flora

(**A**) Pupation curve of larvae reared on a medium complemented with 10^6 CFU of *L*

plantarum and contaminated with Delfin 10X (pink) or not (purple) compared to control

larvae (green) or Delfin 10X only (red). (**B**) Pupae volume derived from larvae raised on a

medium complemented with 10^6 CFU of *L plantarum* and contaminated with Delfin 10X

(pink) or not (purple) compared to control pupae reared on control medium (green) or

medium contaminated with Delfin 10X only (red). (**C**) *Btk* load in the midgut of L3 larvae

reared on a medium contaminated with Delfin 10X and complemented (pink) or not (red) with

10^6 CFU of *L plantarum.* (**D**) Commensal bacterial load in the midgut of L3 larvae of our

Drosophila Canton S strain reared on our conventional medium (left bar) or reared in axenic

condition (right bar). The *Commensalibacter intestini* and *Acetobacter pomorum* grow on

mannitol plate (orange), *L plantarum* and *L brevis* grow on MRS plate (grey), *Enterococcus*

faecalis grows on BHI plate (green) and *Escherichia coli and Staphylococcus aureus* grow on

LB plate (blue). (**E**) Pupation curve of axenic larvae raised on a medium contaminated with

Delfin 10X (dark blue) or not (light blue, control Ax) compared to larvae conventionally

reared (CR, green). (**F**) Pupae volume derived from conventionally reared larvae (green) or

from axenic larvae raised on a sterile medium contaminated with Delfin 10X (dark blue) or

not (light blue).

1030

Figure 5: *Btk* bioinsecticides impacts on gut integrity

L3 larvae raised on a control medium (**A-B'**) or a medium contaminated with $Btk^{\Delta Cry}$ 10X (**C-**

D'), Delfin 10X (**E-F**), purified Crystals 10X (**G-H'**), naphthalene 10X (**I-J'**). (**A-J'**)

Labelling of lipid droplets with ORO (red, A, C, E, G and I) or with Bodipy (green, B, B', D,

D', F, F', H, H', J and J'). DAPI staining in blue marks the nuclei. Midguts are oriented

anterior to the left. Red brackets delineate the acidic region of the midgut. (B',D', F', H' and J')

are x/z section with the apical pole of the epithelium facing the lumen. (**K**) Stored (TAG,

upper panel) and circulating (DAG, lower panel) lipid dosages in control larvae (green bars)

and in larvae raised on Delfin 10X contaminated medium (red bars). (**L-N**)

myo1A>Casp::GFP larvae raised on control a medium (L) or on a medium contaminated with

$Btk^{\Delta Cry}$ 10X (M) or Delfin 10X (N). Arrows point enterocytes in which Caspase 3 is activated.

(**O-P**) L3 larvae raised on a medium complemented with Dextran^{FITC} (O) or with Dextran^{FITC}

and contaminated with *P. entomophila* (P), with $Btk^{\Delta Cry}$ 10X (Q) or Delfin 10X (R).

1044

Figure 6: Larval midgut regeneration upon aggression

(**A-C'**) Immunolabelling of *dlg::GFP* L3 midgut from larvae raised on control medium (A

and A'), on medium contaminated with $Btk^{\Delta Cry}$ 10X (B and B') or Delfin 10X (C and C').

Midguts were stained with the adherens junction marker Armadillo (Arm, purple) that

strongly delineates the Adult Midgut Precursors (AMP) (A, B and C) and marks the

basolateral compartment of the enterocytes (A', B' and C'). DAPI (blue) marks the nuclei and

Phalloidin (red) labels the visceral muscle surrounding the intestine and the apical pole of the

enterocytes facing the lumen (in A', B' and C'). Dlg::GFP (green) marks the sub apical

compartment of the enterocytes. (**D-H**) L3 midgut from larvae raised on control medium

(green bars), on medium contaminated with $Btk^{\Delta Cry}$ 10X (dark blue bars) or Delfin 10X (red

bars). (**D**) Number of AMPs per islets, (**E**) number of dividing AMPs per intestine (mitotic

AMPs are labeled by the Phospho-Histone 3 (PH3) immunostaining), (**F**) Number of AMP

islets (an islet contains more than one diploid cell) per unit of surface, (**G**) enterocyte area and

(**H**) epithelium thickness in the posterior midgut (3 independent measures were taken by

image capture). (**I** and enlargement in **I'**) *dlg::GFP* L3 midgut from larvae raised on medium

contaminated with Delfin 10X and stained for Arm (purple) and DAPI (blue). Arrows point

the cells with intermediate size (smallest than enterocytes but biggest than AMPs). (**J**)

Assessment of enterocyte ploidy in posterior L3 midgut from larvae raised on control

medium, on medium contaminated with $Btk^{\Delta Cry}$ 10X or Delfin 10X (C corresponding to

chromatin copy number). (**K-M**) Lineage tracing of AMP progeny in the *w; esg-Gal4, UAS-*

GFP/+; UAS-H2B::RFP, tub-Gal80^{ts}/+ (esg REDDM) L3 midgut from larvae raised on

control medium (K), on medium contaminated with $Btk^{\Delta Cry}$ 10X (L) or Delfin 10X (M). DAPI

(blue) marks the nuclei. **=P<0.01, ***=P<0.001 compared with controls.

1068

A Pupation curve

152< n < 262

- Control
- Delfin 1X
- Delfin 10X
- Dipel 10X
- ABTS-351 10X

B Pupation curve

143< n < 263

- Control
- $Btk^{\Delta Cry}$ 1X
- $Btk^{\Delta Cry}$ 10X
- Crystals 10X
- Naphthalene 10X
- Delfin 10X

C Pupae volume

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6