Radiation of nitrogen-metabolizing enzymes across the tree of life tracks environmental transitions in Earth history

Chris Parsons[1,2], Eva Stüeken[3,4], Caleb Rosen[1], Katherine Mateos[1] and Rika Anderson[1,4]

[1] *Carleton College, Northfield, MN, USA*
[2] *Massachusetts Institute of Technology, Cambridge, MA, USA*
[3] *University of St. Andrews, St. Andrews, Scotland, UK*
[4] *NASA NExSS Virtual Planetary Laboratory, University of Washington, Box 351580, Seattle, USA*

Abstract

Nitrogen is an essential element to life and exerts a strong control on global biological productivity. The rise and spread of nitrogen-utilizing microbial metabolisms profoundly shaped the biosphere on the early Earth. Here we reconciled gene and species trees to identify birth and horizontal gene transfer events for key nitrogen-cycling genes, dated with a time-calibrated tree of life, in order to examine the timing of the proliferation of these metabolisms across the tree of life. Our results provide new insights into the evolution of the early nitrogen cycle that expand on geochemical reconstructions. We observed widespread horizontal gene transfer of molybdenum-based nitrogenase back to the early Archean, minor horizontal transfer of genes for nitrate reduction in the Archean, and an increase in the proliferation of genes metabolizing nitrite around the time of the Mesoproterozoic (~1.5 Ga). The latter coincides with recent geochemical evidence for a mid-Proterozoic rise in oxygen levels. Our findings provide important constraints for understanding the evolution of the nitrogen cycle over time and provide insights into the bioavailability of various nitrogen sources in the early Earth with possible implications for the emergence of eukaryotic life.

Keywords: nitrogen cycle; horizontal gene transfer; geobiology; microbial evolution; nitrogenase

Introduction

Nitrogen is a critical element to life on Earth, important both for its role in biological molecule synthesis as well as in redox reactions for microbial energy metabolism. It is often a limiting nutrient in marine and terrestrial environments and may therefore have had a significant influence on the evolutionary trajectory of the biosphere over Earth's history. The nitrogen cycle is largely controlled by a variety of microorganisms that enzymatically catalyze the reduction and oxidation of nitrogen at various redox states (Figure 1). Reconstructing the genetic proliferation of these enzymes across the tree of life through horizontal gene transfer can therefore provide novel insights into the evolution of the biosphere and its productivity over time.

Nitrogen enters the biosphere through nitrogen fixation, the process of reducing atmospheric nitrogen (N_2) to ammonium (NH_4^+). This step is catalyzed by the nitrogenase enzyme, of which there are three varieties, distinguished by the metals in their associated active site cofactors: Nif (Fe-Mo), Vnf (Fe-V), and Anf (Fe-Fe). The ability to fix nitrogen is spread across a wide range of archaeal and bacterial lineages, but does not occur in eukaryotes (Dos Santos *et al.*, 2012; Gaby & Buckley, 2014). Importantly, nitrogenase is strongly inhibited by oxygen, forcing nitrogen fixers to develop a means to reduce their intracellular oxygen concentrations or to confine themselves to suboxic environments (Gallon, 1981). Ammonium produced from nitrogen fixation or ammonification (decomposition of biomass) can then either be converted to organic forms of fixed nitrogen, such as amino acids, or, in the presence of oxygen, oxidized to nitrite (NO_2^-) or nitrate (NO_3^-) through the chemoautotrophic nitrification pathway. The resulting nitrate can subsequently be used in the denitrification or DNRA (dissimilatory nitrate reduction to ammonium) pathways as an electron acceptor by microbial heterotrophs in conjunction with the oxidation of organic carbon compounds. Denitrification is a four-step process in which nitrate is reduced stepwise to nitrite, nitric oxide (NO), nitrous oxide (N_2O) and dinitrogen (N_2). Both N_2O and N_2 are gaseous and can escape into the atmosphere. Unlike denitrification, DNRA retains fixed nitrogen as ammonium and may therefore have been a critical metabolism in nutrient-starved ecosystems. Denitrification is often mediated by facultative anaerobes as well as ammonia-oxidizing archaea that use nitrate as an electron acceptor when oxygen levels are low; consequently, denitrification rates are highest in suboxic conditions, including swamps and oxygen minimum zones (Canfield *et al.*, 2005; Löscher *et al.*,

2012; Voss *et al.*, 2013). DNRA, while less understood, has been shown to be a major nitrate sink in a variety of aquatic systems, especially warm intertidal zones (Giblin *et al.*, 2013) and it may be dominant under ferruginous conditions, as suggested by modern analogue studies (Michiels *et al.*, 2017).

Given the importance of nitrogen as a building block of life, as an energy source for microbes and the most abundant element in the Earth's atmosphere, better constraining the evolutionary history of the nitrogen cycle is important for understanding, among other things, variation in global primary productivity and atmospheric pressure over time. The bioavailability and cycling of important limiting nutrients through Earth's history, including nitrogen, would have been important for biological productivity and the rise of early eukaryotic algae (Anbar & Knoll, 2002; Brocks *et al.*, 2017; Isson *et al.*, 2018). The relative abundance of nitrogenous gases in the atmosphere could have had important implications for planetary climate and atmospheric pressure: potential changes in atmospheric pressure may have resulted from biological N_2 drawdown (Som *et al.*, 2016), and the greenhouse gas nitrous oxide (N_2O), produced as part of the nitrogen cycle via nitrification/denitrification, may have contributed to planetary warming during the faint young Sun (Buick, 2007; Roberson *et al.*, 2011; Stanton *et al.*, 2018). Finally, one of the most important questions for the early evolution of life is understanding when fixed nitrogen first became widely available, and through what means. Experimental data suggest that fixed nitrogen can be produced during lightning reactions and under hydrothermal conditions (e.g., Brandes *et al.*, 1998; Navarro-González *et al.*, 2001), and either or both of these sources were likely pivotal for the origin of life. However, the invention of biological N_2 fixation would have made Earth's biosphere less dependent on abiotic reactions and likely spurred primary productivity.

The question of how these metabolisms unfolded over Earth's history has previously been addressed with geochemical approaches (Stüeken *et al.*, 2016a), reconstructions of microbial metabolism based on metal availability (Moore *et al.*, 2017) or with phylogenetics studies targeting the origin and evolution of specific enzymes, which yield conservative estimates for the approximate timing of an enzyme's origin (Raymond *et al.*, 2004; Boyd *et al.*, 2011b, 2011a; Boyd & Peters, 2013; Garcia *et al.*, 2020). However, geochemical data are too insensitive to distinguish between particular enzymatic pathways, and the timing of the origin of an enzyme may not be equivalent to the time when it gained ecological significance on a large

scale. Additional constraints are therefore needed to address key questions about the dynamics of the nitrogen cycle on the early Earth and its evolution over time. The proliferation of sequencing data over the past decade has made vast amounts of genomic data available, which can provide novel insights into the evolution of the nitrogen cycle by providing information about the evolution of nitrogen cycling genes over time.

As a new approach to these questions, we track the timing of birth, speciation, duplication, loss, and horizontal gene transfer events for genes involved in each step of the nitrogen cycle by reconciling and comparing the topology of gene trees and a time-calibrated species tree. We place particular focus on the acquisition of new genes via horizontal gene transfer (HGT), which is common in microbial lineages and is a crucial evolutionary mechanism by which a clade of organisms can develop new and useful phenotypes without the evolutionary cost associated with independently evolving genes (Beiko *et al.*, 2005; Gogarten & Townsend, 2005). The acquisition of new functional genes can allow clades of microbes to invade new ecological niches. The rate of horizontal gene transfer itself is likely to be related to variables like cell density, co-localization of donor and recipient, cell diversity, and the types of organisms involved (Gogarten & Townsend, 2005). Studies of gene gain and loss indicate both HGT and gene loss have been crucial forces molding the evolution of bacterial and archaeal lineages (Koonin *et al.*, 2001; Mirkin *et al.*, 2003), including for crucial genes in the nitrogen cycle (Stolz & Basu, 2002; Kechris *et al.*, 2006; Jones *et al.*, 2008). Although some have argued for a neutral theory of gene transfer in which horizontally acquired genes are not adaptive (Gogarten & Townsend, 2005; Andreani *et al.*, 2017), studies indicating that horizontally transferred genes perform crucial cellular functions and enable adaptation to specific ecological niches suggest that horizontally acquired genes are generally adaptive (Daubin & Ochman, 2004; Coleman & Chisholm, 2010; Burke *et al.*, 2011; Popa *et al.*, 2011; Polz *et al.*, 2013; McInerney *et al.*, 2017; Moulana *et al.*, 2020). Moreover, genes providing selective advantages are more likely to be retained in the genome than genes acquired due to neutral transfer, which are more likely to be purged, thus strengthening the signal of adaptive HGT events in the genomic record. Therefore, any observations of a rise in the relative number of successful horizontal gene transfers for a given gene are likely to give an indication of the relative availability or metabolic importance of a given substrate for that gene. As such, studying the history of HGT of specific genes can provide insights into the points at which possession of such genes provided substantial selective

advantages, which can then be used as a metric for when a specific metabolism became feasible or energetically favorable, or when the substrates of specific enzymes became relatively abundant. A crucial caveat to this method, however, is that it cannot infer relative abundance or population sizes of the organisms carrying these genes. Thus, if a specific strain carrying a nitrogen metabolizing gene grows in abundance with no transfer of genes to other lineages, our method would observe low rates of transfer for genes involved in that metabolism. Additionally, a poor phylogeny or poor phylogenetic signal would make it difficult to precisely identify such horizontal gene transfer events. Therefore, results should be interpreted with such possible scenarios in mind.

Here, we track birth, speciation, duplication, loss, and HGT of genes in the nitrogen cycle over time by comparing the phylogenies for specific nitrogen-metabolizing genes (here called "gene trees") to a phylogeny based on the concatenation of highly-conserved ribosomal proteins (here called a "species tree"). We reconstructed a tree of life based on 308 genomes from isolates and metagenome-assembled genomes (MAGs) (see Methods). Any discrepancies between the species tree and a given gene tree can therefore be attributed to HGT, gene loss, and/or gene duplication. We attribute rough dates to these events by anchoring the species tree to known fossil calibration dates and using molecular clocks to model rates of evolution and estimate divergence times.

Materials and Methods

Genome Selection and Compilation

The construction of both the gene and species trees for this study was based upon the manual curation of a genome database containing 308 genomes (including 254 bacterial and archaeal genomes) that served as the basis for the species tree and was subsequently searched to find genes related to nitrogen metabolism. Assembled genomes were downloaded from ggKBase (Hug *et al.*, 2016) and the NCBI assembly database (Kitts *et al.*, 2016). Additionally, 6 genomes were collected from a recent study identifying novel nitrogen fixers (Delmont *et al.*, 2018). In constructing the tree, we included at least one genome from each bacterial or archaeal phylum represented in the most recent comprehensive tree of life (Hug *et al.*, 2016) in order to create a tree fully representative of our current understanding of microbial diversity. It also includes a set of genomes associated with a database of *nifH* genes (Gaby & Buckley, 2014). Some eukaryotic

genes were included for construction of the tree, but for this study we focused only on archaeal and bacterial genomes for identification of nitrogen cycling genes. Relative to archaea and bacteria, eukaryotes play a more minor role in the nitrogen cycle—while some species of fungi and protists reduce nitrate or nitrite to more reduced forms of nitrogen, there are no known eukaryotes that mediate nitrogen fixation, nitrification, DNRA, or anammox (Stein & Klotz, 2016).

Gene Isolation

To create the species tree, all bacterial and archaeal genomes in our database were mined for single copy ribosomal protein sequences L2, L3, L4, L5, L6, L14, L15, L16, L18, L22, L24, S3, S8, S10, S17, and S19 using Phylosift (Darling *et al.*, 2014), with the isolate and best hit command line flags. These sixteen ribosomal proteins represent the same proteins used to create a recent comprehensive tree of life (Hug *et al.*, 2016) and, in the case of eukaryotes, ribosomal sequences were directly drawn from their dataset. All genomes included in the dataset contained fewer than 50% gaps in the alignment.

Curated gene database queries for each of the nitrogen-cycling genes we investigated were generated based on KEGG orthologies (Ogata *et al.*, 1999) and downloaded from the UniProt database (The Uniprot Consortium, 2017). Nitrogen-cycling amino acid sequences for the gene trees were identified by conducting BLASTP (Altschul *et al.*, 1990) searches of the open reading frames (ORFs) of every genome from the collection of 254 genomes. All ORFs were identified using Prodigal (Hyatt *et al.*, 2010). The maximum e-value cutoff for BLAST hits was 10^{-12} and matches were excluded if the length of the local alignment was less than 50% of the length of the query sequence. BLAST results were compared with results from AnnoTree (Mendler *et al.*, 2019) to verify gene distributions. Given that *nifK* and *nifD* have a shared evolutionary history (Fani *et al.*, 2000), an e-value cutoff of 1e-30 was selected based on close examination of blastp hits in order to distinguish these subunits. For nitrogenase subunits *nifH*, *vnfD, vnfK, anfD, and anfK* we manually curated alignments by identifying key residues that were crucial for enzyme structure and function, as determined through literature searches and visualization using PyMol (Brigle *et al.*, 1987; Kaiser *et al.*, 2011; McGlynn *et al.*, 2012; Howard *et al.*, 2013; Keable *et al.*, 2018). Jalview (Waterhouse *et al.*, 2009) was used for visualization of key residues in alignments. All genes that did not include the key residues were

removed from the alignment. Finally, to ensure that only the genes of interest were included in alignments and to verify annotations, all genes identified in the BLAST search were compared to the KEGG database using Kofam Koala (Aramaki *et al.*, 2019), which assigns KEGG Ortholog numbers to each gene by using a homology search against a database of profile hidden Markov models. Only genes verified to be the gene of interest according to Kofam Koala were retained for downstream analysis. It is important to note that gene identification is necessarily limited by the search tools and databases used for annotation, and the methods used here were chosen to be conservative so as to ensure the removal of non-target genes from the analysis.

Tree Construction

Alignments for the gene trees were created using MUSCLE (Edgar, 2004), and phylogenies were constructed from the resulting alignments using RAxML with 100 rapid bootstraps (Stamatakis, 2014) using an LG substitution matrix, with the GAMMA rate heterogeneity model. Alignments for the species trees were made using the Phylosift pipeline (Darling *et al.*, 2014) and curated to only include the target ribosomal proteins listed above. The CAT model, which calculates site-specific evolutionary rates, was used with an LG substitution matrix to construct the species tree based on reference marker genes (the species tree). The root for the species tree was placed in the Bacterial domain (Fournier & Gogarten, 2010).

Chronogram Construction

The species tree was converted into four species chronograms using PhyloBayes (Lartillot *et al.*, 2009) in order to test the sensitivity of our results to variation in Phylobayes parameters. The root prior was set to 3800 Ma, with standard deviation set to 200 in accordance with previous studies (Magnabosco *et al.*, 2018). Two sets of calibration points were used, one liberal (which represents the earliest date for which there is any evidence of a given event based on the current scientific literature) and one conservative (which represents the earliest date for which there is the most consensus for a given event based on the current scientific literature), to test the sensitivity of methodology (Table 1). Similarly, we tested both a CIR and a lognormal clock model (Lepage *et al.*, 2007) to compare the effects of clock model type on the results. Two clocks were run in parallel for each set of parameters, so that the two concurrent runs could be compared to one another as a test of convergence. Simultaneous runs were run for 40,000 cycles

and generally converged after roughly 10,000 cycles. Chronograms were generated with burn-ins of around 2,500 cycles. Chronograms were visualized using the phytools package in R (Revell, 2012). In order to test for the influence of the priors, we generated additional chronograms in the absence of sequence data using the -prior flag in Phylobayes. These chronograms displayed substantially different node timings, demonstrating that the priors did not overly influence the inferred dates and the sequence data informed the chronogram. Results from molecular clock analyses should be interpreted with caution, given the limitations associated with these analyses, including but not limited to changing generation times, the influence of natural selection, and variation in mutational rates across species (Ayala, 1999; Schwartz & Maresca, 2006; Bromham *et al.*, 2018). Here, we attempted to ameliorate some of these challenges by using methods allowing for variation of rates between and across lineages, and by comparing results produced by both liberal and conservative fossil constraints. Finally, our goal with this analysis was not to pinpoint exact dates for many of the transitions discussed here, but rather to compare the relative timing on broad evolutionary scales.

Gene Tree and Species Chronogram Reconciliation

Gene trees were reconciled with species chronograms using the Analyzer of Gene and Species Trees (AnGST) (David & Alm, 2011). AnGST compares the topology of the gene tree with that of the species tree, rather than rely solely on presence and absence patterns, in order to identify gene birth, transfer, duplication, and loss events. Event penalties were set to hgt: 3, dup: 2, los: 1, and spc: 0. Ultrametric was set to True in order to constrain events temporally. Each run was conducted with 100 gene tree bootstraps in order to increase accuracy (David & Alm, 2011). Individual event timings were defined as the midpoint of the temporal region during which a given event could occur.

Results

Generation of a species tree and fossil-calibrated chronogram

We compared all gene trees to a species tree constructed from an alignment of concatenated sequences of 16 single-copy universal proteins from 308 organisms (Figure 2; genome list available as Supplemental Data). Monophyly was preserved for most major phyla, with three exceptions: (i) Tenericutes is contained within Firmicutes, (ii) the PVC superphylum

contains Omnitrophica, and (iii) Lentisphaerae is nested within Verrucomicrobia. Additionally, in contrast to a recently published comprehensive tree of life (Hug *et al.*, 2016), our tree does not place the recently-discovered Candidate Phyla Radiation (CPR) bacteria as the deepest-rooted bacterial clade; however, the CPR is placed as a sister group to the Cyanobacteria and Melainabacteria, which is consistent with that phylogeny (Hug *et al.*, 2016). For the purposes of this study, placement of these groups should not greatly affect our results due to the high number of duplication/loss/transfer events we inferred overall across all groups. The species tree used for this analysis is a three-domain tree, in contrast to recent studies which have shown the addition of the Asgard Archaea to the tree of life to cause Eukaryotes to group within Archaea (Zaremba-Niedzwiedzka *et al.*, 2017). However, we are agnostic as to the placement of the Asgard superphylum and the eukaryotes on the tree of life, as this was not our focus; the relationship of the three domains to one another should not substantially affect the results shown here due to the relative infrequency of inter-domain HGT relative to interdomain HGT, and the exclusion of eukaryotic nitrogen-cycling genes.

We constructed four different chronograms from our species tree using two different clock models (CIR and lognormal) as well as liberal (representing the earliest date for which there is any evidence of a given event based on the current scientific literature) and conservative (representing the earliest date for which there is the most consensus for a given event based on the current scientific literature) fossil calibration points (see Methods). As expected, the use of liberal calibration points pushed the timing of most divergence events earlier than those resulting from conservative calibration points (Supplementary Figures 1-4; chronograms and Newick files available in Supplemental Data). The only major difference between the chronograms constructed by the CIR and lognormal models is that the lognormal chronograms, especially the one based on conservative calibration points, estimated a longer branch length between LUCA and bacterial diversification than that estimated by the CIR clock model. The timing of LUCA varied slightly according to clock model, and substantially (~500 Ma difference) with liberal versus conservative calibration points. The main results presented here make use of the CIR conservative chronogram (Figure 3), but the results derived from liberal time points and from the lognormal clock model are included as Supplementary Figures 6, 7 and 8, all of which demonstrate the same overall results.

Identification of duplication, speciation, loss, and horizontal gene transfer events for nitrogen-cycling genes

We identified 18 different nitrogen-cycling genes for analysis of horizontal gene transfer, speciation, duplication, and loss events (Table 2). Very few genes from the *hao* and *amo* gene families were identified in these genomes or passed our stringent filtration tests, and thus they did not yield sufficient data for robust conclusions, as this methodology is best suited to genes that have many representatives from a diverse suite of taxa, and so we did not include them in further analyses. We generated maximum likelihood gene trees for each of the remaining genes (Supplementary Data) and compared these gene trees to the fossil-calibrated chronogram in order to identify and infer the timing of birth, duplication, loss, and horizontal gene transfer events for each of these genes. The number of HGT and speciation events inferred for each gene in this study was relatively high compared to the number of inferred duplication and loss events (Table 3), and inferred loss events generally skewed younger than inferred horizontal gene transfer and speciation events. The oldest genes, with horizontal gene transfer and speciation events dating back to before 3 Ga, were the iron-molybdenum nitrogenases (*nifH/nifD/nifK*). Our results indicate that these nitrogenases originated during the Archean era and began to spread across the tree of life via horizontal gene transfer fairly early in Earth history as well (Fig. 3). In contrast, the alternative nitrogenases *anf* and *vnf* were inferred to have arisen much later, but very few *anf* and *vnf* genes were identified among our sample set and therefore these results should be treated with caution (Tables 2 and 3, Supplementary Figure 5). The denitrifying genes *norB, nosZ, nirK* and *nirS* were inferred to have arisen much later in Earth history, approximately 1.5 Ga (Table 4), and began to proliferate across the tree of life shortly thereafter (Fig. 3). These genes encode enzymes that catalyze denitrification processes from nitrite to nitric oxide, nitrous oxide, and dinitrogen gas. The genes *nrfA* and *nirB*, which are involved in the DNRA process, arose fairly early at approximately 2.7-3.0 Ga, with a few horizontal gene transfer and speciation events occurring during the Archean era, but did not proliferate widely across the tree of life until later at approximately ~1.5 Ga. Genes involved in nitrite or nitrate oxidation and reduction, including *narG*, *nasA*, *nxrA* and *napA* were inferred to have arisen relatively early (approximately 3.0 Ga) (Table 4) and we inferred a few speciation and horizontal gene transfer events for these genes between 2-3 Ga, but we did not observe a rise in HGT events for these genes until much later, at approximately 1.5 Ga (Figure 3).

Taken together, we observed that genes related to the fixation of nitrogen from dinitrogen gas to ammonium arose early and proliferated across the tree of life relatively quickly, while genes related to the reduction and oxidation of nitrate arose early, but did not begin to proliferate across the tree of life until later. Genes related to denitrification, particularly downstream from nitrite, arose much later (Figure 3). Our overall results regarding the relative timing of the birth and HGT events for nitrogen-cycling genes were robust to the use of liberal vs. conservative time points as well as the use of the CIR vs. lognormal clock models (Supplementary Figures 6, 7, and 8).

Discussion

On the whole, our results provide support for geochemical data indicating that biological nitrogen fixation was an important source of fixed nitrogen in the Archaean. These results also suggest that local sources of nitrate may have been exploited by denitrifying microbes, while denitrifiers using nitrite or downstream products would not have proliferated until much later in Earth history, until the mid-Proterozoic, well after the Great Oxidation Event. This has important implications for the evolution of the nitrogen cycle and its relationship to the redox state of the Earth.

Nitrogen Fixation

Our data indicate that nitrogen fixation through the use of molybdenum nitrogenase (Nif) is an ancient process, arising by approximately 3.1 Ga at the latest. Our results are consistent with previous work from geochemical analyses suggesting that nitrogen fixation must have arisen early in order to support an expanding biosphere (Stüeken *et al.*, 2016a; Zerkle & Mikhail, 2017). Though some analyses have suggested that nitrogenase arose relatively late (approx. 2.2-1.5 Ga) (Boyd *et al.*, 2011a) and that nitrogen fixers did not become abundant in global oceans until the Neoproterozoic (Sánchez-Baracaldo *et al.*, 2014), our results are consistent with phylogenetics studies suggesting that molybdenum nitrogenase arose early in the evolution of life on earth (Raymond *et al.*, 2004; Weiss *et al.*, 2016; Garcia *et al.*, 2020). The identification of horizontal gene transfer events for nitrogenase subunits during the Archean (Figure 3) suggest that this metabolism may have been abundant and beneficial enough during this time period to have been successfully transferred and retained in microbial genomes. This interpretation is

consistent with recent nitrogen isotope evidence for biological nitrogen fixation back to at least 3.2 Ga (Stüeken *et al.*, 2015).

Our results support the argument that abiotic sources of fixed nitrogen were unlikely to be significant enough to sustain the early biosphere. It has been proposed that early life received its nitrogen from the lightning-catalyzed reaction between N_2 and CO_2 as a source of NO_x, and that a steady reduction in atmospheric CO_2 levels reduced this flux, eventually leading to a nitrogen crisis around 2.2 Ga that would have favored the rise and spread of biological nitrogen fixation (Navarro-González *et al.*, 2001). In contrast, several studies have concluded that abiotic nitrogen fixation on the early Earth would produce 50- to 5000-fold lower rates of fixed nitrogen than what is contained in the modern ocean, making nitrogen highly limiting to the early biosphere, even at higher CO_2 levels (Canfield *et al.*, 2010). Our data suggest that biological nitrogen fixation arose and proliferated early and that acquisition of the nitrogenase enzyme provided enough of a selective advantage to be successfully transferred across lineages during the early Archaean, supporting the notion that fixed nitrogen was not widely available.

Moreover, if the radiation of biological N_2 uptake dates back to the early Archean, it may have had a significant impact on atmospheric pressure. Today, nitrogen makes up approximately 78% of the Earth's modern atmosphere by volume, predominantly as N_2 gas. The atmosphere of the early Earth was probably also N_2-rich, but while some models suggest a 2-4 times larger atmospheric N_2 reservoir in the Archean (Johnson & Goldblatt, 2018), proxy evidence ranges from near modern values (Marty *et al.*, 2013; Avice *et al.*, 2018) to less than half of today's reservoir (Som *et al.*, 2016). If atmospheric N_2 pressure changed over time, it is likely that biological activity would have played a major role in driving these changes by enhancing nitrogen burial in sediments and by accelerating oxidative weathering of nitrogen from continental crust (Stüeken *et al.*, 2016b; Zerkle & Mikhail, 2017). If nitrogen was fixed by nitrogenase early in the Archaean, as indicated by our results, then biological nitrogen burial began long before the onset of oxidative weathering in the Neoarchean (Stüeken *et al.*, 2012) which makes it possible that atmospheric N_2 decreased over the course of the Archean. Although further paleobarometric proxies are needed to verify this trend, our results provide an important anchor point for the onset of biological N_2 drawdown.

Lastly, though we identified very few iron and vanadium nitrogenases (*anf* and *vnf*, respectively) in our datasets, our results tentatively suggest that these nitrogenases arose later

than the iron-molybdenum nitrogenases (*nif*), which supports conclusions from other phylogenetics studies (Garcia *et al.*, 2020). The early rise of Mo-dependent nitrogenase would have required a source of Mo to act as a cofactor for Mo-containing nitrogenases, implying that nanomolar levels of dissolved Mo (Scott *et al.*, 2008; Reinhard *et al.*, 2013), presumably derived from anoxic weathering or hydrothermal sources of molybdenum to the early ocean, were sufficient for the development of Nif and the proliferation of nitrogen fixers. Oxidative weathering was therefore evidently not required for the onset of biological nitrogen fixation (c.f. Boyd *et al.*, 2011b). We speculate that the evolutionary pressure for the diversification of alternative nitrogenases arose in the Neoproterozoic with the rise of eukaryotic algae (Brocks *et al.*, 2017), which may have substantially increased the nitrogen demand in the global ocean. Additional niches for alternative nitrogenases may have arisen with the rise of land plants and associated soil bacteria in the Paleozoic, as recent work has documented the presence of V-based nitrogen fixation in cyanobacterial symbionts of lichens (Darnajoux *et al.*, 2017).

Nitrogen Cycling Prior to the Oxygenation of the Oceans and Atmosphere

Although our results suggest that nitrogen fixation was an important process on the early Earth, microbial metabolisms making use of more oxidized forms of nitrogen, particularly through the process of denitrification downstream from nitrite, do not appear to have arisen and spread across the tree of life until much later. Some early steps in the nitrogen cycle, such as nitrate reduction to nitrite via *nasA*, *narG* or *napA*, appear to have arisen relatively early (in the late Archean or early Proterozoic), though we did not identify many horizontal gene transfer events early in the evolutionary history of these genes, and their proliferation across the tree of life only began in earnest much later, approximately 1.5 Ga. In the anoxic ocean that existed before the development of oxygenic photosynthesis, the most widespread form of dissolved inorganic nitrogen was ammonium (Stüeken *et al.*, 2016a), and as noted above, abiotic sources of fixed nitrogen were probably too minor to sustain the entire biosphere. However, it is likely that a small flux of nitrogen oxides from lightning and volcanism existed (Navarro-González *et al.*, 2001; Mather *et al.*, 2004) and maintained populations of nitrate-reducing denitrifiers. Denitrification is a highly energy-yielding metabolism (Schoepp-Cothenet *et al.*, 2013), and therefore even a small source flux of nitrate to the early ocean would probably have been exploited.

Alternatively, it is possible that the nitrate-reducing genes appeared in locally oxygenated regions of the Archean surface ocean, which have been inferred from isotopic studies back to 3.0 Ga (Olson *et al.*, 2013; Planavsky *et al.*, 2014). In such oxygen oases, oxygen concentrations may have reached micromolar concentrations (Olson *et al.*, 2013), which is high enough for the complete oxidation of ammonium to nitrite and nitrate (Lipschultz *et al.*, 1990). Any nitrate leaving these pockets could have been reduced to nitrite by a variety of microorganisms and could thus have contributed to the observed genetic evidence for nitrate reducers at this time. However, these metabolisms were ecologically not important enough to leave a significant mark in the geochemical record (Ossa Ossa *et al.*, 2019) .

Our results suggest that genes for nitrite-metabolizing enzymes were not frequently transferred during the early Archean, which may imply that nitrite-metabolizing genes were not widespread in Archean ecosystems. It is possible that only specific taxa were responsible for the conversion of nitrite into nitrate or ammonium. Alternatively, the conversion of nitrite to ammonium may have happened abiotically. Fe^{2+}, which is thought to have been abundant in the early anoxic ocean, may have readily reduced nitrite to ammonium or nitrogen gas (Summers & Chang, 1993; Canfield *et al.*, 2010). These results may also indicate that nitrite levels during this time period were too low to cause substantial proliferation of genes for enzymes which take nitrite as a substrate, while nitrate and ammonium concentration were high enough for selection to favor biological metabolism of these molecules by diverse microorganisms.

Effects of Increased Oxygen Levels on the Nitrogen Cycle

Multiple lines of evidence point to increasing oxygenation of surface environments from 2.75 Ga onwards, culminating in the Great Oxidation Event (GOE) at 2.3 Ga (Lyons *et al.*, 2014). Although our data suggest that the genes *nrf* and *nirB,* which reduce a variety of oxidized forms of nitrogen to ammonium through the DNRA pathway, arose much earlier, we observed that the relative number of HGT events for this gene began to rise around the time of the GOE. Our data allow for the possibility that this nitrogen reduction pathway prevailed through the early oxygenation of the ocean, before the more complex denitrification pathway had evolved. Unlike denitrification to N_2 gas, DNRA retains the reduced nitrogen product in the system as aqueous NH_4^+, which may have prevented a potential 'nitrogen crisis' during the wake of the GOE (Falkowski & Godfrey, 2008). The isotopic fingerprint of DNRA can so far not be distinguished

from denitrification in the geochemical record, and thus our results provide the first indication that this metabolism may have been of greater ecological importance than previously proposed. Importantly, geochemical evidence of biological nitrate reduction (either through denitrification or DNRA) dates back to the Neoarchean around 2.7 Ga and becomes more widespread after the GOE around 2.3 Ga (Garvin *et al.*, 2009; Zerkle *et al.*, 2017; Kipp *et al.*, 2018), i.e. long before our observed radiation in denitrification genes. This observation may suggest that these early nitrate-reducing ecosystems were primarily performing DNRA, consistent with iron-rich conditions in the Archean and Paleoproterozoic ocean (Michiels *et al.*, 2017).

After approximately 1.5 Ga, the relative number of HGT events for enzymes involved in the modern denitrification pathway also began to rise. Additionally, our data suggest that the gene *nxr*, which oxidizes nitrite into nitrate, arose quite early (approx. 3 Ga) but did not begin to spread across the tree of life until ~1.5 Ga. One possibility is that increased copper availability facilitated the spread of nitrification/denitrification, which require enzymes that rely on copper cofactors (Moore *et al.*, 2017). However, geochemical evidence of increasing copper concentrations in seawater at 1.5 Ga is so far lacking. Another possibility is that the expansion of these metabolisms is linked to a Mesoproterozoic rise in oxygen, which has recently been proposed in several geochemical studies (Cox *et al.*, 2016; Canfield *et al.*, 2018; Zhang *et al.*, 2018; Shang *et al.*, 2019) Higher oxygen levels would have led to an expansion of the marine nitrate reservoir, which may in turn have been critical for the rise of eukaryotes around this time (Anbar & Knoll, 2002; Knoll & Nowak, 2017).

The primary substrate for many of the enzymes that demonstrated this late rise in the number of HGT events is nitrite, suggesting that nitrite levels may have increased at this time, such that selection favored the spread of nitrite-metabolizing enzymes across the tree of life. Nitrite is thermodynamically unstable in oxic conditions, but it occurs transiently in chemoclines in the modern ocean because it is abundantly produced as an intermediate in nitrification and denitrification pathways (Wada & Hatton, 1971). A more vigorous aerobic nitrogen cycle in the Proterozoic may thus be linked to the establishment of a more permanent dynamic nitrite reservoir. Abiotic reduction of nitrite by Fe^{2+}, which was perhaps dominant during the Archean (see above), may have slowed down with the deepening of the chemocline after the GOE.

However, our results do not support the hypothesis of a N_2O-rich atmosphere in the mid-Proterozoic. The argument for a so-called "laughing gas greenhouse" hinges on the assumption

that widespread euxinia following the Great Oxidation Event would have reduced marine copper levels through sulfide stripping (Buick, 2007). It then follows that NosZ, the terminal enzyme in the denitrification pathway, which requires copper in its active site, would be too difficult to produce in such a copper-limiting environment, leading to a buildup of NorB-generated nitrous oxide. In contrast, our data identified a rise in the number of horizontal gene transfer events in the Mid-Proterozoic for *nosZ*, as well as for *nirK*, which codes for another denitrifying enzyme with copper in the active site, indicating that copper levels were high enough to maintain copper-containing enzymes. However, N_2O may still have been an important greenhouse gas at this time, because even in the presence of copper it is a common "leakage" product during both nitrification and denitrification (e.g. Stanton *et al.*, 2018), which were evidently widespread in the mid-Proterozoic ocean.

Several caveats must be kept in mind while interpreting these results. As mentioned above, the accuracy of molecular clocks is constrained by the models used to infer dates and the accuracy of the fossil-based time points used for clock calibration. We sought to minimize the influence of these parameters by using both liberal and conservative time points and by employing two different clock models, but nevertheless these limitations must be taken into consideration, and thus we emphasize the relative rather than the absolute timing of the events identified here. Moreover, the methods used to identify genes used in the analysis were designed to be conservative so as to reduce the possibility that spurious genes were included in the analysis, but these methods are inherently limited by the quality of the databases used for annotation. These conservative methods precluded the inclusion of *hao* and *amo,* which are involved in the oxidation of ammonia to nitrite and nitrogen gas, in our analysis. Others have observed that archaeal ammonia oxidizers most likely originated during the GOE and began to spread through the shallow ocean approximately 800 Mya (Ren *et al.*, 2019). As new genes and genomes continue to be sequenced and analyzed for gene function, the quality of such analyses will improve over time.

Conclusion

Our results show that biological nitrogen fixation appears to have arisen and proliferated early in the Archaean, and that genes in the denitrification pathway and genes related to the consumption of nitrite and its downstream products began to proliferate across the tree of life

following the oxygenation of Earth's atmosphere and oceans. Moreover, our results support the hypothesis that the molybdenum-based variety of nitrogenase diversified much earlier than alternative nitrogenases, which may only have become important with the increasing nitrogen demand of algae and land plants in the Neoproterozoic and Paleozoic, implying that molybdenum was not a limiting resource in the Archean ocean. Furthermore, our data provide the first indirect evidence for a small source of nitrate to the early ocean, possibly as a result of lightning and volcanism, as well as for nitrate reduction to ammonium in the Proterozoic, which may have helped overcome nitrogen limitation by retaining fixed nitrogen in the system. We cannot confirm the hypothesized suppression of N_2O-metabolizing enzymes in the mid-Proterozoic, but our results are consistent with vigorous nitrification and denitrification after the GOE, in particular from 1.5 Ga onwards, which may have led to leakage of N_2O into the atmosphere and the stabilization of global climate. The proliferation of key nitrogen-metabolizing genes across the tree of life at different points in Earth's redox history provides important insights into the evolution and radiation of microbial metabolisms on the early Earth in response to major environmental transitions and supports the notion that increasing nitrate levels in the mid-Proterozoic may have contributed to the rise of eukaryotic life.

Acknowledgements

We would like to thank Rou-Jia Sung for assistance with structural predictions, as well as Casey Bryce, Colin Goldblatt, Ben Johnson, John Baross, Victoria Meadows, Jaclyn Saunders, and Cara Magnabosco for support and useful discussions. This work was performed as part of NASA's Virtual Planetary Laboratory, supported by the NASA Astrobiology Program under grant 80NSSC18K0829 as part of the Nexus for Exoplanet System Science (NExSS) research coordination network. CWP received a Towsley Fellowship from Carleton College, and REA was also supported by a NASA Postdoctoral Fellowship from the NASA Astrobiology Institute.

References

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *Journal of Molecular Biology* **215**, 403–410.

Anbar AD, Knoll AH (2002) Proterozoic ocean chemistry and evolution: A bioinorganic bridge? *Science* **297**, 1137–1142.

Andreani NA, Hesse E, Vos M (2017) Prokaryote genome fluidity is dependent on effective population size. *The ISME Journal* **11**, 1719–1721.

Avice G, Marty B, Burgess R, Hofmann A, Philippot P, Zahnle K, Zakharov D (2018) Evolution of atmospheric xenon and other noble gases inferred from Archean to Paleoproterozoic rocks. *Geochimica et Cosmochimica Acta* **232**, 82–100.

Ayala FJ (1999) Molecular clock mirages. *BioEssays* **21**, 71–75.

Beiko RG, Harlow TJ, Ragan MA (2005) Highways of gene sharing in prokaryotes. *Proceedings of the National Academy of Sciences* **102**, 14332–14337.

Bekker A, Holland HD, Wang P-L, Rumble D, Stein HJ, Hannah JL, Coetzee LL, Beukes NJ (2004) Dating the rise of atmospheric oxygen. *Nature* **427**, 117–120.

Boyd ES, Anbar AD, Miller S, Hamilton TL, Lavin M, Peters JW (2011a) A late methanogen origin for molybdenum-dependent nitrogenase. *Geobiology* **9**, 221–32.

Boyd ES, Hamilton TL, Peters JW (2011b) An alternative path for the evolution of biological nitrogen fixation. *Frontiers in Microbiology* **2**, 1–11.

Boyd ES, Peters JW (2013) New insights into the evolutionary history of biological nitrogen fixation. *Frontiers in microbiology* **4**, 201.

Brigle KE, Weiss MC, Newton WE, Dean DR (1987) Products of the iron-molybdenum cofactor-specific biosynthetic genes, nifE and nifN, are structurally homologous to the products of the nitrogenase molybdenum-iron protein genes, nifD and nifK. *Journal of bacteriology* **169**, 1547–53.

Brocks JJ, Jarrett AJM, Sirantoine E, Hallmann C, Hoshino Y, Liyanage T (2017) The rise of algae in Cryogenian oceans and the emergence of animals. *Nature* **548**, 578–581.

Bromham L, Duchêne S, Hua X, Ritchie AM, Duchêne DA, Ho SYW (2018) Bayesian molecular dating: Opening up the black box. *Biological Reviews*.

Buick R (2007) Did the Proterozoic "Canfield Ocean" cause a laughing gas greenhouse?

Geobiology **5**, 97–100.

Burke C, Steinberg P, Rusch D, Kjelleberg S, Thomas T (2011) Bacterial community assembly based on functional genes rather than species. *Proceedings of the National Academy of Sciences of the United States of America* **108**, 14288–14293.

Canfield D, Kristensen E, Thamdrup B (2005) *Aquatic Geomicrobiology, Volume 48*.

Canfield DE, Glazer AN, Falkowski PG (2010) The Evolution and Future of Earth's Nitrogen Cycle. *Science* **330**.

Coleman ML, Chisholm SW (2010) Ecosystem-specific selection pressures revealed through comparative population genomics. *Proceedings of the National Academy of Sciences of the United States of America* **107**, 18634–18639.

Darling AE, Jospin G, Lowe E, Matsen FA, Bik HM, Eisen JA (2014) PhyloSift: phylogenetic analysis of genomes and metagenomes. *PeerJ* **2**, e243.

Darnajoux R, Zhang X, McRose DL, Miadlikowska J, Lutzoni F, Kraepiel AML, Bellenger J-P (2017) Biological nitrogen fixation by alternative nitrogenases in boreal cyanolichens: importance of molybdenum availability and implications for current biological nitrogen fixation estimates. *New Phytologist* **213**, 680–689.

Daubin V, Ochman H (2004) Bacterial Genomes as New Gene Homes: The Genealogy of ORFans in E. coli. *Genome Research* **14**, 1036–1042.

David LA, Alm EJ (2011) Rapid evolutionary innovation during an Archaean genetic expansion. *Nature* **469**, 93–6.

Delmont TO, Quince C, Shaiber A, Esen ÖC, Lee ST, Rappé MS, MacLellan SL, Lücker S, Eren AM (2018) Nitrogen-fixing populations of Planctomycetes and Proteobacteria are abundant in surface ocean metagenomes. *Nature Microbiology* **3**, 804–813.

Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research* **32**, 1792–1797.

Eigenbrode JL, Freeman KH (2006) Late Archean rise of aerobic microbial ecosystems. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 15759–15764.

Falkowski PG, Godfrey L V (2008) Electrons, life and the evolution of Earth's oxygen cycle. *Philosophical Transactions of the Royal Society B: Biological Sciences* **363**, 2705–2716.

Fournier GP, Gogarten JP (2010) Rooting the Ribosomal Tree of Life. *Molecular Biology and*

Evolution **27**, 1792–1801.

Gaby JC, Buckley DH (2014) A comprehensive aligned nifH gene database: a multipurpose tool for studies of nitrogen-fixing bacteria. *Database : the journal of biological databases and curation* **2014**, bau001.

Garcia AK, McShea H, Kolaczkowski B, Kaçar B (2019) Reconstructed ancient nitrogenases suggest Mo-specific ancestry. *bioRxiv* 714469.

Garvin J, Buick R, Anbar AD, Arnold GL, Kaufman AJ (2009) Isotopic evidence for an aerobic nitrogen cycle in the latest Archean. *Science* **323**, 1045–8.

Giblin A., Tobias CR, Song B, Weston N, Banta GT, Rivera-Monroy VH (2013) The importance of dissimilatory nitrate reduction to ammonium (DNRA) in the nitrogen cycle of coastal ecosystems. *Oceanography* **26**, 124–131.

Godfrey L V., Falkowski PG (2009) The cycling and redox state of nitrogen in the Archaean ocean. *Nature Geoscience* **2**, 725–729.

Gogarten JP, Townsend JP (2005) Horizontal gene transfer, genome innovation and evolution. *Nature reviews. Microbiology* **3**, 679–87.

Howard JB, Kechris KJ, Rees DC, Glazer AN (2013) Multiple amino acid sequence alignment nitrogenase component 1: insights into phylogenetics and structure-function relationships. *PloS one* **8**, e72751.

Hug LA, Baker BJ, Anantharaman K, Brown CT, Probst AJ, Castelle CJ, Butterfield CN, Hernsdorf AW, Amano Y, Ise K, Suzuki Y, Dudek N, Relman DA, Finstad KM, Amundson R, Thomas BC, Banfield JF (2016) A new view of the tree of life. *Nature Microbiology* **1**, 16048.

Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ (2010) Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* **11**, 119.

Isson TT, Love GD, Dupont CL, Reinhard CT, Zumberge AJ, Asael D, Gueguen B, McCrow J, Gill BC, Owens J, Rainbird RH, Rooney AD, Zhao M-Y, Stueeken EE, Konhauser KO, John SG, Lyons TW, Planavsky NJ (2018) Tracking the rise of eukaryotes to ecological dominance with zinc isotopes. *Geobiology* **16**, 341–352.

Javaux EJ, Marshall CP, Bekker A (2010) Organic-walled microfossils in 3.2-billion-year-old shallow-marine siliciclastic deposits. *Nature* **463**, 934–938.

Johnson BW, Goldblatt C (2018) EarthN: A new earth system nitrogen model. *Geochemistry,*

Geophysics, Geosystems.

Jones CM, Stres B, Rosenquist M, Hallin S (2008) Phylogenetic analysis of nitrite, nitric oxide, and nitrous oxide respiratory enzymes reveal a complex evolutionary history for denitrification. *Molecular Biology and Evolution* **25**, 1955–1966.

Kaiser JT, Hu Y, Wiig JA, Rees DC, Ribbe MW (2011) Structure of precursor bound NifEN: a nitrogenase FeMo cofactor maturase/insertase. *Science (New York, N.Y.)* **331**, 91.

Kechris KJ, Lin JC, Bickel PJ, Glazer AN (2006) Quantitative exploration of the occurrence of lateral gene transfer by using nitrogen fixation genes as a case study. *Proceedings of the National Academy of Sciences of the United States of America* **103**, 9584–9.

Kipp MA, Stüeken EE, Yun M, Bekker A, Buick R (2018) Pervasive aerobic nitrogen cycling in the surface ocean across the Paleoproterozoic Era. *Earth and Planetary Science Letters* **500**, 117-126.

Kitts PA, Church DM, Thibaud-Nissen F, Choi J, Hem V, Sapojnikov V, Smith RG, Tatusova T, Xiang C, Zherikov A, DiCuccio M, Murphy TD, Pruitt KD, Kimchi A (2016) Assembly: a resource for assembled genomes at NCBI. *Nucleic acids research* **44**, D73-80.

Koonin E V., Makarova KS, Aravind L (2001) Horizontal gene transfer in prokaryotes: Quantification and classification. *Annual Review of Microbiology* **55**, 709–742.

Lartillot N, Lepage T, Blanquart S (2009) PhyloBayes 3: a Bayesian software package for phylogenetic reconstruction and molecular dating. *Bioinformatics* **25**, 2286–2288.

Lepage T, Bryant D, Philippe H, Lartillot N (2007) A General Comparison of Relaxed Molecular Clock Models. *Molecular Biology and Evolution* **24**, 2669–2680.

Lipschultz F, Wofsy SC, Ward BB, Codispoti LA, Friedrich G, Elkins JW (1990) Bacterial transformations of inorganic nitrogen in the oxygen-deficient waters of the Eastern Tropical South Pacific Ocean. *Deep Sea Research Part A. Oceanographic Research Papers* **37**, 1513–1541.

Löscher CR, Kock A, Könneke M, LaRoche J, Bange HW, Schmitz RA (2012) Production of oceanic nitrous oxide by ammonia-oxidizing archaea. *Biogeosciences* **9**, 2419–2429.

Lyons TW, Reinhard CT, Planavsky NJ (2014) The rise of oxygen in Earth's early ocean and atmosphere. *Nature* **506**, 307–315.

Magnabosco C, Moore KR, Wolfe JM, Fournier GP (2018) Dating phototrophic microbial lineages with reticulate gene histories. *Geobiology* **16**, 179–189.

Marty B, Zimmermann L, Pujol M, Burgess R, Philippot P (2013) Nitrogen isotopic composition and density of the Archean atmosphere. *Science* **342**, 101–104.

Mather TA, Pyle DM, Allen AG (2004) Volcanic source for fixed nitrogen in the early Earth's atmosphere. *Geology* **32**, 905.

McGlynn SE, Boyd ES, Peters JW, Orphan VJ (2012) Classifying the metal dependence of uncharacterized nitrogenases. *Frontiers in microbiology* **3**, 419.

McInerney JO, McNally A, O'Connell MJ (2017) Why prokaryotes have pangenomes. *Nature Microbiology* **2**, 17040.

Michiels CC, Darchambeau F, Roland FAE, Morana C, Llirós M, García-Armisen T, Thamdrup B, Borges A V., Canfield DE, Servais P, Descy J-P, Crowe SA (2017) Iron-dependent nitrogen cycling in a ferruginous lake and the nutrient status of Proterozoic oceans. *Nature Geoscience* **10**, 217–221.

Mirkin BG, Fenner TI, Galperin MY, Koonin E V (2003) Algorithms for computing parsimonious evolutionary scenarios for genome evolution, the last universal common ancestor and dominance of horizontal gene transfer in the evolution of prokaryotes. *BMC Evolutionary Biology* **3**, 2.

Mojzsis SJ, Arrhenius G, McKeegan KD, Harrison TM, Nutman AP, Friend CRL (1996) Evidence for life on Earth before 3,800 million years ago. *Nature* **384**, 55–59.

Moore EK, Jelen BI, Giovannelli D, Raanan H, Falkowski PG (2017) Metal availability and the expanding network of microbial metabolisms in the Archaean eon. *Nature Geoscience* **10**, 629–636.

Navarro-González R, McKay CP, Mvondo DN (2001) A possible nitrogen crisis for Archaean life due to reduced nitrogen fixation by lightning. *Nature* **412**, 61–64.

Ogata H, Goto S, Sato K, Fujibuchi W, Bono H, Kanehisa M (1999) KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Research* **27**, 29–34.

Olson SL, Kump LR, Kasting JF (2013) Quantifying the areal extent and dissolved oxygen concentrations of Archean oxygen oases. *Chemical Geology* **362**, 35–43.

Pang K, Tang Q, Schiffbauer JD, Yao J, Yuan X, Wan B, Chen L, Ou Z, Xiao S (2013) The nature and origin of nucleus-like intracellular inclusions in Paleoproterozoic eukaryote microfossils. *Geobiology* **11**, 499–510.

Planavsky NJ, Asael D, Hofmann A, Reinhard CT, Lalonde S V, Knudsen A, Wang X, Ossa

Ossa F, Pecoits E, Smith AJB, Beukes NJ, Bekker A, Johnson TM, Konhauser KO, Lyons TW, Rouxel OJ (2014) Evidence for oxygenic photosynthesis half a billion years before the Great Oxidation Event. *Nature Geoscience* **7**, 283–286.

Polz MF, Alm EJ, Hanage WP (2013) Horizontal gene transfer and the evolution of bacterial and archaeal population structure. *Trends in Genetics* **29**, 170–5.

Popa O, Hazkani-Covo E, Landan G, Martin W, Dagan T (2011) Directed networks reveal genomic barriers and DNA repair bypasses to lateral gene transfer among prokaryotes. *Genome research* **21**, 599–609.

R Core T (2013) R: A Language and Environment for Statistical Computing.

Raymond J, Siefert JL, Staples CR, Blankenship RE (2004) The natural history of nitrogen fixation. *Molecular Biology and Evolution* **21**, 541–54.

Reinhard CT, Planavsky NJ, Robbins LJ, Partin CA, Gill BC, Lalonde S V., Bekker A, Konhauser KO, Lyons TW (2013) Proterozoic ocean redox and biogeochemical stasis. *Proceedings of the National Academy of Sciences* **110**, 5357–5362.

Revell LJ (2012) phytools: an R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution* **3**, 217–223.

Roberson AL, Roadt J, Halevy I, Kasting JF (2011) Greenhouse warming by nitrous oxide and methane in the Proterozoic Eon. *Geobiology* **9**, 313–320.

Sánchez-Baracaldo P, Ridgwell A, Raven JA (2014) A neoproterozoic transition in the marine nitrogen cycle. *Current Biology* **24**, 652–657.

Santos PC Dos, Fang Z, Mason SW, Setubal JC, Dixon R (2012) Distribution of nitrogen fixation and nitrogenase-like sequences amongst microbial genomes. *BMC Genomics* **13**, 162.

Schoepp-Cothenet B, Lis R van, Atteia A, Baymann F, Capowiez L, Ducluzeau A-L, Duval S, Brink F ten, Russell MJ, Nitschke W (2013) On the universal core of bioenergetics. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1827**, 79–93.

Schwartz JH, Maresca B (2006) Do Molecular Clocks Run at All? A Critique of Molecular Systematics. *Biological Theory* **1**, 357–371.

Scott C, Lyons TW, Bekker A, Shen Y, Poulton SW, Chu X, Anbar AD (2008) Tracing the stepwise oxygenation of the Proterozoic ocean. *Nature* **452**, 456–459.

Som SM, Buick R, Hagadorn JW, Blake TS, Perreault JM, Harnmeijer JP, Catling DC (2016) Earth's air pressure 2.7 billion years ago constrained to less than half of modern levels. *Nature*

Geoscience **9**, 448–451.

Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics (Oxford, England)* **30**, 1312–1313.

Stanton CL, Reinhard CT, Kasting JF, Ostrom NE, Haslun JA, Lyons TW, Glass JB (2018) Nitrous oxide from chemodenitrification□: A possible missing link in the Proterozoic greenhouse and the evolution of aerobic respiration 1–13.

Stein LY, Klotz MG (2016) The nitrogen cycle. *Current Biology* **26**, R94–R98.

Stolz JF, Basu P (2002) Evolution of Nitrate Reductase: Molecular and Structural Variations on a Common Function. *ChemBioChem* **3**, 198–206.

Stüeken EE, Buick R, Guy BM, Koehler MC (2015) Isotopic evidence for biological nitrogen fixation by molybdenum-nitrogenase from 3.2 Gyr. *Nature*.

Stüeken EE, Catling DC, Buick R (2012) Contributions to late Archaean sulphur cycling by life on land. *Nature Geoscience* **5**, 722–725.

Stüeken EE, Kipp MA, Koehler MC, Buick R (2016a) The evolution of Earth's biogeochemical nitrogen cycle. *Earth-Science Reviews* **160**, 220–239.

Stüeken EE, Kipp MA, Koehler MC, Schwieterman EW, Johnson B, Buick R (2016b) Modeling p N $_2$ through Geological Time: Implications for Planetary Climates and Atmospheric Biosignatures. *Astrobiology* **16**, 949–963.

The Uniprot Consortium (2017) UniProt: the universal protein knowledgebase. *Nucleic Acids Research* **45**, D158–D169.

Ueno Y, Yamada K, Yoshida N, Maruyama S, Isozaki Y (2006) Evidence from fluid inclusions for microbial methanogenesis in the early Archaean era. *Nature* **440**, 516–519.

Voss M, Bange HW, Dippner JW, Middelburg JJ, Montoya JP, Ward B (2013) The marine nitrogen cycle: recent discoveries, uncertainties and the potential relevance of climate change. *Philosophical Transactions of the Royal Society B: Biological Sciences* **368**, 20130121–20130121.

Wada E, Hatton A (1971) Nitrite metabolism in the euphotic layer of the central North Pacific Ocean. *Limnology and Oceanography* **16**, 766–772.

Weiss MC, Sousa FL, Mrnjavac N, Neukirchen S, Roettger M, Nelson-Sathi S, Martin WF (2016) The physiology and habitat of the last universal common ancestor. *Nature Microbiology* **1**, 16116.

Zaremba-Niedzwiedzka K, Caceres EF, Saw JH, Bäckström D, Juzokaite L, Vancaester E, Seitz KW, Anantharaman K, Starnawski P, Kjeldsen KU, Stott MB, Nunoura T, Banfield JF, Schramm A, Baker BJ, Spang A, Ettema TJG (2017) Asgard archaea illuminate the origin of eukaryotic cellular complexity. *Nature* **541**, 353–358.

Zerkle AL, Mikhail S (2017) The geobiological nitrogen cycle: From microbes to the mantle. *Geobiology*.

Zerkle AL, Poulton SW, Newton RJ, Mettam C, Claire MW, Bekker A, Junium CK (2017) Onset of the aerobic nitrogen cycle during the Great Oxidation Event. *Nature* **542**, 465–467.

Tables

Calibration Event	Conservative (Ga)	Refs	Liberal (Ga)	Refs
LUCA (set as root prior)	3.8	Mojzsis *et al.*, 1996	3.8	Mojzsis *et al.*, 1996
Origin of Oxygenic Photosynthesis	>2.45	Bekker *et al.*, 2004	>3	Lyons *et al.*, 2014
Origin of Methanogenesis	>2.7	Eigenbrode & Freeman, 2006	>3.46	Ueno *et al.*, 2006
Origin of Eukaryotes	>1.7	Pang *et al.*, 2013	>3.2	Javaux *et al.*, 2010

Table 1. Fossil calibration points used in Phylobayes runs. Conservative time points reflect the dates around which there is the most consensus; liberal time points reflect the earliest dates for which there is evidence of these events based on the current scientific literature.

Gene	Number of genes identified
amo	4
anfD	11
anfK	10
hao	0
napA	37
narG	34
nasA	28
nifD	165
nifK	92
nifH	233
nirB	39
nirK	10
nirS	16
norB	36
nosZ	21
nrfA	48
nxrA	34
vnfD	6
vnfK	6

Table 2. Number of nitrogen-cycling genes identified within 254 bacterial and archaeal genomes that were included in the analysis. All genes were identified from ORFs using blastp with an e-value cutoff of 1e-12, then filtered using the Kofam Koala tool. Nitrogenases were further refined by identifying key residues (see Methods).

Gene	HGT	Speciation	Loss	Duplication
anfD	7	7	4	0
anfK	6	7	4	0
napA	15	39	18	0
narG	14	34	20	5
nasA	5	28	7	1
nifD	63	162	61	0
nifK	37	109	56	1
nifH	79	214	73	12
nirB	16	31	12	3
nirK	4	9	5	1
nirS	8	10	4	1
norB	26	11	4	2
nosZ	13	8	3	2
nrfA	25	31	12	3
nxrA	16	28	15	4
vnfD	3	3	1	0
vnfK	3	3	1	0

Table 3. Frequencies of events inferred by AnGST for each gene. These results are based on the CIR clock tree of life with the conservative set of calibration points (see Supplementary Tables 1-3 for results under alternate clocks and calibration points).

Gene	Upper	Lower	Midpoint
nirS	1448.64	1436.28	1442.46
vnfD	1452.51	499.84	976.17
vnfK	1452.51	499.84	976.17
norB	1544.32	1436.28	1490.30
nirK	1602.53	1482.46	1542.50
nosZ	2066.05	1544.32	1805.18
anfD	2535.66	1910.29	2222.97
anfK	2535.66	1910.29	2222.97
nasA	2736.69	2225.74	2481.21
nrfA	2665.36	2479.85	2572.60
nirB	2811.92	2684.98	2748.45
nifK	3136.45	2828.7	2982.61
napA	3028.85	2997.31	3013.08
narG	3028.85	2997.31	3013.08
nxrA	3028.85	2997.31	3013.08
nifD	3146.17	3078.86	3112.51
nifH	3240.75	3171.85	3206.30

Table 4. Inferred birth dates for nitrogen-cycling genes using the CIR conservative chronogram. These estimates do not take into account uncertainty in the timings of divergences in the chronogram. Events are inferred to occur along branches, so here we list the inferred timing for the earliest possible timing (upper) and latest possible timing (lower) as well as the midpoint between the two. Times are reported in millions of years ago.

Figure 1. Schematic of the biological nitrogen cycle. Arrows are labelled with the pathway to which they belong. All genes examined in this study are labeled next to the step they catalyze. Genes without sufficient data for subsequent analysis are labeled in grey. Adapted from Canfield et al. (2010).

Figure 2. Species tree used for phylogenetic analysis. Maximum likelihood phylogeny based on an alignment of 308 concatenated single-copy universal proteins. Well-represented bacterial and archaeal phyla are labelled in black text. Bacterial clades are labeled in alternating colors of blue, archaea in green, eukaryotes in red. Bootstrap values (from 100 bootstraps) greater than 50 are shown as transparent gray circles, with larger circles representing higher bootstrap values.

Figure 3. Histograms and density plots of all speciation, loss, duplication and horizontal gene transfer (HGT) events. All events are shown in blue, HGT events alone are shown in green. Y-axis represents proportion of all events (blue) or HGT events only (green). Borders around graphs indicate the metabolic process that the gene is involved in. Histograms are ordered according to the earliest time point at which a speciation, loss, duplication, or HGT event occurred.