Metabolic and haemodynamic resting-state connectivity of the human brain: a high-temporal resolution simultaneous BOLD-fMRI and FDG-fPET multimodality study

Sharna D Jamadar^[1,2,3]*, Phillip GD Ward^[1,2,3], Emma Xingwen Liang[1], Edwina R Orchard[1,2,3], Zhaolin Chen[1,4], Gary F Egan[1,2,3]

1. Monash Biomedical Imaging, 770 Blackburn Rd, Melbourne, Vic, 3800 Australia
2. Turner Institute for Brain and Mental Health, Monash University, Wellington Rd, Melbourne Vic 3800 Australia
3. Australian Research Council Centre of Excellence for Integrative Brain Function, Australia
4. Department of Electrical and Computer Systems Engineering, Monash University, Wellington Rd, Melbourne Vic 3800 Australia

^ These authors contributed equally

ORCiDs:
Jamadar: 0000-0001-7222-7181
Ward: 0000-0001-5934-6944
Liang: 0000-0002-9906-2804
Orchard: 0000-0003-4845-8961
Chen: 0000-0002-0173-6090
Egan: 0000-0002-3186-4026

* Corresponding Author:
Sharna D Jamadar, PhD
770 Blackburn Rd,
Melbourne VIC 3800
Australia
Ph: +61 3 9902 9751
Sharna.jamadar@monash.edu

Abstract

Simultaneous FDG-PET/fMRI ([18F]-fluorodeoxyglucose positron emission tomography functional magnetic resonance imaging) provides the capacity to image two sources of energetic dynamics in the brain – glucose metabolism and haemodynamic response. Functional fMRI connectivity has been enormously useful for characterising interactions between distributed brain networks in humans. Metabolic connectivity based on static FDG-PET has been proposed as a biomarker for neurological disease; but static FDG-PET cannot be used to estimate subject-level measures of *connectivity*, only across-subject *covariance*. Here, we applied high-temporal resolution constant infusion functional PET (fPET) to measure subject-level metabolic connectivity simultaneously with fMRI connectivity. fPET metabolic connectivity was characterised by fronto-parietal connectivity within and between hemispheres. fPET metabolic connectivity showed moderate similarity with fMRI primarily in superior cortex and frontoparietal regions. fPET metabolic connectivity showed little similarity with static FDG-PET metabolic covariance, indicating that metabolic connectivity is a non-ergodic process that cannot be used to infer individual-level connectivity from across-subject covariance. In sum, our results highlight the complementary strengths of fPET and fMRI in measuring the intrinsic connectivity of the brain, and opens up the opportunity for new multivariate metabolic neuroimaging biomarkers for neurological disease.

Keywords functional connectivity, metabolic connectivity, simultaneous MR-PET, FDG-PET, BOLD-fMRI, fPET, resting-state

Introduction

Resting-state connectivity measures the temporal coherence of spontaneous neural activity between spatially distinct regions (Fox and Raichle, 2007). Resting-state connectivity is commonly measured using blood oxygenation level dependent functional magnetic resonance imaging (BOLD-fMRI), which indexes spontaneous large-amplitude changes in blood oxygenation at low frequencies (Biswal et al., 1995). Numerous studies have shown that fMRI resting-state 'functional' connectivity is associated with variability in cognition (e.g., Fox et al., 2007; Jamadar et al., 2016), age (Andrews-Hanna et al., 2007), sex (Jamadar et al., 2018), genetic variants (Fornito et al., 2011), psychiatric conditions (Fornito and Bullmore, 2010; Garrity et al., 2007), and neurodegenerative disease (e.g., Alzheimer's disease, Greicius et al., 2004; Huntington's disease, Poudel et al., 2014). Resting-state fMRI connectivity is under investigation as a potential biomarker of clinical progression for a number of diseases, although its application as a diagnostic biomarker is currently limited (Hohenfeld et al., 2018).

BOLD-fMRI provides a haemodynamic-based surrogate index of neuronal activity with a temporal resolution in the order of seconds (~2sec; sub-second resolution with multiband sequences, Feinberg and Yacoub, 2012), and sub-millimeter spatial resolution. The BOLD-fMRI signal is an indirect index of neuronal function, arising from neurovascular coupling between neuronal activity and cerebral haemodynamics (Buxton et al., 2004; Phillips et al., 2016). While BOLD-fMRI is usually interpreted as arising from neuronal activity, there are a number of non-neuronal contributors to the BOLD signal, including heart rate variability, respiration, head movement, individual variability in haemoglobin and the oxygen-carrying capacity of the blood, etc. (Liu, 2017; Levin et al., 2001; Ward et al., 2019). Signal loss and distortion occurs as a result of magnetic field inhomogeneities, and are particularly pronounced at air-tissue interfaces, including the orbitofrontal and temporal poles (Yeo et al., 2011). The BOLD signal includes contributions from haemodynamic responses to neuronal signals, non-neuronal physiological signals, instrumentational signal artefacts, and motion-related artefacts (Caballero-Gaudes and Reynolds, 2017). As such, BOLD-fMRI is a semi-quantitative non-absolute index of neural activity, and BOLD-fMRI responses cannot be quantitatively compared across brain regions, subjects, or imaging sites, or within the same individual across time (Logothetis, 2008). These characteristics are a major limiting factor in the development of BOLD-fMRI-based biomarkers, particularly when derived from fMRI functional connectivity metrics (Hohenfeld et al., 2018).

Resting-state 'functional' connectivity usually refers to connectivity estimated from BOLD-fMRI data. However, connectivity can be measured using many neuroimaging techniques, with each method providing a unique perspective on information transfer in the brain. For example, connectivity measured using electrophysiological methods such as electroencephalography (EEG) characterises the coherence in electrical signal amplitude and phase across the brain (Srinivasan et al., 2007). In this study, we investigate brain connectivity derived from positron emission tomography (PET) to measure co-variation of glucose uptake throughout the human brain; an index of cerebral metabolism. 18F-fluorodeoxyglucose (FDG)-PET provides a snapshot of cerebral glucose uptake, and cross-sectional connectivity analyses of FDG-PET have characterised the covariation in cerebral glucose uptake across subjects (Horwitz et al., 1984; Moeller et al., 1987). This measure of 'metabolic' connectivity is considered an important complement to BOLD-fMRI functional connectivity, since FDG-PET represents a more direct and quantitative measure of neuronal function than BOLD-fMRI. However, while the interpretation of FDG-PET brain connectivity is not limited in the same manner as BOLD-fMRI, the majority of FDG-PET studies have acquired static images of the brain with an effective temporal resolution of the scan duration – often 10-40mins. Conventionally, FDG-PET based measures of connectivity are referred to as 'metabolic' connectivity whilst BOLD-fMRI based measures of connectivity are 'haemodynamic' connectivity, as both measures are an index of the functional activity of the brain.

The application of connectome methodologies to static FDG-PET resting-state data has revealed both similarities and differences between metabolic FDG-PET connectivity and haemodynamic

BOLD-fMRI connectivity. Using independent component analysis (ICA) network approaches, Di et al., (2017) and Savio et al., (2017) found that brain networks encompassing left-right homologous regions were evident in static FDG-PET resting-state data, but networks involving long-range anterior-posterior connectivity were absent or reduced. This pattern was also obtained using a matrix-based analytical approach (Di et al., 2017). Together, these studies demonstrated only a modest overlap between static FDG-PET connectivity and BOLD-fMRI connectivity. A major limitation of these studies is that since static FDG-PET acquisitions provide a single cumulative measurement per subject, metabolic 'connectivity' was estimated *across-subjects*. As such, these results are more accurately characterised as metabolic *covariance*, rather than *connectivity*. Covariance measures are not dependent on the temporal correlation of glucose uptake and therefore provide patterns that do not necessarily arise from simultaneous activity in brain regions. Furthermore, it is known from BOLD-fMRI that group-level estimates of connectivity do not always accurately predict individual subject-level connectivity (Roberts et al., 2016). Consequently, static FDG-PET images cannot be used to estimate metabolic connectivity within single subjects, greatly limiting their application as a disease biomarker (cf. Yakushev et al., 2017; Veronese et al., 2019).

Recent advances in radiotracer delivery, together with the improved PET signal detection sensitivity of dual-modality MR-PET scanners, has made it possible to study the dynamics of FDG-PET glucose uptake with substantially improved temporal resolution. The method described as 'functional' FDG-PET (FDG-fPET) administers the radiotracer over the course of the scan, either as a constant infusion (e.g., Villien et al., 2014; Hahn et al., 2017; Hahn et al., 2016; Rischka et al., 2018; Jamadar et al., 2019b; Jamadar et al., 2019a; Li et al., 2019), or as a bolus+infusion (e.g. Rischka et al., 2018; Jamadar et al., 2019a). This protocol, adapted from (Carson, 2000), aims to maintain a constant plasma supply of radiotracer to allow examination of glucose uptake with an improved temporal resolution. Task-based FDG-fPET studies have achieved a temporal resolution of 60sec (Hahn et al., 2016; Hahn et al., 2017; Jamadar et al., 2019b; Villien et al., 2014; Li et al., 2019) or less (12sec, Rischka et al., 2018; 16sec, Jamadar et al., 2019a; 30sec, (Hahn et al., 2020).

The goal of the present study was to investigate FDG-fPET within-subject metabolic connectivity with high temporal resolution. We used a constant infusion radiotracer administration protocol (Jamadar et al., 2019a), reconstructed the FDG-fPET data into images of 16sec acquisition duration, and simultaneously acquired BOLD-fMRI data with a temporal resolution of 2.45sec. Two major hypotheses were tested in the present study. Firstly, based on the findings of Roberts et al. (2016) that group-level BOLD-fMRI covariance measures predict seed-based functional connectivity only to a weak-to-moderate degree, we hypothesised that across-subject metabolic covariance, or static FDG-PET (*s*PET), would be only moderately correlated with within-subject metabolic brain connectivity (fPET). Secondly, on the basis of resting-state static FDG-PET ICA results (Di and Biswal, 2012), we predicted that within-subject resting-state metabolic brain connectivity (fPET) would be highly correlated with resting-state BOLD-fMRI connectivity.

Methods and Materials

All methods were reviewed by the Monash University Human Research Ethics Committee, in accordance with the Australian National Statement on Ethical Conduct in Human Research (2007). Administration of ionising radiation was approved by the Monash Health Principal Medical Physicist, in accordance with the Australian Radiation Protection and Nuclear Safety Agency Code of Practice (2005). For participants aged over 18yrs, the annual radiation exposure limit of 5mSv applies and the effective dose equivalent was 4.9mSv. Detailed information on the constant infusion acquisition procedure is available in Jamadar et al. (2019b).

Participants

Participants (n=28) were aged 18-23 years (mean 19 years); 21 female, all right handed (Edinburgh Handedness Inventory). Participants had between 13-18 years of education (mean 14 years), normal or corrected-to-normal vision and no personal history of diagnosed Axis-1 mental illness, diabetes or cardiovascular illness. Participants were screened for claustrophobia, non-MR compatible implants, clinical or research PET scan in the past 12-months, and women were screened for current or suspected pregnancy. Prior to the scan, participants were directed to consume a high protein/low sugar diet for 24-hours, fast for 6 hours, and drink 2-6 glasses of water. Blood sugar level was measured using an Accu-Check Performa (model NC, Mannheim, Germany); all participants had blood sugar levels below 10mmol/L (max in this sample 4.73mmol/L).

Procedure

Prior to the scan, participants completed a brief cognitive battery (30mins, results not reported here). Participants were then cannulated in the vein in each forearm with a minimum size 22-gauge cannula, and a 10mL baseline blood sample was taken at the time of cannulation. For all participants, the left cannula was used for FDG infusion, and the right cannula was used for blood sampling. Primed extension tubing was connected to the right cannula (for blood sampling) via a three-way tap.

Participants underwent a 95-minute simultaneous MR-PET scan in a Siemens (Erlangen) Biograph 3 Tesla molecular MR (mMR) scanner (Figure 1A). Participants were positioned supine in the scanner bore with head in a 16-channel radiofrequency (RF) head coil, and were instructed to lie as still as possible with eyes open, and think of nothing in particular. [18-F] fluroodeoxyglucose (FDG; average dose 233MBq) was infused over the course of the scan at a rate of 36mL/hr using a BodyGuard 323 MR-compatible infusion pump (Caesarea Medical Electronics, Caesarea, Israel). One participant received a lower dose (167MBq) due to infusion pump error. Infusion onset was locked to the onset of the PET scan.

Plasma radioactivity levels were measured throughout the duration of the scan. At 10-mins post-infusion onset, a 10mL blood sample was taken from the right forearm using a vacutainer; the time of the 5mL mark was noted for subsequent decay correction. Subsequent blood samples were taken at 10-min intervals for a total of 10 samples for the duration of the scan. The cannula line was flushed with 10mL of saline after every sample to minimise line clotting. Immediately following blood sampling, the sample was placed in a Heraeus Megafuge 16 centrifuge (ThermoFisher Scientific, Osterode, Germany) and spun at 2000rpm for 5 mins; 1000 µL plasma was pipetted, transferred to a counting tube and placed in a well counter for 4mins. The count start time, total number of counts, and counts per minute were recorded for each sample. Figure 1B shows the interpolated plasma radioactivity concentration over time. The average radioactivity concentration constantly increases over time with the lowest relative slope at the end of the acquisition.

MR-PET protocol

PET data was acquired in list mode. Infusion of FDG radiotracer and PET data acquisition started with the Ultrashort TE (UTE) MRI for PET attenuation correction (Figure 1A). To allow the PET signal to rise to detectable levels, non-functional MRI scans were acquired in the first 30-mins following infusion onset. These scans, in order, were: UTE (TA=1.40mins), T1 3D MPRAGE (TA = 7.01mins, TR=1640ms, TE=2.34ms, flip angle = 8°, FOV=256x256mm^2, voxel size = 1x1x1mm^3, 176 slices; sagittal acquisition), and several scans not reported here: T2 SPACE (TA=5.52min), magnetic resonance spectroscopy (MRS, TA=2.48min), gradient field map (TA=1.02min), pulsed arterial spin labelling (TA=4.21), T2 susceptibility-weighted image (TA=6.50min), left-right phase correction (TA=0.21min). For the remainder of the scan, six consecutive 10min blocks of T2*-weighted echo planar images (EPIs) were acquired (TR=2450ms, TE=30ms, FOV=190mm, 3x3x3mm3 voxels, 44 slices, ascending axial acquisition).

MR image preparation

For the T1 structural image, the brain was extracted, then registered to MNI152 space using Advanced Normalization Tools (ANTs) (Avants et al., 2011). The grey matter, white matter and brain cortex labels of T1 image were segmented by using Freesurfer with Desikan-Killiany Atlas (Diedrichsen et al. 2009).

The six blocks of EPI scans for all subjects (a total of 245 EPIs) underwent a standard fMRI pre-processing pipeline. Specifically, all scans were brain extracted (FSL BET Smith, 2002), corrected for intensity non-uniformity using N4 bias field correction (ANTs Tustison et al., 2010), motion corrected (FSL MCFLIRT Jenkinson et al., 2002), slice timing corrected (AFNI Cox, 1996) and high-pass filtered (Hz > 0.01) to remove low frequency noise (FSL Jenkinson et al., 2012).

PET image reconstruction and preparation

The 5700-second list-mode PET data for each subject was binned into 356 3D sinogram frames each of 16-second interval. The pseudoCT method (Burgos et al., 2014) was used to correct the attenuation for all acquired data. Ordinary Poisson-Ordered Subset Expectation Maximization (OP-OSEM) algorithm (3 iterations, 21 subsets) with point spread function correction was used to reconstruct 3D volumes from the sinogram frames. The reconstructed DICOM slices were converted to NIFTI format with size 344x344x127 (voxel size: 2.09 x 2.09 x2.03 mm^3) for each volume. A 5-mm FWHM Gaussian post-filter was applied to each 3D volume. All 3D volumes were temporally concatenated to form a 4D (344 x 344 x 127 x 356) NIFTI volume. A guided motion correction method using simultaneously acquired MRI images was applied to correct the motion during the PET scan. We retained the 225 16-sec volumes commencing from the 30 minute timepoint, which matched the start of the BOLD-fMRI EPI acquisition, for further analyses. A single static PET image was derived from the sum of the 16-sec volumes.

The 225 PET volumes were motion corrected (FSL MCFLIRT Jenkinson et al., 2002); the mean PET image was brain extracted and used to mask the 4D data. The fPET data was further processed using a spatio-temporal gradient filter to estimate the short-term change in glucose uptake from the cumulative glucose uptake that was measured. This approach intrinsically adjusted for the mean signal whilst avoiding global-signal regression and other approaches that may create spurious anti-correlations in the data (Li et al., 2019; Murphy and Fox, 2017). Due to radiotracer dynamics, it was not expected that the fPET sensitivity would be uniform across the 60minutes of the resting-state data acquisition. As the radiotracer accumulated in the brain, it was anticipated that the signal-to-noise ratio of the PET image reconstruction would progressively improve. The Supplement includes the definition of the spatiotemporal filter (Supplementary Figure 3), the variability of the fPET connectivity matrix with different filter parameters (Supplementary Figure 4), and the variability of the fPET matrix across the scanning duration (Supplementary Figure 5).

Connectivity Analyses

fPET and fMRI timeseries were extracted for each of the 82 regions of interest from the segmentation of the T1-weighted image, interpolated using an ANTs rigid registration (Avants et al., 2011). To construct a connectivity matrix, Pearson's correlation coefficients were estimated between the timeseries from pairs of regions. This produced a single per-subject per-modality 82x82 matrix corresponding to the 60minutes of resting-state in the experimental protocol.

Group-average connectivity for BOLD-fMRI and FDG-fPET: The similarity between the group-average BOLD-fMRI and FDG-fPET matrices was assessed using a Pearson correlation across all edges on the lower triangle of the connectivity matrices (excluding the diagonal and symmetric triangle). The regional variation between the two modalities was assessed by calculating a separate

correlation coefficient for each row of the connectivity matrix, providing a measure of similarity of the connectivity profile for each brain region.

Inter-subject variability of BOLD-fMRI and FDG-fPET: To further characterise and contrast connectivity from the two modalities, the strength and variance of connections to each region were examined. Each connectivity matrix was discretised at the 90^{th} percentile level to provide a binary graph of the 'strongest' connections. We took this approach, rather than a threshold approach based on null-hypothesis testing, as the SNR properties of each modality are substantially different. Thus, a common threshold across all three modalities would lead to many significant regions in some modalities (i.e. sPET) and very few in others (i.e. fPET). We argue that the percentile approach is appropriate in this analysis, as the goal was to examine and compare the distribution of most connected regions between modalities, without the additional concerns of sub-threshold results. The number of these binary edges connected to each region was calculated to provide a map of regional degree for each modality. The inter-subject variation of the connectivity matrix was assessed by calculating the variance in the correlation coefficient for each edge in the matrix across subjects and computing the mean for each region, to provide a map of regional consistency.

Regional connectivity of BOLD-fMRI and FDG-fPET: To summarise the pattern of connectivity, node degree (i.e. the number of connections to each region) for each Desikan-Killaney region was calculated and sorted. For each region, the number of edges in the top 10% of connected regions was counted and projected onto the anatomical surface. To examine the consistency of connections across subjects, the standard deviation of connectivity strength across subjects was also projected onto the anatomical surface.

sPET metabolic covariance: A FDG-sPET metabolic covariance matrix was constructed using the mean signal for each subject from the static PET image. The signal was de-meaned for each subject to remove inter-subject variances, such as dosage and physiology. The across-subject correlation between pairs of ROIs was then estimated to generate the covariance matrix. The similarity of across-subject metabolic covariance of sPET and inter-subject temporal correlation/metabolic connectivity of fPET was examined at a regional and global level. A comparison of the characteristics of the metabolic covariance and the metabolic connectivity was performed, assessing the scale of correlations and the network profile.

Graph-based network analysis: To investigate the patterns of connectivity within and between subnetworks, node degree was calculated and regions were sorted by subnetwork assignment (i.e., frontoparietal, dorsal attention, ventral attention, default mode, somatomotor, limbic, subcortical, and visual; as classified by Yeo et al., 2011). As the regions were not evenly distributed between subnetworks, some subnetworks (e.g. default mode network) contained a higher number of nodes than others. Therefore, there was a higher chance that significant edges would occur within subnetworks with more nodes by random chance. To minimise this bias in our interpretation, we adjusted node degree for the capacity of each network (number of observed edges, divided by number of potential edges between two subnetworks).

Correlation between BOLD and FDG: To investigate the discrepancies and correlations between the two modalities, the relationship between single-modality functional connectivity strength and inter-modality correlation was examined. A non-parametric correlation was calculated between the single-modality strength and cross-modality correlation (Spearman correlation) whilst controlling for ROI size. This provided an assessment of whether agreement between the two modalities was more pronounced for edges with strong fMRI or fPET connectivity. This was explored visually by ordering edges based upon their fMRI connectivity strength and plotting the cross-subject correlation between the two modalities for each edge. A similar plot using the fPET connectivity strength was overlaid on the same axis. The order of connectivity strength, rather than correlational coefficient, was used in this analysis as the sensitivity to and the range of connectivity measured by the two modalities differed substantially.

Results

Metabolic covariance, metabolic connectivity, and haemodynamic connectivity

The across-subject sPET metabolic covariance matrix showed large variability in connection strength across ROIs (Figure 2A). Each cortical sub-division showed strong covariance with neighbouring regions, as evident by the high covariance values along the main diagonal. Furthermore, each cortical sub-division showed strong homotopic covariance, as evident by the high covariance values along the diagonals of the top right and bottom left quadrants of the matrix. High temporo-subcortical covariance was observed both within and between hemispheres; and high fronto-subcortical, fronto-temporal and parieto-occipital covariance was also apparent albeit of smaller magnitude than the temporo-subcortical covariance. FDG-sPET showed the highest region degree in the temporal poles and subcortically (Figure 4A). In the network graphs (Figure 3A), sPET showed strong widespread connectivity between the dorsal attention, frontoparietal, subcortical, sensorimotor and default mode networks.

The most striking effect for the group-averaged within-subject FDG-fPET metabolic connectivity matrix was the strong fronto-parietal connectivity, both within and between hemispheres (Figure 2B). Left-right homotopic connectivity was also evident for frontal and parietal cortices; this was not visually apparent for subcortical, temporal and occipital cortices. fPET showed the highest degree in the frontal poles and the superior cortex (Figure 4). For the network graphs (Figure 3B) fPET showed high connectivity between the frontoparietal and dorsal attention networks, and between the dorsal attention and sensorimotor networks. Moderate connectivity was apparent between the sensorimotor-frontoparietal networks, and between the default mode and dorsal attention, frontoparietal and visual attention networks. The dorsal attention network showed strong within-network connectivity as well.

BOLD-fMRI showed strong connectivity within anatomical sub-divisions, both within and between hemispheres (Figure 2C). This pattern was evident from the four diagonal lines in the fMRI connectivity matrix. A number of long-range connections between anatomical sub-divisions were also evident, including fronto-parietal, parieto-occipital and temporo-parietal regional connectivity. These long-range connections were evident both within and between hemispheres but were of smaller magnitude than the short-range and homotopic connections. Network graphs for the fMRI data (Figure 3C) showed strongest connectivity between the dorsal attention and visual networks, with the visual network having the highest self-connectedness. The visual attention network showed high connectedness with the frontoparietal and sensorimotor network, and the default mode network showed an intermediate level of connectedness with the frontoparietal and dorsal attention networks. The anatomical projections of the most highly connected regions (Figure 4, lower row) indicated that the parietal cortex was most inter-connected region in the fMRI data, with superior frontal and inferior occipital cortices showing intermediate levels of haemodynamic based connectivity. Subcortical and orbitofrontal regions were the least inter-connected regions in the BOLD-fMRI data.

Comparison of FDG-sPET metabolic covariance and FDG-fPET metabolic connectivity

Qualitatively the sPET covariance and fPET connectivity matrices show little similarity (Figure 2A & 2B). The fPET connectivity was dominated by fronto-parietal connectivity, whereas the sPET covariance demonstrated greater variability throughout the connectivity matrix. The distribution of edges for each r value for sPET and fPET showed that fPET is biased towards positive r values, whereas sPET is symmetric around r=0 (Supplementary Figure 1). Comparison of the network graphs (Figure 3) and the anatomical projections of the most connected regions (Figure 4) are consistent with the finding that sPET and fPET show little anatomical similarity across the brain. This result is confirmed by the cross-modal correlations (Figure 5A upper row & Figure 5B left column) which highlights that sPET and fPET connectivity has low correlation across the brain.

Taken together, these results are consistent with the conclusion that across-subject sPET connectivity does not predict within-subject fPET connectivity. These results are not consistent with the hypothesis that static metabolic connectivity would be moderately correlated with within-subject metabolic brain connectivity.

Comparison of FDG-fPET metabolic connectivity and BOLD-fMRI haemodynamic connectivity

Qualitatively the FDG-fPET metabolic connectivity and BOLD-fMRI haemodynamic connectivity matrices showed moderate similarity, with a trend towards stronger connectivity associated with the parietal cortex (Figure 2). Both fPET and fMRI network graphs (Figure 3) showed strong connectivity for the dorsal attention network, with the dorsal attention network strongly connected to the frontoparietal network for fPET, and strongly connected to the visual network for fMRI. The default mode network showed moderate connectedness for both fPET and fMRI. The anatomical projections of the fPET-fMRI correlation showed the strongest correlation in anterior superior frontal and superior parietal regions. Intermediate correlations were obtained in the lateral temporal and temporal pole regions. The lowest correlations were obtained in subcortical regions. These interpretations were confirmed by showing that numerous regions of interest showed strong correlation between fPET and fMRI (Figure 5B right column). Taken together, these results suggest that fPET shows stronger similarity to fMRI than sPET, and is consistent with the hypothesis that within-subject resting-state metabolic brain connectivity would be correlated with resting-state haemodynamic connectivity.

Discussion

In this study, we report for the first time resting-state metabolic brain connectivity measured using a novel FDG-functional PET imaging protocol and methodology in humans. A key strength of this work is the simultaneous acquisition and high temporal resolution that enabled multi-modality within-subject analyses of resting state brain activity. The simultaneous acquisition of fPET and fMRI data is of particular importance, as measures of metabolic and haemodynamic responses to the same neural activity without the confound of intra-individual differences in attention, fatigue, motivation, nutrient intake and blood chemistry that occur in sequential brain imaging experiments (Chen et al., 2018). We compared within-subject functional PET metabolic connectivity with across subject static PET metabolic covariance, and with BOLD-fMRI haemodynamic connectivity. Resting-state functional FDG-PET connectivity showed low similarity with static FDG-PET covariance and greater similarity with BOLD-fMRI connectivity, particularly in fronto-parietal cortical regions.

Metabolic connectivity is predominantly evident in fronto-parietal regions

FDG-fPET metabolic connectivity was dominated by fronto-parietal connectivity both within and between hemispheres. This is consistent with early evidence showing the frontoparietal cortices are highly metabolically active compared to temporo-occipital regions (Sasaki et al., 1986), particularly at rest (Shokri-Kojori et al., 2019; Vaishnavi et al., 2010). This pattern was confirmed by the brain network graphs where the dorsal attention network was strongly connected to the frontoparietal and sensorimotor networks, and showed strong intra-connectedness. Together, the frontoparietal and dorsal attention networks are responsible for flexible, goal directed behaviour (Laird et al., 2011), and the two networks interact closely to subserve action-focused attentional control (Dixon et al., 2018). The inter-connectedness of the frontoparietal and dorsal attention networks with the sensorimotor network is consistent with this interpretation. Interestingly, the dorsal attention and frontoparietal control networks have the highest metastability, update efficiency, and neural synchrony in comparison to other resting-state networks (Alderson et al., 2020). These networks demonstrate a resting-state configuration that is similar to a task-related configuration, which thereby facilitates dynamic and flexible switching between rest and task-active states. Our results

support this interpretation and suggest that glucose metabolism in these networks is maintained at a high level in the resting-state, potentially to facilitate flexible switching to task-positive states when necessary.

The benefits of fPET methodology are twofold. Firstly, FDG uptake is neurophysiologically closer to the process of interest – i.e. neuronal function – than the BOLD response. It is well-known that BOLD-fMRI is an indirect index of neuronal activity that is not only ill-defined (e.g., Phillips et al., 2016), and non-quantitative (Logothetis, 2008), but is also confounded by physiological parameters (e.g., Ward et al., 2019). In contrast, most of the glucose uptake measured by FDG-PET is localised to the synapses (Sokoloff et al., 1977; Sokoloff, 1981; Magistretti and Allaman, 2015). Although technical considerations have led to the current situation where BOLD-fMRI is the more common method for visualising the spatiotemporal dynamics of neural function, FDG-PET should be considered as the gold standard for visualising neuronal activity. The second benefit of fPET is that the method provides information about the dynamic use of the primary energy source of the brain – glucose. Univariate snapshots of glucose hypometabolism are predictive of neurodegeneration (e.g., Jack et al., 2013), however preliminary evidence suggests that changes in the dynamic use of glucose over time may be an earlier and more powerful predictor of metabolic changes in neurodegenerative disease (Sanabria-Diaz et al., 2013). FDG-fPET methodology therefore represents an important development in characterising metabolic connectivity in the human brain and may show promise as a biomarker for disease in the future.

Metabolic covariance is a poor predictor of metabolic connectivity

FDG-sPET metabolic covariance was dominated by left-right homologue, fronto-parietal, temporo-subcortical and occipito-temporal covariance. This pattern is consistent with the earliest evidence of region-to-region correlation of the cerebral metabolic rate of glucose consumption (CMR_{GLC}) reported by Horwitz and colleagues over 30-yrs ago (e.g., (Horwitz et al., 1984; Azari et al., 1994). In their early work, Horwitz et al. noted the paucity of positive correlations between the occipito-temporal and fronto-parietal lobes; a finding which is evident in the present results where fronto-temporal and parieto-occipital edges demonstrated negative correlations. Our sPET metabolic covariance results are also consistent with more recent published findings. Using a static PET acquisition, Di and Biswal (2012) reported that independent component analysis (ICA) decomposition revealed resting-state networks predominantly between left-right homotopic regions, with few networks showing anterior-posterior connectivity. In their follow-up report, Di et al. (2017) again found that sPET covariance is dominated by left-right homotopic connections, and short-range connections within anatomical sub-divisions, with only a few fronto-parietal connections. Similarly, Savio et al., (2017) reported that sPET resting-state networks showed modest overlap with BOLD-fMRI networks, but were sparser and less well-separated from each other compared to BOLD networks (see also Passow et al., 2015). In summary, the static FDG-PET metabolic covariance results in the present study are consistent with the extensive literature of previous resting-state studies.

An examination of within-subject FDG-fPET metabolic connectivity in humans has not been previously reported. We therefore compared fPET connectivity to sPET covariance, since across-subject sPET covariance has (until the present study) been the established method for examining metabolic 'connectivity' in health and disease (Yakushev et al., 2017). On the basis of previous comparisons of across- versus within-subject haemodynamic connectivity (Roberts et al., 2016), we hypothesised that sPET covariance would be moderately correlated with fPET connectivity. Strikingly, we found that group-level FDG-sPET had minimal correlation with individual-level FDG-fPET metabolic connectivity. This was the case both throughout extended brain networks and within brain regions. This somewhat surprising finding is analogous to Simpson's Paradox (Simpson, 1951) whereby a statistical relationship observed at a population level is reversed at the level of the individuals that constitute the population (Kievit et al., 2013). Conceptually, Simpson's Paradox refers to a sign reversal in the statistical relationships obtained at the individual and population (or group) levels. Roberts and co-workers found evidence of Simpson's Paradox for a number of brain

regions in fMRI haemodynamic connectivity in the dorsal attention and default mode networks (Roberts et al., 2016). However, they found that the majority of brain regions did not show a full reversal of the correlation values, as most brain regions showed across-subject correlations in the same direction as the within-subject correlations. Roberts et al. therefore concluded that across-subject fMRI covariance is a reasonable predictor of within-subject fMRI connectivity.

While we did not directly test for Simpson's Paradox in our data, we found that group-level sPET covariance was not correlated with within-subject fPET connectivity. Simpson's Paradox is a special case of the *ecological fallacy* (Robinson, 1950), and the related concept of *ergodicity* (Molenaar, 2008; Kievit et al., 2013). Ergodic processes occur when a group-level result is generalisable to the individuals within the sample. In reality, ergodic processes are quite rare, because two strict criteria must be met: namely, the process must be homogeneous across individuals within a sample, and the statistical parameters that describe the process must be constant or stationary over time. Functional brain connectivity measured using neuroimaging data does not meet the criteria for ergodicity and thus group-level results are not generalisable to the individuals within the group (Fisher et al., 2018). Liegeois and colleagues have provided a complete mathematical examination of ergodicity in the context of BOLD-fMRI functional connectivity data (Liégeois et al., 2017). Due to technical limitations of the PET imaging method, existing studies of resting-state metabolic connectivity has been limited to examinations of group-level covariance rather than within-subject correlation of FDG-PET timeseries (e.g., Horwitz et al., 1984; Di et al., 2017; Savio et al., 2017). Researchers have attempted to draw quite strong conclusions on the basis of these results including attempts to use metabolic covariance as a biomarker for disease (Yakushev et al., 2017). Our results suggest that metabolic covariance cannot be used to predict within-subject connectivity, and that future attempts to explore metabolic connectivity as a biomarker for disease must necessarily use fPET as the only statistically valid approach.

FDG-fPET metabolic connectivity is correlated with BOLD-fMRI haemodynamic connectivity in fronto-parietal regions

FDG-fPET metabolic connectivity showed moderate similarity to BOLD-fMRI, particularly in superior cortex and frontoparietal regions. The similarity between fPET and fMRI was apparent when qualitatively comparing the network graphs and the distribution of the regions with higher degrees of connectivity, as well as the cross-modal correlations. These results complement the recent findings of Amend et al. (2019) who reported that FDG-fPET metabolic connectivity showed similarities to BOLD-fMRI haemodynamic connectivity in a rodent model. Our findings are also consistent with recent results of Shokri-Kojori et al. (2019) who found that frontoparietal regions had the highest concurrent energy use (based on CMR_{GLC}) and BOLD-fMRI activity, in comparison to the rest of the brain.

One puzzling finding was the low degree apparent in subcortical regions for both FDG-fPET and BOLD-fMRI and their cross-modal correlation. Indeed, this effect was also apparent for the FDG-sPET results. These observations are in contrast with the known high degree of inter-connectedness of the subcortical regions with the rest of the brain: the cortico-basal ganglia-cerebellar (Middleton and Strick, 2000; Bostan and Strick, 2010) and cortico-thalamic circuits (Parent and Hazrati, 1995). One possibility is that these results may be attributable to the reduced sensitivity of signal detection in midbrain areas for both PET and fMRI. BOLD-fMRI echo planar images show systematic image artefacts in midbrain areas (Limbrick-Oldfield et al., 2012) and susceptibility artefact at air-tissue interfaces (Bernstein et al., 2006). These effects are evident in the variability maps of raw echo planar images acquired in this study (Supplementary Figure 2). The PET images have lower signal detection in the midbrain, due to the higher attenuation of gamma ray emitted from deep brain regions. The attenuation correction corrects the signal loss but is unable to correct the lower signal to noise in the midbrain region of the images. We observed systematically lower signal to noise and high signal variability in brain regions near the ventricles (Supplementary Figure 2). The low inter-connectedness of the FDG-fPET and BOLD-fMRI measures in the subcortical regions may be attributable to the lower neuronal signals detected in these areas.

Direct comparisons between FDG-fPET and BOLD-fMRI are in part limited by the differences in temporal resolution and signal-to-noise ratio (SNR) of the two imaging modalities. The FDG-fPET protocol involves the administration of low levels of radioactivity over an extended time period, and results in PET images with lower signal and greater noise in comparison to BOLD-fMRI. Neural activity is optimally investigated using imaging techniques that have temporal resolutions in the order of milliseconds, which cannot be achieved by either fPET or fMRI. Nevertheless, the temporal resolution of conventional fMRI has been enormously useful for characterising distributed brain activity involving networks with multiple synapses over large-scale pathways in the brain. Importantly, although our understanding of integrative human brain networks has been developed largely from fMRI experimental results, the technique cannot serve as a 'ground-truth' for brain networks that are identifiable using FDG-fPET. In addition to the inherent methodological limitations of fMRI, the FDG-fPET and BOLD-fMRI experimental techniques detect different physiological targets; namely neural glucose metabolism and deoxygenation of haemoglobin in the cerebrovasculature. In comparison to the large body of literature of fMRI networks and brain connectivity, the evaluation of fPET resting-state networks is based on FDG-PET as a direct and quantitative index of neural metabolic activity without cerebrovascular confounds. Multi-modality imaging including simultaneous MR-PET has motivated important developments in PET detector technology, which have led to improved sensitivity for low signal detection methods including fPET (Chen et al., 2018). Future PET detector technologies may have the potential to further improve the detection sensitivity, which have the potential for the temporal resolution of functional PET to approach that of functional MRI.

Limitations and future directions

The current study has a number of limitations. A limitation of the FDG-fPET methodology is that the biosafety constraints on the dose of the administered radioactivity produces noisy PET images with low SNR. The differing SNR profiles between the sPET, fPET and fMRI modalities motivated the use of a percentile 'thresholding' approach, rather than a statistical threshold which would likely require different thresholds for each modality due to differences in statistical power. The percentile approach allowed comparison of the most highly connected regions between the modalities. The potential of the FDG-fPET technique is that it theoretically provides a more direct measure of neuronal activity than fMRI and does not have the susceptibility artefacts present in BOLD-fMRI. However, this advantage may not be realised in practice as the connectivity estimates in subcortical regions, which are some of the most interconnected regions in the human brain, showed low region degree in both FDG-fPET and BOLD-fMRI. Future work should aim to further characterise and improve the noise properties of the fPET signal, as well as optimise the acquisition and data preparation procedures. An additional technical limitation is that the processing pipeline for FDG-fPET data is immature in comparison to the pre-processing procedures for BOLD-fMRI data. As a nascent technology, FDG-fPET has not had the benefit of many years of work validating acquisition parameters, data preparation and signal detection optimisation. Work in these fields are presently improving the radiotracer administration (e.g., Jamadar et al., 2019a; Rischka et al., 2018), attenuation correction (Baran et al., 2018), motion correction (Chen et al., 2019) and data analysis (Li et al., 2019) techniques. Here, we provide information on the effects of spatiotemporal filtering and the stability of fPET connectivity over the scan period (refer to the Supplement), but further work is required.

An important distinction to make is that the brain connectivity reported here is not comparable to 'dynamic' connectivity as reported in the literature over the past few years. While the fPET metabolic connectivity measures are dynamic in the sense that they cross correlate the regional *time-courses* of FDG uptake over the scan period, the approach is nevertheless temporally-invariant (stationary) as the correlation is expected to be robust to temporal reordering of the measurement time-points. In other words, the Pearson r-values describing the metabolic connectivity would be unchanged by permuting the order of the PET images. Whilst other authors have used the term 'static' to refer to stationary measures of functional connectivity, we do not use the term 'static metabolic connectivity' to avoid confusion with our use of static PET to denote across-subject PET covariance (see Liégeois

et al., 2017 for discussion of static versus dynamic functional connectivity and the mathematical basis of stationarity in haemodynamic connectivity). Our approach to analyse stationary fPET metabolic connectivity was to compare metabolic connectivity with the most common method for estimation of functional connectivity in the BOLD-fMRI literature (Liégeois et al., 2017; Preti et al., 2017), which is the basis of our knowledge of integrative brain connectivity. However, truly 'dynamic' brain connectivity does take into account the temporal fluctuations of functional and metabolic connectivity, since the temporal ordering of the timeseries of brain images is important (Calhoun et al., 2014). Dynamic connectivity approaches include sliding-time window approaches and models of switching between microstates (Preti et al., 2017). Whilst it is important that future studies examine the dynamics of FDG-fPET metabolic connectivity, the current temporal resolution of fPET methodology will present a major challenge. Here, we report fPET images reconstructed with 16-sec acquisition duration whilst other groups have reported results using a 12-sec duration (Rischka et al., 2018). However, to our knowledge attempts to reconstruct fPET images for shorter durations have not yet been successful; we were unable to reliably detect resting state metabolic networks using 4-sec durations (unpublished data; see also Rischka et al., 2018 who reported fPET images using a 2-sec duration). Significant technical, methodological and analytical challenges remain to be solved before FDG-fPET will be used routinely to index dynamic metabolic connectivity.

Conclusion

Simultaneous neuroimaging studies that independently measure brain function are by nature challenging. Data from each imaging modality is optimally acquired using a distinctive experimental protocol with instrumentation that may corrupt the integrity of the other measure of brain function. Simultaneous FDG-PET/BOLD-fMRI represents a significant advance over previous approaches Our experimental approach to simultaneously image two mechanisms that underpin the dynamics of energy consumption in the brain via glucose metabolism and the cerebrovascular haemodynamic response, provides a unique opportunity to investigate the fundamental basis of human brain connectivity. In this study, we have reported the first within-subject FDG-fPET metabolic connectivity in humans, discovered that metabolic connectivity is correlated with BOLD-fMRI haemodynamic connectivity in the frontoparietal cortex, and demonstrated that it is poorly correlated with commonly used group-based FDG-PET metabolic covariance measures. Notably, these findings motivate and provide a basis for future metabolic brain connectivity studies across the human life span, and investigation as a novel biomarker for neurological and neurodegenerative diseases.

Acknowledgements

This work was supported by an Australian Research Council (ARC) Linkage Project (LP170100494) that includes financial support from Siemens Healthineers. SDJ, PGDW & GFE are supported by the ARC Centre of Excellence for Integrative Brain Function (CE140100007). SDJ is supported by an ARC Discovery Early Career Researcher Award (DE150100406) and National Health and Medical Research Council Fellowship (APP1174164).

Author Contributions

SDJ, PGDW & GFE designed the research question, PGDW, EXL & ERO conducted the analysis, SDJ wrote the first draft of the paper with GFE and PGDW edits. All authors contributed to manuscript preparation and review. GFE, SDJ, & ZC sourced funding for the work. The authors acknowledge Richard McIntyre, Alexandra Carey, Disha Sasan and Irene Graafsma from Monash Biomedical Imaging for their assistance in acquiring data.

The authors have no competing or conflicting interests.

Data availability

Data is available from the authors upon reasonable request.

References

Alderson, T.H., Bokde, A.L.W., Kelso, J.A.S., Maguire, L., Coyle, D., 2020. Metastable neural dynamics underlies cognitive performance across multiple behavioural paradigms. Hum Brain Mapp hbm.25009. https://doi.org/10.1002/hbm.25009

Amend, M., Ionescu, T.M., Di, X., Pichler, B.J., Biswal, B.B., Wehrl, H.F., 2019. Functional resting-state brain connectivity is accompanied by dynamic correlations of application-dependent [18F]FDG PET-tracer fluctuations. NeuroImage 196, 161–172. https://doi.org/10.1016/j.neuroimage.2019.04.034

Andrews-Hanna, J.R., Snyder, A.Z., Vincent, J.L., Lustig, C., Head, D., Raichle, M.E., Buckner, R.L., 2007. Disruption of Large-Scale Brain Systems in Advanced Aging. Neuron 56, 924–935. https://doi.org/10.1016/j.neuron.2007.10.038

Avants, B.B., Tustison, N.J., Song, G., Cook, P.A., Klein, A., Gee, J.C., 2011. A reproducible evaluation of ANTs similarity metric performance in brain image registration. NeuroImage 54, 2033–2044. https://doi.org/10.1016/j.neuroimage.2010.09.025

Azari, N.P., Horwitz, B., Pettigrew, K.D., Grady, C.L., Haxby, J.V., Giacometti, K.R., Schapiro, M.B., 1994. Abnormal Pattern of Cerebral Glucose Metabolic Rates Involving Language Areas in Young Adults with Down Syndrome. Brain and Language 46, 1–20. https://doi.org/10.1006/brln.1994.1001

Baran, J., Chen, Z., Sforazzini, F., Ferris, N., Jamadar, S., Schmitt, B., Faul, D., Shah, N.J., Cholewa, M., Egan, G.F., 2018. Accurate hybrid template–based and MR-based attenuation correction using UTE images for simultaneous PET/MR brain imaging applications. BMC Med Imaging 18, 41. https://doi.org/10.1186/s12880-018-0283-3

Bernstein, M.A., Huston, J., Ward, H.A., 2006. Imaging artifacts at 3.0T. J. Magn. Reson. Imaging 24, 735–746. https://doi.org/10.1002/jmri.20698

Biswal, B., Yetkin, F.Z., Haughton, V.M., Hyde, J.S., 1995. Functional connectivity in the motor cortex of resting human brain using echo-planar mri. Magnetic Resonance in Medicine 34, 537–541. https://doi.org/10.1002/mrm.1910340409

Bostan, A.C., Strick, P.L., 2010. The Cerebellum and Basal Ganglia are Interconnected. Neuropsychol Rev 20, 261–270. https://doi.org/10.1007/s11065-010-9143-9

Burgos, N., Cardoso, M.J., Thielemans, K., Modat, M., Pedemonte, S., Dickson, J., Barnes, A., Ahmed, R., Mahoney, C.J., Schott, J.M., Duncan, J.S., Atkinson, D., Arridge, S.R., Hutton, B.F., Ourselin, S., 2014. Attenuation Correction Synthesis for Hybrid PET-MR Scanners: Application to Brain Studies. IEEE Transactions on Medical Imaging 33, 2332–2341. https://doi.org/10.1109/TMI.2014.2340135

Buxton, R.B., Uludağ, K., Dubowitz, D.J., Liu, T.T., 2004. Modeling the hemodynamic response to brain activation. NeuroImage, Mathematics in Brain Imaging 23, S220–S233. https://doi.org/10.1016/j.neuroimage.2004.07.013

Caballero-Gaudes, C., Reynolds, R.C., 2017. Methods for cleaning the BOLD fMRI signal. NeuroImage 154, 128–149. https://doi.org/10.1016/j.neuroimage.2016.12.018

Calhoun, V.D., Miller, R., Pearlson, G., Adalı, T., 2014. The Chronnectome: Time-Varying Connectivity Networks as the Next Frontier in fMRI Data Discovery. Neuron 84, 262–274. https://doi.org/10.1016/j.neuron.2014.10.015

Carson, R.E., 2000. Pet physiological measurements using constant infusion. Nuclear Medicine and Biology 27, 657–660. https://doi.org/10.1016/S0969-8051(00)00138-4

Chen, Z., Jamadar, S.D., Li, S., Sforazzini, F., Baran, J., Ferris, N., Shah, N.J., Egan, G.F., 2018. From simultaneous to synergistic MR-PET brain imaging: A review of hybrid MR-PET imaging methodologies. Hum Brain Mapp 39, 5126–5144. https://doi.org/10.1002/hbm.24314

Chen, Z., Sforazzini, F., Baran, J., Close, T., Shah, N.J., Egan, G.F., 2019. MR-PET head motion correction based on co-registration of multicontrast MR images. Hum Brain Mapp. https://doi.org/10.1002/hbm.24497

Cox, R.W., 1996. AFNI: Software for Analysis and Visualization of Functional Magnetic Resonance Neuroimages. Computers and Biomedical Research 29, 162–173. https://doi.org/10.1006/cbmr.1996.0014

Di, X., Biswal, and Alzheimer's Disease Neu, B.B., 2012. Metabolic Brain Covariant Networks as Revealed by FDG-PET with Reference to Resting-State fMRI Networks. Brain Connectivity 2, 275–283. https://doi.org/10.1089/brain.2012.0086

Di, X., Gohel, S., Thielcke, A., Wehrl, H.F., Biswal, B.B., The Alzheimer's Disease Neuroimaging Initiative, 2017. Do all roads lead to Rome? A comparison of brain networks derived from inter-subject volumetric and metabolic covariance and moment-to-moment hemodynamic correlations in old individuals. Brain Struct Funct 222, 3833–3845. https://doi.org/10.1007/s00429-017-1438-7

Feinberg, D.A., Yacoub, E., 2012. The rapid development of high speed, resolution and precision in fMRI. NeuroImage 62, 720–725. https://doi.org/10.1016/j.neuroimage.2012.01.049

Fisher, A.J., Medaglia, J.D., Jeronimus, B.F., 2018. Lack of group-to-individual generalizability is a threat to human subjects research. Proc Natl Acad Sci USA 115, E6106–E6115. https://doi.org/10.1073/pnas.1711978115

Fornito, A., Bullmore, E.T., 2010. What can spontaneous fluctuations of the blood oxygenation-level-dependent signal tell us about psychiatric disorders? Curr Opin Psychiatry 23, 239–249. https://doi.org/10.1097/YCO.0b013e328337d78d

Fornito, A., Zalesky, A., Bassett, D.S., Meunier, D., Ellison-Wright, I., Yücel, M., Wood, S.J., Shaw, K., O'Connor, J., Nertney, D., Mowry, B.J., Pantelis, C., Bullmore, E.T., 2011. Genetic influences on cost-efficient organization of human cortical functional networks. J. Neurosci. 31, 3261–3270. https://doi.org/10.1523/JNEUROSCI.4858-10.2011

Fox, M.D., Raichle, M.E., 2007. Spontaneous fluctuations in brain activity observed with functional magnetic resonance imaging. Nature Reviews Neuroscience 8, 700–711. https://doi.org/10.1038/nrn2201

Fox, M.D., Snyder, A.Z., Vincent, J.L., Raichle, M.E., 2007. Intrinsic Fluctuations within Cortical Systems Account for Intertrial Variability in Human Behavior. Neuron 56, 171–184. https://doi.org/10.1016/j.neuron.2007.08.023

Garrity, A.G., Pearlson, G.D., McKiernan, K., Lloyd, D., Kiehl, K.A., Calhoun, V.D., 2007. Aberrant "default mode" functional connectivity in schizophrenia. Am J Psychiatry 164, 450–457. https://doi.org/10.1176/ajp.2007.164.3.450

Greicius, M.D., Srivastava, G., Reiss, A.L., Menon, V., 2004. Default-mode network activity distinguishes Alzheimer's disease from healthy aging: evidence from functional MRI. Proc. Natl. Acad. Sci. U.S.A. 101, 4637–4642. https://doi.org/10.1073/pnas.0308627101

Hahn, A., Breakspear, M., Rischka, L., Wadsak, W., Godbersen, G.M., Pichler, V., Michenthaler, P., Vanicek, T., Hacker, M., Kasper, S., Lanzenberger, R., Cocchi, L., 2020. Reconfiguration of functional brain networks and metabolic cost converge during task performance. eLife 9, e52443. https://doi.org/10.7554/eLife.52443

Hahn, A., Gryglewski, G., Nics, L., Hienert, M., Rischka, L., Vraka, C., Sigurdardottir, H., Vanicek, T., James, G.M., Seiger, R., Kautzky, A., Silberbauer, L., Wadsak, W., Mitterhauser, M., Hacker, M., Kasper, S., Lanzenberger, R., 2016. Quantification of Task-Specific Glucose Metabolism with Constant Infusion of 18F-FDG. J Nucl Med 57, 1933–1940. https://doi.org/10.2967/jnumed.116.176156

Hahn, A., Gryglewski, G., Nics, L., Rischka, L., Ganger, S., Sigurdardottir, H., Vraka, C., Silberbauer, L., Vanicek, T., Kautzky, A., Wadsak, W., Mitterhauser, M., Hartenbach, M., Hacker, M., Kasper, S., Lanzenberger, R., 2017. Task-relevant brain networks identified with simultaneous PET/MR imaging of metabolism and connectivity. Brain Struct Funct 1–10. https://doi.org/10.1007/s00429-017-1558-0

Hohenfeld, C., Werner, C.J., Reetz, K., 2018. Resting-state connectivity in neurodegenerative disorders: Is there potential for an imaging biomarker? NeuroImage: Clinical 18, 849–870. https://doi.org/10.1016/j.nicl.2018.03.013

Horwitz, B., Duara, R., Rapoport, S.I., 1984. Intercorrelations of Glucose Metabolic Rates between Brain Regions: Application to Healthy Males in a State of Reduced Sensory Input. J Cereb Blood Flow Metab 4, 484–499. https://doi.org/10.1038/jcbfm.1984.73

Jack, C.R., Knopman, D.S., Jagust, W.J., Petersen, R.C., Weiner, M.W., Aisen, P.S., Shaw, L.M., Vemuri, P., Wiste, H.J., Weigand, S.D., Lesnick, T.G., Pankratz, V.S., Donohue, M.C., Trojanowski, J.Q., 2013. Tracking pathophysiological processes in Alzheimer's disease: an updated hypothetical model of dynamic biomarkers. The Lancet Neurology 12, 207–216. https://doi.org/10.1016/S1474-4422(12)70291-0

Jamadar, S.D., Egan, G.F., Calhoun, V.D., Johnson, B., Fielding, J., 2016. Intrinsic Connectivity Provides the Baseline Framework for Variability in Motor Performance: A Multivariate Fusion Analysis of Low- and High-Frequency Resting-State Oscillations and Antisaccade Performance. Brain Connect 6, 505–517. https://doi.org/10.1089/brain.2015.0411

Jamadar, S.D., Sforazzini, F., Raniga, P., Ferris, N.J., Paton, B., Bailey, M.J., Brodtmann, A., Yates, P.A., Donnan, G.A., Ward, S.A., Woods, R.L., Storey, E., McNeil, J.J., Egan, G.F., 2018. Sexual Dimorphism of Resting-State Network Connectivity in Healthy Ageing. J Gerontol B Psychol Sci Soc Sci. https://doi.org/10.1093/geronb/gby004

Jamadar, S.D., Ward, P.G.D., Carey, A., McIntyre, R., Parkes, L., Sasan, D., Fallon, J., Orchard, E., Li, S., Chen, Z., Egan, G.F., 2019a. Radiotracer Administration for High Temporal Resolution Positron Emission Tomography of the Human Brain: Application to FDG-fPET. JoVE 60259. https://doi.org/10.3791/60259

Jamadar, S.D., Ward, P.G.D., Li, S., Sforazzini, F., Baran, J., Chen, Z., Egan, G.F., 2019b. Simultaneous task-based BOLD-fMRI and [18-F] FDG functional PET for measurement of neuronal metabolism in the human visual cortex. NeuroImage 189, 258–266. https://doi.org/10.1016/j.neuroimage.2019.01.003

Jenkinson, M., Bannister, P., Brady, M., Smith, S., 2002. Improved Optimization for the Robust and Accurate Linear Registration and Motion Correction of Brain Images. NeuroImage 17, 825–841. https://doi.org/10.1006/nimg.2002.1132

Jenkinson, M., Beckmann, C.F., Behrens, T.E.J., Woolrich, M.W., Smith, S.M., 2012. FSL. NeuroImage, 20 YEARS OF fMRI20 YEARS OF fMRI 62, 782–790. https://doi.org/10.1016/j.neuroimage.2011.09.015

Kievit, R.A., Frankenhuis, W.E., Waldorp, L.J., Borsboom, D., 2013. Simpson's paradox in psychological science: a practical guide. Front. Psychol. 4. https://doi.org/10.3389/fpsyg.2013.00513

Laird, A.R., Fox, P.M., Eickhoff, S.B., Turner, J.A., Ray, K.L., McKay, D.R., Glahn, D.C., Beckmann, C.F., Smith, S.M., Fox, P.T., 2011. Behavioral Interpretations of Intrinsic Connectivity Networks. Journal of Cognitive Neuroscience 23, 4022–4037. https://doi.org/10.1162/jocn_a_00077

Levin, J.M., Frederick, B. deB, Ross, M.H., Fox, J.F., von Rosenberg, H.L., Kaufman, M.J., Lange, N., Mendelson, J.H., Cohen, B.M., Renshaw, P.F., 2001. Influence of baseline hematocrit and hemodilution on BOLD fMRI activation. Magnetic Resonance Imaging 19, 1055–1062. https://doi.org/10.1016/S0730-725X(01)00460-X

Li, S., Jamadar, S.D., Ward, P.G.D., Premaratne, M., Egan, G.F., Chen, Z., 2019. Analysis of continuous infusion functional PET (fPET) in the human brain (preprint). Neuroscience. https://doi.org/10.1101/778357

Liégeois, R., Laumann, T.O., Snyder, A.Z., Zhou, J., Yeo, B.T.T., 2017. Interpreting temporal fluctuations in resting-state functional connectivity MRI. NeuroImage 163, 437–455. https://doi.org/10.1016/j.neuroimage.2017.09.012

Limbrick-Oldfield, E.H., Brooks, J.C.W., Wise, R.J.S., Padormo, F., Hajnal, J.V., Beckmann, C.F., Ungless, M.A., 2012. Identification and characterisation of midbrain nuclei using optimised

functional magnetic resonance imaging. NeuroImage 59, 1230–1238.
https://doi.org/10.1016/j.neuroimage.2011.08.016

Liu, T.T., 2017. Reprint of 'Noise contributions to the fMRI signal: An Overview".' NeuroImage,
Cleaning up the fMRI time series: Mitigating noise with advanced acquisition and correction
strategies 154, 4–14. https://doi.org/10.1016/j.neuroimage.2017.05.031

Logothetis, N.K., 2008. What we can do and what we cannot do with fMRI. Nature 453, 869–878.
https://doi.org/10.1038/nature06976

Magistretti, P.J., Allaman, I., 2015. A Cellular Perspective on Brain Energy Metabolism and
Functional Imaging. Neuron 86, 883–901. https://doi.org/10.1016/j.neuron.2015.03.035

Middleton, F., Strick, P., 2000. Basal ganglia and cerebellar loops: motor and cognitive circuits.
Brain Research Reviews 31, 236–250. https://doi.org/10.1016/S0165-0173(99)00040-5

Moeller, J.R., Strother, S.C., Sidtis, J.J., Rottenberg, D.A., 1987. Scaled Subprofile Model: A
Statistical Approach to the Analysis of Functional Patterns in Positron Emission
Tomographic Data. J Cereb Blood Flow Metab 7, 649–658.
https://doi.org/10.1038/jcbfm.1987.118

Molenaar, P.C.M., 2008. On the implications of the classical ergodic theorems: Analysis of
developmental processes has to focus on intra-individual variation. Dev. Psychobiol. 50,
60–69. https://doi.org/10.1002/dev.20262

Murphy, K., Fox, M.D., 2017. Towards a consensus regarding global signal regression for resting
state functional connectivity MRI. NeuroImage 154, 169–173.
https://doi.org/10.1016/j.neuroimage.2016.11.052

Parent, A., Hazrati, L.-N., 1995. Functional anatomy of the basal ganglia. I. The cortico-basal
ganglia-thalamo-cortical loop. Brain Research Reviews 20, 91–127.
https://doi.org/10.1016/0165-0173(94)00007-C

Passow, S., Specht, K., Adamsen, T.C., Biermann, M., Brekke, N., Craven, A.R., Ersland, L.,
Grüner, R., Kleven-Madsen, N., Kvernenes, O.-H., Schwarzlmüller, T., Olesen, R.A.,
Hugdahl, K., 2015. Default-mode network functional connectivity is closely related to
metabolic activity: Metabolic Activity and DMN Connectivity. Hum. Brain Mapp. 36, 2027–
2038. https://doi.org/10.1002/hbm.22753

Phillips, A.A., Chan, F.H., Zheng, M.M.Z., Krassioukov, A.V., Ainslie, P.N., 2016. Neurovascular
coupling in humans: Physiology, methodological advances and clinical implications. J
Cereb Blood Flow Metab 36, 647–664. https://doi.org/10.1177/0271678X15617954

Poudel, G., Egan, G.F., Churchyard, A.J., Chua, P., Stout, J.C., Georgiou-Karistianis, 2014.
Abnormal synchrony of resting state networks in premanifest and symptomatic Huntington
disease: the IMAGE-HD study. J Psychiatry Neurosci 39.
https://doi.org/10.1503/jpn.120226

Preti, M.G., Bolton, T.A., Van De Ville, D., 2017. The dynamic functional connectome: State-of-
the-art and perspectives. NeuroImage 160, 41–54.
https://doi.org/10.1016/j.neuroimage.2016.12.061

Rischka, L., Gryglewski, G., Pfaff, S., Vanicek, T., Hienert, M., Klöbl, M., Hartenbach, M., Haug,
A., Wadsak, W., Mitterhauser, M., Hacker, M., Kasper, S., Lanzenberger, R., Hahn, A.,
2018. Reduced task durations in functional PET imaging with [18F]FDG approaching that of
functional MRI. NeuroImage 181, 323–330.
https://doi.org/10.1016/j.neuroimage.2018.06.079

Roberts, R.P., Hach, S., Tippett, L.J., Addis, D.R., 2016. The Simpson's paradox and fMRI:
Similarities and differences between functional connectivity measures derived from within-
subject and across-subject correlations. NeuroImage 135, 1–15.
https://doi.org/10.1016/j.neuroimage.2016.04.028

Robinson, W.S., 1950. Ecological correlations and the behavior of individuals. International journal
of epidemiology 38, 337–341.

Sanabria-Diaz, G., Martínez-Montes, E., Melie-Garcia, L., the Alzheimer's Disease Neuroimaging
Initiative, 2013. Glucose Metabolism during Resting State Reveals Abnormal Brain

Networks Organization in the Alzheimer's Disease and Mild Cognitive Impairment. PLoS ONE 8, e68860. https://doi.org/10.1371/journal.pone.0068860

Sasaki, H., Kanno, I., Murakami, M., Shishido, F., Uemura, K., 1986. Tomographic Mapping of Kinetic Rate Constants in the Fluorodeoxyglucose Model Using Dynamic Positron Emission Tomography. J Cereb Blood Flow Metab 6, 447–454. https://doi.org/10.1038/jcbfm.1986.78

Savio, A., Fünger, S., Tahmasian, M., Rachakonda, S., Manoliu, A., Sorg, C., Grimmer, T., Calhoun, V., Drzezga, A., Riedl, V., Yakushev, I., 2017. Resting-State Networks as Simultaneously Measured with Functional MRI and PET. J Nucl Med 58, 1314–1317. https://doi.org/10.2967/jnumed.116.185835

Shokri-Kojori, E., Tomasi, D., Alipanahi, B., Wiers, C.E., Wang, G.-J., Volkow, N.D., 2019. Correspondence between cerebral glucose metabolism and BOLD reveals relative power and cost in human brain. Nature Communications 10, 690. https://doi.org/10.1038/s41467-019-08546-x

Simpson, E.H., 1951. The Interpretation of Interaction in Contingency Tables. Journal of the Royal Statistical Society. Series B (Methodological) 13, 238–241.

Smith, S.M., 2002. Fast robust automated brain extraction. Hum. Brain Mapp. 17, 143–155. https://doi.org/10.1002/hbm.10062

Sokoloff, L., 1981. The Deoxyglucose Method for The Measurement of Local Glucose Utilization and The Mapping of Local Functional Activity in The Central Nervous System, in: International Review of Neurobiology. Elsevier, pp. 287–333. https://doi.org/10.1016/S0074-7742(08)60296-2

Sokoloff, L., Reivich, M., Kennedy, C., Rosiers, M.H.D., Patlak, C.S., Pettigrew, K.D., Sakurada, O., Shinohara, M., 1977. The [^{14}C]Deoxyglucose Method For The Measurement Of Local Cerebral Glucose Utilization: Theory, Procedure, And Normal Values In The Conscious And Anesthetized Albino Rat. J Neurochem 28, 897–916. https://doi.org/10.1111/j.1471-4159.1977.tb10649.x

Srinivasan, R., Winter, W.R., Ding, J., Nunez, P.L., 2007. EEG and MEG coherence: Measures of functional connectivity at distinct spatial scales of neocortical dynamics. Journal of Neuroscience Methods 166, 41–52. https://doi.org/10.1016/j.jneumeth.2007.06.026

Thomas Yeo, B.T., Krienen, F.M., Sepulcre, J., Sabuncu, M.R., Lashkari, D., Hollinshead, M., Roffman, J.L., Smoller, J.W., Zöllei, L., Polimeni, J.R., Fischl, B., Liu, H., Buckner, R.L., 2011. The organization of the human cerebral cortex estimated by intrinsic functional connectivity. Journal of Neurophysiology 106, 1125–1165. https://doi.org/10.1152/jn.00338.2011

Tustison, N.J., Avants, B.B., Cook, P.A., Zheng, Y., Egan, A., Yushkevich, P.A., Gee, J.C., 2010. N4ITK: improved N3 bias correction. IEEE Trans Med Imaging 29, 1310–1320. https://doi.org/10.1109/TMI.2010.2046908

Vaishnavi, S.N., Vlassenko, A.G., Rundle, M.M., Snyder, A.Z., Mintun, M.A., Raichle, M.E., 2010. Regional aerobic glycolysis in the human brain. PNAS 107, 17757–17762. https://doi.org/10.1073/pnas.1010459107

Veronese, M., Moro, L., Arcolin, M., Dipasquale, O., Rizzo, G., Expert, P., Khan, W., Fisher, P.M., Svarer, C., Bertoldo, A., Howes, O., Turkheimer, F.E., 2019. Covariance statistics and network analysis of brain PET imaging studies. Scientific Reports 9, 2496. https://doi.org/10.1038/s41598-019-39005-8

Villien, M., Wey, H.-Y., Mandeville, J.B., Catana, C., Polimeni, J.R., Sander, C.Y., Zürcher, N.R., Chonde, D.B., Fowler, J.S., Rosen, B.R., Hooker, J.M., 2014. Dynamic Functional Imaging of Brain Glucose Utilization using fPET-FDG. Neuroimage 100, 192–199. https://doi.org/10.1016/j.neuroimage.2014.06.025

Ward, P.G.D., Orchard, E.R., Oldham, S., Arnatkevičiūtė, A., Sforazzini, F., Fornito, A., Egan, G.F., Jamadar, S.D., 2019. Individual differences in haemoglobin concentration influence BOLD fMRI functional connectivity and its correlation with cognition (preprint). Neuroscience. https://doi.org/10.1101/835660

Yakushev, I., Drzezga, A., Habeck, C., 2017. Metabolic connectivity. Current Opinion in Neurology 30, 677–685. https://doi.org/10.1097/WCO.0000000000000494

Figure Captions

Figure 1: A. Experimental paradigm. Participants underwent a 95-minute simultaneous MRI-PET scan. [18F]-FDG was infused over the course of the scan, with infusion start time locked to the onset of the PET scan (time 0). For the first 30-minutes, non-functional MRI scans were acquired to allow the PET signal to increase to detectable levels (see Figure 2, plasma radioactivity curve). At the 30-minute timepoint, 6 consecutive 10-minute resting-state BOLD-fMRI (T2* echo planar imaging) blocks were completed. Participants rested quietly with eyes open with central fixation (cross) for the full 60-minutes. Abbreviations: UTE, ultrashort echo time; MRS, magnetic resonance spectroscopy; GFM, gradient field map; pASL, pulsed arterial spin labelling; SWI, susceptibility weighted imaging; mins, minutes.

B. Plasma radioactivity curve. Individual subject radioactivity curves are plotted in grey, with the average across all subjects plotted in black.

Figure 2: A. FDG-sPET resting-state covariance matrix thresholded from -0.65 < r < 0.65. **B.** FDG-fPET resting-state connectivity matrix thresholded from 0 < r < 0.12. **C.** BOLD-fMRI resting-state connectivity matrix thresholded from 0 < r < 0.75. Supplementary Figure 1 shows the distribution of the number of edges across r values for each modality.

D. Spatial maps from three exemplar ROIs, i. the posterior cingulate, ii. anterior cingulate, and iii. left premotor cortex. These are spatial maps of the same data presented in the connectivity matrices above, and show the top 10% of regions correlated with the seed region (green).

Figure 3: Connectivity matrices (top panel) and network graphs (bottom panel) for eight canonical resting-state networks. Within-network connections lie on the diagonal (e.g. visual-visual), and between-network connections lie on the off-diagonal (e.g. visual-default mode). **A.** FDG-sPET metabolic covariance, **B.** FDG-fPET metabolic connectivity and **C.** BOLD-fMRI haemodynamic connectivity. Abbreviations: FP, frontoparietal network; DAN, dorsal attention network; VAN, ventral attention network; DMN, default mode network; SM, sensorimotor network; Lim, Limbic network; SC, subcortical network; Vis, visual network.

Figure 4: Spatial representation of the connection strength (region degree) in **A.** FDG-sPET, **B.** FDG-fPET and **C.** BOLD-fMRI.

Figure 5: Cross-modal correlation for (**A**, **C**) FDG-fPET and FDG-sPET, and (**B**, **D**) FDG-fPET and BOLD-fMRI for each of the 82 regions of interest. Boxplots are Tukey plots where the central line indicates the median, and the outer limits of the box represents the lower and upper quartiles.

a. Experimental Paradigm

fMRI & fPET Duration (60 mins)

30 mins

UTE, T1, T2, MRS, GFM,
pASL, SWI

10 — rest
10 — rest
10 — rest
10 — rest
10 — rest
10 — rest

Start
Infusion
PET acquisition

Start
fMRI acquisition
PET data retained from here

b. Subject-Level Plasma Radioactivity

Activity (kBq/cc)

Block #
1 2 3 4 5 6

Infusion start

Start fMRI acquisition
PET data retained

Time (Minutes)

a. FDG-sPET Covariance

b. FDG-fPET Connectivity

c. BOLD-fMRI Connectivity

d. Spatial Maps

i. Posterior Cingulate -2, -62, 28

ii. Anterior Cingulate -48, 6, 28

iii. Left Premotor -48, 6, 28

roi

99th

Percentile

90th

a. fPET-sPET

b. fPET-fMRI

c. fPET - sPET Correlation

d. fPET - fMRI Correlation

LEFT

Frontal

Parietal

Subcortical

Temporal

Occipital

RIGHT

Frontal

Parietal

Subcortical

Temporal

Occipital

Pearson r

-0.4   0   0.5

Metabolic and haemodynamic resting-state connectivity of the human brain: a high-temporal resolution simultaneous BOLD-fMRI and FDG-fPET multimodality study

Sharna D Jamadar Phillip GD Ward, Emma Xingwen Liang, Edwina R Orchard, Zhaolin Chen, Gary F Egan

~ Supplementary Material ~

Supplementary Figure 1: Distribution of edges across r values for each of the 3 modalities.

Supplementary Figure 2: Raw images for one individual subject showing signal intensity variation across the brain. Top panel shows mean image across a 10-minute run; middle panel shows standard deviation (Std Dev) across the same run; and lower panel shows the coefficient of variation (COV, standard deviation divided by the mean) across the run.

Supplementary Figure 3: Voxel weights for the spatiotemporal filter, depicted along the y-axis and the time axis. The filter is symmetrical in the x and z axis directions.

The spatiotemporal filter is defined as the convolution of a 3-dimensional gaussian filter in the spatial domain and a 1-dimensional gaussian filter in the time domain (Supplementary Figure 3). For the filter used in the main results of the paper, the standard deviation of the spatial gaussian was 1 voxel and of the temporal gaussian was 2 frames. The result of the spatiotemporal convolution was then modified to give negative weights on prior frames (negative time), and zero weights for the current frame.

We found highly consistent metabolic connectivity across a range of standard deviations, both temporal and spatial, up to 3 frames and 3 voxels, respectively. We selected the parameters which provided the least amount of smoothing whilst preserving the fidelity of the observed connectivity (Supplementary Figure 4).

The main trends we observed as we explored parameter values was an increase in correlation strength with higher spatial standard deviation, and an increase in mean connectivity with higher temporal standard deviation.

As the spatial domain is de-noising the volumes, the increased correlation strength can be expected for lower values. However, at higher values the filter may begin averaging signals across ROIs. For this reason, we truncated the spatio-temporal filter to a 7x7x7x7 spatio-temporal window to avoid artificial correlation between spatially adjacent ROIs.

In the temporal domain, high values mean estimating the gradient at a longer timescale. In the extreme, this approximates static connectivity as the difference between frames before the scan and after the scan would provide a similar estimate. It is possible the global increase in connectivity, yellow-hue in lower frames of Supplementary Figure 4, are beginning to be contaminated by these longer time-period effects.

Spatial Filter (Varying standard deviation)

Temporal Filter (Varying standard deviation)

Supplementary Figure 4: Variability of fPET connectivity matrix over a range of temporal (t) and spatial (s) filter widths.

Group-Level Per-Block fPET Connectivity

Supplementary Figure 5: Stability of fPET connectivity matrix over the six experimental blocks. Refer to Figure 1 (main document) for experimental design.