Title:

Schizophrenia-associated variation at *ZNF804A* correlates with

altered experience-dependent dynamics of sleep slow-waves and

spindles in healthy young adults

Authors:

Ullrich Bartsch*[1,2,#], Laura J Corbin*[3], Charlotte Hellmich[1], Michelle Taylor[3], Kayleigh E Easey[3],

Claire Durant[4], Hugh M Marston[2], Nicholas J Timpson *[3], Matthew W Jones*[1]

* These authors contributed equally.

Affiliations:

1) School of Physiology, Pharmacology & Neuroscience, University of Bristol, Bristol, UK

2) Translational Neuroscience, Eli Lilly & Co Ltd UK, Erl Wood Manor, Windlesham, UK

3) MRC Integrative Epidemiology Unit, University of Bristol, Bristol, UK

4) Clinical Research and Imaging Centre (CRIC), University of Bristol, Bristol, UK

current address: UK DRI Care Research & Technology, Surrey Sleep Research Centre,

University of Surrey, Clinical Research Building, Egerton Road, Guildford, Surrey, GU2 7XP

Corresponding authors:

n.j.timpson@bristol.ac.uk ; matt.jones@bristol.ac.uk

Key words:

psychosis, schizophrenia, genetics, sleep, slow wave, spindle, motor sequence task, ALSPAC

Short title:

ZNF804A genotype affects learning and sleep

Word count (abstract): 248

Word count (main body): 4221

5 Figures, 2 Tables, 1 Supplemental Information

ABSTRACT

Background

The rs1344706 polymorphism in *ZNF804A* is robustly associated with schizophrenia (SZ), yet

brain and behavioral phenotypes related to this variant have not been extensively

characterized. In turn, SZ is associated with abnormal non-rapid eye movement (NREM) sleep

neurophysiology. To examine whether rs1344706 is associated with intermediate

neurophysiological traits in the absence of disease, we assessed the relationship between

genotype, sleep neurophysiology, and sleep-dependent memory consolidation in healthy

participants.

Methods

We recruited healthy adult males, with no history of psychiatric disorder, from the Avon

Longitudinal Study of Parents and Children (ALSPAC) birth cohort. Participants were

homozygous for either the SZ-associated 'A' allele (N=25) or the alternative 'C' allele (N=22)

at rs1344706. Actigraphy, polysomnography (PSG) and a motor sequencing task (MST) were

used to characterize daily activity patterns, sleep neurophysiology and sleep-dependent

memory consolidation.

Results

Average MST learning and sleep-dependent performance improvements were similar across

genotype groups, but with increased variability in the AA group. CC participants showed

increased slow-wave and spindle amplitudes, plus augmented coupling of slow-wave activity

across recording electrodes after learning. Slow-waves and spindles in those with the AA

genotype were insensitive to learning, whilst slow-wave coherence decreased following MST

training.

Conclusion

We describe evidence that rs1344706 polymorphism in *ZNF804A* is associated with changes

in experience- and sleep-dependent, local and distributed neural network activity that

supports offline information processing during sleep in a healthy population. These findings

highlight the utility of sleep neurophysiology in mapping the impacts of SZ-associated variants

on neural circuit oscillations and function.

INTRODUCTION

Schizophrenia (SZ) is a debilitating psychiatric disorder with a lifetime prevalence of up to 4% (1). SZ etiology is complex and heterogenous, but an estimated heritability of up to 80% reflects critical genetic contributions to SZ liability (2,3). Recent efforts in cataloguing the genetic architecture of SZ have generated a list of over 100 loci thought to contribute in some way to the development of the disease (4,5). Despite most of these risk variants having small individual effects and acting in combination with other genetic and environmental factors, elucidating the neuronal changes downstream of genetic liability remains crucial for understanding normal brain development and the etiology of psychiatric disorders.

The single nucleotide polymorphism (SNP) rs1344706 within the second intron of *ZNF804*A was the first SNP to show genome-wide significant association for psychosis diagnosed in both bipolar disorder and SZ (6). This finding has been replicated in subsequent genome wide association studies (GWAS) (4,7–9) including a fine-mapping study which confirmed rs1344706 as the most strongly associated variant at the locus, with an OR for SZ of 1.10 [1.07 − 1.14] (10). *ZNF804A* is expressed in the brain and is predicted to encode a protein with a C2H2 zinc finger domain, indicating a role in transcriptional regulation (8,10) and thus likely complex biological functions (11). rs1344706 has been linked to a number of behavioral and neuronal phenotypes (12,13), correlating with altered neuroanatomy (14,15) (but see (16) for a null result), abnormal neurophysiology (17–19) and cognitive phenotypes (20–22). In particular, *ZNF804A* genotype has been associated with cortico-hippocampal functional connectivity in healthy control subjects (23,24) and also in SZ patients and their unaffected siblings (17,25).

Whilst cognitive deficits are an established feature of SZ (26,27), links have recently been

made between cognitive symptom dimensions and abnormal sleep. Sleep disturbances are a

core feature of SZ (28,29) and include increased sleep latency and decreased total sleep time,

independent of neuroleptic treatment (30). At the level of neural network activity, sleep in

patients also features changes in characteristic electroencephalography (EEG) oscillations,

particularly during NREM sleep. Thalamo-cortical spindle oscillations are a defining feature of

NREM and are reduced in patients with SZ (31–34). Consistent with the roles of spindle

oscillations in memory consolidation in healthy participants (35–38) spindle deficits in SZ have

been linked to cognitive deficits in patients (39,40). More recently, slow oscillations and their

coordination with spindles have also been implicated in contributing to deficits in sleep-

dependent memory consolidation in patients (41–43).

Overall, there is convergent evidence that circuit abnormalities in SZ are reflected by changes

in sleep physiology that, in turn, may be important for cognitive symptoms (44). In principle,

linking specific genetic variations with sleep neurophysiology phenotypes holds the promise

of illuminating a broader understanding of genetic effects and potential mechanisms of

neural circuit dysfunction in SZ. Here we used a recall-by-genotype approach (45) to recruit

healthy individuals who were homozygous at rs1344706 in order to reduce the issues of

confounding and reverse causality that commonly effect traditional observational

case/control studies. In this case, the availability of genetic data in a large and engaged cohort

study allowed for efficient and balanced recruitment of participants into a detailed

examination of sleep architecture and neurophysiology. We aimed to test the hypothesis

that, in the absence of disease, rs1344706 genotype would associate with the facets of

abnormal sleep neurophysiology and sleep-dependent memory consolidation seen in SZ.

METHODS AND MATERIALS

The study design and protocol was published in advance in (46); raw and processed data and metadata are available upon application to the Avon Longitudinal Study of Parents and Children (ALSPAC) Executive Committee through a standard application process (see http://www.bristol.ac.uk/alspac/researchers/access/).

Participants

Healthy males aged 21-23 years and of European ancestry were recruited from ALSPAC, a prospective birth cohort allowing the study of health and development across the life course (47,48). Participants were invited based on homozygosity either for the rs1344706 allele previously associated with increased liability for SZ (AA group), or for the alternative allele (CC group).

Ethical approval for the study was obtained from the ALSPAC Ethics and Law Committee (ref. 9224). The data collection protocol was previously approved by The University of Bristol Faculty of Science Human Research Ethics Committee as part of a pilot study (ref. 8089). All participants provided informed consent to participate in the study following the recommendations of the ALSPAC Ethics and Law Committee. For a detailed description of the cohort and recruitment see Supplemental Methods.

Procedures

Data were collected from each participant over approximately two weeks, beginning and ending with a night of polysomnography (PSG, including 9-channel EEG) at the Clinical Research and Imaging Centre at the University of Bristol (Figure 1). Both researchers and participants were blind to participant genotype throughout data collection. During visits, participants completed sleep-based questionnaires in order to assess self-rated sleep quality

and collect information about subjective experience of their night in the sleep laboratory (see

a detailed description of all procedures in Supplemental Procedures).

Analysis

Behavioral data

All paper questionnaires were manually scored and transcribed to spreadsheets. MST

performance was quantified using the number of correct sequences and the reaction time

during correct sequences. These variables were used to calculate average values for trial 10-

12 of the evening session ('training performance') and average values for trials 1-3 of the

morning session ('test performance'). Overnight improvement was calculated as the

percentage change in each outcome measure from training to test (49). Actigraphy data were

manually annotated in MotionWare (CamNtech, UK) to derive sleep architecture and

circadian rhythm measures.

Polysomnography data

Polysomnography was scored by an experienced expert (blinded to participant genotype)

based on AASM criteria (50) using REMLogic software (Natus Europe GmbH, Germany). Sleep

architecture was quantified using standard variables including total sleep time (TST) and sleep

onset latency (SOL) (see Polysomnography Analyses in Supplemental Methods). EEG traces

were analyzed using automatic detection of characteristic NREM sleep events - slow waves

(SW), delta waves, slow and fast spindle events as described earlier (43,51). Characteristic

NREM events were further characterized using multitaper spectra and coherence using the

Chronux toolbox (www.chronux.org).

Statistical methods

A detailed description of all analysis and statistical methods can be found in Supplemental

Methods; Tables S1, S2 show a full record of methods and their alignment to analytical

arguments. In brief, behavioral measures were analyzed either by a comparison of means

across groups (two-sample two-sided t-test or Wilcoxon rank-sum test) or by fitting a linear

mixed model with genotype and MST session (training versus testing) fitted as fixed effects

(using Stata v14.2 (52)). PSG-derived sleep architecture measures were analyzed in a linear

mixed model with genotype and recording night (night 1: baseline, night 2: learning) fitted as

fixed effects. PSG-derived event properties, power and coherence measures were compared

across genotype groups, electrodes, recording nights (night 1: baseline, night 2: learning) and

sleep stages (N2, N3) using a linear mixed model framework and a stepwise reduction

procedure implemented using the lme4 (53) and lmerTest (54) packages in R. We built a full

model of the general form [*y ~ genotype + night + electrode + sleep_stage + (genotype **

night) + (1|ID)], where y is any derived sleep variable, and then applied backward elimination

of non-significant model terms using the R function step which is part of the R package

lmerTest (54). Results presented are mean ± standard error (SE) unless stated otherwise.

RESULTS

Data were collected from 47 participants (25 AA and 22 CC). The two genotype groups did not

differ in maternal education, social class, psychosis-like symptoms at age 18, or in the

Wechsler Abbreviated Scale of Intelligence at age 15 (Table S3). Data from seven participants

were excluded due to missing or corrupted data (see Supplemental Results for further

explanation); we therefore present results for 40 participants (Figure S1).

Figure 1: Study design

The study included two study visits to the sleep lab at the Clinical Research & Imaging Centre in Bristol with two
weeks of actigraphy monitoring between visits. Participants visited the sleep lab for a baseline polysomnography
(PSG) recording night on their first visit (night 1). They were also issued an actigraphy watch which they wore
until the end of the second study visit. During the second visit, participants were trained on the motor sequence
task (MST) in the evening and tested in the morning, with an intervening second PSG recording (night 2).

Increased variability in Motor Sequence Task performance in the rs1344706 AA group

Overall performance levels for practice-dependent increases in the number of correct

sequences (NCS) – and corresponding decreases in button press latency within correct

sequences ('reaction time', RT) – were comparable between genotype groups. Figure 2,

panels A, D show the MST learning curves for both genotype groups and Figure 2, panels B, E

show the averages of the last 3 trials in the evening and first 3 trials in the morning that are

used to calculate overnight improvement. Participants in both groups improved overnight in

mean NCS (overnight change in NCS, CC: 16.9% with standard deviation (SD): 9.6, AA: 15.9%

SD: 16.8, Figure 2C, Table 1) and RT (overnight change in absolute RT, CC: 10.5% (SD: 6.2), AA:

8.3% (SD 11.9), Figure 2F, Table 1).

Linear mixed modelling of the MST performance data confirmed effects of session (training

vs. test) on NCS (session: $F_{(1, 39)}= 79.1$, $p = 6.38e-11$) and RT (session: $F_{(1, 39)}= 28.8$, $p =$

3.93e-06, Table 2), suggesting sleep-dependent consolidation of motor memory in both

genotype groups. There was no strong evidence for an effect of genotype on task

performance, but point estimates suggested that AA group participants produced fewer

correct sequences (F(1, 38) = 1.61, *p* = 0.21) and had slower reaction times (F (1, 38)= 3.0, *p* =

0.09, Table 2).

Interestingly, the AA group showed higher variance in overnight improvement in NCS (SD CC:

9.6, AA: 16.8, two-sample variance comparison *p* = 0.02, Table 1) and RT (SD CC: 6.2, AA: 11.9,

two-sample variance comparison *p* = 0.01). This higher variance was particularly pronounced

during the morning test session (SD NCS, CC: 3.6, AA: 5.5, Levene's test *p* = 0.02; SD RT, CC:

41 ms, AA: 62 ms, Levene's test *p* = 0.02).

Figure 2: Average MST learning curves and overnight improvement by genotype group

Black: CC group (N=18); Blue: AA group (N= 22).

A) MST learning curves showing the number of correct sequences per trial. Night 2 is indicated by a dark grey separator, the last 3 and first 3 trials used to calculate the average for the evening (eve) and morning (morn) performance are highlighted in light grey.

B) Box plot showing median number of correct sequences (last 3 trials in the evening v first 3 trials in the morning) for each MST session and genotype group. (Plots indicate the median, with boxes showing the 25th and 75th percentile of data, whiskers indicate the range of values inside 1.5* interquartile range, extreme values (outside 1.5 IQR) are plotted as individual data points, see Statistical Methods for details).

C) Boxplot showing the median of overnight improvement in number of correct sequences/30 s trial as percentage change from evening to morning performance.

D) Learning curves as in A but for the mean reaction time (RT, button press latency within a correct sequence) per trial.

E) Boxplot showing the median RT during correct sequence button presses (last 3 trials in the evening v first 3 trials in the morning) for each MST session and genotype group.

F) Boxplot of median overnight improvement in RT measured as absolute percentage change from evening to morning performance

Sleep timing, architecture and quality appear unaffected by rs1344706 genotype

We found no evidence for consistent effects of genotype on diurnal rhythmicity (Table S4), sleep architecture derived from the PSG (Figure 3 and Table S5) or subjective/objective sleep quality (Table S6).

| **Figure 3: Example hypnograms and sleep architecture variables from polysomnography**

| A) Hypnogram examples from a CC individual for nights 1 and 2, x- axis shows time in hours from lights off. Sleep

| stages are Wake (W), REM (R), NREM1 (N1, NREM2 (N2), NREM3 (N3).

| B) Hypnogram examples from an AA individual for nights 1 and 2, x- axis shows time in hours from lights off

| C-F) Boxplots of PSG derived variables of sleep architecture by genotype group (CC, black; AA, blue) and

| recording night. Outlier (outside the 25th and 75th percentile) are included as individual data points (o symbol).

| We observed no significant effects of night or genotype using linear mixed model approach for C) Total Sleep

| Time (min), D) Percentage of REM sleep, E) Percentage of stage 2 NREM sleep or F) Percentage of stage 3 NREM

| sleep.

rs1344706 genotype is associated with experience dependent changes in NREM sleep EEG
oscillations

To assess potential neurophysiological correlates of variance in MST performance, we used

custom detection algorithms to extract slow and delta waves (0.5-4Hz), and slow (9-12Hz)

and fast spindles (13-16Hz). Figure 4, panels A and B show wave-triggered averages, revealing

the morphologies of slow waves and fast spindles recorded at electrodes F3 and Cz:

waveforms appeared similar across genotype group and recording night at these electrode

locations. Panels C and D show average extracted amplitude values for each event at the

electrode locations F3 and Cz. (For a complementary spectral analysis of power in

corresponding frequency bands see Supplementary Results and Tables S7, S8.)

Slow-wave amplitudes depend on experience, but only in the CC group
We did not detect differences in density of slow-wave events between nights or genotypes

(Table S9). However, a linear mixed model analysis of slow-wave amplitudes from N2 and N3

sleep from all electrode locations suggested a main effect of night and an important

interaction term (night by genotype) in the initial full model. After stepwise reduction, the

night by genotype interaction remained (F(1, 1356.05) = 18.67, p= 1.67e-05, Table S9). Figure

4E shows the differences in estimated marginal means between nights, demonstrating an

increase in slow-wave amplitudes from night 1 (baseline) to night 2 (learning) in CC

participants (night 1: 100±4.80μV; night 2: 105.12±4.78μV, *p* <0.001, Table S10). In contrast,

slow-wave amplitudes in the AA group remained similar across nights (night 1: 112±4.34μV,

night 2: 111±4.34μV, n.s.). These SW event results were supported by very similar results for

delta wave event properties (Tables S11, S12) and separate Fourier analyses of 0.5-1.5Hz

power, which showed the same pattern of experience and genotype-dependent changes

(Tables S7 and S8). Collectively, these analyses suggest that the coordinated firing of cortical

populations during SW events may be modulated following learning in a genotype-dependent

manner.

Figure 4: Slow wave and spindle properties

A) SW wave triggered average at F3 for all genotype groups and both recording nights. CC, baseline night, grey,

CC, learning night, black, AA baseline night, light blue, AA learning night, dark blue. We found no significant

difference for any time bin (p<0.05, Wilcoxon ranksum test, no correction).

B) Spindle wave triggered average at Cz for all genotype groups and both recording nights, same colors as in A.

We found no significant difference for any time bin (p<0.05, Wilcoxon ranksum test, no correction).

C) Estimated marginal means for SW amplitude values for SW recorded at F3, during N3 sleep by genotype group

and by recording night.

D) Estimated marginal means of fast spindle amplitude values for spindles recorded at Cz, during N2 sleep for both genotypes and both recording nights.

E) Estimated marginal means differences for the factors genotype and night estimated from a linear mixed model analysis of all detected SW amplitudes (using the R package lmerTest (see **Supplemental Methods** for details)). CC individuals show an increase in SW amplitude in night 2, but AA's do not. Error bars indicate 95% confidence intervals.

F) Same as E for all detected fast spindle amplitudes. Like SW amplitudes, CC individuals show an increase in fast spindle amplitude in night 2, but AA's do not. Error bars indicate 95% confidence intervals.

Spindle properties depend on experience, with differential effects of genotype

A linear mixed model analysis of 9-12Hz (slow) spindle event properties revealed night and genotype dependent associations with amplitude (Table S13), with a trend for an increase after learning in the CC group (31.1±2.0 µV during night 1 *vs*. 31.7±2.0 µV during night 2, *p*=0.05), but a decrease in the AA group (from 33.8±1.8 µV to 33.2±1.8 µV, *p*=0.03, Table S14). We also observed a main effect for night-dependent associations with slow spindle frequency, with small decreases in slow spindle frequency after motor learning in both groups (from 11.36±0.05 Hz to 11.31±0.05 Hz in CC and from 11.33±0.04 Hz to 11.31±0.04 Hz in AA, p=0.005, Table S14).

We also detected a night by genotype interaction for fast spindle amplitude ($F_{(1, 1356.05)}$=10.24, *p*=0.001, Table S15), with differences in estimated marginal means shown in Figure 4F: amplitudes increased from night 1 to night 2, but again only in the CC group participants (from 31.8±2.0µV to 32.6±2.0µV, *p*=0.007, Table S16). Fast spindle frequency did not vary across nights or genotype, but fast spindle duration showed a similar pattern to fast spindle amplitude: a genotype by night interaction ($F_{(1, 1356.53)}$= 7.24, p= 0.0072), driven by shorter spindles in CC group during night 1 (795±8 ms vs. 819±8 ms in AA, *p*= 0.02, Table S16) and an increase in spindle length from night 1 to night 2 only in the CC group (from 795±8 ms to 800±8 ms, *p*= 0.03, Table S16). We found no strong evidence for an effect of genotype or

night on slow or fast spindle density (Table S13, 15). These fast spindle event-based analyses

were supported by spectral analyses of fast sigma power, which showed differential patterns

of experience-dependent changes between genotypes (Tables S7, S8).

To summarize these NREM EEG event analyses, only the CC genotype group showed slow-

wave and spindle properties – particularly event amplitudes – that were sensitive to

experience, sustaining increases on night 2 (post-MST learning) relative to night 1 (baseline).

It is possible, then, that attenuated experience-dependent changes in thalamocortical activity

contributed to more variable MST performance in the AA group. However, since recent work

has highlighted the importance of temporal interrelationships between these thalamocortical

oscillations for sleep-dependent memory consolidation (42,55–57), we next tested whether

slow-wave coordination also varied across nights and participants.

Slow wave mediated cortical connectivity during NREM sleep

We analyzed slow wave synchronization during NREM sleep for both genotype groups and

nights using multi-taper spectral coherence. Figure 5A illustrates all EEG electrode positions

and Figure 5B shows raw EEG traces surrounding a single SW event detected at electrode Fz.

To illustrate SW-associated temporal covariance in frontal and occipital EEG, we used Fz SW

events (trough times) as triggers to extract +/-2s windows of EEG surrounding each event

across both channels, averaging across all windows for each recording night. Figure 5C shows

Fz SW event triggered averages at Fz and O1 from one participant of the CC group. Here we

can see highly stereotypical SW events detected at Fz (with low variance) during both

recording nights, but different average waveforms at O1. During night 1, Fz SW coincide with

highly variable activity at O1, where a SW-like waveform is hardly separated from surrounding

background activity (Figure 5, C1); in contrast, during night 2 a distinct average SW waveform

coordinated with Fz, manifests at O1 (Figure 5, C2).

Figure 5, panels D1-4 show group-averaged coherograms for the electrode pair Fz-O1 for both

recording nights and genotype groups: the most coherent frequency ranges are 0.5-1.5 Hz

(SW) and fast spindle coherence (12-15 Hz). We used the average SW coherence (0.5-1.5Hz)

during 1s windows surrounding each SW for each electrode pair to construct a cortex-wide

SW connectivity matrix. Figure 5, E1-4 show matrices of group averaged coherence values for

both genotypes and both recording nights during N3 sleep. All matrices show a gradient of

coherence, with highest values between frontal and central electrodes and lowest values

between the most distant pairs, i.e. frontal and occipital electrodes.

We calculated the difference between all coherence values and plotted them in the same

matrix layout to illustrate differences in slow-wave coherence between nights (Figure 5, F1-

2) and genotypes (Figure 5, G1-2). Consistent differences in slow-wave coherence are

apparent between recording nights. Slow wave coherence increases in those in the CC group

from night 1 (baseline) to night 2 (learning, Figure 5, F1) - but a decrease in the AA group can

be seen in the difference matrices for this genotype group (Figure 5, F2). Slow-wave

coherence is higher in the AA group on night 1 (mainly frontal to occipital coupling between

electrodes F3, Fz, F4 to O1, O2), compared to the CC group (Figure 5, G1). Furthermore, a

linear mixed model with subsequent stepwise reduction revealed a genotype by night

interaction, indicating a differential effect of learning on slow-wave coherence in CC vs. AA

genotypes (genotype x night: $F(1, 11182) = 97.37$, $p<2e-16$, Table S17). Both genotypes show

changes in slow-wave coherence upon motor learning but a least squares estimation of group

marginal means reveals that those in the CC group show a post-learning increase in slow-

wave coherence (CC night 1 0.85 ± 0.021, CC night 2 0.87±0.019, p<0.001, Table S18),

whereas the AA group show a decrease in overall slow-wave coherence (genotype by night:

AA night 1 0.89±0.019 - AA night 2 0.86±0.021, p<0.001, Table S18) after learning. These

results are confirmed when using SW coherence of whole noise free NREM sleep epochs (see

Supplemental Results and Tables S19, S20). Thus, the CC group showed the increased SW

coherence predicted by previous studies (43,58), whereas SW coordination was attenuated

following learning in the AA participants.

| **Figure 5: Slow wave triggered coherence**

| A) A head diagram showing sleep EEG electrode positions.

| B) Raw data example showing all EEG traces surrounding a typical detected SW event at Fz.

C) SW triggered averages from one participant from band pass filtered (0.5-4 Hz) and EEG traces. Individual SW event traces were averaged in windows of +/-2 seconds with the SW trough time set t=0. C1) SW triggered average at Fz and O1 for 51 SW events detected during night 1 N2 sleep in one participant (CC, Night 1), C2) SW triggered averages as described in C1 using, 115 SW events during night 2 from the same participant ('CC', Night 2)

D) Average SW triggered coherograms. SW triggered data windows (SW trough time at Fz, t = 0) from the seed electrode (Fz) and target electrode (O1) were used to calculate multitaper coherograms for each data window pair. Coherograms were averaged for each participant and then averaged for each group and recording night. In the coherogram darker colors indicate higher coherence. Overlaid black traces are SW wave triggered averages for seed and target electrode (Fz, top and O1, bottom).

D1) Average coherogram for CC night1, D2) Average coherogram for CC night 2, D3) Average coherogram for AA night 1, D4) Average coherogram for AA night 2.

E) Average SW-triggered coherence values (0.5-1.5 Hz, -0.5 -0.5 s) for all electrode pairs for each genotype group during N2 sleep for both recording nights. E1) CC, during (baseline) night 1; E2) CC, during (learning) night 2, E3) AA, during (baseline) night 1, E4 AA, during (learning) night 2.

F) Differences in SW coherence between nights in each genotype group CC (F1), and AA (F2).

G) Differences between genotype groups on each night, night 1: CC-AA (G1), night 2: CC-AA (G2).

H) Estimated marginal means differences for the factors genotype and night estimated from a linear mixed model analysis of SW event coherence. CC individuals show an increase in SW coherence during night 2, but AA's show an overall decrease in coherence. Error bars indicate 95% confidence intervals.

DISCUSSION

We performed a recall-by-genotype study (45) to investigate the potential contributions of an SZ-associated SNP, rs1344706, to sleep-dependent memory processing and sleep neurophysiology in healthy volunteers. In summary, 1) all participants showed normal wake/sleep rhythms and there was no evidence for an effect of genotype on any variable describing macro sleep architecture; 2) we observed greater variance in learning and sleep-dependent memory consolidation following a motor task in AA participants homozygous for the variant associated with increased SZ liability; 3) we detected genotype- and learning-dependent effects on SW and fast spindle amplitudes, with the AA group showing attenuated

changes in SW and spindle amplitudes after learning; 4) the AA group failed to exhibit the

learning-dependent increase in SW coherence evident in the CC genotype group.

Behavior

Using a motor learning task previously shown to be impaired in SZ patients, we did find

evidence for greater variability in overnight improvement and other variables derived from

MST in those with the AA genotype at rs1344706, suggesting that rs1344706 may associate

with subtle changes motor learning and consolidation.

Motor learning (59) and its sleep-dependent memory consolidation are impaired in patients

diagnosed with SZ (60). The key brain areas involved in motor learning and its sleep

dependent consolidation, including the neocortex, striatum, thalamus, hippocampus and

cerebellum (61,61–64), have all been implicated in the etiology of SZ (65–67). *ZNF804A* has

been shown to be highly expressed in these brain regions, particularly the thalamus,

hippocampus and cortex (68), hence altered *ZNF804A* function may contribute to changes in

brain development and plasticity that influence motor learning and its consolidation (69).

Previous studies have shown that variability between individuals during early phases of

learning a motor task is higher in patients diagnosed with SZ compared to healthy controls

(70), potentially reflecting higher variability in brain anatomy or functional connectivity

patterns (71). Whether this variability and the associations of *ZNF804A* derive from

neurodevelopmental effects or altered adult neural plasticity remains an open question.

NREM sleep and neurophysiology

Although we did not observe consistent evidence for an effect of genotype on diurnal

rhythmicity or sleep architecture, detailed analyses of overnight EEG did unveil relationships

between rs1344706, corticothalamic activity during NREM sleep and neural correlates of

motor memory consolidation. We observed several interaction effects between genotype

and recording night, where NREM sleep activity appears to be differentially affected by the

acquisition of a motor task in AA and CC participants.

Spindle oscillations

Previous studies have shown that sleep dependent motor memory consolidation correlates

with spindle oscillations (56,72–75). Indeed, a substantial body of work has demonstrated

correlations between N2 sleep or spindles with motor memory in healthy participants (76–

79), although contradictory studies do exist (80,81). In particular, the individual contributions

of slow and fast spindles to memory consolidation are still debated.

We found some evidence supporting a role for slow spindle oscillations (9-12 Hz) in motor

memory consolidation, since slow spindle amplitudes appeared to be increased in CC

genotype participants during the learning night, whilst decreasing in the AA group. However,

we found no evidence of an effect of genotype or night on either slow or fast spindle density.

Despite fast spindle density being reduced in first episode (34,82,83) and chronically ill

patients (32,33,39) and their first-degree relatives (29,34,82), and the association of spindle

properties with polygenic risk scores for SZ (84), our results imply that rs1344706 alone does

not impact directly on fast spindle density. Given established polygenic effects in SZ, multiple

genetic variants and their interactions are likely to impact cortico-thalamic circuit

development. In particular, SNPs linked to ion channel genes like *CACAN1C* (84) are likely to

interact with other SNPs to impact corticothalamic development and maturation which might

have causal effects on cortico-thalamic oscillatory signatures or NREM sleep (40).

Slow oscillations

On average, slow-wave amplitudes increased during the sleep after learning only in the CC

genotype group. In addition, SW coherence appears to be differentially modulated after

learning between the genotype groups: those in the CC group show an increase in slow-wave

coherence, but participants with the AA genotype show a decrease in slow wave coherence

during the night that followed motor learning. Previous studies have shown that during early

sleep, after motor learning, slow-wave event amplitudes are locally increased in central and

parietal areas (85). Slow wave coherence has also been shown to increase during sleep after

a declarative memory task (58), and our recent work has demonstrated slow wave coherence

increased after motor learning in a control group, but not in patients diagnosed with SZ (43).

Our results in individuals homozygous for the 'A' allele at rs1344706 seem to be line with

these findings and provide a genetic correlate for slow wave phenotypes related to psychosis

and SZ.

Recent studies on the rodent homologue of *ZNF804A* suggest it has both a role during

development and in adult plasticity (69,86). Our own work in a rodent neurodevelopmental

model of SZ has demonstrated that interference in cortico-thalamic development causes

severe disruption of slow wave coordination between remote cortical areas and simultaneous

desynchronization of spindle and hippocampal ripple oscillations (51). Given the suggested

role of *ZNF804A* in cortical and thalamic development we speculate that rs1344706 may have

a role in corticothalamic development which itself would be related to impaired coordination

of SW activity during sleep. These deep characterizations of genotypic association motivate

future mechanistic studies in animal models that enable high-resolution phenotyping

corticothalamic circuit development and plasticity.

Limitations

Currently, we are only beginning to understand complex genotype-phenotype relationships

related to SZ. Quantifying brain activity and function directly related to identified neuronal

circuits in combination with a Recall-by-Genotype approach is critically advancing our

understanding of underlying biological mechanisms. Here we characterized the impact of the

rs1344706 variant on sleep neurophysiology but have not directly quantified the effect of the

variant on *ZNF804A* gene expression and/or function. Indeed, no consensus has been reached

on the function of *ZNF804A* itself yet, although a recent study suggests that *ZNF804A* may

regulate RNA synthesis (89). The rs1344706 variant may then reduce the expression of a

ZNF804A splice variant during prenatal development (87) leading to reduced spine density in

cortical excitatory neurons (69,88) - although the exact impact of rs1344706 on cortico-

thalamic development remains to be investigated. Moreover, recent studies show that the

impact of rs1344706 on brain function may depend on the carriage of other SZ-associated loci

(e.g. in genes *CACNA1C* (90) and *COMT* (91,92)) and environmental risk factors such as

cannabis use (93). Thus, the relationship between rs1344706 and its downstream

consequences is not easily determined. Furthermore, whilst the intention here was to

understand the role of *ZNF804A* in the absence of disease, further work is needed to show

that the phenotypes we observed here, are indeed relevant to SZ.

Finally, the small sample size and lack of replication within this study (due to the labor and

cost intensive nature of sleep laboratory experiments) naturally limit the strength of our

conclusions. Future work will look to improve the efficiency and reach of genetic sleep studies

by developing protocols that implement wearable technology to monitor sleep

neurophysiology at-home, over extended periods of time and in much larger samples.

Conclusions

Given the complex network of events linking genetics to brain-wide connectivity and function,

how can we best map genomic information to a neurobiological understanding of SZ? Here

we show that sleep neurophysiology presents a uniquely powerful opportunity to bridge

different levels of analysis: relating genotype to sleep-dependent physiology and

environmental factors such as learning, constitutes a rational, neurobiologically-informed

approach to delineating causal mechanisms of thalamocortical circuit dysfunction (44). Future

translational studies should investigate the influence of *ZNF804A* on slow wave and spindle

properties and their coordination in genetic rodent models and patient populations to further

elucidate genetic and circuit mechanisms of psychosis and their impacts on sleep, cognition

and novel therapies.

ACKNOWLEDGEMENTS

We are extremely grateful to all the families who took part in the ALSPAC study, the midwives for their help in recruiting them, and the whole ALSPAC team, which includes interviewers, computer and laboratory technicians, clerical workers, research scientists, volunteers, managers, receptionists and nurses. The UK Medical Research Council (MRC) and Wellcome (Grant ref: 217065/Z/19/Z) and the University of Bristol provide core support for ALSPAC. A comprehensive list of grants funding is available on the ALSPAC website (http://www.bristol.ac.uk/alspac/external/documents/grant-acknowledgements.pdf).

Specifically, GWAS data was generated by Sample Logistics and Genotyping Facilities at Wellcome Sanger Institute and LabCorp (Laboratory Corporation of America) using support from 23andMe; PLIKS data at age 18 was funded by MRC grant G0701503/85179 (PI: Stan Zammit); WASI at age 15 was funded by Wellcome Trust and MRC grant 076467/Z/05/Z (PI: George Davey Smith). This research was specifically funded by Elizabeth Blackwell Institute Catalyst Fund grant (ref: 48474) and the MRC Integrative Epidemiology Unit (IEU) [MC_UU_12013/3]. This publication is the work of the authors and NJ Timpson and MW Jones will serve as guarantors for the contents of this paper.

NJT is a Wellcome Trust Investigator [202802/Z/16/Z to NJT] and works within the University of Bristol National Institute for Health Research (NIHR) Biomedical Research Centre (BRC). LJC is supported by NJT's Wellcome Trust Investigator grant [202802/Z/16/Z]. NJT and LJC work in the MRC Integrative Epidemiology Unit (IEU) at the University of Bristol which is supported by the MRC [MC_UU_00011] and the University of Bristol. NJT is supported by the Cancer Research UK (CRUK) Integrative Cancer Epidemiology Programme [C18281/A19169]. CH was an academic foundation doctor supported by North Bristol NHS Trust during her time on the

project. KE was funded through a PhD studentship at the MRC Integrative Epidemiology Unit.

MT's time on the project was funded through an Elizabeth Blackwell Institute postgraduate

fellowship. CD's time on the project was funded through the Elizabeth Blackwell Institute

Catalyst Fund grant.

MWJ was awarded an MRC Senior Non-Clinical Research Fellowship (G1002064) that part-

funded this work. UB received a Lilly Innovation Fellowship Award through Eli Lilly & Co Ltd

UK.

DISCLOSURES

UB was a full-time contractor for Eli Lilly & Co Ltd UK. HMM was a full-time employee of Eli

Lilly & Co Ltd UK. The remaining authors have no financial disclosures or conflicts of interest.

REFERENCES

1. Bhugra D (2005): The Global Prevalence of Schizophrenia. *PLOS Medicine* 2: e151.

2. Sullivan PF, Kendler KS, Neale MC (2003): Schizophrenia as a Complex Trait: Evidence From a Meta-analysis of Twin Studies. *Arch Gen Psychiatry* 60: 1187–1192.

3. Lichtenstein P, Yip BH, Björk C, Pawitan Y, Cannon TD, Sullivan PF, Hultman CM (2009): Common genetic determinants of schizophrenia and bipolar disorder in Swedish families: a population-based study. *The Lancet* 373: 234–239.

4. Schizophrenia Working Group of the Psychiatric Genomics Consortium (2014): Biological insights from 108 schizophrenia-associated genetic loci. *Nature* 511: 421–427.

5. Pardiñas AF, Holmans P, Pocklington AJ, Escott-Price V, Ripke S, Carrera N, *et al.* (2018): Common schizophrenia alleles are enriched in mutation-intolerant genes and in regions under strong background selection. *Nature Genetics* 50: 381–389.

6. O'Donovan MC, Craddock N, Norton N, Williams H, Peirce T, Moskvina V, *et al.* (2008): Identification of loci associated with schizophrenia by genome-wide association and follow-up. *Nat Genet* 40: 1053–1055.

7. International Schizophrenia Consortium, Purcell SM, Wray NR, Stone JL, Visscher PM, O'Donovan MC, *et al.* (2009): Common polygenic variation contributes to risk of schizophrenia and bipolar disorder. *Nature* 460: 748–752.

8. Riley B, Thiselton D, Maher BS, Bigdeli T, Wormley B, McMichael GO, *et al.* (2010): Replication of association between schizophrenia and ZNF804A in the Irish Case-Control Study of Schizophrenia sample. *Mol Psychiatry* 15: 29–37.

9. Steinberg S, Mors O, Børglum AD, Gustafsson O, Werge T, Mortensen PB, *et al.* (2011): Expanding the range of ZNF804A variants conferring risk of psychosis. *Mol Psychiatry* 16: 59–66.

10. Williams HJ, Norton N, Dwyer S, Moskvina V, Nikolov I, Carroll L, *et al.* (2011): Fine mapping of ZNF804A and genome-wide significant evidence for its involvement in schizophrenia and bipolar disorder. *Mol Psychiatry* 16: 429–441.

11. Donohoe G, Morris DW, Corvin A (2010): The psychosis susceptibility gene ZNF804A: associations, functions, and phenotypes. *Schizophr Bull* 36: 904–909.

12. Sun Y, Hu D, Liang J, Bao Y-P, Meng S-Q, Lu L, Shi J (2015): Association between variants of zinc finger genes and psychiatric disorders: systematic review and meta-analysis. *Schizophr Res* 162: 124–137.

13. Chang H, Xiao X, Li M (2017): The schizophrenia risk gene *ZNF804A*: clinical associations, biological mechanisms and neuronal functions. *Molecular Psychiatry* 22: 944–953.

14. Lencz T, Szeszko PR, DeRosse P, Burdick KE, Bromet EJ, Bilder RM, Malhotra AK (2010): A Schizophrenia Risk Gene, ZNF804A, Influences Neuroanatomical and Neurocognitive Phenotypes. *Neuropsychopharmacology* 35: 2284–2291.

15. Ikuta T, Peters BD, Guha S, John M, Karlsgodt KH, Lencz T, *et al.* (2014): A schizophrenia
risk gene, ZNF804A, is associated with brain white matter microstructure.
Schizophrenia Research 155: 15–20.

16. Cousijn H, Rijpkema M, Harteveld A, Harrison PJ, Fernández G, Franke B, Arias-Vásquez A
(2012): Schizophrenia risk gene ZNF804A does not influence macroscopic brain
structure: an MRI study in 892 volunteers. *Mol Psychiatry* 17: 1155–1157.

17. Paulus FM, Krach S, Bedenbender J, Pyka M, Sommer J, Krug A, *et al.* (2013): Partial
support for ZNF804A genotype-dependent alterations in prefrontal connectivity. *Hum
Brain Mapp* 34: 304–313.

18. del Re EC, Bergen SE, Mesholam-Gately RI, Niznikiewicz MA, Goldstein JM, Woo TU, *et al.*
(2014): Analysis of schizophrenia-related genes and electrophysiological measures
reveals ZNF804A association with amplitude of P300b elicited by novel sounds. *Transl
Psychiatry* 4: e346.

19. O'Donoghue T, Morris DW, Fahey C, Da Costa A, Moore S, Cummings E, *et al.* (2014):
Effects of ZNF804A on auditory P300 response in schizophrenia. *Transl Psychiatry* 4:
e345.

20. Esslinger C, Kirsch P, Haddad L, Mier D, Sauer C, Erk S, *et al.* (2011): Cognitive state and
connectivity effects of the genome-wide significant psychosis variant in ZNF804A.
Neuroimage 54: 2514–2523.

21. Chen M, Xu Z, Zhai J, Bao X, Zhang Q, Gu H, *et al.* (2012): Evidence of IQ-modulated
association between ZNF804A gene polymorphism and cognitive function in
schizophrenia patients. *Neuropsychopharmacology* 37: 1572–1578.

22. Hargreaves A, Morris DW, Rose E, Fahey C, Moore S, Cummings E, *et al.* (2012): ZNF804A
and social cognition in patients with schizophrenia and healthy controls. *Molecular
Psychiatry* 17: 118–119.

23. Esslinger C, Walter H, Kirsch P, Erk S, Schnell K, Arnold C, *et al.* (2009): Neural mechanisms
of a genome-wide supported psychosis variant. *Science* 324: 605.

24. Cousijn H, Tunbridge EM, Rolinski M, Wallis G, Colclough GL, Woolrich MW, *et al.* (2015):
Modulation of hippocampal theta and hippocampal-prefrontal cortex function by a
schizophrenia risk gene. *Hum Brain Mapp* 36: 2387–2395.

25. Rasetti R, Sambataro F, Chen Q, Callicott JH, Mattay VS, Weinberger DR (2011): Altered
Cortical Network Dynamics: A Potential Intermediate Phenotype for Schizophrenia
and Association With ZNF804A. *Arch Gen Psychiatry* 68: 1207–1217.

26. Mohamed S, Paulsen JS, O'Leary D, Arndt S, Andreasen N (1999): Generalized cognitive
deficits in schizophrenia: A study of first-episode patients. *Arch Gen Psychiatry* 56:
749–754.

27. Green MF (2006): Cognitive Impairment and Functional Outcome in Schizophrenia and
Bipolar Disorder. *J Clin Psychiatry* 67: 3–8.

28. Monti JM, Monti D (2005): Sleep disturbance in schizophrenia. *International Review of
Psychiatry* 17: 247–253.

29. Cohrs S (2008): Sleep disturbances in patients with schizophrenia : impact and effect of
antipsychotics. *CNS Drugs* 22: 939–62.

30. Chouinard S, Poulin J, Stip E, Godbout R (2004): Sleep in Untreated Patients With
Schizophrenia: A Meta-Analysis. *Schizophrenia Bulletin* 30: 957–967.

31. Ferrarelli F, Tononi G (2017): Reduced sleep spindle activity point to a TRN-MD thalamus-
PFC circuit dysfunction in schizophrenia. *Schizophr Res* 180: 36–43.

32. Ferrarelli F, Huber R, Peterson MJ, Massimini M, Murphy M, Riedner BA, *et al.* (2007):
Reduced sleep spindle activity in schizophrenia patients. *Am J Psychiatry* 164: 483–
492.

33. Manoach DS, Thakkar KN, Stroynowski E, Ely A, McKinley SK, Wamsley E, *et al.* (2010):
Reduced overnight consolidation of procedural learning in chronic medicated
schizophrenia is related to specific sleep stages. *J Psychiatr Res* 44: 112–120.

34. Manoach DS, Demanuele C, Wamsley EJ, Vangel M, Montrose DM, Miewald J, *et al.*
(2014): Sleep spindle deficits in antipsychotic-naïve early course schizophrenia and in
non-psychotic first-degree relatives. *Front Hum Neurosci* 8.
https://doi.org/10.3389/fnhum.2014.00762

35. Schabus M, Gruber G, Parapatics S, Sauter C, Klösch G, Anderer P, *et al.* (2004): Sleep
spindles and their significance for declarative memory consolidation. *Sleep* 8: 27.

36. Fogel SM, Smith CT (2006): Learning-dependent changes in sleep spindles and Stage 2
sleep. *Journal of Sleep Research* 15: 250–255.

37. Tamaki M, Matsuoka T, Nittono H, Hori T (2008): Fast Sleep Spindle (13-15 Hz) Activity
Correlates with Sleep-Dependent Improvement in Visuomotor Performance. *Sleep* 31:
204–211.

38. Genzel L, Kroes MCW, Dresler M, Battaglia FP (2014): Light sleep versus slow wave sleep
in memory consolidation: a question of global versus local processes? *Trends in
Neurosciences* 37: 10–19.

39. Wamsley EJ, Tucker MA, Shinn AK, Ono KE, McKinley SK, Ely AV, *et al.* (2012): Reduced
sleep spindles and spindle coherence in schizophrenia: mechanisms of impaired
memory consolidation? *Biol Psychiatry* 71: 154–161.

40. Manoach DS, Pan JQ, Purcell SM, Stickgold R (2016): Reduced Sleep Spindles in
Schizophrenia: A Treatable Endophenotype That Links Risk Genes to Impaired
Cognition? *Biological Psychiatry* 80: 599–608.

41. Ramakrishnan M, Sartory G, van Beekum A, Lohrmann T, Pietrowsky R (2012): Sleep-
related cognitive function and the K-complex in schizophrenia. *Behavioural Brain
Research* 234: 161–166.

42. Demanuele C, Bartsch U, Baran B, Khan S, Vangel MG, Cox R, *et al.* (2017): Coordination
of Slow Waves With Sleep Spindles Predicts Sleep-Dependent Memory Consolidation
in Schizophrenia. *Sleep* 40: zsw013.

43. Bartsch U, Simpkin AJ, Demanuele C, Wamsley E, Marston HM, Jones MW (2019):
Distributed slow-wave dynamics during sleep predict memory consolidation and its
impairment in schizophrenia. *npj Schizophr* 5: 1–11.

44. Gardner RJ, Kersanté F, Jones MW, Bartsch U (2014): Neural oscillations during non-rapid
eye movement sleep as biomarkers of circuit dysfunction in schizophrenia. *Eur J*
Neurosci 39: 1091–1106.

45. Corbin LJ, Tan VY, Hughes DA, Wade KH, Paul DS, Tansey KE, *et al.* (2018): Formalising
recall by genotype as an efficient approach to detailed phenotyping and causal
inference. *Nature Communications* 9: 711.

46. Hellmich C, Durant C, Jones MW, Timpson NJ, Bartsch U, Corbin LJ (2015): Genetics, sleep
and memory: a recall-by-genotype study of ZNF804A variants and sleep
neurophysiology. *BMC Medical Genetics* 16: 96.

47. Boyd A, Golding J, Macleod J, Lawlor DA, Fraser A, Henderson J, *et al.* (2013): Cohort
Profile: The 'Children of the 90s'—the index offspring of the Avon Longitudinal Study
of Parents and Children. *Int J Epidemiol* 42: 111–127.

48. Fraser A, Macdonald-Wallis C, Tilling K, Boyd A, Golding J, Davey Smith G, *et al.* (2013):
Cohort Profile: the Avon Longitudinal Study of Parents and Children: ALSPAC mothers
cohort. *Int J Epidemiol* 42: 97–110.

49. Karni A, Meyer G, Rey-Hipolito C, Jezzard P, Adams MM, Turner R, Ungerleider LG (1998):
The acquisition of skilled motor performance: Fast and slow experience-driven
changes in primary motor cortex. *PNAS* 95: 861–868.

50. Silber MH, Ancoli-Israel S, Bonnet MH, Chokroverty S, Grigg-Damberger MM, Hirshkowitz
M, *et al.* (2007): The visual scoring of sleep in adults. *J Clin Sleep Med* 3: 121–131.

51. Phillips KG, Bartsch U, McCarthy AP, Edgar DM, Tricklebank MD, Wafford KA, Jones MW
(2012): Decoupling of sleep-dependent cortical and hippocampal interactions in a
neurodevelopmental model of schizophrenia. *Neuron* 76: 526–533.

52. StataCorp (2017): *Stata Statistical Software: Release 15*, version 15. College Station, TX:
StataCorp LLC.

53. Bates D, Mächler M, Bolker B, Walker S (2015): Fitting Linear Mixed-Effects Models Using
lme4. *Journal of Statistical Software* 67. https://doi.org/10.18637/jss.v067.i01

54. Kuznetsova A, Brockhoff PB, Christensen RHB (2017): lmerTest Package: Tests in Linear
Mixed Effects Models. *Journal of Statistical Software* 82.
https://doi.org/10.18637/jss.v082.i13

55. Latchoumane C-FV, Ngo H-VV, Born J, Shin H-S (2017): Thalamic Spindles Promote
Memory Formation during Sleep through Triple Phase-Locking of Cortical, Thalamic,
and Hippocampal Rhythms. *Neuron* 95: 424-435.e6.

56. Mölle M, Eschenko O, Gais S, Sara SJ, Born J (2009): The influence of learning on sleep
slow oscillations and associated spindles and ripples in humans and rats. *European*
Journal of Neuroscience 29: 1071–1081.

57. Helfrich RF, Mander BA, Jagust WJ, Knight RT, Walker MP (2018): Old Brains Come
Uncoupled in Sleep: Slow Wave-Spindle Synchrony, Brain Atrophy, and Forgetting.
Neuron 97: 221-230.e4.

58. Mölle M, Marshall L, Gais S, Born J (2004): Learning increases human
electroencephalographic coherence during subsequent slow sleep oscillations. *PNAS*
101: 13963–13968.

59. Walther S, Strik W (2012): Motor Symptoms and Schizophrenia. *NPS* 66: 77–92.

60. Manoach DS, Cain MS, Vangel MG, Khurana A, Goff DC, Stickgold R (2004): A failure of
sleep-dependent procedural learning in chronic, medicated schizophrenia. *Biological*
Psychiatry 56: 951–956.

61. Doyon J, Song AW, Karni A, Lalonde F, Adams MM, Ungerleider LG (2002): Experience-
dependent changes in cerebellar contributions to motor sequence learning. *PNAS* 99:
1017–1022.

62. Albouy G, Sterpenich V, Balteau E, Vandewalle G, Desseilles M, Dang-Vu T, *et al.* (2008):
Both the Hippocampus and Striatum Are Involved in Consolidation of Motor Sequence
Memory. *Neuron* 58: 261–272.

63. Heitger MH, Ronsse R, Dhollander T, Dupont P, Caeyenberghs K, Swinnen SP (2012): Motor
learning-induced changes in functional brain connectivity as revealed by means of
graph-theoretical network analysis. *NeuroImage* 61: 633–650.

64. King BR, Hoedlmoser K, Hirschauer F, Dolfen N, Albouy G (2017): Sleeping on the motor
engram: The multifaceted nature of sleep-related motor memory consolidation.
Neuroscience & Biobehavioral Reviews 80: 1–22.

65. Andreasen NC, Pierson R (2008): The Role of the Cerebellum in Schizophrenia. *Biological*
Psychiatry 64: 81–88.

66. Frantseva MV, Fitzgerald PB, Chen R, Möller B, Daigle M, Daskalakis ZJ (2008): Evidence
for Impaired Long-Term Potentiation in Schizophrenia and Its Relationship to Motor
Skill Leaning. *Cereb Cortex* 18: 990–996.

67. Lisman JE, Pi HJ, Zhang Y, Otmakhova NA (2010): A Thalamo-Hippocampal-Ventral
Tegmental Area Loop May Produce the Positive Feedback that Underlies the Psychotic
Break in Schizophrenia. *Biological Psychiatry* 68: 17–24.

68. Chang EH, Kirtley A, Chandon T-SS, Borger P, Husain-Krautter S, Vingtdeux V, Malhotra AK
(2015): Postnatal neurodevelopmental expression and glutamate-dependent
regulation of the ZNF804A rodent homologue. *Schizophrenia Research* 168: 402–410.

69. Deans PJM, Raval P, Sellers KJ, Gatford NJF, Halai S, Duarte RRR, *et al.* (2017): Psychosis
Risk Candidate ZNF804A Localizes to Synapses and Regulates Neurite Formation and
Dendritic Spine Structure. *Biol Psychiatry* 82: 49–61.

70. Adini Y, Bonneh YS, Komm S, Deutsch L, Israeli D (2015): The time course and
characteristics of procedural learning in schizophrenia patients and healthy
individuals. *Front Hum Neurosci* 9. https://doi.org/10.3389/fnhum.2015.00475

71. Mastrandrea R, Piras F, Gabrielli A, Caldarelli G, Spalletta G, Gili T (2019): Brain Network
Topology Maps the Dysfunctional Substrate of Cognitive Processes in Schizophrenia.
arXiv:190108521 [cond-mat, physics:physics, q-bio]. Retrieved July 23, 2019, from
http://arxiv.org/abs/1901.08521

72. Gais S, Mölle M, Helms K, Born J (2002): Learning-Dependent Increases in Sleep Spindle
Density. *J Neurosci* 22: 6830–6834.

73. Mölle M, Bergmann TO, Marshall L, Born J (2011): Fast and Slow Spindles during the Sleep
Slow Oscillation: Disparate Coalescence and Engagement in Memory Processing. *Sleep*
34: 1411–1421.

74. Cox R, Hofman WF, Talamini LM (2012): Involvement of spindles in memory consolidation
is slow wave sleep-specific. *Learn Mem* 19: 264–267.

75. Boutin A, Pinsard B, Boré A, Carrier J, Fogel SM, Doyon J (2018): Transient synchronization
of hippocampo-striato-thalamo-cortical networks during sleep spindle oscillations
induces motor memory consolidation. *NeuroImage* 169: 419–430.

76. Walker MP, Brakefield T, Morgan A, Hobson JA, Stickgold R (2002): Practice with Sleep
Makes Perfect: Sleep-Dependent Motor Skill Learning. *Neuron* 35: 205–211.

77. Morin A, Doyon J, Dostie V, Barakat M, Hadj Tahar A, Korman M, *et al.* (2008): Motor
sequence learning increases sleep spindles and fast frequencies in post-training sleep.
Sleep 31: 1149–1156.

78. Peters KR, Ray L, Smith V, Smith C (2008): Changes in the density of stage 2 sleep spindles
following motor learning in young and older adults. *J Sleep Res* 17: 23–33.

79. Tamaki M, Matsuoka T, Nittono H, Hori T (2009): Activation of fast sleep spindles at the
premotor cortex and parietal areas contributes to motor learning: A study using
sLORETA. *Clinical Neurophysiology* 120: 878–886.

80. Nemeth D, Janacsek K, Londe Z, Ullman MT, Howard DV, Howard JH (2009): Sleep has no
critical role in implicit motor sequence learning in young and old adults. *Exp Brain Res*
201: 351–358.

81. Pan SC, Rickard TC (2015): Sleep and motor learning: Is there room for consolidation?
Psychol Bull 141: 812–834.

82. Schilling C, Schlipf M, Spietzack S, Rausch F, Eisenacher S, Englisch S, *et al.* (2017): Fast
sleep spindle reduction in schizophrenia and healthy first-degree relatives: association
with impaired cognitive function and potential intermediate phenotype. *Eur Arch*
Psychiatry Clin Neurosci 267: 213–224.

83. Tesler N, Gerstenberg M, Franscini M, Jenni OG, Walitza S, Huber R (2015): Reduced sleep
spindle density in early onset schizophrenia: A preliminary finding. *Schizophrenia*
Research 166: 355–357.

84. Merikanto I, Utge S, Lahti J, Kuula L, Makkonen T, Lahti-Pulkkinen M, *et al.* (2019): Genetic
risk factors for schizophrenia associate with sleep spindle activity in healthy
adolescents. *Journal of Sleep Research* 28: e12762.

85. Huber R, Felice Ghilardi M, Massimini M, Tononi G (2004): Local sleep and learning. *Nature*
430: 78–81.

86. Hinna KH, Rich K, Fex-Svenningsen Å, Benedikz E (2015): The Rat Homolog of the
Schizophrenia Susceptibility Gene ZNF804A Is Highly Expressed during Brain
Development, Particularly in Growth Cones. *PLOS ONE* 10: e0132456.

87. Tao R, Cousijn H, Jaffe AE, Burnet PWJ, Edwards F, Eastwood SL, *et al.* (2014): Expression
of ZNF804A in human brain and alterations in schizophrenia, bipolar disorder, and
major depressive disorder: a novel transcript fetally regulated by the psychosis risk
variant rs1344706. *JAMA Psychiatry* 71: 1112–1120.

88. Zhou D, Xiao X, Li M (2020): The schizophrenia risk isoform ZNF804AE3E4 affects dendritic
spine. *Schizophrenia Research*. https://doi.org/10.1016/j.schres.2019.12.038

89. Chapman RM, Tinsley CL, Hill MJ, Forrest MP, Tansey KE, Pardiñas AF, *et al.* (2019):
Convergent Evidence That ZNF804A Is a Regulator of Pre-messenger RNA Processing
and Gene Expression. *Schizophr Bull* 45: 1267–1278.

90. Tecelão D, Mendes A, Martins D, Fu C, Chaddock CA, Picchioni MM, *et al.* (2019): The
effect of psychosis associated CACNA1C, and its epistasis with ZNF804A, on brain
function. *Genes, Brain and Behavior* 18: e12510.

91. Xu Q, Xiong Y, Yuan C, Liu F, Zhao F, Shen J, *et al.* (2018): ZNF804A rs1344706 interacts
with COMT rs4680 to affect prefrontal volume in healthy adults. *Brain Imaging and
Behavior* 12: 13–19.

92. Cui L, Wang F, Chang M, Yin Z, Fan G, Song Y, *et al.* (2019): Spontaneous Regional Brain
Activity in Healthy Individuals is Nonlinearly Modulated by the Interaction of ZNF804A
rs1344706 and COMT rs4680 Polymorphisms. *Neurosci Bull* 35: 735–742.

93. Soler J, Arias B, Moya J, Ibáñez MI, Ortet G, Fañanás L, Fatjó-Vilas M (2019): The interaction
between the ZNF804A gene and cannabis use on the risk of psychosis in a non-clinical
sample. *Progress in Neuro-Psychopharmacology and Biological Psychiatry* 89: 174–
180.