Constitutive activation of cellular immunity underlies the evolution of resistance to infection

Alexandre B. Leitão[*], Ramesh Arunkumar[*], Jonathan P. Day, Emma M. Geldman, Francis M. Jiggins

Department of Genetics, University of Cambridge, Cambridge, United Kingdom

* contributed equally

Abstract: Organisms rely on inducible and constitutive immune defences to

combat infection. Constitutive immunity enables a rapid response to infection

but may carry a cost for uninfected individuals, leading to the prediction that it

will be favoured when infection rates are high. When we exposed populations of

Drosophila melanogaster to intense parasitism by the parasitoid wasp

Leptopilina boulardi, they evolved resistance by developing a more reactive

cellular immune response. Using single-cell RNA sequencing, we found that

immune-inducible genes had become constitutively upregulated. This was the

result of resistant larvae differentiating precursors of specialized immune cells

called lamellocytes that were previously only produced after infection.

Therefore, populations evolved resistance by genetically hard-wiring an induced

immune response to become constitutive.

One Sentence Summary: *Drosophila* populations that frequently encounter

parasites have a constitutively activated immune response.

Main Text: A fundamental division within immune systems is between induced

and constitutive immune defences. Constitutive immunity is always active, even

in the absence of infection. For example, uninfected individuals produce natural

antibodies (*1*), immune cells and antimicrobial peptides (AMPs) (*2*). In contrast,

induced immune responses are triggered after infection, and frequently involve

amplifying existing constitutive factors. This can include immune cells

proliferating and differentiating, and the massive upregulation of AMPs and

cyotoxic molecules (*2*). Constitutive immunity is fast, allowing hosts to avoid

becoming infected altogether or rapidly clear infections. However, immune

defences can be costly, diverting limited resources from other fitness-related

traits (*3*) or damaging the host through immunopathology (*4*). While induced

immune responses only incur these costs after infection, they are always present

for constitutive immunity, even in the absence of infection. This trade-off

between speed and cost is predicted to govern the allocation of host investment

into constitutive and induced immunity (*5*, *6*). Theoretical models predict that

the key parameter determining which type of defence should be favoured is the

frequency that hosts are exposed to infection—more frequent exposure favours

constitutive defences (*5*). In support of this, bacteria tend to evolve phage

resistance by altering surface receptors when exposure is high and rely on

induced CRISPR-Cas immunity when exposure is low (*7*). However, to our

knowledge the key prediction that high rates of parasitism lead to induced

immune responses becoming constitutively expressed has not been

demonstrated.

To investigate these questions, we examined how the constitutive and induced cellular immune defences of *Drosophila melanogaster* evolve under high parasite pressure. Cellular immunity in *D. melanogaster* involves blood cells called hemocytes, which play the equivalent role to leukocytes in vertebrates. In uninfected larvae there are two morphological classes of hemocytes. Plasmatocytes constitute the majority of cells and have diverse functions including phagocytosis and AMP production (*8*). Crystal cells are a less abundant specialized cell type that produces the prophenoloxidase molecules PPO1 and PPO2, which are processed into enzymes required for the melanization of parasites (*9*). Infection can trigger an induced response, where these cell types proliferate, and cells called lamellocytes differentiate from plasmatocytes and prohemocytes in the lymph gland (*8*). Lamellocytes are large flat cells with a specialised role in killing large parasites like parasitic wasps (parasitoids). When a parasitoid injects its egg into *D. melanogaster* larvae, plasmatocytes adhere to the egg. Then, as lamellocytes differentiate they create additional cellular layers known as a 'capsule'. Finally, the capsule is melanised by the activity of phenoloxidases produced in crystal cells and lamellocytes (*9*). The melanin physically encases the parasitoid and toxic by-products of the melanisation reaction likely contribute to parasite killing (*10*).

There is substantial genetic variation in susceptibility to parasitoid infection within *D. melanogaster* populations (*11*), and this is associated with differences in hemocyte composition (*12*, *13*). Here, we used a combination of experimental evolution with physiological assays and single-cell RNA sequencing to examine how a high rate of parasitism by the parasitoid wasp *L. boulardi* affects constitutive and induced immune defences.

The evolution of resistance is associated with a more reactive immune

response

To investigate the evolution of the cellular immune response, we allowed

genetically diverse populations of *D. melanogaster* to evolve under intense levels

of parasitism. We established an outbred population from 377 wild-caught *D.*

melanogaster females, and used this to found six experimental populations that

were maintained at population sizes of 200 flies. In three of these populations,

larvae were infected every generation, and flies that survived by encapsulating

and melanising the wasp were used to establish the next generation (High

Parasitism populations). The other three populations were maintained in the

same conditions but without infection (No Parasitism populations). In line with

previous studies (*12*, *14*), the High Parasitism populations increased in

resistance over the generations (Fig. 1A, Selection regime x Generation: $\chi^2 =$

91.41, d.f.= 1, $p < 10^{-15}$). After 33 generations, a mean of 76% of flies survived

infection in the High Parasitism populations, compared to 10% in the No

Parasitism controls.

As populations evolved resistance, they melanised parasitoids at a higher rate

(Fig. 1A). This could result from increased basal reactivity of the encapsulation

immune response, or because the flies are evolving to escape the effects of

immunosuppressive molecules produced by the parasite. To distinguish these

possibilities, we measured the ability of each population to react to an inert

object. When a small volume of paraffin oil is injected into the larval haemocoel,

it remains as a sphere and can be encapsulated (*15*). Populations that evolved

without parasitism show a very limited reaction to the oil droplets (Fig. 1B). In

contrast, approximately half of larvae from selected populations encapsulate the

oil droplet (Fig. 1B; Selection Regime: $F_{1,4}=24.53$, $p=0.008$). Therefore, high levels

of parasitism led to the cellular immune system evolving a more reactive

encapsulation response.

Figure 1 Encapsulation rate during selection for parasitoid resistance. (A) The

proportion of infected larvae that gave rise to adult flies with visible capsules. An

outbred population of *D. melanogaster* (source, grey square) was used to found six

populations that were parasitized every generation with *L. boulardi* (blue) or maintained

without infection (grey). (B) The proportion of larvae encapsulating oil droplets at

generation 33. Dots represent the proportion of larvae in replicate populations and bars

represent the mean per selection regime.

Constitutive upregulation of immune-induced genes

Underlying induced immune responses is a transcriptional response to infection,

so we examined how the hemocyte transcriptome altered in populations that

evolved under high levels of parasitism. Initially, we pooled the data from

scRNA-seq to investigate global changes in hemocyte gene expression. As

expected, 48 hours post infection there was a strong transcriptional response to

infection (Fig. 2A, X axis; Data S1). Strikingly similar transcriptional changes

occurred in uninfected flies that had evolved under high rates of parasitism for

26 generations (Fig. 2A, Y axis; Data S1). Taking the 173 genes that had at least a

50% change in transcript levels after infection or selection, in every case the

direction of change was the same after infection and adaptation to high

parasitism (Fig. 2A; Data S1). We verified that the evolved response mimicked

the induced response by using quantitative PCR (qPCR) to measure the

expression of the lamellocyte marker *atilla* and the anti-parasitoid effector gene

PPO3 (Fig. S1). Again, we found that high rates of parasitism led to these

immune-induced genes becoming constitutively upregulated. Therefore, as

populations evolved resistance, the induced transcriptional response to infection

has become genetically hard-wired and constitutive.

We took the same approach to examine how the induced transcriptional

response to infection changed as populations evolved resistance. We found that

the transcriptional response to infection involved the same genes in the different

populations (Fig. 2B). However, the magnitude of the changes in gene expression

was substantially less in populations that had evolved under high parasite

pressure (Fig. 2B). Therefore, the increase in constitutive gene expression has

been matched by a decrease in induced expression. Again, this was verified by

qPCR (Fig. S1). Combining the constitutive and induced changes in gene

expression, the hemocyte transcriptome after infection is similar in the

132  populations that evolved under high levels of parasitism and the controls (Fig.

133  2C). Therefore, an induced response has become genetically fixed, as opposed to

134  selection leading to a greater total response.

135  **Figure 2. Changes in gene expression following selection for resistance and**

136  **parasitoid infection.** (A) Constitutive gene expression following 26 generations of

137  selection for resistance to the parasitoid wasp *L. boulardi* (y-axis). (B) The induced

138  change in gene expression following infection with *L. boulardi* in populations selected for

139  resistance (y-axis). (C) The combined induced and constitutive change in gene

140  expression following selection for resistance and infection (y-axis). All are compared to

141  the change in gene expression following infection in populations maintained without

142  parasitoid infection (x-axis). The black diagonal indicates the 1:1 line. The genes *atilla*

143  and *PPO3* are highlighted. Colour represents the density of overlain points. Relative

144  expression is expressed as $\log_2$(fold change).

146 **scRNA-seq reveals novel hemocyte types**

147 The global changes in hemocyte transcriptomes could result from a change in the

148 relative proportion of cell types in the hemocyte population, or from a shift in

149 gene expression within existing cell types. The three classical *Drosophila*

150 hemocyte types were originally defined morphologically, but emerging single

151 cell transcriptomic data suggests that *Drosophila* hemocyte populations are more

complex than this (*16–19*). Excluding cells that did not express the pan-hemocyte

markers *Hemese* or *Serpent* (*20*, *21*)*,* we grouped 19,344 larval hemocytes into

nine clusters based on their transcriptomes (Fig. 3A). Neither infection nor

selection leads to the appearance of new cell states (Fig. 3B).

We identified two large clusters of plasmatocytes. The largest was PLASM1,

which we classify as the cycling plasmatocyte state as it had the greatest number

of replicating cells in the G2M or S phase (Fig. S2). The 5,712 cells in this cluster

had high expression of genes associated with plasmatocytes (*Col4a1*, *Pxn*, *eater*,

Hml, *NimC1* -(*21*)) and actively replicating cells (*stg* – (*22*), *polo* – (*23*)and *scra* –

(*24*)) (Fig. S3). The next largest cluster, PLASM2, contained 2,874 plasmatocytes

that were enriched for the genes encoding rRNA (Fig. S4) or involved in the

ribosome biogenesis (Data S3). There were two smaller plasmatocyte clusters.

The AMP cluster contained 196 cells with elevated expression of genes encoding

antimicrobial and antifungal peptides, including *Drs*, *Mtk* and *CecA1* (Lemaitre

and Hoffmann 2007, Fig. S4; Data S3). The MET cluster contained 283 cells with

high expression of the heat shock proteins *Hsp24* and *Hsp27* and genes involved

in organic acid metabolism (Fig. S4; Data S3). The crystal cells formed a single

cluster of 906 cells (Fig. S6). Three clusters of cells, LAM 1, 2 and 3, had elevated

expression of one or more lamellocyte marker genes (*21*) (Fig. S5) and contained

1,260, 5,104 and 3,000 cells respectively.

Figure 3. Cell states of *Drosophila melanogaster* larval hemocytes. Two-dimensional

UMAP projections and cell state classification. (A) All 19,344 cells combined. PLASM:

plasmatocytes; MET: metabolic plasmatocyte cluster; AMP: antimicrobial peptide

plasmatocyte cluster; LAM: lamellocytes; CC: crystal cells. (B) Cells from populations that

had undergone 26 generations of selection with no parasitism (top panels) or high

parasitism (bottom panels), and were either uninfected (left panels) or 48 hours post

infection (right panels). 64-333 million reads were sequenced in each of the eight 10X

libraries that we generated (Data S2). The number of cells detected per library ranged

from 787 to 3,463 (Fig. S6).

Inducible cell states are present constitutively in resistant populations

Lamellocytes are key effector cells in the anti-parasitoid immune response. In

populations that evolved without exposure to parasitoids, infection resulted in a

large increase in the number of cells in all three lamellocyte clusters (LAM1-3,

Fig. 3B and 4A). In populations that evolved under high levels of parasitism, this

inducible response become constitutive, with uninfected larvae having 3-5 fold

more cells in LAM1 and 2 (Fig. 4A). However, this was matched by a reduction in

the induced response, so the final number of cells in these clusters after infection

was similar regardless of whether the populations had adapted to high or low

parasitism rates (Fig. 4A). These results held when replicate populations were

analyzed separately (Fig. S7A, B) or when the number of cells in each library

were down-sampled to match the library with the least number of cells (Fig.

S7C). Therefore, high infection rates led to the inducible differentiation of

specialised immune cells becoming genetically fixed.

To investigate the developmental origins of this change, we reconstructed the

pathway by which plasmatocytes differentiate into mature lamellocytes. By

subclustering the cells and ordering them along a pseudotemporal scale, we

found a single lineage starting from the self-cycling plasmatocytes (PLASM1) and

ending in LAM3 (Figs. 4B). Along this lineage there was a decrease in cell

replication, with the largest fraction of cells in the G2M and S phase in PLASM1

and the largest fraction in G1 in LAM3 (Fig. S8). Importantly, controlling for the

effects of cell cycle had a negligible impact on the clustering (Table S1) and the

trajectory of lamellocyte differentiation (Fig. S9). The expression of the effector

gene *PPO3,* the lamellocyte marker *atilla* and other marker genes all increased as

cells progressed along this trajectory (Fig. 4C, 4D and S10, Data S4). On the other

hand, genes involved in extracellular matrix organization—a house-keeping

function of plasmatocytes (*25*, *26*)—were down-regulated as cells differentiated

(Data S5). Together, these results suggest that LAM1 cells are immature

lamellocytes, while LAM3 cells are mature lamellocytes. Comparing our

populations, the inducible production of LAM1 cells has become entirely

constitutive in the populations evolved under high rates of parasitism, while

much of the LAM3 response remains inducible. This suggests that high infection

rates result in the cellular immune response becoming 'primed' by constitutively

producing the precursors of the cells required for parasite killing.

Figure 4. Changes in cell state following infection and adaptation to high rates of

parasitism. (A) Proportion of different cell states after infection and selection. (B)

Trajectory of lamellocyte differentiation. Plasmatocyte progenitors and lamellocytes

were subclustered, the trajectories inferred from multi-dimensional principle

components analysis, and the lineage projected onto the two-dimensional UMAP plot. (C-

D) Log$_2$ expression levels of *atilla* and *PPO3*. Relative expression levels were estimated

from normalized and scaled unique molecular identifier counts.

Morphological differentiation of immune cells remains an inducible response

Lamellocytes are classically defined morphologically as large flattened cells. As expected, the proportion of morphologically identifiable lamellocytes increased after infection (Figs 5A and Fig. S11A; Treatment: χ^2= 48.14, d.f= 1, p=3.97x10^{-12}). However, in contrast to our analysis of single cell transcriptomes, there was little difference between selection regimes in either the induced or constitutive proportion of these cells (Figs 5A and Fig. S11A; Selection regime x Treatment: χ^2= 0.16, d.f= 1, p=0.69; Selection regime: χ^2= 0.41, d.f= 1, p=0.52). This suggests that while the transcriptional state of cells has been constitutively activated, the change in cell morphology remains an inducible response. To test this hypothesis, we used microscopy to measure both cell morphology and, as a marker of the transcriptional state of the cells, *atilla* expression (Fig. S12). In line with our scRNAseq data, both infection and adaptation to high parasitism rates increases the proportion of *atilla* positive cells (Fig. 5C; Selection Regime x Treatment: F=16.3, d.f=1,8, p=0.004). However, among the *atilla* positive cells, infection had a large effect on cell size (Fig. 5C; Treatment: F=31.8, d.f.=1,7, p=0.0008), while selection had a small effect (Fig. 5C; Selection Regime: F=7.2, d.f.=1,7, p=0.03). Therefore, while selection has resulted in the constitutive transcriptional activation of cells, changes in cell morphology remain largely an inducible response.

Figure 5. Changes in the morphology and number of cells following infection and adaptation to high parasitism rates. (A) The proportion hemocytes that were morphologically identified as lamellocytes and (B) the concentration of total circulating hemocytes in no infection conditions (grey outline) and 48 hours post infection (red outlines). Samples were collected from populations that evolved 34 generation with high parasitism (blue bars) or no parasitism (grey bars) (C) Cell size and *atilla* expression. The expression of *atilla* was measured as the fluorescence intensity resulting from immunohistochemistry staining. Cell size was measured along the major axis. Replicated populations from each selection regime are represented by different symbols and the proportion of cells in each quarter of the plot is shown.

In addition to the change in the transcriptional state of cells, we found that the populations maintained under strong parasite pressure evolved a constitutive increase in the absolute number of hemocytes in circulation (Fig. 5B and Fig. S11B, Selection regime: χ^2= 22.37, d.f= 1, p=2.25x10^{-6}). This supports results from previous selection experiments (*13*, *27*) and comparative analyses (*28*), and is another aspect of the induced response that has become constitutive.

Conclusions

Our results provide support for the hypothesis that natural selection favours

constitutive defences when infection is common (*5–7*) by demonstrating that an

inducible immune response becomes constitutive when populations frequently

encounter a parasite. The induced immune response to parasitoids relies on the

differentiation of lamellocytes from embryonic-derived plasmatocytes, lymph

gland prohemocytes (*8*) and lymph gland plasmatocytes (*18*). Evidence for

immature lamellocytes has previously come from patterns of marker gene

expression (*29*, *30*) and recent scRNAseq data (*17*, *19*). Our results show that

Drosophila populations evolve to produce these cells constitutively if they are

frequently exposed to infection. However, this change in the transcriptional state

of cells was not accompanied by morphological differentiation. Genes involved in

cytoskeleton dynamics such as *βTub60D* and *shot* are among the genes with the

greatest increases in expression as lamellocytes differentiate (Fig. S10, Data S4)

and these may be required for the final step of changing cell shape (*31*).

Together, our results suggest populations that adapt to high parasitism rates

have their cellular immune defences in a state of readiness, with the final step of

cell differentiation remaining an inducible response.

Evolving resistance to parasitoid wasps is costly in *Drosophila* (*12*, *14*, *32*), so

inducible immunity may be an adaptation that allows uninfected individuals to

avoid this cost under low parasitism conditions. The physiological mechanism

that causes the cost is unknown. It is possible that the differentiation of

immature lamellocytes diverts resources from other fitness related traits.

Lamellocyte differentiation is energy-demanding and results in the increased

expression of glycolysis genes in hemocytes after infection (*3*). This cost may be

borne by uninfected flies in our resistant populations, as we found that genes

involved in trehalose uptake and catabolism (*33*), *Trehalase* and *Tret1-1*, are

upregulated by both infection and selection (Fig. S13). Another possibility is that

a more reactive cellular immune response may increase the effects of

immunopathology.

After infection, whether the host or parasitoid survives is a race between the

parasitoid development and the formation of a cellular capsule by the immune

system (*34*). Constitutive differentiation of immature lamellocytes may allow

lamellocytes to be produced more rapidly upon infection. Furthermore, cytotoxic

molecules could be more readily available. This is supported by an increase in

the constitutive expression of *PPO3* in populations adapted to high rates of

parasitism. *PPO3* encodes a intracellular prophenoloxidase that, when active, is

responsible for the melanization of lamellocytes (*9*). Along with the increase in

the total number of hemocytes, the constitutive activation of the cellular immune

response likely explains the greatly increased resistance of populations adapted

to high rates of parasitism.

References and Notes:
311

1. H. P. Savage, V. M. Yenson, S. S. Sawhney, B. J. Mousseau, F. E. Lund, N.

Baumgarth, Blimp-1-dependent and -independent natural antibody

production by B-1 and B-1-derived plasma cells. *J. Exp. Med.* **214**, 2777–

2794 (2017).

2.   B. Lemaitre, J. Hoffmann, The host defense of Drosophila melanogaster.

Annu. Rev. Immunol. **25**, 697–743 (2007).

3.   A. Bajgar, K. Kucerova, L. Jonatova, A. Tomcala, I. Schneedorferova, J.

Okrouhlik, T. Dolezal, Extracellular Adenosine Mediates a Systemic

Metabolic Switch during Immune Response. *PLoS Biol.* **13**, 1–23 (2015).

4.   A. L. Graham, J. E. Allen, A. F. Read, Evolutionary causes and consequences

of immunopathology. *Annu. Rev. Ecol. Evol. Syst.* **36**, 373–397 (2005).

5.   E. Shudo, Y. Iwasa, Inducible defense against pathogens and parasites:

Optimal choice among multiple options. *J. Theor. Biol.* **209**, 233–247

(2001).

6.   M. Boots, A. Best, The evolution of constitutive and induced defences to

infectious disease. *Proc. R. Soc. B Biol. Sci.* **285** (2018),

doi:10.1098/rspb.2018.0658.

7.   E. R. Westra, S. Van houte, S. Oyesiku-Blakemore, B. Makin, J. M.

Broniewski, A. Best, J. Bondy-Denomy, A. Davidson, M. Boots, A. Buckling,

Parasite exposure drives selective evolution of constitutive versus

inducible defense. *Curr. Biol.* **25**, 1043–1049 (2015).

8.   V. Honti, G. Csordás, É. Kurucz, R. Márkus, I. Andó, The cell-mediated

immunity of Drosophila melanogaster: hemocyte lineages, immune

compartments, microanatomy and regulation. *Dev. Comp. Immunol.* **42**,

47–56 (2014).

9.   J. P. Dudzic, S. Kondo, R. Ueda, C. M. Bergman, B. Lemaitre, Drosophila

innate immunity: regional and functional specialization of

prophenoloxidases. *BMC Biol.* **13**, 81–97 (2015).

10.   A. J. Nappi, M. Poirié, Y. Carton, in *Advances in Parasitology* (2009), pp. 99–
121.

11.   A. R. Kraaijeveld, H. C. J. Godfray, Geographic Patterns in the Evolution of
Resistance and Virulence in Drosophila and Its Parasitoids. *Am. Nat.* **153**,
S61–S74 (1999).

12.   J. E. McGonigle, A. B. Leitão, S. Ommeslag, S. Smith, J. P. Day, F. M. Jiggins,
Parallel and costly changes to cellular immunity underlie the evolution of
parasitoid resistance in three Drosophila species. *PLoS Pathog.* **13** (2017),
doi:10.1371/journal.ppat.1006683.

13.   A. R. Kraaijeveld, E. C. Limentani, H. C. Godfray, Basis of the trade-off
between parasitoid resistance and larval competitive ability in Drosophila
melanogaster. *Proc. Biol. Sci.* **268**, 259–61 (2001).

14.   M. D. Fellowes, A. R. Kraaijeveld, H. C. Godfray, Trade-off associated with
selection for increased ability to resist parasitoid attack in Drosophila
melanogaster. *Proc Biol Sci*. **265**, 1553–1558 (1998).

15.   S. Havard, P. Eslin, G. Prévost, G. Doury, Encapsulation ability: Are all
Drosophila species equally armed? An investigation in the *obscura* group.
Can. J. Zool. **87**, 635–641 (2009).

16.   Y. Fu, X. Huang, P. Zhang, J. Van De Leemput, Z. Han, Single-cell RNA
sequencing identifies novel cell types in Drosophila blood. *J. Genet.*
Genomics (2020), doi:10.1016/j.jgg.2020.02.004.

17.   S. Gopal Tattikota, Y. Hu, Y. Liu, B. Cho, V. Barrera, M. Steinbaugh, S.-H.

Yoon, A. Comjean, F. Li, F. Dervis, R.-J. Hung, J.-W. Nam, S. H. Sui, J. Shim, N.

Perrimon, A single-cell survey of Drosophila blood (2019),

doi:10.1101/2019.12.20.884999.

18. B. Cho, Single-cell transcriptome maps of myeloid blood cell lineages in

Drosophila. *bioRxiv* (2020).

19. P. B. Cattenoz, R. Sakr, A. Pavlidaki, C. Delaporte, A. Riba, N. Molina, N.

Hariharan, T. Mukherjee, A. Giangrande, Temporal specificity and

heterogeneity of Drosophila immune cells . *EMBO J.*, 1–25 (2020).

20. R. B. V. Vivancos, A. Giangrande, glide/gcmIs Expressed and Required in

the Scavenger Cell Lineage. *Dev. Biol.* **191**, 118–130 (1997).

21. U. Banerjee, J. R. Girard, L. M. Goins, C. M. Spratford, *Drosophila* as a Genetic

Model for Hematopoiesis. *Genetics.* **211**, 367–417 (2019).

22. B. A. Edgar, P. H. O'Farrell, The three postblastoderm cell cycles of

Drosophila embryogenesis are regulated in G2 by string. *Cell.* **62**, 469–480

(1990).

23. D. M. Glover, Polo kinase and progression through M phase in Drosophila:

a perspective from the spindle poles. *Oncogene.* **24**, 230–237 (2005).

24. K. Oegema, M. S. Savoian, T. J. Mitchison, C. M. Field, Functional Analysis of

a Human Homologue of the Drosophila Actin Binding Protein Anillin

Suggests a Role in Cytokinesis. *J. Cell Biol.* **150**, 539–552 (2000).

25. B. Olofsson, D. T. Page, Condensation of the central nervous system in

embryonic Drosophila is inhibited by blocking hemocyte migration or

neural activity. *Dev. Biol.* **279**, 233–243 (2005).

26.   S. Bunt, C. Hooley, N. Hu, C. Scahill, H. Weavers, H. Skaer, Hemocyte-

Secreted Type IV Collagen Enhances BMP Signaling to Guide Renal Tubule

Morphogenesis in Drosophila. *Dev. Cell*. **19**, 296–306 (2010).

27.   J. E. Mcgonigle, A. B. Leitão, S. Ommeslag, S. Smith, P. Day, F. M. Jiggins,

Parallel and costly changes to cellular immunity underlie the evolution of

parasitoid resistance in three Drosophila species. *PLoS Pathog.* **13(10)**, 1–

20 (2017).

28.   P. Eslin, G. Prevost, Hemocyte load and immune resistance to Asobara

tabida are correlated in species of the Drosophila melanogaster subgroup.

J. Insect Physiol. **44**, 807–816 (1998).

29.   V. Honti, G. Csordás, R. Márkus, E. Kurucz, F. Jankovics, I. Andó, Cell lineage

tracing reveals the plasticity of the hemocyte lineages and of the

hematopoietic compartments in Drosophila melanogaster. *Mol. Immunol.*

47, 1997–2004 (2010).

30.   I. Anderl, L. Vesala, T. O. Ihalainen, L. M. Vanha-aho, I. Andó, M. Rämet, D.

Hultmark, Transdifferentiation and Proliferation in Two Distinct

Hemocyte Lineages in Drosophila melanogaster Larvae after Wasp

Infection. *PLoS Pathog.* **12**, 1–34 (2016).

31.   T. M. Rizki, R. M. Rizki, Parasitoid-induced cellular immune deficiency in

Drosophila. *Ann. N. Y. Acad. Sci.* **712**, 178–94 (1994).

32.   A. R. Kraaijeveld, H. C. Godfray, Trade-off between parasitoid resistance

and larval competitive ability in Drosophila melanogaster. *Nature*. **389**,

278–280 (1997).

33.   J. Mattila, V. Hietakangas, Regulation of carbohydrate energy metabolism

409   in Drosophila melanogaster. *Genetics* (2017),

410   doi:10.1534/genetics.117.199885.

411   34.   C. Kim-Jo, J. L. Gatti, M. Poirié, Drosophila cellular immunity against

412   parasitoid wasps: A complex and time-dependent process. *Front. Physiol.*

413   **10** (2019), doi:10.3389/fphys.2019.00603.

Acknowledgements: The Genomics Core Facility at Cancer Research UK

Cambridge Institute, G. Raddi and S. Tattikota provided protocols for the

scRNAseq. **Funding:** This work was funded by Natural Environment Research

Council grant NE/P00184X/1 to FJ and AL, an EMBO Postdoctoral Fellowship to

AL (ALT-1556) and a Natural Sciences and Engineering Research Council of

Canada Postdoctoral Fellowship (PDF-516634-2018, Committee: 187) to RA.

Competing Interests: None. **Author Contributions:** AL, RA and FMJ designed

the study and wrote the paper with input from the other authors. AL, JD and EG

performed the *Drosophila* work. AL and JD generated the sequencing data. RA

analysed the sequence data. **Data and materials availability:** Unprocessed

single cell sequence reads were deposited in the Sequence Read Archive

(accession: SRP256887, Bioproject: PRJNA625925). Cell count matrices for all

detected genes, cluster identities and processed scRNA-seq results were

deposited into Gene Expression Omnibus (accession: GSE148826). The R script

used to analyze the scRNA-seq data is available on Github (Repository:

dmel_scRNA_hemocyte).