Shared safety abolishes the recovery of learned threat

Yafeng Pan[1], Andreas Olsson[1], & Armita Golkar[1,2*]

[1] *Department of Clinical Neuroscience, Karolinska Institutet, Stockholm, Sweden*

2 *Department of Psychology, Stockholm University, Stockholm, Sweden*

[*] **Corresponding authors:**

Armita Golkar

armita.golkar@psychology.su.se

Abstract

Humans, like other social animals, learn about threats and safety in the environment through

social cues. Yet, the processes that contribute to the efficacy of social safety learning during

threat transmission remain unknown. Here, we developed an experimental dyadic model

during which live pairs of participants underwent an associative threat learning and extinction

procedure followed by a threat recovery test. In three separate social groups, we manipulated

whether safety information during extinction was acquired via direct conditioned stimulus

(CS) exposure in the presence of another individual (Direct exposure), via observation of

another individual's safety behavior (Vicarious exposure), or via the combination of both

(Shared exposure). As a non-social baseline condition, a fourth group received direct CS

exposure alone (Asocial exposure). Based on skin conductance responses (SCRs), we

observed that compared to Asocial exposure, all three social groups were more successful in

inhibiting the recovery of learned threat responses; in particular, only Shared exposure

abolished threat recovery during early reinstatement. Additionally, synchronicity of dyads'

SCRs was higher for genuine interaction versus surrogate interaction across all three social

groups. These observations indicate that overall, both the mere presence and vicarious safety

transmission alone is sufficient to produce better inhibition of the recovery of learned threat

compared to asocial safety learning, but that abolishing threat recovery requires a

combination of physical presence and vicarious safety signals. This work has relevance for

understanding how social information can optimize standard, asocial safety learning

procedure to augment the effects of exposure on previously acquired fears.

Keywords: social interaction, social learning, shared safety, recovery, threat

1. Introduction

Learning to predict and respond to threats and potential dangers in the environment is critical to adaption and survival. Such learning has been well described using associative threat learning models in both humans and animals [1], and has advanced our understanding of the etiology of threat and anxiety related disorder. These models are however inherently asocial as they rely on individual trial and error learning - neglecting that humans are inherently social. Most of us spend the majority of our daily lives socially connecting with others; through communication, coordination, or observation. Apart from playing an important role in social bonding [2], these behaviors also constitute social forms of learning, facilitating our acquisition of knowledge about the surrounding environment and influence our expectations and behaviors [3–5].

Previous studies on social learning have mainly focused on how information about threats and potential dangers in the environment is transmitted between individuals [6–11] – a process known as *social threat learning* [12,13]. Social threat learning constitute one major route through which humans can acquire fears by observing the aversive experiences of others, is well conserved across a range of species from insects [14] to primates [15] and has been modeled in multiple modalities (behavior: e.g., [16]; physiology: e.g., [11,17,18]; neuroimaging: e.g., [19]). Yet, less is known about how these learned threat responses can be changed and modified through socially acquired safety information. This is critical as deficits in the capacity to adapt and regulate responses to threatening cues when they no longer signal danger is a key feature of threat and anxiety-related disorders [20]. Support for the utility of exploiting social information to regulate aversive responses mainly comes from research demonstrating that the presence of a conspecific attenuates the behavioral, physiological, and neural response to a stressful or threat-provoking event. This "social buffering" phenomenon has been found across social species, including zebrafish [21], rodents [22–24], non-human

primates [25–28], and humans [29,30], suggesting the existence of an evolved system by

social interaction can modulate behavior.

More specifically, previous experimental work in humans have demonstrated that acute

physical [31] and social stress [32] result in significantly smaller elevations in heart rate,

blood pressure, and cortisol in humans socially supported by another partner, compared to

individuals who undergo stress alone. More recently, the mere presence of another partner

was shown to reduce psychophysiological responses to aversive sounds, compared to an alone

condition [33]. Across species, the social buffering research has however almost exclusively

focused on the social regulation of stress responses [34,35] and little work has focused on

how social information modulates learned threat memories or how social safety information is

learned. Support for a regulatory role of social interaction on the expression of learned threat

memories comes from studies in rodents, in which the presence of another animal has been

shown facilitate extinction learning and reduce the expression of conditioned threat memory

[36,37]. Briefly, extinction learning involves the formation of novel, safety associations after

repeated safe exposures to a threatening stimulus; i.e. learning that the stimulus that earlier

predicted danger now predicts safety. However, unlike threat learning, the threat-reducing

effects of extinction learning decline with the lapse of time, do not generalize across different

contexts, and are easily reinstated in behavior following re-exposure to cues associated with

the original fear memory [38]. This recovery of threat responses after extinction also

contributes to relapse of fear that is common after extinction-based exposure therapies that are

used in the treatment of threat and anxiety-related disorders [39]. Consequently, one major

aim of the research on threat learning and regulation is to identify strategies that prevent the

return of fear after extinction in order to guide novel treatments approaches.

To address whether socially acquired safety information can optimize extinction of

learned threat responses in humans we have previously developed an experimental model of

social extinction learning [40]. Using this model, we have shown that social safety learning

accomplished by observing the safety behavior of another individual in the presence of a

threat conditioned stimulus (CS) more efficiently reduced psychophysiological threat

responses compared to asocial safety learning [40] and prevented the return of both short-term

[11,40,41] and long-term [18] threat memory. Moreover, in a related series of experiments,

we have shown that social safety learning prior to threat learning can prevent the social

transmission of threat [42], suggesting that social safety information may serve as a safety

signal that can modulate threat processing.

Previous studies on both social threat and safety learning have commonly adopted video-

based social learning paradigms [12]. In these paradigms, the observer learns about threat or

safety by watching a pre-recorded video depicting the behavior of a demonstrator reacting

either fearfully or calmly in the presence of threat. Although the recent decade has seen a

paradigm shift towards real-time two-person interactions in both social psychological and

neuroscientific research [43–48], few studies in humans have addresses whether social

transmission of threat and safety generalizes to a more naturalistic and ecologically valid

setting with opportunity for interaction (close to real-world situations). In order to model

social threat learning in a two-person setting, we recently examined skin conductance

responses (SCR, an index of threat learning) in dyads composed of observers and

demonstrators [49]. We found that observers successfully "caught" information of threat by

observing the fearful reactions of a demonstrator who sat side-by-side to them. Intriguingly,

synchronization of phasic skin conductance between demonstrator and observer predicted

observer's later threat memories.

In the current study, we aimed at extending our findings to a naturalistic dyadic setting to

examine how social safety learning prevents the recovery of learned threat. Understanding

how social information can optimize the effects of extinction learning and prevent the

recovery of threat is critical as extinction learning serves as a translation model for exposure

therapy that is used to treat symptoms of threat and anxiety-related disorders in humans [50].

Based on our previous work demonstrating that social safety learning outperforms asocial

safety learning in preventing the return of threat using a pre-recorded video-based approach

[40], we developed a dyadic model during which live pairs of participants underwent an

associative threat learning and extinction procedure followed by a threat recovery test. To

better isolate the social processes that are necessary for the efficacy of social safety learning,

we ran three separate social safety learning groups. Thus, we manipulated whether safety

information was acquired via own CS exposure in the mere presence of another individual

who's behavior was unrelated to the task (*Direct exposure*), via observation of another

individual's calm behavior in the absence of own CS exposure (*Vicarious exposure*) or via the

combination of own CS exposure and vicarious information from another individual's calm

behavior (*Shared exposure*). These social safety learning groups were contrasted with a

standard, asocial safety learning group who received direct CS exposure alone (*Asocial*

exposure). Learning was indexed using SCRs throughout all stages.

Our aim was to test two main predictions. First, directly drawn from our previous work

[40], we predicted that (*i*) overall, social safety learning (Direct, Vicarious, or Shared

exposure) would outperform asocial safety learning in *inhibiting* the recovery of threat, and

that (*ii*) in particular, Shared exposure would *abolish* threat recovery during early

reinstatement, paralleling our previous video-based work in which social safety learning

involved both CS and model exposure [11,18,41]. Finally, based on previous research

showing that inter-personal physiological synchrony is associated with social behaviors

[51,52], including threat learning [49], in a two-person situation [48], we carried out an

exploratory analysis to assess the physiological synchrony between two partners in a dyad

during social safety learning.

2. Methods

2.1. Ethics statement

This study was conducted in accordance with guidelines and regulations laid down in the Declaration of Helsinki. Written informed consent was obtained from each participant prior to the experiment. The study procedure was approved by Regional Ethical Review Board in Stockholm (DNR: 2016/249-31/1). Each participant was reimbursed with two movie tickets for their participation.

2.2. Participants

Power calculations were performed based on the obtained effect size (d = 0.75) in our original study comparing social and asocial extinction learning [40]. Setting the threshold to 95% sensitivity (1-beta = 0.95) and a significance level of alpha = 0.05 yielded a required sample size of 40 in each group. Because our original study [40] contained a small sample (20 vs. 16 in each group) and small studies are susceptible to inflated effects size estimates [53], we decided to oversample to reach a minimum of 50 participants in each experimental group. To achieve this, 231 participants were recruited from the student population at Karolinska Institutet. Participants were screened from having previously partaken in conditioning experiments and we confirmed that the participants in each dyad did not already know each other prior to participating in the experiment. Of the 231 participants who completed the experiment, we excluded 1 participant who failed to show any measurable skin conductance response (< 0.02 microsiemens, μS) to any CS during the entire experiment. This left 230 participants (132 females; mean age = 25.28 years, *SD* = 4.07) in the final sample. These participants were randomly assigned to either one of the three different social learning groups (Direct exposure, Vicarious exposure, and Shared exposure) or the asocial learning control group (for group characteristics and demographic information, see **Table1**).

147   **Table 1.** Group characteristics and demographic information.

Group	CS exposure	Social information	N	Gender	Age (SD)
Social safety learning					
Direct exposure	Yes	No	62	36F/26M	25.10 (4.11)
Vicarious exposure	No	Yes	60	33F/27M	25.46 (4.35)
Shared exposure	Yes	Yes	59	41F/18M	25.58 (3.80)
Asocial safety learning	Yes	/	49	22F/27M	24.94 (4.09)

Note. CS, conditioned stimulus; N, number; F, females; M, males; SD, standard deviation.

2.3. Experimental set-up

Social safety learning. Participants were enrolled in pairs of two participants that did not

report previous contact or familiarity with each other. Each pair were seated in a waiting area

during and were given time to read and sign informed consent forms before participation.

Then, each participant underwent a standardized shock calibration alone, with the aim to

individually adjust the level of shock intensity to a level that was highly uncomfortable but

not painful. Next, they were seated next to each other in the experimental room separated by a

moveable wall that occluded any visual information of the other participant. In addition, the

monitors were equipped with privacy screen filters to ensure that the visual presentations

were not visible to the other individual in each pair. Participants were also instructed they

were not allowed to talk to each other during any parts of the experiment. They were also

informed that there would be brief interruptions during which the experimenter would make

some adjustments and that should refrain from asking questions during these breaks.

Prior to the experiment (see **Figure 1** for an overview of the experimental stages), all

participants were allowed to familiarize with the stimuli. Participants were exposed to two

non-reinforced presentations of two different stimuli (one spider image and one snake image)

that would subsequently serve as the conditioned stimuli (CSs). The formal experiment

consisted of the following three stages: acquisition, extinction, and reinstatement test.

During the acquisition stage, all participants underwent a standard threat acquisition

protocol involving two CSs. For each participant, one CS predicted the delivery of a shock

(the unconditioned stimulus; US) on 2/3 of the CS presentations, whereas the other CS (CS-)

never predicted the US. This acquisition stage ended with an instruction screen: "The

experimenter will not make some minor adjustment, please remain seated".

During the subsequent extinction stage, the experimenter removed the movable wall

between the participants and adjusted mirrors that allowed the participants to receive visual

feedback from the face of the other participant. The two mirrors were placed in an angle so

that the reflection of the other participant's face did not require them to move their gaze away

from the visual presentation on their screen. We manipulated safety transmission in three

separate groups by changing both CS exposure and instructions. In the *Direct exposure* group,

participants received the following written instructions: "During the next stage, ONLY you

will be exposed to same pictures as you saw during the preceding stage, and the person sitting

next to you will be exposed to two novel pictures. In the *Vicarious exposure* group,

participants were informed that "During the next stage, you will be exposed to a presentation

of two novel pictures, and ONLY the person sitting next to you will be exposed to the same

pictures as you were exposed to during the preceding stage" and the *Shared exposure* group

were informed "During the next stage, BOTH you and the person sitting next to you will be

exposed to the same pictures as during the preceding stage". For the Direct and Shared

exposure groups, the extinction stage consisted of non-reinforced presentations of each CS. In

the Vicarious exposure group, the CSs were replaced by the presentation of two neutral and

non-related control stimuli (picture of a butterfly and a bird).

During the final reinstatement stage, the experimenter re-positioned the wall between the

participants and removed the mirrors. After this readjustment, all participant received

identical instruction on their screen "The experiment will now be resumed" which was

followed by a standard *reinstatement* test during which participants–received 3 un-signaled

shocks followed by a black screen (34 s) before all CSs were represented to the participants.

Asocial safety learning. Participants underwent the same experimental stages

(acquisition, extinction, and reinstatement) using the same set-up but alone in the absence of

another participant. After the acquisition stage, participants in this *Asocial exposure* group

received the following instructions "During the next stage, you will now be exposed to the

same two pictures as you were during the preceding stage". This stage was followed by the

reinstatement stage.

During all stages of the experiment, each CS was presented six times with a duration of 6

s. Between each CS trial, a black screen (i.e., inter-stimulus interval, ITI) was presented for

12–18 s.

Manipulation check. All pairs reported seeing the other participants face in the mirrors

during extinction, without having to divert gaze from the CS presentation, and none of the

included participants reported being able to see the CS presentation of the other individual.

Figure 1. General set-up. All participants in (A) and (B) underwent an identical acquisition and reinstatement

stage with the exception that in (A), the Asocial learning group performed all stages alone. Participants in the

asocial learning group received exposure to the CSs alone. Participants in the social learning group were further

categorized into three sub-groups (Direct, Vicarious, and Shared exposure) based on different CS exposure and

instructions they received during extinction. Participants in both the Direct exposure and Shared exposure groups

viewed the same CS as during the previous stage but either instructed that the person sitting next to them were

observing novel, irrelevant stimuli (Direct exposure) or the same stimuli (Shared exposure) as themselves.

Participants in the Vicarious exposure group were exposed to two novel, neutral stimuli and instructed that only

the person sitting next to them were exposed to the stimuli from the preceding stage. During all stages, screen

filters occluding all information of what the other participant was exposed to were used.

2.4. Data acquisition

Participants' SCRs were recorded and measured using a Biopac MP150 system and AcqKnowledge 4.0 software (BIOPAC Systems Inc., Goleta, CA, USA). Ag-AgCl electrodes were attached to the middle and index fingers from the hand not receiving shocks. A conducting gel was used to improve signal quality, in accordance with standard recommendations [54]. Electric shocks (aversive stimuli) were delivered using a STM200 module (BIOPAC Systems Inc., Goleta, CA, USA). The sampling rate was set to 2 kHz. Shocks consisted of a single 100ms DC pulse, which were applied to the lower forearm. The intensity of the shocks was individually calibrated, following the rule being "unpleasant but not painful". Stimulus presentation and shock delivery were controlled by the Presentation software (Neurobehavioral Systems, Inc., Albany, CA).

2.5. Data analysis

SCRs were continuously recorded throughout all experimental stages (acquisition, extinction, and reinstatement test). The raw signals were filtered offline with a low-pass filter (1 Hz) and a high-pass filter (0.05 Hz). The SCR was measured for each stimulus trial as the base-to-peak amplitude of the first response (in μS) in the latency window from 0.5 to 4.5 s following stimulus onset [12]. Minimal response criterion was set to 0.02 μS, and responses that did not pass this criterion were scored as 0. SCR scores were then z-transformed to normalize the distributions [54]. Clean SCRs were then used for subsequent statistical comparisons.

In addition, to access the physiological synchrony between two participants in a dyad, clean SCR time series was further submitted to an exploratory analysis using the Pearson correlation coefficient [55,56]. The correlation coefficients between two SCRs from participant # 1 and participant #2 were calculated over the whole extinction stage (**Figure 1**). Resulting values were Fisher-z-transformed. Note that only Social exposure groups (Direct,

Vicarious, and Shared exposure) were included in this exploratory analysis, as physiological

synchrony should be estimated within dyads. One dyad containing a missing participant

(failed in SCR acquisition, see section 2.2) were excluded, resulting in 29 pairs in the Direct

exposure group, 30 pairs in the Vicarious exposure group, and 31 pairs in the Shared exposure

group.

2.6. Statistical analyses

All the statistical analyses were conducted using IBM SPSS (version 21, SPSS Inc., Chicago,

IL, USA). Each stage of the experiment was analyzed separately and significant interactions

were followed up by separate two-tailed t-tests. Significance level was set to $p < 0.05$. η^2 and

Cohen's d are reported as measures of effect size where appropriate.

2.6.1. Confirmatory analysis

When analyzing data from the acquisition stage, a $2 \times 6 \times 4$ mixed-design Analysis of

Variance (ANOVA), with Stimulus (CS+ and CS-) and Trial (#1~6) as two within-subject

factors and Group (Direct, Vicarious, Shared, and Asocial exposure) as a between-subject

factor, was used to analyze all results across conditions. Degrees of freedom were corrected

using Greenhouse-Geisser estimates when assumption of sphericity was violated. We

predicted successful threat acquisition in all experimental groups. Successful acquisition was

defined as higher SCR to the CS+ relative to the CS- across acquisition, which would be

supported by a significant Stimulus $\times$ Trial interaction in the absence of any significant

effects involving groups.

Data from the extinction stage were analyzed using the same mixed-design ANOVA.

Extinction of the conditioned SCR was defined as decreasing SCR responses to the CS+

relative to the CS- across extinction training (Stimulus $\times$ Trial interaction). We anticipated a

decreasing trend of CS+/CS- differentiation (as a marker of diminishing conditioned SCR)

across trials in all groups except in the Vicarious exposure group that did not receive CS

exposure during extinction and was therefore not expected to show significant CS+/CS-

differentiation during this stage (no CS exposure, see **Table 1**).

2.6.2. Hypothesis-driven analysis

When analyzing data from the reinstatement stage, we carried out a series of statistics in a

hypothesis-driven way. To test our first prediction i.e., that social safety learning would

produce less recovery of learned threat compared to asocial safety learning, we compared

CS+/CS- differentiation across reinstatement in a Stimulus (2) × Trial (6) × Group (4) mixed-

design ANOVA, followed by planned comparisons between each social exposure group

(Shared, Vicarious, or Direct) with the Asocial exposure group. To test our second hypothesis

(i.e., that the Shared exposure group would abolish CS+/CS- differentiation during early

reinstatement), we ran a Stimulus (2) × Trial (2: first 2 trials) × Group (4) mixed-design on

SCR. We only focused on the early trials of reinstatement due to rapid re-extinction during

reinstatement [57], and in line with previous research demonstrating the efficacy of social

safety learning in preventing the recovery of threat responses focusing on these early trials

[11,18,41]. Each set of planned contrasts were Bonferroni corrected to account for multiple

comparisons.

2.6.3. Exploratory analysis

Finally, we conducted exploratory analyses to test the potential physiological synchrony

between two participants in a dyad under social settings (i.e. during extinction). Synchrony

was estimated using the Pearson correlational approach, in accordance with recent advances

[55,56]. To evaluate the statistical significance of the Pearson correlation coefficients at the

extinction stage, we created a null distribution based on a surrogate dataset. The surrogate

dataset was obtained by Fourier transforming the original SCR signal, randomizing the phases

of Fourier coefficients, performing an inverse Fourier transform, and re-computing the

correlation coefficients [55,58,59]. This procedure was repeated 1,000 times. Significance

levels (thresholded at 0.05) were evaluated by comparing the correlation coefficients from the

original dataset and those from the surrogate dataset. It was assumed that physiological

synchrony based on original dataset should be higher than that based on surrogate dataset, as

the latter disrupts the temporal structure of the original signals.

3. Results

3.1. Confirmatory results

3.1.1. Acquisition stage

First we confirmed that all group successfully acquired conditioned threat responses as

supported by significantly higher SCR to the CS+ than the CS- across acquisition (Stimulus:

$F_{1,226} = 13.74$, $p < 0.001$, $\eta_{partial}^2 = 0.06$; Stimulus × Trial interaction: $F_{4.71,1063.55} = 37.08$, $p <$

0.001, $\eta_{partial}^2 = 0.14$; see **Figure 2** (left column). There were no significant effects involving

Group (main effect: $F_{3, 226} = 0.14$, $p = 0.94$; Trial × Group: $F_{12.93,974.11} = 1.17$, $p = 0.30$;

Stimulus × Group: $F_{3, 226} = 2.15$, $p = 0.10$; Stimulus × Trial × Group, $F_{14.12,1063.55} = 0.40$, $p =$

0.98).

3.1.2. Extinction stage

Second, we confirmed that CS+/CS- differentiation decreased across extinction. Progressive

extinction was confirmed by a main effect of Trial ($F_{4.19,946.85} = 47.66$, $p < 0.001$, $\eta_{partial}^2 =$

0.17) and an interaction of Stimulus × Trial ($F_{4.58,1034.82} = 3.49$, $p = 0.01$, $\eta_{partial}^2 = 0.02$), with

a generally decreasing trend in CS+/CS- differentiation (**Figure 2**, middle column). However,

as expected, CS+/CS- differentiation differed between the Vicarious exposure group that did

not receive own CS exposure and the other groups. This was evidence by a main effect of

Group ($F_{3,226} = 3.87$, $p = 0.01$, $\eta_{partial}^2 = 0.05$) and a Stimulus $\times$ Group interaction ($F_{3,226} =$

9.14, $p < 0.001$, $\eta_{partial}^2 = 0.11$), indicating that CS+/CS- differentiation was significant in all

groups (Asocial exposure, $t_{48} = 4.66$, $p < 0.001$, Cohen's $d = 2.14$; Direct exposure, $t_{61} = 4.61$.

$p < 0.001$, Cohen's $d = 0.85$; and Shared exposure, $t_{58} = 6.39$, $p < 0.001$, Cohen's $d = 1.03$),

except for the Vicarious exposure group that did not receive own CS exposure ($t_{59} = 0.42$, $p =$

0.68). This analysis did not yield a significant three-way interaction ($F_{13.74,1034.82} = 1.06$, $p =$

0.39).

To further elucidate the extinction effect, we analyzed CS+/CS- differentiation over the

last two trial of the extinction stage. Results indicated that the Direct exposure ($F_{1,61} = 1.92$, p

$= 0.17$) and Vicarious exposure ($F_{1,59} = 0.27$, $p = 0.61$) groups efficiently extinguished

CS+/CS- differentiation, whereas the Shared exposure ($F_{1,58} = 6.12$, $p = 0.02$, $\eta_{partial}^2 = 0.10$)

and Asocial exposure ($F_{1,48} = 5.40$, $p = 0.02$, $\eta_{partial}^2 = 0.10$) groups did not.

3.2. Hypothesis-driven results

To test our first prediction that all social safety learning groups would better inhibit

reinstatement (i.e., less CS+/CS- differentiation) compared to the Asocial exposure group, we

ran a Stimulus (2) $\times$ Trial (6) $\times$ Group (4) mixed design ANOVA. Overall, this analysis

yielded a significant Stimulus $\times$ Group interaction ($F_{3,226} = 3.40$, $p = 0.02$, $\eta_{partial}^2 = 0.04$), a

main effect of Group ($F_{3,226} = 3.22$, $p = 0.02$, $\eta_{partial}^2 = 0.04$) and no interaction with Trial (Fs

< 1.03, $ps > 0.42$). Comparisons of the separate social exposure groups (Shared, Vicarious, or

Direct) vs. the Asocial exposure group using adjusted alpha levels of $p = 0.017$ (0.05/3)

resulted in a significant Stimulus $\times$ Group interaction in the Shared vs. Asocial exposure

comparison ($F_{1,106} = 7.64$, $p = 0.01$, $\eta_{partial}^2 = 0.07$), which, as predicted, was driven by lower

CS+/CS- differentiation in Shared exposure relative to Asocial exposure (see **Figure 2**, right

column). Corresponding analyses for the Vicarious vs. Asocial exposure group ($F_{1,107} = 5.08$,

$p = 0.01$, $\eta_{partial}^2 = 0.05$) and the Direct vs. Asocial exposure ($F_{1,109} = 7.38$, $p = 0.01$, $\eta_{partial}^2 =$

0.06) were also significant at the adjusted alpha level.

Figure 2. Mean skin conductance responses (z-score) across stimuli, groups, and trials over the entire

experiment. The graphs from upper to bottom represent the data from Asocial exposure, Direct exposure,

Vicarious exposure, and Shared exposure groups, respectively. Left column: acq1-acq6 refers to the 6

acquisition trials; middle column: ext1-ext6 refers to the 6 extinction trials; right column: rei1-rei6 refers to the 6

reinstatement trials. Grey shadow highlights the findings from the early reinstatement stage, indicating that

shared safety abolishes the recovery of learned threat. Error bars denote standard errors.

To test our second prediction that the Shared exposure would abolish CS+/CS-

differentiation during early reinstatement, we next analyzed early reinstatement test

performance. This analysis resulted in a significant main effect of Group ($F_{3,226} = 3.64$, $p =$

0.01, $\eta_{partial}^2 = 0.05$), and Trial ($F_{1,226} = 31.00$, $p < 0.001$, $\eta_{partial}^2 = 0.12$) and a significant

Stimulus $\times$ Group interaction ($F_{3,226} = 2.86$, $p = 0.036$, $\eta_{partial}^2 = 0.04$). Using adjusted alpha

levels of $p = 0.017$ (0.05/3), we observed the predicted difference between the Shared

exposure and the Asocial exposure group (Stimulus $\times$ Group, $F_{1,106} = 9.19$, $p = 0.003$, $\eta_{partial}^2$

$= 0.08$), and no significant differences between Vicarious vs. Asocial exposure ($F_{1,107} = 3.55$,

$p = 0.06$), or Direct vs. Asocial exposure ($F_{1,109} = 1.32$, $p = 0.25$). Critically, in contrast to

Asocial exposure (main effect of Stimulus, $F_{1,48} = 23.21$, $p < 0.001$, $\eta_{partial}^2 = 0.33$), Shared

exposure abolished CS+/CS- differentiation during early reinstatement (main effect of

Stimulus, $F_{1,58} = 0.00$, $p = 0.99$, **Figure 3A**) using an adjusted alpha level of $p = 0.025$

(0.05/2). The absence of CS+/CS- differentiation during early reinstatement in the Shared

Exposure group was further evidenced by a Bayesian paired samples *t*-test using JASP

software (https://jasp-stats.org/, JASP Team, 2017), 95% Credible Interval = [-0.30, 0.21],

Bayes　　　Factor　　　(BF_{10})　　　=　　　0.16　　　**(Figure　　3B)**.

Figure 3. (A) Mean skin conductance response (z-score) across stimuli, groups during the early reinstatement

stage. Error bars represent standard errors. (B) Inferential plots for prior and posterior probabilities. Skin

conductance responses to CS+ and CS- during early reinstatement in the Shared Exposure group were compared

using a Bayesian paired samples *t*-test.

3.3. Exploratory results

Our exploratory analysis focused on evaluating the physiological synchrony between two

participants in a dyad at the extinction stage (**Figure 4**), during which the dyads were situated

in a real-time social setting. All three social safety learning groups showed significantly

stronger synchrony (averaged across trials and stimuli) between the SCR signals of two

participants within dyads compared to a surrogate dataset (generated by phase-scrambling of

original signals): Direct exposure (mean ± SD, 0.19 ± 0.05), Vicarious exposure (0.08 ± 0.05),

and Shared exposure (0.20 ± 0.05), *p*s < 0.02.

An additional Stimulus (2) × Trial (6) × Group (3) mixed-design ANOVA on

physiological synchrony revealed a main effect of Trial, $F_{4.46,388.12} = 7.66$, $p < 0.001$, $\eta_{partial}^2 =$

0.08, suggesting that synchrony generally attenuated across trials. The main effect of Stimulus

was marginally significant, $F_{1,87} = 3.11$, $p = 0.08$, $\eta_{partial}^2 = 0.04$, indicating that synchrony was

more pronounced for CS+ vs. CS-. No other significant effects were observed ($Fs < 1.91$, $ps >$

0.15).

Figure 4. Physiological synchrony (average r) between skin conductance responses from two participants in a

dyad across groups during the extinction stage. Ext1-ext6 refers to the 6 extinction trials. Shadow areas represent

± 1 standard errors.

4. Discussion

This study investigated how the recovery of learned threat responses are influenced by

socially acquired safety information. In doing so, the study departed from two main

hypotheses: (*i*) that social safety learning outperforms asocial safety learning in *inhibiting* the

recovery of learned threat responses, and (*ii*) shared social safety learning *abolishes* threat

recovery during early reinstatement. To address these hypotheses, we used a standard

conditioning procedure [40,60], which consists of three stages: (*i*) acquisition –threat

acquisition through direct CS-US exposure; (*ii*) extinction –safety learning in the absence of

aversive outcome (no US); and (*iii*) reinstatement – recovery of learned threat to the CSs after

re-exposure to the US. Critically, in three separate social safety learning groups, we

manipulated whether safety information during extinction was acquired via direct CS

exposure in the presence of another individual (*Direct exposure*), via observation of another

individual's safety behavior (*Vicarious exposure*) or via the combination of direct and

vicarious safety exposure (*Shared exposure*). As a control, we ran a standard, asocial

extinction group who received direct CS exposure alone (*Asocial exposure*). Our results show

that (*i*) compared to asocial safety learning, all three social safety learning groups were more

successful in inhibiting the recovery of learned threat responses, as measured during a

subsequent reinstatement stage, and (*ii*) only shared safety learning abolished the recovery of

learned threat response, as evidenced by the non-significant CS+/CS- differentiation

(confirmed by Bayesian factor analysis) during early stage of reinstatement. For this second

hypothesis, we restricted our analysis to the early trials of reinstatement because previous

research showing that reinstated threat responses typically undergo rapid re-extinction [57].

These results suggest that both the mere presence of another person, and vicarious safety

signals alone is sufficient to produce better inhibition of the recovery of learned threat

responses compared to asocial learning, but that during early reinstatement, the beneficial

effects of socially acquired safety relative to asocial safety learning are particularly

pronounced when safety signals can be acquired from both direct experience and from

another person (*Shared exposure*).

Previous studies supporting a modulatory role of social signals on aversive responses

come mainly from non-human animal work [61] and specifically from rodent models that

have demonstrated that the expression of conditioned threat responses are reduced in the

presence of a non-fearful conspecific [36,37]. This previous work indicates that a key factor

facilitating safety learning is a social modulation mechanism, which requires the presence of a

social conspecific and the trigger of safety signals during safety learning. Such "social

modulation" might be empirically characterized in terms of physiological synchrony [62]. We

observed higher synchrony for all social learning groups, compared to phase-scrambled

controls. Thus, physiological synchrony might support shared experience and/or interpersonal

information transfer by aligning physiological processes between the participants [62]. In

related fields, shared neural representation have been shown to contribute to efficacy of

human learning during both shared understanding [63,64] and shared attention [65,66]. For

example, participants listened to a narrative during which their brain activity was recorded;

interpersonal entrainment of brain activity was detected when participants shared

understanding of a story [63]. These studies suggest that temporally concurrent patterns of

task-related shared experience may lead to entrainment (or phase resetting) of neural

processes across individuals [67,68]. Such neural entrainment might impact upon real-time

acquisition of ongoing information (e,g., safety signals), and eventually upon learning [45,46].

To date, however, no research has investigated this in the field of social safety learning. This

study did not find any significant difference in physiological synchrony between the social

groups; the lack of relationship deserves further investigations, as it may reflect either a lack

of sensitivity compared to linguistics-involved paradigms [63] or analogous effects of social

modulation on interpersonal autonomic coupling during safety learning.

The interpretation of our findings requires some considerations. First, the augmented

inhibition of threat recovery during early reinstatement in the Shared exposure group does not

seem to be related to end extinction performance. Neither Shared exposure nor Asocial

exposure entirely abolished CS+/CS- differentiation during end extinction, but strikingly, they

differed during the subsequent reinstatement stage. This lack of consistency between end

extinction performance and recovery phenomena has been documented in previous research

[69,70]. Second, an inspection of **Figure 3A** shows that whereas both the Shared and

Vicarious extinction group showed relatively low CS+/CS- differentiation during early

reinstatement, the lack of CS+/CS- differentiation in the Shared exposure group was due to

comparably lower CS+ responses. In contrast, the Vicarious exposure group showed high

CS+ *and* high CS- responses, indicating generalization of conditioned threat responses in this

group. The stronger CS- response observed in the Vicarious exposure group might be related

to the fact that this group did not receive CS exposure during the extinction stage and

therefore showed overall stronger SCRs to the re-appearance of the CSs. Third, the current

study did not allow for verbal interaction between two individuals. However, to ensure that

participants could access socially transmitted information, we set-up mirrors so participants

could observe each other's emotional reactions (e.g., facial expression and body language)

during the safety learning (i.e., extinction) stage. Fourth, we observed a similar level of

physiological synchrony among all three social learning groups, which may have contributed

to the increased inhibition of threat recovery in all social groups compared to the asocial

baseline condition. Yet, this also implies that the abolishing learned threat responses during

early reinstatement following Shared exposure cannot be interpreted solely based on

interpersonal autonomic coupling across individuals. The results from these exploratory

analyses should be interpreted with caution, as our power calculations were based on a priori

hypotheses for the main analyses of threat-related SCR responses. An inevitable consequence

for the analyses focusing on physiological synchrony between dyads is that the sample size is

reduced by 50% - which represents a common issue in dyadic studies [71].

Taken together, these observations indicate that both CS exposure in the mere physical

presence of another person, and the mere safety signal from another person is sufficient to

inhibit the recovery of learned threat responses, but that abolishing the recovery of threat

requires the combination of both (or beyond). As we did not have an a priori hypothesis for

differences between the social safety learning groups (Direct vs. Vicarious vs. Shared

exposure; i.e., which group best inhibits the recovery of learned threat), this question should

be further addressed in future research powered enough to detect between-group differences.

Nevertheless, to our knowledge, our work represents the first demonstration of social safety

learning in a two-person situation with high ecological validity. The current set of findings

thus contributes to the emerging field of "two-person physiology" [62] and extends recent

research addressing the influence of physical presence on physiological responding to

aversive stimuli [33] by addressing the influence of social modulation of threat in a safety

learning context. More specifically, by focusing on safety learning, this work has relevance

for understanding how social information can optimize standard, asocial safety learning

procedure to augment the effects of exposure on previously acquired fears. Future research

should more closely address the relative contribution of different social processes to safety

learning and delineate the underlying mechanistic features that characterize these processes.

Data accessibility

The full dataset used is accessible at https://osf.io/h3ckv/.

Authors' contributions

Armita Golkar conceived the study and designed the experiment with input from Andreas

Olsson. Armita Golkar and Yafeng Pan performed the statistical analysis and interpreted the

data and Yafeng Pan drafted the manuscript. All authors critically revised the manuscript and

gave final approval for publication and agree to be held accountable for the work performed

therein.

Competing interests

The authors have declared that no competing interests exist.

Funding

This research was supported by a Consolidator Grant (2018-00877) from the Swedish

Research Foundation (Vetenskapsrådet), the Knut and Alice Wallenberg Foundation (KAW

2014.0237), and a Starting Grant (284366) from the European Research Council to Andreas

Olsson, and an independent project grant to Armita Golkar from the Swedish Research

Council (Vetenskapsrådet: 2017-02210).

Acknowledgements

We would like to thank Lisa Espinosa, Cathelijn Tjaden and Anastasia Dima for help with the

data collection and scoring.

References

1.   LeDoux JE. 2000 Emotion Circuits in the Brain. *Annu. Rev. Neurosci.* **23**, 155–184.

(doi:10.1146/annurev.neuro.23.1.155)

2.   Oullier O, de Guzman GC, Jantzen KJ, Lagarde J, Scott Kelso JA. 2008 Social

coordination dynamics: Measuring human bonding. *Soc. Neurosci.* **3**, 178–192.

(doi:10.1080/17470910701563392)

3.   Askew C, Field AP. 2008 The vicarious learning pathway to fear 40 years on. *Clin.*

Psychol. Rev. **28**, 1249–1265. (doi:10.1016/j.cpr.2008.05.003)

4.   Bandura A. 1977 Self-efficacy: Toward a unifying theory of behavioral change.

Psychol. Rev. **84**, 191–215. (doi:10.1037/0033-295X.84.2.191)

5.   Olsson A, Phelps EA. 2007 Social learning of fear. *Nat. Neurosci.* **10**, 1095–1102.

(doi:10.1038/nn1968)

6.   Beckers T, Krypotos A-M, Boddez Y, Effting M, Kindt M. 2013 What's wrong with

fear conditioning? *Biol. Psychol.* **92**, 90–96. (doi:10.1016/j.biopsycho.2011.12.015)

7.   Debiec J, Olsson A. 2017 Social Fear Learning: from Animal Models to Human

Function. *Trends Cogn. Sci.* **21**, 546–555. (doi:10.1016/j.tics.2017.04.010)

8.   Golkar A, Björnstjerna M, Olsson A. 2015 Learned fear to social out-group members

are determined by ethnicity and prior exposure. *Front. Psychol.* **6**, 123.

(doi:10.3389/fpsyg.2015.00123)

9.   Lau JY *et al.* 2011 Distinct neural signatures of threat learning in adolescents and

adults. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 4500–4505. (doi:10.1073/pnas.1005494108)

10. Lindström B, Selbing I, Olsson A. 2016 Co-evolution of social learning and evolutionary preparedness in dangerous environments. *PLoS One* **11**, e0160245. (doi:10.1371/journal.pone.0160245)

11. Golkar A, Castro V, Olsson A. 2015 Social learning of fear and safety is determined by the demonstrator's racial group. *Biol. Lett.* **11**, 20140817. (doi:10.1098/rsbl.2014.0817)

12. Haaker J, Golkar A, Selbing I, Olsson A. 2017 Assessment of social transmission of threats in humans using observational fear conditioning. *Nat. Protoc.* **12**, 1378–1386. (doi:10.1038/nprot.2017.027)

13. LeDoux JE. 2017 Semantics, Surplus Meaning, and the Science of Fear. *Trends Cogn. Sci.* **21**, 303–306. (doi:10.1016/j.tics.2017.02.004)

14. Leadbeater E, Chittka L. 2009 Bumble-bees learn the value of social cues through experience. *Biol. Lett.* **5**, 310–312. (doi:10.1098/rsbl.2008.0692)

15. Mineka S, Davidson M, Cook M, Keir R. 1984 Observational conditioning of snake fear in rhesus monkeys. *J. Abnorm. Psychol.* **93**, 355–372. (doi:10.1037/0021-843X.93.4.355)

16. Lindström B, Golkar A, Jangard S, Tobler PN, Olsson A. 2019 Social threat learning transfers to decision making in humans. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 4732–4737. (doi:10.1073/pnas.1810180116)

17. Selbing I, Olsson A. 2019 Anxious behaviour in a demonstrator affects observational learning. *Sci. Rep.* **9**, 9181. (doi:10.1038/s41598-019-45613-1)

18. Golkar A, Tjaden C, Kindt M. 2017 Vicarious extinction learning during reconsolidation neutralizes fear memory. *Behav. Res. Ther.* **92**, 87–93. (doi:10.1016/j.brat.2017.02.004)

19. Lindström B, Haaker J, Olsson A. 2018 A common neural network differentially mediates direct and social fear learning. *Neuroimage* **167**, 121–129.

(doi:10.1016/j.neuroimage.2017.11.039)

20. Graham BM, Milad MR. 2011 The Study of Fear Extinction: Implications for Anxiety

Disorders. *Am. J. Psychiatry* **168**, 1255–1265. (doi:10.1176/appi.ajp.2011.11040557)

21. Faustino AI, Tacão-Monteiro A, Oliveira RF. 2017 Mechanisms of social buffering of

fear in zebrafish. *Sci. Rep.* **7**, 1–10. (doi:10.1038/srep44329)

22. Kiyokawa Y, Takeuchi Y, Mori Y. 2007 Two types of social buffering differentially

mitigate conditioned fear responses. *Eur. J. Neurosci.* **26**, 3606–3613.

(doi:10.1111/j.1460-9568.2007.05969.x)

23. Kiyokawa Y, Wakabayashi Y, Takeuchi Y, Mori Y. 2012 The neural pathway

underlying social buffering of conditioned fear responses in male rats. *Eur. J.*

Neurosci. **36**, 3429–3437. (doi:10.1111/j.1460-9568.2012.08257.x)

24. Colnaghi L, Clemenza K, Groleau SE, Weiss S, Snyder AM, Lopez-Rosas M, Levine

AA. 2016 Social involvement modulates the response to novel and adverse life events

in mice. *PLoS One* **11**, e0163077. (doi:10.1371/journal.pone.0163077)

25. Sanchez MM, McCormack KM, Howell BR. 2015 Social buffering of stress responses

in nonhuman primates: Maternal regulation of the development of emotional regulatory

brain circuits. *Soc. Neurosci.* **10**, 512–526. (doi:10.1080/17470919.2015.1087426)

26. Winslow JT, Noble PL, Lyons CK, Sterk SM, Insel TR. 2003 Rearing effects on

cerebrospinal fluid oxytocin concentration and social buffering in rhesus monkeys.

Neuropsychopharmacology **28**, 910–918. (doi:10.1038/sj.npp.1300128)

27. Wittig RM, Crockford C, Weltring A, Langergraber KE, Deschner T, Zuberbühler K.

2016 Social support reduces stress hormone levels in wild chimpanzees across stressful

events and everyday affiliations. *Nat. Commun.* **7**, 13361. (doi:10.1038/ncomms13361)

28. Stanton ME, Patterson JM, Levine S. 1985 Social influences on conditioned cortisol

secretion in the squirrel monkey. *Psychoneuroendocrinology* **10**, 125–134.

(doi:10.1016/0306-4530(85)90050-2)

29.   Gunnar MR, Hostinar CE. 2015 The social buffering of the hypothalamic–pituitary–

adrenocortical axis in humans: Developmental and experiential determinants. *Soc.*

Neurosci. **10**, 479–488. (doi:10.1080/17470919.2015.1070747)

30.   Ditzen B, Heinrichs M. 2014 Psychobiology of social support: The social dimension of

stress buffering. *Restor. Neurol. Neurosci.* **32**, 149–162. (doi:10.3233/RNN-139008)

31.   Roberts MH, Klatzkin RR, Mechlin B. 2015 Social Support Attenuates Physiological

Stress Responses and Experimental Pain Sensitivity to Cold Pressor Pain. *Ann. Behav.*

Med. **49**, 557–569. (doi:10.1007/s12160-015-9686-3)

32.   Heinrichs M, Baumgartner T, Kirschbaum C, Ehlert U. 2003 Social support and

oxytocin interact to suppress cortisol and subjective responses to psychosocial stress.

Biol. Psychiatry **54**, 1389–1398. (doi:10.1016/S0006-3223(03)00465-7)

33.   Qi Y, Herrmann MJ, Bell L, Fackler A, Han S, Deckert J, Hein G. 2020 The mere

physical presence of another person reduces human autonomic responses to aversive

sounds. *Proc. R. Soc. B Biol. Sci.* **287**, 20192241. (doi:10.1098/rspb.2019.2241)

34.   Hostinar CE, Sullivan RM, Gunnar MR. 2014 Psychobiological mechanisms

underlying the social buffering of the hypothalamic-pituitary-adrenocortical axis: A

review of animal models and human studies across development. *Psychol. Bull.* **140**,

256–282. (doi:10.1037/a0032671)

35.   Kikusui T, Winslow JT, Mori Y. 2006 Social buffering: Relief from stress and anxiety.

Philos. Trans. R. Soc. B Biol. Sci. **361**, 2215–2228. (doi:10.1098/rstb.2006.1941)

36.   Brill-Maoz N, Maroun M. 2016 Extinction of fear is facilitated by social presence:

Synergism with prefrontal oxytocin. *Psychoneuroendocrinology* **66**, 75–81.

(doi:10.1016/j.psyneuen.2016.01.003)

37.   Ferreira FF, Rodrigues FS, Schmidt SD, Cavalcante LES, Zinn CG, Farias CP, Furini

CRG, Myskiw JC, Izquierdo I. 2019 Social support favors extinction and impairs

acquisition of both short- and long-term contextual fear conditioning memory.

Neurosci. Lett. **712**, 134505. (doi:10.1016/j.neulet.2019.134505)

38.   Bouton ME. 2002 Context, ambiguity, and unlearning: Sources of relapse after

behavioral extinction. *Biol. Psychiatry.* **52**, 976–986. (doi:10.1016/S0006-

3223(02)01546-9)

39.   Rachman S. 1977 The conditioning theory of fearacquisition: A critical examination.

Behav. Res. Ther. **15**, 375–387. (doi:10.1016/0005-7967(77)90041-9)

40.   Golkar A, Selbing I, Flygare O, Ohman A, Olsson A. 2013 Other people as means to a

safe end: vicarious extinction blocks the return of learned fear. *Psychol. Sci.* **24**, 2182–

90. (doi:10.1177/0956797613489890)

41.   Golkar A, Haaker J, Selbing I, Olsson A. 2016 Neural signals of vicarious extinction

learning. *Soc. Cogn. Affect. Neurosci.* **11**, 1541–1549. (doi:10.1093/scan/nsw068)

42.   Golkar A, Olsson A. 2016 Immunization against social fear learning. *J. Exp. Psychol.

Gen.* **145**, 665–671. (doi:10.1037/xge0000173)

43.   Cheng X, Pan Y, Hu Y, Hu Y. 2019 Coordination Elicits Synchronous Brain Activity

Between Co-actors: Frequency Ratio Matters. *Front. Neurosci.* **13**, 1071.

(doi:10.3389/fnins.2019.01071)

44.   Pan Y, Cheng X, Zhang Z, Li X, Hu YY. 2017 Cooperation in lovers: An fNIRS-based

hyperscanning study. *Hum. Brain Mapp.* **38**, 831–841. (doi:10.1002/hbm.23421)

45.   Pan Y, Novembre G, Song B, Li X, Hu Y. 2018 Interpersonal synchronization of

inferior frontal cortices tracks social interactive learning of a song. *Neuroimage* **183**,

280–290. (doi:10.1016/j.neuroimage.2018.08.005)

46.   Pan Y, Dikker S, Goldstein P, Zhu Y, Yang C, Hu Y. 2020 Instructor-learner brain

coupling discriminates between instructional approaches and predicts learning.

Neuroimage **211**, 116657. (doi:10.1016/j.neuroimage.2020.116657)

47. Babiloni F, Astolfi L. 2014 Social neuroscience and hyperscanning techniques: Past,

present   and   future.   *Neurosci.   Biobehav.   Rev.*   **44**,   76–93.

(doi:10.1016/j.neubiorev.2012.07.006)

48. Redcay E, Schilbach L. 2019 Using second-person neuroscience to elucidate the

mechanisms   of   social   interaction.   *Nat.   Rev.   Neurosci.*   **20**,   495–505.

(doi:10.1038/s41583-019-0179-4)

49. Parnamets P, Espinosa L, Olsson A. 2018 Physiological Synchrony Predicts

Observational Threat Learning in Humans. *bioRxiv* , 454819. (doi:10.1101/454819)

50. Bouton ME, Mineka S, Barlow DH. 2001 A modern learning theory perspective on the

etiology of panic disorder. *Psychol. Rev.* **108**, 4–32. (doi:10.1037/0033-295X.108.1.4)

51. Tschacher W, Meier D. 2019 Physiological synchrony in psychotherapy sessions.

Psychother. Res. , 1–16. (doi:10.1080/10503307.2019.1612114)

52. Woody ML, Feurer C, Sosoo EE, Hastings PD, Gibb BE. 2016 Synchrony of

physiological activity during mother-child interaction: moderation by maternal history

of   major   depressive   disorder.   *J.   Child   Psychol.   Psychiatry*   **57**,   843–850.

(doi:10.1111/jcpp.12562)

53. Button KS, Ioannidis JPA, Mokrysz C, Nosek BA, Flint J, Robinson ESJ, Munafò MR.

2013 Power failure: Why small sample size undermines the reliability of neuroscience.

Nat. Rev. Neurosci. **14**, 365–376. (doi:10.1038/nrn3475)

54. Boucsein W. 2012 Principles of Electrodermal Phenomena. In *Electrodermal Activity*,

pp. 1–86. Boston, MA: Springer US. (doi:10.1007/978-1-4614-1126-0_1)

55. Piazza EA, Hasenfratz L, Hasson U, Lew-Williams C. 2020 Infant and Adult Brains

Are Coupled to the Dynamics of Natural Communication. *Psychol. Sci.* **31**, 6–17.

(doi:10.1177/0956797619878698)

56.   Reddan MC, Young H, Falkner J, López-Solà M, Wager TD. 2020 Touch and social

648   support influence interpersonal synchrony and pain. *Soc. Cogn. Affect. Neurosci.*

649   (doi:10.1093/scan/nsaa048)

650   57.   Haaker J, Golkar A, Hermans D, Lonsdorf TB. 2014 A review on human reinstatement

651   studies: An overview and methodological challenges. *Learn. Mem.* **21**, 424–440.

652   (doi:10.1101/lm.036053.114)

653   58.   Faes L, Pinna GD, Porta A, Maestri R, Nollo G. 2004 Surrogate data analysis for

654   assessing the significance of the coherence function. *IEEE Trans. Biomed. Eng.* **51**,

655   1156–1166. (doi:10.1109/TBME.2004.827271)

656   59.   Zhou G, Bourguignon M, Parkkonen L, Hari R. 2016 Neural signatures of hand

657   kinematics in leaders vs. followers: A dual-MEG study. *Neuroimage* **125**, 731–738.

658   (doi:10.1016/j.neuroimage.2015.11.002)

659   60.   Vervliet B, Geens M. 2014 Fear generalization in humans: Impact of feature learning

660   on conditioning and extinction. *Neurobiol. Learn. Mem.* **113**, 143–148.

661   (doi:10.1016/j.nlm.2013.10.002)

662   61.   Cohen S, Wills TA. 1985 Stress, Social Support, and the Buffering Hypothesis.

663   *Psychol. Bull.* **98**, 310–357. (doi:10.1037/0033-2909.98.2.310)

664   62.   Palumbo R V, Marraccini ME, Weyandt LL, Wilder-Smith O, McGee HA, Liu S,

665   Goodwin MS. 2017 Interpersonal Autonomic Physiology: A Systematic Review of the

666   Literature. *Pers. Soc. Psychol. Rev.* **21**, 99–141. (doi:10.1177/1088868316628405)

667   63.   Liu Y, Piazza EA, Simony E, Shewokis PA, Onaral B, Hasson U, Ayaz H. 2017

668   Measuring speaker-listener neural coupling with functional near infrared spectroscopy.

669   *Sci. Rep.* **7**, 43293. (doi:10.1038/srep43293)

670   64.   Simony E, Honey CJ, Chen J, Lositsky O, Yeshurun Y, Wiesel A, Hasson U. 2016

671   Dynamic reconfiguration of the default mode network during narrative comprehension.

Nat. Commun. **7**, 12141. (doi:10.1038/ncomms12141)

65. Bilek E *et al.* 2015 Information flow between interacting human brains: Identification,

validation, and relationship to social expertise. *Proc. Natl. Acad. Sci. U. S. A.* **112**,

5207–5212. (doi:10.1073/pnas.1421831112)

66. Dikker S *et al.* 2017 Brain-to-Brain Synchrony Tracks Real-World Dynamic Group

Interactions in the Classroom. *Curr. Biol.* **27**, 1375–1380.

(doi:10.1016/j.cub.2017.04.002)

67. Leong V, Byrne E, Clackson K, Georgieva S, Lam S, Wass S. 2017 Speaker gaze

increases information coupling between infant and adult brains. *Proc. Natl. Acad. Sci.*

U. S. A. **114**, 13290–13295. (doi:10.1073/pnas.1702493114)

68. Wass S V., Whitehorn M, Marriott Haresign I, Phillips E, Leong V. 2020 Interpersonal

Neural Entrainment during Early Social Interaction. *Trends Cogn. Sci.* **24**, 329–342.

(doi:10.1016/j.tics.2020.01.006)

69. Golkar A, Bellander M, öhman A. 2013 Temporal properties of fear extinction - Does

time matter? *Behav. Neurosci.* **127**, 59–69. (doi:10.1037/a0030892)

70. Prenoveau JM, Craske MG, Liao B, Ornitz EM. 2013 Human fear conditioning and

extinction: Timing is everything...or is it? *Biol. Psychol.* **92**, 59–68.

(doi:10.1016/j.biopsycho.2012.02.005)

71. Pan Y, Cheng X. 2020 Two-person approaches to studying social interaction in

psychiatry: Uses and clinical relevance. *Front. Psychiatry* **11**, 301.

(doi:10.3389/fpsyt.2020.00301)

693