The electrophysiological index of the central executive determines verbal working memory capacity

Yuri G. Pavlov [1, 2], Boris Kotchoubey [1]

[1] Institute of Medical Psychology and Behavioral Neurobiology, University of Tübingen, 72076, Germany

[2] Department of Psychology, Ural Federal University, Department of Psychology, Ekaterinburg, 620000, Russian Federation

Corresponding author:

Yuri G. Pavlov

Email: pavlovug@gmail.com

Address: Institute of Medical Psychology and Behavioral Neurobiology, University of Tübingen, 72076, Germany

1 Abstract

Working memory (WM) consists of two main components: sensory storage and central executive. We studied cortical oscillatory correlates of these two components in a large sample of 156 participants to assess separately the contribution of them to individual differences in WM. The participants were presented with WM tasks of above-average to high complexity. Some of the tasks required only storage in WM, others required storage and mental manipulations. Our data indicate a close relationship between frontal midline theta, central beta activity and the central executive component of WM. The oscillatory counterparts of the central executive determined individual differences in verbal WM performance. In contrast, alpha activity was not related to the individual differences. The results demonstrate that central executive component of WM, rather than sensory storage capacity, play the decisive role in individual WM capacity limits.

Introduction

Working memory – our ability to maintain and manipulate information in a short term – is not a uniform construct. Baddeley & Hitch (1974) suggested that WM consists of separate storage buffers for memory content, and a central executive, which is responsible for maintenance of the active representation of the memory trace, attentional control, active processing and reorganization, or manipulation of information in the storage buffers. This additional component is what makes WM "working" and differentiates the WM in the narrow sense from mere short-term memory. With a few exceptions most WM tasks in real life require both sensory storage and executive components. For example, keeping a phone number in mind until it is dialed requires transformation of verbal information into a sequence of button presses. The task also involves switching attention between the current number to dial and pressing buttons on the phone panel. A more complex example would be the conversion of a shopping list into the optimal path while shopping. This task, in addition to the translation of the list into a sequence of spatial locations, also involves constant update of information in WM.

Available electrophysiological data converge on the importance of theta oscillations (4-8 Hz) in the medial prefrontal and anterior cingulate cortex for manipulation of information in WM (Berger et al., 2019; Berger, Omer, Minarik, Sterr, & Sauseng, 2014; Eschmann, Bader, & Mecklinger, 2018; Griesmayr, Gruber, Klimesch, & Sauseng, 2010; Itthipuripat, Wessel, & Aron, 2013; Kawasaki, Kitajo, & Yamaguchi, 2010; Kawasaki & Watanabe, 2007; Pavlov & Kotchoubey, 2017; Sauseng et al., 2005). Therefore, theta is thought to reflect engagement of the central executive component of WM (Sauseng, Griesmayr, Freunberger, & Klimesch, 2010). In turn, posterior alpha activity (8-13 Hz) has been associated with sensory storage processes, and may directly underlie maintenance of information in WM (Roux & Uhlhaas, 2014; van Ede, 2018). Alpha activity responds to increasing load reaching asymptote at the levels of individual's WM capacity (Fukuda, Mance, & Vogel, 2015) (although the effect of alpha "saturation" with load is less reliable in verbal WM tasks (Pavlov & Kochoubey, 2020)). Thus, posterior alpha activity can be seen as a reflection of the sensory storage component of WM.

The oscillatory mechanisms supporting individual WM capacity limits remain largely understudied. To study individual differences, large samples are essential. However, most available studies built their conclusions on samples of less than 30 individuals (Bashivan, Bidelman, & Yeasin, 2014; Fukuda et al., 2015; Hsieh, Ekstrom, & Ranganath, 2011; Hu et al., 2019; Kawasaki & Yamaguchi, 2013; Kwon et al., 2015; Maurer et al., 2015; Moran et al., 2010;

Roux, Wibral, Mohr, Singer, & Uhlhaas, 2012; Sghirripa et al., 2019). Better-powered studies (~35 participants per group) yielded mixed findings (Erickson, Smith, Albrecht, & Silverstein, 2019; Pahor & Jausovec, 2017; Pavlov & Kotchoubey, 2017; Zakrzewska & Brzezicka, 2014). For instance, after a median split, only the high-performance group showed an increase of theta with load (Pavlov & Kotchoubey, 2017; Zakrzewska & Brzezicka, 2014), suggesting that theta activity and the related executive functions are important for successful maintenance of WM. But another study did not find this relationship (Pahor & Jausovec, 2017). The relationship between alpha and beta activity and individual WM capacity is even less consistent.

The components of WM may contribute to individual differences in a different scale (Kofler, Rapport, Bolden, Sarver, & Raiker, 2010; Unsworth & Spillers, 2010). Here, we sought to provide solid data to set apart the contribution of oscillatory counterparts of sensory storage and central executive components of WM to individual differences. To reach this goal, we analyzed EEG data from a large number of participants (N=156) in WM tasks of slightly above average to very high complexity, which required either only retention of letter strings (Retention task) or retention and manipulation with the letters (mental recombination to the alphabetical order, Manipulation task) (see Figure 1 for the experimental design). Increasing the complexity of the tasks allowed us to increase the variance, to avoid ceiling effect in performance and to better distinguish between low and high performers. The task with mental manipulations allowed us to assess the contribution of different WM components. We hypothesized that (i) the power of alpha oscillations would be related to the capacity of the sensory storage component of WM, (ii) the power of frontal midline theta would be related to the efficiency of the executive component of WM, and (iii) the latter mechanism, but not the former one, would be related to the overall memory performance.

1   Results

2   Task performance

Figure 1 – (A) The experimental paradigm. (B) Average performance in 6 conditions. Error bars are
standard errors of the mean. (C) Distribution of overall performance stacked into 25 bins.

The accuracy was worse in the manipulation task and monotonically decreased with increasing
WM load (main effect of Task: $F(1, 155) = 431.8$, $\eta^2 = 0.736$, $p < 0.001$; Load: $F(2, 307) = 138.8$,
$\eta^2 = 0.472$, $p < 0.001$). The analyses of simple effects showed no difference between the
performance in the two most difficult tasks Manipulation of 6 and 7 letters as well between
Retention of 7 and Manipulation of 5 letters (Task x Load interaction: $F(2, 305) = 22.1$, $\eta^2 =$
0.125, $p < 0.001$). All the other pairwise differences were significant (see Table 1 for mean
values and Figure 1 for the visual depiction).

13   Table 1 – Descriptive statistics for accuracy

Load	Task	Mean % correct responses	SD
5	Manipulation	79.55	14.33
5	Retention	92.28	9.02
6	Manipulation	66.22	15.51
6	Retention	88.65	10.94
7	Manipulation	65.61	13.52
7	Retention	80.71	13.55

14 *Notes: SD – standard deviation*

Time-frequency analysis

We employed the linear mixed effects models (LME) to explore the relationship between behavioral performance and the spectral power in the theta frequency band. Although we used LME with a continuous Performance variable for statistical inference, for the ease of data interpretation and illustration purposes, the whole sample was median-split into the high performance (N = 78) and low performance (N = 78) groups on the basis of their average accuracy across all conditions.

As can be seen in Figure 2, the high performance group exhibited a stronger theta increase in the Manipulation task than in the Retention task. This observation was confirmed by a positive association between the relative theta power and behavioral performance in the Manipulation task and a negative association with performance in Retention task (Task by Performance interaction ($p = 0.004$), see Figure 2 and Supplementary Table 1 for full statistical output).

Then, we correlated WM performance with the difference between theta in Manipulation and Retention tasks to test the hypothesis on the role of executive components of WM in individual differences. The larger was the difference between baseline normalized theta power in Manipulation and Retention conditions, the better the individual WM performance (Spearman's rho = 0.24, $p = 0.002$) (see Figure 2C and Supplementary Figure 5 for full correlation matrix).

Figure 2 – Relationship between theta power at Fz and WM performance. (A) Time-frequency maps in low (top panels) and high (bottom panels) performance groups in retention and manipulation tasks. Semi-transparent boxes mark time-frequency windows of interest (4-8 Hz, last 6 s of delay). The first vertical bar marks the onset of encoding; the second one marks the onset of delay; and the third one marks the beginning of the time-frequency window of interest; and the fourth one marks the onset of retrieval. (B) The time dynamics of theta activity in Retention, Manipulation tasks and their difference: the result of subtraction of theta relative power in the retention task from the power in the manipulation task (central executive index). The width of the line represents 2 standard errors of the mean. (C) Left panel: correlation of WM performance in the manipulation and retention tasks. Right panel: correlation of WM performance and the central executive index.

No significant main effect or interactions with Performance were found in the alpha frequency band (see Figure 3 and Supplementary Table 2).

Figure 3 – Relationship between alpha power over posterior ROI (P3, T5, O1, P4, T6, O2) and WM performance. (A) Time-frequency maps in low (top panels) and high (bottom panels) performance groups in retention and manipulation tasks. Semi-transparent boxes mark time-frequency windows of interest (9-14 Hz, last 6 s of delay). The first vertical bar marks the onset of encoding; the second one marks the onset of delay; and the third one marks the beginning of the time-frequency window of interest; and the fourth one marks the onset of retrieval. (B) The time dynamics of theta activity in Retention, Manipulation tasks and their difference: the result of subtraction of theta relative power in the retention task from the power in the manipulation task (central executive index). The width of the line represents 2 standard errors of the mean. (C) Left panel: correlation of WM performance in the manipulation and retention tasks. Right panel: correlation of WM performance and the central executive index.

Beta power negatively related to WM performance in the Manipulation task but not in the Retention task (Task by Performance interaction: p<0.001; see Figure 4 and Supplementary Table 3). Like it has been done with the theta activity, a subtraction of Retention beta from

Manipulation beta was taken as an index of executive WM components. The correlation between this index and WM performance had similar magnitude as yielded in the analysis of theta (rho = -0.26, p = 0.001), but with the opposite sign (see Figure 4). Thus, better performance was related to lower beta activity.

Figure 4 – Relationship between beta power over central ROI (Cz, C4, C3) and WM performance. (A) Time-frequency maps in low (top panels) and high (bottom panels) performance groups in retention and manipulation tasks. Semi-transparent boxes mark time-frequency windows of interest (16-22 Hz, last 6 s of delay). The first vertical bar marks the onset of encoding; the second one marks the onset of delay; and the third one marks the beginning of the time-frequency window of interest; and the fourth one marks the onset of retrieval. (B) The time dynamics of theta activity in Retention, Manipulation tasks and their difference: the result of subtraction of theta relative power in the retention task from the power in the manipulation task (central executive index). The width of the line represents 2 standard errors of the mean. (C) Left panel: correlation of WM performance in the manipulation and retention tasks. Right panel: correlation of WM performance and the central executive index.

Then we tested whether the beta and theta central executive indices represent the same underlying mechanism. The correlation between the constructs was in an expected direction but not statistically significant (rho = -0.12, p = 0.124). Thus, theta and beta may play independent roles in individual differences in WM.

Discussion

We found that two EEG indices of the central executive component of WM (calculated as the difference between the relative spectral power in the retention and the manipulation conditions) significantly correlated with WM performance in a large sample of healthy individuals. These findings confirm our hypothesis (iii) stating that oscillatory associates of the executive WM component, but not of the storage component, are related to WM performance. This is congruent with previous behavioral research indicating that individual differences in executive functions were better predictors of WM capacity than differences in sensory storage (Engle, 2002; Shipstead, Lindsey, Marshall, & Engle, 2014; Unsworth & Spillers, 2010).

The positive correlation between WM performance and the theta central executive index, (i.e., more theta, better performance) also confirms our hypothesis (ii). With that, few studies demonstrated a correlation between theta activity and performance in verbal (Zakrzewska & Brzezicka (2014): N=69, r=0.32, Kwon et al. (2015): N=13, r=0.76) and visual (Kawasaki & Yamaguchi (2013): N=14, r=0.51; Maurer et al. (2015): N=24, r=-0.41) short-term memory tasks that require only retention of, but no manipulation with, stimulus material. In contrast, we found no correlation between theta activity in the Retention task and behavioral performance (rho = -0.01, p = 0.88). We hypothesized that the role of theta in retention tasks may be limited to only lower levels of load, but even at the lowest level of 5 letters the correlation between theta during retention and the accuracy was not significant (rho = -0.03, p = 0.67). Thus, theta increase during delay in simple tasks was not predictive to performance in genuine WM tasks.

Contrary to theta, the beta index of the executive control negatively correlated with WM performance (i.e., more beta, worse performance). This finding was not predicted by any hypothesis because the literature concerning this relationship is too scarce. Probably the only comparable study is a spatial WM MEG experiment (Proskovec, Wiesman, Heinrichs-Graham, & Wilson, 2018, 2019). A negative correlation between beta (15-20 Hz) and behavioral performance was reported in both publications. Although the two reports shared the same

sample, the localization of the correlation was different: WM performance correlated with beta activity that was localized in the right superior parietal lobule in one paper, and in the left dorsolateral prefrontal cortex and the bilateral superior temporal gyrus in the other. Moreover, beta activity in the two reports correlated with two different indexes of WM performance. These inconsistencies cast doubt on the reliability of the results.

Beta desynchronization may reflect switching internal attention between the reorganized set of letters and the initial set during the manipulations. Even when attention shifts from one object to another occur without eye movements, the shifts activate the same cortical areas as real eye movements (Grosbras & Paus, 2002). A good example is the activation of frontal eye fields (FEF) found in a spatial manipulation task (Glahn et al., 2002). If the beta activity observed in our study is related to activation of motor control networks, then why was it related to individual differences in WM? Repetitive saccade eye movements during delay were shown to increase episodic memory performance (Christman, Garvey, Propper, & Phaneuf, 2003; Lyle & Edlin, 2015). The movements executed just before blocks of an attentional control task (Flanker) also improved performance in the task (Edlin & Lyle, 2013). The attentional enhancement by preactivation of fronto-parietal network nodes such as FEF was hypothesized to produce the above mentioned effects (Edlin & Lyle, 2013). Consistent with this idea, TMS delivered to FEF improved detection of targets in a visuospatial attention task (Grosbras & Paus, 2002), whereas a suppression of the same region by TMS decreased inhibitory control (Muggleton, Chen, Tzeng, Hung, & Juan, 2010). It is admittedly a speculation at this point, but more efficient use of FEF to execute mental manipulations would potentially facilitate WM performance in our study.

Our hypothesis (i) predicted a significant link between alpha activity and the sensory storage capacity. This hypothesis was not confirmed. Although this link can be expected on the basis of theoretical considerations (as mentioned above in the Introduction), empirical data are not very consistent. Correlations between WM capacity and posterior alpha can be strongly negative (Kawasaki & Yamaguchi,2013: visual WM, N = 14, r = -0.66) or strongly positive (Kwon et al., 2015: verbal WM, N = 13, r = 0.75). Another study found no correlation between WM performance and relative alpha power in the Sternberg task (Sghirripa et al., 2019; N=24). Better WM performance was associated, in some studies, with stronger alpha enhancement in more difficult tasks than in the easy tasks (Hu et al., 2019: N = 20, r = 0.55; N = 23, r = 0.59), in other studies, however, with stronger alpha suppression in more difficult tasks (Erickson et al., 2019: N = 60, r = 0.45; Fukuda et al., 2015: N = 28, r = 0.48). In a yet another study the positive

correlation between the WM capacity and alpha modulation from lower to the higher load did not reach significance (Roux et al., 2012; N=25, r=0.39). We calculated the alpha modulation index (within-subject linear regression coefficients of alpha power by Load), as used in the verbal WM study by Hu et al. (2019). The test attained significance neither for the overall task performance (rho = 0.04, p = 0.64), nor for the retention task only (rho = 0.03, p = 0.72).

Therefore, the correlations obtained in the previous studies are not replicated in a larger sample. One might suggest that our negative result is due to using only high-load conditions, and that stronger correlations would be obtained if we compared alpha between high-load and low-load conditions. Nevertheless, the individuals with higher WM capacity did not show signs of larger sensory storage that would have been reflected in the modulation of alpha at the high levels of load. We suggest that instead it is the executive control as reflected in the theta activity that controls attention and minimizes the interference, thus determining an individual's WM capacity.

The interference hypothesis of WM capacity assumes that WM is affected not only by external distractors but also by mutual interference between the items (Berman, Jonides, & Lewis, 2009; Oberauer, Farrell, Jarrold, & Lewandowsky, 2016; Oberauer et al., 2018). Thus, the detrimental effect on memory precision is a function of the number of items stored in WM and the presence of the concurrent task. When WM load is high and/or manipulations are required, the executive component of WM is needed to cope with interference. Perhaps, the resources that can be deployed to counteract interference are better represented in the higher performing individuals.

There is another possible mechanism to explain how executive processes may cope with increasing WM demands. Camos & Barrouillet (2014) suggested that attention is involved in rapid refreshment of memory traces, thus preventing temporal decay of the WM content. The non-refreshed information fades out and in the worst case cannot be recovered. In extension of this model, beta and theta activity reflect the ability to maintain information in the active state by switching the focus of attention between items, thus defining the individual capacity limit.

Above we interpreted changes in posterior alpha rhythm as a reflection of the demand on the sensory storage component of WM. Another interpretation regards alpha as a mechanism of filtering information by gating only relevant sensory stimuli (Bonnefond & Jensen, 2012; Sauseng et al., 2009). If we accept this view, then either higher WM capacity individuals are no more efficient in the filtering out task-irrelevant distractors than lower WM capacity

individuals, or the alpha filtering mechanism works only under moderate levels of memory load.

To summarize, our data indicate a close relationship between medial frontal theta, central beta activity, and the central executive component of WM. Both oscillatory indices of the central executive, manifested in the theta (as we expected) and beta (unexpectedly) frequency bands, were related to behavioral performance in a verbal WM task. In contrast, we could not find evidence supporting the important role of posterior alpha in WM performance. We can conclude that the ability to control attention plays a larger role in individual differences in WM as compared to the capacity of the sensory storage.

Methods

Participants

186 individuals participated initially in the study. Eight participants with overall performance below 60% were excluded. Furthermore, a subsequent analysis revealed 22 EEG records with an excessive amount of artefacts (less than 12 clean trials in any condition). Thus, 156 participants (82 females, mean age = 21.23, SD=3.22) were included to the final sample. The participants had normal or corrected-to-normal vision and self-reported no history of neurological or mental diseases. All the participants were Russian native speakers. The experimental protocol was approved by the Ural Federal University ethics committee.

Stimuli

Sets of Russian alphabet letters written in capitals were used as stimuli. The letters had been selected randomly from the alphabet, had random order, and no repetitions in the sets. An analogue using Latin letters and English words is shown in Figure 1.

A trial always began with an exclamation mark presented for 200 ms, which was followed by a fixation cross for 3000 ms. Participants were instructed to fixate on the cross whenever it appeared. Next, the word "forward" or "alphabetical", presented for 600 ms, instructed the participants whether they would have to maintain in memory the original set as it was

presented (retention task) or, first, mentally reorganize the letters into the alphabetical order and then maintain the result in memory (manipulation task). After that, sets of 5, 6 or 7 letters were demonstrated for 3000 ms followed by a delay period where a fixation cross was demonstrated for 6700 ms. At the end of the delay period, a randomly chosen letter from the previously presented set appeared on the screen together with a digit that represented the serial number of this letter. The letter-digit probe was presented for 1000 ms. Depending on the task, the participants indicated whether the probe was on the corresponding position either in the original set (retention task), or in the set resulted from alphabetical reordering (manipulation task). The participants were asked to press one of the computer mouse buttons (left or right) if the probe was correct and the other button otherwise. The two buttons were attributed to the correct and wrong probes in a counterbalanced order. The probe was correct in 50 % of the trials, and the order of correct and incorrect probes was randomized. The next trial started after a blank interval that varied between 5000 and 5500 ms.

The experiment entailed six different conditions: maintenance in memory of 5, 6 or 7 letters in the alphabetical (manipulation condition) or forward (retention condition) order. Each condition had 20 consecutive trials. These six blocks of 20 trials were presented in a random order. Two practice blocks with 3 and 6 trials respectively were given shortly before the main experiment.

During the experiment, the participants were seated in a comfortable armchair in front of a computer screen in a dark room. Stimuli were presented in white color on a black background in the center of the screen by using PsyTask software (Mitsar Ltd.). The distance to the screen was 1 m and the size of the letters was 1.2 × 1.2°.

EEG recording and preprocessing

The EEG was recorded from 19 electrodes arranged according to the 10-20 system using Mitsar-EEG-202 amplifier with averaged earlobes reference. Two additional electrodes were used for horizontal and vertical EOG. EEG data were acquired with 500 Hz sampling rate and 150 Hz low-pass filter.

The procedure of EEG artifacts suppression and removal was conducted in two steps. At the first step, in order to suppress ocular activity artifacts, the independent component analysis (ICA) was performed using AMICA algorithm (Palmer, Kreutz-Delgado, & Makeig, 2012). The components clearly related to blinks and eye movements were identified and removed after

1 visual exploration of the data. At the second step, epochs still containing artefacts were visually

2 identified and discarded.

3 EEGLAB toolbox (Delorme & Makeig, 2004) for MATLAB was used for the data preprocessing.

Time-frequency analysis

Before the time-frequency analysis, 1 Hz high-pass, 45 Hz low-pass and 50 Hz notch filters were

applied with the EEGLAB firfilt function. Then, epochs in [-14200 2200 ms] interval where 0 is

the onset of the probe were created.

The EEG time series in each epoch were convolved with a set of complex Morlet wavelets. The

wavelets were defined as the product of a complex sine wave and a Gaussian window –

$e^{-i2\pi t f} e^{-t^2/(2\sigma^2)}$, where t is time, f is frequency. σ is the width of the Gaussian, which set

according to $n/(2\pi f)$, where n is the number of cycles – the parameter defining the time-

frequency precision trade-off. The frequency f increased from 1 to 45 Hz in 45 linearly spaced

steps, and the number of cycles n increased from 3 to 12 in 45 logarithmically spaced steps.

From the resulting complex signal, the power of each frequency at each time point was

obtained. The power was baseline-normalized by computing the percent change of the power

in respect to [-11500 -10500] ms interval (i.e. "Baseline" in Figure 1).

The time-frequency analysis was performed by means of the Fieldtrip toolbox (Oostenveld,

Fries, Maris, & Schoffelen, 2011).

Statistics

In order to decrease the number of factors employed in statistical calculations we defined

frequency-channels regions of interest (ROI) with maximal representation of certain

frequencies in certain group of channels. Thus, theta (4-8 Hz) had the maximal power in Fz,

alpha (9-14 Hz) in posterior channels (left: T5, P3, O1; right: T6, P4, O2) and beta (16-22 Hz) in

central channels (C3, Cz, C4) (see Supplementary materials for time-frequency maps at each

channel).

Then spectral power was averaged in the ROIs in a 6 s time interval corresponding to the delay

period (see Figure 3). We excluded the initial 700 ms from the onset of Delay period. As can be

seen in Figure 2 and Supplementary Figure 1, the first few hundred milliseconds after the onset

of the delay contain evoked response activity that could distort the frequency data if not

rejected (Babu Henry Samuel, Wang, Hu, & Ding, 2018; Ikkai, Blacker, Lakshmanan, Ewen, & Courtney, 2014; van Gerven, Bahramisharif, Heskes, & Jensen, 2009).

For statistical analysis of behavioral data, we employed repeated-measures analysis of variance (RM ANOVA) with factors Task (2 levels: Manipulation, Retention) and Load (3 levels: 5, 6, 7 letters to memorize).

For EEG data analyses we employed linear mixed-effects models (LME). As recommended by Barr et al. (2013), first, we always tried to fit the maximal model. In the case of the convergence problem, we first tried to fit the model with main effect random slopes (no interactions). If it converged, next, we again followed the advice by Barr (2013) and reduced the maximal random effect structure keeping the highest-order interaction slope (e.g., Task x Load x Hemisphere for alpha rhythm) and slopes that showed significant interactions with Performance at the previous step. Before fitting the models, continuous variables were centered around zero, and qualitative variables were effect coded.

For beta and theta rhythms the maximal model included Task and Load, Performance and all possible interactions as fixed effects, Participant as a grouping random intercept effect, and combination of Task, Load and their interactions as random slopes. For alpha activity fixed and random effects of Hemisphere (2 levels: left, right) were additionally included, because alpha frequently shows an asymmetry during delay period (Pavlov & Kotchoubey, 2020). Performance was calculated as the mean percentage of correct answers averaged over all conditions. In this case Performance plays the role of a personal trait and allows a straightforward interpretation of possible interactions with the other fixed effects. Single trial EEG data entered the analysis.

lme4 package (Bates, Maechler, Bolker, & Walker, 2014) and function lmer in *R* was used to fit the models. For the calculation of p-values we used lmerTest package (Kuznetsova, Brockhoff, & Christensen, 2017) with the default Satterthwaite's degrees of freedom approximation.

We decided to use the alpha level of 0.005, as such a threshold exhibits higher evidential value and may also help to improve reproducibility of newly discovered effects (e.g. see Benjamin et al. (2018) but also Miller & Ulrich (2019) for another opinion).

All statistical calculations were performed in *R,* version 3.6.3 (R Core Team, 2013).

CRediT authorship contribution statement

Yuri G. Pavlov: Funding acquisition, Resources, Supervision, Data curation, Formal analysis, Writing - original draft, Writing - review & editing.

Boris Kotchoubey: Writing - review & editing.

Acknowledgements

Study was supported by Russian Foundation for Basic Research (RFBR) #19-013-00027. The RFBR had no role in the writing and no influence on the contents of the manuscript. The authors declare no conflicts of interest.

References

Babu Henry Samuel, I., Wang, C., Hu, Z., & Ding, M. (2018). The frequency of alpha oscillations: Task-dependent modulation and its functional significance. *NeuroImage*, *183*, 897–906. https://doi.org/10/gfjpkr

Baddeley, A. D., & Hitch, G. (1974). Working memory. In *Psychology of learning and motivation* (Vol. 8, pp. 47–89). Elsevier.

Barr, D. J. (2013). Random effects structure for testing interactions in linear mixed-effects models. *Frontiers in Psychology*, *4*. https://doi.org/10.3389/fpsyg.2013.00328

Barr, D. J., Levy, R., Scheepers, C., & Tily, H. J. (2013). Random effects structure for confirmatory hypothesis testing: Keep it maximal. *Journal of Memory and Language*, *68*(3), 255–278.

Bashivan, P., Bidelman, G. M., & Yeasin, M. (2014). Spectrotemporal dynamics of the EEG during working memory encoding and maintenance predicts individual behavioral capacity. *European Journal of Neuroscience*, *40*(12), 3774–3784. https://doi.org/10.1111/ejn.12749

Bates, D., Maechler, M., Bolker, B., & Walker, S. (2014). lme4: Linear mixed-effects models using Eigen and S4. *R Package Version*, *1*(7), 1–23.

Benjamin, D. J., Berger, J. O., Johannesson, M., Nosek, B. A., Wagenmakers, E.-J., Berk, R., … Johnson, V. E. (2018). Redefine statistical significance. *Nature Human Behaviour*, *2*(1), 6–10. https://doi.org/10.1038/s41562-017-0189-z

Berger, B., Griesmayr, B., Minarik, T., Biel, A. L., Pinal, D., Sterr, A., & Sauseng, P. (2019). Dynamic regulation of interregional cortical communication by slow brain oscillations during working memory. *Nature Communications*, *10*(1), 1–11. https://doi.org/10.1038/s41467-019-12057-0

Berger, B., Omer, S., Minarik, T., Sterr, A., & Sauseng, P. (2014). Interacting Memory Systems—Does EEG Alpha Activity Respond to Semantic Long-Term Memory Access in a Working Memory Task? *Biology*, *4*(1), 1–16. https://doi.org/10.3390/biology4010001

Berman, M. G., Jonides, J., & Lewis, R. L. (2009). In Search of Decay in Verbal Short-Term Memory. *Journal of Experimental Psychology. Learning, Memory, and Cognition*, *35*(2), 317–333. https://doi.org/10.1037/a0014873

Bonnefond, M., & Jensen, O. (2012). Alpha Oscillations Serve to Protect Working Memory Maintenance against Anticipated Distracters. *Current Biology*, *22*(20), 1969–1974. https://doi.org/10/f4f3h2

Camos, V., & Barrouillet, P. (2014). Attentional and non-attentional systems in the maintenance of verbal information in working memory: The executive and phonological loops. *Frontiers in Human Neuroscience*, 8. https://doi.org/10.3389/fnhum.2014.00900

Christman, S. D., Garvey, K. J., Propper, R. E., & Phaneuf, K. A. (2003). Bilateral eye movements enhance the retrieval of episodic memories. *Neuropsychology*, *17*(2), 221–229. https://doi.org/10.1037/0894-4105.17.2.221

Delorme, A., & Makeig, S. (2004). EEGLAB: An open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of Neuroscience Methods*, *134*(1), 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Edlin, J. M., & Lyle, K. B. (2013). The effect of repetitive saccade execution on the attention network test: Enhancing executive function with a flick of the eyes. *Brain and Cognition*, *81*(3), 345–351. https://doi.org/10.1016/j.bandc.2012.12.006

Engle, R. W. (2002). Working memory capacity as executive attention. *Current Directions in Psychological Science*, *11*(1), 19–23.

Erickson, M. A., Smith, D., Albrecht, M. A., & Silverstein, S. (2019). Alpha-band desynchronization reflects memory-specific processes during visual change detection. *Psychophysiology*, *56*(11). https://doi.org/10.1111/psyp.13442

Eschmann, K. C. J., Bader, R., & Mecklinger, A. (2018). Topographical differences of frontal-midline theta activity reflect functional differences in cognitive control abilities. *Brain and Cognition*, *123*, 57–64. https://doi.org/10.1016/j.bandc.2018.02.002

Fukuda, K., Mance, I., & Vogel, E. K. (2015). Alpha Power Modulation and Event-Related Slow Wave Provide Dissociable Correlates of Visual Working Memory. *The Journal of Neuroscience*, *35*(41), 14009–14016. https://doi.org/10.1523/jneurosci.5003-14.2015

Griesmayr, B., Gruber, W. R., Klimesch, W., & Sauseng, P. (2010). Human frontal midline theta and its synchronization to gamma during a verbal delayed match to sample task. *Neurobiology of Learning and Memory*, *93*(2), 208–215. https://doi.org/10.1016/j.nlm.2009.09.013

Grosbras, M.-H., & Paus, T. (2002). Transcranial Magnetic Stimulation of the Human Frontal Eye Field: Effects on Visual Perception and Attention. *Journal of Cognitive Neuroscience*, *14*(7), 1109–1120. https://doi.org/10.1162/089892902320474553

Hsieh, L. T., Ekstrom, A. D., & Ranganath, C. (2011). Neural oscillations associated with item and temporal order maintenance in working memory. *The Journal of Neuroscience*, *31*(30), 10803–10810. https://doi.org/10.1523/jneurosci.0828-11.2011

Hu, Z., Barkley, C. M., Marino, S. E., Wang, C., Rajan, A., Bo, K., ... Ding, M. (2019). Working Memory Capacity Is Negatively Associated with Memory Load Modulation of Alpha Oscillations in Retention of Verbal Working Memory. *Journal of Cognitive Neuroscience*, 1–13. https://doi.org/10.1162/jocn_a_01461

Ikkai, A., Blacker, K. J., Lakshmanan, B. M., Ewen, J. B., & Courtney, S. M. (2014). Maintenance of relational information in working memory leads to suppression of the sensory cortex. *Journal of Neurophysiology*, *112*(8), 1903–1915. https://doi.org/10.1152/jn.00134.2014

Itthipuripat, S., Wessel, J. R., & Aron, A. R. (2013). Frontal theta is a signature of successful working memory manipulation. *Experimental Brain Research*, *224*(2), 255–262. https://doi.org/10.1007/s00221-012-3305-3

Kawasaki, M., Kitajo, K., & Yamaguchi, Y. (2010). Dynamic links between theta executive functions and alpha storage buffers in auditory and visual working memory. *European Journal of Neuroscience*, *31*(9), 1683–1689. https://doi.org/10.1111/j.1460-9568.2010.07217.x

Kawasaki, M., & Watanabe, M. (2007). Oscillatory gamma and theta activity during repeated mental manipulations of a visual image. *Neuroscience Letters*, *422*(2), 141–145. https://doi.org/10.1016/j.neulet.2007.04.079

Kawasaki, M., & Yamaguchi, Y. (2013). Frontal theta and beta synchronizations for monetary reward increase visual working memory capacity. *Social Cognitive and Affective Neuroscience*, *8*(5), 523–530. https://doi.org/10.1093/scan/nss027

Kofler, M. J., Rapport, M. D., Bolden, J., Sarver, D. E., & Raiker, J. S. (2010). ADHD and Working Memory: The Impact of Central Executive Deficits and Exceeding Storage/Rehearsal Capacity on Observed Inattentive Behavior. *Journal of Abnormal Child Psychology*, *38*(2), 149–161. https://doi.org/10.1007/s10802-009-9357-6

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest package: Tests in linear mixed effects models. *Journal of Statistical Software*, *82*(13). https://doi.org/10.18637/jss.v082.i13

Kwon, G., Lim, S., Kim, M. Y., Kwon, H., Lee, Y. H., Kim, K., ... Suh, M. (2015). Individual differences in oscillatory brain activity in response to varying attentional demands during a word recall and oculomotor dual task. *Frontiers in Human Neuroscience*, *9*, 381. https://doi.org/10.3389/fnhum.2015.00381

Lyle, K. B., & Edlin, J. M. (2015). Why does saccade execution increase episodic memory

retrieval? A test of the top-down attentional control hypothesis. *Memory*, *23*(2), 187–

202. https://doi.org/10.1080/09658211.2013.877487

Maurer, U., Brem, S., Liechti, M., Maurizio, S., Michels, L., & Brandeis, D. (2015). Frontal midline

theta reflects individual task performance in a working memory task. *Brain Topography*,

28(1), 127–134. https://doi.org/10.1007/s10548-014-0361-y

Miller, J., & Ulrich, R. (2019). The quest for an optimal alpha. *PLOS ONE*, *14*(1), e0208631.

https://doi.org/10.1371/journal.pone.0208631

Moran, R. J., Campo, P., Maestú, F., Reilly, R. B., Dolan, R. J., & Strange, B. A. (2010). Peak

Frequency in the Theta and Alpha Bands Correlates with Human Working Memory

Capacity. *Frontiers in Human Neuroscience*, *4*. https://doi.org/10/fk9dfr

Muggleton, N. G., Chen, C.-Y., Tzeng, O. J. L., Hung, D. L., & Juan, C.-H. (2010). Inhibitory Control

and the Frontal Eye Fields. *Journal of Cognitive Neuroscience*, *22*(12), 2804–2812.

https://doi.org/10.1162/jocn.2010.21416

Oberauer, K., Farrell, S., Jarrold, C., & Lewandowsky, S. (2016). What limits working memory

capacity? *Psychological Bulletin*, *142*(7), 758.

Oberauer, K., Lewandowsky, S., Awh, E., Brown, G. D., Conway, A., Cowan, N., … Hurlstone, M. J.

(2018). Benchmarks for models of short-term and working memory. *Psychological

Bulletin*, *144*(9), 885.

Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J.-M. (2011). FieldTrip: Open source software

for advanced analysis of MEG, EEG, and invasive electrophysiological data.

Computational Intelligence and Neuroscience, *2011*, 1.

Pahor, A., & Jausovec, N. (2017). Multifaceted pattern of neural efficiency in working memory

capacity. *Intelligence*, *65*, 23–34. https://doi.org/10.1016/j.intell.2017.10.001

Palmer, J. A., Kreutz-Delgado, K., & Makeig, S. (2012). AMICA: An adaptive mixture of

independent component analyzers with shared components. *Swartz Center for

Computatonal Neursoscience, University of California San Diego, Tech. Rep*. Retrieved

from https://sccn.ucsd.edu/~jason/amica_a.pdf

Pavlov, Y. G., & Kotchoubey, B. (2017). EEG correlates of working memory performance in

females. *BMC Neuroscience*, *18*(1), 26. https://doi.org/10.1186/s12868-017-0344-5

Pavlov, Y., & Kochoubey, B. (2020). *Oscillatory brain activity and maintenance of verbal working

memory: A systematic review* [Preprint]. https://doi.org/10.31234/osf.io/mn53j

Proskovec, A. L., Wiesman, A. I., Heinrichs-Graham, E., & Wilson, T. W. (2018). Beta Oscillatory Dynamics in the Prefrontal and Superior Temporal Cortices Predict Spatial Working Memory Performance. *Scientific Reports*, *8*(1), 8488. https://doi.org/10/gdp4ww

Proskovec, A. L., Wiesman, A. I., Heinrichs-Graham, E., & Wilson, T. W. (2019). Load effects on spatial working memory performance are linked to distributed alpha and beta oscillations. *Human Brain Mapping.* https://doi.org/10.1002/hbm.24625

R Core Team. (2013). *R: A language and environment for statistical computing.*

Roux, F., & Uhlhaas, P. J. (2014). Working memory and neural oscillations: Alpha-gamma versus theta-gamma codes for distinct WM information? *Trends in Cognitive Sciences*, *18*(1), 16–25. https://doi.org/10.1016/j.tics.2013.10.010

Roux, F., Wibral, M., Mohr, H. M., Singer, W., & Uhlhaas, P. J. (2012). Gamma-Band Activity in Human Prefrontal Cortex Codes for the Number of Relevant Items Maintained in Working Memory. *Journal of Neuroscience*, *32*(36), 12411–12420. https://doi.org/10.1523/jneurosci.0421-12.2012

Sauseng, P., Griesmayr, B., Freunberger, R., & Klimesch, W. (2010). Control mechanisms in working memory: A possible function of EEG theta oscillations. *Neuroscience & Biobehavioral Reviews*, *34*(7), 1015–1022. https://doi.org/10.1016/j.neubiorev.2009.12.006

Sauseng, P., Klimesch, W., Doppelmayr, M., Pecherstorfer, T., Freunberger, R., & Hanslmayr, S. (2005). EEG alpha synchronization and functional coupling during top-down processing in a working memory task. *Human Brain Mapping*, *26*(2), 148–155. https://doi.org/10.1002/hbm.20150

Sauseng, P., Klimesch, W., Heise, K. F., Gruber, W. R., Holz, E., Karim, A. A., … Hummel, F. C. (2009). Brain Oscillatory Substrates of Visual Short-Term Memory Capacity. *Current Biology*, *19*(21), 1846–1852. https://doi.org/10.1016/j.cub.2009.08.062

Sghirripa, S., Graetz, L., Merkin, A., Rogasch, N. C., Semmler, J. G., & Goldsworthy, M. R. (2019). Load-dependent modulation of alpha oscillations during working memory encoding and retention in young and older adults. *BioRxiv*, 848127. https://doi.org/10.1101/848127

Shipstead, Z., Lindsey, D. R. B., Marshall, R. L., & Engle, R. W. (2014). The mechanisms of working memory capacity: Primary memory, secondary memory, and attention control. *Journal of Memory and Language*, *72*, 116–141. https://doi.org/10.1016/j.jml.2014.01.004

Unsworth, N., & Spillers, G. J. (2010). Working memory capacity: Attention control, secondary memory, or both? A direct test of the dual-component model. *Journal of Memory and Language*, *62*(4), 392–406. https://doi.org/10.1016/j.jml.2010.02.001

van Ede, F. (2018). Mnemonic and attentional roles for states of attenuated alpha oscillations in perceptual working memory: A review. *The European Journal of Neuroscience*, *48*(7), 2509–2515. https://doi.org/10.1111/ejn.13759

van Gerven, M., Bahramisharif, A., Heskes, T., & Jensen, O. (2009). Selecting features for BCI control based on a covert spatial attention paradigm. *Neural Networks*, *22*(9), 1271–1277. https://doi.org/10.1016/j.neunet.2009.06.004

Zakrzewska, M. Z., & Brzezicka, A. (2014). Working memory capacity as a moderator of load-related frontal midline theta variability in Sternberg task. *Frontiers in Human Neuroscience*, *8*, 399. https://doi.org/10.3389/fnhum.2014.00399