TITLE: INSIDE-OUT REGULATION OF E-CADHERIN CONFORMATION AND ADHESION

AUTHORS: Ramesh Koirala[1,2,#], Andrew Vae Priest[1,2,#], Chi-Fu Yen[2], Joleen S. Cheah[1], Willem-Jan Pannekoek[3], Martijn Gloerich[3], Soichiro Yamada[1], Sanjeevi Sivasankar[1]*

AFFILIATIONS: [1]Department of Biomedical Engineering, University of California, Davis, CA 95616; [2]Department of Physics and Astronomy, Iowa State University, Ames, IA 50011; [3]Molecular Cancer Research, Center for Molecular Medicine, University Medical Center Utrecht, Utrecht, The Netherlands.

[#] Equal contribution

* CORRESPONDING AUTHOR: ssivasankar@ucdavis.edu

ABSTRACT

Cadherin cell-cell adhesion proteins play key roles in tissue morphogenesis and wound

healing. Cadherin ectodomains bind in two conformations, X-dimers and strand-swap

dimers, with different adhesive properties. However, the mechanisms by which cells

regulate ectodomain conformation are unknown. Cadherin intracellular regions associate

with several actin-binding proteins including vinculin, which are believed to tune cell-cell

adhesion by remodeling the actin cytoskeleton. Here, we show at the single molecule

level, that vinculin association with the cadherin cytoplasmic region allosterically converts

weak X-dimers into strong strand-swap dimers, and that this process is mediated by

myosin II dependent changes in cytoskeletal tension. We also show that in epithelial cells,

~70% of apical cadherins exist as strand-swap dimers while the remaining form X-dimers,

providing two cadherin pools with different adhesive properties. Our results demonstrate,

for the first time, the inside-out regulation of cadherin conformation and establish a

mechanistic role for vinculin in this process.

SIGNIFICANCE STATEMENT

Cadherin cell-cell adhesion proteins play key roles in the formation and maintenance of

tissues. Their adhesion is carefully regulated to orchestrate complex movement of cells.

While cadherin ectodomains bind in two conformations with different adhesive properties,

the mechanisms by which cells regulate the conformation (and consequently adhesion) of

individual cadherins are unknown. Here, we demonstrate that the association of

intracellular vinculin to the cadherin cytoplasmic region, regulates cadherin adhesion by

switching ectodomains from a weak binding to the strongly adhesive conformation. In

contrast with the prevailing view which suggests that vinculin regulates adhesion solely by

remodeling the cytoskeleton, we show that vinculin can directly modulate single cadherin

ectodomain conformation and that this process is mediated by changes in cytoskeletal

tension.

INTRODUCTION

E-cadherins (Ecads) are essential, calcium dependent cell-cell adhesion proteins that play key roles in the formation of epithelial tissue and in the maintenance of tissue integrity. Ecad adhesion is highly plastic and carefully regulated to orchestrate complex movement of epithelial cells and dysregulation of adhesion is a hallmark of numerous cancers [1]. However, little is known about how cells dynamically regulate the biophysical properties of individual Ecads.

The extracellular region of Ecads from opposing cells bind in two distinct *trans* orientations: strand-swap dimers and X-dimers (Figure 1A-B). Strand-swap dimers are the stronger cadherin adhesive conformation and are formed by the exchange of conserved Tryptophan (Trp) residues between the outermost domains of opposing Ecads [2-4]. In contrast, X-dimers, which are formed by extensive surface interactions between opposing Ecads, are a weaker adhesive structure and serve as an intermediate during the formation and rupture of strand swap dimers [5-7]. Using cell-free, single molecule experiments, we previously showed that X-dimers and strand-swap dimers can be distinguished based on their distinctly different response to mechanical force. When a strand-swap dimer is pulled, its lifetime decreases with increasing force, resulting in the formation of a slip bond [8,9] (Figure 1B). In contrast, an X-dimer responds to pulling force by forming a catch bond, where bond lifetime initially increases up to a threshold force, and then subsequently decreases [8,10] (Figure 1B). It has also been shown that wild type Ecad ectodomains in solution can interconvert between X-dimer and strand-swap dimer conformations [9,11]. However, the biophysical mechanisms by which Ecad conformations (and adhesion) are regulated on the cell surface are unknown.

The cytoplasmic region of Ecad associates with the catenin family of proteins:

namely p120-catenin, β-catenin and α-catenin. The Ecad-catenin complex, in turn, links

to filamentous actin (F-actin) either by the direct binding of α-catenin and F-actin or by the

indirect association of α-catenin and F-actin via vinculin [12] (Figure 1A). Adhesive forces

transmitted across intercellular junctions by Ecad induce conformational changes in α-

catenin [13,14], strengthen F-actin binding [15] and recruit vinculin to the sites of force

application [16,17]. However, vinculin and α-catenin do not merely serve as passive

cytoskeletal linkers; they also dynamically modulate cytoskeletal rearrangement and

recruit myosin to cell-cell junctions [13,18-20]. Studies show that α-catenin and vinculin play

important roles in strengthening and stabilizing Ecad adhesion: bead twisting experiments

show force-induced stiffening of Ecad-based junctions and cell doublet stretching

experiments demonstrate reinforcement of cell-cell adhesion in vinculin and α-catenin

dependent manners [18,19,21].

Currently, actin anchorage and cytoskeletal remodeling are assumed to be the

exclusive mechanisms by which α-catenin and vinculin strengthen Ecad adhesion [22-24].

Here, we directly map the allosteric effects of cytoplasmic proteins on Ecad ectodomain

conformation and demonstrate, at the single molecule level, that vinculin association with

the Ecad cytoplasmic region switches X-dimers to strand-swap dimers. We show that

cytoskeletal tension, due to vinculin mediated recruitment of myosin II, regulates Ecad

ectodomain structure and adhesion. Finally, we demonstrate that only ~50% of Ecads are

linked to the underlying cytoskeleton and that while about 70% of Ecads form strand-swap

dimers, the remaining form X-dimers, which provides cells with two Ecad pools with

different adhesive properties.

RESULTS

Single molecule Atomic Force Microscope (AFM) measurements of specific Ecad
trans **adhesion.** We measured interactions between recombinant Ecad extracellular regions immobilized on AFM cantilevers and Ecad endogenously expressed on the apical surface of confluent Madin-Darby Canine Kidney (MDCK) cell monolayers (Figure 2A). To ensure that we measured single Ecad unbinding events, we immobilized Ecad on the AFM tip at low density and pressed on the cell surface with a low initial contact force (mean pressing force = 59 pN; Figure S1). Poisson statistics predict that even the maximum event rate we measure (19%; Figure 2B) corresponds to ~90% probability of measuring single molecule events.

Our experiments were performed with the following MDCK cells (Figure 1C): parental cells (*parental*), vinculin knockout cells (*vinc-KO*), α-catenin knockout cells (*αcat-KO*) [25], αcat-KO cells rescued with α-catenin lacking the vinculin binding site (*ΔVBS*) [26], and Ecad knockout cells (*Ecad-KO*). We also performed measurements with parental cells in the presence of free tryptophan (*parental-Trp*) in order to trap Ecad in an X-dimer conformation [8]. Additionally, we performed experiments in the presence of blebbistatin (*parental-Bleb*) in order to reduce myosin II dependent cytoskeletal contractility. We quantified the specific and nonspecific binding rates for every cell line by measuring single unbinding events using either AFM tips that were decorated with Ecad (specific binding) or AFM tips that lacked Ecad (nonspecific binding). All cell lines, except Ecad-KO, showed a significant increase in single unbinding events when the AFM tips were decorated with Ecad (Figure 2B). In contrast, in the Ecad-KO cells, event rates using AFM tips with/without Ecad were similar to each other and were comparable to the nonspecific

event rates measured with the other cell lines (Figure 2B). Taken together, this confirmed

that the measured specific events corresponded to Ecad-Ecad binding interactions.

Ecad conformation and cytoskeletal-linkage in each cell line were determined from

previously reported 'signatures' in the measured force curves. Since force measurements

have shown that pulling on a cell surface protein that is not linked to the cytoskeleton

results in the formation of a *membrane-tether* [27,28] (Figure 3A), we used membrane-tethers

to identify Ecad that are uncoupled from the actin cytoskeleton (Figure 3B). Previous

studies have also shown that pulling on a transmembrane protein that is firmly bound to

the underlying cytoskeleton results in a linearly elastic response to pulling force [29,30]

(Figure 3C). Since these *jump* events occur when interactions between proteins on the

AFM tip and cell surface are weaker than the protein-cytoskeleton linkage, we interpreted

jumps as a signature of actin-linked Ecads that form weak *trans* dimers, which rupture

before failure of the Ecad-cytoskeletal linkage (Figure 3D). Finally, if the Ecad *trans* dimers

adhere robustly compared to the Ecad-cytoskeletal linkage, an initial steep increase in

force is measured which relaxes viscoelastically when the Ecad-cytoskeletal bond

ruptures, resulting in the formation of a *cyto-tether* (Figure 3E-F). To further validate tether

formation, we fit the membrane-tether and cyto-tether force-extension curves to the

standard linear solid (SLS) model (SI Note 1) [31]. The SLS fitting parameters we obtained

for the tethers, were consistent with previous values obtained for tether formation on

Jurkat cells and T-cells (Table S1) [31,32].

About 50% of non-junctional Ecads are coupled to the actin cytoskeleton. We first

compared the fraction of Ecad uncoupled from the cytoskeleton (membrane-tethers) to

the fraction of Ecads that are coupled to cortical actin (cyto-tethers and jumps). Our data

showed that in parental cells, 55% ± 9% of Ecad on the apical surface of the cell are

uncoupled from the actin cytoskeleton (Figure 4A). Since α-catenin is believed to be vital

in mediating Ecad linkage to F-actin, we next measured the cytoskeletal linkage of Ecad

in αcat-KO cells. As anticipated from previous biochemical and cell biological results [33,34],

our data showed that the vast majority of Ecad (92% ± 12%) in αcat-KO cells are

uncoupled from the actin cytoskeleton (Figure 4A). In contrast, the fraction of uncoupled

Ecads in both vinc-KO and ΔVBS cells, where Ecads are presumably weakly linked to F-

actin by α-catenin, but where this linkage is not reinforced by vinculin [13,35], was comparable

to parental cells (65% ± 9% for vinc-KO and 45% ± 4% for ΔVBS cells; Figure 4A). Taken

together, this data confirms the obligatory role of α-catenin in mediating Ecad linkage to

F-actin. These direct, single molecule measurements of Ecad-actin coupling on the apical

surface of live cells are in excellent agreement with previous single particle tracking of

Ecad on the cell surface that inferred approximately 50% of Ecads on the apical surface

of cells are coupled to the actin cytoskeleton [36].

Absence of vinculin binding weakens adhesion of individual Ecad. We next

compared the fraction of weak (jumps) and strong (cyto-tethers) *trans* dimers, formed by

cytoskeleton-coupled Ecad. Our data showed that in parental cells that contain both α-

catenin and vinculin, 77% ± 16% of cytoskeleton-coupled Ecad bind robustly while the

remaining 23% ± 14% of Ecad interact weakly (Figure 4B). In contrast, in vinc-KO cells,

only 5% ± 3% of cytoskeleton-coupled Ecad bind strongly (Figure 4B). Similarly, in ΔVBS

cells where vinculin cannot bind to α-catenin, only 2% ± 1% of cytoskeleton-linked Ecad

bind in a robust binding conformation (Figure 4B). Taken together, this indicates that

vinculin association with the Ecad cytoplasmic region is required for robust Ecad *trans*

dimerization. It is important to point out that the strength of *trans* dimer adhesion in the

absence of α-catenin could not be determined by comparing the fraction of cyto-tethers

and jumps because these events were extremely rare in the αcat-KO cells.

163   Since an Ecad *trans* dimer can exist in two different conformations with different

adhesive strengths — weaker X-dimer and stronger strand-swap dimer — we asked

whether the weak (jumps) and strong (cyto-tethers) adhesive states of cytoskeleton-

coupled Ecad represent different Ecad conformations. Since we have previously shown

that Ecad ectodomains can be trapped in an X-dimer conformation simply by incubating

them in free Trp [8], we performed AFM measurements on parental-Trp cells to identify the

signature of X-dimer formed by cytoskeleton-coupled Ecad. Interestingly, almost all (99%

± 13%) cytoskeleton-bound Ecads in parental-Trp formed jump events (Figure 4B),

suggesting that the jump events represent the Ecads coupled to the cytoskeleton that form

X-dimers. This, along with exclusive presence of jumps in vinc-KO and ΔVBS (Figure 4B),

indicates that the weakening of Ecad adhesion in the absence of vinculin binding is

consistent with Ecad being trapped in an X-dimer conformation. Importantly, switching

ectodomains into an X-dimer conformation did not decouple Ecad from the cytoskeleton

and the fraction of cytoskeleton-coupled Ecad in the Trp treated cells (52% ± 5%)

remained the same as parental cells (Figure 4A).

178   We also inferred the *trans* dimer binding conformation of cytoskeleton-coupled

Ecad in the parental-Trp, vinc-KO and ΔVBS cells using dynamic force spectroscopy

analysis (DFS) [37]. Since the number of jump events in vinc-KO cells were insufficient to

obtain statistically reliable results, we combined the vinc-KO jumps with the jumps

measured with ΔVBS cells. The measured k_{off} for jump events for vinc-KO and ΔVBS (k_{off}

= 1.77 ± 0.51 s^{-1}; Figure 4C) was comparable to parental-Trp X-dimers (k_{off} = 1.73 ± 0.62

s^{-1}; Figure 4D), and higher than previously published values for strand-swap dimers [38-41].

Since a higher k_{off} corresponds to a weaker binding, these data further suggest that in the

absence of vinculin association, cytoskeleton-coupled Ecad adopts a weaker X-dimer
conformation.

Next, since cadherins trapped in an X-dimer conformation have delayed junction
formation [42], we monitored the formation of cell-cell barriers in parental-Trp and ΔVBS
cells using Electric Cell-Substrate Impedance Sensing (ECIS), in order to determine if the
junction formation in these cell lines are consistent with Ecad adopting an X-dimer
conformation. Our experiments measuring the barrier formation half time (T50), showed a
significant delay in calcium-induced barrier formation in parental-Trp and ΔVBS cells
compared to parental cells (parental-Trp 1.23 ± 0.65 hr delay in T50, ΔVBS 3.66 ± 2.26 hr
delay in T50; Figure 4F). Importantly, delayed barrier formation with ΔVBS was consistent
with previous reports [26]. Taken together, our ECIS results indicate that both Ecad
conformation and vinculin binding to α-catenin are important for the formation of cadherin
junctions, and further suggest that vinculin is responsible for regulating Ecad
conformation.

Ecad ectodomain conformation is regulated by vinculin. Next, we proceeded to
directly measure the binding conformation of Ecad ectodomains in the different cell lines
by measuring if the Ecads on the cell surface form catch bonds (X-dimers) or slip bonds
(strand-swap dimers). We measured the force *vs.* lifetimes of Ecad-Ecad bonds, using a
method described previously [43]. Briefly, since the force required to rupture membrane-
tethers and cyto-tethers scales linearly with pulling velocity [28,44] (Figure S2), and because
these tethers extend at a constant force (after any initial viscoelastic relaxation) when
pulled at a constant velocity [27,31] (Figure 3A, 3E), we determined bond lifetimes at a range
of forces from the persistence time of membrane-tethers and cyto-tethers.

We first measured the force *vs.* lifetime of Ecad endogenously expressed on the

apical surface of parental cells. Fits of our data to the composite of a slip bond [45] and catch

bond [46] model showed that 69% of Ecad in the parental cells form strand-swap dimers

while the remaining 31% of Ecad occupy an X-dimer conformation (Figure 5A; Methods;

Table S2). Upon Trp incubation, however, Ecads in parental cells switched their

conformation and formed catch bonds characteristic of an X-dimer (Figure 5B).

Since the bond-lifetimes for parental cells (Figure 5A) were obtained from all the

tethers (a mix of cyto-tethers and membrane-tethers), we separately analyzed the force-

dependent lifetimes of cyto-tether and membrane-tether events (Figure S3A, B).

Consistent with cyto-tethers corresponding to strong Ecad-Ecad bonds, our analysis

showed that cyto-tethers (which correspond to Ecad initially coupled to the cytoskeleton)

exclusively formed slip-bonds. Ecad that were decoupled from the cytoskeleton

(membrane-tether events) formed X-dimers and strand-swap dimers with almost equal

probabilities (45% catch bonds: 55% slip bonds) (Figure S3A, B). This provided further

evidence that the association of vinculin with the Ecad cytoplasmic domain is required for

ectodomains to efficiently form strand-swap dimers.

As anticipated from our DFS experiments, Ecads in the vinc-KO cells formed catch

bonds characteristic of an X-dimer conformation (Figure 5C) suggesting that vinculin

association is necessary for Ecads to form robust strand-swap dimers. Similarly, Ecads in

αcat-KO cells, where vinculin cannot associate with the Ecad cytoplasmic region [12], also

formed catch bonds (Figure 5D). Finally, eliminating vinculin binding to α-catenin in ΔVBS

cells also resulted in catch bonds (Figure 5E), confirming that association of vinculin with

the Ecad cytoplasmic region via α-catenin is obligatory for transitioning from a weak X-

dimer conformation to a robust strand-swap dimer conformation.

Regulation of Ecad ectodomain conformation is myosin II dependent. Having

identified vinculin as a key protein involved in allosterically regulating Ecad ectodomain

conformation, we proceeded to identify the mechanism by which vinculin regulates Ecad

structure. Since vinculin localizes to force bearing sites in the cell-cell junction and

mediates the recruitment of myosin II [18], we hypothesized that Ecad ectodomain

conformation could be regulated by myosin II mediated cytoskeletal contractile force. We

therefore tested the effect of myosin II mediated force on Ecad conformation in parental

cells by reducing cytoskeleton contractility using the myosin II inhibitor, blebbistatin

(*parental-Bleb*). Strikingly, in parental-Bleb cells, the fraction of Ecad in strand-swap dimer

conformation (43%; Figure 5F) decreased compared to parental cells (69% strand-swap

dimer; Figure 5A). This indicated that myosin II dependent cytoskeletal contractility plays

a role in driving strand-swap dimerization of Ecad ectodomains.

Additionally, unlike parental cells where the majority of cytoskeleton-coupled Ecad

existed in the robust binding conformation (cyto-tethers), most of the cytosekeleton-

coupled Ecad in parental-Bleb formed weak binding structures (jumps) (27% ± 7% cyto-

tethers, 73% ± 25% jumps; Figure 4B). Importantly, the fraction of cytoskeleton-coupled

Ecad in parental-Bleb (42% ± 9%; Figure 4A) remained the same as in parental cells.

Finally, the measured k_{off} for jump events in parental-Bleb (2.37 ± 0.58 s^{-1}; Figure 4E) was

comparable to the values measured for X-dimers formed by parental-Trp, vinc-KO and

ΔVBS cells (Figure 4C-D). Taken together, these data confirm that reduced myosin II

dependent cytoskeletal contractility reduces the ability of Ecad to form robust strand-swap

dimers.

Since forces generated by the actomyosin cytoskeleton can transmit to the Ecad

cytoplasmic domain [47], we used steered molecular dynamics (SMD) simulations to test if

force could convert X-dimers to strand-swap dimers (Figure 6A-B). We modeled

cytoskeletal forces by pulling on the C-terminus of the EC1-2 X-dimer crystal structure

(PDB ID: 3LNH) at five angles (0°, ±30°, ±40°), chosen to account for different orientations

of actin filaments (Figure 6B). Our simulations showed that regardless of the pulling angle,

all X-dimers converted to strand-swap dimers after a period of force application (Figure

6B-C). Root mean squared deviations (RMSDs) of the SMD structures, calculated relative

to the strand-swap dimer crystal structure (PDB ID: 2QVF), converged to an average of

0.67 ± 0.12 nm confirming that the structures closely resembled strand-swap dimers

(Figure 6C). These simulations confirm that when Ecad dimers are firmly coupled to the

actin cytoskeleton by α-catenin and vinculin, myosin-II dependent force can convert X-

dimers to strand-swap dimers.

DISCUSSION

Using live-cell, single molecule AFM measurements, we directly demonstrate

inside-out regulation of cadherin conformation and show that cytoplasmic proteins

regulate Ecad adhesive properties, similar to inside-out regulation of integrins [48]. Our data

establish a distinct, mechanistic role for vinculin in allosterically regulating Ecad

conformation and adhesion at the single molecule level. Our results also demonstrate that

previous interpretations of how vinculin strengthens Ecad adhesion are incomplete. While

these studies solely focused on vinculin mediated remodeling of the junctional actomyosin

cytoskeleton [22-24], we demonstrate that vinculin also changes the adhesive properties of

individual Ecad ectodomains. Furthermore, our data suggests that vinculin mediates the

transmission of cytoskeletal contractile force to the Ecad ectodomain and drives the

conversion of X-dimers to strand-swap dimers. Consequently, when Ecads decouple from

vinculin and/or α-catenin, they are trapped in an X-dimer structure. Force induced changes

in Ecad dimer conformation has been previously suggested [7], but until now has not been

directly measured.

While previous AFM measurements with α-catenin knockdown cells suggest that

reducing α-catenin levels decrease the unbinding force of Ecad ectodomains, the

molecular mechanism by which α-catenin influences Ecad adhesion was not determined

[49,50]. Our data shows that vinculin association and myosin II dependent cytoskeletal

contractility allosterically trigger strand-swap dimerization of Ecad and suggests that α-

catenin affects Ecad adhesion by serving as the scaffold protein that binds vinculin to

Ecad-catenin complex.

It is important to point out that our experiments measure the allosteric

conformational regulation of non-junctional Ecads present on the apical surface of cells.

While apical Ecads may be organized differently from Ecads localized to cell-cell junctions

[51,52], the intracellular proteins that couple Ecad to the actin cytoskeleton are similar for

both Ecad pools. Previous studies show that both apical and junctional Ecads are under

similar levels of constitutive actomyosin tension [47], suggesting that both Ecad pools

interact with the actin cytoskeleton in similar manners. Moreover, magnetic twisting

cytometry show force-induced, vinculin-dependent stiffening of Ecad coated beads placed

on the apical cell surface, suggesting that vinculin also associates with the extra-junctional

Ecad [18]. More recently, 3D super-resolution microscopy of MDCK cells attached to an

Ecad-coated substrate, show that the intracellular organization of many Ecad-associating

proteins including p-120 catenin, β-catenin, α-catenin and vinculin, resemble native cell-

cell junctions [53]. These studies suggest that the inside-out regulation that we measure with

apical Ecads, also occurs with junctional Ecads.

Our ECIS experiments show that while barrier formation half time for both parental-

Trp and ΔVBS cells was longer compared to parental cells (Figure 4F), the maximum

impedance after barrier formation was completed for parental and ΔVBS cells were similar

to each other and slightly higher than parental-Trp cells (Figure S4). This finding is

explained by the fact that the maximum impedance is a measure of the steady state of

barrier integrity after the dynamic process of junction formation is completed and is

principally determined by formation of tight junctions. Consequently, given enough time,

the clustering of Ecads within ΔVBS junctions could orient them to promote formation of

strand-swap dimers, similar to parental cells. However, when Trp is added to the parental

cells, it blocks strand-swapping and traps Ecad in an X-dimer conformation even when

junction formation is completed and reaches steady-state. Previous studies show that tight

junctions in Ecad deficient mice have increased permeability [54]. This suggests that

inhibiting strand-swapping in Ecad may affect tight junction formation resulting in a lower

maximum impedance in parental-Trp cells.

Despite the large number of proteins expressed on the cell surface, we were able

to measure single molecule unbinding by limiting Ecad density on the AFM tip and

controlling the force with which the AFM tip contacts the cell. However, we were also aided

by a low density of Ecads expressed on the apical surface of confluent MDCK cell

monolayers. While the precise oligomeric state of Ecad on apical cell surfaces is unclear,

with super-resolution imaging suggesting that apical Ecads in A431 cells are present as

well separated monomers [51] while Ecads in A431D cells organize into small nanoclusters

[52], our low unbinding event rates nonetheless suggest that only single Ecad on the AFM

tip interact with Ecad on the cell surface even if a direct comparison between MDCK and

A431/A431D cells cannot be made potentially due to their different properties.

Interestingly, the total Ecad event rate measured with αcat-KO cells is higher than with

parental, ΔVBS or vinc-KO cells (Figure 2B). This likely occurs because the αcat-KO cells

do not form stable cell-cell junctions [25]. Consequently, Ecads that would otherwise be

corralled within cell-cell junctions are now redistributed on the apical cells surface and

subsequently interact with Ecads on the AFM tip with a higher probability. The same may

be true for other membrane proteins which contributes to an increase in non-specific event

rate in control experiments with αcat-KO cells (Figure 2B).

Although AFM is uniquely suited to study the biomechanics of membrane proteins,

as with any experiment, there are limitations. The cell surface is a complex environment

with many molecules and it is difficult to determine which cell surface protein is interacting

with Ecad on the AFM tip. For this reason, we engineered the Ecad-KO cell lines and show

that we are not observing any significant heterophilic interactions. For instance, while Ecad

can interact heterophilically with several transmembrane proteins, including Desmoglein-

2 (Dsg2) [55,56], our control measurements with Ecad-KO cells show that there is no

significant binding of Ecad with other cell-surface proteins (Figure 2B). The binding rates

measured for the interaction between an AFM tip functionalized with Ecad and the Ecad-

KO cells (which corresponds to heterophilic binding between Ecad on the AFM tip and

other proteins expressed in MDCK cells) is similar to the binding rate for a bare AFM tip

lacking Ecad and parental MDCK cells expressing all transmembrane proteins. This

demonstrates that heterophilic Ecad interactions on the apical cell surface occur at such

a low rate that we are unable to detect them. One possible reason for the low heterophilic

binding of Ecad is that other transmembrane binding partners may have low expression

on the apical surface of MDCK cells. Alternatively, the affinity of Ecad heterophilic binding

may be lower compared to Ecad-Ecad interactions. In addition, since the structures of

Ecad bound to different heterophilic binding partners are unknown, our experimental

design may also preclude some of these interactions from occurring. For instance, since

Dsg2 binds to the *cis* interface of Ecad [55], this would require tethered Ecads on the AFM

cantilever to flip their orientation and interact with Dsg2 on the cell surface, an event that

has a low probability of occurring. Finally, it is important to note that western blot analysis

shows similar level of Dsg2 expression for all cell lines compared to Ecad-KO cells (Figure

S5), suggesting that the lack of heterophilic Ecad-Dsg2 binding in Ecad-KO cells would

be true for other cell lines as well. However, we cannot completely eliminate the possibility

of some heterophilic binding in our experiments as it is possible that the knock-out cells

have different localization of Ecad and its binding partners. It is also important to note that

since our cells are grown on non-porous glass coverslips for up to 72 hours, they are, at

best, only partially polarized. Consequently, while some microvilli or primary cilia may be

present on the apical cell surface, the probability of probing Ecad on these structures is

quite low.

Previous studies also suggest that α-catenin is not the sole linker of Ecad and the

cortical actin cytoskeleton. The binding of β-catenin to vinculin is believed to serve as an

alternate, secondary bypass connection between Ecad and F-actin [57]. However, this

alternate coupling is believed to be initiated by α-catenin [58]. Our AFM data is consistent

with these findings. The fraction of cytoskeletal-linked Ecad in both parental and ΔVBS

cells, which presumably contain the primary α-catenin/F-actin and the secondary β-

catenin/vinculin linkages, are similar (Figure 4A). In contrast, in vinc-KO cells, which

presumably contain α-catenin/actin linkages but do not contain β-catenin/vinculin

linkages, we see a reduced coupling of Ecad to the actin cytoskeleton (Figure 4A). Finally,

in αcat-KO cells we see practically no coupling between Ecad and F-actin, presumably

because both the primary α-catenin/actin and the alternate β-catenin/vinculin linkages

380 cannot be formed (Figure 4A). However, our data do not eliminate the possibility that the

381 greater fraction of Ecad-actin linkages in ΔVBS cells compared to vinc-KO cells occurs

382 because the ΔVBS cells contain higher amounts of cytoplasmic α-catenin (Figure 1C).

383  It is also worth noting that measurements in αcat-KO cells — where Ecad cannot

384 link to actin cytoskeleton — mainly resulted in membrane-tethers (Figure 4A), validating

385 the use of membrane-tethers as a signature of cytoskeleton-uncoupled Ecad.

386 Furthermore, the very low occurrence of jumps and cyto-tethers in αcat-KO cells (Figure

387 4A) implies that jumps and cyto-tethers indeed correspond to the pulling of Ecad coupled

388 to the underlying cytoskeleton. Additionally, the presence of only jumps in the parental-

389 Trp, vinc-KO, and ΔVBS cells (Figure 4B), where Ecad exclusively formed weaker X-

390 dimers (Figure 5B-C, E), supports that jumps represent X-dimers linked to the actin

391 cytoskeleton. Finally, the decrease in fraction of cyto-tethers (Figure 4B) with the

392 decreased probability of strand-swap dimer formation in parental-Bleb (Figure 5F) strongly

393 suggests that cyto-tethers correspond to strand-swap dimers linked to the actin

394 cytoskeleton. While Ecad in parental cells that are coupled to the cytoskeleton via α-

395 catenin and vinculin (cyto-tethers) exclusively formed strand-swap dimers, we were

396 surprised to observe that Ecad which are decoupled from the cytoskeleton (membrane-

397 tethers) formed X-dimers and strand-swap dimers with coequal probabilities (Figure S3A,

398 B). A likely explanation for this result is that the Ecad-cytoskeletal linkage is dynamic and

399 that even transient interactions with vinculin are sufficient to drive strand-swap dimer

400 formation. Since the AFM-tip and the cell surface remain in contact for about 0.45s in our

401 experiments (Figure S6), this may allow enough time from Ecads on the parental cells to

402 convert to a strand-swap dimer conformation and detach from the cytoskeleton (thereby

403 forming membrane-tethers), before the AFM-tip is withdrawn from the cell surface.

When parental cells were treated with blebbistatin, the reduced actomyosin tension

resulted in a decrease in the fraction of cyto-tethers (Figure 4B). While these cyto-tethers

behaved like parental cyto-tethers and exclusively formed slip-bonds (Figure S3C), a

larger fraction of parental-Bleb membrane-tethers formed X-dimers compared to parental

cells (69% in parental-Bleb compared to 55% in parental cells), further demonstrating that

myosin generated contractile force is important for cadherin strand-swap dimer formation

(Figure S3D).

Given the higher lifetimes for X-dimers compared to strand-swap dimers at forces

> 50 pN (Figure 5A), it might be expected that force would convert Ecad to X-dimers

instead of strand-swap dimers. However, since contractile force applied by actomyosin on

a single Ecad is only around 1-2 pN [47] and the lifetime of the strand swap dimer is higher

than the X-dimer at forces < ~50 pN, (Figure 5A), the strand-swap dimer is the preferred

binding conformation in a physiological context. Furthermore, as shown by our simulations

(Figure 6A-C), the geometry of these conformations do not permit a transition from strand-

swap dimer to X-dimer since pulling on the strand-swap dimer moves the X-dimer interface

further apart.

Consistent with a previous study that used optical tweezers and single particle

tracking to infer the fraction of cytoskeleton-bound Ecad [36], our data shows that only ~50%

of Ecads on the apical cell surface are coupled to the actin cytoskeleton. Studies also

show that cells expressing Ecad/α-catenin fusion constructs that constitutively bind to the

actin cytoskeleton, exhibit higher cell-cell adhesion and reduced intercellular motility

compared to cells expressing wild type Ecad [36,59]. These findings are supported by our

observation that two rapidly exchangeable pools of Ecad structures (X-dimers and strand-

swap dimers) exist on the cell surface. It is likely that these different pools of Ecads are

crucial in reorganizing cell-cell junctions during collective cell migration: as the cell-cell

contacts rapidly rearrange, strand-swap dimers anchor the contact while X-dimers probe

for formation of new contacts.

Based on the results presented in this manuscript, we propose a biophysical model

for the inside-out regulation of Ecad conformation. Our data suggests that vinculin

mediated myosin II recruitment generates a contractile force which propagates to the Ecad

ectodomain and drives the conversion of X-dimers to strand-swap dimers. (Figure 6D).

Consequently, treating parental cells with blebbistatin, inhibits myosin II dependent

cytoskeletal contractility and reduces the ability of Ecad to form robust strand-swap dimers

(Figure 5F). All atom computer simulations also demonstrate that a force applied to the

membrane proximal region of X-dimers results in formation of strand-swap dimers (Figure

6B, 6C). Previous studies suggest that Ecad adhesion is altered by changing the

phosphorylation state of cytoplasmic p120-catenin [40,60,61]; however, the mechanism by

which this occurs and the exact conformational states Ecads adopt, are unknown. It is

possible that p120-catenin also indirectly modulates vinculin association with the Ecad

cytoplasmic region, which in turn regulates Ecad ectodomain conformation and adhesion.

Recent studies have suggested that regulation of Ecad adhesion plays an

important role in cancer progression. For instance, it has been shown that up-regulation

of Ecad adhesive activity, rather than its amount, reduces the number of cells

metastasized from the mammary gland to the lung [62]. Additionally, several point mutations

in Ecad ectodomains that are prevalent in hereditary diffuse gastric carcinoma selectively

interfere with the inside-out regulation of Ecad adhesion, rather than the ability of Ecad to

adhere [62]. Furthermore, studies have shown that while vinculin expression is significantly

reduced in colorectal cancer (CRC), overexpressing vinculin reduces metastasis in CRC

[63] and other cancer cell lines [64]. Consequently, understanding how cytoplasmic proteins

induce conformational changes of Ecad ectodomains could be crucial to understanding

the role of Ecad in cancer progression and metastasis.

METHODS

Generating mutant cell lines. Generation of αcat-KO cells (ΔαE-catenin MDCK cells) [25]

has been described previously. For generation of ΔVBS cells, αcat-KO cells were stably

rescued [26] with GFP-α-catenin–ΔVBS mutants [35] as described previously. Vinc-KO and

Ecad-KO cells were generated by stably expressing a PiggyBac plasmid encoding Cas9

with a cumate inducible promoter, and gRNA expression cassette for canine Ecad or

vinculin specific gRNA, which resulted in complete knockout cells (see Figure 1C). The

mutations were sequence verified by PCR amplifying the target sequences from genomic

DNA of knockout cell lines and inserted into TOPO vectors (Invitrogen) for sequencing.

Ecad:
gRNA: ACAGACCAGTAACTAACGA
KO: ACAGACCAGT----AACGA
Vinculin:
gRNA: GGA GCACCGAGTAATGTTGG
KO: GGAtGCACCGAGTgATGTTGG

Western Blotting. Cell lysates for western blotting were prepared in 1x SDS sample

buffer (10 mM Tris pH 6.8, 1% SDS, 10% glycerol, 0.005% Bromophenol Blue, 2% β-

mercaptoethanol). Primary antibodies against E-cadherin (clone 36; BD Biosciences), α-

catenin (clone 15D9; Enzo Life Sciences), vinculin (clone hVIN-1; Milipore Sigma), Dsg2

(clone 141409; R&D Systems), and α-tubulin (clone DM1A; Cell Signaling) were used in

manufacturer recommended dilutions in Phosphate-Buffered Saline with 0.1% Tween 20

(PBST). Horse Radish Peroxidase (HRP) conjugated secondary antibody (catalog no.

1706516; Bio-Rad) was used in 1:1000 dilution in PBST. Protein samples were detected

with a WesternBright Chemiluminescence Kit (catalog no. K-12045; Advansta). Images

were acquired using Image Lab software from Bio-Rad.

Purification of Ecad ectodomain. Generation of Ecad monomer plasmids containing a

C-terminal Avi tag has been described previously [65]. The plasmids were incorporated into

pcDNA3.1(+) vectors and were transiently transfected into HEK 293T cells using PEI

(Milipore Sigma) as previously described [66]. Three days post transfection, conditioned

media was collected for protein purification. Purification of Ecad were performed using

methods described previously [8,65]. Media containing his-tagged Ecads was passed

through a chromatography column containing Ni-NTA agarose beads (Qiagen). Beads

were then washed with a pH 7.5 biotinylation buffer (25mM HEPES, 5mM NaCl, and 1mM

$CaCl_2$). Ecads bound to the Ni-NTA beads were biotinylated with BirA enzyme (BirA 500

kit; Avidity) for 1hr at 30^0C. Following biotinylation, free biotins were removed using the

Ni-NTA column and biotinylated Ecads bound to Ni-NTA beads were eluted using a pH

7.5 buffer containing 200mM Imidazole, 20mM Na_2HPO_4, 500mM NaCl, and 1mM $CaCl_2$.

Epithelial barrier formation measurements. Ecad-dependent formation of epithelial

barriers was assessed by real-time impedance sensing. To this end, parental or ΔVBS

cells were trypsinized, re-suspended in calcium-free DMEM medium (Immunosource,

D9800-10) supplemented with 10% dialyzed fetal calf serum, and plated onto L-cysteine-

reduced, collagen I-coated 8WE10 electrodes (Applied Biophysics) at a density of 2×10^5

cells per well and in the presence or absence of 2 mM Trp. Impedance measurements

were performed at 37°C and 6% CO2 using a 1600R Electrical Cell Impedance Sensing

(ECIS) system (Applied Biophysics) at a frequency of 4000 Hz. $CaCl_2$ was added at a final

concentration of 2 mM to allow for adherens junction formation, and measurements were

continued until stable impedance levels were reached in all conditions. Within each

individual experiment, 4 technical replicates for each condition were averaged to assess

the time-dependent impedance for that condition. The half times of barrier formation were

calculated as the timepoint at which the impedance signal surpassed 50% of the ultimate

maximum impedance.

Ecad functionalization on AFM cantilever. Purified Ecad monomers were immobilized

on Si tip of AFM cantilevers (Hydra 2R-100N; AppNano) as described previously [8,9,55,65].

Briefly, the AFM cantilevers were cleaned by immersing in 25% H_2O_2/ 75% H_2SO_4 solution

overnight. The cantilevers were then sequentially washed with deionized water and

acetone. The cantilevers were functionalized with amine groups by immersing in 2%

(vol/vol) 3-aminopropyltriethoxysilane (Millipore Sigma) solution dissolved in acetone.

Polyethylene glycol (PEG) (MW 5000, Lysan Bio) spacers containing amine-reactive N-

hydroxysuccinimide ester group at one end were covalently attached to the cantilever (100

mg/ml in 100mM $NaHCO_3$ dissolved in 600mM K_2SO_4, for 4 hrs); 20% of the PEG spacers

presented biotin molecules at the other end. Unreacted amine groups on the silane

molecules were capped using NHS-sulfo acetate (10 mg/ml for 30 mins; Thermo Fisher).

PEG-functionalized AFM cantilevers were incubated in 1 mg/ml Bovine Serum Albumin [55]

overnight. The cantilevers were then sequentially incubated with streptavidin (0.1 mg/ml

for 30 mins) and biotinylated Ecads (200 nM for 90 mins). Finally, free biotin-binding sites

of streptavidin were blocked by using 0.02 mg/ml free biotin for 20 mins. All steps, except

the incubation with Ecad, were kept the same for cantilever functionalization in control

experiments.

Cell culture for AFM experiment. Cells were cultured in low glucose Dulbecco's modified

Eagle's medium (DMEM) (Milipore Sigma) supplemented with 10% FBS (Milipore Sigma)

and 1% antibiotics solution (penicillin + streptomycin + kanamycin) at 37^0C and 5% CO_2.

Cells were trypsinized 48-72 hrs prior to the experiment and plated on an ethanol-cleaned

glass coverslip at a low density. The confluent monolayer of cells was washed 5 times

with a pH 7 DMEM and antibiotics solution before loading onto the AFM setup with the

same medium.

AFM force measurements. Force measurements were performed using two Agilent 5500

AFMs with closed loop scanners. An AFM cantilever functionalized with purified Ecad was

brought in contact with a confluent monolayer of cells, pressed against the cell surface

with a low force (typical pressing force was about 50 pN) for 0.1s, and then retracted at

one of five constant velocities (3, 5, 7, 9, 11 µm/s). Control experiments used cantilevers

decorated with streptavidin but lacking Ecad. Cantilever spring constants were measured

using the thermal fluctuation method [67]. All the experiments were performed in pH 7 DMEM

and antibiotics solution in a custom-built environmental chamber supplied with 5% CO_2 at

room temperature. Parental-Bleb and parental-Trp experiments were performed with

100µM ($\pm$) -Blebbistatin (Milipore Sigma) and 2mM L-Tryptophan (Alfa Aesar) in the

solution respectively. A typical experiment lasted for ~10 hours yielding ~1200 force-

distance traces.

Analysis of AFM data. AFM-generated force-distance traces were analyzed using

custom MATLAB scripts. Traces containing single rupture events were classified into

membrane-tether, jump, and cyto-tether categories based on the following criteria:

• Membrane-tethers: force plateau at least 100nm long and slope less than 25

pN/µm.

• Cyto-tethers: have a difference between the peak force and the plateau force that

is greater than the noise, calculated as 2 standard deviations of 500 data points

after the force drop.

• Jumps: have a linear force-distance relationship with a slope of more than 25

pN/µm.

Rupture events having unbinding force less than the noise of each respective force

curve or a force >150 pN were not considered for analysis. Tethers were further fit to the

SLS model as described in [31] while jumps were fit to a Hookean spring model. We only

retained events that fit either the SLS or spring model with a Root Mean Squared Error

(RMSE) of less than the mean plus two standard deviations of all fits for each respective

model.

To obtain the fractions of membrane-tether, cyto-tether, and jump events for each

cell line, we first measured the binding probability of each type of event for both the Ecad

experiment and the control. The binding probabilities obtained from the control experiment

were subtracted from the corresponding event rate of the Ecad experiment.

Loading rates for the jump events used for dynamic force spectroscopy analysis

(Figure 4C-E) were calculated by multiplying the slope of the jump by the pulling velocity.

K-means clustering was used to group the loading rates as described previously [68]. Mean

unbinding force and mean loading rate for the groups were fit using a weighted non-linear

least square fit to the Bell-Evans model [37]. Raw data was bootstrapped with replacement,

grouped using k-means, and the mean values were fit to the Bell-Evans model 1000 times;

the fitting parameters and errors were determined from the mean and standard deviations

of the resulting distributions. The data used in each fit was weighted by the fraction of data

points in a particular bin relative to the total number of data points. Force-dependent Ecad-

Ecad bond mechanics were studied by analyzing the forces and lifetimes of membrane-

tethers and cyto-tethers. Bond lifetimes were determined from the force curves starting at

the beginning of the plateau region to the bond rupture. The unbinding force was

calculated as the mean of the force values spanning that same region. This data was used

to create the plots shown in Figure 5. The unbinding forces were fit to a Gaussian mixture

model (GMM) using an expectation maximization algorithm to determine the bins; the

means and standard deviations used in the GMM fit were initialized using the k-means

clustering algorithm. We determined the characteristic lifetime of each bin using maximum

likelihood estimation (MLE) on bootstrapped samples. The likelihood function, $\mathcal{L}(\theta; X) =$

$\prod_{i=1}^{N} f_i(x_i; \theta)$, was formulated using an exponential distribution f.

Where, $f(x; \lambda) = \begin{cases} \lambda e^{-\lambda x} & x \geq 0, \\ 0 & x < 0. \end{cases}$

The MLE lifetimes were fit by Chi-squared minimization to one of two models: a composite

slip-catch bond model or a catch bond model [46]. Assuming the errors in the measurements

are Gaussian distributed, fitting by minimizing Chi-squared is the same as maximizing the

log likelihood [46]. The composite slip-catch model used to fit the parental and parental-Bleb

data was the sum of a slip bond model and a catch bond model. The slip bond model [45],

$\tau(F) = \frac{1}{k_{off}(F)} = \tau(0)e^{-\frac{Fx_\beta}{k_BT}}$, was used as the first component of the composite slip-catch

model to fit the parental and parental-Bleb data (Figure 5A, F), where τ is the characteristic

bond lifetime in the absence of force and x_β is the reaction coordinate distance. The catch

bond model used in the composite model and to fit all other cell lines is given by,

$$\tau(F) = \frac{\sqrt{\pi}r_0(E_1 - 2F(d + r_0))e^{\beta(E_0+dF)}(e^{\beta Fr_0} - 1)}{4D(\beta E_0)^{\frac{3}{2}}F\left(1 + \frac{r_0}{d}\right)^2 (1 - e^{\beta(2F(d+r_0)-E_1)})}$$

where r_0 is the natural bond length, d is the reaction coordinate (same as x_β for the slip

bond model), E_0 and E_1 are components of the bond energy, $\beta = \frac{1}{k_BT}$ (k_B is the Boltzmann

constant), and D is a diffusion constant ($D = \frac{k_BT}{6\pi\eta r_0}$). The number of bins used was

established by the minimum weighted RMSE of the composite slip-catch or catch bond

model relative to the mle lifetimes when fitting the forces with the GMM using 5 to 10 bins.

(7-10 for parental and parental-bleb) for each cell line. The weights were determined by

the fraction of data points in a bin relative to the total number of data points for that cell

line. The composite slip-catch and catch bond model fitted parameters are shown in Table

S2. The Chi-squared fitting function was acquired from the Mathworks File Exchange

website (Generalized Nonlinear Non-analytic Chi-Square Fitting by Nathaniel Brahms,

May 26, 2006).

The fraction of strand-swap and X-dimers calculated from Figure 5A and 5F were

determined by integrating each component of the composite model in 5 pN bins from 0 to

150 pN to determine the slip/catch area of that bin. The number of data points in each bin

that corresponded to the slip bond model, for example, was determined by,

$$\Delta n_{slip} = \left(\frac{\Delta A_{slip}}{\Delta A_{slip} + \Delta A_{catch}}\right)\Delta n$$

where ΔA_{slip} and ΔA_{catch} are the area under slip and catch bond fit respectively for the

5pN bin, and Δn is the total number of data points in the 5pN bin. As expected, the sum of

the slip and catch bond fits—and therefore the sum of their areas—is equal to the

composite fit. The total fraction of data corresponding to the slip bond (strand-swap

dimers) is then,

621
$$Fraction_{slip} = \frac{\sum \Delta n_{slip}}{\sum \Delta n}$$

Steered Molecular Dynamics Simulations and Structural Analysis. Steered molecular

dynamics simulations were performed with GROMACS 2020.1 using the FARM high

performance computing cluster at UC Davis. The X-dimer crystal structure (PDB ID:

3LNH) was used in all simulations. Missing residues were added using PDBFixer and the

N-terminal β-strand with Trp2 was positioned near the strand-swap interface using the

sculpting tool in PyMOL. All simulations were performed with the OPLS-AA/L force field

and the TIP4P water model.

The X-dimer crystal structure was equilibrated by performing a 20 ns molecular

dynamics (MD) simulation. To setup the simulation, the X-dimer was placed in the center

of a dodecahedral box such that no atom of the X-dimer was closer than 1 nm to any

boundary. The box was solvated by adding water molecules and charge-neutralized by

adding Na$^+$ ions. Additionally, NaCl, KCl, and CaCl$_2$ were added to the box at

concentrations of 150 mM, 4 mM, and 2 mM, respectively, for a total system size of

148,965 atoms. Energy minimization, using the steepest decent algorithm, was performed

until the force on any atom in the system was less than 1000 kJ/mol/nm. The temperature,

pressure, and density of the system were stabilized by equilibrating under *isothermal-*

isochoric and *isothermal-isobaric* conditions using a modified Berendsen thermostat and

Berendsen barostat. Following equilibration, a 20 ns MD simulation was performed with 2

fs integration steps, LINCS constraints on bond length, and long-range interactions were

evaluated using the particle mesh Ewald method. The stability of the X-dimer was

monitored by calculating the root mean squared deviation (RMSD) of the X-dimer relative

to the initial structure.

Conformational switching was modeled using Steered molecular dynamics (SMD)

simulations performed on the fully equilibrated X-dimer by pulling on a group of residues

(residues 151-166, 174-186, and 208-213) — to avoid unfolding—at the C-terminal end of

one Ecad monomer in five directions (0°, ±30°, ±40°; *see Figure 6A*). The C-terminal of

the opposing Ecad was constrained to simulate a robust linkage to the actin cytoskeleton.

The X-dimer structure from the final frame of the MD simulation was placed at the center

of a 12 x 30 x 8 nm^3 box, and solvated and equilibrated as described above to an average

of 376,743 atoms. After centering the X-dimer in the box, the pulling angle was achieved

by rotating the X-dimer about the z-axis (short axis) by the corresponding pulling angle.

For each simulation, the X-dimer was pulled with a constant force (force constant = 498.2

pN/nm; 300 kJ/mol/nm^2) along the long axis of the box.

The conversion of the X-dimer to strand-swap dimer was determined by aligning

the simulated structures to the X-dimer (PDB ID: 3LNH) and strand-swap dimer (PDB ID:

2QVF) crystal structures and calculating the RMSD of the alpha carbon atoms before and

after pulling, respectively. The strand-swap dimer was formed from 2QVF by

crystallographic symmetry operations.

Acknowledgments: Research in the SS lab was supported in part by the National

Science Foundation (PHY-1607550) and National Institute of General Medical Sciences

of the National Institutes of Health (R01GM121885 and R01GM133880). Research in the

SY lab was supported by NIH R03 EB021636, NSF 1562095 (SY, REU supplement to

JC), and UC Davis Provost's Undergraduate Fellowship (to JC). JC is a recipient of the

Beckman Scholars Award. Research in the MG lab was supported by the Netherlands

Organization for Scientific Research (NWO; 016.Vidi.189.166 and NWO gravitational

program CancerGenomiCs.nl).

Data availability: All data produced by this research are made available in the Supporting

Information.

Author contributions: R.K., A.V.P. and S.S. designed research; S.S. directed the

research; R.K., A.V.P. and C.F.Y. performed AFM experiments; A.V.P. performed

computer simulations; W.J.P and M.G. performed ECIS experiments; J.S.C., M.G., and

S.Y., generated cell lines; R.K., A.V.P., and S.S. wrote the paper.

References

1   Mendonsa, A. M., Na, T.-Y. & Gumbiner, B. M. E-cadherin in contact inhibition

and cancer. *Oncogene* **37**, 4769-4780 (2018).

2   Boggon, T. J. *et al.* C-cadherin ectodomain structure and implications for cell

adhesion mechanisms. *Science* **296**, 1308-1313 (2002).

3   Haussinger, D. *et al.* Proteolytic E-cadherin activation followed by solution NMR

and X-ray crystallography. *EMBO J.* **23**, 1699-1708 (2004).

4   Parisini, E., Higgins, J. M. G., Liu, J. H., Brenner, M. B. & Wang, J. H. The crystal

structure of human E-cadherin domains 1 and 2, and comparison with other

cadherins in the context of adhesion mechanism. *J. Mol. Biol.* **373**, 401-411,

doi:10.1016/j.jmb.2007.08.011 (2007).

5   Sivasankar, S., Zhang, Y., Nelson, W. J. & Chu, S. Characterizing the initial

encounter complex in cadherin adhesion. *Structure* **17**, 1075-1081 (2009).

6   Harrison, O. J. *et al.* Two-step adhesive binding by classical cadherins. *Nat.*

Struct. Mol. Biol. **17**, 348-U121, doi:10.1038/nsmb.1784 (2010).

7   Hong, S. J., Troyanovsky, R. B. & Troyanovsky, S. M. Cadherin exits the junction

by switching its adhesive bond. *J. Cell Biol.* **192**, 1073-1083,

doi:10.1083/jcb.201006113 (2011).

8   Rakshit, S., Zhang, Y., Manibog, K., Shafraz, O. & Sivasankar, S. Ideal, catch,

and slip bonds in cadherin adhesion. *Proceedings of the National Academy of*

Sciences of the United States of America **109**, 18815-18820,

doi:10.1073/pnas.1208349109 (2012).

9   Manibog, K. *et al.* Molecular determinants of cadherin ideal bond formation:

Conformation-dependent unbinding on a multidimensional landscape.

Proceedings of the National Academy of Sciences of the United States of

America **13**, E5711-E5720 doi:10.1073/pnas.1604012113 (2016).

10   Manibog, K., Li, H., Rakshit, S. & Sivasankar, S. Resolving the molecular
706   mechanism of cadherin catch bond formation. *Nat. Commun.* **5**, 3941,
707   doi:10.1038/ncomms4941 (2014).

708   11   Li, Y. *et al.* Mechanism of E-cadherin dimerization probed by NMR relaxation
709   dispersion. *Proceedings of the National Academy of Sciences of the United*
710   *States of America* **110**, 16462-16467, doi:10.1073/pnas.1314303110 (2013).

711   12   Ratheesh, A. & Yap, A. S. A bigger picture: classical cadherins and the dynamic
712   actin cytoskeleton. *Nature Reviews Molecular Cell Biology* **13**, 673-679 (2012).

713   13   Yonemura, S., Wada, Y., Watanabe, T., Nagafuchi, A. & Shibata, M. alpha-
714   Catenin as a tension transducer that induces adherens junction development.
715   *Nat. Cell Biol.* **12**, 533-U535, doi:10.1038/ncb2055 (2010).

716   14   Kim, T.-J. *et al.* Dynamic visualization of α-catenin reveals rapid, reversible
717   conformation switching between tension states. *Curr. Biol.* **25**, 218-224 (2015).

718   15   Buckley, C. D. *et al.* The minimal cadherin-catenin complex binds to actin
719   filaments under force. *Science* **346**, 1254211, doi:10.1126/science.1254211
720   (2014).

721   16   Choi, H.-J. *et al.* αE-catenin is an autoinhibited molecule that coactivates vinculin.
722   *Proc. Natl. Acad. Sci. U. S. A.* **109**, 8576-8581, doi:10.1073/pnas.1203906109
723   (2012).

724   17   Yao, M. *et al.* Force-dependent conformational switch of α-catenin controls
725   vinculin binding. *Nat. Commun.* **5** (2014).

18   le Duc, Q. *et al.* Vinculin potentiates E-cadherin mechanosensing and is recruited
to actin-anchored sites within adherens junctions in a myosin II-dependent
manner. *J Cell Biol* **189**, 1107-1115, doi:10.1083/jcb.201001149 (2010).

19   Barry, A. K. *et al.* α-catenin cytomechanics–role in cadherin-dependent adhesion
and mechanotransduction. *J. Cell Sci.* **127**, 1779-1791 (2014).

20   Seddiki, R. *et al.* Force-dependent binding of vinculin to α-catenin regulates cell–
cell contact stability and collective cell behavior. *Mol. Biol Cell* **29**, 380-388
(2018).

21   Thomas, W. A. *et al.* α-Catenin and vinculin cooperate to promote high E-
cadherin-based adhesion strength. *J. Biol. Chem.* **288**, 4957-4969 (2013).

22   Huveneers, S. & de Rooij, J. Mechanosensitive systems at the cadherin-F-actin
interface. *J. Cell Sci.* **126**, 403-413, doi:10.1242/jcs.109447 (2013).

23   Leckband, D. & de Rooij, J. Cadherin adhesion and mechanotransduction. *Annu.*
Rev. Cell Dev. Biol. **30**, 291-315 (2014).

24   Mège, R. M. & Ishiyama, N. Integration of cadherin adhesion and cytoskeleton at
adherens junctions. *Cold Spring Harbor Perspect. Biol.* **9**, a028738 (2017).

25   Ortega, F. E. *et al.* Adhesion to the host cell surface is sufficient to mediate
Listeria monocytogenes entry into epithelial cells. *Mol. Biol Cell*, E16-12-0851
(2017).

26   Twiss, F. *et al.* Vinculin-dependent Cadherin mechanosensing regulates efficient
epithelial barrier formation. *Biology open* **1**, 1128-1140 (2012).

27   Hochmuth, R. & Evans, E. Extensional flow of erythrocyte membrane from cell

body to elastic tether. I. Analysis. *Biophys. J.* **39**, 71-81 (1982).

28   Hochmuth, R., Wiles, H., Evans, E. & McCown, J. Extensional flow of erythrocyte

membrane from cell body to elastic tether. II. Experiment. *Biophys. J.* **39**, 83-89

(1982).

29   Evans, E., Heinrich, V., Leung, A. & Kinoshita, K. Nano-to microscale dynamics

of P-selectin detachment from leukocyte interfaces. I. Membrane separation from

the cytoskeleton. *Biophys. J.* **88**, 2288-2298 (2005).

30   Sariisik, E. *et al.* Decoding cytoskeleton-anchored and non-anchored receptors

from single-cell adhesion force data. *Biophys. J.* **109**, 1330-1333 (2015).

31   Schmitz, J., Benoit, M. & Gottschalk, K.-E. The viscoelasticity of membrane

tethers and its importance for cell adhesion. *Biophys J* **95**, 1448-1459 (2008).

32   Heinrich, V., Leung, A. & Evans, E. Nano-to microscale dynamics of P-selectin

detachment from leukocyte interfaces. II. Tether flow terminated by P-selectin

dissociation from PSGL-1. *Biophys. J.* **88**, 2299-2308 (2005).

33   Rimm, D. L., Koslov, E. R., Kebriaei, P., Cianci, C. D. & Morrow, J. S. Alpha 1

(E)-catenin is an actin-binding and-bundling protein mediating the attachment of

F-actin to the membrane adhesion complex. *Proc. Natl. Acad. Sci. U. S. A.* **92**,

8813-8817 (1995).

34   Desai, R. *et al.* Monomeric [alpha]-catenin links cadherin to the actin

cytoskeleton. *Nat. Cell Biol.* **15**, 261-273 (2013).

35   Huveneers, S. *et al.* Vinculin associates with endothelial VE-cadherin junctions to

769   control force-dependent remodeling. *J. Cell Biol.* **196**, 641-652 (2012).

770   36   Sako, Y., Nagafuchi, A., Tsukita, S., Takeichi, M. & Kusumi, A. Cytoplasmic

771   regulation of the movement of E-cadherin on the free cell surface as studied by

772   optical tweezers and single particle tracking: corralling and tethering by the

773   membrane skeleton. *J. Cell Biol.* **140**, 1227-1240 (1998).

774   37   Evans, E. & Ritchie, K. Dynamic strength of molecular adhesion bonds. *Biophys*

775   *J* **72**, 1541-1555 (1997).

776   38   Panorchan, P. *et al.* Single-molecule analysis of cadherin-mediated cell-cell

777   adhesion. *J. Cell Sci.* **119**, 66-74, doi:10.1242/jcs.02719 (2006).

778   39   Shi, Q. M., Maruthamuthu, V., Li, F. & Leckband, D. Allosteric Cross Talk

779   between Cadherin Extracellular Domains. *Biophys. J.* **99**, 95-104,

780   doi:10.1016/j.bpj.2010.03.062 (2010).

781   40   Shashikanth, N. *et al.* Allosteric regulation of E-cadherin adhesion. *J. Biol. Chem.*

782   **290**, 21749-21761 (2015).

783   41   Shashikanth, N., Kisting, M. A. & Leckband, D. E. Kinetic Measurements Reveal

784   Enhanced Protein-Protein Interactions at Intercellular Junctions. *Scientific*

785   *Reports* **6**, 11, doi:10.1038/srep23623 (2016).

786   42   Bunse, S. *et al.* Role of N-Cadherin cis and trans Interfaces in the Dynamics of

787   Adherens Junctions in Living Cells. *PLOS ONE* **8**, e81517,

788   doi:10.1371/journal.pone.0081517 (2013).

789   43   Krieg, M., Helenius, J., Heisenberg, C. P. & Muller, D. J. A bond for a lifetime:

790   employing membrane nanotubes from living cells to determine receptor–ligand

791   kinetics. *Angew. Chem. Int. Ed.* **47**, 9775-9777 (2008).

792   44   Hosu, B. G., Sun, M., Marga, F., Grandbois, M. & Forgacs, G. Eukaryotic

793   membrane tethers revisited using magnetic tweezers. *Phys. Biol.* **4**, 67 (2007).

794   45   Bell, G. I. Models for specific adhesion of cells to cells. *Science* **200**, 618-627

795   (1978).

796   46   Chakrabarti, S., Hinczewski, M. & Thirumalai, D. Plasticity of hydrogen bond

797   networks regulates mechanochemistry of cell adhesion complexes. *Proc. Natl.*

798   *Acad. Sci. U. S. A.* **111**, 9048-9053, doi:10.1073/pnas.1405384111 (2014).

799   47   Borghi, N. *et al.* E-cadherin is under constitutive actomyosin-generated tension

800   that is increased at cell-cell contacts upon externally applied stretch. *Proceedings*

801   *of the National Academy of Sciences of the United States of America* **109**,

802   12568-12573, doi:10.1073/pnas.1204390109 (2012).

803   48   Kim, M., Carman, C. V. & Springer, T. A. Bidirectional transmembrane signaling

804   by cytoplasmic domain separation in integrins. *Science* **301**, 1720-1725 (2003).

805   49   Bajpai, S. *et al.* alpha-Catenin mediates initial E-cadherin-dependent cell-cell

806   recognition and subsequent bond strengthening. *Proc. Natl. Acad. Sci. U. S. A.*

807   **105**, 18331-18336, doi:10.1073/pnas.0806783105 (2008).

808   50   Bajpai, S., Feng, Y., Krishnamurthy, R., Longmore, G. D. & Wirtz, D. Loss of α-

809   catenin decreases the strength of single E-cadherin bonds between human

810   cancer cells. *J. Biol. Chem.* **284**, 18252-18259 (2009).

51   Indra, I. *et al.* Spatial and temporal organization of cadherin in punctate adherens

junctions. *Proceedings of the National Academy of Sciences of the United States*

of America **115**, E4406-E4415, doi:10.1073/pnas.1720826115 (2018).

52   Wu, Y., Kanchanawong, P. & Zaidel-Bar, R. Actin-delimited adhesion-

independent clustering of E-cadherin forms the nanoscale building blocks of

adherens junctions. *Dev Cell* **32**, 139-154, doi:10.1016/j.devcel.2014.12.003

(2015).

53   Bertocchi, C. *et al.* Nanoscale architecture of cadherin-based cell adhesions. *Nat*

Cell Biol **19**, 28-37, doi:10.1038/ncb3456 (2017).

54   Tunggal, J. A. *et al.* E-cadherin is essential for in vivo epidermal barrier function

by regulating tight junctions. *EMBO J* **24**, 1146-1156,

doi:10.1038/sj.emboj.7600605 (2005).

55   Shafraz, O. *et al.* E-cadherin binds to desmoglein to facilitate desmosome

assembly. *eLife* **7**, e37629 (2018).

56   Shafraz, O., Xie, B., Yamada, S. & Sivasankar, S. Mapping transmembrane

binding partners for E-cadherin ectodomains. *Proceedings of the National*

Academy of Sciences **117**, 31157, doi:10.1073/pnas.2010209117 (2020).

57   Peng, X., Cuff, L. E., Lawton, C. D. & DeMali, K. A. Vinculin regulates cell-

surface E-cadherin expression by binding to β-catenin. *J. Cell Sci.*, jcs. 056432

(2010).

831 58 Bertocchi, C., Ravasio, A., Ong, H. T., Toyama, Y. & Kanchanawong, P.

832   Mechanical Roles of Vinculin/β-catenin interaction in Adherens Junction. *BioRxiv*,

833   doi:10.1101/770735 (2019).

834 59 Nagafuchi, A., Ishihara, S. & Tsukita, S. The roles of catenins in the cadherin-

835   mediated cell adhesion: functional analysis of E-cadherin-alpha catenin fusion

836   molecules. *J. Cell Biol.* **127**, 235-245 (1994).

837 60 Petrova, Y. I., Spano, M. M. & Gumbiner, B. M. Conformational epitopes at

838   cadherin calcium-binding sites and p120-catenin phosphorylation regulate cell

839   adhesion. *Mol Biol Cell* **23**, 2092-2108, doi:10.1091/mbc.E11-12-1060 (2012).

840 61 Maiden, S. L., Petrova, Y. I. & Gumbiner, B. M. Microtubules inhibit E-cadherin

841   adhesive activity by maintaining phosphorylated p120-catenin in a colon

842   carcinoma cell model. *PLoS One* **11**, e0148574 (2016).

843 62 Petrova, Y. I., Schecterson, L. & Gumbiner, B. M. Roles for E-cadherin cell

844   surface regulation in cancer. *Mol. Biol Cell* **27**, 3233-3244 (2016).

845 63 Li, T. *et al.* Loss of vinculin and membrane-bound β-catenin promotes metastasis

846   and predicts poor prognosis in colorectal cancer. *Molecular Cancer* **13**, 263,

847   doi:10.1186/1476-4598-13-263 (2014).

848 64 Rodriguez Fernandez, J. *et al.* Suppression of tumorigenicity in transformed cells

849   after transfection with vinculin cDNA. *J. Cell Biol.* **119**, 427-438 (1992).

850 65 Zhang, Y., Sivasankar, S., Nelson, W. J. & Chu, S. Resolving cadherin

851   interactions and binding cooperativity at the single molecule level. *Proc. Natl.*

852   *Acad. Sci. U. S. A.* **106**, 109-114 (2009).

66   Longo, P. A., Kavran, J. M., Kim, M.-S. & Leahy, D. J. in *Methods Enzymol.* Vol.

529   227-240 (Elsevier, 2013).

67   Hutter, J. L. & Bechhoefer, J. Calibration of atomic-force microscope tips. *Rev.*

Sci. Instrum. **64**, 1868-1873 (1993).

68   Yen, C.-F. & Sivasankar, S. Improving estimation of kinetic parameters in

dynamic force spectroscopy using cluster analysis. *J. Chem. Phys.* **148**, 123301

(2018).

Figures

Figure 1. Overview of experiment. (A) The extracellular region of Ecad from opposing

cells mediate adhesion. The cytoplasmic region of Ecad associates either directly or

indirectly with p120 catenin, β-catenin, α-catenin, vinculin, and F-actin. **(B)** Strand-swap

dimers form slip bonds (blue) while X-dimers form catch bonds (red). Ecads interconvert

between these two dimer conformations. Structures were generated from the crystal

structure of mouse Ecad (PDB: 3Q2V); the X-dimer was formed by alignment to an X-

dimer crystal structure (PDB: 3LNH). **(C)** Graphics showing the cell lines used in

experiments and western blot analysis of corresponding cell lysates.

Figure 2. Measuring specific and non-specific interaction using AFM. (A) Biotinylated

Ecad monomers were immobilized on AFM cantilevers functionalized with Polyethylene

glycol (PEG) and Streptavidin as described in the methods section. Interactions between

Ecads on the cantilever and Ecads expressed on the apical surface of MDCK cells were

measured. **(B)** Binding probabilities were measured using AFM cantilevers lacking Ecad (control experiment, light gray) or AFM cantilevers decorated with Ecad (Ecad, dark blue). Total number of Ecad/control measurements performed on each cell line were 6793/3361 measurements on parental cells, 8593/3719 measurements on parental-Bleb, 7268/4858 measurements on parental-Trp, 4291/3666 measurements on vinc-KO cells, 4518/3381 measurements on αcat-KO cells, 6124/6070 measurements on ΔVBS cells, and 3672/2449 measurements on Ecad-KO cells. Error bars are bootstrapped standard deviations.

Figure 3. Force 'signatures' used to identify cytoskeletal coupling and strength of *trans* adhesion. (A) Typical force curve showing a membrane-tether which extends at a constant force. **(B)** Membrane-tethers are measured when an Ecad, which is not linked to the actin cytoskeleton is pulled. **(C)** Jump events exhibit a linear increase in force and are measured upon pulling **(D)** an Ecad *trans* dimer that is weaker than its cytoskeleton coupling. **(E)** Typical force curve showing a cyto-tether with an initial increase in force that viscoelastically relaxes as a membrane-tether is formed. **(F)** Cyto-tethers are measured when a robust Ecad *trans* dimer that is linked to the actin cytoskeleton is pulled. Raw data (gray) was acquired at 25 kHz and smoothed (blue) to 111 Hz. Green curve is the Standard Linear Solid model fit to membrane-tethers and cyto-tethers. Black bars indicate bond lifetimes determined from membrane-tethers and cyto-tethers.

Figure 4. Role of α-catenin and vinculin in Ecad-cytoskeleton linkage and adhesion strength. (A) Fraction of uncoupled (membrane-tether) and coupled (cyto-tether + jump)

events. Fractions were calculated after subtracting the binding probability of the control experiment for each cell line. **(B)** Fractions of strong (cyto-tether) and weak (jump) Ecad *trans* dimer events. Error bars are the propagated errors obtained from bootstrapped standard deviations. Dynamic force spectroscopy of jump events in **(C)** vinc-KO + ΔVBS, **(D)** parental-Trp, and **(E)** parental-Bleb cells. Loading rates and peak forces in jump events were bootstrapped 1000 times then clustered using k-means and fit to the Bell-Evans model (green line) using weighted nonlinear least-square fitting. Data points represent single unbinding events, and colors correspond to data points from the same k-means cluster. The data from ΔVBS (filled circles) and vinc-KO (circles) cells in (C) were combined because the number of data points were insufficient for a statistically reliable calculation. Intrinsic off rate (k_{off}) and distance to the transition energy barrier (x_β) were obtained from the mean of the fitted parameters acquired by fitting the k-means clusters that were calculated from the bootstrapped samples. Errors in the fitted parameters correspond to standard deviations of the bootstrapped means. **(F)** Trans-epithelial resistance measurement of parental cells in the absence (Parental; n = 9) and presence (Parental-Trp; n = 8) of 2 mM tryptophan, or ΔVBS cells (ΔVBS; n = 5), following addition of 2 mM $CaCl_2$ to induce Ecad dependent formation of the epithelial barrier. Barrier formation half times (T50) were calculated as the time-point at which the impedance signal surpassed 50% of the ultimate maximum impedance. * p = 0.0025, ** p = 0.0011; Paired t-test.

Figure 5. Measuring conformation of Ecad *trans* dimers using an AFM. Force *vs* lifetime profile for **(A)** parental cells show that 69% of Ecad form slip bonds (strand-swap dimer structure shown in blue) while the remaining 31% form catch bonds (X-dimer structure shown in red). **(B)** Parental-Trp, **(C)** vinc-KO, **(D)** αcat-KO, and **(E)** ΔVBS cells

form catch bonds. **(F)** Inhibition of Myosin II using Blebbistatin reduces the fraction of slip

bond forming Ecad. Parental-Bleb shows that 43% of Ecad form slip bonds while the

remaining 57% form catch bonds. Solid lines represent fits to the corresponding models.

Data was binned using a Gaussian mixture model and the lifetimes were determined using

maximum likelihood estimation. The colored data points correspond to the data points in

each respective bin. The data was fit by χ^2 minimization and errors were obtained by

bootstrapping. The y-axes were limited to 1s for clarity.

Figure 6. Cytoskeletal tension converts X-dimers to strand-swap dimers. (A) X-dimer

formed by Ecad on opposing cells (left). X-dimer crystal structure (PDB ID: 3LNH) used in

SMD simulations (right). The pulling directions are indicated by colored arrows. The full X-

dimer (left) was simulated by aligning 3Q2V to 3LNH. **(B)** X-dimer conversion to strand-

swap dimer. Simulated structures remain in an X-dimer conformation after stabilizing MD

simulations (left). X-dimers become strand-swap dimers after pulling on the C-terminal of

one Ecad at a range of angles (right). The transparent blue structure is the strand-swap

crystal structure (PDB ID: 2QVF). The transparent red structure is the X-dimer crystal

structure (PDB ID: 3LNH). The remaining, opaque structures are the simulated structures.

(C) Root mean squared deviations (RMSD) of the pulled structures relative to the strand-

swap dimer crystal structure as a function of the time it took to converge to the strand-

swap crystal structure (minimum RMSD). RMSDs were determined by least-squares fitting

to the strand-swap dimer crystal structure and calculating the distance between alpha

carbons of the two structures. **(D)** Model for vinculin-dependent, force-induced conversion

of X-dimers to strand-swap dimers. In the absence of vinculin, force applied is insufficient

to convert the X-dimer to a strand-swap dimer (left). Cytoskeletal tension is transmitted in

a vinculin dependent manner to the Ecad extracellular bond to promote conversion to a

strand-swap dimer (right).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6