Network Analysis Reveals Different Cellulose Degradation Strategies across *Trichoderma harzianum* Strains Associated with XYR1 and CRE1

Rafaela Rossi Rosolen[1,2], Alexandre Hild Aono[1,2], Déborah Aires Almeida[1,2], Jaire Alves
Ferreira Filho[1,2], Maria Augusta Crivelente Horta[1], Anete Pereira de Souza[1,3*]

[1]Center for Molecular Biology and Genetic Engineering (CBMEG), University of Campinas
(UNICAMP), Cidade Universitária Zeferino Vaz, Campinas, SP, Brazil

[2]Graduate Program in Genetics and Molecular Biology, Institute of Biology, UNICAMP, Campinas,
SP, Brazil

[3]Department of Plant Biology, Institute of Biology, UNICAMP, Cidade Universitária Zeferino Vaz,
Rua Monteiro Lobato, Campinas, SP, Brazil

* Correspondence:
Anete Pereira de Souza
anete@unicamp.br

Number of words: 9022

The number of figures: 7

The number of tables: 1

Keywords: *Trichoderma harzianum*, *Trichoderma atroviride*, cellulose, transcription factors,
coexpression networks

Abstract

Trichoderma harzianum, whose gene expression is tightly controlled by the transcription factors
(TFs) XYR1 and CRE1, is a potential candidate for hydrolytic enzyme production. Here, we
performed a network analysis of *T. harzianum* IOC-3844 and *T. harzianum* CBMAI-0179 to explore
how the regulation of these TFs varies between these strains. In addition, we explored the
evolutionary relationships of XYR1 and CRE1 protein sequences among *Trichoderma* spp. The
results of the *T. harzianum* strains were compared with those of *Trichoderma atroviride* CBMAI-
0020, a mycoparasitic species. Although transcripts encoding carbohydrate-active enzymes
(CAZymes), TFs, transporters, and proteins with unknown functions were coexpressed with *cre1* or
xyr1, other proteins indirectly related to cellulose degradation were identified. The enriched GO
terms describing the transcripts of these groups differed across all strains, and several metabolic
pathways with high similarity between both regulators but strain-specific differences were identified.
In addition, the CRE1 and XYR1 subnetworks presented different topology profiles in each strain,
likely indicating differences in the influences of these regulators according to the fungi. The hubs of
the *cre1* and *xyr1* groups included transcripts not yet characterized or described as being related to
cellulose degradation. The first-neighbor analyses confirmed the results of the profile of the
coexpressed transcripts in *cre1* and *xyr1*. The analyses of the shortest paths revealed that CAZymes
upregulated under cellulose degradation conditions are most closely related to both regulators, and

new targets between such signaling pathways were discovered. Although the evaluated *T. harzianum*
strains are phylogenetically close and their amino acid sequences related to XYR1 and CRE1 are
very similar, the set of transcripts related to *xyr1* and *cre1* differed, suggesting that each *T.*
harzianum strain used a specific regulation strategy for cellulose degradation. More interestingly, our
findings may suggest that XYR1 and CRE1 indirectly regulate genes encoding proteins related to
cellulose degradation in the evaluated *T. harzianum* strains. An improved understanding of the basic
biology of fungi during the cellulose degradation process can contribute to the use of their enzymes
in several biotechnological applications and pave the way for further studies on the differences across
strains of the same species.

1   Introduction

Lignocellulosic biomass is a complex recalcitrant structure that requires a consortium of
carbohydrate-active enzymes (CAZymes) for its complete depolymerization. Due to their unique
ability to secrete these proteins efficiently, filamentous fungi, such as *Trichoderma* spp. and
Aspergillus spp., are widely explored for the industrial production of CAZymes (Bischof, et al. 2016;
de Assis, et al. 2015). In the genus *Trichoderma*, *Trichoderma reesei* is the primary fungal industrial
source of cellulases and hemicellulases (Martinez, et al. 2008), while *Trichoderma harzianum* and
Trichoderma atroviride have been widely explored by examining their biocontrol capacity against
plant pathogenic fungi (Medeiros, et al. 2017; Saravanakumar, et al. 2017). However, hydrolytic
enzymes from *T. harzianum* strains have demonstrated great potential in the conversion of
lignocellulosic biomass into fermentable sugars (Delabona, et al. 2020; Zhang, et al. 2020).

The production of CAZymes in filamentous fungi is controlled at the transcriptional level by several
positive and negative transcription factors (TFs) (Benocci, et al. 2017). In *T. reesei*, the Zn_2Cys_6-type
TF xylanase regulator 1 (XYR1) is described as the most important activator of cellulase and
xylanase gene expression (Stricker, et al. 2006). During growth on an induction carbon source,
XYR1 was shown to be synthesized *de novo* and degraded at the end of induction (Lichius, et al.
2014). Although XYR1 orthologs are present in almost all filamentous ascomycete fungi, the
molecular mechanisms triggered by this regulator depend on the species (Klaubauf, et al. 2014). In *T.*
atroviride, the induction of genes that encode cell wall-degrading enzymes considered relevant for
mycoparasitism, such as *axe1* and *swo1*, is influenced by XYR1 (Reithner, et al. 2014). In *T.*
harzianum, the overexpression of *xyr1* increases the levels of the reducing sugars released in a
shorter time during saccharification (Delabona, et al. 2017). Overall, in the genus *Trichoderma*,
XYR1 evolved by vertical gene transfer (Druzhinina, et al. 2018).

In the presence of easily metabolizable carbon sources, such as glucose, the expression of *xyr1* and
genes encoding lignocellulose-degrading enzymes is repressed by carbon catabolite repression
(CCR), which is regulated by the C_2H_2-type TF carbon catabolite repressor 1 (CRE1) (Alazi and Ram
2018; Mach-Aigner, et al. 2008; Strauss, et al. 1995). Upon glucose depletion, the concentration of
CRE1 in the nucleus rapidly decreases, and CRE1 is recycled into the cytoplasm (Lichius, et al.
2014). Furthermore, the phosphorylation of CRE1 plays an essential role in signal transduction to
achieve CCR (Han, et al. 2020; Horta, et al. 2019). CRE1 is the only conserved TF throughout the
fungal kingdom, suggesting a conserved mechanism for CCR in fungi (Adnan, et al. 2017).
Interestingly, the effect of *cre1* deletion on the gene expression profile of its targets varies among
species. In *T. reesei* RUT-C30, the deletion of a full *cre1* can lead to pleiotropic effects and strong
growth impairment (Mello-de-Sousa, et al. 2014; Portnoy, et al. 2011), whereas in *T. harzianum*
P49P11, it can increase the expression levels of CAZyme genes and *xyr1* (Delabona, et al. 2021).

Although *T. harzianum* strains have shown high cellulolytic activity (Horta, et al. 2018; Li, et al.
2020b), most studies investigating this species have focused on biological control (Maruyama, et al.
2020; Yan and Khan 2021). Thus, the regulatory mechanisms underlying hydrolytic enzyme
production by these fungi are still poorly explored at the transcriptional level (Delabona, et al. 2017,
2020, 2021). In addition, given the high degree of genetic variation observed within the genus
Trichoderma (Kubicek, et al. 2019) along with the complex speciation observed in the *T. harzianum*
species (Druzhinina, et al. 2010), it is necessary to explore how the main regulators XYR1 and CRE1
behave across *T. harzianum* strains.

Previously, the transcriptomes of *T. harzianum* IOC-3844 (Th3844), *T. harzianum* CBMAI-0179
(Th0179), and *T. atroviride* CBMAI-0020 (Ta0020) were investigated under cellulose degradation
conditions (Almeida, et al. 2021). Different types of enzymatic profiles were reported, and both *T.*
harzianum strains had higher cellulase activity than *T. atroviride*. Using this dataset, we aimed to
investigate how the regulation of the TFs CRE1 and XYR1 varies among *T. harzianum* strains. Based
on the assumption that coexpressed genes tend to share similar expression patterns and that they
could be coregulated by the same elements, we modeled a network of Th3844, Th0179, and Ta0020
using a weighted correlation network analysis (WGCNA) (Langfelder and Horvath 2008). The last
strain, which is distantly related to *T. reesei* (Druzhinina, et al. 2006) and represents a well-defined
phylogenetic species (Dodd, et al. 2003), was used to assess the differences across *Trichoderma*
species. In addition, phylogenetic analyses of XYR1 and CRE1 protein sequences were performed to
clarify the evolutionary relationships of these regulators among the evaluated strains.

In this study, we identified and compared modules, hub genes, and metabolic pathways associated
with CRE1 and XYR1 under cellulose degradation conditions. To deeply investigate their regulatory
activities, we also performed first neighbor and shortest-path network analyses. Although the
evaluated *T. harzianum* strains are phylogenetically close, by comparing their coexpressed transcript
profiles, functional diversity was observed. This difference was accentuated when associating the
results of Th0179 and Th3844 with those of Ta0020. Thus, we observed a specific transcriptional
pattern related to CRE1 and XYR1 in each strain. Our study could contribute to improving our
understanding of the regulation of cellulose degradation in *T. harzianum* and paves the way for
further studies evaluating differences across strains within the same species. Addressing these
questions by investigating the genetic regulatory mechanisms involved in cellulose degradation is
important for enhancing both our basic understanding and biotechnological industrial applications of
fungal abilities.

2   Materials and methods

2.1   Fungal strains, culture conditions and transcription profiling

The species were obtained from the Brazilian Collection of Environment and Industry
Microorganisms (CBMAI) located in the Chemical, Biological, and Agricultural Pluridisciplinary
Research Center (CPQBA) of the University of Campinas (UNICAMP), Brazil. The identity of the
Trichoderma isolates was authenticated by CBMAI based on phylogenetic studies of their internal
transcribed spacer (ITS) region and translational elongation factor 1 (*tef1*) marker gene. The *T.*
harzianum CBMAI-0179 (Th0179), *T. harzianum* IOC-3844 (Th3844), and *T. atroviride* CBMAI-
0020 (Ta0020) strains were cultivated in simple carbon sources, including cellulose and glucose, to
induce a clear genetic response. Details on the culture conditions and transcription profiling of the
described experiments are provided in Supplementary Material 1, while the profile of enzymatic

activities (cellulase, beta-glucosidase, and xylanase), the total protein content, and the constitution of
the secreted proteome has been published previously (Almeida, et al. 2020; Horta, et al. 2018).

2.2   Phylogenetic analyses

The ITS nucleotide sequences of *Trichoderma* spp. were retrieved from the NCBI database
(https://www.ncbi.nlm.nih.gov/). Additionally, ITS nucleotide sequences amplified from the genomic
DNA of Th3844, Th0179, and Ta0020 via PCR were kindly provided by the CBMAI and included in
the phylogenetic analysis. The ITS region has been found to be among the markers with the highest
probability of correctly identifying a very broad group of fungi (Schoch, et al. 2012). *T. harzianum*
T6776 was used as a reference genome to retrieve the CRE1 and XYR1 protein sequences belonging
to *Trichoderma* spp. Additionally, the CRE1 and XYR1 sequences were obtained from the *T.*
harzianum T6776 (Baroncelli, et al. 2015) reference genome of Th3844 and Th0179 and the *T.*
atroviride IMI206040 (Kubicek, et al. 2011) genome of Ta0020 and used as references for the TF
consensus sequences for RNA-Seq read mapping. These sequences were included in the phylogenetic
analyses.

The multiple sequence alignment was performed using ClustalW (Thompson, et al. 1994), and a
phylogenetic tree was created using Molecular Evolutionary Genetics Analysis (MEGA) software
v7.0 (Kumar, et al. 2016). The maximum likelihood (ML) (Jones, et al. 1992) method of inference
was used based on (I) a Kimura two-parameter (K2P) model (Kimura 1980), (II) a Dayhoff model
with frequencies (F^+), and (III) a Jones-Taylor-Thornton (JTT) model for ITS, CRE1, and XYR1,
respectively. We used 1,000 bootstrap replicates (Felsenstein 1985) in each analysis. The trees were
visualized and edited using Interactive Tree of Life (iTOL) v6 (https://itol.embl.de/).

2.3   Weighted gene coexpression network analysis

The gene coexpression networks of Th3844, Th0179, and Ta0020 were modeled using transcripts per
million (TPM) value data, which were validated through qPCR and described by Almeida et al.
(2021), of three biological replicates with the R (R Core Team 2018) WGCNA package. Transcripts
showing null values for most replicates under different experimental conditions were excluded. The
network was assembled by calculating the Pearson's correlation coefficient of each pair of genes. A
soft power β was chosen for each network using *pickSoftThreshold* to fit the signed network to a
scale-free topology. Then, an adjacency matrix in which the nodes correspond to transcripts and the
edges correspond to the strength of their connection was obtained. To obtain a dissimilarity matrix,
we built a topological overlap matrix (TOM) as implemented in the package.

To identify groups of transcripts densely connected in the gene coexpression network, we applied
simple hierarchical clustering to the dissimilarity matrix. From the acquired dendrogram, the
dynamicTreeCut package (Langfelder, et al. 2008) was used to obtain the ideal number of groups and
the respective categorization of the transcripts. According to the functional annotation performed by
Almeida, et al. (2021), groups containing the desired TFs XYR1 and CRE1 were identified and
named the *xyr1* and *cre1* groups, respectively.

2.4   Functional annotation of transcripts in the xyr1 and cre1 groups

We functionally annotated the transcripts in the *xyr1* and *cre1* groups. By conducting a Fisher's exact
test to extract the overrepresented terms (p-value < 0.05), all identified Gene Ontology (GO)
(Ashburner, et al. 2000) categories were used to identify enriched GO terms with the topGO package
in R (Alexa and Rahnenfuhrer 2021). To visualize the possible correlated enriched categories in the

Degradation

dataset caused by cellulose degradation, we created a treemap using the REVIGO tool (Supek, et al.
2011). Then, the metabolic pathways related to the Kyoto Encyclopedia of Genes and Genomes
(KEGG) (Kanehisa and Goto 2000) Orthology (KO) identified in *T. reesei* were selected due the high
number of annotations correlated with this species in the KEGG database. To identify the pathways
related to the enzymes identified as belonging to different groups, we used the Python 3
programming language (Sanner 1999) along with the BioPython library (Cock, et al. 2009). The
automatic annotation was manually revised using the UniProt databases (UniProt Consortium 2019).

2.5   Assessing the groups' network topologies

To provide a visual network characterization, we modeled additional gene coexpression networks of
all strains using the highest reciprocal rank (HRR) approach (Mutwil, et al. 2010). Using an R
Pearson correlation coefficient threshold of 0.8, we assessed the 30 strongest edges, which were
coded as ▢ (top 10), 1/15 (top 20), and 1/30 (top 30). To visualize the behavior of the groups
modeled by WGCNA in the HRR networks, we used Cytoscape software v3.7.0 (Shannon, et al.
2003). Using the HRR methodology, we were able to infer the possible biological correlations
depending on the network topology. First, by performing a topological analysis based on the degree
distribution, we identified and compared the hub nodes in the *cre1* and *xyr1* groups among all
evaluated strains. Second, to identify the more indirect associations of both TFs studied, we
considered HRR global networks to evaluate the first neighbors of the *xyr1* and *cre1* transcripts.
Finally, because XYR1 and CRE1 are described as regulatory proteins in CAZyme gene expression,
we were also interested in evaluating the minimum pathway between these regulatory proteins and
these hydrolytic enzymes. Based on the classification proposed by Almeida, et al. (2021), we selected
CAZymes with higher expression levels under cellulose growth conditions (upregulated transcripts)
that were present in both Th0179 and Th3844. To obtain the corresponding homologs of Ta0020,
BLASTp was performed using *T. atroviride* IMI206040 as the reference genome. The numbers of the
shortest paths between both TFs, namely, CRE1 and XYR1, and the selected CAZymes were
identified using the Pesca v. 3.0 plugin (Scardoni, et al. 2016) in Cystoscope. The results were
visualized using the R package pheatmap (Kolde 2019).

3   Results

3.1   Molecular phylogeny of the evaluated *T. harzianum* strains

Although Kubicek, et al. (2019) deeply investigated the phylogenetic relationships in the genus
Trichoderma, no results were reported for Th3844, Th0179, and Ta0020. Here, we modeled a
phylogenetic tree based on the ITS sequence of 14 *Trichoderma spp.*, including those of our study
strains (Supplementary Material 1: Supplementary Figure 1). According to the results, high genetic
proximity between Th3844 and Th0179 was observed. In contrast, both strains were phylogenetically
distant from Ta0020, which grouped with other *T. atroviride* strains. *Neurospora crassa* and
Fusarium oxysporum were used as outgroups.

Additionally, to represent the evolutionary relationships of CRE1 and XYR1 among Th3844,
Th0179, and Ta0020, a phylogenetic analysis was performed while considering their amino acid
sequences (Figure 1). The phylogenetic trees were modeled based on 26 and 21 protein sequences
related to CRE1 and XYR1, respectively, in the *Trichoderma* genus, including those of our studied
strains. *Fusarium* spp. was used as an outgroup. The NCBI accession number of the sequences used
to model the phylogenetic trees is available in Supplementary Material 2: Supplementary Table 2.

The CRE1 phylogenetic tree indicated a close genetic proximity between Th3844 and Th0179 (with
bootstrap support of 67%) (Figure 1A). These results were supported by the alignment of both
protein sequences, which showed high conservation in the alignment of their amino acid sequences
(with a percent identity of 99.76%) (Supplementary Material 1: Supplementary Figure 2). In contrast,
the CRE1 protein sequence of Ta0020 shared closer affinity with two other *T. atroviride* strains (T23
and IMI206040), with bootstrap support of 99% (Figure 1A). Furthermore, the C_2H_2 domain is
highly conserved in the Th3844, Th0179, and Ta0020 sequences of CRE1, with only one amino acid
change from *T. atroviride* compared to both *T. harzianum* strains (Supplementary Material 1:
Supplementary Figure 2).

In the XYR1 phylogenetic tree, Th3844, Th0179, and Ta0020 were closest to *T. harzianum* CBS
226.95 (with bootstrap support of 97%), *T. harzianum* T6776 (with bootstrap support of 51%), and *T.
atroviride* IMI206040 (with bootstrap support of 88%), respectively (Figure 1B). Compared to the
results of CRE1, the alignment of their amino acid sequences showed a lower percentage of identity,
supporting the phylogenetic results observed (Supplementary Material 1: Supplementary Figure 2).
In addition, the multiple alignments of the XYR1 protein sequence of Th3844, Th0179, and Ta0020
showed reasonable conservation of the Zn2Cys6 and fungal-specific TF domains (Supplementary
Material 1: Supplementary Figure 2).

3.2   Weighted gene coexpression network analysis

Recently, the transcriptomes of Th3844, Th0179, and Ta0020 grown on crystalline cellulose and
glucose after a time course of 96 h were investigated (Almeida, et al. 2021). By using these
transcriptome data and applying the described filters in the WGCNA package, we obtained (I) 11,050
transcripts (Th3844), (II) 11,105 transcripts (Th0179), and (III) 11,021 transcripts (Ta0020)
(Supplementary Material 3: Supplementary Table 3). Based on these transcripts, we calculated
Pearson's correlation matrices. Then, different soft power β values were chosen (48 for Th3844, 8 for
Th0179, and 27 for Ta0020) to obtain the scale-free topology, reaching fit indexes of (I) 0.8 for
Th3844 (mean connectivity of 40), (II) 0.9 for Th0179 (mean connectivity of 842), and (III) 0.85 for
Ta0020 (mean connectivity of 406). Then, the networks were partitioned into manageable groups to
explore the putative coregulatory relationships. In total, we identified 87 groups in Th3844, 75
groups in Th0179, and 100 groups in Ta0020 (Supplementary Material 4: Supplementary Table 4).
Transcripts with expression patterns correlated with XYR1 and CRE1 were identified in the Th3844,
Th0179, and Ta0020 coexpression networks and grouped (described in Supplementary Material 5:
Supplementary Table 5). Among the strains, Ta0020 presented the highest number of transcripts
coexpressed with *cre1* and the lowest number of transcripts coexpressed with *xyr1*; the other strains
showed the opposite profile.

3.3   Functional characterization of the transcripts in the *xyr1* and *cre1* groups

We performed an enrichment analysis of the transcripts in each *cre1* and *xyr1* group of all evaluated
strains using the GO categories (Supplementary Material 1: Supplementary Figures 3-5,
respectively). The *cre1* group of Th3844, Th0179, and Ta0020 presented 50, 38, and 50 enriched GO
terms, respectively, while in the *xyr1* group, 34, 86, and 50 enriched GO terms were found in
Th3844, Th0179, and Ta0020, respectively. Overall, the GO terms describing the transcripts of these
groups were diverse across all strains, indicating a high degree of differences among the biological
processes related to XYR1 and CRE1.

For example, the enrichment analyses of Ta0020 indicated that the carbohydrate metabolic process
might be directly affected by CRE1 (high percentage of enriched GOs). Furthermore, GO terms

related to fungal growth and nucleoside transmembrane transport were observed. In contrast, in
Th0179 and Th3844, organic substance metabolic processes and regulation processes were
pronounced, respectively. Interestingly, response to light stimulus was an enriched GO term in the
cre1 group of Th3844, and fungal-type cell wall organization was an enriched GO term in Th0179. In
the XYR1 group, Ta0020 also presented carbohydrate metabolic process as an enriched GO term.
However, few genes corresponding to this term were observed, while terms related to the regulation
of DNA transcription were pronounced. Interestingly, response to external stimulus and oxidation-
reduction process were notable terms in Th0179 and Th3844, respectively. Furthermore, our results
suggest that the regulation of biological processes was a common enriched term in all the evaluated
strains, with some particularities across the *T. harzianum* strains, including regulation of cell
population proliferation (Th0179) and mRNA metabolic process (Th3844).

The KO functional annotation of the transcripts encoding enzymes and unknown proteins with
enzymatic activity in the *cre1* and *xyr1* groups was also investigated. In the 14 pathway classes
observed in each TF, 13 were shared between the two regulators (Figure 2). Only two pathway
classes, i.e., metabolism of other amino acids and lipid metabolism, were exclusive to CRE1 and
XYR1, respectively. However, the number of identified enzymes in a given pathway class differed in
each strain. This difference stresses the diversity of proteins with several functions, which could
result in exclusive enzymatic performance in cellulose degradation according to the strain.

Next, we summarize some important aspects of the identified metabolic pathways. For example,
Ta0020 showed a greater number of enriched pathway classes in the *cre1* group, while both *T.*
harzianum strains presented the opposite profile, with more enzymatic activity pathways enriched in
the *xyr1* group. Carbohydrate metabolism was a pathway class enriched in both *T. harzianum* strains
in the *xyr1* group, while Th0179 and Ta0020 presented such a profile in the *cre1* group. In the *xyr1*
group, nucleotide metabolism and metabolism of cofactors and vitamins were enriched terms in
Th0179, while xenobiotic degradation and metabolism of terpenoids and polyketides were enriched
terms exclusively in Ta0020. The pathway classes related to secondary metabolite compounds were
also enriched in all evaluated strains in the *cre1* and *xyr1* groups, and Ta0020 presented the highest
number of enzymes related to this term in the *cre1* group.

In this study, network analyses were conducted to investigate how the molecular basis of XYR1 and
CRE1 for cellulose-degrading enzyme production varies among *T. harzianum* strains. Thus, after
analyzing the functional composition of the transcripts in the *cre1* and *xyr1* groups of all evaluated
strains, particular attention was given to CAZymes, TFs, and transporters (Figure 3). We observed
that in the *cre1* group, Ta0020 showed the highest number of transcripts encoding CAZymes, TFs,
and transporters, while in the *xyr1* group, both *T. harzianum* strains presented a great number of
coexpressed transcripts encoding TFs, followed by those encoding transporters. Furthermore, Th3844
showed a high number of CAZymes in the *xyr1* group.

We identified the main CAZyme classes, including glycoside hydrolases (GHs), carbohydrate
esterases (CEs), glycosyltransferases (GTs), and polysaccharide lyases (PLs), coexpressed with *xyr1*
and *cre1* transcripts. To determine the similarities and differences across the strains, their CAZyme
profiles were compared (Figure 4). In the *xyr1* and *cre1* groups, the GH family, which encompasses
enzymes that hydrolyze glycosidic bonds, was identified in all the evaluated strains with different
numbers of transcripts (Figure 4A and B). In the *cre1* group, Ta0020 presented the highest number of
classified transcripts of the GH family, followed by Th0179 and Th3844. In the *xyr1* group, Th3844
presented a higher number of GHs than Th0179 and Ta0020. Furthermore, transcripts encoding CEs,

which hydrolyze ester bonds, were associated with the *cre1* transcript in Ta0020 and Th0179,
whereas transcripts of the PL family, which cleave bonds in uronic acid-containing polysaccharide
chains, were coexpressed with the *xyr1* transcript in only Th3844. In addition, transcripts of the GT
family, which synthesizes glycosidic bonds from phosphate-activated sugar donors, were present in
the *cre1* group of Th0179 and the *xyr1* groups of Th3844 and Ta0020. We also investigated the
quantification of each CAZyme family in the *cre1* and *xyr1* groups of all evaluated strains (Figure 4C
and D). Overall, Ta0020 presented a high number of GHs belonging to family 18, coexpressed with
the *cre1* transcript, while Th0179 exhibited a significant amount of GT90 (Figure 4C). Different
CAZyme profiles were observed in the *xyr1* groups, and Th3844 showed the highest diversity of
CAZyme families (Figure 4D).

Several positive and negative regulators are involved in the expression of CAZyme-coding genes and
other proteins required for lignocellulose breakdown. Thus, we sought to identify transcripts
encoding TFs coexpressed in the *cre1* and *xyr1* groups (Figure 5). In both groups, transcripts
encoding Zn_2Cys_6-type TFs were identified in all the evaluated strains with different numbers of
transcripts. In the *cre1* group of Ta0020, the number was higher than that observed in Th0179 and
Th3844 (Figure 5A), while in the *xyr1* group, we observed the opposite profile (Figure 5B). Most
Zn_2Cys_6 proteins also contain a fungal-specific TF domain, which was coexpressed with both *cre1*
and *xyr1* transcripts in Th3844 and Ta0020. In this last strain, a C6 zinc finger domain TF was
coexpressed with *xyr1*. Transcripts encoding C_2H_2-type TFs were identified in Ta0020 and *T.*
harzianum strains in the *cre1* and *xyr1* groups, respectively. Furthermore, in the *xyr1* group, we
found a transcript encoding the SteA regulator, which is involved in the regulation of fungal
development and pathogenicity (Hoi and Dumas 2010), in Th0179.

Transporters are also important players in the degradation of plant biomass. Here, transcripts
encoding transport proteins coexpressed with *cre1* and *xyr1* were selected (Figure 5). Among them,
transcripts encoding major facilitator superfamily (MFS) transport proteins were coexpressed with
cre1 in Th0179 and Ta0020 (Figure 5C) and *xyr1* in Th0179 and Th3844 (Figure 5D). Amino acid
transporters were present in the *xyr1* groups of all strains and the *cre1* group only of Th0179.
Transcripts encoding the ATP-binding cassette (ABC) were coexpressed with *xyr1* only in Th3844.
Th0179 showed transcripts encoding several types of transporters coexpressed with *xyr1*, e.g., vesicle
transporter SEC22, cation transporting ATPase, and calcium-dependent mitochondrial carrier, while
Ta0020 coexpressed with *cre1*, e.g., UDP-galactose transporter and CNT. Transcripts encoding ion
transporters were present in both the *cre1* and *xyr1* groups of Ta0020 and Th0179. In the *cre1* group,
such transcripts included zinc and magnesium transporter proteins in Ta0020 and potassium
transporter proteins in Th0179. In the *xyr1* group, such transcripts include $Cd^{2+}Zn^{2+}$ transporters in
Ta0020 and Ca^{2+} transporters in Th0179. Th0179 and Th3844 also present a G protein coexpressed
with the *cre1* transcript.

To better understand the regulation mediated by the studied TFs in the evaluated *T. harzianum*
strains, we also investigated other classes of proteins coexpressed with the *cre1* and *xyr1* transcripts.
For example, phosphatases, kinases and TFs are key components in cellular signaling networks.
Transcripts encoding kinase proteins were coexpressed with *cre1* transcripts in Ta0020 and Th3844
and *xyr1* transcripts in Ta0020 and Th0179. In addition, all strains showed transcripts encoding
phosphatases coexpressed with *cre1,* and in Th0179 and Th3844, such transcripts were coexpressed
with *xyr1*. In addition, transcripts encoding cytochrome P450 genes, which constitute an important
group of enzymes involved in xenobiotic degradation and metabolism, were found in the *cre1* groups
of all the evaluated strains.

Degradation

Furthermore, different quantities of differentially expressed transcripts were found in Ta0020 (78)
and Th0179 (6) (described in Supplementary Material 6: Supplementary Table 6 and Supplementary
Material 1: Supplementary Table 7). In Ta0020, most upregulated transcripts under cellulose growth
(70) were associated with the *cre1* group, including *cre1*, whereas *xyr1* demonstrated a potentially
significant expression level under glucose growth. In Th3844, *cre1* significantly modulated the
expression of the two carbon sources, with a high expression level under glucose growth. Even with
the reduced differentiation observed (-1.13), this phenomenon was considered in other studies
(Antonieto, et al. 2014; Castro, et al. 2016). In Th0179, *xyr1* and *cre1* revealed a low modulation in
transcript expression between growth under cellulose or glucose. However, *xyr1* had a higher
transcript expression level under glucose than *cre1*, which had a higher transcript expression level
under cellulose growth conditions. In Th3844, *xyr1* exhibited similar results.

3.4   Assessing the groups' network topologies

To identify the main aspects of the network topologies, we applied the HRR methodology to the
transcriptome data, which were also used to model the networks using the WGCNA methodology
(Figure 6A-C).

In the CRE1 and XYR1 subnetworks, different connection profiles were observed in each strain
(Figure 8). While the transcripts in the *cre1* groups showed less ranked connections in the WGCNA
groups of the *T. harzianum* strains Figure 6D and E), Ta0020 showed the opposite profile. In this
strain, the transcripts in the *cre1* group were more strongly connected (Figure 6F). In contrast, in the
xyr1 groups, the transcripts were more densely connected in the *T. harzianum* strains (Figure 6G and
H), while in Ta0020, the transcripts were separated (Figure 6I).

Transcripts at the top of the degree distribution, which are defined here as hubs, are topologically
important to the network structure and are often functionally relevant (Luscombe et al. 2004). Thus,
the hub nodes were sought in the *cre1* and *xyr1* groups. In the *cre1* groups, transcripts encoding a
SET-domain protein, an ATP synthase, and a transcript not yet annotated were hub nodes of Th3844,
Th0179, and Ta0020, respectively. In Ta0020, we found that such an uncharacterized transcript had a
degree value equal to 52, which differs from the *T. harzianum* strains in which the hub nodes had
degree values of 4 (Th3844) and 9 (Th0179) (Supplementary Material 7: Supplementary Table 8). In
the *xyr1* groups, transcripts encoding an ATP-dependent RNA helicase, a cytokinesis sepA, and
hypothetical proteins were found as hub nodes of Th3844, Th0179, and Ta0020, respectively
(Supplementary Material 7: Supplementary Table 8). Here, we show that the hub nodes in the *cre1*
group of Ta0020 had a degree value of 1 and all encoded hypothetical proteins, while in the *T.*
harzianum strains, the hub nodes had degree values of 13 (Th3844) and 7 (Th0179). Among the other
transcripts with a degree value lower than that of those at the top of the degree distribution, in the
xyr1 group, GH76 and Zn_2Cys_6-type TFs (degree value of 6) were identified as candidate hub genes
of Th0179, while regulator-nonsense transcripts (degree value of 9) and Zn_2Cys_6-type TFs (degree
value of 8) were identified as candidate hub genes of Th3844. In the *cre1* group, the proteasome
component PRE3 (degree value of 2) in Th3844 was found as a hub node, and replication factor C
and phosphoglucomutase (degree value of 5) were found in Th0179. In addition, in Ta0020, we
identified calcium calmodulin-dependent kinase (degree value of 9) and GH93 (degree value of 8) as
hub nodes.

In all evaluated strains, as a reduced number of differentially expressed transcripts was found in the
cre1 and *xyr1* groups, we expanded our investigation of the studied groups by selecting the first

neighbors of XYR1 and CRE1 in global HRR networks. Different quantities of *cre1* neighbors were
found in Th3844 (59), Th0179 (21), and Ta0020 (25), and different quantities of *xyr1* were found in
Th3844 (46), Th0179 (59), and Ta0020 (70) (described in Supplementary Material 8: Supplementary
Table 9). Regarding the *cre1* neighbors, these transcripts were distributed among 24 groups in
Th3844, 15 groups in Th0179, and 16 groups in Ta0020. In contrast, considering the *xyr1* neighbors,
such transcripts were distributed among 19 groups in Th3844, 32 groups in Th0179, and 42 groups in
Ta0020. The direct coexpressed neighbors of *xyr1* included downregulated transcripts in Th3844 (4),
Th0179 (5), and Ta0020 (22) and upregulated transcripts in Th3844 (1), Th0179 (1), and Ta0020 (1).
Interestingly, only Ta0020 showed differentially expressed transcripts as the first neighbor of the
cre1 transcript (4 downregulated and 4 upregulated). Overall, these differentially expressed
transcripts included transporters, CAZymes, kinases, and other proteins related to the regulation
process.

For simplification, only a few neighboring transcripts of *xyr1* and *cre1* in Th3844, Th0179, and
Ta0020 are shown in Table 1. In Th3844, TFs, transporters, a cytochrome, and a phosphatase were
found to be first neighbors of the *cre1* transcript, while TFs, transporters, and CAZymes were found
to be the first neighbors of the *xyr1* transcript. In Th0179, several types of proteins, including a
kinase, were found to be first neighbors of *cre1*, while transporters, CAZymes, and a TF were found
to be first neighbors of the *xyr1* transcript. In Ta0020, several types of proteins, including kinases,
were found to be first neighbors of *cre1*, while transporters, CAZymes, TFs, and a DNA ligase were
found to be first neighbors of the *xyr1* transcript. Hypothetical proteins were also found to be
neighbors of the *cre1* and *xyr1* transcripts of all strains, indicating that both regulators might have a
regulatory influence on uncharacterized proteins.

We also applied a shortest-path network analysis to identify possible transcripts outside of the *xyr1*
and *cre1* groups that may be influenced by CRE1 and XYR1 (described in Supplementary Material
9: Supplementary Table 10). As the shortest path, we considered the minimal number of edges that
need to be traversed from a node to reach another node (Koutrouli, et al. 2020). As both regulators
act on the gene expression of hydrolytic enzymes related to plant biomass degradation, we sought to
determine the number of shortest paths from XYR1 and CRE1 to the selected CAZymes. In all
evaluated strains and both TFs, we found a similar quantity of shortest paths (Figure 9A and
Supplementary Material 1: Supplementary Figures 6 and 7). However, some aspects should be noted.
For example, Th0179 presented transcripts encoding CAZymes from the GH1, GH5, and GH6
families, which are more distantly related to CRE1, while Ta0020 showed only a CAZyme from the
GH6 family satisfying this requirement. In Th3844, we found a transcript encoding a CAZyme from
GH3 strongly related to XYR1, and CAZymes with the CBM1 domain were also closely connected
with such a regulator. In Th0179, a CAZyme from GH55 exhibited a close relationship with XYR1,
while in Ta0020, CAZymes from the GH7, GH72, and GH1 families and the CBM1 domain also
presented such a profile. In addition, given a determined shortest path, we also considered the
number of possibilities that may have occurred (Figure 9B and Supplementary Material 1:
Supplementary Figure 6 and 7). Across all strains and in both TFs, significant differences were
observed in the number of chances. For instance, a significant number of possible minimum paths
between CRE1 and a CAZyme from the GH6 family were observed in Ta0020, likely suggesting a
strong influence of this repressor regulator on enzyme activity. Similarly, CAZymes from the AA9
and GH16 families appeared to be affected by XYR1. In Th0179, CAZymes from GH10, GH6, and
GH1 presented a higher number of possible shortest paths related to CRE1, potentially indicating an
impact of this TF on these enzyme activities.

We also performed a shortest-path analysis to identify transitive transcripts between *cre1* or *xyr1* and
the selected CAZymes, which allowed us to discover new transcripts that may be involved in the
same biological process. For simplification, we only chose a few shortest paths to explore such
transitive transcripts. In Th3844, NADH:ubiquinone oxidoreductase (THAR02_08259), which is the
largest multiprotein complex of the mitochondrial respiratory chain (Whitehouse, et al. 2019), was
found between GH45 with a CBM1 domain (THAR02_02979) and *cre1*. We also found another
transcript involved in ATP synthesis, NAD(P) transhydrogenase beta subunit (THAR02_08260),
between GH1 (THAR02_05432) and *cre1*. Furthermore, a transcript encoding a component of the
endoplasmic reticulum quality control system called ER-associated degradation (THAR02_08220)
(Phillips, et al. 2020) was found between GH16 (THAR02_03302) and *cre1*. Interestingly, in
Ta0020, a phosphotyrosine phosphatase was found in the shortest path between GH1
(TRIATDRAFT_135426), which is a homolog of GH1 (THAR02_05432) in *T. harzianum* strains,
and *cre1*. This phosphatase is involved in protein phosphorylation, which is a key posttranslational
modification critical for the control of many cellular functions (Nasa and Kettenbach 2018). In
Th3844, we found a transcript encoding a mitochondrial outer membrane porin (THAR02_07078)
between CBM1 (THAR02_02133) and *xyr1*. These protein transporters transport small molecules
and play significant roles in diverse cellular processes, including the regulation of mitochondrial ATP
and calcium flux (Grevel and Becker 2020). In Ta0020, methyltransferase domain-containing
(TRIATDRAFT_292180), which is important for the regulation of chromatin and gene expression
(Kouzarides 2007), was found between GH1 (TRIATDRAFT_150220) and *xyr1*.

4   Discussion

Although the importance of XYR1 and CRE1 in the expression of CAZyme-encoding genes and
other proteins required for lignocellulose degradation is evident, the transcriptional regulation
mediated by both proteins in *T. harzianum* strains remains poorly explored (Delabona, et al. 2017,
2021). A previous study demonstrated that some genes encoding regulatory proteins, such as *xyr1*,
evolved by vertical gene transfer in *Trichoderma* spp. (Druzhinina, et al. 2018). Here, we inferred the
evolutionary relationships of XYR1 and CRE1 in the evaluated strains. In both regulatory proteins, a
great genetic distance was observed between the amino acid sequences of Ta0020 and those of
Th3844 and Th0179. These findings supported the genetic tree of the evaluated species in which
Ta0020 was grouped with other *T. atroviride* strains. However, among the *T. harzianum* strains, we
noted that the CRE1 amino acid sequences were more highly conserved than those of XYR1. These
results are consistent with previous studies reporting that while the function of CRE1 is conserved
throughout the fungal kingdom (Adnan, et al. 2017), the role of XYR1 greatly differs across
ascomycete fungi (Klaubauf, et al. 2014).

Recently, different types of enzymatic profiles across *Trichoderma* species were reported, and
Th3844 and Th0179, which have hydrolytic potential, have higher cellulase activity during growth
on cellulose than Ta0020 (Almeida, et al. 2021). Because such diversity in the enzyme response can
be affected by the specific functionalities of regulatory proteins and evolutionary divergence between
XYR1 and CRE1 (Benocci, et al. 2017; Klaubauf, et al. 2014), we aimed to investigate how both TFs
could affect the activities of the transcripts in response to cellulose degradation in the *T. harzianum*
strains. Because transcripts that share the same function or are involved in the same regulatory
pathway tend to present similar expression profiles and, hence, form modules (Wolfe, et al. 2005),
networks were modeled for Th3844, Th0179, and Ta0020, and the last strain was used to assess the
differences across *Trichoderma* species. In fungi, such methodology has been successfully used to

provide insight into the regulatory mechanisms of hydrolysis (Arntzen, et al. 2020; Borin, et al. 2018;
Li, et al. 2020a).

In recent years, network methodologies have been employed for the discovery of new targets
associated with biological processes (Cortijo, et al. 2020; Liu, et al. 2021; Zhang, et al. 2021).
Because such approaches might provide mechanisms for inferring new gene relationships and
functions, experimental validation by the construction of mutants is usually a way of corroborating
these insights. In this context, the construction of mutants can show the influence of a unique missing
gene on genome regulation, helping to corroborate the hypotheses raised in the present study.
However, due to the inefficient homologous recombination machinery of filamentous fungi, once a
fungus reproduces asexually, a low frequency of correct genomic integration has been observed for
Trichoderma spp. (Liu, et al. 2017; Vieira, et al. 2018; Zeilinger 2004). These challenges can be
overcome by using the CRISPR/Cas9 system, which represents a good alternative in *Trichoderma*
spp. (Fonseca, et al. 2020; Vieira, et al. 2021) and may enable the future establishment of efficient
protocols for each strain described here. Thus, by predicting the profiles of the transcripts
coexpressed with XYR1 and CRE1, the results can be used to engineer mutant strains with the
objective to investigate the influence of the regulation of such TFs *in vitro*.

We observed that each network had a different configuration with distinct module profiles of XYR1
and CRE1, providing insight into how these TFs might act in specific ways in different *Trichoderma*
species. These functional differences in the *xyr1* and *cre1* groups can be attributed to previously
reported differences in the regulatory mechanisms of hydrolysis (Almeida, et al. 2021). By analyzing
both groups, we observed that more transcripts were coexpressed with XYR1 in the *T. harzianum*
strains than *T. atroviride*, which had more transcripts coexpressed with CRE1. However, although
the phylogenetic similarity between the evaluated *T. harzianum* strains shows that they share a
largely common genetic background, their profiles of transcripts coexpressed with *xyr1* and *cre1*
differed.

4.1 Insight into the genetic impacts of XYR1 and CRE1

To obtain insight into the functional profile of the *cre1* and *xyr1* groups, GO enrichment analyses of
transcripts from these groups of all evaluated strains were performed. Here, response to external
stimulus was a notable GO term in Th0179 in the *xyr1* group, suggesting that the transcripts of these
group likely respond to external environmental conditions, such as carbon sources. Furthermore,
interconnections between nutrient and light signaling pathways have been reported in filamentous
fungi, such as *N. crassa* and *T. reesei*, with substantial regulation by photoreceptors (Schmoll 2018).
More interestingly, the influence of light on CRE1 functions has been reported (Monroy, et al. 2017),
supporting our findings showing that the response to a light stimulus was an enriched GO term in the
cre1 group of Th3844. In contrast, the enrichment analysis of Ta0020 indicates that the activity of
CRE1 may be stronger in such fungi, directly repressing genes related to plant cell wall-degrading
enzymes, which was not observed in the *T. harzianum* strains. In these strains, organic substance
metabolic processes (Th3844) and regulation processes (Th0179) were enriched GO terms. In
addition, fungal-type cell wall organization was an enriched term of Th0179 in the *cre1* group.

By investigating the KO functional annotation of the *cre1* and *xyr1* groups, we suggest that both
regulators act on the same enzymatic pathways, which was expected due to antagonism in their
function, i.e., while CRE1 is the main repressor of genes encoding proteins related to lignocellulose
degradation, XYR1 is the main activator of such genes. However, each triggers a specific metabolic
pathway according to the strain; therefore, some aspects are noteworthy. For example, it has been

reported that the *Trichoderma* species display several mechanisms during their antagonistic action
against plant pathogens, including the production of secondary metabolites (Malmierca, et al. 2015).
According to our findings, due to the highest number of pathways related to secondary metabolites,
including the metabolism of terpenoids and polyketides (Mukherjee, et al. 2008), we suggest that
CRE1 and XYR1 participate in the regulation of secondary metabolism compounds in *T. atroviride.*
Furthermore, XYR1 in *T. harzianum* strains appeared to be deeply connected with enzyme pathways
related to the metabolism of carbohydrates, especially in the Th3844 strain.

Through a network analysis, we identified transcripts encoding CAZymes coexpressed with *xyr1* and
cre1 in all evaluated strains. We identified CAZyme families responsible for cellulose degradation
(e.g., GH12); hemicellulose degradation (e.g., GH16, GH17, CE1, and CE5); pectin degradation
(e.g., GH28 and GH93); and other CAZy families with multiple activities or minor activities on
lignocellulosic substrates, such as GH2, GH5, GH43, GH95, GH30, and GH39 (de Vries, et al. 2017;
Kameshwar, et al. 2019). Although Ta0020 presented the highest number of GHs coexpressed with
CRE1, it was found to be less efficient than the other *Trichoderma* strains in degrading plant biomass
(Almeida, et al. 2021). In the GH class, enzymes belonging to the GH18 family, which is mainly
represented by chitinase-like proteins, are directly related to fungal cell wall degradation in
mycoparasite species of *Trichoderma* (Gruber and Seidl-Seiboth 2012). Four GH18 enzymes were
coexpressed with CRE1 in *T. atroviride*, which is widely used as a biocontrol agent in agriculture.
However, in Ta0020, enzymes from the GH75 family of chitosanases were coexpressed with *xyr1*.
Therefore, the degradation of chitosan, which is a partially deacetylated derivative of chitin (Hahn, et
al. 2020), is a relevant aspect of mycoparasitism that may be influenced by the activity of XYR1.
Recently, transcripts encoding GHs were found to be coexpressed with some TFs involved in
biomass degradation, e.g., XYR1 (Borin, et al. 2018). In the present study, the transcripts encoding
hydrolytic enzymes could be sorted into other groups formed in the networks mainly because their
expression patterns differed from those of the *xyr1* and *cre1* transcripts.

Several fungal TFs have been described to be directly involved in the regulation of plant biomass
utilization (Benocci, et al. 2017). Most TFs belong to the zinc cluster family, including Zn_2Cys_6- and
C_2H_2-type TFs, which are characterized by the presence of zinc finger(s) in their binding domains.
Most positive regulators appear to belong to the Zn_2Cys_6 class, while repressors belong to the C_2H_2
class (Benocci, et al. 2017). Here, we found the highest number of transcripts encoding the Zn_2Cys_6
and C_2H_2 classes in the *xyr1* and *cre1* groups of Th0179 and Ta0020, respectively. Both *T. harzianum*
strains had a similar profile of transcripts coexpressed with CRE1 and XYR1. However, SteA, a
C_2H_2-type TF, was coexpressed with the *xyr1* transcript of Th0179. This regulator has been described
as an important player in fungal environmental adaptation in response to nutrient deprivation, the
production of extracellular proteins involved in the degradation of complex substrates (Hoi and
Dumas 2010), and the mediation of the regulatory role of mitogen-activated protein kinase (MAPK)
during mycoparasitic responses (Gruber and Zeilinger 2014). In *Aspergillus nidulans*, SteA is
required for sexual development (Vallim, et al. 2000).

During plant biomass degradation, fungi secrete extracellular enzymes to decompose polysaccharides
into small molecules, which are then imported into cells through transporters (Sloothaak, et al. 2016).
One of the most relevant sugar transporter families in filamentous fungi is the MFS family (Zhang, et
al. 2013). Here, we found the highest number of transcripts encoding MFS coexpressed with *cre1* and
xyr1 in Ta0020 and Th3844, respectively. Among the *T. harzianum* strains, compared with Th3844,
Th0179 showed transcripts encoding several types of transporter proteins, such as transporter
proteins of calcium ions (THAR02_04202, THAR02_03989, and THAR02_01350). It has already

been reported that metal ions, such as Ca^{2+}, have a positive effect on the mycelial growth of *T. reesei*
and cellulase production (Chen, et al. 2016). This molecular signaling mechanism is mediated by
cations transporting ATPase and calcium-dependent mitochondrial carriers, which are both
components of $Ca^{2+/}$calmodulin signal transduction, including the TF Crz1 (Chen, et al. 2016;
Martins-Santana, et al. 2020). Therefore, investigating the role of proteins related to calcium
transporters in the induction of genes responsive to lignocellulose degradation in Th0179 cells is
important since these proteins could transport cations that activate gene expression. Furthermore,
transcripts encoding G proteins were coexpressed with *cre1* in Th0179 and Ta0020. Heterotrimeric G
proteins have been well studied in several *Trichoderma* species. In saprophytic species, such proteins
are involved in the nutrient signaling pathway in connection with a light response, triggering the
posttranscriptional regulation of cellulase expression (Hinterdobler, et al. 2021); in mycoparasitic
species, G protein-coupled receptors are involved in the regulation of processes related to
mycoparasitism (Zeilinger and Atanasova 2020).

Although CAZymes, TFs, and transporters play an important role in cellulose degradation, these
types of proteins represented only a small percent of the transcripts coexpressed with XYR1 and
CRE1 compared to the other protein classes as follows: in the *cre1* group, (I) 10.4% (Th0179), (II)
5.7% (Th3844), and (III) 17.9% (Ta0020), and in the *xyr1* group, (IV) 11.8% (Th0179), (V) 11.9%
(Th3844), and (VI) 16% (Ta0020). Since the regulation of genes involved in biomass breakdown is a
complex process that involves several signaling pathways, we expected to find proteins with a great
range of functions coexpressed with *xyr1* and *cre1*. For example, various kinases and phosphatases
were coexpressed with *xyr1* and *cre1* in all evaluated strains. It has been reported that cellulase gene
expression can be regulated by the dynamics of protein phosphorylation and dephosphorylation,
which involve protein kinases and phosphatases, respectively (Schmoll, et al. 2016). Furthermore, in
filamentous fungi, phosphorylation is a prerequisite for CRE1 activity (Cziferszky, et al. 2002; de
Assis, et al. 2021; Han, et al. 2020). Therefore, in *T. harzianum*, it is important to elucidate the role
of kinases and phosphatases in the regulation of CRE1 function. In all evaluated strains, cytochrome
P450 coding genes represented another class of proteins coexpressed in the *cre1* group. It has been
reported that these enzymes are important for cells to perform a wide variety of functions, including
primary and secondary metabolism, xenobiotic degradation, and cellular defense against plant
pathogenic fungi (Chadha, et al. 2018; Fan, et al. 2013; Siewers, et al. 2005). To expand previous
findings concerning *Trichoderma* spp. (Chadha, et al. 2018), their similar expression pattern with the
cre1 transcript makes them important candidates for an extensive investigation in the cellulose
degradation context.

We also investigated the expression profiles of the transcripts in the *cre1* and *xyr1* groups, including
xyr1 and *cre1*, under cellulose and glucose growth. Castro, et al. (2016) reported that in *T. reesei*, the
expression level of *xyr1* was minimal in the presence of glucose; this phenomenon was not observed
in Ta0020 in the present study. In addition, in Ta0020, *cre1* was upregulated in the presence of
cellulose, which was not expected due to the role of CRE1 in the repression of cellulolytic and
hemicellulolytic enzymes when an easily metabolizable sugar, i.e., glucose, is available in the
environment (Benocci, et al. 2017). In contrast, in Th3844, the *cre1* expression level in the presence
of glucose was higher than that in the presence of cellulose. The gene coexpression networks were
modeled based on the transcript expression level under two sets of conditions (cellulose and glucose).
Considering that the expression of the *cre1* transcript in Ta0020 was upregulated under cellulose
growth, upregulated transcripts were expected and found in the *cre1* group. Recently, Almeida et al.
(Almeida, et al. 2021) reported that Ta0020 presented the highest number of differentially expressed
transcripts under cellulose growth conditions relative to glucose, followed by Th3844 and Th0179.
Here, the quantity of upregulated transcripts was low in the *cre1* and *xyr1* groups of the *T. harzianum*

Degradation

strains. These results may indicate that both TFs have a basal expression level at 96 h of
fermentation; thus, the transcripts grouped with such regulatory proteins presented the same
expression pattern and, mostly, were not differentially expressed transcripts.

4.2   Examining the network's topology

To extract additional information from the networks, we characterized the network topologies of the
modules identified separately. We might infer that many transcripts were under the influence of the
repressor CRE1 in Ta0020, while in the *T. harzianum* strains, the transcripts seem to have been
affected by the activator XYR1. Therefore, the described profiles may indicate that in *T. harzianum*,
the transcripts, including those encoding XYR1, act together to perform a determined biological
function that is favorable to the expression of genes related to cellulose degradation. In contrast, in *T.*
atroviride, the transcripts in the *cre1* group appeared to act on the same biological process, which
may be related to the repression of genes encoding hydrolytic enzymes and other proteins required
for cellulose degradation.

Hub transcripts were identified in the *cre1* and *xyr1* groups, and new targets were discovered. In the
cre1 group, a transcript with a SET-domain coding gene was found as a hub node of Th3844. Such
SET-domain proteins participate in chromatin modifications by methylating specific lysines on the
histone tails (Kouzarides 2007). In filamentous fungi, the epigenetic regulation of holocellulase gene
expression has already been reported (Zeilinger, et al. 2003), and CRE1 plays an important role in
nucleosome positioning (Ries, et al. 2014). Interestingly, ATP synthase, which synthesizes ATP from
ADP and inorganic phosphate on mitochondria (Burger, et al. 2003), was found to be a hub node of
Th0179. Thus, we might infer that a great number of transcripts were coexpressed with a hub node
related to mitochondrial ATP production, which is the main energy source for intracellular metabolic
pathways (Neupane, et al. 2019). In the *xyr1* group, a transcript encoding an ATP-dependent RNA
helicase was found as a hub node of Th3844. Such enzymes catalyze the ATP-dependent separation
of double-stranded RNA and participate in nearly all aspects of RNA metabolism (Jankowsky 2011).
Additionally, the sepA transcript was identified as a hub node of Th0179. In *Aspergillus nidulans*, the
sepA gene encodes a member of the FH1/2 protein family, which is involved in cytokinesis and the
maintenance of cellular polarity, i.e., related to the cell division process (Harris, et al. 1997). Overall,
our findings suggest that the hub genes were not necessarily the most effective genes related to
lignocellulose deconstruction, confirming the indirect action of XYR1 and CRE1 in *T. harzianum* on
regulatory hydrolysis mechanisms. In Ta0020, transcripts encoding hypothetical proteins were
identified as potential hub nodes in both groups, providing new targets for further studies evaluating
their functions in fungal physiology.

We also investigated the first neighbors of the *cre1* and *xyr1* transcripts in the modeled global HRR
networks of all evaluated strains. While Ta0020 showed differentially expressed transcripts as the
first neighbor of the *cre1* transcript, interestingly, both *T. harzianum* strains presented the opposite
profile in which only the first neighbors of the *xyr1* transcript were differentially expressed. In
addition, the evaluated strains showed different numbers of neighbors of the *cre1* and *xyr1*
transcripts, which were distributed among several groups. Such a divergent profile may indicate that
each TF affects a set of transcripts in a specific way that varies among the strains. Overall, we
identified CAZymes, TFs, and transporters as first neighbors of *cre1* and *xyr1*, confirming the results
obtained in this study using the WGCNA approach.

Another property of a network's topology is the shortest paths connecting two transcripts
(Pavlopoulos, et al. 2011). Here, the significant shortest paths between both studied TFs, i.e., CRE1
and XYR1, and the CAZymes with a higher-level expression under cellulose growth conditions were
investigated in all strains. The average shortest pathway distance between the transcripts encoding
CRE1 or XYR1 and all selected CAZymes was relatively short, which may be attributed to a network
phenomenon called the small-world effect, i.e., networks can be highly clustered with a small number
of necessary steps to reach one node from another (Maier 2019). However, several aspects should be
highlighted. We identified CAZymes responsible for cellulose degradation, such as GH6 (Th0179
and Ta0020), and other CAZy families with multiple activities or minor activities on lignocellulosic
substrates, such as GH1 and GH5 (Th0179) (de Vries, et al. 2017; Kameshwar, et al. 2019) with a
low number of minimum pathways to CRE1 in Ta0020. In contrast, CAZymes responsible for
cellulose degradation, such as CBM1 domain (Th3844 and Ta0020) and GH7 (Ta0020), for
hemicellulose degradation, such as GH55 (Th0179), and CAZy families with multiple activities or
minor activities on lignocellulosic substrates, such as GH3 (Th3844) and GH1 (Ta0020) (de Vries, et
al. 2017; Kameshwar, et al. 2019), were identified with a high number of minimum pathways to
XYR1. Because certain shortest paths are not necessarily unique, we also considered the number of
possibilities that the paths may occur. Our results suggest that the number of possible minimum paths
between some CAZymes and CRE1 or XYR1 was higher than of others. For example, a great
number of possible shortest paths was observed between CRE1 and GH6 in Ta0020 and between
CRE1 and GH10, GH6, or GH1 in Th0179. In contrast, in Ta0020, such profiles were observed
between XYR1 and CAZymes with cellulolytic (AA9) and hemicellulolytic activities (GH16) and
between CRE1 and GH6 with a CBM1 domain.

Furthermore, we performed a shortest path analysis to explore transitive transcripts between two
nodes. Considering that the lowest number of transitive transcripts between two nodes may indicate
the need for fewer signal pathways, representing a more direct relation, we chose to investigate the
shortest paths with only one transcript between the desired targets. Interestingly, transcripts encoding
proteins related to ATP synthesis were found between GH45 with a CBM1 domain and *cre1* and
between GH1 and *cre1* in Th3844, which could be explained by the demand for energy required for
CRE1 to exercise its repressor activity on these hydrolytic enzymes. However, in Th3844, a
transcript encoding a protein involved in quality control processes that center on the endoplasmic
reticulum was found between GH16 and *cre1*, indicating that a signaling pathway is triggered by
CRE1 to repress the expression of such an enzyme. Curiously, in Ta0020, a phosphotyrosine
phosphatase was found in the shortest path between GH1, which is a homolog of GH1 in the *T.
harzianum* strains, and *cre1*. Such proteins are involved in posttranslational modification, including
the phosphorylation process, which plays an essential role in signal transduction to achieve CCR by
CRE1 (Han, et al. 2020; Horta, et al. 2019). In Th3844, we found a transcript encoding a protein
related to the transport of substances across the mitochondrial membrane between CBM1 and *xyr1*,
which may indicate high cellular activity and, therefore, a high demand for energy for gene
expression. In Ta0020, a transcript encoding a protein involved in the regulation of chromatin and
gene expression was found between GH1 and *xyr1*, which may indicate an intermediated process for
the expression of such an enzyme.

In conclusion, biological networks represent a powerful approach to accelerate the elucidation of the
molecular mechanisms underlying important biological processes. Our findings suggest that the set
of transcripts related to XYR1 and CRE1 varies among the studied *T. harzianum* strains, suggesting
regulatory differences in enzymatic hydrolysis. These findings corroborate previous studies in which
differences in biomass degradation and enzyme production between strains of the same species have
been reported (de Vries, et al. 2017; Thanh, et al. 2019; Tolgo, et al. 2021). Furthermore, such

Degradation

transcripts were not limited to CAZymes and other proteins related to biomass degradation. Thus, we
suggest that both TFs play a role in the undirected regulation of gene encoding proteins related to
cellulose degradation, and multiple pathways related to gene regulation, protein expression, and
posttranslational modifications may be triggered by the studied TFs. We expect that our results could
contribute to a better understanding of fungal biodiversity, especially regarding the transcription
regulation involved in hydrolytic enzyme expression in *T. harzianum*. By describing new potential
targets involved in the cellulose degradation pathway, this knowledge can be used to develop genetic
manipulation strategies and expand the use of *T. harzianum* as an enzyme producer in
biotechnological industrial applications.

Conflicts of interest

706 *The authors declare that the research was conducted in the absence of any commercial or financial*
707 *relationships that could be construed as potential conflicts of interest.*

Author contributions

709 **RRR:** Writing - original draft, Methodology, and Conceptualization. **AHA:** Methodology, Software,
710 Writing - review & editing, and Formal analysis. **DAA:** Resources, and Writing - review & editing.
711 **JAFF:** Writing - review & editing. **MACH:** Resources and Writing - review & editing. **APS:**
712 Supervision, review & editing, and funding acquisition.

Funding

Abbreviations

ABC: ATP-binding cassette
CAZymes: carbohydrate-active enzymes
CCR: carbon catabolite repression
CEs: carbohydrate esterases
CRE1: carbon catabolite repressor 1
GHs: glycoside hydrolases
GO: Gene Ontology
GTs: glycosyltransferases
HRR: highest reciprocal rank
iTOL: Interactive Tree of Life
ITS: internal transcribed spacer
JTT: Jones-Taylor-Thornton
K2P: Kimura two-parameter

Degradation

KEGG: Kyoto Encyclopedia of Genes and Genomes
KO: KEGG Orthology
MEGA: Molecular Evolutionary Genetics Analysis
MFS: major facilitator superfamily
ML: Maximum likelihood
PLs: polysaccharide lyases
Ta0020: *Trichoderma atroviride* CBMAI-0020
tef1: translational elongation factor 1
TFs: transcription factors
Th0179: *Trichoderma harzianum* CBMAI-0179
Th3844: *Trichoderma harzianum* IOC-3844
TOM: topological overlap matrix
TPM: transcripts per million
WGCNA: weighted correlation network analysis
XYR1: xylanase regulator 1

Acknowledgments

We are grateful to the Brazilian Biorenewables National Laboratory (LNBR), Campinas – SP for
conducting the fermentation experiments; the Center of Molecular Biology and Genetic Engineering
(CBMEG) at the University of Campinas, SP for the use of the center and laboratory space; and the
São Paulo Research Foundation (FAPESP), the Coordination of Improvement of Higher Education
Personnel (CAPES, Computational Biology Program), and the Brazilian National Council for
Technological and Scientific Development (CNPq) for supporting the project and researchers.

Data availability statement

All data generated or analyzed in this study are included in this published article (and its
supplementary information files). The raw RNA-Seq datasets were deposited at the NCBI Sequence
Read Archive and can be accessed under the BioProject number PRJNA336221.

References

Adnan, M., Zheng, W., Islam, W., Arif, M., Abubakar, Y.S., Wang, Z., et al. (2017). Carbon
catabolite repression in filamentous fungi. Int. J. Mol. Sci. 19, 48. doi: 10.3390/ijms19010048

Alazi, E., and Ram, A.F.J. (2018). Modulating transcriptional regulation of plant biomass degrading
enzyme networks for rational design of industrial fungal strains. Front. Bioeng. Biotechnol. 6,
133. doi: 10.3389/fbioe.2018.00133

Alexa, A., and Rahnenfuhrer, J. (2021). topGO: Enrichment Analysis for Gene Ontology. R Package
Version 2.44.0.

Almeida, D.A., Horta, M.A.C., Ferreira Filho, J.A., Murad, N.F., and de Souza, A.P. (2021). The
synergistic actions of hydrolytic genes reveal the mechanism of Trichoderma harzianum for
cellulose degradation. J. Biotechnol. 334, 1-10. doi: 10.1016/j.jbiotec.2021.05.001

Almeida, D.A., Horta, M.A.C., Filho, J.A.F., Murad, N.F., and de Souza, A.P. (2020). The
synergistic actions of hydrolytic genes in coexpression networks reveal the potential of
Trichoderma harzianum for cellulose degradation. bioRxiv 2020.01.14.906529.

Antonieto, A.C., Castro, L.D.S., Silva-Rocha, R., Persinoti, G.F., and Silva, R.N. (2014). Defining
the genome-wide role of CRE1 during carbon catabolite repression in Trichoderma reesei
using RNA-Seq analysis. Fungal Genet. Biol. 73, 93-103. doi: 10.1016/j.fgb.2014.10.009

Arntzen, M.O., Bengtsson, O., Varnai, A., Delogu, F., Mathiesen, G., and Eijsink, V.G.H. (2020).
Quantitative comparison of the biomass-degrading enzyme repertoires of five filamentous
fungi. Sci. Rep. 10, 20267. doi: 10.1038/s41598-020-75217-z

Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., et al. (2000). Gene
ontology: tool for the unification of biology. The gene ontology consortium. Nat. Genet. 25,
25-29. doi: 10.1038/75556

Baroncelli, R., Piaggeschi, G., Fiorini, L., Bertolini, E., Zapparata, A., Pe, M.E., et al. (2015). Draft
whole-genome sequence of the biocontrol agent *Trichoderma harzianum* T6776. Genome
Announc. 3, e00647-15. doi: 10.1128/genomeA.00647-15

Benocci, T., Aguilar-Pontes, M.V., Zhou, M., Seiboth, B., and de Vries, R.P. (2017). Regulators of
plant biomass degradation in ascomycetous fungi. Biotechnol. Biofuels 10, 152. doi:
10.1186/s13068-017-0841-x

Bischof, R.H., Ramoni, J., and Seiboth, B. (2016). Cellulases and beyond: the first 70 years of the
enzyme producer Trichoderma reesei. Microb. Cell Fact. 15, 106. doi: 10.1186/s12934-016-
0507-6

Borin, G.P., Carazzolle, M.F., Dos Santos, R.A.C., Riano-Pachon, D.M., and Oliveira, J.V.C. (2018).
Gene Co-expression network reveals potential new genes related to sugarcane bagasse
degradation in Trichoderma reesei RUT-30. Front. Bioeng. Biotechnol. 6, 151. doi:
10.3389/fbioe.2018.00151

Burger, G., Gray, M.W., and Lang, B.F. (2003). Mitochondrial genomes: anything goes. Trends
Genet. 19, 709-716. doi: 10.1016/j.tig.2003.10.012

Castro, L.D.S., de Paula, R.G., Antonieto, A.C., Persinoti, G.F., Silva-Rocha, R., and Silva, R.N.
(2016). Understanding the role of the master regulator XYR1 in Trichoderma reesei by global
transcriptional analysis. Front. Microbiol. 7, 175. doi: 10.3389/fmicb.2016.00175

Chadha, S., Mehetre, S.T., Bansal, R., Kuo, A., Aerts, A., Grigoriev, I.V., et al. (2018). Genome-
wide analysis of cytochrome P450s of Trichoderma spp.: annotation and evolutionary
relationships. Fungal Biol. Biotechnol. 5, 12. doi: 10.1186/s40694-018-0056-3

Chen, L., Zou, G., Wang, J., Wang, J., Liu, R., Jiang, Y., et al. (2016). Characterization of the Ca(2+)
-responsive signaling pathway in regulating the expression and secretion of cellulases in
Trichoderma reesei Rut-C30. Mol. Microbiol. 100, 560-575. doi: 10.1111/mmi.13334

Cock, P.J., Antao, T., Chang, J.T., Chapman, B.A., Cox, C.J., Dalke, A., et al. (2009). Biopython:
freely available Python tools for computational molecular biology and bioinformatics.
Bioinformatics 25, 1422-1423. doi: 10.1093/bioinformatics/btp163

Cortijo, S., Bhattarai, M., Locke, J. C. W., & Ahnert, S. E. (2020). Co-expression Networks From
Gene Expression Variability Between Genetically Identical Seedlings Can Reveal Novel
Regulatory Relationships. Frontiers in Plant Science, 11. doi:10.3389/fpls.2020.599464

Cziferszky, A., Mach, R.L., and Kubicek, C.P. (2002). Phosphorylation positively regulates DNA
binding of the carbon catabolite repressor Cre1 of Hypocrea jecorina (Trichoderma reesei). J.
Biol. Chem. 277, 14688-14694. doi: 10.1074/jbc.M200744200

de Assis, L.J., Ries, L.N., Savoldi, M., Dos Reis, T.F., Brown, N.A., and Goldman, G.H. (2015).
Aspergillus nidulans protein kinase A plays an important role in cellulase production.
Biotechnol. Biofuels 8, 213. doi: 10.1186/s13068-015-0401-1

de Assis, L.J., Silva, L.P., Bayram, O., Dowling, P., Kniemeyer, O., Kruger, T., et al. (2021). Carbon
catabolite repression in filamentous fungi is regulated by phosphorylation of the transcription
factor CreA. mBio 12, e03146-20. doi: 10.1128/mBio.03146-20

de Vries, R.P., Riley, R., Wiebenga, A., Aguilar-Osorio, G., Amillis, S., Uchima, C.A., et al. (2017).
Comparative genomics reveals high biological diversity and specific adaptations in the
industrially and medically important fungal genus Aspergillus. Genome Biol. 18, 28. doi:
10.1186/s13059-017-1151-0

Delabona, P.D.S., Codima, C.A., Ramoni, J., Zubieta, M.P., de Araujo, B.M., Farinas, C.S., et al.
(2020). The impact of putative methyltransferase overexpression on the *Trichoderma
harzianum* cellulolytic system for biomass conversion. Bioresour. Technol. 313, 123616. doi:
10.1016/j.biortech.2020.123616

Delabona, P.D.S., Lima, D.J., Codima, C.A., Ramoni, J., Gelain, L., de Melo, V.S., et al. (2021).
Replacement of the carbon catabolite regulator (cre1) and fed-batch cultivation as strategies to
enhance cellulase production in Trichoderma harzianum. Bioresour. Technol. Rep. 13,
100634. doi: 10.1016/j.biteb.2021.100634

Delabona, P.D.S., Rodrigues, G.N., Zubieta, M.P., Ramoni, J., Codima, C.A., Lima, D.J., et al.
(2017). The relation between xyr1 overexpression in *Trichoderma harzianum* and sugarcane
bagasse saccharification performance. J. Biotechnol. 246, 24-32. doi:
10.1016/j.jbiotec.2017.02.002

Dodd, S.L., Lieckfeldt, E., and Samuels, G.J. (2003). Hypocrea atroviridis sp. nov., the teleomorph of
Trichoderma atroviride. Mycologia 95, 27-40. doi: 10.1080/15572536.2004.11833129

Druzhinina, I.S., Chenthamara, K., Zhang, J., Atanasova, L., Yang, D., Miao, Y., et al. (2018).
Massive lateral transfer of genes encoding plant cell wall-degrading enzymes to the
mycoparasitic fungus Trichoderma from its plant-associated hosts. PLoS Genet. 14,
e1007322. doi: 10.1371/journal.pgen.1007322

Druzhinina, I.S., Kopchinskiy, A.G., and Kubicek, C.P. (2006). The first 100 Trichoderma species
characterized by molecular data. Mycoscience 47, 55-64. doi: 10.1007/s10267-006-0279-7

Druzhinina, I.S., Kubicek, C.P., Komon-Zelazowska, M., Mulaw, T.B., and Bissett, J. (2010). The
Trichoderma harzianum demon: complex speciation history resulting in coexistence of
hypothetical biological species, recent agamospecies and numerous relict lineages. BMC
Evol. Biol. 10, 94. doi: 10.1186/1471-2148-10-94

Fan, J., Urban, M., Parker, J.E., Brewer, H.C., Kelly, S.L., Hammond-Kosack, K.E., et al. (2013).
Characterization of the sterol 14alpha-demethylases of Fusarium graminearum identifies a
novel genus-specific CYP51 function. New Phytol. 198, 821-835. doi: 10.1111/nph.12193

Felsenstein, J. (1985). Confidence limits on phylogenies: an approach using the bootstrap. Evolution
39, 783-791. doi: 10.2307/2408678

Fonseca, L. M., Parreiras, L. S., & Murakami, M. T. (2020). Rational engineering of the Trichoderma
reesei RUT-C30 strain into an industrially relevant platform for cellulase production.
Biotechnology for Biofuels, 13(1), 93. doi:10.1186/s13068-020-01732-w

Grevel, A., and Becker, T. (2020). Porins as helpers in mitochondrial protein translocation. Biol.
Chem. 401, 699-708. doi: 10.1515/hsz-2019-0438

Gruber, S., and Seidl-Seiboth, V. (2012). Self versus non-self: fungal cell wall degradation in
Trichoderma. Microbiology (Reading) 158, 26-34. doi: 10.1099/mic.0.052613-0

Gruber, S., and Zeilinger, S. (2014). The transcription factor Ste12 mediates the regulatory role of the
Tmk1 MAP kinase in mycoparasitism and vegetative hyphal fusion in the filamentous fungus
Trichoderma atroviride. PLoS One 9, e111636. doi: 10.1371/journal.pone.0111636

Hahn, T., Tafi, E., Paul, A., Salvia, R., Falabella, P., and Zibek, S. (2020). Current state of chitin
purification and chitosan production from insects. J. Chem. Technol. Biotechnol. 95, 2775-
2795. doi: 10.1002/jctb.6533

Han, L., Tan, Y., Ma, W., Niu, K., Hou, S., Guo, W., et al. (2020). Precision engineering of the
transcription factor Cre1 in Hypocrea jecorina (Trichoderma reesei) for efficient cellulase

production in the presence of glucose. Front. Bioeng. Biotechnol. 8, 852. doi:
10.3389/fbioe.2020.00852

Harris, S.D., Hamer, L., Sharpless, K.E., and Hamer, J.E. (1997). The Aspergillus nidulans sepA
gene encodes an FH1/2 protein involved in cytokinesis and the maintenance of cellular
polarity. EMBO J. 16, 3474-3483. doi: 10.1093/emboj/16.12.3474

Hinterdobler, W., Li, G., Turrà, D., Schalamun, M., Kindel, S., Sauer, U., et al. (2021). Integration of
chemosensing and carbon catabolite repression impacts fungal enzyme regulation and plant
associations. bioRxiv 2021.05.06.442915.

Hoi, J.W.S., and Dumas, B. (2010). Ste12 and Ste12-like proteins, fungal transcription factors
regulating development and pathogenicity. Eukaryot. Cell 9, 480-485. doi:
10.1128/EC.00333-09

Horta, M.A.C., Filho, J.A.F., Murad, N.F., Santos, E.D.O., Dos Santos, C.A., Mendes, J.S., et al.
(2018). Network of proteins, enzymes and genes linked to biomass degradation shared by
Trichoderma species. Sci. Rep. 8, 1341. doi: 10.1038/s41598-018-19671-w

Horta, M.A.C., Thieme, N., Gao, Y., Burnum-Johnson, K.E., Nicora, C.D., Gritsenko, M.A., et al.
(2019). Broad Substrate-Specific Phosphorylation Events Are Associated With the Initial
Stage of Plant Cell Wall Recognition in Neurospora crassa. Front. Microbiol. 10, 2317. doi:
10.3389/fmicb.2019.02317

Jankowsky, E. (2011). RNA helicases at work: binding and rearranging. Trends Biochem. Sci. 36,
19-29. doi: 10.1016/j.tibs.2010.07.008

Jones, D.T., Taylor, W.R., and Thornton, J.M. (1992). The rapid generation of mutation data matrices
from protein sequences. Comput. Appl. Biosci. 8, 275-282. doi:
10.1093/bioinformatics/8.3.275

Kameshwar, A.K.S., Ramos, L.P., and Qin, W. (2019). CAZymes-based ranking of fungi (CBRF): an
interactive web database for identifying fungi with extrinsic plant biomass degrading abilities.
Bioresour. Bioprocess. 6, 51. doi: 10.1186/s40643-019-0286-0

Kanehisa, M., and Goto, S. (2000). KEGG: kyoto encyclopedia of genes and genomes. Nucleic Acids
Res. 28, 27-30. doi: 10.1093/nar/28.1.27

Kimura, M. (1980). A simple method for estimating evolutionary rates of base substitutions through
comparative studies of nucleotide sequences. J. Mol. Evol. 16, 111-120. doi:
10.1007/BF01731581

Klaubauf, S., Narang, H.M., Post, H., Zhou, M., Brunner, K., Mach-Aigner, A.R., et al. (2014).
Similar is not the same: differences in the function of the (hemi-)cellulolytic regulator XlnR
(Xlr1/Xyr1) in filamentous fungi. Fungal Genet. Biol. 72, 73-81. doi:
10.1016/j.fgb.2014.07.007

Kolde, R. (2019). Pheatmap: Pretty Heatmaps. R Package Version 1.0.12.

Koutrouli, M., Karatzas, E., Paez-Espino, D., and Pavlopoulos, G.A. (2020). A guide to conquer the
biological network era using graph theory. Front. Bioeng. Biotechnol. 8, 34. doi:
10.3389/fbioe.2020.00034

Kouzarides, T. (2007). Chromatin modifications and their function. Cell 128, 693-705. doi:
10.1016/j.cell.2007.02.005

Kubicek, C.P., Herrera-Estrella, A., Seidl-Seiboth, V., Martinez, D.A., Druzhinina, I.S., Thon, M., et
al. (2011). Comparative genome sequence analysis underscores mycoparasitism as the
ancestral life style of *Trichoderma*. Genome Biol. 12, R40. doi: 10.1186/gb-2011-12-4-r40

Kubicek, C.P., Steindorff, A.S., Chenthamara, K., Manganiello, G., Henrissat, B., Zhang, J., et al.
(2019). Evolution and comparative genomics of the most common Trichoderma species.
BMC Genom. 20, 485. doi: 10.1186/s12864-019-5680-7

Kumar, S., Stecher, G., and Tamura, K. (2016). MEGA7: molecular evolutionary genetics analysis
version 7.0 for bigger datasets. Mol. Biol. Evol. 33, 1870-1874. doi: 10.1093/molbev/msw054

Langfelder, P., and Horvath, S. (2008). WGCNA: an R package for weighted correlation network
analysis. BMC Bioinform. 9, 559. doi: 10.1186/1471-2105-9-559

Langfelder, P., Zhang, B., and Horvath, S. (2008). Defining clusters from a hierarchical cluster tree:
the Dynamic Tree Cut package for R. Bioinformatics 24, 719-720. doi:
10.1093/bioinformatics/btm563

Li, C.X., Zhao, S., Luo, X.M., and Feng, J.X. (2020a). Weighted gene co-expression network
analysis identifies critical genes for the production of cellulase and xylanase in penicillium
oxalicum. Front. Microbiol. 11, 520. doi: 10.3389/fmicb.2020.00520

Li, J.X., Zhang, F., Jiang, D.D., Li, J., Wang, F.L., Zhang, Z., et al. (2020b). Diversity of cellulase-
producing filamentous fungi from tibet and transcriptomic analysis of a superior cellulase
producer Trichoderma harzianum LZ117. Front. Microbiol. 11, 1617. doi:
10.3389/fmicb.2020.01617

Lichius, A., Seidl-Seiboth, V., Seiboth, B., and Kubicek, C.P. (2014). Nucleo-cytoplasmic shuttling
dynamics of the transcriptional regulators XYR1 and CRE1 under conditions of cellulase and
xylanase gene expression in Trichoderma reesei. Mol. Microbiol. 94, 1162-1178. doi:
10.1111/mmi.12824

Liu, J., Liu, S., & Yang, X. (2021). Construction of Gene Modules and Analysis of Prognostic
Biomarkers for Cervical Cancer by Weighted Gene Co-Expression Network Analysis.
Frontiers in Oncology, 11. doi:10.3389/fonc.2021.542063

Liu, R., Chen, L., Jiang, Y., Zou, G., & Zhou, Z. (2017). A novel transcription factor specifically
regulates GH11 xylanase genes in Trichoderma reesei. Biotechnology for Biofuels, 10(1),
194. doi:10.1186/s13068-017-0878-x

Lombard, V., Ramulu, H.G., Drula, E., Coutinho, P.M., and Henrissat, B. (2014). The carbohydrate-
active enzymes database (CAZy) in 2013. Nucleic Acids Res. 42, D490-D495. doi:
10.1093/nar/gkt1178

Luscombe, N.M., Babu, M.M., Yu, H., Snyder, M., Teichmann, S.A., Gerstein, M. (2004). Genomic
analysis of regulatory network dynamics reveals large topological changes. Nature. Sep
16;431(7006):308-12. doi: 10.1038/nature02782. PMID: 15372033

Mach-Aigner, A.R., Pucher, M.E., Steiger, M.G., Bauer, G.E., Preis, S.J., and Mach, R.L. (2008).
Transcriptional regulation of xyr1, encoding the main regulator of the xylanolytic and
cellulolytic enzyme system in *Hypocrea jecorina*. Appl. Environ. Microbiol. 74, 6554-6562.
doi: 10.1128/AEM.01143-08

Maier, B.F. (2019). Generalization of the small-world effect on a model approaching the Erdos-Renyi
random graph. Sci. Rep. 9, 9268. doi: 10.1038/s41598-019-45576-3

Malmierca, M.G., McCormick, S.P., Cardoza, R.E., Alexander, N.J., Monte, E., and Gutierrez, S.
(2015). Production of trichodiene by Trichoderma harzianum alters the perception of this
biocontrol strain by plants and antagonized fungi. Environ. Microbiol. 17, 2628-2646. doi:
10.1111/1462-2920.12506

Martinez, D., Berka, R.M., Henrissat, B., Saloheimo, M., Arvas, M., Baker, S.E., et al. (2008).
Genome sequencing and analysis of the biomass-degrading fungus Trichoderma reesei (syn.
Hypocrea jecorina). Nat. Biotechnol. 26, 553-560. doi: 10.1038/nbt1403

Martins-Santana, L., Paula, R.G., Silva, A.G., Lopes, D.C.B., Silva, R.D.N., and Silva-Rocha, R.
(2020). CRZ1 regulator and calcium cooperatively modulate holocellulases gene expression
in *Trichoderma reesei* QM6a. Genet. Mol. Biol. 43, e20190244. doi: 10.1590/1678-4685-
GMB-2019-0244

Maruyama, C.R., Bilesky-Jose, N., de Lima, R., and Fraceto, L.F. (2020). Encapsulation of
Trichoderma harzianum preserves enzymatic activity and enhances the potential for biological
control. Front. Bioeng. Biotechnol. 8, 225. doi: 10.3389/fbioe.2020.00225

Medeiros, H.A., Filho, J.V.A., Freitas, L.G., Castillo, P., Rubio, M.B., Hermosa, R., et al. (2017).
Tomato progeny inherit resistance to the nematode *Meloidogyne javanica* linked to plant
growth induced by the biocontrol fungus *Trichoderma atroviride*. Sci. Rep. 7, 40216. doi:
10.1038/srep40216

Mello-de-Sousa, T.M., Gorsche, R., Rassinger, A., Pocas-Fonseca, M.J., Mach, R.L., and Mach-
Aigner, A.R. (2014). A truncated form of the Carbon catabolite repressor 1 increases cellulase
production in *Trichoderma reesei*. Biotechnol. Biofuels 7, 129. doi: 10.1186/s13068-014-
0129-3

Monroy, A.A., Stappler, E., Schuster, A., Sulyok, M., and Schmoll, M. (2017). A CRE1- regulated
cluster is responsible for light dependent production of dihydrotrichotetronin in Trichoderma
reesei. PLoS One 12, e0182530. doi: 10.1371/journal.pone.0182530

Mukherjee, P.K., Nautiyal, C.S., and Mukhopadhyay, A.N. (2008). "Molecular mechanisms of
biocontrol by Trichoderma spp," in Molecular Mechanisms of Plant and Microbe

Coexistence, eds. C.S. Nautiyal & P. Dion (Berlin, Heidelberg: Springer Berlin Heidelberg),
243-262.

Mutwil, M., Usadel, B., Schutte, M., Loraine, A., Ebenhoh, O., and Persson, S. (2010). Assembly of
an interactive correlation network for the Arabidopsis genome using a novel heuristic
clustering algorithm. Plant Physiol. 152, 29-43. doi: 10.1104/pp.109.145318

Nasa, I., and Kettenbach, A.N. (2018). Coordination of protein kinase and phosphoprotein
phosphatase activities in mitosis. Front. Cell Dev. Biol. 6, 30. doi: 10.3389/fcell.2018.00030

Neupane, P., Bhuju, S., Thapa, N., and Bhattarai, H.K. (2019). ATP synthase: structure, function and
inhibition. Biomol. Concepts 10, 1-10. doi: 10.1515/bmc-2019-0001

Pavlopoulos, G.A., Secrier, M., Moschopoulos, C.N., Soldatos, T.G., Kossida, S., Aerts, J., et al.
(2011). Using graph theory to analyze biological networks. BioData Min. 4, 10. doi:
10.1186/1756-0381-4-10

Phillips, B.P., Gomez-Navarro, N., and Miller, E.A. (2020). Protein quality control in the
endoplasmic reticulum. Curr. Opin. Cell Biol. 65, 96-102. doi: 10.1016/j.ceb.2020.04.002

Portnoy, T., Margeot, A., Linke, R., Atanasova, L., Fekete, E., Sandor, E., et al. (2011). The CRE1
carbon catabolite repressor of the fungus Trichoderma reesei: a master regulator of carbon
assimilation. BMC Genom. 12, 269. doi: 10.1186/1471-2164-12-269

R Core Team (2018). R: A Language and Environment for Statistical Computing. Vienna, Austria: R
Foundation for Statistical Computing.

Reithner, B., Mach-Aigner, A.R., Herrera-Estrella, A., and Mach, R.L. (2014). *Trichoderma
atroviride* transcriptional regulator Xyr1 supports the induction of systemic resistance in
plants. Appl. Environ. Microbiol. 80, 5274-5281. doi: 10.1128/AEM.00930-14

Ries, L., Belshaw, N.J., Ilmen, M., Penttila, M.E., Alapuranen, M., and Archer, D.B. (2014). The role
of CRE1 in nucleosome positioning within the cbh1 promoter and coding regions of
Trichoderma reesei. Appl. Microbiol. Biotechnol. 98, 749-762. doi: 10.1007/s00253-013-
5354-3

Sanner, M.F. (1999). Python: a programming language for software integration and development. J.
Mol. Graph. Model. 17, 57-61. doi: 10.1186/1471-2105-4-2

Saravanakumar, K., Li, Y., Yu, C., Wang, Q.Q., Wang, M., Sun, J., et al. (2017). Effect of
Trichoderma harzianum on maize rhizosphere microbiome and biocontrol of Fusarium Stalk
rot. Sci. Rep. 7, 1771. doi: 10.1038/s41598-017-01680-w

Scardoni, G., Tosadori, G., Pratap, S., Spoto, F., and Laudanna, C. (2016). Finding the shortest path
with PesCa: a tool for network reconstruction. F1000Research 4, 484. doi:
10.12688/f1000research.6769.2

Schmoll, M. (2018). Light, stress, sex and carbon - The photoreceptor ENVOY as a central
checkpoint in the physiology of Trichoderma reesei. Fungal Biol. 122, 479-486. doi:
10.1016/j.funbio.2017.10.007

Schmoll, M., Dattenbock, C., Carreras-Villasenor, N., Mendoza-Mendoza, A., Tisch, D., Aleman,
M.I., et al. (2016). The genomes of three uneven siblings: footprints of the lifestyles of three
trichoderma species. Microbiol. Mol. Biol. Rev. 80, 205-327. doi: 10.1128/MMBR.00040-15

Schoch, C.L., Seifert, K.A., Huhndorf, S., Robert, V., Spouge, J.L., Levesque, C.A., et al. (2012).
Nuclear ribosomal internal transcribed spacer (ITS) region as a universal DNA barcode
marker for Fungi. Proc. Natl. Acad. Sci. U. S. A. 109, 6241-6246. doi:
10.1073/pnas.1117018109

Shannon, P., Markiel, A., Ozier, O., Baliga, N.S., Wang, J.T., Ramage, D., et al. (2003). Cytoscape: a
software environment for integrated models of biomolecular interaction networks. Genome
Res. 13, 2498-2504. doi: 10.1101/gr.1239303

Siewers, V., Viaud, M., Jimenez-Teja, D., Collado, I.G., Gronover, C.S., Pradier, J.M., et al. (2005).
Functional analysis of the cytochrome P450 monooxygenase gene bcbot1 of Botrytis cinerea
indicates that botrydial is a strain-specific virulence factor. Mol. Plant-Microbe Interact. 18,
602-612. doi: 10.1094/MPMI-18-0602

Sloothaak, J., Tamayo-Ramos, J.A., Odoni, D.I., Laothanachareon, T., Derntl, C., Mach-Aigner,
A.R., et al. (2016). Identification and functional characterization of novel xylose transporters
from the cell factories Aspergillus niger and Trichoderma reesei. Biotechnol. Biofuels 9, 148.
doi: 10.1186/s13068-016-0564-4

Strauss, J., Mach, R.L., Zeilinger, S., Hartler, G., Stöffler, G., Wolschek, M., et al. (1995). Crel, the
carbon catabolite repressor protein fromTrichoderma reesei. FEBS Lett. 376, 103-107. doi:
10.1016/0014-5793(95)01255-5

Stricker, A.R., Grosstessner-Hain, K., Wurleitner, E., and Mach, R.L. (2006). Xyr1 (xylanase
regulator 1) regulates both the hydrolytic enzyme system and D-xylose metabolism in
Hypocrea jecorina. Eukaryot. Cell 5, 2128-2137. doi: 10.1128/EC.00211-06

Supek, F., Bosnjak, M., Skunca, N., and Smuc, T. (2011). REVIGO summarizes and visualizes long
lists of gene ontology terms. PLoS One 6, e21800. doi: 10.1371/journal.pone.0021800

Thanh, V.N., Thuy, N.T., Huong, H.T.T., Hien, D.D., Hang, D.T.M., Anh, D.T.K., et al. (2019).
Surveying of acid-tolerant thermophilic lignocellulolytic fungi in Vietnam reveals
surprisingly high genetic diversity. Sci. Rep. 9, 3674. doi: 10.1038/s41598-019-40213-5

Thompson, J.D., Higgins, D.G., and Gibson, T.J. (1994). CLUSTAL W: improving the sensitivity of
progressive multiple sequence alignment through sequence weighting, position-specific gap
penalties and weight matrix choice. Nucleic Acids Res. 22, 4673-4680. doi:
10.1093/nar/22.22.4673

Tolgo, M., Huttner, S., Rugbjerg, P., Thuy, N.T., Thanh, V.N., Larsbrink, J., et al. (2021). Genomic
and transcriptomic analysis of the thermophilic lignocellulose-degrading fungus Thielavia
terrestris LPH172. Biotechnol. Biofuels 14, 131. doi: 10.1186/s13068-021-01975-1

UniProt Consortium. (2019). UniProt: a worldwide hub of protein knowledge. Nucleic Acids Res. 47,
D506-D515. doi: 10.1093/nar/gky1049

Vallim, M.A., Miller, K.Y., and Miller, B.L. (2000). Aspergillus SteA (sterile12-like) is a
homeodomain-C2/H2-Zn+2 finger transcription factor required for sexual reproduction. Mol.
Microbiol. 36, 290-301. doi: 10.1046/j.1365-2958.2000.01874.x

Vieira, A. A., Vianna, G. R., Carrijo, J., Aragão, F. J. L., & Vieira, P. M. (2021). Generation of
Trichoderma harzianum with pyr4 auxotrophic marker by using the CRISPR/Cas9 system.
Scientific Reports, 11(1), 1085. doi:10.1038/s41598-020-80186-4

Vieira, P. M., Zeilinger, S., Brandão, R. S., Vianna, G. R., Georg, R. C., Gruber, S., . . . Ulhoa, C. J.
(2018). Overexpression of an aquaglyceroporin gene in Trichoderma harzianum affects stress
tolerance, pathogen antagonism and Phaseolus vulgaris development. Biological Control, 126,
185-191. doi:https://doi.org/10.1016/j.biocontrol.2018.08.012

Whitehouse, D.G., May, B., and Moore, A.L. (2019). "Respiratory chain and ATP synthase," in
Reference Module in Biomedical Sciences (Elsevier). doi: 10.1016/B978-0-12-801238-
3.95732-5

Wolfe, C.J., Kohane, I.S., and Butte, A.J. (2005). Systematic survey reveals general applicability of
"guilt-by-association" within gene coexpression networks. BMC Bioinform. 6, 227. doi:
10.1186/1471-2105-6-227

Yan, L., and Khan, R.A.A. (2021). Biological control of bacterial wilt in tomato through the
metabolites produced by the biocontrol fungus, Trichoderma harzianum. Egypt. J. Biol. Pest
Control 31, 5. doi: 10.1186/s41938-020-00351-9

Zeilinger, S., and Atanasova, L. (2020). "Chapter 2 - Sensing and regulation of mycoparasitism-
relevant processes in Trichoderma," in New and Future Developments in Microbial
Biotechnology and Bioengineering, eds. V.K. Gupta, S. Zeilinger, H.B. Singh & I. Druzhinina
(Amsterdam: Elsevier), 39-55.

Zeilinger, S., Schmoll, M., Pail, M., Mach, R.L., and Kubicek, C.P. (2003). Nucleosome transactions
on the Hypocrea jecorina (Trichoderma reesei) cellulase promoter cbh2 associated with
cellulase induction. Mol. Genet. Genom. 270, 46-55. doi: 10.1007/s00438-003-0895-2

Zeilinger, S. (2004). Gene disruption in Trichoderma atroviride via Agrobacterium-mediated
transformation. Current Genetics, 45(1), 54-60. doi:10.1007/s00294-003-0454-8

Zhang, W., Kou, Y., Xu, J., Cao, Y., Zhao, G., Shao, J., et al. (2013). Two major facilitator
superfamily sugar transporters from Trichoderma reesei and their roles in induction of
cellulase biosynthesis. J. Biol. Chem. 288, 32861-32872. doi: 10.1074/jbc.M113.505826

Zhang, Y., Yang, J., Luo, L., Wang, E., Wang, R., Liu, L., et al. (2020). Low-cost cellulase-
hemicellulase mixture secreted by Trichoderma harzianum EM0925 with complete
saccharification efficacy of lignocellulose. Int. J. Mol. Sci. 21, 371. doi:
10.3390/ijms21020371

Zhang, Z.-Q., Wu, W.-W., Chen, J.-D., Zhang, G.-Y., Lin, J.-Y., Wu, Y.-K., . . . Si, T.-M. (2021).
Weighted Gene Coexpression Network Analysis Reveals Essential Genes and Pathways in
Bipolar Disorder. Frontiers in Psychiatry, 12. doi:10.3389/fpsyt.2021.553305

1105

1106

1107

1108

1109

1110

1111

1112

1113

1114

1115

1116

1117

1118

1119

1120

1121

1122

1123

1124

1125

1126

1127

1128

Figure legends

Figure 1. Molecular phylogenies of CRE1 and XYR1 in *Trichoderma* spp. The 26 and 21
complete protein sequences related to CRE1 (**A**) and XYR1 (**B**), respectively, were used to infer the
phylogenetic relationships of the studied strains in the genus *Trichoderma*. Close genetic proximity
between Th3844 and Th0179 was observed in the CRE1 phylogenetic tree, while Ta0020 shared a
closer affinity with other *T. atroviride* strains. In the XYR1 phylogenetic tree, Th3844, Th0179, and
Ta0020 were closest to *T. harzianum* CBS 226.95, *T. harzianum* T6776, and *T. atroviride*
IMI206040, respectively. *Fusarium* spp. was used as an outgroup. The bootstrap value is represented
in each node. The phylogenetic trees were constructed using MEGA7 software and edited using the
iTOL program.

Figure 2. KO functional classification of the transcripts identified in the *cre1* and *xyr1* groups.
The sequences related to the enzymes coexpressed with *cre1* (**A**) and *xyr1* (**B**) were annotated
according to the main KO functions of Th3844, Th0179, and Ta0020. For all evaluated strains, 14
pathway classes were observed in each TF, and 13 classes were shared between the two regulators.
However, the number of identified enzymes in a given pathway class differed in each strain. Each
strain is represented by a different color. Ta0020: *T. atroviride* CBMAI-0020; Th0179: *T. harzianum*
CBMAI-0179; Th3844: *T. harzianum* IOC-3844; KO: Kyoto Encyclopedia of Genes and Genomes
Orthology.

Figure 3. Comparisons of TFs, transporters, and CAZymes among strains and groups. Number
of transcripts encoding TFs, transporters, and CAZymes in Ta0020, Th0179, and Th3844 distributed
among the *cre1* (**A**) and *xyr1* (**B**) groups. The number of transcripts encoding these proteins
associated with *cre1* was more accentuated in Ta0020 than those observed for the other strains, while
in the *xyr1* group, both *T. harzianum* strains presented a great number of coexpressed transcripts
encoding TFs, followed by those encoding transporters. Th3844 also showed a high number of
CAZymes in the *xyr1* group. TFs: transcription factors; CAZymes: carbohydrate-active enzymes.
Ta0020: *T. atroviride* CBMAI-0020; Th0179: *T. harzianum* CBMAI-0179; Th3844: *T. harzianum*
IOC-3844.

Figure 4. Distribution of CAZyme families in the *cre1* and *xyr1* groups. Classification of
CAZyme families coexpressed with *cre1* (**A**) and *xyr1* (**B**) in Th3844, Th0179, and Ta0020 and
quantification of each CAZyme family in the *cre1* groups (**C**) and *xyr1* groups (**D**). Illustration
showing that the *xyr1* and *cre1* groups presented CAZymes from the GH family in all the evaluated
strains with different numbers of transcripts. Overall, Ta0020 had a high number of GHs belonging
to family 18, coexpressed with the *cre1* transcript, while Th0179 exhibited a significant amount of
GT90. Different CAZyme profiles were observed in the *xyr1* groups, and Th3844 showed the highest
diversity of CAZyme families. PL: polysaccharide lyase; GH: glycoside hydrolase; GT:
glycosyltransferase; CE: carbohydrate esterase; Ta0020: *T. atroviride* CBMAI-0020; Th0179: *T.*
harzianum CBMAI-0179; Th3844: *T. harzianum* IOC-3844.

Figure 5. Distribution of TFs and transporters in the *cre1* and *xyr1* groups in *Trichoderma* spp.
Classification of TFs coexpressed with *cre1* (**A**) and *xyr1* (**B**) and transporters coexpressed with *cre1*
(**C**) and *xyr1* (**D**) in Th3844, Th0179, and Ta0020. The TFs and the transporters were classified
according to their class and quantity. MFS: major facilitator superfamily; ABC: ATP-binding
cassette; Th3844: *T. harzianum* IOC-3844; Th0179: *T. harzianum* CBMAI-0179; Ta0020: *T.*
atroviride CBMAI-0020.

Figure 6. Modeled global networks and modeled subnetworks of the *cre1* and *xyr1* groups of
the evaluated strains. The transcriptome datasets were used to infer the global HRR networks of (**A**)
Th3844, (**B**) Th0179, and (**C**) Ta0020. The networks were modeled and edited using Cytoscape
software. The global networks were partitioned, and subnetworks of Th3844 (**D**), Th0179 (**E**) and
Ta0020 (**F**) related to *cre1* were constructed, as well as Th3844 (**G**), Th0179 (**H**) and Ta0020 (**I**)
related to *xyr1*. Each color represents a category of transcripts identified in the global networks and
subnetworks. Th3844: *T. harzianum* IOC-3844; Th0179: *T. harzianum* CBMAI-0179; Ta0020: *T.*
atroviride CBMAI-0020; TF: transcription factor.

Figure 7. Heatmap plotted based on the shortest pathway between XYR1 or CRE1 and
CAZymes in all evaluated *Trichoderma* spp. Number of shortest pathways between XYR1 or
CRE1 and the selected CAZymes (**A**) and number of occurrences of such an event (**B**), as indicated
by the accentuation of the purple color in the subtitle of the figures. Th3844: *T. harzianum* IOC-
3844; Th0179: *T. harzianum* CBMAI-0179; Ta0020: *T. atroviride* CBMAI-0020.

1185

1186

1187

1188

1189

1190

1191

1192

1193

1194

1195

1196

1197

1198

1199

1200

1201

1202 **Table**

1203 **Table 1.** First neighbors of the transcripts *xyr1* and *cre1* identified in Th3844, Th0179, and Ta0020.

First neighbors						
Th3844						
CRE1			**XYR1**			
Gene ID	**Description**	**Group**	**Gene ID**	**Description**	**Group**
THAR02_06363	Zn2Cys6 transcriptional regulator	29	THAR02_07076	regulator-nonsense transcripts 1	9
THAR02_10232	phosphate:H+ symporter	13	THAR02_00049	polysaccharide lyase family 7	11
THAR02_07487	Mg2+ transporter	78	THAR02_09951	MFS permease	70
THAR02_07586	cytochrome P450 CYP2 subfamily	39	THAR02_01555	MFS permease	0
THAR02_07548	C2H2 transcriptional regulator	13	THAR02_00890	glycoside hydrolase family 3	0
THAR02_04042	acid phosphatase	8	THAR02_09118	fungal specific transcription factor	26
Th0179						
THAR02_03655	proteasome subunit alpha type-5	70	THAR02_00084	MFS transporter	63
THAR02_06292	restriction of telomere capping 5	53	THAR02_02333	MFS transporter	30
THAR02_11314	xaa-Pro aminopeptidase	46	THAR02_11077	glycosyltransferase family 2	35
THAR02_03482	kinase domain	32	THAR02_00098	glycoside hydrolase family 43	67
THAR02_03668	adenine phosphoribosyltransferase	31	THAR02_03812	G-coupled receptor	39
THAR02_00283	NADH dehydrogenase	8	THAR02_04897	C2H2 transcriptional regulator	19
Ta0020						
013948456.1_10935	SH3 domain-containing	52	013943820.1_6616	Zn2Cys6 transcriptional regulator	14
013938015.1_374	serine threonine kinase	7	013943207.1_5652	replication factor A1	17

013937789.1_299	ribosomal S18	35	013941180.1_3604	MFS transporter	1
013949023.1_11526	pleiotropic drug resistance	2	013943062.1_5470	MFS transporter	90
013946760.1_9261	MYB DNA-binding domain-containing	13	013941580.1_4080	glycoside hydrolase family 76	1
013941398.1_3866	aspartate aminotransferase	28	013945761.1_8223	DNA ligase 1	16

1204

Global networks

A — Th3844
B — Th0179
C — Ta0020

- Downregulated
- XYR1 group
- XYR1 + Downregulated
- XYR1 + Upregulated
- CRE1 + Downregulated
- CRE1 + Upregulated
- CRE1 group
- Upregulated

Subnetworks

D — Th3844
E — Th0179
F — Ta0020

G
H
I

CRE1

XYR1

- Downregulated
- Upregulated
- Up + CAZyme
- Up + TF
- Up + Transporter
- XYR1
- CAZyme
- CRE1
- TF
- Transporter

CRE1

A

CRE1

C

Strain	GH2	GH5	GH12	GH16	GH18	GH23	GH28	GH39	GH43	GH47	GH55
Ta0020	1	1	1	1	4	0	2	1	1	0	2
Th0179	0	0	0	0	0	1	0	0	0	1	0
Th3844	0	0	0	0	0	0	1	0	0	0	0

Strain	GH64	GH76	GH81	GH92	GH93	GH95	GT90	CE1	CE3	CE5
Ta0020	1	1	1	2	2	1	0	0	1	1
Th0179	0	0	0	0	0	0	3	1	0	0
Th3844	0	0	0	0	0	0	0	0	0	0

XYR1

B

XYR1

D

Strain	GH16	GH17	GH18	GH30	GH39	GH75	GH76	GT-	GT22	PL8
Ta0020	0	0	0	0	0	1	0	0	1	0
Th0179	1	0	0	0	0	1	1	0	0	0
Th3844	0	1	1	1	1	1	0	1	0	1