Predicting amphibian intraspecific diversity with machine learning: Challenges and

prospects for integrating traits, geography, and genetic data

Lisa N. Barrow[1,2*] (https://orcid.org/0000-0001-7081-2432)

Emanuel Masiero da Fonseca[1] (https://orcid.org/0000-0002-2952-8816)

Coleen E. P. Thompson[1] (https://orcid.org/0000-0003-0591-7654)

Bryan C. Carstens[1] (https://orcid.org/0000-0002-1552-227X)

[1]Department of Evolution, Ecology and Organismal Biology, The Ohio State University, 318 W.

12[th] Ave, Columbus, OH 43210

[2]Museum of Southwestern Biology and Department of Biology, University of New Mexico,

Albuquerque NM 87131

[*]Corresponding author: lnbarrow@unm.edu

Running Title: AMPHIBIAN BIOGEOGRAPHY AND RANDOM FORESTS

Abstract

The growing availability of genetic datasets, in combination with machine learning frameworks, offer great potential to answer long-standing questions in ecology and evolution. One such question has intrigued population geneticists, biogeographers, and conservation biologists: What determines intraspecific genetic diversity? This question is challenging to answer because many factors may influence genetic variation, including life history traits, historical influences, and geography, and the relative importance of these factors varies across taxonomic and geographic scales. Furthermore, interpreting the influence of numerous, potentially correlated variables is difficult with traditional statistical approaches. To address these challenges, we combined repurposed data with machine learning and investigated predictors of genetic diversity, focusing on Nearctic amphibians as a case study. We aggregated species traits, range characteristics, and >42,000 genetic sequences for 299 species using open-access scripts and various databases. After identifying important predictors of nucleotide diversity with random forest regression, we conducted follow-up analyses to examine the roles of phylogenetic history, geography, and demographic processes on intraspecific diversity. Although life history traits were not important predictors for this dataset, we found significant phylogenetic signal in genetic diversity within amphibians. We also found that salamander species at northern latitudes contain lower genetic diversity. Data repurposing and machine learning provide valuable tools for detecting patterns with relevance for conservation, but concerted efforts are needed to compile meaningful datasets with greater utility for understanding global biodiversity.

Keywords

Caudata, data repurposing, latitude, random forests, nucleotide diversity, phylogenetic signal

Introduction

Genetic variation within species is a fundamental aspect of diversity that has long intrigued population geneticists, biogeographers, and conservation biologists (Avise, 2000; Lande, 1988; Leffler et al., 2012). Within species, genetic diversity includes both diversity within populations and genetic differences among populations, typically described as population genetic structure. Species with sufficient genetic diversity may have the ability to develop resistance to disease or adapt to environmental change (Ekroth, Rafaluk-Mohr, & King, 2019; Frankham, 1995; Jamieson & Allendorf, 2012). In contrast, species with limited genetic diversity can experience detrimental effects due to inbreeding depression and may be at higher risk of extinction from catastrophic events such as disease outbreaks (Frankham, 2005; Hedrick & Kalinowski, 2000; Lande, 1988). While the importance of conserving intraspecific diversity is well-established (Moritz, 2002; Moritz & Faith, 1998; Paz-Vinas et al., 2018), it remains to be fully integrated as standard practice in global conservation strategies (Laikre et al., 2020). In the past few decades, the availability of genetic datasets has grown dramatically, and large amounts of genetic data are now available to investigate diversity within species (Garrick et al., 2015). Furthermore, modern researchers have access to troves of data including species traits, genetic information, and geographic distributions from museum collections (e.g., iDigBio), genetic data repositories (e.g., NCBI GenBank), biodiversity portals (e.g., GBIF) and phenotypic databases (e.g., MorphoBank). Predicting and understanding the determinants of intraspecific genetic diversity broadly across taxonomic groups is an increasingly attainable goal.

Many potential predictors of intraspecific genetic diversity have been implicated. Effective population size and mutation rate are two key determinants of genetic diversity, each of which is governed by life history traits, which are expected to influence the accumulation and

maintenance of genetic variation (Ellegren & Galtier, 2016). For example, Romiguier et al.,

(2014) investigated genome-wide diversity of 76 animal species and found a correlation between

genetic diversity and life history strategy; long-lived species with few, large offspring and thus

higher parental investment ("K-strategists") had lower genetic diversity than species with many,

small offspring ("r-strategists"). Chen, Glémin, & Lascoux, (2017) confirmed these results for

animals and also found that longevity and mating system explained genomic diversity in plants.

Body size has been negatively correlated with genetic diversity in various taxonomic groups

including mammals (Brüniche-Olsen, Kellner, Anderson, & DeWoody, 2018), bees (López-

Uribe, Jha, & Soro, 2019), and butterflies (Mackintosh et al., 2019), a relationship that is

presumed to correspond with limits to population abundance in larger-bodied species (White,

Ernest, Kerkhoff, & Enquist, 2007). Ecological traits associated with dispersal propensity,

including body size, larval period, philopatry, or habitat specialization, could also influence

whether or not species exhibit genetic structure (Mims, Hauser, Goldberg, & Olden, 2016; Paz,

Ibáñez, Lips, & Crawford, 2015; Semlitsch, 2008). However, empirical support for associations

between species traits and genetic diversity across taxonomic groups has been mixed. Within

butterflies, Mackintosh et al., (2019) found no relationship between genetic diversity and

longevity, propagule size, abundance, or host range. In Australian lizards, Grundler, Singhal,

Cowan, & Rabosky, (2019) found no relationship between genetic diversity and body size or

habitat specialization, and Singhal et al., (2017) found the only significant predictor of

intraspecific diversity to be the number of museum occurrence records, a possible proxy for

abundance. Investigations such as these exemplify the possibility that the determinants of genetic

diversity may not be easily predictable and are likely to vary among taxonomic groups.

In addition to life history traits, geographic range characteristics can be useful metrics for

predicting intraspecific genetic diversity. Total range size is sometimes treated as a proxy for

census population size, which should scale positively with genetic diversity, because species that

occupy more area or that have historically been able to expand to new areas are expected to have

higher abundances (Leffler et al., 2012; Singhal et al., 2017). Additionally, total range size should

be related to genetic structure because larger ranges typically encompass more habitat variation or

physical barriers, leading to patterns of isolation by distance or environment, assuming the

species are not capable dispersers (Sexton, Hangartner, & Hoffmann, 2014; Wright, 1943).

Incorporating geographic variables such as elevational, topographic, or climatic variation has

illustrated the influence of these factors on population connectivity and phylogeographic structure

(Rodríguez et al., 2015; Wang, 2012). In a global study mapping amphibian genetic diversity,

Miraldo et al., (2016) identified a latitudinal gradient of genetic diversity with higher diversity in

the tropics. Similarly, at the intraspecific level, latitude has been negatively associated with

genetic structure and diversity for a broad range of taxa (Brüniche-Olsen et al., 2018; Pelletier &

Carstens, 2018; Smith, Seeholzer, Harvey, Cuervo, & Brumfield, 2017). A plausible explanation

is that the climatic and environmental instability experienced by taxa at higher latitudes during

glacial cycles contributed to this pattern by reducing suitable habitat and population sizes

(Hewitt, 2000). The continued integration of intraspecific genetic variation, species traits, and

geographic variables is needed to better understand the drivers of genetic diversity across

different taxonomic groups and spatial scales.

Meta-analysis and data repurposing represent two strategies for integrating and analyzing

complex data types from multiple species. Meta-analyses combine results from many studies to

increase power and improve our understanding of key phenomena (e.g., Field et al., 2009; Soltis,

Morris, McLachlan, Manos, & Soltis, 2006). Meta-analyses are able to draw from individual

studies, each tailored to a focal taxon, but it is not easy to synthesize results across studies that

differ in their aims, types of data, inference methods, and sampling design. It is often more

efficient to proceed via data repurposing (Sidlauskas et al., 2010), where data from multiple

studies are reanalyzed in a common framework. Repurposed data can be effectively combined

with machine learning techniques to investigate questions of interest (e.g., Pelletier & Carstens,

2018). This strategy is consistent with broader goals in ecology and evolution to make data

accessible and standardized (Blanchet, Prunier, & De Kort, 2017; Carpenter et al., 2009; Parr,

Guralnick, Cellinese, & Page, 2012; Peters, 2010; Pope, Liggins, Keyse, Carvalho, & Riginos,

2015; Reichman, Jones, & Schildhauer, 2011; Sidlauskas et al., 2010; Soltis & Soltis, 2016).

Furthermore, repurposing published data enhances their value and generates a greater return on

investment in these data by funding agencies.

Of the many machine learning techniques available, we utilize the random forest approach

introduced by Breiman (2001) because it is fast, easy to implement, can accommodate very large

datasets including many predictor variables, and enables an intuitive evaluation of the importance

of these variables for predicting the response. Random forest is a machine learning ensemble

approach that uses multiple decision trees (a forest) to predict a user-defined response based on

many potential predictor variables. Each individual decision tree consists of a subset of the data

and a random subset of predictor variables at the nodes. Individually, each tree is a weak

predictor, but when many trees are combined, random forest results in high predictive accuracy

(Breiman, 2001). The importance of each variable is determined by examining the increase in

prediction error after randomly permuting the data for that variable while all others are left

unchanged. A review of random forest classification and regression is provided by (Liaw &

Wiener, 2002). Here, we apply random forest regression to address a long-standing question in

evolutionary biology (Leffler et al., 2012): What factors determine genetic diversity within

species?

Nearctic amphibians as a case study

Amphibians have long been considered the most imperiled class of vertebrates (Stuart et

al., 2004). More than 40% of data-sufficient amphibian species are considered threatened by the

International Union for Conservation of Nature (IUCN, 2019), and considerable efforts have

focused on addressing the issue that a large proportion of species (~25%) are data deficient,

meaning there is insufficient information to assign them to an IUCN threat category (Nori,

Villalobos, & Loyola, 2018). In a global analysis of amphibians, González-del-Pliego et al.,

(2019) found that body size and range size were correlated with threat status, estimated that half

of data-deficient species are threatened, and identified regions of the world (e.g., the Neotropics

and Southeast Asia) where a high number of data-deficient species are predicted to be threatened.

Using machine learning, Howard & Bickford (2014) predicted that >63% of data-deficient

species are threatened and identified similar geographic regions with high predicted extinction

risk. However, potentially useful information is lacking from these studies. Notably, intraspecific

genetic diversity and the factors that influence this measure are often absent in conservation

assessments, despite their relevance to understanding the long-term persistence and adaptive

capacity of species.

Our goal was to aggregate existing trait, geographic, and genetic data to investigate the

predictors of genetic diversity in amphibians. We focused on Nearctic amphibians as an initial

test case given the long history of study, tractable number of species, and presumed availability of

natural history and genetic information. This dataset included all species native to Canada and the

U.S., a subset of which have ranges extending into Mexico. First, we compiled species' range

variables, natural history characteristics, and gene alignments using a series of open-access

databases and scripts. Second, we used random forest regression to identify predictors of

intraspecific nucleotide diversity based on *Cytochrome b* (*cytb*), the gene with the most data

available for amphibians and one that is commonly used for studying genetic variation within

species (e.g., Vences, Thomas, Bonett, & Vieites, 2005). Finally, we investigated the relevance of

phylogenetic history and demographic processes in contributing to current patterns of genetic

diversity identified across amphibian species. We found evidence that genetic diversity within

species is phylogenetically conserved, species traits do not predict genetic diversity for this set of

species, and geographic range characteristics correspond with genetic diversity in Nearctic

salamanders.

Materials and Methods

Species traits

We compiled trait data for amphibian species focusing on traits related to life history and

ecology that may influence intraspecific genetic diversity. The following traits were initially

extracted from the AmphiBIO database (Oliveira, São-Pedro, Santos-Barrera, Penone, & Costa,

2017): development mode (larval or direct), maximum body size (mm), age at maturity (min and

max years), longevity (max years), clutch size (min and max number of eggs), and egg size (min

and max in mm). We retrieved additional body size information from the Peterson field guides to

reptiles and amphibians of the United States (Powell, Conant, & Collins, 2016; Stebbins &

McGinnis, 2018). We used maximum snout-vent length (SVL) for Anura (frogs), and maximum

total length for Caudata (salamanders; note that SVL was not always available for Caudata). The

remaining traits of interest (neoteny, breeding habitat, larval period, time to hatching, dispersal

distance, and home range size) were compiled primarily from species accounts in

(AmphibiaWeb, 2018) <https://amphibiaweb.org>. Neoteny, or the retention of juvenile

characteristics as adults, was categorized as 'yes' (a species is always neotenic), 'no' (neoteny

has never been reported), or 'some' (neoteny has been reported in some populations, but not all).

Breeding habitat, including the habitat in which eggs and/or larvae develop, was categorized as

'terrestrial', or 'aquatic', with aquatic species further categorized as 'permanent', 'ephemeral', or

'aquatic generalist'. Larval period was first recorded numerically (min and max days) and then

binned into categories: 'none' (direct developing), 'short' (0–90 days), 'mid' (91–365 days), or

'long' (>1 year). Time to hatching (min and max days), dispersal distance (max recorded in

meters), and home range size (max recorded in meters squared) were recorded when available,

but were unavailable for >40% of species and were thus excluded from further analyses in the

present study (Fig. S1).

Range characteristics

Species range maps for amphibians were downloaded as shapefiles from the IUCN Red

List of Threatened Species Version 6.1 (www.iucnredlist.org/) on 22 January 2019. Range maps

were manually edited in QGIS v. 2.18.2 as needed to exclude non-native ranges originally

included in IUCN range maps, and to generate range maps for species that have been described

recently from a portion of a former species' range (e.g., *Acris blanchardi* split from *Acris*

crepitans; Gamble, Berendzen, Bradley Shaffer, Starkey, & Simons, 2008). Revised shapefiles

are available on Dryad (https://doi.org/10.5061/dryad.0cfxpnvzh; Barrow, Fonseca, Thompson, &

Carstens, 2020). Altitude and bioclimatic variables were downloaded from WorldClim v. 1.4

(Hijmans, Cameron, Parra, Jones, & Jarvis, 2005) at a spatial resolution of 2.5 minutes (~21

km^2). We extracted the following information from the list of focal amphibian species: total

range size (km^2), latitude (min, max, extent = max-min, midpoint = (max+min)/2), altitude (min,

max, average, standard deviation (SD)), slope (min, max, average, SD, extent), and bioclimatic

variables (min, max, average, SD) using a custom script in R and the packages 'rgdal' (Bivand,

Keitt, & Rowlingson, 2019), 'raster' (Hijmans, 2019b), and 'geosphere' (Hijmans, 2019a).

Analyses were conducted in R v. 3.6 (R Core Team, 2019) and the scripts and datasets are

available on Dryad.

Genetic sequences

Sequences were downloaded from GenBank in two ways and were processed using a

series of R and Python scripts modified from Pelletier & Carstens (2018). First, we obtained

georeferenced sequences for all Nearctic amphibian species by querying accession numbers

linked to records downloaded from (GBIF.org, 28 November 2018). Second, we downloaded all

GenBank sequences for the 19 focal amphibian families, regardless of whether they were

georeferenced, and only retained sequences from species within these families that have native

ranges in the Nearctic. For both the georeferenced and non-georeferenced sequence sets

separately, we sorted sequences by species and gene and generated multiple sequence alignments

for every species by gene combination using Mafft v. 7.402 (Katoh & Standley, 2013). After

summarizing the number of species and locus lengths for each species, we determined that

cytochrome b (*cytb*) was the best-represented gene (Fig. S2). Alignments for *cytb* were then

visually inspected in Geneious v. 8.1.9 (https://www.geneious.com) and edited to remove

misaligned sequences prior to further analysis (<0.3% removed). Alignments for six additional

anuran species were added from previously published datasets (Barrow, Bigelow, Phillips, &

Lemmon, 2015; Barrow, Soto-Centeno, Warwick, Lemmon, & Moriarty Lemmon, 2017)

available on Dryad or as unannotated mitochondrial genomes on GenBank (Accession numbers

MF198257–MF198403; note these were missed by our scripts because gene names are not

defined on GenBank, but sequences could be manually aligned to known *cytb* sequences).

For each species, we calculated nucleotide diversity using the nuc.div() function in the

'pegas' R package (Paradis, 2010). To assess sensitivity of nucleotide diversity to sample size, or

the number of individuals sequenced, we randomly subsampled sequences from each *cytb* dataset

with replacement to generate 100 datasets for each sample size between two and 25 sequences.

We calculated nucleotide diversity for each dataset and examined the variance in the estimates

from the 100 replicate datasets for each sample size. Given the steep decline in variance observed

as sample size increased above five (Fig. S3), we conducted analyses of genetic diversity with

species that included at least five sequences. Analyses were repeated using (1) the original value

of nucleotide diversity for the full datasets and (2) the median value of nucleotide diversity from

the 100 datasets with five randomly sampled sequences. Given the consistent results between the

two datasets, we report results only for the latter. The R code and results for all analyses are

available on Dryad.

Random forest regression

We used random forest regression in the R package 'randomForest' (Liaw & Wiener,

2002) to predict intraspecific nucleotide diversity based on species traits (body size, development

type, breeding habitat, neoteny, clutch size, and larval period), range characteristics (total range

size, latitude, altitude, slope, and bioclimatic variables), and three additional predictors:

taxonomic family, the number of GBIF occurrences, and the number of sequences for each

species. We grew 2000 trees and used the defaults of randomly sampling one third of the

predictors at each node ('m_{try}') and minimum of five terminal nodes ('nodesize'). Variable

importance was determined based on the percent increase in mean squared error (IncMSE) of

prediction when that predictor variable was randomly permuted (removing its effect) while all

others were held constant. Model performance was assessed as the percent of variance in the

response explained by the models. We conducted analyses for the dataset including species with

at least five sequences and then conducted separate analyses for Anura and Caudata because

some species traits are not comparable or applicable to both orders. For example, body size is

measured differently between the two orders, only salamanders exhibit neoteny, and very few

frog species native to the region have direct development. To examine the consistency of variable

importance, we conducted 100 iterations with 2000 trees each, then averaged the percent variance

explained and IncMSE for each variable across iterations. After conducting analyses with all

variables of interest, we repeated analyses with reduced sets of predictors for ease of visualizing

and interpreting the results.

Phylogenetic comparative methods

Random forest analyses pointed to the importance of taxonomy for predicting nucleotide

diversity (see Results). To better understand these results, we downloaded phylogeny subsets

from VertLife.org for the Nearctic amphibian species in our dataset. Phylogenetic relationships

are from Jetz & Pyron (2018) and are based on a time-calibrated posterior tree distribution for

nearly all extant amphibian species, generated using a combination of phylogenetic inference and

taxonomic assignment. We tested for phylogenetic signal in continuous life history traits and

nucleotide diversity using the phylosig() function and one tree randomly sampled from the

posterior. To visualize the differences in genetic diversity among species, we mapped median

cytb nucleotide diversity onto the tree as a continuous trait using the contMap() function in the R

package 'phytools' (Revell, 2012). This function estimates maximum likelihood ancestral traits at

internal nodes and interpolates the states along the branches based on Felsenstein (1985).

Additional analyses and tree pruning were conducted using the R package 'ape' (Paradis &

Schliep, 2019). To account for phylogenetic relatedness between important continuous variables

(see Results), we calculated phylogenetically independent contrasts using the pic() function,

which implements the method described by Felsenstein (1985). We then assessed relationships

between these continuous variables using linear models.

Species distribution models

284   Random forest analyses indicate that the minimum latitude of a species was an important

predictor of genetic diversity (see Results). We hypothesized that the signature of past

demographic events, including population bottlenecks or post-glacial population expansion,

would be more pronounced in species at higher (more northern) latitudes. We predicted that: (1)

species at northern latitudes would have been more likely to experience range contraction during

the last glacial maximum (LGM) than species at southern latitudes, and (2) species at northern

latitudes would have low niche stability, or minimal overlap between the LGM and present

suitability models. To test these predictions independent of the genetic data, we modeled species

distributions based on current climate and hindcasted these to the LGM climate conditions.

Occurrence records for each species were downloaded from (GBIF.org, 04 July 2019) using a

polygon to encompass the ranges of all Nearctic amphibians. Custom scripts (available from

Dryad, Barrow et al., 2020) were used to generate a file of occurrence localities for each species

with *cytb* data, removing any points that fell outside the IUCN range map (0.5 degree buffer

width) for that species, and thinning occurrence points to retain unique localities at a threshold of

10 km using the R package 'spThin' (Aiello-Lammens, Boria, Radosavljevic, Vilela, &

Anderson, 2015). For species with a very large number of occurrences, we randomly sampled

3,000 localities prior to thinning to save computational time, and for species with small ranges

and <25 localities, we did not thin datasets. We estimated species distribution models (SDMs) for

138 species with *cytb* data.

We downloaded 19 bioclimatic layers from WorldClim v. 1.4 (Hijmans et al. 2005) for

current climate at a resolution of 30 seconds (~1 km^2) and for the LGM (CCSM4) at the highest

resolution available (2.5 min). We conducted SDM analyses in R using the packages 'biomod2'

(Thuiller, Georges, Engler, & Breiner, 2019), 'raster' (Hijmans, 2019b), 'rgdal' (Bivand, Keitt, &

Rowlingson, 2019), 'rgeos' (Bivand & Rundel, 2019), and 'HDInterval' (Meredith & Kruschke,

2018). Bioclimatic layers for each time period were stacked, clipped to the extent of North

America (-170, -52, 12, 72), and masked by current land borders. We tested for correlations

between current climate layers and removed highly correlated variables (>0.8; Peterson, 2011),

retaining five temperature (bio2, bio3, bio7, bio8, and bio10) and five precipitation variables

(bio13, bio14, bio15, bio18, and bio19). Ensemble models were generated with the Generalized

Linear Models ('GLM'), Random Forest ('RF'), and Maxent ('MAXENT.phillips') modeling

options in 'biomod2'. We used five model replicates for cross-validation, each with 80% of the

data for training and 20% for testing and evaluated model accuracy with the 'ROC' evaluation

metric. Models were hindcasted onto the LGM climate and were projected onto the current

climate with the same resolution as the LGM (2.5 min).

For each time period, the five replicate models were averaged prior to calculating niche

stability metrics. We converted models to binary presence/absence layers using a 95% occurrence

probability threshold. The resulting models include the total area in which the model probability

is equal to or greater than the probability threshold where 95% of the occurrence points are

contained. For each threshold, we calculated the total suitable area for current and LGM models

and determined whether the suitable area in the LGM was smaller (range contraction in the LGM)

or larger (range expansion in the LGM) than the suitable area in the current model. We also

determined the area of overlap between the two time periods and calculated the overlap index

(OI) metric described by Hijmans & Graham (2006). The OI describes the proportion of the

current range that was also suitable during the LGM, with higher values indicating higher niche

stability. We compared the minimum latitude of species that were (1) predicted to undergo range

contraction versus expansion and (2) species that had no niche overlap versus some niche overlap

using Student's t-tests in R.

Results

Data summary

We compiled data for 299 amphibian species native to the U.S. and Canada, 44 of which

have ranges extending into Mexico. The dataset consists of 197 salamanders in nine families and

102 frogs in 10 families. Complete trait data were available for body size, development, breeding

habitat, and neoteny, with various degrees of missing data for the remaining traits (Fig. S1).

Range characteristics derived from IUCN range maps were available for 268 species, with data

unavailable for recently described species (since 2008) or those with very small ranges. We

assembled 4,759 georeferenced GenBank sequences that were linked to GBIF records, but only

37 species (20 salamanders, 17 frogs) were represented by at least five sequences from at least

two localities. Preliminary analyses indicated there was no predictive power in models of genetic

structure with this small dataset. We therefore focused analyses on non-spatial genetic datasets

and used metrics of overall intraspecific genetic diversity. We assembled 42,067 GenBank

sequences from 263 species, which consisted primarily of mitochondrial genes (Fig. S2). The best

represented gene was *cytb*, with at least five sequences available for each of 147 species.

Predictors of intraspecific genetic diversity

Full models with all predictors included 137 amphibian species with *cytb* data. These

models explained an average of 24.9% variance in median nucleotide diversity based on 100

sequence datasets with five randomly sampled sequences. The most important predictors were

consistently taxonomic family and the number of sequences, indicating that nucleotide diversity

is influenced by phylogenetic relatedness and sample size. Some bioclimatic variables, including

the average precipitation of the wettest month (bio13), average precipitation of the warmest

quarter (bio18), average precipitation of the driest quarter (bio17), and the standard deviation of

mean temperature of the driest quarter (bio9), were also ranked as important. Models with a

subset of predictors explained an average of 30.1% variance in median nucleotide diversity.

Taxonomic family, the number of sequences, and average precipitation of the wettest month

(bio13) remained the most important predictors of nucleotide diversity, while species traits were

not predictive of nucleotide diversity for this set of species (Fig. 1a).

When we considered each order separately, the Anura dataset included only 39 species

with *cytb* data and there was no predictive power in these models (variance explained was

negative). Full models for Caudata, however, included 98 species and explained an average of

20.5% of the variance in median nucleotide diversity. The most important predictors of

nucleotide diversity were consistently the number of sequences and the minimum latitude of a

species range. Models with a subset of predictors explained an average of 28.9% of the variance

in median nucleotide diversity. The number of sequences and minimum latitude were the most

important predictors (Fig. 1b). Models with fewer species (n=67) and additional trait information

available explained an average of 28.3% variance in median nucleotide diversity. Minimum

latitude remained the most important predictor and species traits were not important predictors of

nucleotide diversity (Fig. 1c).

Phylogenetic signal, minimum latitude, and climatic niche stability

We found phylogenetic signal in species-wide nucleotide diversity (lambda = 0.152, p =

0.008). Phylogenetic signal is also present in several traits related to reproduction and ecology

including clutch size, larval period, and body size (Table S1), although these traits do not predict

nucleotide diversity (described above and further tested with phylogenetic generalized linear

mixed models; results not shown). The salamander families Sirenidae and Plethodontidae

included multiple species with high nucleotide diversity (Fig. 2), but within Caudata,

phylogenetic signal in nucleotide diversity was not significant (lambda = 0.105, p = 0.605).

Notably, there was no phylogenetic signal in the number of sequences sampled, suggesting that

the phylogenetic signal present in nucleotide diversity is not an artefact of sampling.

After accounting for phylogenetic relationships there was a negative correlation between

species-wide nucleotide diversity and minimum latitude in Caudata (Fig. 3). Species with more

northern ranges tended to have lower nucleotide diversity (98-species dataset: $R^2 = 0.057$, p =

0.0102; 67-species dataset: $R^2 = 0.115$, p = 0.003). Plots of the raw values of nucleotide diversity

and minimum latitude are shown in Fig. S4. Results from SDMs did not, however, support the

hypothesis that species at higher latitudes were more heavily impacted by glaciation at the LGM.

At the 95% occurrence probability threshold, we inferred LGM contraction of 76 Caudata species

and LGM expansion for 20 species, but there was no difference in minimum latitude of the two

groups (t = -1.59, p = 0.12). At the same threshold, 49 Caudata species had no overlap between

the LGM and current models, indicating low niche stability, while 47 species had at least some

overlap, indicating predicted areas of stability. There was no difference in minimum latitude of

the two groups (t = 0.83, p = 0.41).

Discussion

Combining repurposed data with machine learning is a powerful strategy for addressing

broad questions in evolutionary biology and conservation (Howard & Bickford, 2014; Sidlauskas

et al., 2010). Here, we apply these approaches to investigate a classic question: What factors

determine intraspecific genetic variation? We discuss results from our case study, including novel

insights in Nearctic amphibian biogeography and broader implications that demonstrate the

relevance of continued studies in this framework. We then highlight the prospects and challenges

of combining data repurposing and machine learning to address questions that involve many

complex data types and predictors.

Predictors of genetic diversity in Nearctic amphibians

Amphibians play critical ecological roles as consumers (Whiles et al., 2006), prey

(Zipkin, DiRenzo, Ray, Rossman, & Lips, 2020), and indicators of aquatic and terrestrial

environmental health (Kerby, Richards-Hrdlicka, Storfer, & Skelly, 2010; Vitt, Caldwell, Wilbur,

& Smith, 1990; Welsh & Ollivier, 1998). Given these roles, additional efforts to synthesize

diverse datasets are warranted to inform conservation priorities. We found that minimum latitude

predicted genetic diversity within salamanders, with generally lower intraspecific genetic

diversity for species at higher (i.e., more northern) latitudes. Intraspecific genetic diversity also

appears to be phylogenetically conserved, a finding perhaps explained by conserved life history

traits such as clutch size or body size that are associated with intraspecific genetic diversity (e.g.,

(Paz et al., 2015; Romiguier et al., 2014). While life history traits are not predictors of genetic

diversity for this dataset, taxonomic family was identified as important. This is likely driven by

two salamander families, Sirenidae and Plethodontidae, which contain most species with the

highest intraspecific diversity. Since more than half of the amphibian species in our dataset are

plethodontid salamanders, the lack of a relationship between traits and genetic diversity might be

explained by the limited trait variation (e.g., all are relatively small-bodied and have small clutch

sizes) but considerable variation in genetic diversity within the group (Fig. 2).

The relevance of geography for understanding patterns of genetic diversity is well known

(Hewitt, 2000; Wright, 1943). The negative association between latitude and intraspecific genetic

diversity likely relates to population history since high intraspecific diversity at lower latitudes

may be attributed to long-term stability (Miraldo et al., 2016) and low diversity at northern

latitudes may be explained by population bottlenecks during glacial periods (Hewitt, 2000).

Published phylogeographic studies on some of the species included in our study support this

prediction, attributing lower genetic diversity at higher latitudes to postglacial range expansion

(Carstens, Brunsfeld, Demboski, Good, & Sullivan, 2005; Highton & Webster, 1976; Radomski,

Hantak, Brown, & Kuchta, 2020). However, the effect of past climate cycles on intraspecific

genetic diversity is clearly dependent on refugial dynamics, as a species that is restricted into a

single glacial refugium would exhibit different patterns than one where populations were isolated

in separate refugia. One complicating factor that we did not address here is whether current

species taxonomy is an appropriate unit of comparison, or whether some species include cryptic

lineages that have not been formally described. Future studies with georeferenced localities and

assignment of sequences to lineages to estimate within-population diversity should help clarify

these issues further.

Prospects and challenges for applying machine learning to repurposed data

Data repurposing can meet some of the same goals as meta-analysis, such as establishing what we (as a field) understand about key patterns and pinpointing underlying mechanisms(Carstens, Morales, Field, & Pelletier, 2018). Both types of investigation play an important role by potentially identifying connections among many variables. However, machine learning techniques make data repurposing more appealing than meta-analysis as a strategy for extracting the most information out of publicly available data. While meta-analyses certainly provide important insights, genetic datasets collected and analyzed separately across studies, which may use disparate sample designs and report incongruent metrics, can limit the number of useable datasets and make interpretation difficult (Emel & Storfer, 2012; López-Uribe et al., 2019). Once data are assembled, machine learning enables researchers to analyze very large datasets that incorporate many complex variables. By including a large suite of potential predictors, random forests can identify potential links among variables in an unbiased manner. This feature is useful for complex topics such as the determinants of genetic diversity, where researchers might be biased by previous findings and focus only on a subset of predictor variables. One caveat is that machine learning analysis is only as useful as the response variable used for a particular question. For example, while predictive models of binary questions (e.g., 'LC' versus 'nonLC' IUCN status; Fig. S5) will likely be able to explain more variation than those that rely on regression techniques, not all questions can easily be reduced to a binary outcome (e.g., a continuous response such as nucleotide diversity). In this case, the best random forest regression models explained ~30% of the variance in nucleotide diversity, and the

importance of the number of sequences per species points to sampling issues that may only be

solved with the collection of additional data.

The application of machine learning techniques to repurposed data offers great potential

for addressing key questions in ecology and evolutionary biology, but several challenges remain

(Table 1). Our focus on assembling a curated set of species allowed us to identify several errors

that would have been missed by automated pipelines. While useful for future work, this effort

limited both the taxonomic and geographic scale of our study. For example, it seems plausible

that the associations between life history traits and genetic diversity identified previously may

only be apparent across broader taxonomic scales (e.g., Romiguier et al., 2014) or when more

species can be included that span substantial trait variation (e.g., Brüniche-Olsen et al., 2018).

Expanding the set of taxa from this study to a broader regional or global scale is a clear next step

that will require addressing two substantial challenges. Compiling meaningful trait datasets for

large numbers of species is time-consuming, and for many species, natural history information is

not easily available. Solving this challenge will require collaborative efforts by the research

community to standardize trait information, contribute to trait datasets, and make these datasets

easily available (e.g., Oliveira et al., 2017). In the future, increasing efforts to digitize natural

history collections and develop informatics tools will ultimately enable the incorporation of

individual and population-level trait variation into these types of studies (Guralnick et al., 2016;

Hedrick et al., 2020). The second major challenge to extending the scope of this study is the

limited amount of georeferenced genetic data available for most species.

Two main axes of sampling effort are important to interpreting intraspecific genetic

diversity: the number (and location) of individuals sampled and the number (and type) of loci

sampled. Researchers have typically had to make a trade-off between these two for logistical

reasons. Although the number of loci that can feasibly be sequenced is growing rapidly (Garrick

et al., 2015), the vast majority of available sequences are not directly linked to localities

(Marques, Maronna, & Collins, 2013). Increasing the amount of georeferenced genetic data

available per species will enable within-population variation to be disentangled from among-

population variation and will foster a better understanding of the mechanisms that influence

intraspecific genetic diversity. Currently, the genetic sequences available for most animal species

are mitochondrial genes and it remains unclear how representative mitochondrial DNA actually is

for describing intraspecific genetic diversity (Bazin, Glémin, & Galtier, 2006; Galtier, Nabholz,

GlÉmin, & Hurst, 2009; Mulligan, Kitchen, & Miyamoto, 2006). Furthermore, it is likely that

available molecular markers do not accurately reflect adaptive variation (Mittell, Nakagawa, &

Hadfield, 2015), which may be important for designing conservation strategies that promote

species persistence (Moritz, 2002). Despite these issues, comparing genetic diversity among

species yields important insights, and new findings will be enhanced by continued efforts to link

existing genetic data to localities and to collect genome-scale data from a large number of non-

model species encompassing variation in geographic complexity and life history traits.

Conclusions

Identifying the predictors of intraspecific genetic diversity and the scale for which they

are predictive are important goals for both evolutionary biology and applied conservation. We

demonstrated the potential for combining repurposed data with machine learning techniques to

predict genetic diversity in Nearctic amphibians. Our findings indicate that life history traits do

not predict genetic diversity in this dataset, but future studies should incorporate additional

species and trait variation on a global scale. Total range size is an important criterion for

assessing species conservation risk, but it does not predict genetic diversity within Nearctic

amphibians. Instead, we found that minimum latitude was an important predictor of genetic

diversity within salamanders, suggesting that this aspect of a species' range represents an

important component to consider when assessing species risk. Northern latitude species that

harbor low genetic diversity may be even more vulnerable to future climate change scenarios or

disease outbreaks. Identifying these potential risks by combining new and existing datasets can

lead to proactive management strategies that preserve remaining genetic diversity.

Acknowledgements

We thank Drew Duckett, Flavia Mol Lanna, Danielle Parsons, Megan Smith, Tara Pelletier,

Jamin Wieringa, and Tamaki Yuri for helpful discussions, and the many contributors to

AmphibiaWeb, GenBank, GBIF, and the IUCN Red List for archiving the data that made this

study possible. This work was supported by NSF DBI 1910623, The Ohio State University

(OSU) President's Postdoctoral Scholars Program, and the OSU Department of Evolution,

Ecology and Organismal Biology. EMF thanks the Coordenação de Aperfeiçoamento de Pessoal

de Nível Superior (CAPES) for his doctoral fellowship (process #88881.170016/2018).

References

Aiello-Lammens, M. E., Boria, R. A., Radosavljevic, A., Vilela, B., & Anderson, R. P. (2015). spThin: An R package for spatial thinning of species occurrence records for use in ecological niche models. *Ecography*, *38*(5), 541–545. doi: 10.1111/ecog.01132

AmphibiaWeb. (2018). <https://amphibiaweb.org>. Retrieved December 18, 2018, from University of California, Berkeley, CA, USA. Accessed 18 Dec 2018. website: https://amphibiaweb.org

Avise, J. C. (2000). *Phylogeography: the history and formation of species*. Cambridge, Massachussetts: Harvard University Press.

Barrow, L. N., Bigelow, A. T., Phillips, C. A., & Lemmon, E. M. (2015). Phylogeographic inference using Bayesian model comparison across a fragmented chorus frog species complex. *Molecular Ecology*, *24*(18), 4739–4758. doi: 10.1111/mec.13343

[dataset] Barrow, L. N., Fonseca, E. M., Thompson, C. E. P., & Carstens, B. C. (2020). Data from: Predicting amphibian intraspecific diversity with machine learning: Challenges and prospects for integrating traits, geography, and genetic data. *Dryad Digital Repository*, https://doi.org/10.5061/dryad.0cfxpnvzh.

Barrow, L. N., Soto-Centeno, J. A., Warwick, A. R., Lemmon, A. R., & Moriarty Lemmon, E. (2017). Evaluating hypotheses of expansion from refugia through comparative phylogeography of south-eastern Coastal Plain amphibians. *Journal of Biogeography*, *44*, 2692–2705. doi: 10.1111/jbi.13069

Bazin, E., Glémin, S., & Galtier, N. (2006). Mitochondrial Genetic Diversity in Animals. *Science*, *312*(April), 570–572.

Bivand, R., Keitt, T., & Rowlingson, B. (2019). *rgdal: Bindings for the "Geospatial" Data Abstraction Library. R package version 1.4-8.* (p. https://cran.r-project.org/package=rgdal.). p. https://cran.r-project.org/package=rgdal.

Bivand, R., & Rundel, C. (2019). *rgeos: Interface to Geometry Engine - Open Source ('GEOS'). R package version 0.4-3. https://CRAN.R-project.org/package=rgeos.*

Blanchet, S., Prunier, J. G., & De Kort, H. (2017). Time to Go Bigger: Emerging Patterns in Macrogenetics. *Trends in Genetics*, *33*(9), 579–580. doi: 10.1016/j.tig.2017.06.007

Breiman, L. (2001). Random forests. *Machine Learning*, *45*, 5–32. doi: 10.1201/9780367816377-11

Brüniche-Olsen, A., Kellner, K. F., Anderson, C. J., & DeWoody, J. A. (2018). Runs of homozygosity have utility in mammalian conservation and evolutionary studies. *Conservation Genetics*, *19*(6), 1295–1307. doi: 10.1007/s10592-018-1099-y

Carpenter, S. R., Armbrust, E. V., Arzberger, P. W., Chapin, F. S., Elser, J. J., Hackett, E. J., … Zimmerman, A. S. (2009). Accelerate Synthesis in Ecology and Environmental Sciences. *BioScience*, *59*(8), 699–701. doi: 10.1525/bio.2009.59.8.11

Carstens, B. C., Brunsfeld, S. J., Demboski, J. R., Good, J. M., & Sullivan, J. (2005). Investigating the evolutionary history of the Pacific Northwest mesic forest ecosystem: hypothesis testing within a comparative phylogeographic framework. *Evolution*, *59*(8), 1639–1652. doi: 10.1111/j.0014-3820.2005.tb01815.x

Carstens, B. C., Morales, A. E., Field, K., & Pelletier, T. A. (2018). A global analysis of bats using automated comparative phylogeography uncovers a surprising impact of Pleistocene glaciation. *Journal of Biogeography*, *45*(8), 1795–1805. doi: 10.1111/jbi.13382

Chamberlain, S. A., & Szöcs, E. (2013). taxize: taxonomic search and retrieval in R.
F1000Research, *2*(191).
Chen, J., Glémin, S., & Lascoux, M. (2017). Genetic diversity and the efficacy of purifying
selection across plant and animal species. *Molecular Biology and Evolution*, *34*(6), 1417–
1428. doi: 10.1093/molbev/msx088
Ekroth, A. K. E., Rafaluk-Mohr, C., & King, K. C. (2019). Host genetic diversity limits parasite
success beyond agricultural systems: A meta-analysis. *Proceedings of the Royal Society*
B: Biological Sciences, *286*(1911). doi: 10.1098/rspb.2019.1811
Ellegren, H., & Galtier, N. (2016). Determinants of genetic diversity. *Nature Reviews Genetics*,
17(7), 422–433. doi: 10.1038/nrg.2016.58
Emel, S. L., & Storfer, A. (2012). A decade of amphibian population genetic studies: Synthesis
and recommendations. *Conservation Genetics*, *13*(6), 1685–1689. doi: 10.1007/s10592-
012-0407-1
Felsenstein, J. (1985). Phylogenies and the comparative method. *American Naturalist*, *125*(1), 1–
15.
Field, R., Hawkins, B. A., Cornell, H. V., Currie, D. J., Diniz-Filho, J. A. F., Guégan, J. F., …
Turner, J. R. G. (2009). Spatial species-richness gradients across scales: A meta-analysis.
Journal of Biogeography, *36*(1), 132–147. doi: 10.1111/j.1365-2699.2008.01963.x
Frankham, R. (1995). Effective population size/adult population size ratios in wildlife: A review.
Genetics Research, *66*, 95–107. doi: 10.1017/S0016672308009695
Frankham, R. (2005). Genetics and extinction. *Biological Conservation*, *126*(2), 131–140. doi:
10.1016/j.biocon.2005.05.002
Galtier, N., Nabholz, B., GlÉmin, S., & Hurst, G. D. D. (2009). Mitochondrial DNA as a marker
of molecular diversity: A reappraisal. *Molecular Ecology*, *18*(22), 4541–4550. doi:
10.1111/j.1365-294X.2009.04380.x
Gamble, T., Berendzen, P. B., Bradley Shaffer, H., Starkey, D. E., & Simons, A. M. (2008).
Species limits and phylogeography of North American cricket frogs (Acris: Hylidae).
Molecular Phylogenetics and Evolution, *48*(1), 112–125. doi:
10.1016/j.ympev.2008.03.015
Garrick, R. C., Bonatelli, I. A. S., Hyseni, C., Morales, A., Pelletier, T. A., Perez, M. F., …
Carstens, B. C. (2015). The evolution of phylogeographic datasets. *Molecular Ecology*,
24, 1164–1171. doi: 10.1111/mec.13108
GBIF.org (28 November 2018) GBIF Occurrence Download https://doi.org/10.15468/dl.r6bshz.
GBIF.org (04 July 2019) GBIF Occurrence Download https://doi.org/10.15468/dl.rk2srg.
González-del-Pliego, P., Freckleton, R. P., Edwards, D. P., Koo, M. S., Scheffers, B. R., Pyron,
R. A., & Jetz, W. (2019). Phylogenetic and Trait-Based Prediction of Extinction Risk for
Data-Deficient Amphibians. *Current Biology*, *29*(9), 1557-1563.e3. doi:
10.1016/j.cub.2019.04.005
Grundler, M. R., Singhal, S., Cowan, M. A., & Rabosky, D. L. (2019). Is genomic diversity a
useful proxy for census population size? Evidence from a species-rich community of
desert lizards. *Molecular Ecology*. doi: 10.1111/mec.15042
Guralnick, R. P., Zermoglio, P. F., Wieczorek, J., LaFrance, R., Bloom, D., & Russell, L. (2016).
The importance of digitized biocollections as a source of trait data and a new VertNet
resource. *Database*, *2016*. doi: 10.1093/database/baw158

Hedrick, B. P., Heberling, J. M., Meineke, E. K., Turner, K. G., Grassa, C. J., Park, D. S., …
Davis, C. C. (2020). Digitization and the Future of Natural History Collections.
BioScience, *XX*(X), 1–9. doi: 10.1093/biosci/biz163

Hedrick, P. W., & Kalinowski, S. T. (2000). Inbreeding depression in conservation biology.
Annual Review of Ecology and Systematics, *31*, 139–162.

Hewitt, G. (2000). The genetic legacy of the Quaternary ice ages. *Nature*, *405*(6789), 907–913.
doi: 10.1038/35016000

Highton, R., & Webster, T. P. (1976). Geographic protein variation and divergence in populations
of the salamander Plethodon cinereus. *Evolution*, *30*(1), 33–45.

Hijmans, R. J. (2019a). *geosphere: Spherical Trigonometry. R package version 1.5-10.*
https://CRAN.R project.org/package=geosphere.

Hijmans, R. J. (2019b). *raster: Geographic Data Analysis and Modeling. R package version 3.0-*
7. https://CRAN.R-project.org/package=raster.

Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high
resolution interpolated climate surfaces for global land areas. *International Journal of*
Climatology, *25*(15), 1965–1978. doi: 10.1002/joc.1276

Hijmans, R. J., & Graham, C. H. (2006). The ability of climate envelope models to predict the
effect of climate change on species distributions. *Global Change Biology*, *12*(12), 2272–
2281. doi: 10.1111/j.1365-2486.2006.01256.x

Howard, S. D., & Bickford, D. P. (2014). Amphibians over the edge: Silent extinction risk of
Data Deficient species. *Diversity and Distributions*, *20*(7), 837–846. doi:
10.1111/ddi.12218

IUCN. (2019). The IUCN Red List of Threatened Species. Retrieved March 6, 2020, from
Version 2019-3 website: https://www.iucnredlist.org/resources/summary-statistics#Figure
2

Jamieson, I. G., & Allendorf, F. W. (2012). How does the 50/500 rule apply to MVPs? *Trends in*
Ecology and Evolution, *27*(10), 578–584. doi: 10.1016/j.tree.2012.07.001

Jetz, W., & Pyron, R. A. (2018). The interplay of past diversification and evolutionary isolation
with present imperilment across the amphibian tree of life. *Nature Ecology and Evolution*,
2(5), 850–858. doi: 10.1038/s41559-018-0515-5

Katoh, K., & Standley, D. M. (2013). MAFFT multiple sequence alignment software version 7:
Improvements in performance and usability. *Molecular Biology and Evolution*, *30*(4),
772–780. doi: 10.1093/molbev/mst010

Kerby, J. L., Richards-Hrdlicka, K. L., Storfer, A., & Skelly, D. K. (2010). An examination of
amphibian sensitivity to environmental contaminants: Are amphibians poor canaries?
Ecology Letters, *13*(1), 60–67. doi: 10.1111/j.1461-0248.2009.01399.x

Laikre, L., Hoban, S., Bruford, M. W., Segelbacher, G., Allendorf, F. W., Gajardo, G., …
Vernesi, C. (2020). Post-2020 goals overlook genetic diversity. *Science*, *367*(6482),
1083–1085.

Lande, R. (1988). Genetics and demography in biological conservation. *Science*, *241*(4872),
1455–1460. doi: 10.1126/science.3420403

Leffler, E. M., Bullaughey, K., Matute, D. R., Meyer, W. K., Ségurel, L., Venkat, A., …
Przeworski, M. (2012). Revisiting an Old Riddle: What Determines Genetic Diversity
Levels within Species? *PLoS Biology*, *10*(9). doi: 10.1371/journal.pbio.1001388

Liaw, A., & Wiener, M. (2002). Classification and Regression by randomForest. *R News*, *2*(3), 18–22.

López-Uribe, M. M., Jha, S., & Soro, A. (2019). A trait-based approach to predict population genetic structure in bees. *Molecular Ecology*, *28*(8), 1919–1929. doi: 10.1111/mec.15028

Mackintosh, A., Laetsch, D. R., Hayward, A., Charlesworth, B., Waterfall, M., Vila, R., & Lohse, K. (2019). The determinants of genetic diversity in butterflies. *Nature Communications*, *10*(1), 1–9. doi: 10.1038/s41467-019-11308-4

Marques, A. C., Maronna, M. M., & Collins, A. G. (2013). Putting GenBank Data on the Map. *Science*, *341*(September), 1341.

Meredith, M., & Kruschke, J. (2018). *HDInterval: Highest (Posterior) Density Intervals. R package version 0.2.0. https://CRAN.R-project.org/package=HDInterval.*

Mims, M. C., Hauser, L., Goldberg, C. S., & Olden, J. D. (2016). *Genetic Differentiation , Isolation-by- Distance , and Metapopulation Dynamics of the Arizona Treefrog (Hyla wrightorum) in an Isolated Portion of Its Range*. 1–23. doi: 10.1371/journal.pone.0160655

Miraldo, A., Li, S., Borregaard, M. K., Flórez-Rodríguez, A., Gopalakrishnan, S., Rizvanovic, M., … Nogués-Bravo, D. (2016). An Anthropocene map of genetic diversity. *Science*, *353*(6307), 1532–1535. doi: 10.1126/science.aaf4381

Mittell, E. A., Nakagawa, S., & Hadfield, J. D. (2015). Are molecular markers useful predictors of adaptive potential? *Ecology Letters*, *18*(8), 772–778. doi: 10.1111/ele.12454

Moritz, C. (2002). Strategies to protect biological diversity and the evolutionary processes that sustain it. *Systematic Biology*, *51*(2), 238–254.

Moritz, C., & Faith, D. P. (1998). Comparative phylogeography and the identification of genetically divergent areas for conservation. *Molecular Ecology*, *7*(4), 419–429. doi: 10.1046/j.1365-294x.1998.00317.x

Mulligan, C. J., Kitchen, A., & Miyamoto, M. M. (2006). Comment on "Population size does not influence mitochondrial genetic diversity in animals." *Science*, *314*(5804), 1390. doi: 10.1126/science.1122033

Nori, J., Villalobos, F., & Loyola, R. (2018). Global priority areas for amphibian research. *Journal of Biogeography*, *45*(11), 2588–2594. doi: 10.1111/jbi.13435

Oliveira, B. F., São-Pedro, V. A., Santos-Barrera, G., Penone, C., & Costa, G. C. (2017). AmphiBIO, a global database for amphibian ecological traits. *Scientific Data*, *4*, 1–7. doi: 10.1038/sdata.2017.123

Paradis, E. (2010). pegas: an R package for population genetics with an integrated-modular approach. *Bioinformatics*, *26*, 419–420.

Paradis, E., & Schliep, K. (2019). Ape 5.0: An environment for modern phylogenetics and evolutionary analyses in R. *Bioinformatics*, *35*(3), 526–528. doi: 10.1093/bioinformatics/bty633

Parr, C. S., Guralnick, R., Cellinese, N., & Page, R. D. M. (2012). Evolutionary informatics: Unifying knowledge about the diversity of life. *Trends in Ecology and Evolution*, *27*(2), 94–103. doi: 10.1016/j.tree.2011.11.001

Paz, A., Ibáñez, R., Lips, K. R., & Crawford, A. J. (2015). Testing the role of ecology and life history in structuring genetic variation across a landscape: A trait-based phylogeographic approach. *Molecular Ecology*, *24*(14), 3723–3737. doi: 10.1111/mec.13275

Paz-Vinas, I., Loot, G., Hermoso, V., Veyssière, C., Poulet, N., Grenouillet, G., & Blanchet, S. (2018). Systematic conservation planning for intraspecific genetic diversity. *Proceedings of the Royal Society B: Biological Sciences*, *285*(1877). doi: 10.1098/rspb.2017.2746

Pelletier, T. A., & Carstens, B. C. (2018). Geographical range size and latitude predict population genetic structure in a global survey. *Biology Letters*, *14*(1). doi: 10.1098/rsbl.2017.0566

Peters, D. P. C. (2010). Accessible ecology: Synthesis of the long, deep, and broad. *Trends in Ecology and Evolution*, *25*(10), 592–601. doi: 10.1016/j.tree.2010.07.005

Peterson, A. T. (2011). Ecological niche conservatism: a time-structured review of evidence. *Journal of Biogeography*, *38*(5), 817–827. doi: 10.1111/j.1365-2699.2010.02456.x

Pope, L. C., Liggins, L., Keyse, J., Carvalho, S. B., & Riginos, C. (2015). Not the time or the place: The missing spatio-temporal link in publicly available genetic data. *Molecular Ecology*, *24*(15), 3802–3809. doi: 10.1111/mec.13254

Powell, R., Conant, R., & Collins, J. T. (2016). *Peterson Field Guide to Reptiles and Amphibians of Eastern and Central North America 4th Edition* (4th Editio). Boston, MA: Houghton Mifflin Harcourt.

R Core Team. (2019). *R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.* Vienna, Austria: R Foundation for Statistical Computing.

Radomski, T. P., Hantak, M. M., Brown, A. D., & Kuchta, S. R. (2020). Multilocus phylogeography of eastern red-backed salamanders: cryptic appalachian diversity and postglacial range expansion. *Herpetologica*, *76*(1), 61–73.

Reichman, O. J., Jones, M. B., & Schildhauer, M. P. (2011). Challenges and opportunities of open data in ecology. *Science*, *331*(6018), 703–705. doi: 10.1126/science.1197962

Revell, L. J. (2012). phytools: an R package for phylogenetic comparative biology (and other things). *Methods in Ecology and Evolution*, *3*(2), 217–223. doi: 10.1111/j.2041-210X.2011.00169.x

Rodríguez, A., Börner, M., Pabijan, M., Gehara, M., Haddad, C. F. B., & Vences, M. (2015). Genetic divergence in tropical anurans: deeper phylogeographic structure in forest specialists and in topographically complex regions. *Evolutionary Ecology*, *29*(5), 765–785. doi: 10.1007/s10682-015-9774-7

Romiguier, J., Gayral, P., Ballenghien, M., Bernard, A., Cahais, V., Chenuil, A., … Galtier, N. (2014). Comparative population genomics in animals uncovers the determinants of genetic diversity. *Nature*, *515*(7526), 261–263. doi: 10.1038/nature13685

Semlitsch, R. D. (2008). Differentiating Migration and Dispersal Processes for Pond-Breeding Amphibians. *Journal of Wildlife Management*, *72*(1), 260–267. doi: 10.2193/2007-082

Sexton, J. P., Hangartner, S. B., & Hoffmann, A. A. (2014). Genetic isolation by environment or distance: Which pattern of gene flow is most common? *Evolution*, *68*(1), 1–15. doi: 10.1111/evo.12258

Sidlauskas, B., Ganapathy, G., Hazkani-Covo, E., Jenkins, K. P., Lapp, H., McCall, L. W., … Kidd, D. M. (2010). linking big: The continuing promise of evolutionary synthesis. *Evolution*, *64*(4), 871–880. doi: 10.1111/j.1558-5646.2009.00892.x

Singhal, S., Huang, H., Title, P. O., Donnellan, S. C., Holmes, I., & Rabosky, D. L. (2017). Genetic diversity is largely unpredictable but scales with museum occurrences in a species-rich clade of Australian lizards. *Proceedings of the Royal Society B: Biological Sciences*, *284*(1854). doi: 10.1098/rspb.2016.2588

Smith, B. T., Seeholzer, G. F., Harvey, M. G., Cuervo, A. M., & Brumfield, R. T. (2017). A
latitudinal phylogeographic diversity gradient in birds. *PLoS Biology*, *15*(4), 1–25. doi:
10.1371/journal.pbio.2001073
Soltis, D. E., Morris, A. B., McLachlan, J. S., Manos, P. S., & Soltis, P. S. (2006). Comparative
phylogeography of unglaciated eastern North America. *Molecular Ecology*, *15*(14), 4261–
4293. doi: 10.1111/j.1365-294X.2006.03061.x
Soltis, D. E., & Soltis, P. S. (2016). Mobilizing and integrating big data in studies of spatial
and phylogenetic patterns of biodiversity. *Plant Diversity*, *38*(6), 264–270. doi:
10.1016/j.pld.2016.12.001
Stebbins, R. C., & McGinnis, S. M. (2018). *Peterson Field Guide to Western Reptiles and*
Amphibians 4th Edition (4th Editio). Boston, MA: Houghton Mifflin Harcourt.
Stuart, S. N., Chanson, J. S., Cox, N. A., Young, B. E., Rodrigues, A. S. L., Fischman, D. L., &
Waller, R. W. (2004). Status and trends of amphibian declines and extinctions worldwide.
Science, *306*(October), 1783–1786. doi: 10.1126/science.1103538
Thuiller, W., Georges, D., Engler, R., & Breiner, F. (2019). *biomod2: Ensemble Platform for*
Species Distribution Modeling. R package version 3.3-7.1. https://CRAN.R-
project.org/package=biomod2.
Vences, M., Thomas, M., Bonett, R. M., & Vieites, D. R. (2005). Deciphering amphibian
diversity through DNA barcoding: chances and challenges. *Philosophical Transactions of*
the Royal Society of London. Series B, Biological Sciences, *360*(1462), 1859–1868. doi:
10.1098/rstb.2005.1717
Vitt, L. J., Caldwell, J. P., Wilbur, H. M., & Smith, D. C. (1990). Amphibians as harbingers of
decay. *BioScience*, *40*(6), 418.
Wang, I. J. (2012). Environmental and topographic variables shape genetic structure and effective
population sizes in the endangered Yosemite toad. *Diversity and Distributions*, *18*(10),
1033–1041. doi: 10.1111/j.1472-4642.2012.00897.x
Welsh, H. H., & Ollivier, L. M. (1998). Stream amphibians as indicators of ecosystem stress: A
case study from California's redwoods. *Ecological Applications*, *8*(4), 1118–1132. doi:
10.1890/1051-0761(1998)008[1118:SAAIOE]2.0.CO;2
Whiles, M. R., Lips, K. L., Pringle, C. M., Kilham, S. S., Bixby, R. J., Brenes, R., … Peterson, S.
(2006). The effects of amphibian population declines on the structure and function of
Neotropical stream ecosystems. *Frontiers in Ecology and the Environment*, *4*(1), 27–34.
doi: 10.1017/CBO9781107415324.004
White, E. P., Ernest, S. K. M., Kerkhoff, A. J., & Enquist, B. J. (2007). Relationships between
body size and abundance in ecology. *Trends in Ecology and Evolution*, *22*(6), 323–330.
doi: 10.1016/j.tree.2007.03.007
Wright, S. (1943). Isolation by Distance. *Genetics*, *28*(2), 114–138. doi: Article
Zipkin, E. F., DiRenzo, G. V., Ray, J. M., Rossman, S., & Lips, K. R. (2020). Tropical snake
diversity collapses after widespread amphibian loss. *Science*, *367*(6479), 814–816. doi:
10.1126/science.aay5733

Data Accessibility

Author Contributions

LNB and BCC conceived the ideas; LNB, EMF, and CEPT gathered and analyzed the data; LNB and BCC wrote the paper with input from all authors.

794   **Tables and Figures**
795
796
797   **Figure Captions**
798
799
800
801   **Figure 1**
802
803   Predictors of intraspecific *cytb* nucleotide diversity for a) 137 species of Nearctic amphibians, b)
804   98 species of Caudata, and c) 67 species of Caudata with additional species traits. Average
805   variable importance (increase in mean squared error) for each predictor from 100 iterations is
806   shown.
807
808
809   **Figure 2**
810
Phylogeny of Nearctic amphibian species with *cytb* nucleotide diversity mapped as a continuous
trait. Phylogenetic relationships are based on Jetz and Pyron (2018) and were downloaded from
VertLife.org, with length indicating time in millions of years. The median nucleotide diversity
from 100 datasets with five randomly-sampled sequences was mapped onto the tree using the
contMap() function on phytools (Revell, 2012). Warmer (brown/tan) colors indicate species with
higher *cytb* nucleotide diversity.
817
818
819   **Figure 3**
820
821   Minimum latitude and *cytb* nucleotide diversity for 98 Caudata species. a) Nucleotide diversity
822   has a negative relationship with the minimum latitude of a species range. The median nucleotide
823   diversity from 100 datasets with five randomly-sampled sequences is shown. b) Phylogenetic
824   independent contrasts for the same metrics in a) using the pruned phylogeny from VertLife.org
825   (Jetz and Pyron 2018).
826
827

828   **Table 1**. Summary of challenges for repurposed data projects. Examples of challenges
829   encountered in the present study are provided with potential avenues for correction in the future.

Challenge	Example or Description	Potential for Correction
Trait Data		
Lack of natural history information	Difficult to measure traits such as dispersal, longevity, development time	• Promote and fund these efforts. • Focus on traits that can be attained from natural history collections.
Recorded differently across sources	Body size for salamanders either SVL or TL depending on source	• Develop standards within a taxonomic group. • Carefully check your data.
Time consuming, difficult to automate	Three of these authors spent weeks compiling data	• More collaborative efforts. • Make trait tables open access and extendable by research community. • Incorporate individual variation using museum specimens and open access databases.
Deciding how to treat quantitative traits	Min/max attainable for the most species, but may not capture variation of interest	
Deciding how to treat categorical traits	Some ambiguity; variation across ranges for habitat type	
Range Data		
Taxonomy mismatches across databases	e.g., *Dryophytes*/*Hyla*, *Hyliola*/*Pseudacris*	• Check for synonyms manually. • Use 'taxize' R package[1]
Species without IUCN range maps	Newly described or split species, e.g., *Acris blanchardi*	• Revise manually. • Use occurrence records.
Introduced localities or errors in range maps or occurrence records	e.g., *Hyla cinerea* in Puerto Rico, *Rana pipiens* historical records throughout Mexico	• Check data; Revise range maps. • Use script to remove records outside of range maps.
Genetic Data		
Most sequences are not georeferenced	4,759 GenBank accessions linked to GBIF localities of 42,067 assembled (<12%)	• Link sequences to georeferenced voucher specimens in databases. • Include lat/lon when uploading.
Different genes sequenced, most mitochondrial	Fig. S1; 217 species represented by at least one gene; only 147 with cytb	• Standardize loci collected. • Identify comparable metrics. • Shift to genome-scale datasets.
Inconsistencies during GenBank upload	Different gene names (cytb, cytochrome b, Cyt-b, etc.)	• Use script to relabel names. • Standardize names in databases.
Data not uploaded to GenBank correctly (or at all)	"UNVERIFIED" sequences without genes annotated properly were missed	• Manually add data from other sources (e.g., Dryad). • Correct GenBank records.
Errors in automated alignments	Mis-aligned sequences identified, particularly when lengths differed	• Inspect alignments. • Test different software and optimize settings.
Small or uneven sample sizes across species	N individuals sampled was associated with diversity	• Check sensitivity to sampling. • Collect more data!

830   [1](Chamberlain & Szöcs, 2013). SVL = snout-vent length; TL = total length.

  Figure 1

834   **Figure 2**

    Figure 3

(a)

(b)