Large scale genomic analysis of 3067 SARS-CoV-2 genomes reveals a clonal geo-distribution and a rich genetic variations of hotspots mutations

Meriem LAAMARTI[1]ʃ, Tarek ALOUANE[1]ʃ, Souad KARTTI[1], M.W. CHEMAO-ELFIHRI[1], Mohammed HAKMI[1], Abdelomunim ESSABBAR[1], Mohamed LAAMART[1], Haitam HLALI[1], Loubna ALLAM[1], Naima EL HAFIDI[1], Rachid EL JAOUDI[1], Imane ALLALI[2], Nabila MARCHOUDI[3], Jamal FEKKAK[3], Houda BENRAHMA[4], Chakib NEJJARI[5], Saaid AMZAZI[2], Lahcen BELYAMANI[6] and Azeddine IBRAHIMI[1]*

[1] Medical Biotechnology Laboratory (MedBiotech), Bioinova Research Center, Rabat Medical & Pharmacy School, Mohammed Vth University in Rabat, Morocco

[2] Laboratory of Human Pathologies Biology, Department of Biology, Faculty of Sciences, and Genomic Center of Human Pathologies, Faculty of Medicine and Pharmacy, Mohammed V University in Rabat, Morocco.

[3] Anoual Laboratory of Radio-Immuno Analysis, Casablanca, Morocco.

[4] Faculty of Medicine, Mohammed VI University of Health Sciences (UM6SS), Casablanca, Morocco.

[5] International School of Public Health, Mohammed VI University of Health Sciences (UM6SS), Casablanca, Morocco.

[6] Emergency Department, Military Hospital Mohammed V, Rabat Medical & Pharmacy School, Mohammed Vth University in Rabat, Morocco.

* Corresponding author: a.ibrahimi@um5s.net.ma

ʃ These authors contributed equally to this work

Abstract

In late December 2019, an emerging viral infection COVID-19 was identified in Wuhan, China, and became a global pandemic. Characterization of the genetic variants of SARS-CoV-2 is crucial in following and evaluating their spread across countries. In this study, we collected and analyzed 3,067 SARS-CoV-2 genomes isolated from 59 countries during the first three months after the onset of this virus. Using comparative genomics analysis, we traced the profiles of the whole-genome mutations and compared the frequency of each mutation in the studied population. The accumulation of mutations during the epidemic period with their geographic locations was also monitored. The results showed 716 site mutations, of which 457 (64%) had a non-synonymous effect. Frequencies of mutated alleles revealed the presence of 39 recurrent non-synonymous mutations, including 10 hotspot mutations with a prevalence higher than 0.10 in this population and distributed in six genes of SARS-CoV-2. The distribution of these recurrent mutations on the world map revealed certain genotypes specific to the geographic location. We also found co-occurring mutations resulting in the presence of several haplotypes. Thus, evolution over time has shown a mechanism of co-accumulation and the phylogenetic analysis of this population indicated that this virus can be divided into 3 clades, including a subgroup-specific to the genomes of the United States. On the other hand, analysis of the selective pressure revealed the presence of several negatively selected residues that could be useful for considerations as therapeutic target design.

We have also created an inclusive unified database (http://moroccangenomes.ma/covid/) that lists all of the genetic variants of the SARS-CoV-2 genomes found in this study with phylogeographic analysis around the world.

Keywords: SARS-CoV-2, Hotspots mutations, Dissemination, Genomic analysis.

Introduction

The recent emergence of the novel, human pathogen Severe Acute Respiratory Syndrome Coronavirus 2 (SARS-CoV-2) in China with its rapid international spread poses a global health emergency. On March 11, 2020, World Health Organization (WHO) publicly announced the SARS-CoV-2 epidemic as a global pandemic. As of March 23, 2020, the COVID-19 pandemic had affected more than 190 countries and territories, with more than 464,142 confirmed cases and 21,100 deaths (1).

The new SARS-CoV-2 coronavirus is an enveloped positive-sense single-stranded RNA virus (2) and member of a large family named coronavirus which have been classified under three groups (2) two of them are responsible for infections in mammals (2), such us: bat SARS-CoV-like; Middle East respiratory syndrome coronavirus (MERS-CoV). Many recent studies have suggested that SARS-CoV-2 was diverged from bat SARS-CoV-like (3-4).

The size of the SARS-CoV2 genome is approximately 30 kb and its genomic structure has followed the characteristics of known genes of Coronavirus; the polyprotein ORF1ab also known as the polyprotein replicase covers more than 2 thirds of the total genome size, structural proteins, including spike protein, membrane protein, envelope protein and nucleocapsid protein. There are also six ORFs (ORF3a, ORF6, ORF7a, ORF7b, ORF8 and ORF10) predicted as hypothetical proteins with no associated function (5).

Characterization of viral mutations can provide valuable information for assessing the mechanisms linked to pathogenesis, immune evasion and viral drug resistance. In addition, viral mutation studies can be crucial for the design of new vaccines, antiviral drugs and diagnostic tests. A previous study (6) based on an analysis of 103 genomes of SARS-CoV-2 indicates that this virus has evolved into two main types. Type L being more widespread than type S, and type S representing the ancestral version. In addition, another study (7) conducted on 32 genomes of strains sampled from China, Thailand and the United States between December 24, 2019 and January 23, 2020 suggested increasing tree-like signals from 0 to 8.2%, 18.2% and 25,4% over time, which may indicate an increase in the genetic diversity of SARS-CoV-2 in human hosts.

Therefore, analyzing viral mutations and monitoring of the evolution capacity over time of SARS-CoV-2 in a large population is necessary. In this study, we characterized the genetic

variants in 3067 complete genomes of SARS-CoV-2 to assess the genetic diversity of

SARS-CoV-2 and to track mutations accumulation over time geographic location. On the

other hand, we established selective pressure analysis to predict negatively selected

residues which could be useful for the design of therapeutic targets. We have also created

a database containing the comparative genomic analysis of the SARS-CoV-2 genomes, in

order to facilitate comparison and research for the COVID-19 scientific community.

Materials and Methods

Data collection and Variant calling analysis

3067 sequences of SARS-CoV-2 were collected from the GISAID EpiCovTM (update: 02-

04-2020) and NCBI (update: 20-03-2020) databases. Only complete genomes were used

in this study. **Table S1** illustrating detailed information on the genomes downloaded.

Genomes were mapped to the reference sequence Wuhan-Hu-1/2019 using Minimap

v2.12-r847-dirty (8).The BAM files were sorted by SAMtools sort (9). The final sorted

BAM files were used to call the genetic variants in variant call format (VCF) by SAMtools

mpileup (9) and bcftools (9). The final call set of the 3067 genomes, was annotated and

their impact was predicted using SnpEff v4.3t (10). First, the SnpEff databases were built

locally using annotations of the reference genome NC_045512.2 obtained in GFF format

from NCBI database. Then, the SnpEff database was used to annotate SNPs and InDels

with putative functional effects according to the categories defined in the SnpEff manual

(http://snpeff.sourceforge.net/SnpEff_manual.html).

Phylogentic analysis and geo-distribution

The downloaded full-length genome sequences of coronaviruses isolated from different

hosts from public databases were subjected to multiple sequence alignments using Muscle

v 3.8 (11). Maximum-likelihood phylogenetic trees with 1000 bootstrap replicates were

constructed using RaxML v 8.2.12 (39), Tree was done using figtree v 1.4.4(12)

Selective pressure and modelling

We used Hyphy v2.5.8 (13) to estimate synonymous and non-synonymous ratio dN / dS

(ω). Two datasets of 191 and 433 for orf1ab and genes respectively were retrieved from

Genbank (http://www.ncbi.nlm.nih.gov/genbank/). After deletion of duplicated and

cleaning the sequences, only 92 and 35 for orf1ab and spike proteins, respectively, were

used for the analysis. The selected nucleotide sequences of each dataset were aligned using

Clustalw codon-by-codon and the phylogenetic tree was obtained using ML (maximum

likelihood) available in MEGA X (14). For this analysis, four Hyphy's methods were used

to study site-specific selection: SLAC (Single-Likelihood Ancestor Counting (15), FEL

(Fixed Effects Likelihood) (15) , FUBAR (Fast, Unconstrained Bayesian AppRoximation)

(16) and MEME (Mixed Effects Model of Evolution) (17). For all the methods, values

supplied by default were used for statistical confirmation and the overall ω value was

calculated according to ML trees under General Time Reversible Model (GTR model). The

CI- TASSER generated models (https: //zhanglab.ccmb.med.umich.edu / COVID-19 /) of

ORF1ab nonstructural proteins (NSP) 3,4,6,12,13,14 and 16 were used to highlight the

sites under selective pressure on the protein, due to the lack of their experimental structures

and the absence of suitable templates for homology modelling. On the other hand, the cryo-

EM structure with PDB id 6VSB was used as a model for the gene S in its prefusion

conformation. Structure visualization and image rendering were performed in PyMOL 2.3

(Schrodinger LLC).

Pangenome construction

115 proteomes of the genus *Betacorononavirus* were obtained from the NCBI database

(update: 20-03-2020), of which 83 genomes belonged to the SARS-CoV-2 species and the

rest distributed to other species of the same genus publicly available. These proteomes were

used for the construction of pangenome at the inter-specific scale of *Betacoronavirus* and

intra-genomic of SARS-CoV-2. The strategy of best reciprocal BLAST results (18) was

implemented to identify all of the orthologous genes using Proteinortho v6.0b (19).

Proteins with an identity above 60% and sequence coverage above 75% with an e-value

threshold below 1e-5 were used to be considered as significant hits.

Results

SARS-CoV-2 genomes used in this study

In this study, we used 3,067 SARS-CoV2 complete genomes collected from GISAID

EpiCovTM (update: 02-04-2020) and NCBI (update: 20-03-2020) databases. These strains

were isolated from 59 countries (**Fig 1A**). The most represented origin was American

strains with 783 (25.53%), followed by strains from England, Iceland, and China with 407

(13.27%), 343 (11.18%), 329 (10.73%), respectively. The date of isolation was during the

first three months after the appearance of the SARS-CoV-2 virus, from December 24, 2019,

to March 25, 2020 (**Fig 1B**). Likewise, about two-thirds of these strains collected in this

work were isolated during March.

Allele frequencies revealed a diversity of genetic variants in six SARS-Cov-2 genes

To study and follow the appearance and accumulation of mutations, we traced the profiles

of non-synonymous mutations, and we compared their frequency in the studied population.

Remarkably, compared to the reference sequence Wuhan-Hu-1/2019, a total of 716

mutations were identified, including 457 (64%) non-synonymous and 188 (26%)

synonymous mutations. The rest (10%) distributed to the intergenic regions. Focusing on

non-synonymous mutations, the analysis of the frequencies of mutated alleles revealed the

presence of 39 mutations with a prevalence greater than 0.006 (0.06% of the population),

which corresponds to at least 20 / 3,067 SARS-CoV-2 genomes (**Fig 2**).

These recurring mutations were distributed in six genes with variable frequencies, of which

the gene coding for replicase polyprotein (ORF1ab), spike protein, membrane

glycoprotein, nucleocapsid phosphoprotein, ORF3a, and ORF8. Overall, ORF1ab harbored

more non-synonymous mutations compared to the other five genes with 22 mutations,

including three mutations located in NSP12-RNA- dependent RNA polymerase (M4555T,

T4847I and T5020I), three in NSP13-helicase (V5661A, P5703L and M5865V), two in

NSP5-main proteinase (G3278S and K3353R), two in NSP15-EndoRNAse (I6525T,

Ter6668W), two in NSP3-multi domains (A876T and T1246I), one in NSP14-exonuclease

(S5932F) and one in NSP4-transmembrane domain 2 (F3071Y). Likewise, spike protein

harbored three frequent mutations, including V483A in receptor-binding domain (RBD).

The rest of the mutations were found in nucleocapsid phosphoprotein (S193I, S194L,

S197L, S202N, R203K and G204R), ORF3a (S193I, S194L, S197L, S202N, R203K and

G204R), membrane glycoprotein (D3G and T175M) and ORF8 (V62L and L84S).

Identification of ten hyper-variable genomic hotspot in SARS-CoV-2 genomes

Interestingly, among all recurrent mutations, ten were found as hotspot mutations with a

frequency greater than 0.10 in this study population (**Fig 2**). The most represented was

D614G mutation at spike protein with 43.46% (n = 1.333) of the genomes, the second was

L84S at ORF8 found in 23.21% (n = 712). Thus, the gene coding for orf1ab had four

mutations hotspots, including S5932F of NSP14-exonuclease, M5865V of NSP13-helicase

L3606F of NSP6- transmembrane domain and T265I of NSP2 found with 17.02%, 16.56%,

14.38% and 10.66% of the total genomes, respectively. For the four other hotspot mutations

were distributed in ORF3a (Q57H and G251V) and nucleocapsid phosphoprotein (R203K

and G204R).

Geographical distribution and origin of mutations worldwide

Out of 3067 genomes, only 105 were found to be wild type and were particularly of Chinese

origin. However, mutant strains were dispersed in different countries and represented up

to 2962 strains with different genotype profiles. We performed a geo-referencing mutation

analysis to identify region-specific loci. Remarkably, USA was the country with the highest

number of mutations, with 316 (44% of the total number of mutations), including 173

(24%) singleton mutation (specific to USA genomes). While the Chinese genomes

containing 22% of the total number of mutations, followed by France and New Zealand

with 4% and 2%, respectively. It is interesting to note that among the 59 countries, 26

harbored singleton mutations. **Table S2** illustrates the detailed singleton mutations found

in these countries.   The majority of the genomes analyzed carried more than one mutation.

However, among the recurrent non-synonymous, synonymous and intergenic mutations

(**Fig 3**). We found G251V (in ORF3a), L84S (in ORF8) and S5932F (in ORF1ab) present

on all continents except Africa and Austria (**Fig 4**). While F924F, L4715L (in orf1ab),

D614G (in spike) and an intergenic variant at position 241 appeared in all strains except

those from Asia. In Algeria, the genomes harbored mutations very similar to those in

Europe, including two recurrent mutations T265I and Q57H of the ORF3a. Likewise, the

European and Dutch genomes also shared ten recurrent mutations. On the other hand,

continent-specific mutations have also been observed, for example in America, we found

seven mutations shared in almost all genomes. In addition, two mutations at positions

28117 and 28144 were shared by the Asian genomes, while four different positions 1059,

14408, 23403, 25563 and 1397, 11083, 28674, 29742 were shared by African and

Australian genomes (Supplementary material). The majority of these mutations are

considered to be transition mutations with a high ratio of A substituted by G. The genome

variability was more visible in Australia, New Zealand and America than in the rest of the

world.

Evolution of mutations over time

We selected the genomes of the SARS-CoV-2 virus during the first three months after the

appearance of this virus (December 24 to March 25). We have noticed that the mutations

have accumulated at a relatively constant rate (**Fig 5**). The strains selected at the end of

March showed a slight increase in the accumulation of mutations with an average of 11.34

mutations per genome, compared to the gnomes of February, December and January with

an average number of mutations of 9.26, 10.59 10.34 respectively. The linear curve in

Figure 5 suggests a continuous accumulation of SNPs in the SARS-CoV-2 genomes in the

coming months. This pointed out that many countries had multiple entries for this virus

that could be claimed. Thus in the deduced network demonstrated transmission routes in

different countries.

The study of the accumulation of mutations over time showed a higher number of

mutations in the middle of the outbreak (end of January). At the same time, an increase in

the number of mutations in early April was also observed. The first mutations to appear

were mainly located in the intergenic region linked to the nucleocapsid phosphoprotein and

the orf8 protein. The T265I, D614G and L84S hotspot mutations located in orf1ab and

Spike proteins respectively were introduced into the virus for the first time in late February.

Phylogeographical analysis of SARS-CoV-2 genomes

The phylogenetic tree based on the whole genome alignment showed the presence of 3

different clades (**Fig S1**) and demonstrate that SARS-CoV-2 is wildly disseminated across

distinct geographical location. The results showed that several strains are closely related

even though they belong to different countries. Which indicate likely transfer events and

identify   routes   for   geographical   dissemination.   For   phylogenetic   tree

(http://moroccangenomes.ma/covid/) showed multiple introduction dates of the virus

inside the USA with the first haplotype introduced related to the second epidemic wave in

china.

Selective pressure analysis

Selective pressure on ORF1ab, gene harbored a high rate of mutations and on the Spike

gene, indicated a single alignment-wide ω ratio of 0.571391 and 0.75951 for spike and

or1ab, respectively. Most sites for both genes had ω <1 values, indicating purifying
selection. We estimated eight sites under negative selection pressure (696, 1171, 2923,
3003, 3715, 5221, 5704 and 6267) and three sites under positive selection pressure (1473,
2244 and 3090) in ORF1ab. For spike, we found seven sites under negative selection
pressure (215, 474, 541, 809, 820, 921 and 1044), while only site 5 was found under
negative selection pressure (**Table 1**).

The modelling results of ORF1ab showed that the sites with positive selections were
distributed in NSP3 and NSP4. While the negatively selected codons were located in NSP3,
NSP4, NSP6, NSP12, NSP13, NSP14 and NSP16 (**Fig 6**). Thus, only one negatively
selected amino acid residue was observed on the receptor binding domain, suggesting that
there was no strong selective pressure in the region (**Fig 7**).

Inter and intra-specific pan-genome analysis
In order to highlight the structural proteins shared at the inter-specific scale between the
isolates of the genus *Betacoronavirus*, thus at the intra-genomic scale of SARS-CoV-2, we
have constructed a pan-genome by clustering the sets of proteins encoded in 115 genomes
available publicly in NCBI (update: 20-03-2020), including 83 genomes of SARS-Cov-2
and the rest distributed to other species of the same genus. Overall, a total of 1,190 proteins
were grouped into a pangenome of 94 orthologous cluster proteins, of which ten proteins
cluster were shared between SARS-CoV-2 and only three species of the genus
Betacoronavirus (BatCoV RaTG13, SARS-CoV and Bat Hp-betacoronavirus/
Zhejiang2013). Of these, BatCoV RaTG13 had more orthologous proteins shared with
SARS-CoV-2, followed by SARS-CoV with ten and nine orthologous proteins,
respectively (**Fig 8A**).

It is interesting to note that among all the strains used of *BetaCoronavirus*, the ORF8
protein was found in orthology only between SARS-RATG13 and SARS-CoV-2. In
addition, the ORF10 protein was found as a singleton for SARS-CoV-2. While, the analysis
of the pangenome at the intra-genomic scale of 83 isolates of SARS-CoV-2 (**Fig 8B**),
showed that ORF7b and ORF10 were two accessory proteins (proteins variable) in SARS-
CoV- 2 genomes, while the other proteins belonged to the core proteins of SARS-CoV-2
(conserved in all genomes).

Discussion

The rate of mutations results in viral evolution and variability in the genome, thus allowing

viruses to escape host immunity, as well as drugs (20). Initial published data suggests that

SARS-CoV-2 is genetically stable (21) which may increase the effectiveness of vaccines

under development. The study on the genomic variation of SARS- CoV- 2 is very

important for the investigation of pathogenesis, disease course, prevention, and treatment

of SARS- CoV- 2 infection. In this study, we characterized the genetic variations in a large

population of SARS-CoV-2 genomes (3,067). Our results showed a diversity of mutations

detected with different frequencies. Overall, more than 450 non-synonymous mutations in

SARS-CoV-2 genomes have been identified. The orf1ab gene contains 16 non-structural

proteins (NSP1-NSP16) (22), and harbored more than 50 % of the frequent mutations, with

a high mutation number 117, 61, and 61 in the NSP3, NSP12, and NSP2 Respectively.

Both NSP2 and NSP3 were reported to be essential for correcting virus replication errors

(23). On the other, hand recent studies have suggested that mutations falling in the

endosome- associated- protein- like domain of the NSP2 protein, could explain why this

virus is more contagious than SARS (24). The Replication enzymes NSP12 to NSP16 have

been reported as antiviral targets for SARS-CoV (25). In the SARS-CoV-2 genomes, we

found that NSP12 to NSP15 harbored nine recurrent non-synonymous mutations. Among

them, eight identified as new mutations, including three (M4555T, T4847I, T5020I) in

NSP12-RNA-dependent RNA polymerase, three (V5661A, P5703L and M5865V) in

NSP13-Helicase, two (I6525T, Ter6668W) in NSP15-EndoRNAse. However, these new

mutations must be taken into account when developing a vaccine using the ORF1ab protein

sequences as a therapeutic target.

A high number of mutations were identified in the spike protein, an important determinant

in pathogenicity that allows the virion attachment to the cell membrane by interacting with

the host ACE2 receptor (26). Among all the frequent mutations in this protein, the V483A

mutation has been identified in this receptor and found mainly in SARS-CoV-2 genomes

isolated from USA. This result is consistent with the study of Junxian et al. (27). Eight

stains from china, USA and France harbored V367F mutation previously described to

enhance the affinity with ACE2 receptor (27).

Among the 716 mutations, ten were considered hypervariable genomic hotspots with high
frequencies of mutated alleles such us the L3606F mutation detected in NSP6. The NSP6
protein works with NSP3 and NSP4 by forming double-membrane vesicles and convoluted
membranes involved in viral replication (28). Besides, three previously reported mutations
(M5865V, S5932F) and (R203K) were identified in ORF1ab and N respectively (20).
Moreover, intraspecies pangenome analysis of SARS- CoV-2 showed that the six of the
genes harboring hotspot mutations belong to the core genome.

Thus, under normal circumstances genomic variation increase the viruses spread and
pathogenicity. This happens when the virus accumulated mutation enables its virulence
potential (29). Genomic comparison of the studied population allowed us to gain insights
into virus mutations occurrence over time and within different geographic areas. In the
SARS-CoV virus, the SNP distribution is not random, and it is more dominant in critical
genes for the virus. As it was already found in many studies (20,30). Our results confirmed
what was previously described and elucidate the presence of numerous hotspot mutations.
Besides, Co-occurrence mutations were also common in different countries all along with
singleton mutations. In the case of the USA, the singleton mutations are driven by the single
group that diverged differently due to the environment, the host, and the number of
generations. These mutations are due to the low fidelity of reverse transcriptase (29, 31).

Spain, Italy, and the US contain a high number of specific mutations which may be the
cause of a rapid transmission, especially in the US. These specific mutations may also be
correlated with the critical condition in Italy and Spain.

In this study, we have defined 3 major clades. The clustering of these genomes revealed
the spread of clades to diverse geographical regions. We observed an increase of mutations
over time following the first dissemination event from China. Specific haplotypes were
also predominant to a geographical location, especially in the US. This study opens up new
perspectives to determine whether one of these frequent mutations will lead to biological
differences and their correlation with different mortality rates.

Among the 7 proteins of orf1ab that harbored sites under selective pressure, only NSP3
and NSP4 contains both residues under positive and negative selection. The modelling of
NSP3 domains shows that only the negative selection site 1171 (Thr- 353), was located at

the conserved macro domain Mac1 (previously X or ADP-ribose 1" phosphatase) (32).
This domain has been previously shown to be dispensable for RNA replication in the
context of a SARS-CoV replicon (33). However, it could counteract the host's innate
immune response (34). It was proposed that the 3Ecto luminal domain of NSP3 interacts
with the large luminal domain of NSP4 (residues 112-164) to "zip" the ER membrane and
induce discrete membrane formations as an important step in the generation of ER viral
replication organelles (35, 36). As we have shown previously by the FEL, MEME and
FUBAR methods, the orf1ab 2244 site coding for ILE-1426 is under positive selection
pressure and since it is located on the luminal 3ecto domain of the NSP3 protein, this can
be explained by a possible host influence on the virus in this domain.The results of selective
pressure analysis revealed the presence of several negatively selected residues, one of
which is located at the receptor-binding domain (GLN-474) and which is known by its
interaction with the GLN24 residue of the human ACE2 (Angiotensin-converting enzyme
2) receptor (37). In general, it is well-known that negatively selected sites could indicate a
functional constraint and could be useful for drug or vaccine target design, given their
conserved nature and therefore less likely to change (38).

Conclusion
The SARS-CoV-2 pandemic has caused a very large impact on health and economy
worldwide. Therefore, understanding genetic diversity and virus evolution become a
priority in the fight against the disease. Our results show several molecular facets of the
relevance of this virus. We identified ten non-synonymous hotspot mutations distributed
in six of the virus genes with high frequencies of mutated alleles. We also were able to
highlight an increase in mutations accumulation overtime and revealed the existence of
three major clades in various geographic regions. Finally, the study allowed us to identify
specific haplotypes by geographic location and provides a list of sites under selective
pressure that could serve as an interesting avenue for future studies.

Conflict of interest

The authors declare that they have no competing interests.

Acknowledgments

We sincerely thank the authors and laboratories around the world who have sequenced and

shared the full genome data for SARS-CoV-2 in the GISAID database. All data authors

can be contacted directly via www.gisaid.org

This work was carried out under National Funding from the Moroccan Ministry of Higher

Education and Scientific Research (PPR program) to AI. This work was also supported, by

a grant to AI from Institute of Cancer Research of the foundation Lalla Salma.

References

1. World Health Organization. Infection prevention and control during health care
when COVID-19 is suspected: interim guidance, 19 March 2020. World Health
Organization. 2020. Available from:
https://apps.who.int/iris/handle/10665/331495
2. Enjuanes LD. Cavanagh K. Holmes MMC, Lai H. Laude P. Masters P et al.
(2000) Coronaviridae. In: Virus taxonomy. Classification and nomemclature
of viruses (M. H. V. van Regenmortel, C. M. Fauquet, D. H. L. Bishop, E. B.
Carstens, M. K. Estes, S. M. Lemon, J. Maniloff, M. A. Mayo, D. J. McGeoch,
C. R. Pringle, and R. B. Wickner eds.) Academic Press, San Diego. pp 835-
849.
3. Yeşilbağ K, Aytoğu G. Coronavirus host divergence and novel coronavirus
(Sars-CoV-2) outbreak. Clinical and Experimental Ocular Trauma and
Infection. 2020 Apr 23;2(1):1-9.
4. Andersen KG, Rambaut A, Lipkin WI, Holmes EC, Garry RF. The proximal
origin of SARS-CoV-2. Nat Med. 2020;26: 450–452. DOI:10.1038/s41591-
020-0820-9

5. Wu F, Zhao S, Yu B, Chen YM, Wang W, Song ZG, et al. A new coronavirus
associated with human respiratory disease in China. Nature. 2020;579:265-269.

6. Tang X, Wu C, Li X, Song Y, Yao X, Wu X, et al. On the origin and continuing
evolution of SARS-CoV-2. Natl Sci Rev. 2020. DOI: 10.1093/nsr/nwaa036.

7. Li LQ, Huang T, Wang YQ, Wang ZP, Liang Y, Huang TB, et al. COVID-19
patients' clinical characteristics, discharge rate, and fatality rate of meta-
analysis. J Med Virol. 2020 Mar 12. Doi: 10.1002/jmv.25757.

8. Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics.
2018;34: 3094–3100 .DOI: 10.1093/bioinformatics/bty191.

9. Li H, Hansaker B, Wysoker A, Fennell T, Ruan J, Homer N, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25: 2078–9.
DOI:10.1093/bioinformatics/btp352

10. Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, et al. A program
for annotating and predicting the effects of single nucleotide polymorphisms,
SnpEff: SNPs in the genome of Drosophila melanogaster strain w1118; iso-2;
iso-3. Fly (Austin). 2012;6: 80-92 .DOI: 10.4161/fly.1969

11. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and
high throughput. Nucleic Acids Re. 2004;32: 1792–1797. DOI:
10.1093/nar/gkh340

12. Rambaut, Andrew. "FigTree v1. 4." (2012).

13. Pond SL, Frost SD, Muse SV. HyPhy: hypothesis testing using phylogenies.
Bioinformatics. 2005;21: 676–679. DOI: 10.1093/bioinformatics/bti079

14. Kumar S, Stecher G, Li M, Knyaz C, Tamura. MEGA X: Molecular
Evolutionary Genetics Analysis across Computing Platforms. Mol Biol Evol.
2018;35: 1547-1549. DOI: 10.1093/molbev/msy096.

15. Pond SL, Frost SD. Not so different after all: a comparison of methods for
detecting amino-acid sites under selection. Mol Biol Evol. 2005;22: 1208-1222.
Doi:10.1093/molbev/msi105.

16. Murrell B, Moola S, Mabona A, Weighill T, Sheward D, Pond SL, et al. Fubar:
a fast, unconstrained Bayesian approximation for inferring selection. Mol Biol
Evol. 2013;30: 1196-1205. DOI: 10.1093/molbev/mst030

17. Murrell B, Wertheim JO, Moola S, Weighill T, Scheffler K, Pond SL. Detecting
Individual Sites Subject to Episodic Diversifying Selection. PLoS Genet.
2012;8: e1002764. Doi: 10.1371/journal.pgen.1002764

18. Jordan IK, Rogozin IB, Wolf YI, Koonin EV. Essential genes are more
evolutionarily conserved than are nonessential genes in bacteria. Genome Res.
2002;12:962–968.

19. Lechner M, Findeiß S, Steiner L, Marz M, Stadler PF, Prohaska SJ.
Proteinortho: detection of (Co-)orthologs in large-scale analysis. BMC
Bioinformatics. 2011;12:1–9. DOI:10.1186/1471-2105-12-124.

20. Pachetti M, Marini B, Benedetti F, Giudici F, Mauro E, Storici P, et al.
Emerging SARS-CoV-2 Mutation Hot Spots Include a Novel RNA-dependent-
RNA Polymerase Variant. J Transl Med. 2020;18: 179. DOI: 10.1186/s12967-
020-02344-6.

21. Su Y, Anderson D, Young B, Zhu F, Linster M, Kalimuddin S, et al. Discovery
of a 382-nt deletion during the early evolution of SARS-CoV-2. BioRxiv
[Preprint]. 2020 [cited 2020 March 25]. Available from:
https://www.biorxiv.org/content/10.1101/2020.03.11.987222v1

22. Khailany RA, Safdar M, Ozaslan M. Genomic characterization of a novel
SARS-CoV-2.   Gene   Rep.   2020;19:100682.
DOI:10.1016/j.genrep.2020.100682

23. Harcourt BH, Jukneliene D, Kanjanahaluethai A, Bechill J, Severson KM,
Smith CM, et al. Identification of severe acute respiratory syndrome
coronavirus replicase products and characterization of papain-like protease
activity. J Virol. 2004;78:13600-13612. DOI:10.1128/JVI.78.24.13600-
13612.2004

24. Angeletti S, Benvenuto D, Bianchi M, Giovanetti M, Pascarella S, Ciccozzi
M. . COVID 2019: the role of the NSP2 and NSP3 in its pathogenesis. Journal
of Medical Virology, 2020;92:584-588.

25. Subissi L, Imbert I, Ferron F, Collet A, Coutard B, Decroly E, et al. SARS-
CoV ORF1b-encoded nonstructural proteins 12-16: replicative enzymes as

antiviral targets. Antiviral Res. 2014;101:122-30. DOI: 10.1016/j.antiviral.2013.11.006.

26. Hoffmann, Markus, et al. SARS-CoV-2 cell entry depends on ACE2 and TMPRSS2 and is blocked by a clinically proven protease inhibitor.*Cell* (2020).

27. Ou J, Zhou Z, Dai R, Zhang J, Lan W, Zhao S. Emergence of RBD mutations from circulating SARS-CoV-2 strains with enhanced structural stability and higher human ACE2 receptor affinity of the spike protein. bioRxiv 2020.03.15.991844; doi: https://doi.org/10.1101/2020.03.15.991844

28. Angelini MM, Akhlaghpour M, Neuman BW, Buchmeier MJ. Severe acute respiratory syndrome coronavirus nonstructural proteins 3, 4, and 6 induce double-membrane vesicles. mBio. 2013 Aug 13. pii: e00524-13. DOI: 10.1128/mBio.00524-13.

29. Stern A, Yeh MT, Zinger T, et al. The Evolutionary Pathway to Virulence of an RNA Virus. Cell. 2017;169(1):35- 46.e19. doi:10.1016/j.cell.2017.03.013.

30. Wen F, Yu H, Guo J, Li Y, Luo K, Huang S. Identification of the hyper-variable genomic hotspot for the novel coronavirus SARS-CoV-2. J Infect. 2020. pii: S0163-4453(20)30108-0. DOI: 10.1016/j.jinf.2020.02.027.

31. Denison MR, Graham RL, Donaldson EF, Eckerle LD, Baric RS. Coronaviruses: an RNA proofreading machine regulates replication fidelity and diversity. RNA Biol. 2011;8: 270–279. DOI: 10.4161/rna.8.2.15013.

32. Neuman BW. Bioinformatics and functional analyses of coronavirus nonstructural proteins involved in the formation of replicative organelles. Antiviral Res. 2016;135: 97-107. DOI: 10.1016/j.antiviral.2016.10.005.

33. Kusov Y, Tan J, Enrique A, Luis E, Hilgenfeld R. A G-quadruplex-binding macrodomain within the "SARS-unique domain" is essential for the activity of the SARS-coronavirus replication-transcription complex. Virology. 2015;484: 313–322. DOI: 10.1016/j.virol.2015.06.016.

34. Fehr AR, Channappanavar R, Jankevicius G, Fett C, Zhao J, Athmer J. The Conserved Coronavirus Macrodomain Promotes Virulence and Suppresses the Innate Immune Response during Severe Acute Respiratory Syndrome

Coronavirus Infection. mBio. 2016 Dec 13. pii: e01721-16. DOI: 10.1128/mBio.01721-16.

35. Hagemeijer MC, Monastyrska I, Griffith J, van der Sluijs P, Voortman J, en Henegouwen PM, et al. Membrane rearrangements mediated by coronavirus nonstructural proteins 3 and 4. Virology. 2014;458: 125-135. DOI: 10.1016/j.virol.2014.04.027.

36. Qiu Y, Xu K. Functional studies of the coronavirus nonstructural proteins. STEMedicine. 2020;1:e39. DOI: 10.37175/stemedicine.v1i2.39.

37. Yan R, Zhang Y, Li Y, Xia L, Guo Y, Zhou Q. Structural basis for the recognition of SARS-CoV-2 by full-length human ACE2. Science. 2020;367:1444-1448. DOI: 10.1126/science.abb2762.

38. Mazumder R, Hu ZZ, Vinayaka CR, Sagripanti JL, Frost SD, Pond SL, et al. Computational analysis and identification of amino acid sites in dengue E proteins relevant to development of diagnostics and vaccines. Virus Genes. 2007;35:175-186. DOI: 10.1007/s11262-007-0103-2.

39. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. Bioinformatics. 2014;30: 1312–1313. Doi: 10.1093/bioinformatics/btu033.

Figure 1: Distribution of the genomes of the 3,067 genomes used in this study by county and date of isolation. **A)** The pie chart represents the percentage of genomes used in this study according to their geographic origins. The colors indicate different countries. **B)** Number of genomes of complete pathogens, distributed over a period of 3 months from the end of December to the end of March.

Figure 2: The linear diagrams represent genes distribution in the SARS-CoV-2 genome. Diagrams in garnet and brown represent the protein subunits of ORF1ab and S respectively. The presence of a mutation is represented by vertical lines. under each line, the most frequent variants are annotated as the amino-acid change at that specific site, and the frequency of mutations is presented by color-coded circles.

528
529
530 **Figure 3: Map showing Geographical distribution of hotspot mutation in the studied**
531 **population worldwide.** The pie charts show the relative frequencies of haplotype for each
532 population. The haplotypes are color coded as shown in the key.

547
548 **Figure 4 : The graph represents substitutions accumulation in a three months period**.
549 The accumulation of mutations increases linearly with time. The dots represent the number of
550 mutations in a single genome. All substitutions were included non-synonymous, synonymous,
551 intergenic.

564
Figure 5 : Lollipop plots showing mutations distribution and frequency and time of
the appearance in SARS-CoV-2 genome. A) The presence of a mutation is shown on the x-
axis (lollipop), and the frequency of mutations is shown on the y-axis and correlates with the heights
of the vertical lines representing each lollipop. Non-synonymous, synonymous, STOP Codon and
intergenic mutations are presented as green, blue, red, and pink circles respectively. Predicted
amino acid change has been represented for each hotspot mutation at the top of the circle. **B)** The
diagram represents genes encoded in SARS-CoV-genome. The mutations are represented by a
color-coded vertical line, each color represents a time period corresponding to the date of the first
appearance of the mutation.

Figure6 : Structural view of selective pressure in orf1ab gene. The residue under the positive and negative selection is highlighted in blue and red respectively. The modeling of orf1ab non-structural proteins (NSP3, NSP4, NSP6, NSP12, NSP13, NSP14, and NSP16) harboring residues under pressure selection was produced using CI-TASSER. **A**. The NSP3 domains MAC1, Ubl1, Ubl2-PLpro, and SUD-C are color-coded in the 3D representation. The residues Ile-1426 and Ala-655 under negative selection are located respectively on 3Eco and SUD-C domains while Thr-353 residue under positive selection is shown on the MAC1 domain, **B**. 3D representation of the NSP4 protein, **C**. 3D representation of the NSP6 protein, **D**. 3D representation of the NSP12 protein, **E**. 3D representation of the NSP13 protein, **F**. 3D representation of the NSP14 protein, **G**. 3D representation of the NSP16 protein.

629 **Figure 7:** S**tructural view of selective pressure in spike gene**. The negatively selected site
630 in spike protein is highlighted in red. The only amino acid residue selected negatively on the
631 receptor-binding domain corresponds to GLN-474. The cryo-EM structure with PDB id 6VSB was
632 used as a model for the gene S in its prefusion conformation.

Figure 8 : Pangenome analysis of 32 from different *Betacoronavirus* species and 83 of SARS-CoV-2. (**A**) The Venn diagram represents the number of core, accessory, and unique proteins inside the *Betacoronavirus* genus. (**B**) The pie chart illustrates the core and accessory protein inside the SARS-CoV-2 specie.

Table 1 : Selective pressure analysis on the spike and orf1ab genes of SARS-CoV-2

Genes	ω	FEL method		MEME method	SLAC method		FUBAR method	
		PS	NS	PS	PS	NS	PS	NS
Spike	0.571391	-	Codons 215, 474, 809, 820, 921,	-	-	-	Codon 5	Codons 215, 474, 541,
		PS	NS	PS	PS	NS	PS	NS
orf1ab	0.75951	Codon 2244	Codons 1171, 2923, 3003, 3715, 5221, 5704, 6267, 6961	Codon2244	-	-	Codons 1473, 2244, 3090	-

653