1
9 # The LINC complex transmits integrin-dependent tension to the nuclear lamina and represses epidermal differentiation
10
11
12

13 Emma Carley[1]*, Rachel K. Stewart[1]*, Abigail Zieman[2]*, Iman Jalilian[1], Diane. E. King[3], Amanda
14 Zubek[4], Samantha Lin[2], Valerie Horsley[2,4, +], and Megan C. King[1,2, +]
15
16 [1] Department of Cell Biology, Yale School of Medicine
17 [2] Department of Molecular, Cell and Developmental Biology, Yale University
18 [3] Sunnycrest Bioinformatics
19 [4] Department of Dermatology, Yale School of Medicine
20
21 * These authors contributed equally to this work
22
23 + Co-corresponding authors: valerie.horsley@yale.edu or megan.king@yale.edu

Abstract:

While the mechanisms by which chemical signals control cell fate have been well studied, how mechanical inputs impact cell fate decisions are not well understood. Here, using the well-defined system of keratinocyte differentiation in the skin, we examine whether and how direct force transmission to the nucleus regulates epidermal cell fate. Using a molecular biosensor, we find that tension on the nucleus through Linker of Nucleoskeleton and Cytoskeleton (LINC) complexes requires integrin engagement in undifferentiated epidermal stem cells, and is released during differentiation concomitant with decreased tension on A-type lamins. LINC complex ablation in mice reveals that LINC complexes are required to repress epidermal differentiation *in vivo* and *in vitro* and influence accessibility of epidermal differentiation genes, suggesting that force transduction from engaged integrins to the nucleus plays a role in maintaining keratinocyte progenitors. This work reveals a direct mechanotransduction pathway capable of relaying adhesion-specific signals to regulate cell fate.

Introduction:

Physical forces and the architecture of the extracellular environment are an emerging area of cell fate regulation (Discher et al., 2009). Cells sense changes in the physical environment through cell-cell and cell-extracellular matrix (ECM) adhesions, providing a mechanism by which mechanical inputs can contribute to the control of cell fate. While several studies have linked geometrical and physical inputs to changes in cellular signaling through mechanoresponsive transcription factors such as YAP/TAZ (Totaro et al., 2018) or MKL/MRTF (Connelly et al., 2010), whether direct transmission of mechanical force to the nucleus can influence cell fate is not known.

The interfollicular epidermis is an excellent model to explore how mechanical inputs regulate cell fate. Epidermal stem cells adhere to the underlying basal lamina through β1 integrin-based adhesions and β4 integrin-based hemidesmosomes (Raghavan et al., 2000), which maintain stem cell fate, proliferation, and inhibit differentiation (Hotchin et al., 1995; Levy et al., 2000; Rippa et al., 2013; Watt et al., 1993). Upon differentiation, basal keratinocytes release their integrin-based adhesions, initiate gene expression changes to activate terminal differentiation (Tsuruta et al., 2011), and move upward to enter the stratified epithelium until they are shed at the skin's surface. Despite the well-established role for integrin adhesions in regulating epidermal differentiation, how integrin signals are propagated to the nucleus to regulate the keratinocyte differentiation program is not well understood.

Here, we explore whether mechanical cues can regulate keratinocyte cell fate via tension on the nucleus via the Linker of Nucleoskeleton and Cytoskeleton (LINC) complex. The LINC complex, composed of tail-anchored Nesprins integrated into the outer nuclear membrane and SUN proteins integrated into the inner nuclear membrane, spans the nuclear envelope to mechanically integrate the cytoplasmic cytoskeleton and the nuclear interior – specifically the nuclear lamina and its associated chromatin (Chang et al., 2015). While the LINC complex has been postulated to act either as a direct mechanosensor or as a conduit for mechanotransduction, to influence gene expression (Alam et al., 2016; Wang et al., 2009), whether the LINC complex controls gene expression *in vivo* remains largely untested. Indeed, to date direct gene targets of the LINC complex that regulate genetic programs *in vivo*, for example during differentiation, remain to be identified.

Several lines of evidence suggest that epidermal differentiation is regulated by the nuclear lamina, yet the mechanisms remain unknown. First, during basal stem cell differentiation, a large chromosomal region termed the epidermal differentiation complex (EDC), which consists of sixty consecutive genes necessary for epidermal stratification and the production of the cornified envelope, relocates away from the nuclear lamina towards the nuclear interior to be transcriptionally activated (Gdula et al., 2013; Mardaryev et al., 2014; Williams et al., 2002)). Further, the AP-1 transcription factor complex, itself regulated by A-type lamins, influences EDC gene expression in both proliferating and differentiating keratinocytes *in vitro* (Oh et al., 2014) and coordinates with EZH2 in the polycomb complex to regulate epidermal differentiation (Ezhkova et al., 2009). Lastly, a skin-specific lamin-null mouse model (Lamin B1/B2/A/C triple-

94 knockout) exhibits a thickened epidermis attributed to precocious differentiation (Jung et al.,
95 2014).
96
Here, we provide evidence that the LINC complex regulates epidermal differentiation *in vitro* and
in vivo. Using a novel molecular biosensor to measure forces exerted on LINC complex
molecules, we find that tension is high on the LINC complex and A-type lamins in epidermal
stem cells in an integrin dependent manner and is reduced upon differentiation. In mouse
keratinocytes lacking LINC complexes we observe precocious differentiation *in vitro* and
expansion of the differentiated, suprabasal layers of the skin *in vivo*. This work suggests that
tension from integrins is communicated through the LINC complex to the nuclear lamina to
maintain the progenitor state of basal keratinocytes, providing a potential mechanism by which
cell fate is regulated directly by mechanical cues.
106
107

Results and Discussion:

A tension sensor in Nesprin-2 is sensitive to integrin engagement
In order to visualize tension on the LINC complex in living cells, we generated a molecular
biosensor to measure force on individual Nesprin proteins. To this end, we inserted a tension
sensor module composed of the fluorescence resonance energy transfer (FRET) pair mTFP
and Venus connected by an elastic flagelliform linker (Grashoff et al., 2010) into the
juxtamembrane region of a mini-Nesprin2 construct (Luxton et al., 2011), called hereafter "N2G-
JM-TSMod". This TSMod has been shown previously to be sensitive to forces in the single pN
range (Grashoff et al., 2010). In this construct, the TSMod lies between the transmembrane
domain and the entire cytosolic domain of mini-Nesprin2, which contains both the N-terminal
calponin homology domains that engage actin and a portion of the spectrin repeat region (**Fig.**
1A). This construct is distinct from previously-described Nesprin tension sensors in its design
(Arsenovic et al., 2016; Déjardin et al., 2020). We also generated a tension insensitive construct
by inserting the TSMod at the N-terminus of mini-Nesprin2 (**Fig. 1A**, "NoT_TSMod") as well as
"dark" controls that allow for bleed through between fluorescence channels to be assessed **(Fig.**
1-figure supplement 1).

We transfected primary undifferentiated mouse keratinocytes (MKCs) cultured on fibronectin-
coated glass bottom dishes with either the N2G-JM-TSMod or NoT_TSMod construct; both
were targeted efficiently to the nuclear envelope when expressed at moderate levels (**Fig. 1B,**
Fig. 1D, Fig. 1F, Fig. 1H, Fig. 1J). We anticipated that the tension insensitive NoT_TSMod
would exhibit constitutively high FRET indexes, reflecting the conformational freedom at the N-
terminus of the mini-Nesprin2 protein. By contrast, we expected that intramolecular tension
exerted on the N2G-JM-TSMod would lead to a decrease in the FRET index (**Fig. 1A**). Indeed,
relative to the mean NoT_TSMod FRET ratio (set to a value of 1), the N2G-JM-TSMod exhibited
lower FRET indexes when expressed in undifferentiated MKCs as assessed qualitatively by the
pseudocolored images reflecting the relative FRET index (**Fig. 1B**) or quantitatively for across
populations of MKCs (**Fig. 1C**). To determine if the actin cytoskeleton is required for tension on
the N2G-JM-TSMod in undifferentiated MKCs, we pharmacologically disrupted filamentous actin
with latrunculin A (Lat A). After 5 hours of Lat A treatment, the N2G-JM-TSMod displayed
relaxed tension compared to the vehicle control (set to a value of 1) as indicated by an increase
in the FRET indexes (**Fig. 1D, Fig. 1E**). By contrast, Lat A treatment had no effect on the
NoT_TSMod, as expected **(Fig. 1-figure supplement 1B)**. Based on these data, we conclude
that 1) the tension reported by the N2G-JM-TSMod reflects intramolecular tension exerted on
the juxtamembrane region; and 2) the N2G-JM-TSMod is sensitive to actin-dependent tension
at the nuclear envelope.

Next, we examined how the properties of cell-substrate engagement influenced tension on the
LINC complex. We plated undifferentiated MKCs on glass or 3 kPa PDMS substrates coated
with fibronectin (Mertz et al., 2013). We observed high tension on the N2G-JM-TSMod on stiff
glass substrates (set to a value of 1) relative to cells plated on the 3 kPa substrates, which
displayed lower tension (higher FRET ratio; **Fig. 1F, Fig. 1G**). Proliferative keratinocyte
progenitor cells bind to the basement membrane via several integrin based adhesions with the

laminin receptors α3β1 and the hemidesmosome-specific α6β4 being the chief epidermal
integrins (reviewed in (Burgeson and Christiano, 1997)). Thus, we also explored how substrate
composition affects tension on the LINC complex by plating MKCs on either fibronectin- or
laminin-coated glass coverslips or those coated with poly-L-lysine, which allows cell adhesion
through its positive charge rather than specific cellular adhesions. MKCs expressing the N2G-
JM-TSMod exhibited high tension (lower relative FRET ratio) at the nuclear envelope when
plated on fibronectin or laminin relative to when MKCs were plated on poly-L-lysine (**Fig. 1H**
and Fig. 1I). Furthermore, *β1* integrin null MKCs plated on glass coverslips coated with
fibronectin had higher N2G-JM-TSMod FRET ratios (lower tension) than WT controls (**Fig. 1J**
and **Fig. 1K**). Taken together, these data demonstrate that integrin adhesions are required for
high tension on LINC complexes at the nuclear envelope in epidermal progenitor cells.
163
Cell-intrinsic integrin engagement predicts N2G-JM-TSMod tension and differentiation in
cohesive MKC colonies
Primary MKCs can be induced to differentiate *in vitro* by elevating extracellular calcium levels,
leading to the formation of cohesive colonies that engage E-cadherin-based cell-cell adhesions
and subsequently differentiate as assessed by up-regulation of markers expressed in epidermal
suprabasal layers *in vivo* (Mertz et al., 2013). In our previous work, we demonstrated that this
transition leads to a reorganization of focal adhesions and traction stresses to cells at the colony
periphery, while cells in the colony interior solely engage cell-cell adhesions (Mertz et al.,
2013)(**Fig. 2A**). We also documented that cells at the colony periphery display a biased nuclear
position towards the colony center while cells within the colony interior display a central nuclear
position (**Fig. 2A**), suggesting that LINC complex tension might be distinct between MKCs at the
colony periphery and interior (Stewart et al., 2015). To test if tension on the N2G-JM-TSMod is
sensitive to the presence of cell intrinsic focal adhesion engagement, we segmented cells in
cohesive MKC colonies into "periphery" and "interior" cells. We find that the FRET ratio of the
N2G-JM-TSMod for interior cells is significantly higher than that observed for cells at the colony
periphery (**Fig. 2B**), indicating that tension on the LINC complex is lost in interior cells that
contain solely cell-cell adhesions. These data are consistent with the ability of focal adhesions
to drive LINC complex tension in a cell intrinsic manner in undifferentiated MKCs.
182
Given that the LINC complex has been shown to transmit tension from the cell substrate to the
lamin network at the nuclear periphery (Ihalainen et al., 2015), we examined whether tension on
lamin A/C within cohesive MKC colonies mirrored that observed for the LINC complex with the
N2G-JM-TSMod. To this end, we employed a conformationally sensitive lamin A/C antibody,
which has been used to examine changes in lamin tension due to substrate stiffness (Ihalainen
et al., 2015). When A-type lamins are under tension, the epitope of lamin A/C becomes
inaccessible, resulting in a decrease in fluorescence. We immunostained differentiated,
cohesive MKC colonies with a conformationally sensitive lamin A/C antibody and examined cells
at the periphery and interior. We imaged nuclei in which we could resolve the apical and basal
regions of the nuclear envelope and quantified the fluorescent signal in the z-plane as described
(Ihalainen et al., 2015) (**Fig. 2C**). We fitted the signal with two gaussian curves and extracted
the ratio of the basal to apical intensity across many colonies (n=17). In cells at the colony
periphery, we observe a strong bias of fluorescence at the apical nuclear surface compared to

the basal surface (**Fig. 2D**), indicating that tension on the lamin network is high on the basal
surface in cells at the periphery of colonies. By contrast, cells in the colony interior display far
more uniform conformationally sensitive lamin A/C staining (**Fig. 2E**). Taken together, these
results suggest that cells at the colony periphery that engage focal adhesions have high tension
on both LINC complexes and the nuclear lamina, while cells at the colony interior possess
relaxed LINC complexes and lamin A/C.
202
To examine whether the cells in the colony interior are more likely to express differentiation
markers, we performed RNA fluorescence in-situ hybridization (FISH) for *Sprr1b* mRNA, which
is upregulated in differentiated MKCs and is found in the EDC. We find that MKCs at the colony
interior express more *Sprr1b* and *Ivl* (involucrin) mRNA 24 and 48 hours after induction of
differentiation compared to cells at the colony periphery (**Fig. 2F-G** and **Fig. 2-figure**
supplement 1). These data indicate that tension on LINC complexes and the nuclear lamina is
more prominent in cells that maintain integrin adhesions and a progenitor fate.
210
The epidermis of Sun dKO mice displays precocious differentiation despite normal
adhesion To determine whether tension on LINC complexes controls MKC differentiation, we
analyzed the skin of mouse models lacking the ubiquitously expressed SUN proteins, SUN1 and
SUN2, which are expected to lack all LINC complexes in somatic tissues (Zhang et al., 2009;
Zhang et al., 2007). Similar to our prior analysis of mouse postnatal mouse skin (Stewart et al.,
2015), SUN1 and SUN2 are expressed at the nuclear envelope in all cellular layers of the E15.5
developing epidermis (**Fig. 3-figure supplement 1A**). Although *Sun1$^{-/-}$/Sun2$^{-/-}$* (hereafter Sun
dKO) mice die after birth (Zhang et al., 2009), we were able to analyze the developing epidermis
of Sun dKO mice at E15.5. Western blot analysis revealed that SUN1 and SUN2 proteins were
absent in the skin of Sun dKO compared to WT mice that express both SUN proteins (**Fig. 3-**
figure supplement 1B).
222
At E15.5 histological analysis of skin tissue revealed a thickening of the epidermis in the Sun
dKO mouse compared to WT littermates (**Fig. 3A**). Immunostaining of skin sections of WT and
Sun dKO mice with antibodies against proteins expressed in differentiated keratinocytes
revealed an expansion of keratin-10 (K10) and K1 in the spinous layer, involucrin in the spinous
and granular layers, and fillagrin in the granular and cornified layers (**Fig. 3B, Fig. 3C, and Fig.**
3-figure supplement 2). These observations suggest that LINC complex ablation in the
epidermis could enhance keratinocyte differentiation.
230
To further examine whether the thickened epidermis of Sun dKO mice resulted from changes in
differentiation and/or proliferation, we performed an EdU pulse chase experiment. We labelled
proliferating cells in embryonic WT and Sun dKO mice with EdU and examined the skin after 24
hrs (**Fig. 3D**). Immunostaining skin sections with K10 antibodies and staining for EdU
incorporation revealed that the total number of EdU-positive cells was similar between WT and
Sun dKO epidermis, indicating that proliferation was not altered in the absence of LINC
complexes in skin (**Fig. 3E**). However, while 40% of EdU-positive cells had moved to the basal,
K10-positive layer in WT tissue (**Fig. 3F**), 60% of EdU-positive cells were observed in the
differentiated layers in Sun dKO skin (**Fig. 3D** and **3F**), indicating an increase in keratinocyte

differentiation in the absence of LINC complexes. Thus, Sun dKO keratinocytes precociously
differentiate *in vivo* without changes in proliferation.

As integrin signals are required to repress the differentiation of basal keratinocytes (Adams and
Watt, 1989; Levy et al., 2000; Zhu et al., 1999), we explored the possibility that cell-matrix
adhesions are altered in number and/or size in the epidermis of Sun dKO mice. We therefore
performed electron microscopy on skin sections from heterozygous (*Sun1/2+/-*) and Sun dKO
mice at P0.5. The basal layer-basal lamina interface in *Sun1/2+/-* and Sun dKO skin appeared
indistinguishable, displaying tight association in mice from both genotypes (**Fig. 3G**). Moreover,
hemidesmosomes (which can be observed in electron micrographs, arrows) were of equivalent
number and size (**Fig. 3H**). These observations suggest that Sun dKO mice display precocious
epidermal differentiation *in vivo* that is uncoupled from changes in adhesion between basal
keratinocytes and the basal lamina.

Sun dKO MKCs exhibit precocious differentiation *in vitro*
To further analyze keratinocyte differentiation, we isolated WT and Sun dKO MKCs from
newborn mice and performed global transcriptional profiling on WT and Sun dKO MKCs grown
in either low calcium media to maintain a progenitor state or high calcium media to induce
differentiation. Gene ontology analysis revealed striking differences in the expression of genes
associated with keratinocyte differentiation between WT and Sun dKO MKCs in both low and
high calcium, including those that reside in the EDC (**Fig. 4A, Fig. 4B and Supplementary
Files 1-2**). Further examination of these RNAseq data revealed evidence of precocious
differentiation in Sun dKO MKCs. For example, numerous members of the gene family in the
SPRR region of the EDC (**Fig. 4A**) were de-repressed in Sun dKO MKCs grown in the low
calcium condition (**Fig. 4C**). We validated this global analysis by RT-qPCR analysis of *Sprr*
genes (**Fig. 4D** and **Fig. 4-figure supplement 1A**). Not only do we observe precocious
expression of genes such as *Sprr1b* in Sun dKO MKCs cultured in low calcium media, but we
also observe much higher expression in the presence of calcium in Sun dKO MKCs (**Fig. 4E**
and **Fig. 4-figure supplement 1A**). RNA FISH analysis of the level of *Sprr1b* (**Fig. 4F**) and *Ivl*
(**Fig. 4-figure supplement 1B-C**) transcripts further confirmed that Sun dKO MKCs displayed
higher expression levels of differentiation genes compared to WT MKCs.

Since we found that the differentiated, interior MKCs within cohesive colonies displayed low
tension on the LINC complex and A-type lamins (**Fig. 2**), we hypothesized that the lack of LINC
complexes may result in differentiation of progenitor cells at the colony periphery despite
integrin engagement, which is normally associated with maintenance of the progenitor fate. To
address this hypothesis, we performed RNA FISH on WT and Sun dKO cohesive MKC colonies.
While we demonstrated that WT MKCs biased expression of differentiation markers such as
Sprr1b and *Ivl* at the colony interior over the colony periphery (**Fig. 2F and Fig. 2G**) in
agreement with repressive signals from integrin engagement in cells at the colony periphery,
Sun dKO MKC colonies instead expressed *Sprr1b* mRNAs in MKCs residing in both the interior
and periphery (**Fig. 4E**), despite still having focal adhesions in peripheral cells (**Fig. 4-figure
supplement 2**). As a consequence, expression of differentiation markers is both elevated at the
colony periphery and is random with respect to colony position in the absence of LINC

complexes (**Fig. 4G**). Finally, we directly tested if β1 integrins are required for proper
keratinocyte differentiation by performing RT-PCR to analyze *Sprr1b*, Involucrin (*Ivl*) and
S100a14 mRNA levels in β1 integrin (*Itgb1*) KO MKCs. β1 integrin KO MKCs displayed
precocious expression of all three of these epidermal differentiation genes (**Fig. 4I**), mirroring
our observations in Sun dKO MKCs. Taken together, these results suggest that LINC
complexes are required to repress epidermal differentiation genes when integrins are actively
engaged.
291
Sun dKO MKCs exhibit an increase in chromatin accessibility at genes within the
Epidermal Differentiation Complex
The EDC moves away from the nuclear lamina during epidermal differentiation (Gdula et al.,
2013; Mardaryev et al., 2014; Williams et al., 2002) with an associated loss of repressive
H3K27me3 chromatin marks on differentiation-specific genes (Lien et al., 2011). Given that Sun
dKO MKCs display a precocious differentiation phenotype, we hypothesized that the EDC may
be more accessible when LINC complexes are ablated. To test this, we applied Assay for
Transposase-Accessible Chromatin using sequencing (ATAC-seq)(**Fig. 5-figure supplement 1**
and Supplementary File 3) to determine if chromatin accessibility was changed in Sun dKO
MKCs with a focus on the EDC. Strikingly, Sun dKO MKCs displayed almost twice as many
ATAC-seq peaks within the EDC (34 peaks) relative to WT MKCs (19 peaks)(**Fig. 5**).
Specifically, in Sun dKO MKCs we observed 18 additional peaks, 16 maintained peaks and a
loss of just 3 peaks compared to WT MKCs (**Fig. 5A)**. This net gain of peaks is observed
throughout all regions of the EDC, with the SPRR region displaying the most striking change,
gaining five novel ATAC-seq peaks upon loss of SUN proteins (**Fig. 5B, Fig. 5C**). Importantly,
while there is a trend towards increased chromatin accessibility in regions flanking the EDC
(**Fig. 5-figure supplement 2A**), the gain of ATAC-seq peaks is more pronounced in the EDC
compared to chromatin regions 5' and 3' (**Fig. 5D**). We also noted examples in which the
additional ATAC-seq peak was located near the promoter region of epidermal differentiation
genes such as *Sprr2a3* (**Fig. 5-figure supplement 2B**).
312
To test if this mode of regulation is specific to keratinocyte differentiation, we also looked at the
chromatin state upon loss of SUN proteins at other master transcriptional regulators that have
previously been shown to be repressed in basal keratinocytes (Ezhkova et al., 2009). Indeed,
chromatin at the master transcriptional regulators for muscle (*Myod1)* and the HOX (*Hoxd10*,
Hoxb13, and *Hoxa11*) gene families are in a closed chromatin state in both WT and Sun dKO
MKCs (**Fig. 5-figure supplement 2C**). Two of four neuronal master transcription factors (*Olig2*
and *Neurog2*) also remained in a closed state while the others (*Olig3* and *Neurog2*) revealed a
novel peak in Sun dKO cells (**Fig. 5-figure supplement 1**). Taken together, our observations
suggest that LINC complexes are required to maintain a closed, inhibited state of the EDC in
undifferentiated keratinocytes, providing a mechanism by which loss of LINC complexes leads
to precocious keratinocyte differentiation.
324
325
326

Discussion:

Here we demonstrate that LINC complex tension in MKCs responds specifically to integrin-engagement in a cell-intrinsic manner. While integrin-dependent signals normally repress the differentiation of keratinocytes, in the absence of functional LINC complexes these basal progenitors instead differentiate precociously both *in vitro* and *in vivo*. Based on our findings we propose a model in which LINC complexes transmit forces from the integrin-engaged actin network to the nuclear lamina to directly regulate the expression of epidermal differentiation genes (**Fig. 6**), although an indirect role for the LINC complex in an alternative signaling pathway remains possible. In particular, as our electron microscopy data strongly suggest that adhesion between basal keratinocytes and the basal lamina remains normal in Sun dKO animals, we strongly favor this direct model (**Fig. 6**). Of note, lower tension on a distinct Nesprin TSMod variant similar in design to the one described here was also observed in cells migrating through constrictions or plated at high versus low packing density, suggesting that tension on the LINC complex may respond to a number of mechanical inputs depending on the experimental system (Déjardin et al., 2020). However, high tension on the LINC complex at the edge of cell monolayer upon wounding compared to the interior of the monolayer (Déjardin et al., 2020) is in line with the integrin-dependence observed in this study.

How might tension on the nuclear lamina regulate epidermal differentiation? As a skin-specific mouse model lacking A- and B-type lamin expression also demonstrated precocious epidermal differentiation (Jung et al., 2014) and we find that MKC differentiation coincides with a relaxation of lamin A/C tension (**Fig. 2D and Fig. 2E**), one attractive model is that the LINC complex acts through tension-dependent remodeling of the nuclear lamina. The chromosome region housing the EDC moves away from the nuclear lamina during epidermal differentiation (Gdula et al., 2013; Mardaryev et al., 2014; Williams et al., 2002) and is associated with the loss of H3K27me3 chromatin marks on differentiation-specific genes (Lien et al., 2011). If and how lamin tension influences this change in nuclear compartmentalization remains to be defined. We favor a model in which stretching of lamins drives remodeling of the composition of the nuclear lamina to maintain the epidermal progenitor state. Interestingly, the conformational epitope in lamin A/C examined in this work (**Fig. 2**) overlaps with binding sites for DNA, histones, SUN proteins, and the integral inner nuclear membrane protein emerin (Haque et al., 2010; Ihalainen et al., 2015). Thus, the force-dependent differential exposure of these binding sites may influence the interaction of lamin A/C with the genome, either directly or through modulation of the activity of its binding partners. Indeed, stretching of MKCs leads to the release of emerin from the nuclear lamina; abrogating this mechanical response also leads to precocious differentiation (Le et al., 2016). Moreover, this response involves the formation of a perinuclear actin cage, a structure which we found previously to be perturbed in MKCs lacking SUN2 (Stewart et al., 2015).

Further work will be required to determine if the levels and geometry of forces exerted on LINC complexes differ between integrin-based adhesions and cell-cell adhesions. Another open question is whether building tension on the nuclear lamina requires a force opposing that mediated by the LINC-actin-integrin axis. In our prior work we found that control over nuclear position in MKCs at the periphery of cohesive colonies was dictated by a "tug-of-war" between

the actomyosin and microtubule networks (Stewart et al., 2015). Whether coupling of
microtubules to LINC complexes contributes to building tension on the LINC-actin-integrin axis,
and the magnitude of these intramolecular forces, remains an open question, as all Nesprin
TSMod constructs devised to date (Arsenovic et al., 2016; Déjardin et al., 2020), including the
one we describe here, lack the spectrin repeats responsible for mediating interactions with
microtubules and microtubule motors (Schneider et al., 2011). Examining whether the geometry
of exogenous force application to cells influences nuclear lamina-dependent regulation of gene
expression by modulating the extent of chromatin stretch (Tajik et al., 2016) represents another
exciting direction.
380
381
382
383

Materials and Methods

Cell culture and plasmid transfection

Mouse keratinocytes were isolated from skin from E18.5 embryos or newborn whole body *Sun1⁻/⁻/Sun2⁻/⁻* or WT pups as previously described (Mertz et al., 2013). Under sterile conditions, pups were sacrificed, and back skin was excised, washed in PBS, and floated on dispase at 4°C for 16–20 h. The epidermis was separated from the dermis with forceps and incubated in 0.25% trypsin for 15 min at RT. Cells were liberated by trituration, filtered using a 40–70-µm strainer, and plated on mitomycin-C–treated J2 fibroblasts in medium-calcium medium (0.3 mM $CaCl_2$). After two to four passages, keratinocytes were plated on plastic dishes without feeder cells and maintained in media containing 0.05 mM $CaCl_2$ (E-low calcium media). Cells at low passage were stored under liquid nitrogen, and thawed cells were only used to passage 16. *Integrin β1* KO keratinocytes and corresponding WT controls (Bandyopadhyay et al., 2012) (kind gift from David Calderwood and Srikala Raghavan) were cultured at 32ºC with 7.5% CO_2 in E media (DMEM/F12 in a 3:1 ratio with 15% fetal bovine serum supplemented with insulin, transferrin, hydrocortisone, cholera toxin, triidothyronone, and penicillin-streptomycin).

Construction and application of N2G-JM-TSMod constructs

The N2G-JM-TSMod construct was derived from pEGFP-C1 harboring mini-Nesprin2 (Luxton et al., 2010) (kind gift from Gant Luxton and Gregg Gundersen). The GFP was removed by QuikChange mutagenesis. The mTFP-Venus tension sensor module (Grashoff et al., 2010) (kind gift from Martin Schwartz) was inserted just prior to the transmembrane domain. Dark controls were generated using QuikChange mutagenesis (Dark Venus: Y67L; Dark mTFP: Y72L). The no-tension (NoT_TSmod) construct was generated by inserting the mTFP-Venus tension sensor module 5' of mini-Nesprin2 using the AgeI and XhoI restriction sites. For FRET experiments, cells were plated in E-low calcium media (or E media for Integrin B1 KO cells) 16 hours before transfections on glass-bottomed dishes (MatTek Corporation) coated with 50ug/ml fibronectin (20 minutes at room temperature, Sigma-Aldrich), Poly-L-Lysine (2 hours at room temperature, Sigma P9155) or Laminin (2 hours at 37ºC, Thermo Cat#CB-40232). Cells were transfected using Jetprime reagent (Polyplus) according to the manufacturer's instructions and imaged the following day. Donor bleed-through samples consisted of cells transfected with mini-Nesprin2G-mTFP1 alone (Fig. 1. supplement 1). Acceptor cross-excitation samples consisted of cells transfected with mini-Nesprin2G-Venus alone (Fig. 1, supplement 1). In addition, untransfected cells were also used as dark controls (Fig. 1. supplement 1). FRET samples consisted of Nesprin2G-JM-TSMod. Actin depolymerization was achieved by the addition of 0.5 µM latrunculin A (or vehicle control) for 5 hours prior to imaging.

Generation of hydrogels
The soft 3 kPa silicone gels were prepared by mixing a 1:1 ratio (w/w) of CY 52-276A and CY 52-276B (Dow Corning). Components were mixed thoroughly and immediately degassed for 15 mins in a desiccator. The surface of 35 mm glass bottomed culture dishes (MatTek corporation) were covered with 200 µl of the mixed gel. Each dish was then spin-coated at 1000 rpm for 60

428 seconds using a spin coater instrument (Headway Research, PWM32). Gels were cured at
429 room temperature overnight. The next morning, gels were sterilized for 30 minutes under UV
430 light and then washed once with sterile PBS. Rheometry analysis (ARES-LS1) confirmed that
431 the Young's modulus of the silicone gels was ~3 kPa.

432 **RNA FISH experiments**
433
WT and Sun dKO MKCs were plated at low density (15,000 cells/well of 24 well dish) onto glass
coverslips coated with 50 µg/ml fibronectin (Sigma Aldrich), MKCs were cultured in E-low
calcium media overnight before switching to 1.2 mM calcium media to induce differentiation for
24 or 48 hours. RNA FISH was performed using the ViewRNA ISH Cell Assay kit (ThermoFisher
Cat# QVC0001) with probes for *Involucrin* (ThermoFisher Cat# VB1-3030396-VC) and *Sprr1b*
(ThermoFisher Cat# VB4-3117172-VC) according to manufacturer's instructions. Coverslips
were mounted onto glass slides with Prolong Gold with DAPI (Invitrogen P36935) and sealed
with clear nail polish. Images were acquired with the Zeiss Imager M1 using Zen software.
Images for the same marker were acquired at the same exposure, pixel range, and gamma
values. Acquired images were equally brightened, contrasted and cropped using ImageJ/Fiji
(version 1.50e) software (Schindelin et al., 2012). For quantitation, signal threshold was
determined using no probe controls for each individual experiment. Colony outlines were drawn
and then cells were counted and scored as positive or negative for gene expression and interior
or periphery based on location within colony. Statistical analysis was performed using Prism 8
software.
449
450 **Mouse tissue isolation, histology and immunofluorescence staining**
451
This study was performed in strict accordance with the recommendations in the Guide for the
Care and Use of Laboratory Animals of the National Institutes of Health (IACUC protocol
number 2018-11248). All animal care and experimental procedures were conducted in accord
with requirements approved by the Institutional Animal Care and Use Committee of Yale
University. At harvest, embryos were submerged in O.C.T compound (Tissue-Tek) and
sectioned using a cryostat (CM3050S; Leica). Sections were cut in a specific and consistent
orientation relative to embryo morphology and stained with hematoxylin and eosin for routine
histopathology or incubated with primary antibodies and Alexa Fluor-conjugated secondary
antibodies for indirect immunofluorescence as previously described (Stewart et al., 2015).
Primary antibodies against SUN1 (1:100, Abcam, ab124770) and SUN2 (1:100; Abcam
ab124916) were used. Primary antibodies to keratin 10 (K10), keratin 1 (K1), involucrin, and
filaggrin were gifts from the Segre lab and are described in (Harmon et al., 2013). Images were
acquired with the Zeiss Imager M1 using Zen software. Images for the same marker were
acquired at the same exposure, pixel range, and gamma values. Acquired images were equally
brightened, contrasted and cropped using ImageJ/Fiji. Spinous layer thickness was quantitated
by making 5-8 measurements per image using ImageJ software. Statistical analysis was
performed using Prism 8 software.
469
470 **Western Blot Analysis**
471

472   Whole-cell lysates of WT, *Sun2-/-*, and Sun dKO MKCs were prepped as previously described
473   (Stewart et al., 2015). Primary antibodies against SUN1 (1:100, Abcam, ab124770)
474   SUN2 (1:100; Abcam ab124916), and β-Actin (1:1,000; mouse; Abcam) were used.
475
476   **EdU incorporation and quantitation**
477
Pregnant females were pulsed with EdU via IP injection when embryos were age E14.5. 24
hours after injection, embryos were isolated and embedded in O.C.T. media (Tissue Tek) and
stored at -80° C until sectioning using a cryostat (CM3050S; Leica). After sectioning, the Click-
iTEdU Cell Proliferation Kit for Imaging, Alexa Fluor 488 dye (Invitrogen C10337) was used
according to manufacturer's instructions to evaluate EdU incorporation. Tissue sections were
then co-stained for K10 (see mouse tissue isolation, histology, and immunofluorescence
staining). Before imaging, coverglass was mounted onto slides using Prolong Gold with DAPI
(Invitrogen P36935) and sealed with clear nail polish. Images were acquired with the Zeiss
Imager M1 using Zen software. Total Edu positive cells were counted and normalized to the
total number of epidermal cells (determined by DAPI staining). Location of proliferating cells was
determined by K10 staining (basal keratinocytes (EdU+/K10-), suprabasal keratinocytes
(EdU+/K10+)) and then normalized to total EdU positive cells. Statistical analysis was
performed using Prism 8 software.
491
492   **Electron Microscopy**
493
Transmission electron microscopy was performed in the Yale School of Medicine Center for
Cellular and Molecular Imaging Electron Microscopy core facility. Back skin sections from *Sun
1+/-/Sun2-/-* and Sun dKO pups was isolated at age P0.5; three mice were examined for each
genotype. Tissue blocks were fixed in 2.5% glutaraldehyde/2% paraformaldehyde in 0.1 M
sodium cacodylate buffer, pH 7.4, for 30 min at RT and 1.5 h at 4°C. The samples were rinsed
in sodium cacodylate buffer and were postfixed in 1% osmium tetroxide for 1 h. The samples
were rinsed and en bloc stained in aqueous 2% uranyl acetate for 1 h followed by rinsing,
dehydrating in an ethanol series to 100%, rinsing in 100% propylene oxide, infiltrating with
EMbed 812 (Electron Microscopy Sciences) resin, and baking overnight at 60°C. Hardened
blocks were cut using an ultramicrotome (UltraCut UC7; Leica). Ultrathin 60-nm sections were
collected and stained using 2% uranyl acetate and lead citrate for transmission microscopy.
Carbon-coated grids were viewed on a transmission electron microscope (TecnaiBioTWIN; FEI)
at 80 kV. Images were taken using a CCD camera (Morada; Olympus) and iTEM (Olympus)
software. The length and number of hemidesmosomes was quantitated from sections from 3
mice of each genotype.
509
510   **Immunofluorescence**
511
WT and Sun dKO keratinocytes were plated at low density (15,000 cells/well of 24 well dish)
onto glass coverslips coated with 50 µg/ml fibronectin (Sigma Aldrich), MKCs were cultured in
E-low calcium media overnight before switching to 1.2 mM calcium media to induce
differentiation for 48 hours. Cells were fixed with methanol at -20° C for 5 minutes and washed

with PBS. Cells were permeabilized using 0.5% Triton X-100 in PBS at RT for 20 minutes and
blocked using 10% goat serum, 5% BSA, and 0.5% Tween 20 in PBS for 1 hour. Cells were
incubated in primary antibody, conformationally sensitive lamin A/C (1:200; ab8984; Abcam),
diluted in blocking buffer at 4C overnight. Coverslips were washed with PBS for three 5 minute
intervals and incubated with AlexaFluor 488 conjugated secondary antibody (1:1000; Invitrogen
Cat# A-11029) diluted in blocking buffer at RT for 1 hour. Coverslips were then costained with
Hoechst 33342 (1:2000; ThermoFisher Scientific) and Alexa Fluor 594-conjugated Wheat Germ
Agglutinin (1:1000; Life Technologies) diluted in PBS for 5 minutes at RT. Coverslips were
washed with PBS, mounted using Fluoromount-G and sealed with clear nail polish.
525
526 **Imaging and image analysis**
527
Live FRET imaging was performed on a Zeiss LSM 710 DUO NLO confocal microscope using a
100x, 1.4 NA oil objective with a stage maintained at 37°C and 7.5% CO_2. Images were
acquired using Zen software. FRET imaging was performed in a similar manner as previously
described (Grashoff et al., 2010; Kumar et al., 2016). Three sequential images were acquired:
the donor mTFP1 channel using a 458 nm laser line (ex), 458 nm MBS filter, and PMT detector
set for mTFP1 emission; the acceptor Venus channel using a 514 nm laser line (ex), 458/514
nm MBS filter, and PMT detector set for Venus emission; and the FRET channel using a 458
nm laser line (ex), 458/514 nm MBS filter, and PMT detector set for Venus emission. In all
cases, the nuclear midplane was imaged. Acquisition settings were standardized and
maintained during experiments. FRET image analysis was performed using the intensity-based
FRET method, implemented as previously described (Kumar et al., 2016). Non-linear spectral
bleed-through corrections were first determined using the PixFRET plugin (Feige et al., 2005)
for ImageJ/Fiji (Schindelin et al., 2012). Donor mTFP1 leakage was quantified using cells
transfected with mini-Nesprin2G-mTFP1 alone (dark Venus control), while acceptor Venus
cross-excitation was quantified using cells transfected with mini-Nesprin2G-Venus alone (dark
mTFP control). At least 10 cells each were used for bleed-through corrections. For FRET index
determination, masks consisting of a 3 pixel-wide band encompassing the nuclear envelope
were used to segment the nuclear envelope. Mean FRET index per nucleus was then
determined using previously published software, either PixFRET (Figure 1) (Feige et al., 2005)
or in-house MatLab scripts (Figure 2) (Kumar et al., 2016).
548
Imaging of conformationally sensitive Lamin A/C was performed on a Leica SP5 confocal
microscope using the LAS-AF software. Lamin A/C (AF488) was imaged using a 488nm laser
line (ex) and PMT detector set for AF488 emission; and the WGA (AF594) using a 594nm laser
line (ex) and PMT detector set for AF594 emission. PMT gain was adjusted for each sample to
avoid over/under exposure. Acquisition settings were standardized and maintained during
experiments. Single plane images to determine the location of each cell within a colony were
acquired using a 40x 1.25 NA air objective, imaging the AF488 and AF594 channels
sequentially. Next, Z-stacks of each images were acquired using a 63x, 1.4 NA oil objective in
the AF488 channel. Image zoom and size was adjusted such that voxel size was 55-65 nm in
the xy-plane and 130 nm in the z-plane. Image stacks were deconvolved using Huygens
Professional software (Scientific Volume Imaging, The Netherlands, http://svi.nl) as described

(Ihalainen et al., 2015). A theoretical point spread function was used for iterative deconvolution.
Deconvolution was performed using the following software parameters: image signal to noise
was set to 5, the quality threshold was 0.01, and maximum iterations was 50 (however usually
fewer than 20 iterations were required to reach the quality threshold). Images were then
analyzed using ImageJ/Fiji (Schindelin et al., 2012). The apical-basal Lamin A/C intensity was
measured for a single XZ or YZ slice from the middle of each individual nucleus of interest. The
intensity of antibody staining was measured from the apical to the basal side of the nucleus
using a straight line along the z-axis that is approximately half the width of the nucleus of
interest. The fluorescence intensity across the z-axis was plotted and fit to two gaussian curves,
one corresponding to the apical side of the nuclear envelope and the second corresponding to
the basal side. The area under each curve and the ratio of these areas were calculated using
MATLAB R2019b and the software can be found on GitHub at the following url:
https://github.com/LusKingLab/GaussianFit.
573
574 **RNAseq**
575
WT and Sun dKO cells were grown in E-low calcium media for 24 h to 100% confluency
(undifferentiated) or were switched to high calcium media to induce adhesion formation and
differentiation for 48 h. Total RNA was isolated using the RNeasy Plus kit (QIAGEN) according
to the manufacturer's instructions for three biological replicates for each condition. cDNA was
synthesized using reagents from the TruSeq RNA sample preparation kit (Illumina) according to
the manufacturer's instructions. cDNA libraries were sequenced (paired end 75 nts) at the Yale
Stem Cell Center Genomics and Bioinformatics Core on the HiSeq4000 platform. Reads were
mapped using BowTie/TopHat2 (Kim et al., 2013) to the mm10 genome build. Differentially
expressed genes between the WT and Sun dKO conditions in the undifferentiated and
differentiated states were identified using DESeq2 (Love et al., 2014). All sequencing data can
be accessed at NCBI under BioProject Accession PRJNA636991.
587
588 **RT-qPCR**
589
WT and Sun dKO keratinocytes were plated in a fibronectin-coated (50ng/ml; Sigma Aldrich) 6-
well dish such that they were 70-80% confluent. They were cultured in E-low calcium media
overnight before switching to 1.2 mM calcium media to induce differentiation for 48 hours. Total
RNA was isolated using the RNeasy Plus kit (QIAGEN) according to the manufacturer's
instructions. The iScript cDNA Synthesis Kit (BioRad) was used to generate cDNA from equal
amounts of total RNA (1mg) according to the manufacturer's instructions. Quantitative real-time
PCR was performed with a BioRad CFX96 using iTaq Universal SYBR Green Supermix
(BioRad) for 40 cycles. Primers used include: GAPDH forward:
AGGTCGGTGTGAACGGATTTG and reverse: TGTAGACCATGTAGTTGAGGTCA; Sprr1b
forward: GATCCCAGCGACCACAC and reverse: GCTGATGTGAACTCATGCTTC; Sprr2d
forward: GTGGGCACACAGGTGGAG and reverse: GCCGAGACTACTTTGGAGAAC;
Involucrin forward: GCAGGAGAAGTAGATAGAG and reverse: TTAAGGAAGTGTGGATGG;
S100a14 forward: GGCAGGCTATAGGACA and reverse: CCTCAGCTCCGAGTAA.. PCR
product levels were normalized to GAPDH mRNA levels.

604
ATAC-seq

ATAC-seq libraries were generated as previously described (Buenrostro et al., 2015). Briefly,
50,000 nuclei were purified from WT or Sun dKO cells and DNA was tagmented using the
Nextera Library Prep Kit (Illumina). Tagmented DNA was purified using the Qiagen MinElute
PCR Purification kit, then amplified using PCR. RT-qPCR was used to determine the
appropriate number of cycles of amplification. The DNA was then purified twice using Sera-Mag
Select beads (GE). Library purity and quality was assessed using an Agilent Bioanalyzer, then
was sequenced on the Illumina HiSeq 4000 using paired end reads. Sequencing was performed
at the Yale Stem Cell Center Genomics Core facility. Sequences were processed using
the ENCODE ATAC-seq pipeline version 1.8.0, an automated end-to-end quality control and
processor of ATAC-seq data. The minimum template seeded the input JSON onto the mm10
genome build with the auto detect adapter parameter set to true. All sequencing data can be
accessed at NCBI under BioProject Accession PRJNA636991.

Acknowledgements: We would like to thank the laboratories of G.W. Gant Luxton (University
of California, Davis), Gregg Gundersen (Columbia University), Julia Segre (NHGRI), Srikala
Raghavan (Institute for Stem Cell Science and Regenerative Medicine), David Calderwood
(Yale University), Martin Schwartz, (Yale University), and Chinedum Osuji (Yale University) for
sharing of reagents, instrumentation and/or expertise. The high-throughput sequencing was
conducted by the Yale Stem Cell Center Genomics and Bioinformatics Core facility, which was
supported by the Connecticut Regenerative Medicine Research Fund. This work would not have
been possible without the support of the Ludwig Family Foundation and the Physical
Engineering Biology Program at Yale University.

Figure Legends
632
Figure 1
A tension sensor in mini-Nesprin2 is under actin-dependent tension when WT mouse
keratinocytes engage the extracellular matrix through integrins. (A) Cartoon of the N2G-
JM-TSMod, in which the mTFP-Venus tension sensor module is inserted between the
cytoplasmic domain and the C-terminal KASH domain/transmembrane domain of mini-Nesprin2.
Tension leads to a reduction in the FRET index. Cartoon of the NoT_TSMod control, in which
the TSMod resides at the N-terminus of the N2G, and therefore cannot experience
intramolecular tension. **(B,C)** The N2G-JM-TSMod displays higher tension (lower FRET index)
than the no-tension NoT_TSMod control (higher FRET index) when expressed in mouse
keratinocytes (MKCs) plated on fibronectin-coated glass. Representative images show that, at
low expression levels, both the N2G-JM-TSMod and NoT_TSMod are successfully targeted to
the nuclear envelope. Images are pseudocolored according to the normalized FRET index
(Feige et al., 2005). The median FRET index value of cells expressing the NoT_TSMod was set
to a value of 1 in (C) and used to scale to relative values for the N2G-JM-TSMod. **(D, E)**
Disruption of actin filaments with 0.5 μM latrunculin A (Lat A) decreases the tension on the N2G-
JM-TSMod, leading to a higher FRET index. The median FRET index of cells expressing the
N2G-JM-TSMod was set to a value of 1 in (E). **(F,G)** Tension on the N2G-JM-TSMod is
sensitive to substrate mechanics. Plating of MKCs on fibronectin-coated compliant substrates (3
kPa PDMS) leads to increased FRET compared to MKCs plated on fibronectin-coated glass.
(H,I) Plating MKCs on extracellular matrix that engages integrins drives high tension on the
N2G-JM-TSMod. MKCs grown on glass coated with fibronectin or laminin drives a higher
tension state (low FRET index) than for cells grown on glass coated with poly-L-lysine (high
FRET index). The median FRET index value of cells expressing the N2G-JM-TSMod plated on
fibronectin-coated glass was set to a value of 1 in (G and I). Data for FN replotted from (I). **(J,K)**
MKCs lacking β1 integrin fail to drive high tension on the N2G-JM-TSMod. Representative
images demonstrate that the N2G-JM-TSMod is successfully targeted to the nuclear envelope
in WT and β1 integrin KO MKCs. Higher FRET indexes at the nuclear envelope in β1 integrin
KO MKCs demonstrates that β1 integrin engagement with the extracellular matrix is required for
high tension on the N2G-JM-TSMod. The median FRET index value of cells expressing the
N2G-JM-TSMod plated on fibronectin-coated glass was set to a value of 1 in (K). For all plots
errors reflect SD, $n \geq 30$ cells for each condition measured from $n = 3$ experiments. **** $p <$
0.0001 as determined by unpaired t-test (C, E, G, K) or one-way ANOVA (I). All scale bars = 5
μm.
666
Figure 1-figure supplement 1
Validation controls for the N2G-JM-TSMod. (A, C, D) Representative images of the "dark"
control constructs (mTFP donor or Venus acceptor fluorescence only) used for bleed through
assessment as required for FRET experiments. For comparison, representative images of the
N2G-JM-TSMod construct in which fluorescent signals can be readily detected in the donor,
acceptor and FRET channels are included. **(B)** The FRET index for the NoT_TSMod control is
unaffected by actin depolymerization. In WT MKCs expressing the NoT_TSMod control there is
no significant (ns) change in FRET index as assessed by unpaired t-test upon addition of 0.5

μM latrunculin A (Lat A) compared to vehicle controls. Data for vehicle alone replotted from Fig.
1C. Errors reflect SD, *n* ≥ 30 cells for each condition measured from *n* = 3 experiments. Scale
bar = 5 μm.
678
Figure 2
Tension on the N2G-JM-TSMod and nuclear lamina is released at the interior of cohesive
mouse keratinocyte (MKC) colonies concomitant with differentiation. (A) Cartoon of cell
junction reorganization upon differentiation. In single MKCs grown in low calcium media focal
adhesions and traction forces are found at the periphery of each cell (red). In response to high
calcium, MKCs form cohesive colonies and engage cell-cell junctions, which leads to a
reorganization of focal adhesions and traction forces to the colony periphery. Cells in the colony
interior have cell-cell but not cell-matrix adhesions. **(B)** The FRET index is higher in cells at the
interior of cohesive MKC colonies compared to cells at the periphery, suggesting that tension on
the N2G-JM-TSMod requires cell-intrinsic focal adhesion engagement. **(C)** Cartoon of method
used to measure changes in the tension state of A-type lamins using a conformationally-
sensitive lamin A/C antibody whose epitope is lost when lamins are under tension. Cells under
tension (left) lose staining on the basal side of the nuclear envelope. The profile of fluorescence
intensity from the apical to basal side of each nuclei was measured for a region of interest in XZ
and YZ slices from individual nuclei at varying locations within a colony. **(D)** Tension at the basal
surface of nuclei is lost at the interior of cohesive MKC colonies and is maintained in cells at the
colony periphery in response to differentiation. Examples of xz and yz confocal sections of
individual nuclei and corresponding Z-intensity profiles measured in the region of interest,
indicated by the white box (left). Intensity profiles were fit to two gaussians (right), representing
staining of the apical and basal sides of the nuclear envelope. The intensity of antibody staining
was defined as the area under the curve of the gaussian distribution corresponding to each side
of the nuclear envelope. A ratio of the intensity of antibody staining of the basal relative to the
apical side of the nuclear envelope was calculated. This analysis revealed that tension on the
basal nuclear surface is relaxed at the colony interior relative to the colony periphery. Scale bar
= 5 μm. **(E)** A histogram of the ratio of basal to apical intensity calculated as described in (C) for
all cells analyzed (n = 47 for periphery and n = 28 for interior) shows that the ratio of basal to
apical intensity for interior cells is shifted to higher values relative to cells at the periphery,
indicating the lamina is under less tension. Values were binned every 0.05 arbitrary units and
the central value of each bin is labeled. Representative images for the low (magenta triangle)
and high (blue triangle) bins, and bins corresponding to the highest percentage of periphery
(yellow triangle) and interior (green triangle) are shown in (C). **(F-G)** The differentiation marker
Sprr1b is expressed at higher levels in the colony interior than at the colony periphery. **(F)**
Representative image of RNA FISH for *Sprr1b* 24 hours after addition of calcium to induce
differentiation. Inset 1 shows *Sprr1b*-positive cells at the colony interior. Inset 2 shows *Sprr1b*-
positive cells at the colony periphery. Dotted lines are colony outline. Scale bar = 100 μm. **(G)**
Quantitation of the percent of *Sprr1b*-positive cells that are located at the interior and periphery
of WT MKC colonies normalized to the total *Sprr1b*-positive cells. 24h = 24 hours calcium
treatment. 48h = 48 hours calcium treatment. * p value < 0.05. ** p value <0.01 as determined
by unpaired t-test. Error bars are SD. N = 3 biological replicates.
718

Figure 2-figure supplement 1
The differentiation marker involucrin (*Ivl*) is expressed at higher levels in the colony
interior than at the colony periphery. (A) Representative image of RNA FISH in WT MKCs for
Ivl 24 hours after addition of calcium to induce differentiation. Inset 1 shows *Ivl*-positive cells at
the colony interior. Inset 2 shows *Ivl*-positive cells at the colony periphery. Dotted lines are
colony outline. Scale bar = 100 µm. **(B)** Quantitation of the percent of *Ivl*-positive cells that are
located at the interior and periphery of WT MKC colonies normalized to the total *Ivl*-positive
cells. 24h = 24 hours calcium treatment. 48h = 48 hours calcium treatment. Statistical
significance determined by unpaired t-test. * p value < 0.05. ** p value <0.01. Error bars are SD.
N = 3 biological replicates.
729
Figure 3
LINC complex ablation leads to precocious epidermal differentiation *in vivo*. (A) Increased
epidermal thickness in Sun dKO mice. Representative Hematoxylin and Eosin staining of WT
and Sun dKO skin at age E18.5. Dotted line denotes dermal/epidermal junction. Vertical line
denotes epidermal thickness. Scale bar = 20 µm.**(B)** Expansion of the spinous layer in Sun dKO
mice. Representative immunostaining for keratin 10 (K10) in WT and Sun dKO skin at age
E15.5. Dotted line denotes dermal/epidermal junction. Vertical line denotes spinous layer
thickness. Scale bar = 20 µM. Nuclei are stained with DAPI. **(C)** Quantitation of average spinous
layer thickness as determined by immunostaining for K10 in WT and Sun dKO epidermis at
E15.5. Unpaired t-test was used to determine statistical significance. ** p value <0.01. Error
bars are SD. N = 5-8 biological replicates per genotype. **(D-F)** Pulse chase analysis reveals an
increase in EdU-positive cells in the suprabasal layers of Sun dKO mice (and decrease in EdU-
positive basal cells) but no increase in overall proliferation. **(D)** Representative images of E15.5
WT and Sun dKO skin 24 hours after EdU pulse. Spinous layer keratinocytes are marked by
K10 staining. Dotted line denotes dermal/epidermal junction. Asterisks denote suprabasal
keratinocytes marked with EdU. Scale bar = 20 µm. Nuclei are stained with DAPI. **(E)**
Quantitation of total EdU positive cells normalized to total epidermal cells in both WT and *Sun*
dKO epidermis at E15.5. Statistical significance was determined using unpaired t-test. n.s. = not
significant. N = 3 mice/genotype. **(F)** Quantitation of location of EdU positive cells in the
epidermis of E15.5 WT and Sun dKO epidermis 24 hours after EdU pulse. K10 staining was
used to differentiate suprabasal from basal keratinocytes. Statistical significance was
determined by performing unpaired t-test. ** p value < 0.01. N = 3 mice/genotype. **(G-H)**
Adhesion between basal keratinocytes and the basal lamina is normal in the Sun dKO skin. **(G)**
Representative electron microscopy images of P0.5 WT and Sun dKO basal keratinocytes *in*
vivo. Scale bar = 10 µm. Dotted line denotes basement membrane. Arrows point to
hemidesmosomes. **(H)** Quantitation of hemidesmosome length (left) and the number of
hemidesmosomes per surface area (right) in control and Sun dKO epidermis at age P0.5. ns =
not statistically significant as determined by unpaired t-test.
758
Figure 3-figure supplement 1
SUN1 and SUN2 are expressed throughout the epidermis. (A-B) Representative
immunostaining for proteins SUN1 (**A**) and SUN2 (**B**) in WT skin at age E15.5. Dotted line
denotes dermal/epidermal junction. Scale bar = 20 µm. **(C)** Western blot analysis of WT, *Sun2* -

/- and Sun dKO primary keratinocytes confirming loss of SUN1 and SUN2 expression as
expected. β-actin was used as a loading control.
765

Figure 3-figure supplement 2
Thickening of the differentiated layers in the Sun dKO epidermis. (A) Representative
immunostaining for keratin 1 (K1), a marker of spinous keratinocytes, in WT and Sun dKO skin
at age E15.5. Dotted line denotes dermal/epidermal junction. Vertical line denotes spinous layer
thickness. Scale bar = 20 μm. Nuclei are stained with DAPI. **(B)** Quantitation of average spinous
layer thickness as determined by immunostaining for K1 in WT and Sun dKO epidermis at
E15.5. Unpaired t-test was used to determine statistical significance. ** p value <0.01. Error
bars are SD. N = 4 biological replicates per genotype. **(C)** Representative immunostaining for
involucrin (Ivl), a marker of spinous and granular keratinocyte layers, in WT and Sun dKO skin
at age E15.5. Dotted line denotes dermal/epidermal junction. Vertical line denotes spinous layer
thickness. Scale bar = 20 μm. Nuclei are stained with DAPI. **(D)** Quantitation of average
spinous/granular layer thickness as determined by immunostaining for Ivl in WT and Sun dKO
epidermis at E15.5. Unpaired t-test was used to determine statistical significance. ** p value
<0.01. Error bars are SD. N = 4 biological replicates per genotype. **(E)** Representative
immunostaining for Filaggrin (Flg), a marker for granular and cornified keratinocyte layers, in
WT and Sun dKO skin at age E15.5. Dotted line denotes dermal/epidermal junction. Vertical line
denotes spinous layer thickness. Scale bar = 20 μm. Nuclei are stained with DAPI. **(F)**
Quantitation of average granular/cornified layer thickness as determined by immunostaining for
Flg in WT and Sun dKO epidermis at E15.5. Unpaired t-test was used to determine statistical
significance. ** p value <0.01. Error bars are SD. N = 4 biological replicates per genotype.
786

Figure 4
LINC complex ablation leads to precocious MKC differentiation *in vivo*, which is
associated with aberrant up-regulation of differentiation markers at the colony periphery.
(A) Cartoon of the epidermal differentiation complex (EDC), a large genic region that is
coordinately up-regulated upon epidermal differentiation. **(B)** Comparative transcriptome
analysis reveals that Sun dKO MKCs display precocious expression of epidermal differentiation
genes when cultured in low calcium media and higher levels of expression of epidermal
differentiation genes in high calcium media. See also Supplementary Files 1 and 2. **(C)**
Examples of EDC genes that are precociously expressed in Sun dKO MKCs cultured in low
calcium media from the RNAseq data, expressed as fragments per kb of transcript per million
reads. **(D)** Real-time qPCR analysis of *Sprr1b* in WT and Sun dKO MKCs in the presence and
absence of calcium validates precocious *Sprr1b* expression in Sun dKO MKCs without calcium
stimulation. Ct values were normalized to GAPDH. Fold change in expression was determined
by calculating the $2^{\Delta\Delta Ct}$ relative to the mean ΔCt of WT MKCs cultured in low calcium media. n=3
biological replicates for all conditions. Statistical significance was determined by performing
multiple t-tests. * = p < 0.05. The Holm-Sidak method was used to correct for multiple
comparisons. Error bars are SD. N = 3 biological replicates. **(E-G)** Cohesive Sun dKO MKCs
express elevated levels of differentiation markers and lose the relationship between position in
the colony and EDC gene expression. **(E)** Representative images of RNA FISH for *Sprr1b* in
WT and Sun dKO MKCs after 24 hours calcium treatment. Dotted lines are colony outline. Scale

bar = 100 μm. **(F)** Quantitation of RNA FISH for *Sprr1b* in WT and Sun dKO MKCs after calcium
treatment. *Sprr1b*-positive cells were counted and normalized to total cells in each field. 24h =
24 hours calcium treatment. 48h = 48 hours calcium treatment. Statistical significance was
determined by performing unpaired t-test. * p value < 0.05. n.s. = not statistically significant.
Error bars are SD. N = 3 biological replicates. **(G)** Quantitation of the percent *Sprr1b*-positive
cells that are located at the periphery of WT and Sun dKO MKC colonies normalized to the total
Sprr1b-positive cells. 24h Ca = 24 hours calcium treatment. 48h Ca = 48 hours calcium
treatment. n.s. = not statistically significant as determined by unpaired t-test. Error bars are SD.
N = 3 biological replicates. **(H)** Quantitation of the percent of *Sprr1b*-positive cells that are
located at the interior and periphery of Sun dKO MKC colonies normalized to the total *Sprr1b*-
positive cells. 24h = 24 hours calcium treatment. 48h = 48 hours calcium treatment. n.s. = not
statistically significant as determined by unpaired t-test. Error bars are SD. N = 3 biological
replicates. **(I)** Real-time qPCR analysis of *Sprr1b*, Involucrin (*Ivl*), and *S100a14* demonstrate
precocious differentiation in β1 integrin knock out MKCs. Ct values were normalized to GAPDH.
Fold change in expression was determined by calculating the $2^{\Delta\Delta Ct}$ relative to the mean of WT
ΔCt. Statistical significance was determined by student's t-tests. * = p < 0.05. Error bars are SD.
N = 3 biological replicates.
824
Figure 4-figure supplement 1
Additional evidence for precocious differentiation in Sun dKO MKCs. (A) Real-time qPCR
demonstrating precocious expression of *Sprr2b* in Sun dKO MKCs. Fold change in expression
was determined by calculating the $2^{\Delta\Delta Ct}$ relative to the mean of WT no calcium ΔCt. n=3
biological replicates for all conditions. Statistical significance was determined by performing
multiple t-tests. * = p < 0.05. The Holm-Sidak method was used to correct for multiple
comparisons. Error bars are SD. N = 2 biological replicates. **(B)** Representative images of RNA
FISH for involucrin (*Ivl*) in WT and Sun dKO MKCs after 24 hours of calcium treatment. Dotted
lines are colony outline. Scale bar = 100 μm. **(C)** Quantitation of RNA FISH for involucrin in WT
and Sun dKO MKCs after calcium treatment. Involucrin positive cells were counted and
normalized to total cells in each field. 24h = 24 hours calcium treatment. 48h = 48 hours calcium
treatment. Error bars are SD. N = 3 biological replicates. **(D)** Quantitation of the percent
involucrin-positive cells that are located at the periphery of WT and Sun dKO MKC colonies
normalized to the total involucrin-positive cells. 24h Ca = 24 hours calcium treatment. 48h Ca =
48 hours calcium treatment. Error bars are SD. N = 3 biological replicates. **(E)** After 48 hours
Sun dKO MKCs fail to display a bias in involucrin expression towards the colony interior (as
seen in WT MKCs (Fig. 2-figure supplement 1B)). Statistical significance was determined by
paired t-test in (C-E).
843
844 **Figure 4-figure supplement 2**
845 **Cohesive Sun dKO MKC colonies have focal adhesions at the colony periphery.**
846 Representative images of WT and Sun dKO MKCs cultured in calcium for 48 hours, fixed and
847 stained with anti-paxillin antibodies. DNA was stained with Hoechst 33342. Scale bar = 25 μm.
848
Figure 5.

LINC complex ablation leads to an increase in chromatin accessibility of genes residing
in the Epidermal Differentiation Complex. ATAC-seq tracks generated from either WT (top;
N=2) or Sun dKO (bottom; N=2) MKCs grown in low calcium media. **(A)** There is an increase in
the number of accessible peaks (n=38) within the Epidermal Differentiation Complex (EDC) in
Sun dKO MKCs compared to WT MKCs (n=19). Peaks gained in Sun dKO MKCs are
highlighted in blue and peaks lost are highlighted in red. Unchanged peaks are not highlighted.
(B) Expanded view of ATAC-seq peaks in each indicated region of the mouse EDC. Peaks
gained in Sun dKO MKCs are highlighted in blue and peaks lost are highlighted in red.
Unchanged peaks are not highlighted. **(C)** Quantitation of novel ATAC-seq peaks in WT and
Sun dKO MKCs. The number of unique peaks over the indicated regions are plotted for the two
genotypes. **(D)** Quantitation of the percentage of total ATAC-seq peaks that are novel in WT
and Sun dKO MKCs for the EDC and surrounding chromatin regions (ATAC-seq tracks in Fig.
5-figure supplement 1). While all gains in ATAC-seq peaks across all regions is equivalent for
WT MKCs and Sun dKO MKCs generally demonstrate an increase in accessibility across all
three regions, the gain of peaks at the EDC is more pronounced than in equivalently sized
chromatin regions 5' and 3' to the EDC. See also Supplementary File 3.
866
867 **Figure 5-figure supplement 1**
868 **Quality assessment for each replicate of ATAC-seq data.** Both replicates of WT and Sun
869 dKO MKCs cultured in low calcium media demonstrate enrichment of ATAC-seq peaks with
870 transcriptional start sites (TSSs) (>6-fold). See also Supplementary File 3.
871
Figure 5-figure supplement 2
Increased accessibility within the EDC is specific as assessed by ATAC-seq in Sun dKO
MKCs cultured in low calcium media. (A) Tracks generated from either WT (top; n=2) or Sun
dKO (bottom; n=2) MKCs cultured in low calcium media at equivalently EDC-sized genomic
regions of chromatin directly 5' (top) and 3' (bottom) to the Epidermal Differentiation Complex.
Peaks gained upon loss of SUN proteins are highlighted in blue and peaks lost are highlighted
in red. Unchanged peaks are not highlighted. **(B)** Example of a novel ATAC-seq peak within the
promoter region of the *Sprr2a3* gene in Sun dKO MKCs cultured in low calcium media,
indicating increased chromatin accessibility of this region. **(C)** ATAC-seq revealed a closed
chromatin state in both WT and Sun dKO MKCs cultured in low calcium media at most master
transcription factors tied to other differentiation programs. *Myod1* (muscle) and the HOX gene
master transcription factors *Hoxd10*, *Hoxb13*, and *Hoxa11* and at two of four neuronal master
transcription factors (*Olig2* and *Neurog1*) show no changes between WT and Sun dKO MKCs.
Two additional neuronal master transcription factors (*Olig3* and *Neurog2*) revealed a novel peak
in Sun dKO cells.
887
Figure 6.
Model for the mechanism of precocious differentiation in the Sun dKO epidermis. WT
progenitor cells adhere to the basal lamina through integrins, leading to tension on LINC
complexes that is transmitted to the basal surface of the nuclear lamina to maintain repression
of the EDC. As cells move into the suprabasal layer they lose integrin engagement. In
suprabasal cells that only engage cell-cell junctions, LINC complexes and the nuclear lamina

are relaxed, leading to up-regulation of EDC gene expression. In the Sun dKO epidermis,
tension from engaged integrins is not propagated to the nucleus in progenitor cells despite
normal adhesion, leading to precocious expression of EDC genes.
897
898   **Supplementary File 1**
899   Table of RNAseq (Tab 1) and GO Term analysis (Tab 2) for Sun dKO versus WT mouse
900   keratinocytes grown in low calcium media (undifferentiated).
901
902   **Supplementary File 2**
903   Table of RNAseq (Tab 1) and GO Term analysis (Tab 2) for Sun dKO versus WT mouse
904   keratinocytes grown in high calcium media (differentiated).
905
906   **Supplementary File 3**
907   Quality control metrics for ATAC-seq experiments.
908
909
910

Key Resources Table				
Reagent type (species) or resource	**Designation**	**Source or reference**	**Identifiers**	**Additional information**
strain, strain background (*M. musculus*)	WT C57Bl/6J	Jackson Laboratories, Bar Harbor, ME	Stock number 000664	
genetic reagent (*M. musculus*)	*Sun1⁻/⁻/Sun2⁻/⁻* (Sun dKO)	Jackson Laboratories, Bar Harbor, ME	B6;129S6-*Sun1^{tm1Mhan}*/J Stock No: 012715 crossed to B6;129S6-*Sun2^{tm1Mhan}*/J Stock No: 012716	
Cell line (*M. musculus*)	Primary WT Keratinocyte	This paper		Isolated from WT pups
Cell line (*M. musculus*)	Primary Sun dKO keratinocyte	This paper		Isolated from Sun dKO pups
Cell line (*M. musculus*)	*Integrin β1* KO	(Bandyopadhyay et al., 2012)		
Antibody	SUN1 antibody (Rabbit monoclonal)	Abcam	ab124770	IHC (1:1000) WB (1:100)
Antibody	SUN2 antibody (rabbit monoclonal)	Abcam	ab124916	IHC (1:1000) WB (1:100)
Antibody	keratin 10 antibody (Rabbit polyclonal)	Gift from Julia Segre (Harmon et al., 2013)		IHC (1:500)

Antibody	keratin 1 antibody (Chicken polyclonal)	Gift from Julia Segre (Harmon et al., 2013)		IHC (1:500)
Antibody	Involucrin antibody (Rabbit polyclonal)	Gift from Julia Segre (Harmon et al., 2013)		IHC (1:500)
Antibody	Filaggrin antibody (Chicken polyclonal)	Gift from Julia Segre (Harmon et al., 2013)		IHC (1:500)
Antibody	Beta actin antibody (mouse monoclonal)	Abcam	ab13772	WB: 1:1000
Antibody	Conformationally sensitive Lamin A/C antibody	Abcam	ab8984	IF: 1:200
Recombinant DNA reagent	N2G-JM-TSMod (plasmid)	This paper		Constructed from pEGFP-C1 containing mini-Nesprin-2G (Luxton et al., 2010)
Recombinant DNA reagent	N2G-JM-TSMod Dark Venus (plasmid)	This paper		Mutation in Venus Y67L
Recombinant DNA reagent	N2G-JM-TSMod Dark mTFP (plasmid)	This paper		Mutation in mTFP Y72L
Recombinant DNA reagent	NoT_TSMod (Plasmid)	This paper		Constructed from pEGFP-C1 containing N2G-JM-TSMod
Sequence-based reagent	*Involucrin* RNA FISH probe	Thermo Fisher	VB1-3030396-VC	
Sequence-based reagent	*Sprr1b* RNA FISH probe	Thermo Fisher	VB4-3117172-VC	

Sequence-based reagent	GAPDH_F	This paper	qPCR Primer	AGGTCGGTGTGAACGGATTTG
Sequence-based reagent	GAPDH_R	This paper	qPCR Primer	TGTAGACCATGTAGTTGAGGTCA
Sequence-based reagent	Sprr1b_F	This paper	qPCR Primer	GATCCCAGCGACCACAC
Sequence-based reagent	Sprr1b_R	This paper	qPCR Primer	GCTGATGTGAACTCATGCTTC
Sequence-based reagent	Sprr2d_F	This paper	qPCR Primer	GTGGGCACACAGGTGGAG
Sequence-based reagent	Sprr2d_R	This paper	qPCR Primer	GCCGAGACTACTTTGGAGAAC
Sequence-based reagent	Involucrin_F	This paper	qPCR Primer	GCAGGAGAAGTAGATAGAG
Sequence-based reagent	Involucrin_R	This paper	qPCR Primer	TTAAGGAAGTGTGGATGG
Sequence-based reagent	S100a14_F	This paper	qPCR Primer	GGCAGGCTATAGGACA
Sequence-based reagent	S100a14_R	This paper	qPCR Primer	CCTCAGCTCCGAGTAA
peptide, recombinant protein	Fibronectin	Sigma-Aldrich	F4759	50 ug/mL
peptide, recombinant protein	Poly-L-ysine	Sigma-Aldrich	P9155	50 ug/mL

peptide, recombinant protein	Laminin	Thermo Fisher	CB-40232	50 ug/mL
Chemical compound, drug	Latrunculin A	Cayman Chemical Company	10010630	0.5 uM
Commercial assay, kit	ViewRNA ISH Cell Assay kit	Thermo Fisher	QVC0001	
Commercial assay, kit	Click-iTEdU Cell Proliferation Kit for Imaging	Invitrogen	C10337	Alexa Fluor 488 dye
Commercial assay, kit	RNeasy Plus mini kit	QIAGEN	74134	
Commercial assay, kit	TruSeq RNA sample preparation kit	Illumina	RS-122-2001	
Commercial assay, kit	iScript cDNA synthesis kit	BioRad	1708890	
Commercial assay, kit	SYBR Green Supermix	BioRad	170-8882	
Commercial assay, kit	Nextera Library Prep Kit	Illumina	15028212	
Commercial assay, kit	MinElute PCR Purification Kit	QIAGEN	28004	
Software, algorithm	Image J / FIJI	National Institutes of Health		Version 1.50e
Software, algorithm	GraphPad Prism 8.0	GraphPad		Version 8.0

Software, algorithm	PixFRET ImageJ Plugin	(Feige et al., 2005)		
Software, algorithm	Gaussian fit	This paper		https://github.com/LusKingLab/GaussianFit
Software, algorithm	BowTie/TopHat2	(Kim et al., 2013)		
Software, algorithm	DESeq2	(Love et al., 2014)		
Software, algorithm	ENCODE ATAC-seq pipeline	Kundaje Lab	version 1.8.0	https://github.com/ENCODE-DCC/atac-seq-pipeline
other	Prolong Gold with DAPI	Invitrogen	P36935	
other	Sera-Mag Select Beads	GE	29343052	
other	CY 52-276	Dow Corning	52-276	To make 3kPa hydrogels
other	Gil 2 Haematoxylin	Richard Allan Scientific	Cat # 72504	
other	Eosin-Y Alcoholic	Richard Allan Scientific	Cat # 71204	
other	JetPrime	Polyplus transfection	114-07	Transfection reagent

912
913

References:

Adams, J.C., and F.M. Watt. 1989. Fibronectin inhibits the terminal differentiation of human
keratinocytes. *Nature*. 340:307-9.10.1038/340307a0

Alam, S.G., Q. Zhang, N. Prasad, Y. Li, S. Chamala, R. Kuchibhotla, B. Kc, V. Aggarwal, S.
Shrestha, A.L. Jones, S.E. Levy, K.J. Roux, J.A. Nickerson, and T.P. Lele. 2016. The
mammalian LINC complex regulates genome transcriptional responses to substrate rigidity. *Sci*
Rep. 6:38063.10.1038/srep38063

Arsenovic, P.T., I. Ramachandran, K. Bathula, R. Zhu, J.D. Narang, N.A. Noll, C.A. Lemmon,
G.G. Gundersen, and D.E. Conway. 2016. Nesprin-2G, a Component of the Nuclear LINC
Complex, Is Subject to Myosin-Dependent Tension. *Biophys J.* 110:34-
43.10.1016/j.bpj.2015.11.014

Bandyopadhyay, A., G. Rothschild, S. Kim, D.A. Calderwood, and S. Raghavan. 2012.
Functional differences between kindlin-1 and kindlin-2 in keratinocytes. *Journal of Cell Science*.
125:2172-2184.10.1242/jcs.096214

Buenrostro, J.D., B. Wu, H.Y. Chang, and W.J. Greenleaf. 2015. ATAC-seq: A Method for
Assaying Chromatin Accessibility Genome-Wide. *Curr Protoc Mol Biol.* 109:21.29.1-
21.29.9.10.1002/0471142727.mb2129s109

Burgeson, R.E., and A.M. Christiano. 1997. The dermal-epidermal junction. *Curr Opin Cell Biol*.
9:651-8.10.1016/s0955-0674(97)80118-4

Chang, W., H.J. Worman, and G.G. Gundersen. 2015. Accessorizing and anchoring the LINC
complex for multifunctionality. *J Cell Biol.* 208:11-22.10.1083/jcb.201409047

Connelly, J.T., J.E. Gautrot, B. Trappmann, D.W. Tan, G. Donati, W.T. Huck, and F.M. Watt.
2010. Actin and serum response factor transduce physical cues from the microenvironment to
regulate epidermal stem cell fate decisions. *Nat Cell Biol.* 12:711-8.10.1038/ncb2074

Déjardin, T., P.S. Carollo, F. Sipieter, P.M. Davidson, C. Seiler, D. Cuvelier, B. Cadot, C. Sykes,
E.R. Gomes, and N. Borghi. 2020. Nesprins are mechanotransducers that discriminate
epithelial-mesenchymal transition programs. *J Cell Biol.* 219.10.1083/jcb.201908036

Discher, D.E., D.J. Mooney, and P.W. Zandstra. 2009. Growth factors, matrices, and forces
combine and control stem cells. *Science*. 324:1673-7.10.1126/science.1171643

Ezhkova, E., H. Pasolli, J. Parker, N. Stokes, I. Su, G. Hannon, A. Tarakhovsky, and E. Fuchs.
2009. Ezh2 orchestrates gene expression for the stepwise differentiation of tissue-specific stem
cells. *Cell.* 136:1122-35

Feige, J.N., D. Sage, W. Wahli, B. Desvergne, and L. Gelman. 2005. PixFRET, an ImageJ plug-
in for FRET calculation that can accommodate variations in spectral bleed-throughs. *Microsc*
Res Tech. 68:51-8.10.1002/jemt.20215

Gdula, M.R., K. Poterlowicz, A.N. Mardaryev, A.A. Sharov, Y. Peng, M.Y. Fessing, and V.A.
Botchkarev. 2013. Remodeling of three-dimensional organization of the nucleus during terminal

keratinocyte differentiation in the epidermis. *J Invest Dermatol*. 133:2191-
201.10.1038/jid.2013.66

Grashoff, C., B.D. Hoffman, M.D. Brenner, R. Zhou, M. Parsons, M.T. Yang, M.A. McLean, S.G.
Sligar, C.S. Chen, T. Ha, and M.A. Schwartz. 2010. Measuring mechanical tension across
vinculin reveals regulation of focal adhesion dynamics. *Nature*. 466:263-6.10.1038/nature09198

Haque, F., D. Mazzeo, J.T. Patel, D.T. Smallwood, J.A. Ellis, C.M. Shanahan, and S.
Shackleton. 2010. Mammalian SUN protein interaction networks at the inner nuclear membrane
and their role in laminopathy disease processes. *J Biol Chem*. 285:3487-
98.10.1074/jbc.M109.071910

Harmon, R.M., C.L. Simpson, J.L. Johnson, J.L. Koetsier, A.D. Dubash, N.A. Najor, O. Sarig, E.
Sprecher, and K.J. Green. 2013. Desmoglein-1/Erbin interaction suppresses ERK activation to
support epidermal differentiation. *The Journal of Clinical Investigation*. 123:1556-
1570.10.1172/JCI65220

Hotchin, N.A., A. Gandarillas, and F.M. Watt. 1995. Regulation of cell surface beta 1 integrin
levels during keratinocyte terminal differentiation. *J Cell Biol*. 128:1209-
19.10.1083/jcb.128.6.1209

Ihalainen, T.O., L. Aires, F.A. Herzog, R. Schwartlander, J. Moeller, and V. Vogel. 2015.
Differential basal-to-apical accessibility of lamin A/C epitopes in the nuclear lamina regulated by
changes in cytoskeletal tension. *Nat Mater*. 14:1252-61.10.1038/nmat4389

Jung, H.J., A. Tatar, Y. Tu, C. Nobumori, S.H. Yang, C.N. Goulbourne, H. Herrmann, L.G. Fong,
and S.G. Young. 2014. An absence of nuclear lamins in keratinocytes leads to ichthyosis,
defective epidermal barrier function, and intrusion of nuclear membranes and endoplasmic
reticulum into the nuclear chromatin. *Mol Cell Biol*. 34:4534-44.10.1128/mcb.00997-14

Kim, D., G. Pertea, C. Trapnell, H. Pimentel, R. Kelley, and S.L. Salzberg. 2013. TopHat2:
accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions.
Genome Biology. 14:R36.10.1186/gb-2013-14-4-r36

Kumar, A., M. Ouyang, K. Van den Dries, E.J. McGhee, K. Tanaka, M.D. Anderson, A.
Groisman, B.T. Goult, K.I. Anderson, and M.A. Schwartz. 2016. Talin tension sensor reveals
novel features of focal adhesion force transmission and mechanosensitivity. *J Cell Biol*.
213:371-83.10.1083/jcb.201510012

Le, H.Q., S. Ghatak, C.Y. Yeung, F. Tellkamp, C. Günschmann, C. Dieterich, A. Yeroslaviz, B.
Habermann, A. Pombo, C.M. Niessen, and S.A. Wickström. 2016. Mechanical regulation of
transcription controls Polycomb-mediated gene silencing during lineage commitment. *Nat Cell
Biol*. 18:864-75.10.1038/ncb3387

Levy, L., S. Broad, D. Diekmann, R.D. Evans, and F.M. Watt. 2000. beta1 integrins regulate
keratinocyte adhesion and differentiation by distinct mechanisms. *Mol Biol Cell*. 11:453-66

Lien, W.H., X. Guo, L. Polak, L.N. Lawton, R.A. Young, D. Zheng, and E. Fuchs. 2011.
Genome-wide maps of histone modifications unwind in vivo chromatin states of the hair follicle
lineage. *Cell Stem Cell*. 9:219-32.10.1016/j.stem.2011.07.015

Love, M.I., W. Huber, and S. Anders. 2014. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. *Genome Biology*. 15:550.10.1186/s13059-014-0550-8

Luxton, G.W., E.R. Gomes, E.S. Folker, E. Vintinner, and G.G. Gundersen. 2010. Linear arrays
of nuclear envelope proteins harness retrograde actin flow for nuclear movement. *Science*.
329:956-9.10.1126/science.1189072

Luxton, G.W., E.R. Gomes, E.S. Folker, H.J. Worman, and G.G. Gundersen. 2011. TAN lines: a
novel nuclear envelope structure involved in nuclear positioning. *Nucleus*. 2:173-
81.10.4161/nucl.2.3.16243

Mardaryev, A.N., M.R. Gdula, J.L. Yarker, V.U. Emelianov, V.N. Emelianov, K. Poterlowicz, A.A.
Sharov, T.Y. Sharova, J.A. Scarpa, B. Joffe, I. Solovei, P. Chambon, V.A. Botchkarev, and M.Y.
Fessing. 2014. p63 and Brg1 control developmentally regulated higher-order chromatin
remodelling at the epidermal differentiation complex locus in epidermal progenitor cells.
Development. 141:101-11.10.1242/dev.103200

Mertz, A.F., Y. Che, S. Banerjee, J.M. Goldstein, K.A. Rosowski, S.F. Revilla, C.M. Niessen,
M.C. Marchetti, E.R. Dufresne, and V. Horsley. 2013. Cadherin-based intercellular adhesions
organize epithelial cell-matrix traction forces. *Proc Natl Acad Sci U S A*. 110:842-
7.10.1073/pnas.1217279110

Oh, I.Y., D.M. Albea, Z.A. Goodwin, A.M. Quiggle, B.P. Baker, A.M. Guggisberg, J.H. Geahlen,
G.M. Kroner, and C.d.G. Strong. 2014. Regulation of the Dynamic Chromatin Architecture of the
Epidermal Differentiation Complex Is Mediated by a c-Jun|[sol]|AP-1-Modulated Enhancer.
Journal of Investigative Dermatology. 134:2371-2380.doi:10.1038/jid.2014.44

Raghavan, S., C. Bauer, G. Mundschau, Q. Li, and E. Fuchs. 2000. Conditional ablation of
beta1 integrin in skin. Severe defects in epidermal proliferation, basement membrane formation,
and hair follicle invagination. *J Cell Biol*. 150:1149-60

Rippa, A.L., E.A. Vorotelyak, A.V. Vasiliev, and V.V. Terskikh. 2013. The role of integrins in the
development and homeostasis of the epidermis and skin appendages. *Acta Naturae*. 5:22-33

Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch, C.
Rueden, S. Saalfeld, B. Schmid, J.-Y. Tinevez, D.J. White, V. Hartenstein, K. Eliceiri, P.
Tomancak, and A. Cardona. 2012. Fiji: an open-source platform for biological-image analysis.
Nature Methods. 9:676-682.10.1038/nmeth.2019

Schneider, M., W. Lu, S. Neumann, A. Brachner, J. Gotzmann, A.A. Noegel, and I.
Karakesisoglou. 2011. Molecular mechanisms of centrosome and cytoskeleton anchorage at
the nuclear envelope. *Cellular and Molecular Life Sciences*. 68:1593-1610.10.1007/s00018-010-
0535-z

Stewart, R.M., A.E. Zubek, K.A. Rosowski, S.M. Schreiner, V. Horsley, and M.C. King. 2015.
Nuclear-cytoskeletal linkages facilitate cross talk between the nucleus and intercellular
adhesions. *J Cell Biol*. 209:403-18.10.1083/jcb.201502024

Tajik, A., Y. Zhang, F. Wei, J. Sun, Q. Jia, W. Zhou, R. Singh, N. Khanna, A.S. Belmont, and N.
Wang. 2016. Transcription upregulation via force-induced direct stretching of chromatin. *Nat
Mater*. 15:1287-1296.10.1038/nmat4729

Totaro, A., T. Panciera, and S. Piccolo. 2018. YAP/TAZ upstream signals and downstream
responses. *Nat Cell Biol*. 20:888-899.10.1038/s41556-018-0142-z

Tsuruta, D., T. Hashimoto, K.J. Hamill, and J.C. Jones. 2011. Hemidesmosomes and focal
contact proteins: functions and cross-talk in keratinocytes, bullous diseases and wound healing.
J Dermatol Sci. 62:1-7.10.1016/j.jdermsci.2011.01.005

Wang, N., J.D. Tytell, and D.E. Ingber. 2009. Mechanotransduction at a distance: mechanically
coupling the extracellular matrix with the nucleus. *Nat Rev Mol Cell Biol*. 10:75-
82.10.1038/nrm2594

Watt, F.M., M.D. Kubler, N.A. Hotchin, L.J. Nicholson, and J.C. Adams. 1993. Regulation of
keratinocyte terminal differentiation by integrin-extracellular matrix interactions. *J Cell Sci*. 106 (
Pt 1):175-82

Williams, R.R.E., S. Broad, D. Sheer, and J. Ragoussis. 2002. Subchromosomal positioning of
the epidermal differentiation complex (EDC) in keratinocyte and lymphoblast interphase nuclei.
Experimental Cell Research. 272:163-175.10.1006/excr.2001.5400

Zhang, X., K. Lei, X. Yuan, X. Wu, Y. Zhuang, T. Xu, R. Xu, and M. Han. 2009. SUN1/2 and
Syne/Nesprin-1/2 complexes connect centrosome to the nucleus during neurogenesis and
neuronal migration in mice. *Neuron*. 64:173-87.10.1016/j.neuron.2009.08.018

Zhang, X., R. Xu, B. Zhu, X. Yang, X. Ding, S. Duan, T. Xu, Y. Zhuang, and M. Han. 2007.
Syne-1 and Syne-2 play crucial roles in myonuclear anchorage and motor neuron innervation.
Development. 134:901-8.10.1242/dev.02783

Zhu, A.J., I. Haase, and F.M. Watt. 1999. Signaling via beta1 integrins and mitogen-activated
protein kinase determines human epidermal stem cell fate in vitro. *Proc Natl Acad Sci U S A*.
96:6728-33.10.1073/pnas.96.12.6728
1054

Figure 1 A tension sensor in mini-Nesprin2 is under actin-dependent tension when WT mouse keratinocytes engage the extracellular matrix through integrins.

Figure 2 Tension on the N2G-JM-TSMod and nuclear lamina is released at the interior of cohesive mouse keratinocyte (MKC) colonies concomitant with differentiation.

Figure 3
LINC complex ablation leads to precocious epidermal differentiation *in vivo*.

Figure 4 LINC complex ablation leads to precocious MKC differentiation in vivo, which is associated with aberrant up-regulation of differentiation markers at the colony periphery.

Figure 5

LINC complex ablation leads to an increase in chromatin accessibility of genes residing in the Epidermal Differentiation Complex.

Figure 6
Model for the mechanism of precocious differentiation in the Sun dKO epidermis