Single Cell RNA-Sequencing-based Analysis of CD4[+] T-Cell Subset-Specific Susceptibility

to Transcriptional Modulation by HIV-1 Latency-Reversing Agents

Julia Kazmierski[1,2], Dylan Postmus[1,2], Emanuel Wyler[3], Cornelius Fischer[4], Jenny Jansen[1,2],

Karolin Meixenberger[5], Sarah Nathalie Vitcetz[4], Madlen Sohn[4], Sascha Sauer[4], Norbert

Bannert[1,5], Markus Landthaler[3,6], and Christine Goffinet[1,2*]

[1]Institute of Virology, Campus Charité Mitte, Charité - Universitätsmedizin Berlin, Charitéplatz

1, 10117 Berlin, Germany

[2]Berlin Institute of Health, 10178 Berlin, Germany

[3]Berlin Institute for Medical Systems Biology, Max-Delbrück-Center for Molecular Medicine

in the Helmholtz Association, Robert-Rössle-Strasse 10, 13125 Berlin, Germany

[4]Scientific Genomics Platforms, Laboratory of Functional Genomics, Nutrigenomics and

Systems Biology, Max Delbrück Center for Molecular Medicine, 13125 Berlin, Germany.

[5]Division of HIV and Other Retroviruses, Robert Koch Institute, Berlin, Germany

[6]IRI Life Sciences, Institute für Biologie, Humboldt Universität zu Berlin, Philippstraße 13,

10115 Berlin, Germany

*Corresponding author:

Christine Goffinet, Institute of Virology, Campus Charité Mitte, Charité - Universitätsmedizin

Berlin, Charitéplatz 1, 10117 Berlin, Germany

Phone: +0049 30 450 525 489

e-mail: christine.goffinet@charite.de

Keywords: HIV-1, latency-reversing agents, CD4$^+$ T-cells, interferon, interferon-regulated

genes, single cell-RNA sequencing

Abstract

Shock-and-kill is one of the conceptually most advanced strategy towards establishment of an

HIV-1 cure. Treatment with latency-reversing agents (LRAs), including histone deacetylase

inhibitors with chromatin-remodeling capabilities, combined with anti-retroviral therapy,

reactivates HIV-1 transcription *in vivo*. However, LRA treatment fails to significantly reduce

the HIV-1 reservoir in HIV-1-positive individuals, indicating that it is probably insufficient to

eliminate latently infected cells. The global and T-cell subset-specific impact of individual

LRAs on the transcriptome of CD4$^+$ T-cells, the main HIV-1 reservoir containing cell type *in*

vivo, remains understudied. Here, using single cell RNA-sequencing, we characterize LRA

treatment-induced alterations of CD4$^+$ T-cell subset composition and of subpopulation-specific

transcriptomes, using Vorinostat and Panobinostat as two prototypic HDAC inhibitors. *Ex vivo*

exposure of CD4$^+$ T-cells from an aviremic HIV-1-positive individual to Panobinostat

markedly reduced the percentage of T_{REG} cells. Furthermore, it altered expression of a multitude

of interferon-regulated genes, resulting in suppression of several well-characterized antiviral

genes, and in enhancement of selected interferon-regulated genes with proviral activities. These

changes were most pronounced in T_N, T_{CM}, T_{TM} and T_{EM}, and less pronounced in T_{REG}.

Exposure to Vorinostat resulted in a comparably mild change of cellular transcriptomic profile,

regarding both the number of deregulated genes and their fold change of expression.

Nevertheless, selected interferon-regulated genes exhibited a subset-specific expression profile

upon Vorinostat treatment. Finally, some genes were deregulated by both treatments in a subset-

specific manner. We conclude that treatment by both individual HDAC inhibitors induces an

overall proviral milieu in CD4$^+$ T-cells subsets. While this proviral state might be favorable for

efficient HIV-1 reactivation, we hypothesize that it may impede the instruction of activation of

cellular and adaptive immunity required for effective killing of reactivated cells.

Introduction

The development and implementation of highly active antiretroviral therapy (HAART) is

established as an effective tool for suppressing viremia and improving the quality and duration

of life for individuals living with HIV/AIDS. However, though these patients may exhibit

undetectable plasma levels of the virus during optimal treatment, latently infected cells persist

and expand (Vanhamel et al., 2019). In addition, cessation of treatment ultimately results in a

return to a viraemic state within weeks in the vast majority of HIV-1-positive individuals (Hurst

et al., 2015).

The HIV-1 reservoir is maintained by long-lived, latently infected $CD4^+$ T-cells and

cells from the monocyte/macrophage lineage. Among resting $CD4^+$ T-cells carrying latent HIV-

1, T_{CM} and T_{TM} predominate, but latent HIV-1 also resides in T_N and T_{EM} , albeit to a lower

percentage (Chomont et al., 2009). The viral reservoir displays a very long half-life and slow

decay rate, with predicted 73 years of antiretroviral therapy that would be required for

eradication (Finzi et al., 1999; Siliciano et al., 2003). A main mechanism of persistence of the

viral reservoir during antiretroviral treatment is clonal expansion through homeostatic

proliferation, as indicated by a vast majority of $CD4^+$ T-cells in treated patients sharing identical

HIV-1 genome sequence and integration site (Cohn et al., 2015; von Stockenstrom et al., 2015;

Wagner et al., 2014) or TCR sequence (Cohn et al., 2018). Of recent interest as a potential cure

strategy is the so-named shock-and-kill approach. It is based on the administration of latency-

reversing agents (LRAs), including histone deacetylase (HDAC) inhibitors, combined with

antiretroviral treatment, and is anticipated to reactivate viral transcription in these reservoirs

and thus target virus-infected cells for eradication by immunological or pharmacological means

(Kim et al., 2018; Margolis et al., 2016). Clinical trials addressing the ability of individual LRAs

to purge latent HIV-1 *in vivo* demonstrated reactivation of latent HIV-1 but no detectable

reduction of the viral reservoir (Archin et al., 2012; Rasmussen et al., 2014; Sogaard et al.,

2015). It has been suggested that individual LRAs display distinct specificities and potencies

with regards to different T-cell sub-populations (Grau-Exposito et al., 2019), resulting in

suboptimal responses in some, but not other subsets. Furthermore, PHA-reactivated CD4$^+$ T-

cells have been reported to display differential expression of genes related to cellular antiviral

immunity (Cohn et al., 2018). Here, we exposed CD4$^+$ T-cells isolated from an aviremic HIV-

1-positive individual to Vorinostat and Panobinostat, respectively, and investigated, in

individual T-cell subsets and by single cell RNA-sequencing, LRA-specific global

transcriptomic changes and modulation of expression of cellular genes implicated in cell-

intrinsic innate immunity.

Methods & Materials

CD4$^+$ T-Cells from an Aviremic HIV-1 Patient

EDTA blood was collected in the context of the German HIV-1 Seroconverter Study

(Machnowska et al., 2019) with approval of the local ethics committee (Ethical review

committee of Charité University Medicine Berlin, Germany, EA 2/105/05). All study patients

sign an informed consent form prior to enrollment. CD4$^+$ T-cells were isolated using the

EasySep Direct Human CD4$^+$ T Cell Isolation Kit (STEMCELL Technologies). Cells were

cultured at 2×10^6/ml in RPMI 1640 containing 10% heat-inactivated fetal calf serum (Sigma-

Aldrich), 1% penicillin-streptomyin (Thermo Fisher Scientific), 2 mM L-glutamine (Thermo

Fisher Scientific), 1x MEM Non-essential Amino Acid Solution (Thermo Fisher Scientific) and

1mM Sodium Pyruvate (Thermo Fisher Scientific).

Reagents

IL-2 and PHA were purchased from Roche and Oxoid, respectively. Panobinostat was obtained

from Cayman Chem. Vorinostat was obtained through the NIH AIDS Research & Reference

Reagent Program.

IL-2/PHA Stimulation and Treatment with Latency-Reversing Agents

CD4$^+$ T-cells were mock-treated or treated individually with Vorinostat (500 nM), Panobinostat

(50 nM) or IL-2 (20 U/ml)/PHA (1 µg/ml) for 48 hours.

Single cell RNA sequencing

Single Cell RNA-Seq libraries were prepared with the 10x Genomics platform using the

Chromium Next GEM Single Cell 3' Reagent Kits v.3.1 following manufacturer's instructions.

Quality control of the libraries was performed with the KAPA Library Quantification Kit

(KK4854, Roche) and Agilent TapeStation (5067-5584, Agilent). Libraries were sequenced on

a HiSeq 4000 (Illumina) device using the following sequencing mode: read 1: 28 bp, read 2:

91-100 bp, Index i7: 8 bp. The libraries were sequenced to reach ~20 000 reads per cell.

Data Presentation and Statistical Analysis

Cell clustering, cell subset type identification and differential expression analysis

Analysis of the scRNA data was performed using R Studio v3.6 (R Core Team, 2017) and the

Seurat v3.1.4 package (Butler et al., 2018). Filtered Cell Ranger (v 3.1.0) matrices were

imported into R and analysed according to the guidelines set out by the Satija Lab website. The

data were initially filtered to remove cells with fewer than 500 unique detected genes to ensure

adequate sequencing depth, as well as cells with unusually high numbers of gene reads, which

likely represent cell multiplets, and cells with more than 10% mitochondrial DNA reads, as

these represent dead cells. After filtration, a total of 22715 cells remained. The data was log-

normalised and scaled using the LogNormalize and ScaleData functions. Principal component

analysis and linear dimensional reduction analysis using a Uniform Manifold Approximation

and Projection (UMAP) plot were performed, where cells were clustered by means of a Shared

Nearest Neighbor (SNN) graph, based on a K-Nearest Neighbor graph. In general, a resolution

parameter of 1.0 or higher was used to cluster cells in order to retrieve a maximum amount of

clusters. Identities of clusters were determined using a graph-based clustering approach and the

list of pre-determined cell marker genes. Integration of datasets was performed using Seurat's

in-built toolset for finding 'anchors' between datasets (FindIntegrationAnchors) via a canonical

correspondence analysis (CCA). Testing for differential gene expression was performed using

a Wilcoxon sum rank test with a Bonferroni correction. Only genes with a natural log fold-

change of at least ± 0.25 (1.28 or 0.77 fold change) and that were expressed in at least 10% of

cells in either the mock treated or LRA-treated sample were included, where an adjusted p-

value of lower than 0.05 was considered significant. As the analysis focussed on $CD4^+$ T cells,

any clusters identified as $CD8^+$ cells, NK cells ($CD3D^-$, $NKG7^{hi}$, $GNLY^{hi}$) or dead cells (where

the most significantly differentially regulated genes were mitochondrial in origin) were

manually removed.

Cell trajectory and GSEA analysis

Cell trajectory/Pseudotime analysis was performed using the Monocle v2.14.0 package

(Trapnell et al., 2014) according to the guidelines on the Monocle website. Data from the

created Seurat object was imported into Monocle as raw counts (UMI). For each cell subset,

genes identified to be significantly differentially regulated based on the above-mentioned

criteria were used as the ordering genes for the Pseudotime analysis. The minimum spanning

tree was generated using the DDRTree algorithm and the cells arranged along the resulting

Pseudotime trajectory. This was also performed using all genes differentially expressed

between treated and mock-treated cells across all cell subsets. As very few (<10) Treg cells in

the Panobinostat treatment could be identified, these were omitted from the trajectory analysis.

Genes defining the branchpoint observed in the trajectory analysis were determined using a

branch expression analysis model (BEAM). Gene set enrichment analysis (GSEA) was

performed on the same genes used for ordering in the trajectory analysis using the fGSEA

v1.12.0 package for R (Korotkevich et al., 2019). The analysis was performed using 10000

permutations, with the REACTOME curated pathway dataset (Jassal et al., 2020) obtained from

the MSigDB R package (msigdbr) v7.0.1 used as the input pathway list. Of the full lists of

differentially regulated genes for each cell subset and treatment, genes that represented

interferon-regulated genes (IRGs) were identified by cross-referencing these lists with the

Interferome 2.01 database (Rusinova et al., 2013). Differential expression testing was then

performed as described.

Additional data visualisation

Venn diagrams depicting the intersection of genes differentially expressed amongst different

cell types were generated using the venn R package v1.9 (Dusa, 2020), while those depicting

the intersection in different LRA treatments were generated using the VennDiagram R package

v1.6.20 (Chen, 2018). Heatmaps were generated using GraphPad Prism (Version 8.3.0). All

other plots were generated using the ggplot2 R package (Wickham, 2016).

Results

In order to characterize LRA-specific effects and CD4$^+$ T-cell subpopulation-specific susceptibilities to individual LRAs, we exposed purified CD4$^+$ T-cells isolated from an aviremic, HIV-1 positive individual (Table S1) to Vorinostat, Panobinostat, IL-2/PHA or left them mock-treated for 48 hours, and subjected them to single cell RNA-sequencing. We chose to test Vorinostat and Panobinostat due to their demonstrated efficacy to reactivate HIV-1 mRNA expression *in vivo* (Archin et al., 2012; Rasmussen et al., 2014). The employed concentrations and duration of Vorinostat and Panobinostat treatment correspond to the documented ability of the drugs to reactivated HIV-1 *in vitro* (Rasmussen et al., 2013; Tsai et al., 2016). We included IL-2/PHA treatment as a reference that we expected to result in maximum and relatively unspecific activation of T-cells. Finally, we decided to restrict our study on CD4$^+$ T-cells from an aviremic HIV-1-positive individual, as opposed to T-cells from a healthy individual for two reasons. First, this strategy allowed us to consider potential changes of the T-cell composition and transcriptome in the context of a preceding HIV-1 infection and long-term antiretroviral treatment (Wang et al., 2019). Second, we aim to correlate, at the single cell level, identified transcriptomic signatures to patterns and efficiencies of HIV-1 mRNA expression.

We used a graph-based clustering approach and T-cell subset-specific marker genes to assign cells into different subsets (Fig. 1A-C). The cells were grouped into naive CD4$^+$ T-cells (T$_N$; CD8A$^-$, CD3D$^+$, CCR7$^+$, IL7R^{hi}), central memory CD4$^+$ T-cells (T$_{CM}$; CD8A$^-$, CD3D$^+$, CCR7$^+$, S100A4^{int}, IL7R$^+$, SELL^{hi}) (Golubovskaya and Wu, 2016; MacLeod et al., 2009), transitional memory T-cells (T$_{TM}$; CD8A$^-$, CD3D$^+$, CCR7^{lo}, IL7R$^+$, S100A4^{hi}, SELL^{lo}, GZMA$^-$, CD27$^+$) (Mahnke et al., 2013), effector memory CD4$^+$ T-cells (T$_{EM}$; CD8A$^-$, CD3D$^+$, CCR7$^-$, IL7R$^+$, S100A4^{hi}, CD27$^-$, GZMA$^+$) (Golubovskaya and Wu, 2016; Weatherly et al., 2015), CD4$^+$ regulatory T-cells (T$_{REG}$; CD8A$^-$, CD3D$^+$, FOXP3$^+$, IL2RA$^+$) (Golubovskaya and Wu,

2016) and a subset we defined as effector lo CD4$^+$ T-cells (CD8A$^-$, CD3D$^+$, CCR7$^-$, IL7R$^+$,

S100A4^{hi}, CD27$^-$, GZMA^{lo}), which defined a separate cluster expressing low levels of GZMA

but relatively high levels of other effector molecules (GNLY, GZMB).

While exposure to Vorinostat did not grossly alter the CD4$^+$ T-cell subtype composition,

individual treatment with Panobinostat and IL-2/PHA substantially altered relative percentages

of CD4$^+$ T-cell subsets (Fig. 1B-C). Panobinostat treatment induced an almost entire eclipse of

T$_{REG}$ cells (0.1%, versus 6% in mock-treated cells). Furthermore, it enlarged the pool of T$_N$

cells (38.4% versus 26.2% in mock-treated cells). IL-2/PHA treatment resulted in appearance

of a separate cluster of cells we referred to as effector lo CD4$^+$ T$_{EM}$-cells. We hypothesize that

this additional cell cluster represents a transitory state of differentiation between T$_{TM}$ and T$_{EM}$-

cells, which would be in line with the canonical role of IL- 2 as an inducer of T-cell

differentiation towards effector memory status (Ross and Cantrell, 2018) (Figure 1A-C).

LRA- and CD4$^+$ T-cell subset specific modulation of the transcriptome

Integrated analysis of the composition of CD4$^+$ T-cells in the absence and presence of a specific

treatment revealed a noticeably altered topography of the UMAP in the case of treatment with

Panobinostat and IL-2/PHA relative to Vorinostat. Indeed, though Seurat's toolkit for integrated

analysis is robust in promoting cell subset-specific clustering despite treatment-specific effects,

we observed the emergence of several clusters comprising almost entirely of cells from the

LRA treatment (T$_N$ cells in the case of Panobinostat, as well as T$_N$, T$_{CM}$ and T$_{TM}$ cells in the

case of IL2-PHA treatment). Though this observation alone is insufficient to make a solid

conclusion about gene expression profiles, it is in agreement with our observations that the

transcriptomes of Panobinostat and IL-2/PHA treated cells appear to be radically altered in

comparison to the Vorinostat treatment (Fig. 2A).

The number of statistically significantly ($p < 0.05$, $0.78 \leq$ fold-change ≥ 1.28) deregulated

genes varied between individual treatments and among CD4$^+$ T-cell subsets. Exposure to

Vorinostat induced deregulation of relatively few genes in all five subpopulations (Fig. 2B,

Table S2). Panobinostat treatment resulted in altered expression of several hundred genes in

most CD4$^+$ T-cell subsets, except T$_{REG}$ CD4$^+$ T-cells, which presented a lower degree of

responsiveness to this HDAC inhibitor treatment compared to the other subsets. IL-2/PHA

treatment changed the transcriptome in all five sub-populations (Fig. 2B). In all CD4$^+$ T-cell

subsets except T$_{REG}$, most genes deregulated by Vorinostat were also modulated by the two

other treatments (Fig. 2C).

The relatively few genes modulated by Vorinostat were evenly distributed among the

CD4$^+$ T-cell sub-populations, with T$_{REG}$ presenting the highest number of unique differentially

expressed genes (DEGs) (29 DEGs), and T$_{TM}$ displaying not a single specific DEG (Fig. 2D).

All individual T-cell subsets expressed a relatively unique set of 43-170 different genes in the

context of treatment with Panobinostat, and presented some overlap between T$_N$ and T$_{CM}$ (85

DEGs), T$_{TM}$ and T$_{EM}$ (43 DEGs), and T$_{TM}$ and T$_{CM}$ (51 DEGs), indicating that treatment induces

a T-cell subset-specific transcriptomic change (Fig. 2D). All subsets excluding T$_{REG}$ shared 593

DEGs upon Panobinostat treatment. IL-2/PHA treatment induced a large number of DEGs in

all subsets, 876 deregulated in all five sub-populations (Fig. 2D).

In order to investigate whether the altered gene expression profile in the Panobinostat

treated sample induced a single or multiple cell trajectories and which pathways were most

associated with this profile at cell subset level, we performed Pseudotime and gene set

enrichment analysis (GSEA). Genes significantly differentially regulated in the treated versus

mock cells for each subset were used as ordering and input genes, respectively. As there were

very few T$_{REG}$ cells in the Panobinostat treated sample, this cell subset was omitted from the

analysis as we concluded that the results would likely not be conclusive (Fig. 2E). Cell

trajectory analysis revealed that for all of the subsets, except T$_N$ cells, the altered gene

expression profile defines a single cell fate. For T$_N$ cells, branched expression analysis modeling

(BEAM) revealed that the differentially expressed genes defining the branchpoint are

comprised largely of genes encoding ribosomal structural proteins (Supplementary Figure S1).

Interestingly, the positively and negatively enriched pathways in all subsets were similar (Fig.

2F). Generally, positively enriched pathways related to either cell stress, as in the DNA repair,

cellular response to stress and cellular senescence pathways, or to cell proliferation and

productivity, as in the case of the cilium assembly, nonsense mediated decay (NMD), rRNA

processing and organelle biogenesis and maintenance pathways (Han et al., 2018; Sanchez and

Dynlacht, 2016). In contrast, the negatively enriched pathways relate largely to immune

function, such as the TCR signaling, adaptive immune signaling, cytokine immune signaling

and immunoregulatory interactions between a lymphoid and non-lymphoid cell, suggesting that

Panobinostat-treated cells may exhibit an impaired immune response capability. This migration

of cells towards a single cell fate defined by Panobinostat treatment was recapitulated in the

Pseudotime analysis combining all CD4$^+$ T-cell subsets. Interestingly, as the path through

Pseudotime is followed, the root state (containing largely T$_N$ cells) branches off into two cell

fates – a branch containing entirely mock treated cells with a high representation of T$_{EM}$ cells

and a branch containing all of the Panobinostat treated cells. This branching implies that the

trajectory of T$_{EM}$ cells is largely contrarian to the transcriptomic profile induced by treatment,

which is in line with the observation that Panobinostat appears to downregulate immune

response pathways. Simultaneously, despite some cells having a very different transcriptomic

profile to begin with, it becomes clear that Panobinostat induces a single and similar cell fate

across all cell subsets.

LRA treatment alters the expression of IRGs in CD4$^+$ T-cells in a treatment- and subset-

specific manner

In order to better understand how LRA treatment modulates the landscape of interferon-

regulated genes (IRGs), we determined the fraction of all DEGs representing IRGs according

to and using Interferome v2.01 (Rusinova et al., 2013). Interestingly, the majority of DEGs

represented IRGs (80%, 68%, 46% for Vorinostat, Panobinostat and IL-2/PHA, respectively).

Consequently, due to the overall higher degree of gene expression alteration, treatment with IL-

2/PHA and Panobinostat led to a much broader panel of differentially expressed IRGs than did

treatment with Vorinostat (Fig. 3A). While Vorinostat treatment tended to result in a balanced

number of up- and downregulated IRGs, Panobinostat treatment provoked more than twice the

number of downregulated versus upregulated IRGs in T_{CM}, T_{TM} and T_{EM} cells. IL-2/PHA, in

contrast, consistently resulted in a considerably larger ratio (generally at least two- to three-

fold) of upregulated, compared to downregulated, IRGs (Figure 3A).

While Vorinostat treatment resulted in 75 differentially regulated IRGs across all CD4$^+$

T-cell subsets, individual exposure to Panobinostat and IL-2/PHA resulted in 1139 and 1342

differentially expressed IRGs, respectively. Of these, only 5, 29 and 509 for the Vorinostat,

Panobinostat and IL-2/PHA exposure, respectively, were identified as differentially regulated

across all T-cell sub-populations within the individual treatments (Fig. 3B), implying that

alterations in the cellular IRG expression profile are predominantly subset-specific for all

LRAs. Furthermore, the altered IRG profile observed in each cell subtype was largely

determined by the specific LRA. Generally, while there was a sizeable overlap in IRGs

differentially regulated by both Panobinostat and IL-2/PHA, we observed a substantial number

of IRGs whose expression was only affected by treatment with either Panobinostat or IL-

2/PHA. For example, in T_N cells, only 18 IRGs are differentially expressed upon all three

individual treatments, while 412 IRGs were affected by Panobinostat and Il-2/PHA but

Vorinostat and a further 363 and 891 genes were uniquely affected by Panobinostat and IL-

2/PHA, respectively (Figure 3C).

In the context of Vorinostat and Panobinostat treatment, we focused our attention on

significantly deregulated IRGs whose gene products have been characterized to serve as

antiviral restriction factors, to be implicated in downstream signaling upon PRR triggering

through viral PAMPs, and to be otherwise implicated in HIV-1 biology. Strikingly, analysis of

their expression level upon treatment (in comparison to mock treatment), revealed a marked

downregulation in most, but not all subsets. Importantly, some key anti-HIV-1 factors, such as

APOBEC3G (Sheehy et al., 2002), *IFITM1-3* (Compton et al., 2014; Foster et al., 2016; Tartour

et al., 2014), *MX2* (Goujon et al., 2013; Kane et al., 2013; Liu et al., 2013) and *SAMHD1*

(Hrecka et al., 2011; Laguette et al., 2011) were downregulated in the context of Panobinostat

treatment for the majority of cell subsets. *GBP5*, a gene encoding a cellular factor that interferes

with HIV-1 Env processing and maturation (Braun et al., 2019; Krapp et al., 2016) displayed a

subset-specific regulation of expression, with most pronounced downregulation in T_{REG}, weak

downregulation in T_N and T_{TM}, virtually no alteration in T_{CM} and upregulation in T_{EM}.

Interesting exceptions to this trend were *CDC37* and *HSP90AB1*, whose gene products act in

complex on regulation of HIV-1 latency (Anderson et al., 2014) and *LGALS1*, an IRG encoding

a protein facilitating HIV-1 attachment to target cells (Mercier et al., 2008; Ouellet et al., 2005;

St-Pierre et al., 2011). The latter and all three factors were markedly upregulated by Vorinostat

and Panobinostat, respectively, in most T-cell subsets (Fig. 3D-E). Importantly, some key

antiviral factors, such as *TRIM56* (Kane et al., 2016 PMID 27631702), *SUN2* (Donahue *et al.*,

2016 PMID: 26865710   SUN2) and *IFITM1-3* (Tartour et al., 2014 PMID 25422070; Foster

et al., 2016; Compton et al., 2014 PMID:25464829) were downregulated by Panobinostat

treatment for the majority of cell subsets (Figure 3D-E). Taken together, these data imply that

Vorinostat and Panobinostat treatments may result in an abrogated innate immune response and

potentially a more proviral milieu in CD4[+] T-cells.

Discussion

The eminent aim of a LRA treatment in the context of HIV-1 cure is reactivation of HIV-1 mRNA and protein expression. This is supposed to occur through relatively unspecific remodulation of cellular chromatin by currently tested LRAs, boosting transcriptional reactivation of latent HIV-1 genomes but concomitantly resulting in expression changes of a multitude of cellular genes. How those transcriptomic changes of cellular genes indirectly impact HIV-1 reactivation and susceptibility of cells to re-infection remains understudied. An optimal shock-and-kill regimen must be, on the one hand, proviral with regards to all post-integrational steps. It should support or even boost Tat-mediated trans-activation of transcription, protein expression, and cellular trafficking of viral proteins in order to turn previously latent cells in immunologically visible cells which display Env and MHC-presented viral epitopes at the surface. On the other hand, an optimal shock-and-kill strategy must also trigger antiviral immune mechanisms, including cell-mediated immunity by CD8- and NK-cell-mediated cytotoxicity. Along this line, it has been suggested that a reduction of HIV-1 DNA levels during latency reversal is associated with distinct patterns of ISG expression (Olesen et al., 2015). HDAC inhibitors have clearly demonstrated their ability to promote reactivation of HIV-1 mRNA synthesis and protein expression *in vivo* and *in vitro* (Archin et al., 2012; Rasmussen et al., 2013; Rasmussen et al., 2014; Sogaard et al., 2015; Tsai et al., 2016). In *ex vivo*-treated CD4$^+$ T-cells, Vorinostat, Panobinostat and Romidepsin enhanced HIV-1 *de novo* infection (Lucera et al., 2014), raising concerns on potential spread of HIV-1 during shock-and-kill regime in body compartments insufficiently targeted by HAART. On the contrary, another report presented rather an antiviral and no effect of Romidepsin and Panobinostat, respectively (Jonsson et al., 2015). In addition to the missing or controversial knowledge on the net cellular state that is overall induced in LRA-treated CD4$^+$ T-cell cultures, there is no information on subset-specific impact of individual LRAs, and also not if, within a given T-cell subset, there are individual susceptibilities to LRA treatment at the single cell level. Given the preferential

occurrence of latent HIV-1 in T_{CM} and T_{TM}, and potential subtle particularities of cells harboring

latent HIV-1 DNA as opposed to their HIV-1 negative counterparts, the characterization of the

susceptibility to LRA treatment at the greatest possible level of detail is urgently required.

Vorinostat induced rather subtle changes in the CD4$^+$ T-cell subset composition and in

the cellular transcriptome of individual T-cell subpopulations. A previous report using a higher

concentration and shorter treatment duration suggested that genes promoting and inhibiting

HIV-1 transcription were both upregulated in Vorinostat-treated CD4$^+$ T-cells (Beliakova-

Bethell et al., 2019). Based on the current status of our analyses, we observe that several IRGs

with documented anti-HIV-1 activity are downregulated in most T-cell subsets, and one

interferon-modulated gene with a proviral function (*LGALS1*) is upregulated. Based on these

observations, Vorinostat seems to promote an overall rather proviral state. Interestingly, not all

subtypes were equally susceptible to this pattern, pointing to the idea that T_{EM} may be slightly

more resistant to the Vorinostat-mediated suppression of antiviral IRGs. Future studies are

required to identify if this correlates to a less efficient reactivation of latent HIV-1 in this

particular CD4$^+$ T-cell subtype. Interestingly, Vorinostat has been reported to reactivate

endogenous retroviruses (White et al., 2018), and investigation of this issue in our dataset is

imminent.

Exposure of CD4$^+$ T-cells to Panobinostat for 48 hours resulted in an almost entire

disappearance of T_{REG} cells. This finding appears to contradict results obtained upon *in vivo*

administration of Panobinostat to HIV-1 patients in the context clinical latency reversal trials,

which demonstrated a profound and reversible expansion of T_{REG} cells (Brinkmann et al., 2018).

One possibility to reconcile those findings is that Panobinostat´s effect may not be restricted to

CD4$^+$ T-cells *in vivo* and may have triggered expansion of T_{REG} via indirect effects, involving

other cell types than CD4$^+$ T-cells. Importantly, another study suggested that Panobinostat

treatment *in vitro*, and in the context of co-administration with the cytidine analog azacitidine

in vivo, lowered the frequency of a specific T_{REG} subset that is elevated in frequency in the

peripheral blood of acute myeloid leukemia patients (Govindaraj et al., 2014).

Panobinostat treatment clearly modulated the transcriptional landscape of all $CD4^+$

subsets. We currently cannot exclude that the apparently smaller pool of deregulated genes in

T_{REG} may be related to the reduction of frequency of this subset itself. Nevertheless, the other

four T-cell subsets generally shared the downregulation of IRGs with antiviral activity, with

some notable exceptions for individual genes in individual subsets. We are currently analyzing

the T-cell subpopulation-specific expression pattern for HIV-1 latency modulating genes.

The preliminary findings of this study must be interpreted with caution. They are based

on findings obtained from cells from a single donor, and currently performed analyses on cells

from additional donors will be included very soon. Furthermore, the correlation analysis of the

identified, partially subtype-specific cellular transcriptomic patterns to HIV-1 mRNA

expression is an important ongoing task. Interestingly, preliminary experiments indicate that

despite the limited number of cells analyzed, and despite the relative rarity of HIV-1-positive

cells in the peripheral blood of aviremic individuals, this experimental pipeline allows the

identification of reads that map to the HIV-1 genome.

Acknowledgments

We thank the HIV-1-positive individual who agreed to participate to the Seroconverter Study

and provide his blood for research. We thank NIH AIDS Research & Reference Reagent

Program for providing Vorinostat. This work was supported by funding of Berlin Institute of

Health (BIH) to C.G. We thank Christian Drosten for constant support. J.K. is supported by the

Center of Infection Biology and Immunity (ZIBI) and Charité PhD Program.

Author Contributions

J.K. and C.G. conceived and designed the experiments; J.K., J.J., S.N.V. and M.S. performed

the experiments, J.K., D.P., E.W., C.F., K.M. and C.G. analyzed the data; J.K., D.P. and C.G.

drafted the manuscript.; J.K., D.P. and C.G. wrote the manuscript.; S.S., N.B., M.L. and C.G.

supervised the research.

Declaration of Interests

None

Legends

Figure 1. LRA treatment modulates CD4$^+$ T-subset composition

(**A**) UMAP plots of mock-, Vorinostat-, Panobinostat- and IL-2/PHA-treated CD4$^+$ T cells. The

individual CD4$^+$ T-cell subsets are indicated by colouration.

(**B**) Pie charts indicating the relative percentage of individual CD4$^+$ T cell subset for indicated

conditions.

(**C**) Raw counts and relative percentages of individual CD4$^+$ T cell subsets for indicated

conditions.

Figure 2. LRA- and CD4$^+$ T-cell subset specific modulation of the transcriptome

(**A**) UMAP plots of the integrated analysis Vorinostat, Panobinostat and IL-2/PHA treated

samples with the mock treated sample.

(**B**) Volcano plots of all significantly differentially regulated genes in each LRA treatment

relative to the mock sample (minimum log e fold change of ±0.25, Bonferroni corrected p-value

<0.05, Wilcoxon rank-sum test). Upregulated genes are indicated in red, downregulated genes

in blue.

(**C**) Overlap in differentially regulated genes between treatments within each cell subset.

(**D**) Overlap of differentially regulated genes between cell subsets within each treatment.

(**E**) Pseudotime cell trajectory analysis and GSEA analysis using genes differentially regulated

between the mock and Panobinostat treatments for T_N, T_{CM}, T_{TM} and T_{EM} cells, and for total

CD4$^+$ T-cells. Cells are represented as arrayed along the Pseudotime trajectory, coloured by

both treatment and their pseudotime scores.

(**F**) The top five most significantly positively and negatively enriched pathways (based on the

associated p-value) for each cell subset are indicated, along with the corresponding normalised

enrichment score (NES).

Figure 3. LRA treatment alters the expression of IRGs in CD4$^+$ T-cells in a treatment-

and subset-specific manner

(**A**) Volcano plots of all significantly differentially regulated IRGs in each LRA treatment

relative to the mock sample (minimum log e fold change of ±0.25, Bonferronni corrected p-

value <0.05, Wilcoxon rank-sum test). Upregulated genes are indicated in red, downregulated

genes in blue.

(**B**) Overlap of differentially regulated IRGs between cell subsets within each treatment.

(**C**) Overlap in differentially regulated IRGs between treatments within each cell subset.

(**D**) Heat maps indicating fold change in expression between mock and treated samples

for selected IRGs with potential implications for anti-viral activity in the Vorinostat and

Panobinostat treatments. The normalised fold change in IRG expression is shown (expression

in mock treatment = 1). Crossed-out (X) fields represent genes that were not expressed in at

least 10% of either the mock or treated sample. Average fold changes in gene expression for

significantly differentially expressed IRGs are indicated above the corresponding violin plot for

each cell type (Wilcoxon rank-sum test, Bonferroni corrected p-value <0.05).

(**E**) Violin plots showing normalised expression for selected IRGs with potential implications

for anti-viral activity in the Vorinostat and Panobinostat treatments. Numbers indicate fold

change expression of significantly deregulated IRGs.

References

Anderson, I., Low, J.S., Weston, S., Weinberger, M., Zhyvoloup, A., Labokha, A.A., Corazza, G., Kitson, R.A., Moody, C.J., Marcello, A.*, et al.* (2014). Heat shock protein 90 controls HIV-1 reactivation from latency. Proc Natl Acad Sci U S A *111*, E1528-1537.

Archin, N.M., Liberty, A.L., Kashuba, A.D., Choudhary, S.K., Kuruc, J.D., Crooks, A.M., Parker, D.C., Anderson, E.M., Kearney, M.F., Strain, M.C.*, et al.* (2012). Administration of vorinostat disrupts HIV-1 latency in patients on antiretroviral therapy. Nature *487*, 482-485.

Beliakova-Bethell, N., Mukim, A., White, C.H., Deshmukh, S., Abewe, H., Richman, D.D., and Spina, C.A. (2019). Histone deacetylase inhibitors induce complex host responses that contribute to differential potencies of these compounds in HIV reactivation. J Biol Chem *294*, 5576-5589.

Braun, E., Hotter, D., Koepke, L., Zech, F., Gross, R., Sparrer, K.M.J., Muller, J.A., Pfaller, C.K., Heusinger, E., Wombacher, R.*, et al.* (2019). Guanylate-Binding Proteins 2 and 5 Exert Broad Antiviral Activity by Inhibiting Furin-Mediated Processing of Viral Envelope Proteins. Cell Rep *27*, 2092-2104 e2010.

Brinkmann, C.R., Hojen, J.F., Rasmussen, T.A., Kjaer, A.S., Olesen, R., Denton, P.W., Ostergaard, L., Ouyang, Z., Lichterfeld, M., Yu, X.*, et al.* (2018). Treatment of HIV-Infected Individuals with the Histone Deacetylase Inhibitor Panobinostat Results in Increased Numbers of Regulatory T Cells and Limits Ex Vivo Lipopolysaccharide-Induced Inflammatory Responses. mSphere *3*.

Butler, A., Hoffman, P., Smibert, P., Papalexi, E., and Satija, R. (2018). Integrating single-cell transcriptomic data across different conditions, technologies, and species. Nat Biotechnol *36*, 411-420.

Chen, H. (2018). VennDiagram: Generate High-Resolution Venn and Euler Plots. R package version 1.6.20. https://CRANR-projectorg/package=VennDiagram.

Chomont, N., El-Far, M., Ancuta, P., Trautmann, L., Procopio, F.A., Yassine-Diab, B., Boucher, G., Boulassel, M.R., Ghattas, G., Brenchley, J.M.*, et al.* (2009). HIV reservoir size and persistence are driven by T cell survival and homeostatic proliferation. Nat Med *15*, 893-900.

Cohn, L.B., da Silva, I.T., Valieris, R., Huang, A.S., Lorenzi, J.C.C., Cohen, Y.Z., Pai, J.A., Butler, A.L., Caskey, M., Jankovic, M.*, et al.* (2018). Clonal CD4(+) T cells in the HIV-1 latent reservoir display a distinct gene profile upon reactivation. Nat Med *24*, 604-609.

Cohn, L.B., Silva, I.T., Oliveira, T.Y., Rosales, R.A., Parrish, E.H., Learn, G.H., Hahn, B.H., Czartoski, J.L., McElrath, M.J., Lehmann, C.*, et al.* (2015). HIV-1 integration landscape during latent and active infection. Cell *160*, 420-432.

Compton, A.A., Bruel, T., Porrot, F., Mallet, A., Sachse, M., Euvrard, M., Liang, C., Casartelli, N., and Schwartz, O. (2014). IFITM proteins incorporated into HIV-1 virions impair viral fusion and spread. Cell Host Microbe *16*, 736-747.

Dusa, A. (2020). venn: Draw Venn Diagrams. R package version 1.9. https://CRANR-projectorg/package=venn.

Finzi, D., Blankson, J., Siliciano, J.D., Margolick, J.B., Chadwick, K., Pierson, T., Smith, K.,
Lisziewicz, J., Lori, F., Flexner, C., *et al.* (1999). Latent infection of CD4+ T cells provides a
mechanism for lifelong persistence of HIV-1, even in patients on effective combination
therapy. Nat Med *5*, 512-517.

Foster, T.L., Wilson, H., Iyer, S.S., Coss, K., Doores, K., Smith, S., Kellam, P., Finzi, A.,
Borrow, P., Hahn, B.H., *et al.* (2016). Resistance of Transmitted Founder HIV-1 to IFITM-
Mediated Restriction. Cell Host Microbe *20*, 429-442.

Golubovskaya, V., and Wu, L. (2016). Different Subsets of T Cells, Memory, Effector
Functions, and CAR-T Immunotherapy. Cancers (Basel) *8*.

Goujon, C., Moncorge, O., Bauby, H., Doyle, T., Ward, C.C., Schaller, T., Hue, S., Barclay,
W.S., Schulz, R., and Malim, M.H. (2013). Human MX2 is an interferon-induced post-entry
inhibitor of HIV-1 infection. Nature *502*, 559-562.

Govindaraj, C., Tan, P., Walker, P., Wei, A., Spencer, A., and Plebanski, M. (2014).
Reducing TNF receptor 2+ regulatory T cells via the combined action of azacitidine and the
HDAC inhibitor, panobinostat for clinical benefit in acute myeloid leukemia patients. Clin
Cancer Res *20*, 724-735.

Grau-Exposito, J., Luque-Ballesteros, L., Navarro, J., Curran, A., Burgos, J., Ribera, E.,
Torrella, A., Planas, B., Badia, R., Martin-Castillo, M., *et al.* (2019). Latency reversal agents
affect differently the latent reservoir present in distinct CD4+ T subpopulations. PLoS Pathog
15, e1007991.

Han, X., Wei, Y., Wang, H., Wang, F., Ju, Z., and Li, T. (2018). Nonsense-mediated mRNA
decay: a 'nonsense' pathway makes sense in stem cell biology. Nucleic Acids Res *46*, 1038-
1051.

Hrecka, K., Hao, C., Gierszewska, M., Swanson, S.K., Kesik-Brodacka, M., Srivastava, S.,
Florens, L., Washburn, M.P., and Skowronski, J. (2011). Vpx relieves inhibition of HIV-1
infection of macrophages mediated by the SAMHD1 protein. Nature *474*, 658-661.

Hurst, J., Hoffmann, M., Pace, M., Williams, J.P., Thornhill, J., Hamlyn, E., Meyerowitz, J.,
Willberg, C., Koelsch, K.K., Robinson, N., *et al.* (2015). Immunological biomarkers predict
HIV-1 viral rebound after treatment interruption. Nat Commun *6*, 8495.

Jassal, B., Matthews, L., Viteri, G., Gong, C., Lorente, P., Fabregat, A., Sidiropoulos, K.,
Cook, J., Gillespie, M., Haw, R., *et al.* (2020). The reactome pathway knowledgebase.
Nucleic Acids Res *48*, D498-D503.

Jonsson, K.L., Tolstrup, M., Vad-Nielsen, J., Kjaer, K., Laustsen, A., Andersen, M.N.,
Rasmussen, T.A., Sogaard, O.S., Ostergaard, L., Denton, P.W., *et al.* (2015). Histone
deacetylase inhibitor romidepsin inhibits de novo HIV-1 infections. Antimicrob Agents
Chemother *59*, 3984-3994.

Kane, M., Yadav, S.S., Bitzegeio, J., Kutluay, S.B., Zang, T., Wilson, S.J., Schoggins, J.W.,
Rice, C.M., Yamashita, M., Hatziioannou, T., *et al.* (2013). MX2 is an interferon-induced
inhibitor of HIV-1 infection. Nature *502*, 563-566.

Kim, Y., Anderson, J.L., and Lewin, S.R. (2018). Getting the "Kill" into "Shock and Kill":
Strategies to Eliminate Latent HIV. Cell Host Microbe *23*, 14-26.

Korotkevich, G., Sukhov, V., and Sergushichev, A. (2019). Fast gene set enrichment analysis.
biorxiv doi: 101101/060012.

Krapp, C., Hotter, D., Gawanbacht, A., McLaren, P.J., Kluge, S.F., Sturzel, C.M., Mack, K.,
Reith, E., Engelhart, S., Ciuffi, A., *et al.* (2016). Guanylate Binding Protein (GBP) 5 Is an
Interferon-Inducible Inhibitor of HIV-1 Infectivity. Cell Host Microbe *19*, 504-514.

Laguette, N., Sobhian, B., Casartelli, N., Ringeard, M., Chable-Bessia, C., Segeral, E., Yatim,
A., Emiliani, S., Schwartz, O., and Benkirane, M. (2011). SAMHD1 is the dendritic- and
myeloid-cell-specific HIV-1 restriction factor counteracted by Vpx. Nature *474*, 654-657.

Liu, Z., Pan, Q., Ding, S., Qian, J., Xu, F., Zhou, J., Cen, S., Guo, F., and Liang, C. (2013).
The interferon-inducible MxB protein inhibits HIV-1 infection. Cell Host Microbe *14*, 398-
410.

Lucera, M.B., Tilton, C.A., Mao, H., Dobrowolski, C., Tabler, C.O., Haqqani, A.A., Karn, J.,
and Tilton, J.C. (2014). The histone deacetylase inhibitor vorinostat (SAHA) increases the
susceptibility of uninfected CD4+ T cells to HIV by increasing the kinetics and efficiency of
postentry viral events. J Virol *88*, 10803-10812.

Machnowska, P., Meixenberger, K., Schmidt, D., Jessen, H., Hillenbrand, H., Gunsenheimer-
Bartmeyer, B., Hamouda, O., Kucherer, C., Bannert, N., and German, H.I.V.S.S.G. (2019).
Prevalence and persistence of transmitted drug resistance mutations in the German HIV-1
Seroconverter Study Cohort. PLoS One *14*, e0209605.

MacLeod, M.K., Clambey, E.T., Kappler, J.W., and Marrack, P. (2009). CD4 memory T
cells: what are they and what can they do? Semin Immunol *21*, 53-61.

Mahnke, Y.D., Brodie, T.M., Sallusto, F., Roederer, M., and Lugli, E. (2013). The who's who
of T-cell differentiation: human memory T-cell subsets. Eur J Immunol *43*, 2797-2809.

Margolis, D.M., Garcia, J.V., Hazuda, D.J., and Haynes, B.F. (2016). Latency reversal and
viral clearance to cure HIV-1. Science *353*, aaf6517.

Mercier, S., St-Pierre, C., Pelletier, I., Ouellet, M., Tremblay, M.J., and Sato, S. (2008).
Galectin-1 promotes HIV-1 infectivity in macrophages through stabilization of viral
adsorption. Virology *371*, 121-129.

Olesen, R., Vigano, S., Rasmussen, T.A., Sogaard, O.S., Ouyang, Z., Buzon, M., Bashirova,
A., Carrington, M., Palmer, S., Brinkmann, C.R., *et al.* (2015). Innate Immune Activity
Correlates with CD4 T Cell-Associated HIV-1 DNA Decline during Latency-Reversing
Treatment with Panobinostat. J Virol *89*, 10176-10189.

Ouellet, M., Mercier, S., Pelletier, I., Bounou, S., Roy, J., Hirabayashi, J., Sato, S., and
Tremblay, M.J. (2005). Galectin-1 acts as a soluble host factor that promotes HIV-1
infectivity through stabilization of virus attachment to host cells. J Immunol *174*, 4120-4126.

Rasmussen, T.A., Schmeltz Sogaard, O., Brinkmann, C., Wightman, F., Lewin, S.R.,
Melchjorsen, J., Dinarello, C., Ostergaard, L., and Tolstrup, M. (2013). Comparison of HDAC
inhibitors in clinical development: effect on HIV production in latently infected cells and T-
cell activation. Hum Vaccin Immunother *9*, 993-1001.

Rasmussen, T.A., Tolstrup, M., Brinkmann, C.R., Olesen, R., Erikstrup, C., Solomon, A.,
Winckelmann, A., Palmer, S., Dinarello, C., Buzon, M., *et al.* (2014). Panobinostat, a histone

deacetylase inhibitor, for latent-virus reactivation in HIV-infected patients on suppressive
antiretroviral therapy: a phase 1/2, single group, clinical trial. Lancet HIV *1*, e13-21.

Ross, S.H., and Cantrell, D.A. (2018). Signaling and Function of Interleukin-2 in T
Lymphocytes. Annu Rev Immunol *36*, 411-433.

Rusinova, I., Forster, S., Yu, S., Kannan, A., Masse, M., Cumming, H., Chapman, R., and
Hertzog, P.J. (2013). Interferome v2.0: an updated database of annotated interferon-regulated
genes. Nucleic Acids Res *41*, D1040-1046.

Sanchez, I., and Dynlacht, B.D. (2016). Cilium assembly and disassembly. Nat Cell Biol *18*,
711-717.

Sheehy, A.M., Gaddis, N.C., Choi, J.D., and Malim, M.H. (2002). Isolation of a human gene
that inhibits HIV-1 infection and is suppressed by the viral Vif protein. Nature *418*, 646-650.

Siliciano, J.D., Kajdas, J., Finzi, D., Quinn, T.C., Chadwick, K., Margolick, J.B., Kovacs, C.,
Gange, S.J., and Siliciano, R.F. (2003). Long-term follow-up studies confirm the stability of
the latent reservoir for HIV-1 in resting CD4+ T cells. Nat Med *9*, 727-728.

Sogaard, O.S., Graversen, M.E., Leth, S., Olesen, R., Brinkmann, C.R., Nissen, S.K., Kjaer,
A.S., Schleimann, M.H., Denton, P.W., Hey-Cunningham, W.J.*, et al.* (2015). The
Depsipeptide Romidepsin Reverses HIV-1 Latency In Vivo. PLoS Pathog *11*, e1005142.

St-Pierre, C., Manya, H., Ouellet, M., Clark, G.F., Endo, T., Tremblay, M.J., and Sato, S.
(2011). Host-soluble galectin-1 promotes HIV-1 replication through a direct interaction with
glycans of viral gp120 and host CD4. J Virol *85*, 11742-11751.

Tartour, K., Appourchaux, R., Gaillard, J., Nguyen, X.N., Durand, S., Turpin, J., Beaumont,
E., Roch, E., Berger, G., Mahieux, R.*, et al.* (2014). IFITM proteins are incorporated onto
HIV-1 virion particles and negatively imprint their infectivity. Retrovirology *11*, 103.

R Core Team, (2017). R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria; https://wwwR-projectorg/.

Trapnell, C., Cacchiarelli, D., Grimsby, J., Pokharel, P., Li, S., Morse, M., Lennon, N.J.,
Livak, K.J., Mikkelsen, T.S., and Rinn, J.L. (2014). The dynamics and regulators of cell fate
decisions are revealed by pseudotemporal ordering of single cells. Nat Biotechnol *32*, 381-
386.

Tsai, P., Wu, G., Baker, C.E., Thayer, W.O., Spagnuolo, R.A., Sanchez, R., Barrett, S.,
Howell, B., Margolis, D., Hazuda, D.J.*, et al.* (2016). In vivo analysis of the effect of
panobinostat on cell-associated HIV RNA and DNA levels and latent HIV infection.
Retrovirology *13*, 36.

Vanhamel, J., Bruggemans, A., and Debyser, Z. (2019). Establishment of latent HIV-1
reservoirs: what do we really know? J Virus Erad *5*, 3-9.

von Stockenstrom, S., Odevall, L., Lee, E., Sinclair, E., Bacchetti, P., Killian, M., Epling, L.,
Shao, W., Hoh, R., Ho, T.*, et al.* (2015). Longitudinal Genetic Characterization Reveals That
Cell Proliferation Maintains a Persistent HIV Type 1 DNA Pool During Effective HIV
Therapy. J Infect Dis *212*, 596-607.

Wagner, T.A., McLaughlin, S., Garg, K., Cheung, C.Y., Larsen, B.B., Styrchak, S., Huang,
H.C., Edlefsen, P.T., Mullins, J.I., and Frenkel, L.M. (2014). HIV latency. Proliferation of

cells with HIV integrated into cancer genes contributes to persistent infection. Science *345*,
570-573.

Wang, S., Zhang, Q., Hui, H., Agrawal, K., Karris, M.A.Y., and Rana, T. (2019). An Atlas of
Immune Cell Exhaustion in HIV-Infected Individuals Revealed by Single-Cell
Transcriptomics. bioRxiv doi: 101101/678763.

Weatherly, K., Bettonville, M., Torres, D., Kohler, A., Goriely, S., and Braun, M.Y. (2015).
Functional profile of S100A4-deficient T cells. Immun Inflamm Dis *3*, 431-444.

White, C.H., Beliakova-Bethell, N., Lada, S.M., Breen, M.S., Hurst, T.P., Spina, C.A.,
Richman, D.D., Frater, J., Magiorkinis, G., and Woelk, C.H. (2018). Transcriptional
Modulation of Human Endogenous Retroviruses in Primary CD4+ T Cells Following
Vorinostat Treatment. Front Immunol *9*, 603.

A

B

C

Treatment	Mock		Vorinostat		Panobinostat		IL-2/PHA	
	counts	ratio	counts	ratio	counts	ratio	counts	ratio
T_N	1544	26.2%	1815	26.7%	1854	31.5%	1607	27.5
T_{CM}	1594	27.1%	1885	27.8%	1824	26.9%	1895	32.4%
T_{TM}	1854	31.5%	1824	26.9%	1295	30.9%	1158	19.8%
T_{EM}	535	9.1%	782	11.5%	1158	19.8%	450	7.7%
T_{EM} effector lo	0	0%	0	0%	0	0%	493	8.4%
T_{REG}	355	6.0%	484	7.1%	4	0.1%	247	4.2%
Total counts	5882		6790		4193		5850	