Widespread misregulation of inter-species hybrid transcriptomes due to sex-specific and sex-chromosome regulatory evolution

Santiago Sánchez-Ramírez[1*], Jörg G. Weiss[1], Cristel G. Thomas[1,2], Asher D. Cutter[1*]

[1]Department of Ecology and Evolutionary Biology, University of Toronto,

Toronto, ON, M5S3B2, Canada

[2]Los Alamos National Laboratory, Los Alamos, NM, USA

[*]Corresponding authors: santiago.snchez@gmail.com, asher.cutter@utoronto.ca

Santiago Sánchez-Ramírez ORCID = 0000-0002-5963-9410

Asher D. Cutter ORCID = 0000-0001-7141-0013

Abstract

When gene regulatory networks diverge between species, their dysfunctional expression in inter-species hybrid individuals can create genetic incompatibilities that generate the developmental defects responsible for intrinsic post-zygotic reproductive isolation. Both *cis*- and *trans*-acting regulatory divergence can be hastened by directional selection through adaptation, sexual selection, and inter-sexual conflict, in addition to inconspicuous evolution under stabilizing selection. Dysfunctional sex-biased gene expression, in particular, may provide an important source of sexually-dimorphic genetic incompatibilities. Here, we characterize and compare male and female/hermaphrodite transcriptome profiles for sibling nematode species *Caenorhabditis briggsae* and *C. nigoni*, along with allele-specific expression in their F_1 hybrids, to deconvolve features of expression divergence and regulatory dysfunction. Despite evidence of widespread stabilizing selection on gene expression, misexpression of sex-biased genes pervades F_1 hybrids of both sexes. This finding implicates greater fragility of male genetic networks to produce dysfunctional organismal phenotypes. Spermatogenesis genes are especially prone to high divergence in both expression and coding sequences, consistent with a "faster male" model for Haldane's rule and elevated sterility of hybrid males. Moreover, underdominant expression pervades male-biased genes compared to female-biased and sex-neutral genes and an excess of *cis-trans* compensatory regulatory divergence for X-linked genes underscores a "large-X effect" for hybrid male expression dysfunction. Extensive regulatory divergence in sex determination pathway genes likely contributes to feminization of XX hybrids. The evolution of genetic incompatibilities due to regulatory versus coding sequence divergence, however, are expected arise in an uncorrelated fashion. This study identifies important differences between the sexes in how regulatory networks diverge to contribute to sex-biases in how genetic incompatibilities manifest during the speciation process.

Keywords: *Caenorhabditis*, regulatory evolution, sex-biased expression, genome evolution, genetic incompatibilities, hybridization, *cis-trans* regulation, Haldane's rule

Author's summary

As species diverge, many mutations that affect traits do so by altering gene expression. Such gene regulatory changes also accumulate in the control of static traits, due to compensatory effects of mutation on multiple regulatory elements. Theory predicts many of these changes to cause inter-species hybrids to experience dysfunctional gene expression that leads to reduced fitness, disproportionately affecting genes biased toward expression in one sex and that localize to sex chromosomes. Our analyses of genome-wide gene expression from *Caenorhabditis* nematode roundworms support these predictions. We find widespread rewiring of gene regulation, despite the extensive morphological stasis and conserved expression profiles that are hallmarks of these animals. Misregulation of expression in inter-species hybrids of both sexes is most severe for genes linked to the X-chromosome, but male organismal phenotypes are most disrupted in hybrids. This fragility of male genetic networks and sex differences in regulatory evolution of local versus distant elements may underlie feminized and sterile phenotypes among hybrids. Our work clarifies how distinct components of regulatory networks evolve and contribute to sex differences in the manifestation of genetic incompatibilities in the speciation process.

Introduction

Many kinds of reproductive barriers can contribute to speciation (1,2), with genetically intrinsic post-zygotic barriers a kind that makes speciation irreversible. Such intrinsic barriers result from disrupted developmental programs due to divergence in the regulatory controls of and functional activity within genetic networks. Consequently, research for decades has aimed to decipher the identity and general features of genetic changes that accumulate by selection and genetic drift to lead to Dobzhansky-Muller (DM) incompatibilities in hybrids of diverging populations, due to non-additive, negatively-epistatic interactions among genes (1,3). It is therefore crucial to decipher how genes and gene expression evolve to understand how gene regulation influences post-zygotic reproductive isolation through misregulated gene interactions in hybrids (3–5).

Evolution of the regulatory controls over gene expression influences much phenotypic evolution (5,6), despite stabilizing selection as a prevailing force acting to preserve expression profiles (7–11). Expression differences between species accrue in predictable ways. Regulatory differences between species disproportionately involve the evolutionary accumulation of mutations to *cis*-regulatory elements, facilitated by such changes being predisposed to additivity and low pleiotropy of effects on traits and fitness (12,13). In contrast, larger, more pleiotropic effects can result from *trans*-regulatory changes that occur at distant genomic positions, such as to transcription factors, chromatin regulators, and small RNA genes. Consequently, theory predicts *trans*-regulatory mutations to fix less readily and to contribute fewer differences between species, despite their large mutational target size and disproportionate contribution to genetic variation within a species (12–15). Studies nevertheless commonly find both *cis*- and *trans*-regulatory differences between species (16–19). Indeed, the coevolution of changes to both *cis*- and *trans*-acting factors represents one plausible outcome of stabilizing selection on expression level. The compensatory effects of such coevolved *cis*- and *trans*-regulatory changes yield an overall conserved expression profile (10,14,20,21), but this multiplicity of changes are predisposed to generating misexpression in F_1 hybrids due to dysfunctional *cis*-by-*trans* regulatory interactions (5,22).

Decomposing the changes of gene networks into their *cis-* and *trans*–regulatory components,

however, presents a challenge to studying gene regulatory evolution. One way to address this

problem is with hybrid cross experiments that assess differential expression between two

closely related species and from allele-specific expression (ASE) in their F_1 hybrids (5,10,23).

Differences in gene expression between species reveal the joint effects of *cis-* and *trans*-

regulatory divergence, whereas differences in ASE within F_1 hybrids typically represent the

effects of *cis*-regulatory divergence alone (23). Studies of this kind have unveiled broad

empirical patterns of regulatory evolution, whether carried out in flies (16,24), mice (25), plants

(17,18), or yeast (19,26). Overall, previous work has shown substantial regulatory divergence in

both *cis* and *trans*, extensive non-additivity, and disrupted regulation and misexpression in F_1

hybrids. Whether these patterns hold for the nematode phylum is, as yet, unknown, and the

links between regulatory mechanisms and sex-biases in expression remain incompletely

resolved.

Hybrid dysfunction of developmental programs, organismal traits, and fitness may often trace

their origins to gene misregulation, from transcriptional to post-translational levels (5,27). Sex-

biased misregulation, therefore, should underlie sex-biased developmental and fitness effects

in hybrid individuals. Misexpression of male-biased genes and genes related to

spermatogenesis links regulatory disruption to male sterility in hybrids, with supporting

empirical evidence in several kinds of animals (25,28–30). In organisms with chromosomal sex

determination, more severe defects typically occur in hybrid individuals carrying heterogametic

sex chromosomes (i.e., XO males in *Caenorhabditis* nematodes). This Haldane's rule pattern can

arise from dominance effects (31), faster molecular evolution of genes with male-biased

expression (32), greater sensitivity of male developmental programs to perturbation (32), and

faster evolution of sex-linked loci (33), among other causes (34,35). Because of the prominent

role that the X-chromosome plays in reproductive isolation (36–38), we might also expect to

find greater expression divergence and misexpression for X-linked genes compared to

autosomes (5,39–41), with the caveat that genes with male-specific expression might not

necessarily be abundant on the X-chromosome (42–44). Thus, distinguishing between abnormal

expression in hybrids for X-linked genes overall and for sex-biased autosomal genes is

important to decipher the genetic mechanisms that underpin Haldane's rule in particular and

hybrid dysfunction in general.

Caenorhabditis nematode roundworms provide an especially tractable system to study

speciation genomics (35). The growing number of *Caenorhabditis* species known to science

conform to the biological species concept, with a few cases where sibling species can produce

some viable and fertile adult hybrid offspring (45,46). The *C. briggsae* × *C. nigoni* species pair is

one such case, where recent divergence (~3.5 Ma (47)) allows them to form hybrids of both

sexes. In this system, *C. nigoni* has the ancestral gonochoristic reproductive mode (XX females

and X0 males) whereas XX hermaphrodites in the androdioecious *C. briggsae* (also with XO

males) are morphologically female and capable of self-fertilization due to each gonad arm

producing a set of sperm cells prior to an irreversible switch to oocyte production. In hybrids,

both Haldane's rule and Darwin's corollary are fulfilled: F_1 male hybrids are always sterile with

near complete inviability when arising from *C. briggsae* mothers (48–50). In contrast, F_1 females

from both reciprocal crosses are fertile, retaining the "female" condition (i.e., are not

hermaphrodites) (48). This transition in reproductive mode led to the evolution of a suite of

genomic and phenotypic traits in *C. briggsae* that define a "selfing syndrome" due to relaxed

selection on male function and adaptation to a selfing lifestyle (51,52), and has led to the

disproportionate loss of male-biased genes in the *C. briggsae* genome (47,51,53–56). Moreover,

extensive inclusion of X-linked loci in hybrid male sterility implicates a large-X effect (57).

Further analyses revealed that X-autosome incompatibilities involve misregulation of the 22G

class of small RNAs, leading to down-regulation of spermatogenesis genes and contributing to

hybrid male sterility (30,58). The full extent of hybrid misexpression throughout the genome of

each sex and its root causes in regulatory divergence, however, remain to be characterized.

Given the extensive phenotypic stasis within the genus (59) and previous work showing

prevalent action of *cis* and *trans* compensatory evolution, even between more distantly related

species (20), we expect to find substantial developmental systems drift within this system,

where sex-biased genetic networks will be disproportionately prone to misregulation and

misexpression. In particular, we expect greater dysfunctional expression in hybrid males, which

are sterile, compared to hybrid females, which are fertile. These effects also are expected to

associate most strongly with genetic networks that most depend on X-linked genes. To test

these ideas, we analyze mRNA transcriptome expression for each sex from each of *C. briggsae*,

C. nigoni, and their F_1 hybrids. Using ASE profiles, we then characterize and quantify *cis-* and

trans-acting regulatory causes of expression divergence linking genomic change to sex-biased

expression, chromosomal features, and hybrid dysfunction.

Results

Extensive expression divergence between species and between the sexes involve the X-chromosome

Each species and sex show distinctive overall transcriptome profiles that are further distinguishable from each sex of F_1 inter-species hybrids (**Supplementary Fig. S1A**). *C. briggsae* hermaphrodites resemble females phenotypically except that they are able to produce sperm in addition to eggs, and therefore show masculinized expression of some genes (**Supplementary Fig. S1A,2,3**). To more appropriately contrast "female" expression profiles between hermaphrodites and females, we identified 1,238 orthologous genes that are male-biased and up-regulated in hermaphrodites relative to sperm-less *C. briggsae* "pseudo-female" mutants (*she-1(v4)*; (53)) (9% of the 13,636 one-to-one orthologs analyzed). These 1,238 genes were then analyzed separately from most sex-based analyses that, for simplicity, we refer to hereafter as "female" expression contrasts.

The transcriptomes of pure *C. briggsae* and *C. nigoni* differed in expression for more than half of their genes, with slightly more genes differentially expressed for "females" than for males. Females had a total of 61% (7,598) of genes differentially expressed between species, compared to 54% (7,385) for males (**Fig. 1A**). The X-chromosome shows the most extreme differences in number of differentially expressed genes between species, with both males and "females" showing significantly higher ratios of upregulated X-linked genes in *C. nigoni* than in *C. briggsae* (**Fig. 1B**). Autosomes, by contrast, showed greater abundance of genes with higher expression in *C. briggsae*, albeit only significantly for Chromosome I in males (**Fig. 1B**; Fisher's exact test; $P < 0.05$). Similarly, we found more genes with significant ASE for autosomes among female hybrids (6,070 or 61%; n=9,932) than among hybrid males (5,402 or 53%; n=10,155). Which allele gets upregulated in hybrids was largely consistent between the autosomes of hybrid males and females (**Fig. 1B**), albeit slightly more consistent for males (higher expression of *C. briggsae* allele: 2,793 or 27%; *C. nigoni* allele: 2,609 or 26%) than for females (higher expression of *C. briggsae* allele: 3,215 or 32%; *C. nigoni* allele: 2,855 or 29%). The slightly

greater tendency in females for *C. briggsae* alleles to be upregulated could suggest more

regulatory changes fixed in the *C. briggsae* lineage that lead to higher expression, or some

degree of mapping bias in *C. briggsae*-derived reads compared to those from *C. nigoni*

(simulations suggest that mapping biases are an unlikely cause; see Methods).

Within each species, approximately 60% of genes showed significant sex biases in expression

(**Fig. 1C**). Male-biased and female-biased genes occurred with similar incidence in both species:

30% *C. nigoni* male-biased, 29% *C. briggsae* male-biased, 29% *C. nigoni* female-biased, 27% *C.*

briggsae "female"-biased. In stark contrast, over 80% of genes in F_1 hybrids were significantly

sex-biased, with a slightly higher incidence of male-biased genes (43% vs 40%). In *C. briggsae*

and F_1 hybrid transcriptomes, Chromosomes I and III were enriched for "female"-biased genes,

whereas Chromosomes V and X were enriched for male-biased and sex-neutral genes (**Fig. 1D;**

Supplementary Fig. S2). No *C. nigoni* chromosomes exhibited strong enrichment of sex-biased

genes and genes with "female"-biased expression in both species and in hybrids showed strong

depletions from the X-chromosome (**Fig. 1D; Supplementary Fig. S2**).

Expression dominance in F_1 hybrids differs between males and females

We contrasted expression profiles of F_1 hybrids with their parent species to infer the expression

inheritance of genes, i.e., to identify genes that exhibited additive, dominant (*C. briggsae*- and

C. nigoni-like expression), or transgressive (overdominant and underdominant) expression

patterns for each sex (**Fig. 2A**). Gene sets with distinct expression inheritance profiles revealed

substantial differences between the sexes in terms of expression distance (**Fig. 2B**), number of

genes (**Fig. 2C**), and enrichment across the genome (**Fig. 2D**).

The sexes differed most strikingly in their total number of transgressive genes: 26% (3,494) of

genes show transgressive profiles in males (1,788 overdominant and 1,706 underdominant)

versus 55% (6,881) in females (4,121 overdominant and 2,760 underdominant) (**Fig. 2A,C**).

Transgressive genes are thought to be associated with hybrid dysfunction, as such

underdominant and overdominant expression profiles represent misexpression phenotypes

that manifest values beyond the range of both parents (29,60). Given the pronounced sterility

of hybrid males, we were surprised that hybrid male transcriptomes showed only half the

incidence of misexpression as hybrid females. This finding suggests that the genetic networks

that control fitness are more robust to expression perturbation in female hybrids than in males.

In addition, genes classified as underdominant, overdominant, and additive had significantly

higher Euclidean expression distances from the centroid of expression space than genes with no

change in expression or with simple dominance (Gamma-distributed multiple generalized linear

least square regression [GLM], $P < 0.001$) (**Fig. 2A,B**), suggesting that genes with these

expression inheritance profiles are more prone to deviant expression phenotypes. However,

except for transgressively expressed genes, "females" had on average significantly larger

magnitude in expression distance from the centroid expression space than males (GLM, all $P <$

0.001 except transgressive $P > 0.1$) (**Fig. 2A,B**). These results are consistent with our

multidimensional scaling analysis that showed shorter expression distances for F_1 males to

parental males (**Supplementary Fig. S1A**), in contrast to more dissimilar expression profiles of

F1 females to parental females.

Genes with additive expression of alleles from both parental species were relatively rare in F_1

hybrids of both sexes (8% or 691 genes in females; 8% or 946 genes in males) (**Fig. 2C**). Genes

showing simple dominance were up to four-times more common, with approximately 20-30%

of genes expressed by each sex either matching *C. briggsae* or *C. nigoni* expression (23% or

2,869 genes in females; 30% or 3,698 genes in males). Expression dominance matching *C.

briggsae* was consistently more frequent in hybrids of both sexes (15% or 1,876 genes in

females and 18% or 2,228 genes in males), however, compared to expression dominance

matching *C. nigoni* (**Fig. 2C**; 8% or 993 genes in females and 12% or 1,470 in males), being more

extreme in female hybrids across autosomes (mean ratio *C. briggase* vs *C. nigoni* dominant

expression = 1.75 vs 1.14 in males) (**Fig. 2C,D**). The disproportionate dominance of the *C.

briggsae* copy in hybrid females was even more pronounced for the X-Chromosome (F_1 female

X ratio = 2.66 vs F_1 male X ratio = 1.13).

Expression dominance on the X-chromosome is distinct in hybrids of both sexes

Genes and traits with dysfunctional expression are often associated with the X-chromosome, and we expect differences between the sexes due to Haldane's rule (25,28,39,40). Consistent with these expectations, we found inheritance profiles associated with misexpression to differ between F_1 males and females across the genome, and to differ most conspicuously for the X-chromosome. In particular, the X-chromosome was enriched for underdominant genes in both F_1 males and females (**Fig. 2C,D**), but only significantly so in females (Fisher's exact test, $P <$ 0.05). The X-chromosome was also significantly enriched for overdominant genes in F_1 males, whereas females had significant depletions of such genes on the X (autosomal enrichments: V in males, and I and III in females) (**Fig. 2C,D**; Fisher's exact test, $P < 0.05$). These data show clear differences in expression inheritance across chromosomes between the sexes and reflect distinct hybrid expression dynamics between autosomal and X-linked genes.

Given the distinct misexpression of genes linked to the X-chromosome, we evaluated whether disrupted dosage compensation in hybrids might have systematically perturbed X-linked expression profiles. X-chromosome dosage compensation in *Caenorhabditis* acts to halve expression of the two copies in females and hermaphrodites (61). The magnitude of expression that we observed for the X-linked genes in hybrid females, however, does not differ on average from either parental species (**Supplementary Fig. S1B**). Consistent with the implications of conserved mechanistic features of the dosage compensation complex and its regulation across multiple *Caenorhabditis* species, including *C. briggsae* and *C. nigoni* (62), this suggests that dysfunctional dosage compensation is not a key driver of X-linked underdominant expression in hybrid females.

cis and *trans* regulatory divergence modulates differences between sexes

Identifying the spectrum of changes to *cis-* and *trans*-acting regulators is important to understand how selection influences the evolution of gene expression and its effects on hybrid phenotypes. Correspondingly, we classified the types of regulatory changes and examined how

they perturbed gene expression in F_1 hybrids for each sex. Consistent with ASE studies in flies

(16,24), mice (25), plants (17,18) and yeast (19,26), we found substantial expression divergence

due to *cis-only* or *trans*-only regulatory changes in addition to joint effects of *cis* and *trans*

changes, with changes in *trans* and *cis + trans* conferring larger magnitudes of expression

divergence (**Fig. 2E,F**). Genes with *cis*-only and *trans*-only effects were not significantly enriched

on any autosome for either sex, although Chromosome V exhibited a trend toward enrichment

of genes with *trans*-only effects for males (**Fig. 2H**). Comparing regulatory divergence between

sexes, we found nearly 60% more genes involving *trans*-only changes in females (18% or 2,195

genes) compared to males (12% or 1,391 genes) (**Fig. 2G**). For genes with *cis*-only divergence,

however, the sexes showed a reciprocal pattern: *cis*-only divergence was more prevalent in

male than in female transcriptomes (17% or 1,944 genes in males; 14% or 1,647 genes in

females). Thus, the expectation that *cis*-regulatory divergence will be more prevalent than

trans-regulatory divergence holds true only when expression is measured in males, and points

to fundamental differences between the sexes in trends of regulatory evolution.

Furthermore, we categorized genes into 13 groups based on distinct combinations of species

differences and sex differences in expression, including their interactions, and looked at the

proportion of genes with different *cis-* and *trans*-effects (**Fig. 5A,B**). Our results are consistent

with the idea that *cis* and *trans* changes each play distinct roles in sex-biased expression and

sexual dimorphism (63). We find, on one hand, that *trans*-only changes in females are more

often associated with genes that have male-biased expression (37% *trans*-only in male-biased

genes vs. 15% in female-biased genes) and, on the other hand, that female-biased genes show

more conserved regulation (32% conserved in female-biased genes vs. 9% in male-biased

genes) when expressed in males (**Fig. 5D**; **Supplementary Fig. S2**). This pattern suggests that

distinct sex-specific regulatory controls may be asymmetric in this system and may have

evolved to repress expression of genes biased for the opposite sex. Together, these

observations highlight how the evolution sex-specific regulatory controls can underlie sexually

dimorphic expression phenotypes, with implications for sex biases in the speciation process.

Because of codominant effects of *trans*-acting factors controlling the expression of both alleles,

trans-only regulatory divergence is often associated with deviations from additivity (64).

Consistent with this idea, we found *trans*-regulatory changes more often associated with genes

having dominant expression patterns and that genes with additive expression more often

associated with genes having significant *cis*-effects (*cis*-only and *cis* + *trans*) (**Fig. 4**). However,

genes with *cis*-only effects often showed dominant patterns of expression. Genes that show *cis*-

only divergence on autosomes for male expression often have *C. briggsae* or *C. nigoni* dominant

expression in hybrids whereas, in females, they are more typically overdominantly expressed

(**Fig. 4**). We can think of three potential mechanisms that might contribute to this sex difference

and to the *cis*-dominant effect in general: (1) more extensive post-transcriptional regulation in

females (65) might limit transcript abundance through epistatic interactions; (2) greater

degradation or turnover of *cis*-acting binding sites for male-expressed genes (54,55) could

reduce affinity in transcription factor binding activity; and (3) the presence of species-specific

trans regulators that originate from lineage-specific gene duplication and/or loss resulting in

allele-specific *trans*-regulation.

Despite the limitations posed by the hemizygous condition of the X-chromosome in males, we

devised a strategy to confidently assign one of three categories of regulatory divergence to X-

linked genes in males based on their expression inheritance. In short, we inferred genes to have

i) *cis*-only* divergence (inferred from expression dominance with respect to *C. nigoni*, which

may include genes with X-linked *trans* regulators, with this distinction denoted by *), ii) *trans*-

only* divergence with recessive *C. nigoni* *trans*-acting factors (expression dominance with

respect to *C. briggsae*, which may exclude X-linked genes with codominant *trans* regulators,

denoted by *), which provides a lower-bound estimate of *trans*-acting regulatory divergence

affecting the X, and iii) *cis-trans* compensatory changes (no difference in expression between

parent species with hybrids showing over- or underdominance). We did not find enrichments of

cis-only* divergent genes on the X, as might be expected under a "large-X" effect model for

regulatory contributions to hybrid dysfunction. Similarly, we found only a non-significant trend

in both sexes toward enrichment of *trans*-only regulatory divergence affecting X-linked genes

(**Fig. 3**; Fisher exact test, *P* > 0.05). Instead, however, *cis-trans* compensatory changes were 1.9-

fold enriched on the X chromosome for males and 1.7-fold underrepresented on the X for

females (**Fig. 3**; Fisher exact test, *P* < 0.05). These results for regulatory divergence

"regroupings" are consistent with the full set of categories that could be defined for females

(**Fig. 2H**). This finding implicates stabilizing selection on X-linked male-specific networks as

being especially prone to disruption in F_1 hybrids, and contrasts with the 1.6-fold enrichment of

conserved regulatory controls for X-linked genes in females (**Fig. 3**; Fisher exact test, *P* < 0.05).

Hybrid misexpression commonly involves genes with joint *cis-trans* regulatory

changes

Genes exhibiting transgressive expression profiles in F_1 hybrids are often associated with

dysfunctional traits, due to radically different expression from those of parent species.

cis and *trans* changes with opposing effects can interact epistatically in hybrids to induce

transgressive expression and allelic imbalance (22). Such regulatory evolution can arise through

coevolutionary fine-tuning even when overall expression level is subject to stabilizing selection

(6,11). Consistent with this idea, we found that genes with transgressive effects in hybrids often

are associated with *cis-trans* regulatory changes (33% or 877 genes in males and 42% or 2,872

genes in females; **Fig. 4**). In both sexes, F_1 hybrids revealed a higher fraction of compensatory

changes (27% in males and 31% in females) compared to enhancing *cis-trans* changes (6% in

males and 11% in females). These results highlight how stabilizing selection can act differently

on sex-specific transcript abundance, leading to opposing *cis* and *trans* effects that are

dysfunctional in F_1 hybrids. In addition, we also found substantial *cis-trans* compensatory

changes among genes that show no change in expression between species (**Supplementary Fig.**

S3) and in those that do not show sex-biased expression (i.e., C-N-N group in **Fig. 5D**),

implicating extensive developmental systems drift in regulatory processes.

Genes with additive expression inheritance also may generate hybrid dysfunction by generating

intermediate expression profiles in F_1 hybrids, and are thought to commonly reflect *cis*-only

regulatory divergence (23,39,66). In line with this idea, we found that genes classified as

additive associated more often with significant *cis*-acting divergence in both sexes (*cis*-only:

43% in males, 26% in females; and enhancing *cis-trans* effects: 30% in males, 42% in females;

Fig. 4). However, expression additivity is not abundant in our analysis (**Fig. 2C**), suggesting that

it is not a major source of phenotypic dysfunction in hybrids of this system.

To further assess the role that different regulatory controls play in the origin and maintenance

of divergent sex-biased expression, we contrasted expression inheritance and patterns of *cis*-

and *trans*-regulatory divergence for male-biased and female-biased genes (**Fig. 5C,D**). We found

that male-biased genes expressed in F_1 hybrids of both sexes frequently show underdominant

transgressive misexpression compared to sex-neutral and female-biased genes (**Fig. 5A-C**).

Examining male-biased genes when expressed in females, we find that they often show

expression dominance matching the species with lower expression (**Fig. 5A-C**; N-M-I, B-M-I, N-

M-N, B-M-N; also M5, M8, and M14 in **Supplementary Fig. S2**). Additionally, genes that show

conserved expression between species, but with significant sex-bias (C-M-N for male-biased

and C-F-N for female-biased genes), also often are either misexpressed (over- or

underdominant) or have had no significant change in expression in hybrids (**Fig. 5C**). These

groups also have the highest proportion of genes with *cis-trans* compensatory changes (**Fig.

5D**), suggesting that many "conserved" sex-biased networks have undergone substantial

developmental systems drift.

Interestingly, genes expressed in F_1 males are more commonly underdominant when they

correspond to male-biased genes than to female-biased genes (957 genes among male-biased

genes vs 431 genes among female-biased genes), suggesting that male-biased regulatory

networks are more fragile. This idea is supported further by male-biased genes having a higher

proportion of genes with enhancing or compensatory *cis-trans* changes (1,086 or 23% genes in

male-biased genes vs 702 or 18% genes in female-biased genes). In addition, we find that male-

biased genes define more distinct expression profile modules than do female-biased genes (6

versus 3 co-expression modules), including one with male-specific underdominance

(**Supplementary Fig. S2**; M15). By contrast, female-biased genes expressed in F_1 females were

predominantly overdominant and are more often associated with *cis*-only and enhancing *cis-*

trans changes (**Fig. 5D; Fig. 5C, Supplementary Fig. S2**). These observations suggest that female

gene regulatory networks can be more resilient to regulatory divergence, potentially translating

into similar resilience for organismal traits such as fertility (48).

Faster regulatory and molecular evolution of male-biased and spermatogenesis

genes

Sexual selection and sexual conflict are predicted to drive faster rates of molecular evolution

and expression divergence (37,67,68). Consistent with these predictions, we found that male-

biased genes have higher average expression divergence ($|\log_2$-fold-change$|$, GLM, t=-23.639, P

< 0.001) and faster rates of molecular evolution in the two groups of male-biased genes that

are rare on the X-chromosome (K_a/K_s for B-M-I: GLM, t=-8.947, $P < 0.001$; B-M-N: GLM, t=-

8.764, $P < 0.001$) (**Fig. 5E,F**). Male-biased genes show elevated expression divergence compared

to sex-neutral genes and female-biased genes as a whole, though the signal for faster sequence

evolution was weak (K_a/K_s male-biased vs. sex-neutral GLM, t=-0.77, $P = 0.44$; male-biased vs.

female-biased GLM, t=-0.78, $P = 0.43$). Rapid sequence evolution for male-biased genes was not

associated with enrichment in chromosomal arms (**Fig. 5G**, Fisher's exact test, $P < 0.05$), regions

which are known to show higher divergence (47). The groups of male-biased genes with

enrichments on the X-chromosome, however, have either conserved expression between *C.*

briggsae and *C. nigoni* or have higher expression in *C. nigoni* males (**Fig. 5G**; C-M-N, N-M-I, N-M-

N; **Supplementary Fig. S2**; Fisher's exact test, $P < 0.05$), indicating that the X-chromosome is

home to a subset of genes that reflect ancestral male-biased gene networks.

We observed the highest expression divergence as well as high rates of molecular evolution in

the distinctive set of genes that combine male-biased expression, higher expression in *C.*

briggsae than *C. nigoni*, and a species-by-sex interaction (B-M-I; **Fig. 5E,F**). The species-by sex

interaction in this B-M-I group indicates a masculinized expression profile for *C. briggsae*

hermaphrodites, implicating a role for them in sperm production (**Supplementary Fig. S4**; and

M5 in **Supplementary Fig. S2**). To test this idea, we looked at *C. elegans* genes previously

identified as spermatogenesis genes (44) and found overlapping orthologs in *C. briggsae* and *C.*

nigoni to be 11-fold enriched in the B-M-I group (**Fig. 6B and 6C**; Fisher's exact test, *P* < 0.05)

and depleted from the X-chromosome (**Fig. 5G**), consistent with previous observations for

sperm genes in *Caenorhabditis* (35,42,43) and *Drosophila* (69). Further consistent with sperm-

related function, male-biased genes that show upregulated expression in *C. briggsae*

hermaphrodites relative to *C. briggsae* pseudo-females (**Fig. 6A-C**) were highly enriched (46-

fold) and overlapped extensively with the B-M-I group (**Fig. 6B and 6C**; Fisher's exact test, *P* <

0.05). Thus, the high divergence in both expression and sequence for these genes suggests

distinctive selection pressure on them, potentially reflecting the outcome of sexual selection

and sexual conflict on sperm.

The collection of male-biased genes that also show elevated expression in *C. briggsae*

hermaphrodites, putatively linked to spermatogenesis, showed expression phenotypes in

hybrids often resembling *C. nigoni*: extensive expression dominance for the *C. nigoni* copy and

underdominant expression in F_1 hybrids as well as high expression divergence (**Figs. 6D,E;**

Supplementary Fig. S5). They also show an abundance of *trans*-only regulatory divergence (**Fig.**

6F,G).

Regulatory and expression divergence within the sex determination cascade

Sex determination of somatic and germline development involves a negative regulatory cascade

(70), including the secreted protein HER-1 that inhibits transcription factor TRA-2, and further

downstream, TRA-1 represses genes such as *fog-3* in controlling spermatogenesis (71,72).

Consistent with what is known about this pathway (70), our data shows male-biased expression

of *her-1* in both parental species and that it has evolved enhancing *cis* + *trans* regulatory

changes that implicate lineage-specific regulatory changes that promote its expression in males

of both species (**Supplementary Fig. S6**). Without HER-1, TRA-3 then cleaves the intracellular

domain of TRA-2 in XX individuals, which then interacts with FEM proteins preventing FEM–

TRA-1 interactions (72). We found the ortholog of *tra-3* to be similarly expressed and regulated

(*cis*-only) between sexes of pure species and hybrids. However, *fem* genes and *tra-2* seem to be

female-biased and overdominantly expressed (**Supplementary Fig. S6**), suggesting sex-specific

co-regulation; and at least two have potentially evolved *cis* and *trans* regulatory changes (*fem-*

2, fem-3). In addition, antagonistic *cis* x *trans* regulatory effects on *tra-1* seem to have evolved

in females, suggesting that that opposing regulatory changes have evolved on an important

transcription factor controlling germline and somatic sexual differentiation. However, the story

for *tra-1* regulatory divergence in males differs from females and is also less clear: our analysis

categorizes it as "ambiguous" (significant ASE, but non-significant regulatory divergence and

non-significant *trans* effects). In contrast, orthologs of *fog-3*, which is involved in

spermatogenesis, get upregulated in *C. briggsae* hermaphrodites and in males of both species

(B-M-I), as expected. In F_1 males, however, *fog-3* is misexpressed (underdominant) having a

non-significant *trans* effect ($P = 0.28$) and a marginally-significant *cis* ASE effect ($P = 0.04$). Its

expression in F_1 females shows *C. nigoni* dominant expression that is due to *trans*-only

regulatory divergence between the species (**Supplementary Fig. S6**). The regulatory divergence

between *C. briggsae* and *C. nigoni* for genes involved with sex differentiation and development

point to plausible mechanisms for suppression of spermatogenesis in XX hybrids to produce a

"female" rather than a "hermaphrodite" phenotype, as well as yielding hybrid male sterility.

Genome architecture only modestly affects regulatory divergence

Given that protein-coding sequence evolution and gene composition vary non-randomly along

chromosomes in many *Caenorhabditis* species in association with the chromosomal

recombination landscape, we asked whether distinct chromosomal domains would also

associate with the degree of *cis*-regulatory divergence. We find higher molecular divergence

between the genomes of *C. briggsae* and *C. nigoni* in chromosomal arms compared to centers

in noncoding sequences upstream of protein-coding genes (**Supplementary Fig. S7A**), in

addition to protein-coding sequence divergence (**Supplementary Fig. S8**; also see (47)). These

observations are consistent with the idea of stronger purifying selection on mutations to genes

and their *cis*-regulatory regions when linked to chromosome centers or more effective positive

selection when linked to chromosome arms. Despite the elevated molecular divergence in arm

regions, we only found modest elevation of ASE divergence for genes on arms (**Supplementary**

Fig. S7B), as well as modest but significant differences in the magnitude of regulatory

divergence in *cis* and *trans* between chromosome arms and centers (**Supplementary Fig. S7C**;

cis-only GLM [females], coeff=0.06, *t*=1.99, *P* = 0.04; *trans*-only GLM [females], coeff=0.05,

t=4.45, *P* < 0.001). Notably, we observed only a weak positive correlation between ASE

divergence and rates of molecular evolution (**Fig. 7A,B**; linear regression for K_a/K_s': adjusted R =

0.035, *m* = 0.019, *P* < 0.0001; 1-P_{cons}: adjusted R = 0.01, *m* = 0.017, *P* < 0.0001). Overall, these

patterns indicate that rates of divergence for gene expression and their *cis*-regulatory controls

are largely decoupled from protein-coding sequence evolution.

Discussion

Regulatory control over gene expression is an important component of phenotypic evolution (12). As species diverge and accumulate mutations, selection will permit regulatory changes that maintain transcript levels as well as changes that allow exploration of new phenotypic space when they confer a fitness advantage. Sexual selection and sexual conflict can further promote such genomic divergence, both in terms of molecular evolution (e.g., rapid coding or regulatory sequence evolution for male-biased genes) and in terms of gene expression (e.g., divergence in sex-biased gene expression levels) (37,67,68). In interspecies hybrids, sexually driven sources of genomic divergence can disrupt gene networks to create negative epistatic interactions that manifest as sex-biased hybrid sterility or inviability and generate reproductive isolation (5). Here we document extensive regulatory divergence in the face of both conserved and divergent gene expression, with prominent influences of sex-biases and genomic location on the potential to induce misexpression in interspecies hybrids.

Compensatory regulatory evolution implicates pervasive developmental system drift

C. briggsae and *C. nigoni* acquired substantial divergence at the DNA level since they diverged from their common ancestor ~35 million generations ago (3.5 Mya assuming 10 generations per year), including ~20% sequence divergence for synonymous sites, changes to genome size, and disproportionate loss of short male-biased genes in *C. briggsae* since its transition in reproductive mode to androdioecy (47,55). Despite this genomic divergence, hybridization between these species yields viable and fertile F_1 hybrid females, as well as viable hybrid males that suffer complete sterility (35,48).

Despite observing substantial expression divergence, we find that 39% of genes expressed in "females" (4,783 genes) and 46% in males (6,236 genes) show no differential expression between species. Conserved expression between species may result from stabilizing selection, recognized as a common force acting on transcript abundance (7–9,11). Mechanistically,

conservation of the expression phenotype can occur, despite sequence evolution, with co-

evolutionary changes to both *cis-* and *trans*-regulatory elements; for example, if a *trans*-acting

mutation fixes due to a pleiotropic benefit on other loci, selection would favor fixation of any

subsequent compensatory mutation in *cis* that returns expression to optimal levels at the focal

locus (5,20). We find evidence of widespread compensatory *cis-trans* divergence in gene

regulation between *C. briggsae* and *C. nigoni*. Such coevolution represents just one mechanism

leading to "developmental system drift," in which the molecular controls over developmental

pathways can diverge while resulting in little or no change to their phenotypic outputs

(14,21,73). In *Caenorhabditis* nematodes, developmental system drift and stabilizing selection

have been invoked to explain the high degree of phenotypic stasis and morphological constraint

among species (59,74–76). Gene network conservation despite *cis*-regulatory divergence has

been demonstrated by inter-species promoter swaps in *Caenorhabditis*, showing both

robustness in regulatory networks and neo-functionalization in specific cell types (20,77,78), as

well concerted action of *cis* and *trans* compensatory regulation between more distantly related

Caenorhabditis species (20). Our results reinforce this view of pervasive developmental system

drift: we show a high incidence of transgressively expressed genes (overdominant and

underdominant in hybrids) and *cis-trans* compensatory changes in the sexually-dimorphic

transcriptomes of each sex (**Fig. 4**), in addition to an abundance of transgressive hybrid

expression and *cis-trans* compensatory changes among genes with conserved sex-neutral

expression profiles (C-N-N) (**Fig. 5A-D**).

Sequence divergence and developmental system drift in regulatory pathways can lead hybrids

to experience misregulation due to the conflicting regulatory signals from the divergent

genomes, yielding misexpression in hybrid transcriptomes (27). This situation could present a

Dobzhansky-Muller incompatibility if the misexpression leads to reduced fitness; genetic

interactions like those experience by hybrids have been untested by natural selection and will

likely be detrimental (1). The most striking signal of misexpression in our hybrids is the sharp

contrast in the fraction of sex-biased genes: ~83% in hybrids vs ~60% in each parental species

(**Fig. 1C**). The degree of sex-biased gene expression is more extreme for the X chromosome in F_1

hybrids due in part to transgressive underdominance effects (**Figs. 1D, 2C,D**), though the

fraction of male-biased genes likely involved with spermatogenesis is, in fact, depleted on the X

(**Figs. 5G, 6C,E,G**), consistent with previous findings (42,43). The X-chromosome in hybrid males

is enriched, however, for genes showing overdominant expression, which can be linked directly

to *cis-trans* compensatory changes (**Figs. 2D, 3**; **Supplementary Fig. S2**). Overall, these results

implicate extensive sex-limited developmental systems drift that generates extensive

misexpression of genetic networks distinctly in each sex for hybrids of *C. briggsae* and *C. nigoni*.

While expression differences between species are often biased towards one species in systems

such as fruit files and plants (16,18,24), we do not observe much asymmetry toward one

species (**Fig. 1A**; binomial test: males, ratio = 0.5, P = 0.26; females, ratio = 0.47, $P < 0.001$). This

symmetry in expression suggests that demographic effects that exacerbate genetic drift are not

likely to bias regulatory changes toward either increased or decreased expression

disproportionately for one species, as could occur if regulatory changes fix more rapidly in

species like *C. briggsae* with lower effective population sizes due to selfing.

Sex differences in regulatory divergence expose sexual dimorphism of genetic

networks

Transgressive expression in F_1 hybrids beyond the bounds of parental expression levels is a

signature of misexpression, which we observe in abundance. Given that *C. briggsae* × *C. nigoni*

hybrids obey Haldane's rule (48–50), we expected more misexpression in hybrid males. Our

analyses revealed, however, that it is hybrid females that experience more extensive

transgressive gene misexpression (**Fig. 2A-C**). Studies in mice have shown that hybrid

misexpression of X-linked genes confers male sterility (5,28) and *Drosophila* demonstrates an

important influence of X-autosome incompatibilities for Haldane's rule and hybrid male sterility

(36,38), though with a more tenuous link to misexpression of X-linked genes (29,79). We found

enrichment of misexpressed loci on the X-chromosome for both sexes (**Fig. 2D**) and no

indication of compromised dosage compensation. Most female-biased transgressive genes

show overdominant misexpression, however, compared to more underdominant misexpression

in hybrid males (**Fig. 5A-C**). We hypothesize that *cis-* and/or *trans-*regulatory changes acquired

after speciation experienced selection to sustain upregulated expression of female-biased

genes, with those changes behaving in an overdominant manner in hybrids. Indeed,

overdominant genes with *cis-trans* divergence in females have disproportionately evolved

"enhancing" regulatory changes compared to males (**Figs. 2G, 3, 5D**). In spite of the extensive

overdominance in female-biased expression (**Figs. 2B,C, 5C**), the fact that hybrid females are

fertile suggests that overdominant expression does not impact fitness as negatively as does

regulatory divergence that leads to underdominance in hybrids. Consequently, this sexual

dimorphism in gene regulatory networks may contribute to the greater sensitivity of males to

manifesting hybrid dysfunction.

Interestingly, we find that regulatory controls over male-biased genes when they are expressed

in F_1 females largely reveals *trans-*only regulatory changes (**Fig. 5D,E, 6F,G**). These *trans-*acting

regulatory changes are more strongly associated with *C. nigoni* dominant expression in females

among autosomes, which contrasts with these same genes when expressed in males that reveal

extensive *cis-*regulatory changes and *C. briggsae* dominant expression (**Fig. 4**). Previous studies

have shown that *trans-*acting regulation often manifests expression dominantly in hybrids,

possibly as a result of masking effects between dominant and recessive *trans* alleles, whereas

*cis-*acting regulation often generates additive contributions in expression (64). However, *cis-*

dominant expression patterns are not uncommon in ASE studies (18) and may arise, potentially,

from cases where *cis* changes in one species decrease transcription factor binding affinity or

where significant post-transcriptional regulation in pure species is compromised in hybrids.

Merritt et al. (65) showed that *C. elegans* oogenic germline gene networks depend strongly on

3'UTR post-transcriptional regulation, but that genes showing spermatogenic expression rely

primarily on upstream transcriptional regulation. Curiously, disruption of endogenous 22G

smallRNA regulation of spermatogenesis in hybrids of *C. briggsae* x *C. nigoni* also is implicated

in male sterility (30). It may be that sex differences in the dominant mode of regulation

(transcriptional versus post-transcriptional) contributes to the sex differences in the *cis-*

dominance that we report. These results also align with observations of downregulation of

spermatogenesis genes, such as *fog-1* and *fog-3*, by specific transcription factors (i.e., *tra-1*),

and sperm-specific expression depending more on upstream promoter regions than 3-UTRs in

C. elegans (65,71,80).

We observed distinct signatures of regulatory divergence for genes with sex-biased expression

when expressed in the opposite sex, raising the possibility that such divergence may reflect a

history of sexual selection and sexual conflict (51,56). One possible explanation invokes Rice's

hypothesis (81): the fixation of regulatory changes that create sex-biased expression serve as

evolutionary solutions to inter-sexual genomic conflict. To avoid traits that are detrimental to

females but improve male performance, genomic conflict resolution by means of sex-biased

expression may be attained faster through *trans*-regulatory changes, which are more

pleiotropic, downregulating male-biased genes in females. This logic aligns with the hypothesis

that sex-biased expression is partly driven by selection acting to resolve sexual conflict by

means of modifier alleles or regulators (63,81). However, the fact that *trans*-only regulatory

changes do not predominate in the control of female-biased genes in males (**Fig. 4E**), suggests

that regulatory mechanism to resolve genomic sexual conflict act in different ways for the two

sexes. Direct manipulative experiments would prove valuable in testing these ideas further.

The fact that the egg-bearing sex in the *C. briggsae* parent is actually a hermaphrodite may help

explain the presence of underdominant genes in hybrid females. Many of the genes in hybrid

females that show underdominant effects would otherwise show male-biased expression (**Fig.**

5C, 6E), suggesting that they may compromise spermatogenesis to effectively convert F_1

hermaphrodites into females. This perspective provides a complementary view to the idea that

hermaphroditism is 'recessive' to femaleness in a simple Mendelian manner (48). Our

expression data characterizes *tra-1*, a master regulator of sexual development programs

through its repression of genes such as *fog-3* that would promote spermatogenesis and male

development (71), as a conserved sex-neutral gene that is overdominant in both male and

female F_1 hybrids (C-N-N; **Supplementary Fig. S6**). Furthermore, *tra-2* and *fem* genes, which

interact to facilitate TRA-1 activity as a repressor, are all female-biased and overdominantly

expressed in hybrids, which suggest an additional mechanism for hybrid XX "feminization".

Consequently, repression of male-biased genes involved with the *tra-1* pathway is likely to

cause both the "female" phenotype in hybrid females (i.e. absence of male function normally

seen in hermaphrodites) and reduced fertility in hybrid males.

Gene expression in hybrid males predominantly shows either simple dominance or no change

(**Fig. 2C**). While it is tempting to speculate that it might be a byproduct of males of different

species sharing the same reproductive role, reduced sexual selection in *C. briggsae* males

(52,56), genomic divergence (54,55), and clear sex differences in hybrid fertility (48,49), suggest

otherwise. If most transgressive expression occurs in the gonad, then the small and defective

gonad development of F_1 males may have led to their observed paucity of transgressive

expression. Distinct relative sizes of tissues between species, sexes, and hybrids could

potentially influence differences in expression and their interpretation (82). Several aspects of

these *Caenorhabditis* minimize the potential influence of allometric effects between species.

The morphology of *Caenorhabditis* species is highly conserved even between very distantly

related species, with body size being very similar for *C. briggsae* and *C. nigoni* in particular (83);

body size differences within and between species reflect differences in cell size not cell

numbers (59). Our data show symmetry in upregulated genes between species (**Fig. 1A**),

consistent with limited potential role of allometric bias between species. While hybrid females

experience delayed germline development (48), we analyzed adult animals with complete

development. However, the gonad of hybrid males is anatomically deformed, potentially

contributing to differences in expression due to tissue allometry. If disproportionately small

gonad size of hybrid males were to skew expression patterns, then we would expect a large

signal of genes showing low expression specifically in hybrid males. We observe two modules of

co-expression with such a pattern (M3 and M15 in **Supplementary Fig. S2**), but M15 comprises

just 437 (8.7%) of male-biased genes and M3 contains genes with female-biased expression

overall (672 genes, 15% of female-biased genes). Moreover, we observed a high number of

genes with conserved expression across males, including hybrids (**Fig. 2C**), counter to what

would be expected if tissue allometry introduced substantial bias.

Implications for Haldane's rule and the large-X effect

Male-biased genes are expected to evolve fast because of sexual selection and sexual conflict,

resulting in higher rates of protein-coding and gene expression divergence (37,67,68). Faster

evolution of male-biased genes is the premise behind the "faster male" hypothesis to explain

the high incidence of hybrid male sterility in Haldane's rule (32). We find faster evolving coding

sequences for the subset of male-biased genes that also show exceptionally high expression

divergence (B-M-I and B-M-N; **Fig. 5E,F**), many of which are implicated in spermatogenesis (**Fig.**

6A-C; **Supplementary Fig. S4,5**). This result is consistent with previous reports of faster

molecular evolution of spermatogenesis and male germline genes of *C. elegans* (56,84,85), and

suggests that "faster male" evolution may contribute to Haldane's rule in *Caenorhabditis*. To

the extent that rapid evolution predisposes genes to forming genetic incompatibilities,

however, the fact that we rarely observe such genes on the X-chromosome suggests that the

"faster X" model does not provide a compelling explanation for Haldane's rule for hybrid male

sterility (33) (**Fig. 5G**).

The "large-X" effect, where hybrid male sterility results from a disproportionate count of X-

linked loci involved in genetic incompatibilities (86), is evinced from deletion screen

experiments in *Drosophila* (36,38) and introgression experiments in *C. briggsae* x *C. nigoni*

(57,58). An analogous effect can result from X-linked regulatory divergence that causes

misexpression and hybrid dysfunction (25,28,87). Consistent with a large-X effect due to

regulatory evolution in *Caenorhabditis*, our analyses show that the X chromosome contains an

excess of genes that have undergone *cis-trans* compensatory changes causing misexpression in

hybrid males (**Fig. 3**). This pattern implicates a disproportionate role for regulatory divergence

of the X-chromosome in mediating misexpression in *Caenorhabditis* hybrids.

Two non-mutually exclusive ways in which hybrid male dysfunction (i.e., sterility) can arise are:

1) through misexpression and misregulation of X-linked genes involved with male function, and

2) through negative epistatic interactions (i.e., incompatibilities) between X-linked and

autosome genes involved in male-specific pathways. Our results suggest that both cases are

plausible.

First, the paucity of X-linked sex-biased genes in parental genotypes of *Caenorhabditis* species

suggests that any misregulation and misexpression on the X might exert little downstream

impact (**Fig. 1D**; (42,43)). However, misexpression of X-linked genes in hybrids is relatively

common compared to autosomes (with the exception of Chromosome V) (**Fig. 2D**), with hybrid

males having higher relative incidence of effectively misregulated genes (*trans*-only and

compensatory *cis-trans* changes) compared to female hybrids (**Fig. 3**). In fact, we find that

trans-acting factors often contribute to misexpression in both sexes (**Fig. 3, 4**). In hybrid

females, *trans*-acting factors largely drive the expression of X-linked genes with *C. briggsae*

dominant and underdominant expression, unlike autosomes (except Chromosome V) (**Fig. 4**). In

hybrid males, this effect is even more pronounced, in part due to our inference that all X-linked

genes with *C. briggsae* dominant expression arise from *trans*-only* effects that implicate

recessive *C. nigoni trans* regulators (**Supplementary Fig. S14**), which must be at minimum an

underestimation of all *trans*-acting regulatory divergence affecting the X-chromosome. These

findings are consistent with previous observations, particularly in *Drosophila*, of *trans*-acting

sex-specific changes causing misregulation of X-linked genes (39,88,89).

Second, the extensive expression dominance in F_1 males that disproportionately matches the *C.*

briggsae expression level due to *cis* effects have the potential to disrupt gene networks as they

may interact negatively with *C. nigoni* X-linked genes in hybrid males (**Fig. 4; Supplementary**

Fig. S10). Autosomal spermatogenesis genes, by contrast, tend to show *C. nigoni*-dominant

expression in F_1 hybrid males (**Supplementary Fig. S5B,C**), consistent with previous work

showing recessive effects of the *C. briggsae* autosomal portions of genetic incompatibilities

(57). In addition, this prior work also showed that sterility in *C. nigoni* x *C. briggsae* hybrid males

may not require many X-autosome incompatibilities (58). Despite their low abundance on the

X-chromosome, X-linked spermatogenesis genes are often enriched for both misexpression and

misregulation (**Fig. 3; Supplementary Fig. S5B,C**), potentially enhancing their role in hybrid

dysfunction. Our genome-wide transcriptome analysis of *cis-* and *trans*-regulatory divergence

therefore sheds new light on Haldane's rule, reinforcing some previous key inferences about

hybrid dysfunction associated with males, spermatogenesis, and the X-chromosome.

Decoupled coding vs regulatory divergence and the evolution of hybrid dysfunction

Genes that evolve fast may be predisposed to creating genetic incompatibilities, which could

result either from dysfunctional structural activities of the encoded proteins or controls over

the timing or location of gene expression. Studies to date indicate that rates of evolution of

coding sequences and regulatory regions do not correlate strongly (90,91). Supporting the idea

that regulatory divergence due to *cis*-acting elements is largely decoupled from rates of

molecular evolution, we found only weak positive correlations between rates of molecular for

protein structure and gene regulation (**Fig. 7**). *cis*-regulatory divergence also showed only a

weak elevation in chromosome arms compared to centers (**Supplementary Fig. S7B**), genomic

regions with marked differences in recombination rates, gene density, sequence conservation

within and between species (**Supplementary Fig. S7A,8**) (47,92) that influence the rate at which

mutations, especially weakly-selected regulatory mutations, can get fixed as a result of direct

selection and linked selection (93). The decoupled rates of evolution for protein structure and

gene regulation imply that genetic incompatibilities mediated through protein activity versus

gene regulation may follow different rules in the evolution of reproductive isolation.

Conclusion

We contrasted sex-specific transcriptomic profiles between *C. briggsae* and *C. nigoni* and their hybrids to understand how the evolution of *cis-* and *trans*-regulatory elements can contribute to F_1 hybrid dysfunction. Regulatory evolution underlies divergent expression as well as conserved expression subject to compensatory effects of changes to multiple elements. The sharp contrast of extensive misexpression in F_1 hybrids with the morphological stasis and expression conservation between *Caenorhabditis* species indicates substantial developmental system drift of regulatory networks that destabilize in hybrids to enforce reproductive isolation between species. Despite more extensive transgressive expression in hybrid females, they are fertile but unable to produce self-sperm and hybrid males are entirely sterile, suggesting that 1) genetic networks controlling "male" developmental pathways are more fragile in the face of genetic perturbation and 2) hybrid females may represent "demasculinized" hermaphrodites through the disruption of sperm-specific regulatory networks. Despite the rarity of sex-biased genes on the X-chromosome, the X is home to disproportionate misexpression in both sexes, with misregulation in hybrid males largely occurring through *trans*-acting factors and *cis-trans* compensatory changes. X-autosome incompatibilities in hybrid males likely result from the propensity for divergent *cis*-acting factors to drive *C. briggsae*-dominant autosomal expression to yield allele-specific expression biases, which then interact negatively with *C. nigoni* X-linked genes. Moreover, *C. nigoni*-dominant *trans*-acting factors may act to downregulate male-biased genes in both males and females, through the misregulation of master controllers of sexual development such as *tra-1*. Finally, we find only weak correlations of *cis*-regulatory divergence with chromosome architecture and protein-coding and non-coding sequence divergence, indicating that regulatory and protein evolution are largely decoupled. Consequently, Dobzhansky-Muller incompatibilities involving regulatory and coding sequences may accumulate independently of one another, and in distinct ways in the regulatory networks of each sex, in the build-up of reproductive isolation in the speciation process.

Material and Methods

Samples, RNA isolation, and sequencing

We cultured triplicate populations of isofemale *C. briggsae* (AF16) and *C. nigoni* (JU1421) on NGM-agar plates with *Escherichia coli* OP50 at 25°C, isolating total RNA via Trizol-chloroform-ethanol extraction from groups of approximately 500 individual age-synchronized young adult males or females (hermaphrodites) for each replicate sample. *C. briggsae* hermaphrodites are treated as the female sex for the purposes of this study (see below), as their soma is phenotypically female despite the gonad producing a small number of sperm in addition to abundant oocytes. We also crossed in triplicate virgin *C. nigoni* females to male *C. briggsae* (isolated as L4 larvae) to produce F_1 hybrid progeny, with RNA isolated from male and female F_1 hybrids as for the parental pure species genotypes.

The triplicate mRNA samples for each sex and genetic group (*C. briggsae*, *C. nigoni*, F_1 hybrid) underwent 100bp read length, single-ended Illumina HiSeq sequencing at GenomeQuebec according to their standard TruSeq3 protocol. A total of ~250 million reads from these 18 barcoded samples spread across 4 lanes were cleaned for quality control using TRIMMOMATIC v0.38 (with arguments: ILLUMINACLIP:TruSeq3-SE:2:30:10 LEADING:3 TRAILING:3 SLIDINGWINDOW:4:15 MINLEN:36) (94).

Reference alignment and allele-specific read assignment

Following quality control trimming and filtering, we mapped sequence reads from each sample to the chromosome-level genome assembly and annotation of each species (*C. briggsae* WS271 https://osf.io/a4e8g/, *C. nigoni* 2018-01-WormBase https://osf.io/dkbwt/; (55)) using STAR v2.6 (https://github.com/alexdobin/STAR; (95)) with default parameters and adjusting for intron size (--alignIntronMin 40 --alignIntronMax 15600). Reads from *C. briggsae* and *C. nigoni* were mapped to their own reference, while reads from F_1 hybrids were mapped to both reference genomes. Reference genomes and GFF files can also be found online (https://github.com/santiagosnchez/competitive_mapping_workflow/tree/master/references).

To obtain allele-specific read counts in F_1 hybrids, we applied a competitive read mapping

approach by developing a Python implementation for competitive read-mapping we call

COMPMAP (https://github.com/santiagosnchez/CompMap). This approach uses the PYSAM library

(https://github.com/pysam-developers/pysam) internally for reading and indexing BAM

alignments by read name. Using BED files with transcript-level coordinates consistent between

references, read overlaps were then counted for each feature in both alignments. At each

feature, the alignment score (AS) and number of mismatches (nM) of each read to both

references were compared, assigning best aligned to reference-specific counts. Ambiguous

reads (i.e., those being equally good alignments) were also counted and redistributed

proportionally to the number of reads assigned to each reference. We validated our method

with simulated RNA-seq data using the R package POLYESTER (96) finding high correlation

between true ASE and COMPMAP counts, as well as low type 1 and type 2 error rates among *cis*

and *trans* regulatory divergence categories (<5%; **Supplementary Fig. S11,S12**). We expected to

have high power to detect ASE, given ~20% neutral sequence divergence between *C. briggsae*

and *C. nigoni* (47) conferring on average ~5 nucleotide differences for every 100 bp of coding

sequence (0.2 divergence * 0.25 fraction of synonymous sites * 100 bp). Additionally, we did

not expect significant mapping bias (97,98) given that our data was strain-specific (i.e. our C.

briggsae RNA-seq data comes from strain AF16, which is the same as the reference genome and

our *C. nigoni* data comes from strain JU1421 which derives from the same wild isolate as the *C.*

nigoni reference genome; see supplementary data in (55)). Scripts, programs, and commands

used for bioinformatic analyses can be found online

(https://github.com/santiagosnchez/competitive_mapping_workflow).

Ortholog identification and read abundance quantification

The chromosomes in the *C. briggsae* and *C. nigoni* genomes are largely colinear, with only a few

small inversions and translocations reported (99). Therefore, we quantified gene expression

abundance for a set of 13,636 genes that we inferred to be one-to-one reciprocal orthologs

between *C. briggsae* and *C. nigoni*, for which upstream regulatory regions should also be

syntenic. To identify orthologs, we applied a phylogenetic approach using ORTHOFINDER v2.2.6

(100,101), based on longest-isoform peptide sequence translations for gene annotations of 28

Caenorhabditis species (102) (http://caenorhabditis.org/). BLASTp all-by-all searches were done

separately on SciNet's Niagara supercomputer cluster. ORTHOFINDER was run with default

options, which included: -M dendroblast (gene tree reconstruction) and -I 1.5 (MCL inflation

point). In further analysis of the final set of 13,636 orthologs, we excluded from a preliminary

set of 15,461 orthologs those genes for *C. briggsae* and *C. nigoni* that could not be assigned to

any of their six chromosomes (688 genes), that were associated with inter-chromosomal

translocations (370 genes), that we could not estimate K_a/K_s reliably (275 genes), or that

exhibited low mRNA-seq read abundance (492 genes; see below). A list of all the orthologs can

be found online

(https://github.com/santiagosnchez/competitive_mapping_workflow/blob/master/orthologs.tx

t).

We quantified gene expression in parent species for each ortholog with FEATURECOUNTS v2.0.1

(103) as we found its read-counting method to be compatible with COMPMAP. Raw read counts

were combined into a single table and imported into R together with ASE counts (104) for

normalization and statistical analyses. All raw data counts can be found online

(https://github.com/santiagosnchez/competitive_mapping_workflow/tree/master/counts).

Identification of male-biased genes in *C. briggsae* hermaphrodites

C. briggsae hermaphrodites have the ability to produce low amounts of sperm in addition to

oocytes, while their soma phenotypically resembles that of a female. Therefore, to make more

biologically realistic contrasts between *C. briggsae* hermaphrodites, *C. nigoni* females, and

female hybrids, we used an *in silico* approach to identify male-biased genes, which are also

upregulated in hermaphrodites, through comparisons using previously published RNA-seq data.

We used RNA-seq read data derived from *C. briggsae* pseudo-females (AF16-derived *she-1(v47)*

mutant strain) (53), which are unable to produce self-sperm. We conducted differential

expression analyses using a similar pipeline as described below to identify genes with significant

sex-biased expression in our *C. briggsae* RNA-seq data and the one from pseudo-females in

Thomas et al. (53), in addition to being differentially expressed between datasets. These genes

where then excluded from analyses comparing hermaphrodites and females together and were

analyzed separately. Data for these analyses can be found online

(https://github.com/santiagosnchez/competitive_mapping_workflow/tree/master/analyses/ta

bles/Thomas_et_al_data).

Differential expression analyses: contrasts between species, hybrids, and sexes

We used the R Bioconductor package DESEQ2 (105) to assess differential expression. Before

statistically assessing differential expression, we summed the allele-specific counts from F_1

hybrids to yield a single count of transcripts per gene. After calculating library size factors, we

filtered out genes that did not meet the criterion of having at least 3 samples with more than

10 library-size scaled counts. We visualized the overall expression distance between samples

using a non-metric multi-dimensional scaling plot, which showed all three biological replicates

to cluster consistently within their corresponding treatment (**Supplementary Fig. S1A**). We

inferred sex-biased gene expression by comparing differential expression profiles between

males and females (or hermaphrodites) in each genetic group (*C. briggsae*, *C. nigoni*, F_1

hybrids). We also quantified differential expression between the genetic groups in a pairwise

manner (*C. briggsae* vs F_1, *C. briggsae* vs *C. nigoni*, *C. nigoni* vs F_1) for each sex separately. We

then contrasted expression patterns between species (*C. briggsae* and *C. nigoni*) by looking at

sex differences (sex-biased expression) and their interaction (expression ~ species * sex). Within

hybrid males and females, we compared allele-specific counts to measure ASE. For all of these

contrasts, we used the negative binomial generalized linear model fitting and Wald statistics to

determine differentially expressed genes, as implemented in DESEQ2. We used FDR-adjusted *P*-

values at the 5% level to assess significance. DESeq2 results can be found online

(https://github.com/santiagosnchez/competitive_mapping_workflow/tree/master/analyses/ta

bles/DESeq2).

Co-expression clustering

To identify groups of genes with shared co-expression trends, we averaged $\log_2$-transformed

normalized read counts with the *rlog* function from DESEQ2 for each sample. Then we

standardized gene-wise expression data by calculating Z-score values and used *K*-means to

cluster co-expression groups. We chose a sensible *k* value (*k*=15) approaching the asymptote by

plotting the within-group sum of squares for a range of *k* values (from 2 to 100). We then

calculated centroid expression levels by estimating the mean relative expression across samples

within each group. Our co-expression clustering results can be found online

(https://github.com/santiagosnchez/competitive_mapping_workflow/tree/master/analyses/ta

bles/clustering).

Mode of expression inheritance in F_1 hybrids

Based on patterns of expression in F_1 hybrids relative to parent species, we classified genes into

those having **additive** (intermediate), **dominant** (matching either of the species), **overdominant**

(higher that both parents), and **underdominant** (lower than both parents) profiles following the

logic in McManus et al. (16). Genes with no significant differences in expression between F_1s

and their parent species were deemed to have conserved regulatory controls resulting in **no**

change in expression in F_1s. Genes with additive effects had intermediate expression in F_1s

compared to both parental species, meaning that there were significant differences in

expression between all groups in a manner where expression levels in F_1s fall in between both

species. Genes with dominant allelic effects were those with expression levels in F_1s matching

either one of the parent species (i.e. no significant differential expression), but with significant

differential expression between species. Finally, genes with significant differential expression

from both parents, but that were either significantly underexpressed (underdominant) or

overexpressed (overdominant) compared to both species were regarded as transgressive.

Genes falling outside any these specific categories were considered **ambiguous**. A per-gene

summary table with all expression inheritance classification can be found online

(https://github.com/santiagosnchez/competitive_mapping_workflow/tree/master/analyses/ta

bles/expression_inheritance).

We also measured absolute Euclidean distances in expression relative to the centroid or origin

in expression space of F_1 hybrids relative to both parent species. For example, for every gene

we took the expression difference between F_1s and *C. briggsae* and between F_1s and *C. nigoni*

as an xy coordinate system. Then, we measured the Euclidean distance from that point in

expression space to the origin (0,0), reflecting no change in expression:

908
$$d = \sqrt{(\Delta_{F1/Cbr})^2 + (\Delta_{F1/Cbr})^2}$$

Where $\Delta_{F1/Cbr}$ and $\Delta_{F1/Cni}$ are coefficients of differential expression between F_1 hybrids and

each parent species. This metric allowed us to visualize the magnitude of expression distance

from a "conserved" expression profile.

cis- and *trans*- regulatory divergence

We used ASE in F_1s to quantify the extent and type of *cis*- and *trans*-regulatory differences

between species. Expression divergence between parent species results from both *cis*- and

trans-regulatory changes, whereas significant differential expression between alleles in F_1s

results from *cis*-regulatory divergence only (16). To quantify the extent of *trans* effects, we

applied a linear model to test for differences in gene expression between parent species (P) and

between alleles in F_1 hybrids (ASE) using the following model: expression ~ species/group,

where "group" represented categorical variables pointing to data from P and ASE. The division

operator of the function "/" measures expression ratios independently for each category in

"group". We then used a post-hoc Wald-type test (*linearHypothesis* from the CAR package) to

test for significant differences between both coefficients (**P**[*C. nigoni/C. briggsae*] = **ASE**[*C.*

nigoni/C. briggsae]). *P* values were considered significant after a 5% FDR analysis (106).

We inferred the influence of *cis*- and *trans*-regulatory divergence on genes linked to autosomes,

as well as to the X-chromosome in females, following the criteria in McManus et al. (16). This

procedure allowed us to designate genes having undergone significant regulatory divergence

due to **cis-only**, **trans-only**, and **cis-trans** effects. Genes with significant *cis* and *trans* effects

were split into those with synergistic (**cis + trans**), antagonistic (**cis x trans**), and **compensatory**

cis-trans. Genes expressed with no significant differences in expression between parents, ASE,

or *trans* effects were deemed as **conserved** and those that did not strictly fit into any of the

previous groups were considered **ambiguous (Supplementary Fig. S13)**. A table with all gene-

wise classifications for expression inheritance in all chromosomes for females and for

autosomes in males can be found online

(https://github.com/santiagosnchez/competitive_mapping_workflow/tree/master/analyses/ta

bles/cis_trans).

Given the hemizygous condition of the X-chromosome in males, we cannot use F_1 ASE of X-

linked genes to assess *cis* and *trans* regulatory divergence. However, we devised a scheme to

assign different types of regulatory divergence to X-linked genes, with some limitations, based

on the differences in expression between male F_1 hybrids and parent species (**Supplementary**

Fig. S14; Wayne et al. 2004). F_1 males of our study carry their maternal *C. nigoni* X-

chromosome. Therefore, assuming that the bulk of the *trans* environment derives from

autosomes, we would expect that X-linked genes that differ in expression between parental

species but display *C. nigoni* dominant expression owe their regulatory divergence mostly to

cis-regulatory (**cis-only***) changes, because any significant deviation from *C. nigoni* expression

levels would indicate *trans*-regulatory changes. An exception would involve the action of "local"

trans regulators found on the X, either way still considered "local". Genes with significant

regulatory divergence between species where hybrids display *C. briggsae* dominant expression

(**trans-only***) would indicate no significant changes due to *cis* regulation, with the condition

that autosomal *C. nigoni trans* regulators affecting those genes are recessive, and therefore not

expressed. We consider these a subset of total *trans* regulation and is likely an underestimation

compared to autosomes and X-chromosomes expressed in females. The last category that we

can assign confidently is **compensatory *cis-trans* changes**. Genes with no differential expression

between species, but with significant up- or down-regulation in F1 males (i.e., overdominant or

underdominant) were considered as having *cis-trans* compensatory changes. For the purpose of

comparing regulatory divergence between autosomes and the X in males and females, we

lumped all other genes with significant expression divergence into "other", as they would be

difficult to disambiguate without additional data. These include genes with *trans*-only (i.e., with

codominant *trans* regulation), *cis* + *trans* (enhancing), *cis* x *trans* (compensatory), and

ambiguous. Genes with no expression change were kept as "conserved" (**Supplementary Fig.**

S12).

Molecular evolution in coding sequences

Orthologs in the genomes of both *C. briggsae* and *C. nigoni* were first aligned as protein coding

sequences using MAFFT v7.407 (107). These alignments were then back-translated to coding

sequence (CDS) alignments using the python program CODONALIGN

(https://github.com/santiagosnchez/CodonAlign). We estimated rates of synonymous site (K_s)

and non-synonymous site divergence (K_a) between the two aligned sequences using a custom

Python script (https://github.com/santiagosnchez/DistKnKs) applying the Yang and Nielsen

(2000) model implemented in BioPython (Cock et al. 2009). We also corrected K_s values for

selection on codon usage using the effective number of codons (ENC; (108,109)) as a predictor

in a linear model. In short, we fitted a linear regression model (K_s ~ ENC), which we used to

predict K_s at the maximum value of ENC (=60). Then, we corrected the bias in K_s by adding the

residuals of the linear model to that idealized value of K_s at ENC = 60. We refer to these

corrected set of K_s estimates as K_s'. A table with these estimates of molecular evolution for each

gene can be found online

(https://github.com/santiagosnchez/competitive_mapping_workflow/tree/master/analyses/ta

bles/molecular_evolution).

Upstream non-coding sequence conservation

Chromosome-level FASTA sequences for *C. briggsae* and *C. nigoni* were aligned using LASTZ

(110), outputting alignment files for each chromosome in MAF format. We used BEDTOOLS's

v2.27 (111) *flank* function to generate 500 bp intervals of the 5' upstream flanking regions of

each orthologous gene. We then used *maf_parse* from PHAST (112) to extract overlapping

alignment blocks of at least 500 bp long. We quantified sequence conservation as the average

number of identical 5 bp non-overlapping windows between aligned DNA in both sequences.

Spermatogenesis genes

To infer genes involved with spermatogenesis, we downloaded a list of *C. elegans* genes

previously identified as spermatogenesis-related based on tissue-specific transcript abundance

(44) ([Additional File 4](#)). We then used the BioMart tool of the WormBase Parasite website (113)

to retrieve *C. briggsae* orthologs from the list of *C. elegans* genes. We cross-referenced *C.*

briggsae orthologs to our own set of orthologs between *C. briggsae* and *C. nigoni* and

annotated the 1,089 gene matches with a spermatogenesis tag. The data obtained for *C.*

elegans spermatogenesis orthologs can be found online

(https://github.com/santiagosnchez/competitive_mapping_workflow/tree/master/analyses/ta

bles/spermatogenesis).

1000

1001 Data availability

1002 Raw sequencing data can be found on the Short Read Archive under accessions SRR12532505-

1003 SRR12532522 and BioProject PRJNA659254. Raw gene counts, workflow scripts, and main data

1004 tables and files can be found on GitHub under the following repository:

1005 https://github.com/santiagosnchez/competitive_mapping_workflow.

1006 Acknowledgements

We thank Katja Kasimatis and Karl Grieshop for comments on previous versions of the

manuscript. Computations were performed on the Niagara supercomputer at the SciNet HPC

Consortium. SciNet is funded by the Canada Foundation for Innovation, the Government of

Ontario, the Ontario Research Fund - Research Excellence, and the University of Toronto. S.S.R.

is supported by an EEB Departmental Postdoctoral Fellowship. A.D.C. is supported by a

Discovery Grant from the Natural Sciences and Engineering Research Council of Canada.

1013

References

1. Coyne JA, Orr HA. Speciation. Sinauer Associates Incorporated: Sinauer Associates Incorporated; 2004.

2. Rundle HD, Nosil P. Ecological speciation. Ecol Lett. 2005;8(3):336–52.

3. Orr HA. The population genetics of speciation: the evolution of hybrid incompatibilities. Genetics. 1995;139(4):1805–13.

4. Gordon KL, Ruvinsky I. Tempo and mode in evolution of transcriptional regulation. PLoS Genet. 2012;8(1):e1002432.

5. Mack KL, Nachman MW. Gene regulation and speciation. Trends Genet. 2017;33(1):68–80.

6. Lemos B, Meiklejohn CD, Cáceres M, Hartl DL. Rates of divergence in gene expression profiles of primates, mice, and flies: stabilizing selection and variability among functional categories. Evolution. 2005;59(1):126–37.

7. Denver DR, Morris K, Streelman JT, Kim SK, Lynch M, Thomas WK. The transcriptional consequences of mutation and natural selection in *Caenorhabditis elegans*. Nat Genet. 2005;37(5):544–8.

8. Rifkin SA, Houle D, Kim J, White KP. A mutation accumulation assay reveals a broad capacity for rapid evolution of gene expression. Nature. 2005;438(7065):220–3.

9. Gilad Y, Oshlack A, Rifkin SA. Natural selection on gene expression. Trends Genet. 2006;22(8):456–61.

10. Signor SA, Nuzhdin S V. The evolution of gene expression in *cis* and *trans*. Trends Genet. 2018;34(7):532–44.

11. Bedford T, Hartl DL. Optimization of gene expression by natural selection. Proc Natl Acad Sci U S A. 2009;106(4):1133–8.

12. Wray GA. The evolutionary significance of *cis*-regulatory mutations. Nat Rev Genet. 2007;8(3):206–16.

13. Stern DL, Orgogozo V. The loci of evolution: how predictable is genetic evolution? Evolution. 2008;62(9):2155–77.

14. Gordon KL, Ruvinsky I. Tempo and mode in evolution of transcriptional regulation. PLoS Genet. 2012;8(1):e1002432.

15. Wittkopp PJ. Genomic sources of regulatory variation in *cis* and in *trans*. Cell Mol Life Sci. 2005;62(16):1779–83.

16. McManus CJ, Coolon JD, Duff MO, Eipper-Mains J, Graveley BR, Wittkopp PJ. Regulatory divergence in *Drosophila* revealed by mRNA-seq. Genome Res. 2010;20(6):816–25.

17. Shi X, Ng DW-K, Zhang C, Comai L, Ye W, Chen ZJ. *cis*- and *trans*-regulatory divergence between progenitor species determines gene-expression novelty in *Arabidopsis* allopolyploids. Nat Commun. 2012;3(1):950–9.

18. Bell GDM, Kane NC, Rieseberg LH, Adams KL. RNA-seq analysis of allele-specific expression, hybrid effects, and regulatory divergence in hybrids compared with their parents from natural populations. Genome Biol Evol. 2013;5(7):1309–23.

19. Tirosh I, Reikhav S, Levy AA, Barkai N. A yeast hybrid provides insight into the evolution of gene expression regulation. Science. 2009;324(5927):659–62.

20.   Barrière A, Gordon KL, Ruvinsky I. Coevolution within and between regulatory loci can
preserve promoter function despite evolutionary rate acceleration. PLoS Genet.
2012;8(9):e1002961-13.
21.   True JR, Haag ES. Developmental system drift and flexibility in evolutionary trajectories.
Evol Dev. 2001;3(2):109–19.
22.   Landry CR, Wittkopp PJ, Taubes CH, Ranz JM, Clark AG, Hartl DL. Compensatory *cis-trans*
evolution and the dysregulation of gene expression in interspecific hybrids of *Drosophila*.
Genetics. 2005;171(4):1813–22.
23.   Wittkopp PJ, Haerum BK, Clark AG. Evolutionary changes in *cis* and *trans* gene regulation.
Nature. 2004;430(6995):85–8.
24.   Wittkopp PJ, Haerum BK, Clark AG. Regulatory changes underlying expression differences
within and between *Drosophila* species. Nat Genet. 2008;40(3):346–50.
25.   Mack KL, Campbell P, Nachman MW. Gene regulation and speciation in house mice.
Genome Res. 2016;26(4):451–61.
26.   Emerson JJ, Hsieh LC, Sung HM, Wang TY, Huang CJ, Lu HHS, et al. Natural selection on *cis*
and *trans* regulation in yeasts. Genome Res. 2010;20(6):826–36.
27.   Ortiz-Barrientos D, Counterman BA, Noor MAF. Gene expression divergence and the
origin of hybrid dysfunctions. Genetica. 2006;129(1):71–81.
28.   Turner LM, White MA, Tautz D, Payseur BA. Genomic networks of hybrid sterility. PLoS
Genet. 2014;10(2):e1004162--23.
29.   Gomes S, Civetta A. Hybrid male sterility and genome- wide misexpression of male
reproductive proteases. Sci Rep. 2015;5(1):1–11.
30.   Li R, Ren X, Bi Y, Ho VWS, Hsieh C-L, Young A, et al. Specific down-regulation of
spermatogenesis genes targeted by 22G RNAs in hybrid sterile males associated with an
X-Chromosome introgression. Genome Res. 2016;26(9):1219–32.
31.   Turelli M, Orr HA. The dominance theory of Haldane's rule. Genetics. 1995;140(1):389–
402.
32.   Wu CI, Davis AW. Evolution of postmating reproductive isolation: the composite nature
of Haldane's rule and its genetic bases. Am Nat. 1993;142(2):187–212.
33.   Charlesworth B, Coyne JA, Barton NH. The relative rates of evolution of sex
chromosomes and autosomes. Am Nat. 1987;130(1):113–46.
34.   Delph LF, Demuth JP. Haldane's rule: genetic bases and their empirical support. J Hered.
2016;107(5):383–91.
35.   Cutter AD. X exceptionalism in *Caenorhabditis* speciation. Mol Ecol. 2018;27(19):3925–
34.
36.   Masly JP, Presgraves DC. High-resolution genome-wide dissection of the two rules of
speciation in *Drosophila*. PLoS Biol. 2007;5(9):1890–8.
37.   Koevoets T, Beukeboom LW. Genetics of postzygotic isolation and Haldane's rule in
haplodiploids. Heredity. 2008;102(1):16–23.
38.   Presgraves DC. Evaluating genomic signatures of "the large X-effect" during complex
speciation. Mol Ecol. 2018;27(19):3822–30.
39.   Wayne ML, Pan YJ, Nuzhdin S V, McIntyre LM. Additivity and *trans*-acting effects on gene
expression in male *Drosophila simulans*. Genetics. 2004;168(3):1413–20.
40.   Llopart A. The rapid evolution of X-linked male-biased gene expression and the large-X

effect in *Drosophila yakuba*, *D. santomea*, and their hybrids. Mol Biol Evol.
2012;29(12):3873–86.

41.   Meisel RP, Malone JH, Clark AG. Faster-X evolution of gene expression in *Drosophila*.
PLoS Genet. 2012;8(10):e1003013.

42.   Reinke V, Gil IS, Ward S, Kazmer K. Genome-wide germline-enriched and sex-biased
expression profiles in *Caenorhabditis elegans*. Development. 2004;131(2):311–23.

43.   Ortiz MA, Noble D, Sorokin EP, Kimble J. A new dataset of spermatogenic vs. oogenic
transcriptomes in the nematode *Caenorhabditis elegans*. G3. 2014;4(9):1765–72.

44.   Ma X, Zhu Y, Li C, Xue P, Zhao Y, Chen S, et al. Characterisation of *Caenorhabditis elegans*
sperm transcriptome and proteome. BMC Genomics. 2014;15(1):1–13.

45.   Baird SE, Seibert SR. Reproductive isolation in the Elegans-Group of *Caenorhabditis*. Nat
Sci. 2013;05(04):18–25.

46.   Félix M-A, Braendle C, Cutter AD. A streamlined system for species diagnosis in
Caenorhabditis (Nematoda: Rhabditidae) with name designations for 15 distinct
biological species. PLoS One. 2014;9(4):e94723-9.

47.   Thomas CG, Wang W, Jovelin R, Ghosh R, Lomasko T, Trinh Q, et al. Full-genome
evolutionary histories of selfing, splitting, and selection in *Caenorhabditis*. Genome Res.
2015;25(5):667–78.

48.   Woodruff GC, Eke O, Baird SE, Félix M-A, Haag ES. Insights into species divergence and
the evolution of hermaphroditism from fertile interspecies hybrids of *Caenorhabditis*
nematodes. Genetics. 2010;186(3):997–1012.

49.   Kozlowska JL, Ahmad AR, Jahesh E, Cutter AD. Genetic variation for postzygotic
reproductive isolation between *Caenorhabditis briggsae* and *Caenorhabditis* sp. 9.
Evolution. 2011;66(4):1180–95.

50.   Bundus JD, Alaei R, Cutter AD. Gametic selection, developmental trajectories, and
extrinsic heterogeneity in Haldane's rule. Evolution. 2015;69(8):2005–17.

51.   Cutter AD. Reproductive transitions in plants and animals: selfing syndrome, sexual
selection and speciation. New Phytol. 2019;224(3):1080–94.

52.   Palopoli MF, Peden C, Woo C, Akiha K, Ary M, Cruze L, et al. Natural and experimental
evolution of sexual conflict within Caenorhabditis nematodes. Int J Evol Biol. 2015;1–13.

53.   Thomas CG, Li R, Smith HE, Woodruff GC, Oliver B, Haag ES. Simplification and
desexualization of gene expression in self-fertile nematodes. Curr Biol.
2012;22(22):2167–72.

54.   Fierst JL, Willis JH, Thomas CG, Wang W, Reynolds RM, Ahearne TE, et al. Reproductive
mode and the evolution of genome size and structure in *Caenorhabditis* nematodes.
PLoS Genet. 2015;11(6):e1005323-25.

55.   Yin D, Schwarz EM, Thomas CG, Felde RL, Korf IF, Cutter AD, et al. Rapid genome
shrinkage in a self-fertile nematode reveals sperm competition proteins. Science.
2018;359(6371):55–61.

56.   Cutter AD, Morran LT, Phillips PC. Males, outcrossing, and sexual selection in
Caenorhabditis nematodes. Genetics. 2019;213(1):27–57.

57.   Bi Y, Ren X, Yan C, Shao J, Xie D, Zhao Z. A genome-wide hybrid incompatibility landscape
between Caenorhabditis briggsae and C. nigoni. PLoS Genet. 2015;11(2):e1004993-26.

58.   Bi Y, Ren X, Li R, Ding Q, Xie D, Zhao Z. Specific interactions between autosome and X-

chromosomes cause hybrid male sterility in *Caenorhabditis* species. Genetics.
2019;212(3):801–13.

59.   Woodruff GC, Willis JH, Phillips PC. Dramatic evolution of body length due to
postembryonic changes in cell size in a newly discovered close relative of *Caenorhabditis*
elegans. Evol Lett. 2018;2(4):427–41.

60.   Renaut S, Nolte AW, Bernatchez L. Gene expression divergence and hybrid misexpression
between lake whitefish species pairs *Coregonus* spp. Salmonidae). Mol Biol Evol.
2009;26(4):925–36.

61.   Meyer BJ. X-Chromosome dosage compensation. WormBook. 2005;1–14.

62.   Schartner CM. Evolutionary comparison of X-chromosome dosage compensation across
Caenorhabditis species. [University of California, Berkeley]: University of California,
Berkeley; 2016.

63.   Meiklejohn CD, Coolon JD, Hartl DL, Wittkopp PJ. The roles of *cis*- and *trans*-regulation in
the evolution of regulatory incompatibilities and sexually dimorphic gene expression.
Genome Res. 2014;24(1):84–95.

64.   Lemos B, Araripe LO, Fontanillas P, Hartl DL. Dominance and the evolutionary
accumulation of cis- and trans-effects on gene expression. Proc Natl Acad Sci.
2008;105(38):14471–6.

65.   Merritt C, Rasoloson D, Ko D, Seydoux G. 3'UTRs are the primary regulators of gene
expression in the *C. elegans* germline. Curr Biol. 2008;18(19):1476–82.

66.   Gibson G, Weir B. The quantitative genetics of transcription. Trends Genet.
2005;21(11):616–23.

67.   Kasimatis KR, Nelson TC, Phillips PC. Genomic signatures of sexual conflict. J Hered.
2017;108(7):780–90.

68.   Rowe L, Chenoweth SF, Agrawal AF. The genomics of sexual conflict. Am Nat.
2018;192(2):274–86.

69.   Sturgill D, Zhang Y, Parisi M, Oliver B. Demasculinization of X chromosomes in the
Drosophila genus. Nature. 2007;450(7167):238–41.

70.   Kelleher DF, de Carvalho CE, Doty A V, Layton M, Cheng AT, Mathies LD, et al.
Comparative genetics of sex determination: masculinizing mutations in *Caenorhabditis*
briggsae. Genetics. 2008;178(3):1415–29.

71.   Ellis RE. Sex determination: TRA-1 Is a non-binary regulator of sexual identity. Curr Biol.
2020;30(18):R1036--R1038.

72.   Goodwin EB, Biology REEC, 2002. Turning clustering loops: sex determination in
Caenorhabditis elegans. Curr Biol. 2002;12(3):R111--R120.

73.   Verster AJ, Ramani AK, McKay SJ, Fraser AG. Comparative RNAi screens in *C. elegans* and
C. briggsae reveal the impact of developmental system drift on gene function. PLoS
Genet. 2014;10(2):e1004077-15.

74.   Kiontke K, Barrière A, Kolotuev I, Podbilewicz B, Sommer R, Fitch DHA, et al. Trends,
stasis, and drift in the evolution of nematode vulva development. Curr Biol.
2007;17(22):1925–37.

75.   Cutter AD. Divergence times in *Caenorhabditis* and *Drosophila* inferred from direct
estimates of the neutral mutation rate. Mol Biol Evol. 2008;25(4):778–86.

76.   Mallard F, Noble L, Guzella T, Afonso B, Baer CF, Teotónio H. Selection and drift

determine phenotypic stasis despite genetic divergence. 2019;

77.   Barrière A, Ruvinsky I. Pervasive divergence of transcriptional gene regulation in
Caenorhabditis nematodes. PLoS Genet. 2014;10(6):e1004435-15.

78.   Barkoulas M, Vargas Velazquez AM, Peluffo AE, Félix M-A. Evolution of new cis-
regulatory motifs required for cell-specific gene expression in *Caenorhabditis*. PLoS
Genet. 2016;12(9):e1006278-23.

79.   Civetta A. Misregulation of gene expression and sterility in interspecies hybrids: causal
links and alternative hypotheses. J Mol Evol. 2016;82(4):176–82.

80.   Lamont LB, Kimble J. Developmental expression of FOG-1/CPEB protein and its control in
the *Caenorhabditis elegans* hermaphrodite germ line. Dev Dyn. 2007;236(3):871–9.

81.   Rice WR. Sex chromosomes and the evolution of sexual dimorphism. Evolution.
1984;38(4):735–42.

82.   Ranz JM, Namgyal K, Gibson G, Hartl DL. Anomalies in the expression profile of
interspecific hybrids of *Drosophila melanogaster* and *Drosophila simulans*. Genome Res.
2004;14(3):373–9.

83.   Vielle A, Callemeyn-Torre N, Gimond C, Poullet N, Gray JC, Cutter AD, et al. Convergent
evolution of sperm gigantism and the developmental origins of sperm size variability in
Caenorhabditis nematodes. Evolution. 2016;70(11):2485–503.

84.   Cutter AD, Ward S. Sexual and temporal dynamics of molecular evolution in *C. elegans*
development. Mol Biol Evol. 2005;22(1):178–88.

85.   Artieri CG, Haerty W, Gupta BP, Singh RS. Sexual selection and maintenance of sex:
evidence from comparisons of rates of genomic accumulation of mutations and
divergence of sex-related genes in sexual and hermaphroditic species of *Caenorhabditis*.
Mol Biol Evol. 2008;25(5):972–9.

86.   Turelli M, Moyle LC. Asymmetric Postmating Isolation: Darwin's Corollary to Haldane's
Rule. Genetics. 2007;176(2):1059–88.

87.   Lu X, Shapiro JA, Ting CT, LI Y, Li C, Xu J, et al. Genome-wide misexpression of X-linked
versus autosomal genes associated with hybrid male sterility. Genome Res.
2010;20(8):1097–102.

88.   Graze RM, McIntyre LM, Morse AM, Boyd BM, Nuzhdin S V, Wayne ML. What the X has
to do with it: differences in regulatory variability between the sexes in *Drosophila*
simulans. Genome Biol Evol. 2014;6(4):818–29.

89.   Stocks M, Dean R, Rogell B, Friberg U. Sex-specific *trans*-regulatory variation on the
Drosophila melanogaster X chromosome. PLoS Genet. 2015;11(2):e1005015-19.

90.   Castillo-Davis CI, Hartl DL, Achaz G. *cis*-regulatory and protein evolution in orthologous
and duplicate genes. Genome Res. 2004;14(8):1530–6.

91.   Tirosh I, Barkai N. Evolution of gene sequence and gene expression are not correlated in
yeast. Trends Genet. 2008;24(3):109–13.

92.   Cutter AD, Choi JY. Natural selection shapes nucleotide polymorphism across the genome
of the nematode *Caenorhabditis briggsae*. Genome Res. 2010;20(8):1103–11.

93.   Barton NH. The reduction in fixation probability caused by substitutions at linked loci.
Genet Res. 1994;64(3):199–208.

94.   Bolger AM, Lohse M, Usadel B. Trimmomatic: A flexible trimmer for Illumina sequence
data. Bioinformatics. 2014;30(15):2114–20.

95.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast
1233   universal RNA-seq aligner. Bioinformatics. 2012;29(1):15–21.
1234   96.   Frazee AC, Jaffe AE, Langmead B, Leek JT. Polyester: Simulating RNA-seq datasets with
1235   differential transcript expression. Bioinformatics. 2015;31(17):2778–84.
1236   97.   Stevenson KR, Coolon JD, Wittkopp PJ. Sources of bias in measures of allele-specific
1237   expression derived from RNA-seq data aligned to a single reference genome. BMC
1238   Genomics. 2013;14(1):513–36.
1239   98.   Degner JF, Marioni JC, Pai AA, Pickrell JK, Nkadori E, Gilad Y, et al. Effect of read-mapping
1240   biases on detecting allele-specific expression from RNA-sequencing data. Bioinformatics.
1241   2009;25(24):3207–12.
1242   99.   Ren X, Li R, Wei X, Bi Y, Ho VWS, Ding Q, et al. Genomic basis of recombination
1243   suppression in the hybrid between *Caenorhabditis briggsae* and *C. nigoni*. Nucleic Acids
1244   Res. 2018;46(3):1295–307.
1245   100.   Fang G, Bhardwaj N, Robilotto R, Gerstein MB. Getting started in gene orthology and
1246   functional analysis. PLOS Comput Biol. 2010;6(3):e1000703-8.
1247   101.   Emms DM, Kelly S. OrthoFinder: solving fundamental biases in whole genome
1248   comparisons dramatically improves orthogroup inference accuracy. Genome Biol.
1249   2015;16:157.
1250   102.   Stevens L, Félix M-A, Beltran T, Braendle C, Caurcel C, Fausett S, et al. Comparative
1251   genomics of 10 new *Caenorhabditis* species. Evol Lett. 2019;3(2):217–36.
1252   103.   Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for
1253   assigning sequence reads to genomic features. Bioinformatics. 2014;30(7):923–30.
1254   104.   R Core Team. R: A Language and Environment for Statistical Computing. R Foundation for
1255   Statistical Computing: R Foundation for Statistical Computing; 2019.
1256   105.   Anders S, Huber W. Differential expression analysis for sequence count data. Genome
1257   Biol. 2010;11(10):1–12.
1258   106.   Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful
1259   approach to multiple testing. J R Stat Soc Ser B. 1995;57(1):289–300.
1260   107.   Katoh K. MAFFT version 5: improvement in accuracy of multiple sequence alignment.
1261   Nucleic Acids Res. 2005;33(2):511–8.
1262   108.   Wright F. The "effective number of codons" used in a gene. Gene. 1990;87(1):23–9.
1263   109.   Fuglsang A. The "effective number of codons" revisited. Biochem Biophys Res Commun.
1264   2004;317(3):957–64.
1265   110.   Harris RS. Improved pairwise alignment of genomic DNA. [Ph.D. Thesis, Pennsylvania
1266   State University]: Ph.D. Thesis, Pennsylvania State University; 1997.
1267   111.   Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic
1268   features. Bioinformatics. 2010;26(6):841–2.
1269   112.   Hubisz MJ, Pollard KS, Siepel A. PHAST and RPHAST: phylogenetic analysis with
1270   space/time models. Brief Bioinform. 2011;12(1):41–51.
1271   113.   Howe KL, Bolt BJ, Shafie M, Kersey P, Berriman M. WormBase ParaSite – a
1272   comprehensive resource for helminth genomics. Mol Biochem Parasitol. 2017;215:2–10.
1273

Figures

1275
1276

Figure 1. Incidences of differentially expressed genes for 13,636 orthologs between species and sexes. (**A**) Histogram of $\log_2$ expression divergence between species and allele-specific expression (ASE) in hybrids (*C. nigoni*/*C. briggsae*) for female (top row) and male transcriptomes (bottom row). Colored bars indicate significantly differentially expressed genes, while grey bars indicate counts of genes with non-significant expression differences. (**B**) Enrichment of differentially expressed genes between species and alleles for females and males ($\log_2$ odds ratio, i.e. observed/expected). Asterisks mark significant enrichments (positive values) or depletions (negative values) on chromosomes (*P* value < 0.05 and |$\log_2$ odds ratio| > 0.5). On the legend, "no DE" denotes genes that are not significantly differentially expressed. (**C**) Number and percentage of genes that show significant sex-bias is greater in F_1 hybrids than parent species. In *C. briggsae*, male-biased genes also expressed in hermaphrodites (dark green) are shown separately from male-biased only (brown) and from female-biased genes expressed in hermaphrodites (light green, "female") (see Figure 6). (**D**) Enrichment of genes with significant sex-bias and sex-neutrality in parent species and F_1 hybrids for each chromosome. Same legend as D and asterisks denote same statistics as C.

1292

1293
1294

Figure 2. **Sex-specific differences in expression divergence and regulatory controls identify expression inheritance profiles between species**. (**A**) Per-gene biplot of $\log_2$ expression differences between F_1s and each parent species. (**B**) Box- and density plots of expression distance from the origin or centroid of F_1 hybrids for genes within a given expression inheritance category (see Materials and Methods). (**C**) Stacked barplot of gene counts in each expression inheritance group for each chromosome. (**D**) Per-chromosome enrichment ($\log_2$ odds ratio, i.e. observed/expected) of genes in a given expression inheritance group. Asterisks mark significant enrichments (positive values) or depletions (negative values) on chromosomes (*P* value < 0.05 and |$\log_2$ odds ratio| > 0.5). (**E**) Biplot of expression divergence between species (x-axis) versus allele-specific expression (ASE) in hybrids that indicates the magnitude *cis*-acting expression difference between alleles (y-axis). (**F**) Box- and density plots of the magnitude of absolute expression divergence between species for each type of *cis* and *trans* regulatory changes. (**G**) and (**H**) as for C and D, but indicating different types of *cis* and *trans* regulatory-change profiles. Colors indicate different groups of genes with different expression inheritance (see legend for A-D) and *cis* and *trans* regulatory changes (see legend for E-H). Top panels in each of A-H correspond to female transcriptomes, bottom panels to male transcriptomes.

1311

1312

Figure 3. The X-chromosome is enriched for compensatory *cis-trans* changes in males and *trans*-only* regulatory changes in both sexes. (A) Positive values of the log$_2$ odds ratios for females and males denote enrichment for the X-chromosome and negative values indicate enrichment for autosomes. (B) Gene counts for autosomes and the X-chromosome for regulatory divergence categories inferred from expression in males and females. In males, X-linked regulatory divergence was classified following Supplementary Fig. S12 (also see Material and Methods). Briefly, X-linked compensatory *cis-trans* changes in males represent genes that are not differentially expressed between species, but hybrids show over- or underdominance; *cis*-only* represent genes with expression divergence between species and hybrids matching *C. nigoni* expression; and *trans*-only* denote genes with expression divergence with hybrids matching *C. briggsae* expression. The * differentiates these categories from the *cis*-only and *trans*-only categories inferred from ASE in autosomes and XX individuals.

1325

1326
1327 **Figure 4**. **Differences in regulatory controls between chromosomes and between sexes**.
1328 Heatmap of the number of genes in each expression inheritance group (x-axis) for each type of
1329 *cis* and *trans* regulatory changes (y-axis) for each chromosome (I-V, X) and each sex. Excludes
1330 conserved or ambiguous expression and regulation categories (Supplementary Figure S3).
1331

Figure 5. **Male-biased genes show higher expression divergence, molecular evolution, sex-specific regulatory divergence, and X enrichments**. (**A**) Thirteen distinct species-by-sex gene expression gene groups with respective gene counts. Matrix color indicates state with higher relative expression (N=*C. nigoni*, B=*C. briggsae*; M=male, F=female; I=significant interaction; gray cell indicates no difference between treatment groups: C=conserved, N=non-significant interaction). (**B**) Centroid relative expression reaction norm plots for each group indicating expression differences between species and sexes (Supplementary Figure S4). (**C**) Proportion of genes within each species-by-sex gene group differ in the relative representation of expression inheritance categories and (**D**) types of *cis*- and *trans*-regulatory divergence, distinctly for females (left panels) and males (right panels). (**E**) Absolute expression divergence and (**F**) protein sequence divergence (K_a/K_s') differs across species-by-sex expression profiles most strongly for male-biased genes. (**G**) Male-biased genes are not enriched in chromosome arms but are more abundant on the X-chromosome relative to autosomes (except for the BMI group that is rare on X, see Figure 6). * Fisher's exact test, $P < 0.05$ and |log₂ odds ratio > 0.5|.

1350
Figure 6. Male-biased genes that are upregulated in hermaphrodites are enriched for
spermatogenesis genes and have distinct expression and regulatory patterns in hybrids. (**A**)
Biplot of log-fold-change values in male:female contrasts (sex-biased expression) between our
C. briggsae data and data from (53), which includes *C. briggsae* "pseudo-females". Green dots
denote genes that were differentially expressed between hermaphrodites and "pseudo-
females". (**B and C**) Both male-biased genes upregulated in hermaphrodites and orthologs to *C.*
elegans spermatogenesis genes (44) are enriched for the B-M-I group that shows a species-by-
sex interaction (Fig. 5; Supplementary Fig. S4). Divergence in expression and regulation (**D and**
E) is driven largely by *trans*-only regulatory changes (**F**) resulting in abundant *C. nigoni*
dominant expression and underdominant misexpression (**G**).
1361

1362

1363
Figure 7. **The magnitude of *cis*-regulatory divergence shows positive but weak correlations**
with rates of molecular evolution. (**A**) Scatter plot and linear regression of mutation-adjusted
protein evolution rates (K_a/K_s', Spearman's $\rho = 0.17$, $P < 0.0001$; adjusted $R = 0.035$, $m = 0.019$,
$P < 0.0001$) and (**B**) the proportion of non-conserved 5 bp windows within 500 bp upstream of
each gene (1-P_{cons}, Spearman's $\rho = 0.12$, $P > 0.05$; 1-P_{cons}: adjusted $R = 0.01$, $m = 0.017$, $P <$
0.0001) as a function of *cis*-regulatory divergence (allele-specific expression in hybrids, ASE).
Color scale indicates density of points with brighter colors denoting higher density.

Supplementary figures

1372

Supplementary Figure S1. Distinct expression profiles encompass species, sexes, F1 hybrids. (A)
Multi-dimensional scaling plot showing overall expression distance between samples. (B)
magnitude of expression (rlog-transformed counts) for autosomes and the X-chromosome
across *C. briggsae* hermaphrodites, *C. nigoni* females, and F1 females. The X-chromosome has,
on average, lower magnitude of gene expresion than autosomes, which is expected with
dosage compensation. However, F1 hybrids do not show a strong pattern of over-active dosage
compensation leading to generalized lower levels of expression.

1380
1381

1382
1383
1384 Supplementary Figure S2. Centroid relative expression of co-expression clusters and
1385 chromosomal enrichments grouped by sex-biased expression. The naming scheme for each of
1386 the clusters is arbritary. Numbers inside circles indicate the number of genes in each cluster.
1387 Enrichment is represented by the $\log_2$ odds ratio (i.e., observed/expected), with positive values
1388 indicating enrichment and negative values depletion.
1389
1390

1391

1392
1393 Supplementary Figure S3. X-autosome differences in regulatory controls in females and
1394 between sexes for autosomes underlie hybrid transcriptomic profiles relative to parent species.
1395 Heatmap of the number of genes in each expression inheritance group (x-axis) for each type of
1396 *cis* and *trans* regulatory changes (y-axis) for each chromosome (I-V, X) and each sex. Includes all
1397 categorizations.
1398
1399
1400
1401

1402
1403 Supplementary Figure S4. Centroid of normalized expression and 95% CI of species-by-sex
1404 groups (see Fig. 5) plotted as reaction norms. Rows show relative expression of sex-neutral,
1405 male-biased, and female-biased genes. Groups that include an "I" at the end in the name code
1406 have significant species-by-sex interactions.
1407
1408

1409
Supplementary Figure S5. Male-biased, hemaphrodite upregulated genes (see Fig. 6) in males
have similar expression inheritance, (B) but not expression distance (A), to females (Fig. 6). The
X-chromosome is depleted compared to autosomes (B) but has distinct relative enrichments of
genes with *C. briggsae* expression dominance in both males and "females" in contrast to
autosomes, and presents different misexpression categories between males (overdominant)
and females (underdominant).
1416

1417
Supplementary Figure S6. Genes involved with sex determination and germline development
have significant divergence in regulatory networks. The left panel shows relative (log-
transformed and normalized) expression for males (brown) and females (light green) in *C.*
briggsae (Cbr), F$_1$ hybrids (HF1), and *C. nigoni* (Cni). The right panel shows a biplot between the
log-fold change in allele-specific expression (y-axis) and the log-fold change in expression
divergence between species (x-axis). Dotted lines mark expected trajectories for *trans*-only
(horizontal), *cis*-only (diagonal), and *cis-trans* compensatory (vertical) changes. Grey dots
represent the regulatory space of other orthologs in our dataset.
1426
1427

1428
Supplementary Figure S7. Chromosomal arm (green) and center (purple) regions differ strongly
in (A) upstream sequence divergence (1-P_{cons}; proportion of non-conserved 5 bp windows
within 500 bp upstream of each gene) but only moderately in (B) *cis*-regulatory divergence (log$_2$
allele-specific expression, ASE; females on top). (**A**) Proportion of non-conserved 5 bp windows
within 500 bp upstream of each gene (1-P_{cons}) for each chromosome. (**B**) Absolute magnitude of
log2 allele-specific expression or *cis*-regulatory divergence for each chromosome in females and
for autosomes only in males. Black lines in A and B indicate general additive regression (GAM)
trendlines. (**C**) Arm and center regions do not differ greatly in magnitude of regulatory
divergence. Box- and density plots of total (absolute) *cis*-regulatory divergence (left two panels)
and *trans* regulatory divergence (right two panels) for either male or female gene expression.
Allele-specific expression values are square-root-normalized in C to facilitate visual comparison.
1440

1441

1442 Supplementary Figure S8. Coding sequence evolutionary rates for replacement sites (K_a) and

1443 synonymous sites (K_s', adjusted for selection on codon usage) for 13,636 orthologs between *C.*

1444 *briggsae* and *C. nigoni* along the chromosome positions of the *C. briggsae* genome. Colors mark

1445 chromosome arms (green) and center (purple); black lines indicate general additive regression

1446 (GAM) trends.

1447

Supplementary Figure S9. Biplot of *cis* vs *trans* effects for genes with significant sex-bias. *trans* divergence was measured as the $\log_2$ expression difference between parent species (*cis* and *trans*) - the $\log_2$ expression difference between alleles in F1 hybrids (*cis*). Genes with *cis*-only and *trans*-only effects were kept as *cis* and *trans* respectively, whereas genes with significant joint *cis-trans* effects in this analysis were deemed *cis* if abs(cis) > abs(*trans*) and vice versa to be deemed *trans*. Error bars were calculated using non-parametric bootstrapping (n=1000). Female transcriptomes in top panels; males in bottom panels.

1458
1459

1460
1461 Supplementary Figure S10. Proportion of autosomal genes with significant allele-specific
1462 expression (y-axis) showing either additive expression (first panel, top row), simple expression
1463 dominance of one species (second and third panels, top row) and transgressive over- or under-
1464 dominant expression (bottom panels) in hybrid males. Genes with conserved regulation or
1465 *trans*-only effects were excluded for not having significant allele specific expression.
1466

1467

1468
Supplementary Figure S11. Scatter plot and linear regressions (top 4 panels) between
regulatory divergence (log-fold-change) due to allele-specific expression or *cis* changes and
their FDR corrected *P*-values, regulatory divergence due to *trans* effects and their FDR
corrected *P*-values, and volcano plots (lower panels) comparing results from simulated RNA-seq
allele-specific expression data generated by *polyester* and allele-specific read counts by
CompMap.

Supplementary Figure S12. Type 1 (false positive) and type 2 (false negative) error rates for each category of regulatory divergence. Expected values were drawn from RNA-seq allele-specific expression data simulated with *polyester*. Observed values were drawn based on inferences drawn from CompMap allele-specific count data.

1481
1482 Supplementary Fig. S13. Qualitative examples of allele-specific expression with their
1483 classification of gene regulation type changes.
1484

1485
1486 Supplementary Fig. S14. Qualitative examples to diagram regulatory divergence scoring for the
1487 X-chromosome in males that are hemizygous for the X (i.e., X0). Categories not fitting into
1488 either *cis*-only, *trans*-only, or compensatory *cis-trans* were lumped into "other".
1489
1490
1491