Low Intensity Vibration Restores Nuclear YAP Levels and Acute YAP Nuclear Shuttling in

Mesenchymal Stem Cells Subjected to Simulated Microgravity

Thompson M[1], Woods K[2], Newberg, J[1], Oxford JT[2], Uzer G[1†]

[1] Mechanical and Biomedical Engineering, Boise State University

[2] Biomolecular Sciences Graduate Program, Boise State University

**[†] Corresponding Author**

**Running title:** LIV rescues SMG-inhibited YAP nuclear entry

Funding support: NASA ISGC NNX15AI04H, NIH R01AG059923, and 5P2CHD086843-03,

P20GM109095, P20GM103408 and NSF 1929188.

**[†] Corresponding author:**

Gunes Uzer PhD

Boise State University

Department of Mechanical and Biomedical Engineering

1910 University Drive, MSd-2085

Boise, ID 83725-2085

Ph. (208) 426-4461

**Email: gunesuzer@boisestate.edu**

**Abstract**

Reducing the bone deterioration that astronauts experience in microgravity requires countermeasures that can improve the effectiveness of rigorous and time-expensive exercise regimens under microgravity. The ability of low intensity vibrations (LIV) to activate force-responsive signaling pathways in cells suggests LIV as a potential countermeasure to improve cell responsiveness to subsequent mechanical challenge. Mechanoresponse of mesenchymal stem cells (MSC) which maintain bone-making osteoblasts is in part controlled by the "mechanotransducer" protein YAP (Yes-associated protein) which is shuttled into the nucleus in response cyto-mechanical forces. Here, using YAP nuclear shuttling as a measure of MSC mechanoresponse, we tested the effect of 72 hours of simulated microgravity (SMG) and daily LIV application ($LIV_{DT}$) on the YAP nuclear entry driven by either acute LIV ($LIV_{AT}$) or Lysophosphohaditic acid (LPA), applied at the end of the 72h period. We hypothesized that SMG-induced impairment of acute YAP nuclear entry will be alleviated by daily application of $LIV_{DT}$. Results showed that while both acute $LIV_{AT}$ and LPA treatments increased nuclear YAP entry by 50% and 87% over the basal levels in SMG-treated MSCs, nuclear YAP levels of all SMG groups were significantly lower than non-SMG controls. Daily dosing of $LIV_{DT}$, applied in parallel to SMG, restored the SMG-driven decrease in basal nuclear YAP to control levels as well as increased the LPA-induced but not $LIV_{AT}$-induced YAP nuclear entry over the non-$LIV_{DT}$ treated, SMG only, counterparts. These cell level observations suggest that utilizing daily LIV treatments is a feasible countermeasure for increasing the YAP-mediated anabolic responsiveness of MSCs to subsequent mechanical challenge under SMG.

**Key Words:** Simulated Microgravity, Low Intensity Vibrations, YAP, Nucleus, Mesenchymal Stem Cells, Mechanosignaling

**Introduction**

The musculoskeletal deterioration which astronauts experience on long-term space missions and the resulting increase of traumatic physical injury risk is in part due to the reduction of mechanical loading on the musculoskeleton [1]. To alleviate the detrimental effects of unloading, astronauts undergo intensive regimens of running and resistance training in orbit [2]. Despite these efforts, astronauts lose an average bone density of 1% for each month they spend in space [3]. This loss necessitates new non-pharmacologic therapies in addition to exercise to keep bones healthy during long-term space missions. In bone, tissue level response to mechanical challenge is in part regulated by osteoblasts and osteocytes [4]. Both osteoblasts and osteocytes in turn share a common progenitor: the mesenchymal stem cell (MSC). Therefore, the growth and differentiation of MCSs in response to mechanical stimulation is required for the maintenance and repair of bone [5]. It is for this reason that the MSCs are a potential target for mechanical therapies aiming to alleviate bone loss in astronauts, injured service personnel with long periods of bedrest, and physically inactive aged individuals [6].

To maintain healthy bone making cell populations, MSCs rely on environmental mechanical signals inside the bone marrow niches and near bone surfaces. While the exact characteristics of the mechanical environment in which MSCs exist remains to be quantified, it is known that during habitual activities, our bones are subjected to combinations of complex loads including strain, fluid shear, and acceleration, each of which is inseparable [7]. For example, during moderate running, cortical bone can experience strains up to 2000$\mu\epsilon$ [8,9], which also generates coupled fluid flow within canaliculi of up to 100$\mu$m/s [10]. The interior of bone is filled with bone marrow with viscosities in the range of 400-800cP [11]. During moderate running, tibial accelerations are within the 2-5g range [12] (1g = 9.81 m/s$^2$), creating a complex loading at the bone-marrow interface that depends on many factors including frequency, amplitude, and viscosity [13]. *In silico* studies reveal that when exposed to vibrations (0.1-2g), marrow-filled

trabecular compartments generate fluid shear stresses up to 2Pa [13,14], capable of driving bone

cell functions [15]. Interestingly, while these high magnitude forces are only experienced a few

times during the day, bones are bombarded by smaller mechanical signals arising from muscle

contractions that generate bone strains ranging between 2 to 10με [16]. Exogenous application of

small magnitude mechanical regimes in the form of low intensity vibrations (LIV) ranging

between 0.1-2g acceleration magnitudes and 20-200Hz frequencies were shown to be effective

in improving bone and muscle indices in clinical and preclinical studies [17]. At the cellular level,

our group has reported that application of LIV increases MSC contractility [18], activates RhoA

signaling [19], and results in increased osteogenic differentiation and proliferation of MSCs [20].

One of the most actively investigated signaling pathways that regulate the MSC

mechanoresponse is the Yes-associated protein (YAP) signaling pathway. YAP depletion in

stem cells results in reduced proliferation and osteogenesis [21,22]. Similarly, depleting YAP from

osteoblast progenitors decreases both bone quality and quantity in mice [23]. Functionally, in

response to cytomechanical forces and substrate stiffness, YAP moves from the cytoplasm to

the nucleus where it interacts with its co-transcriptional activators such as TEAD to regulate

gene expression related to proliferation [24]. For example, application of substrate strain induces

YAP nuclear entry and YAP transcriptional activity which is required to activate proliferation [25].

While it has been shown that YAP nuclear entry is triggered by soluble factors that increase F-

actin contractility such as Lysophosphatidic acid (LPA) [26,27], large changes in substrate stiffness

[22], or substrate stretch ranging from 3% to 15% [25,28], it is not known if low magnitude signals like

LIV also trigger acute YAP nuclear entry.

Research aimed at studying the effects of microgravity at the cellular level often relies on

simulated microgravity (SMG) devices designed to alter the gravitational conditions that cells

experience by rotating on one or multiple axes at low speed [29-31]. SMG decreases MSC

proliferation [32] and cytoskeletal contractility [29,33,34]. In this way, application of physical or soluble

factors that induce cytoskeletal contractility are commonly used as countermeasures for SMG

[35,36]. SMG also alters nuclear structure. Research from our group has shown that SMG results in

reduced levels of integral nuclear proteins such as Lamin A/C and LINC (Linker of

Nucleoskeleton and Cytoskeleton) complex elements Sun-2 [37]. As mechanically induced

nuclear shuttling of YAP and its paralog TAZ have been associated with LINC complex function

[38], SMG also results in decreased nuclear levels of the YAP paralog protein TAZ [35].

Interestingly, twice daily application of LIV for 20 minutes during SMG recovers both MSC

proliferation and levels of nuclear envelope proteins Lamin A/C and Sun-2, suggesting LIV as a

potential countermeasure to improve YAP-mediated mechanosignaling in MSCs under SMG.

In this study, using YAP nuclear shuttling as a measure of MSC mechanoresponse, we tested

the effect of 72 hours of simulated microgravity (SMG) and daily LIV application ($LIV_{DT}$) on the

YAP nuclear entry driven by either acute LIV ($LIV_{AT}$) or Lysophosphohaditic acid (LPA), applied

at the end of the 72h period. We hypothesized that SMG-induced impairment of YAP nuclear

entry in response to mechanical and soluble factors would be alleviated by daily application of

LIV.

**Results**

*Acute $LIV_{AT}$ application increases nuclear YAP levels*

To quantify the acute YAP nuclear entry in response to LIV, MSCs were plated at density of

5,200 cells/cm$^2$ and were allowed to attach for 24hr. Following this, MSCs were subjected to

treatment in two groups: control and acute LIV treatment regimen ($LIV_{AT}$). The $LIV_{AT}$ regimen

consisted of 5x 20min vibration periods separated by 1hr in between each repetition at room

temperature while control samples were treated identically (also taken out of the incubator) but

were not vibrated. Immediately after $LIV_{AT}$, samples were immunostained for YAP and DAPI.

MATLAB was used to quantify the changes in the nuclear YAP levels. As shown in **Fig.1a**,

confocal images showed increased nuclear YAP following the $LIV_{AT}$ treatment. Analysis of

confocal images to quantify nuclear YAP intensity shown in **Fig.1b** revealed a 32% increase in

the nuclear YAP levels in the $LIV_{AT}$ samples as compared to the control samples ($p<0.0001$).

We also used C2C12 myoblasts to confirm the $LIV_{AT}$ induced YAP nuclear entry on a second

cell line, quantitative analysis of confocal images showed a 40% increase of nuclear YAP in

$LIV_{AT}$ samples compared to controls (**Fig.S1**). As both LIV-induced focal adhesion signaling,

initiated by focal adhesion kinase (FAK) phosphorylation at Tyr 397 residue [19], and YAP nuclear

entry in response to substrate strain [28] requires intact LINC function, disabling LINC function via

a dominant negative overexpression of Nesprin KASH (Klarsicht, ANC-1, Syne homology)

fragment both decreased basal nuclear YAP levels by 34% ($p<0.0001$) and impeded the LIV-

induced YAP nuclear entry when compared to empty plasmid (**Fig.S2**). FAK phosphorylation at

Tyr 397 residue (pFAK) was blocked via a FAK inhibitor (FAKi) PF573228 (3µM) 1hr prior to

$LIV_{AT}$ treatment as previously described [19] and stained against DAPI and YAP (**Fig.S3a**). FAKi

inhibited the $LIV_{AT}$ induced pFAK and decreased its basal levels (**Fig.S3b**). As shown in

**Fig.S3c**, measuring nuclear YAP levels showed that $LIV_{AT}$ induced YAP nuclear entry was not

affected by FAKi when compared DMSO treated controls.

*Basal nuclear YAP levels decreased by SMG were rescued by daily application of $LIV_{DT}$*

We next tested whether SMG decreases basal YAP levels and whether a daily LIV treatment

regimen ($LIV_{DT}$), applied in parallel with SMG, could alleviate decreased YAP in the nucleus. As

we reported previously, $LIV_{DT}$ consisting of 2x 20min vibrations applied every 24 hours during

the 72h period of SMG. This $LIV_{DT}$ regimen was effective at restoring MSC proliferation and

whole cell YAP levels when applied in conjunction with SMG [37]. MSCs were plated at a density

of 1,700 cells/cm$^2$ in 9cm$^2$ tissue culture SlideFlasks (Nunc, #170920) and were allowed to

attach for 24h, after which point, the flasks were filled completely with growth medium, sealed,

and subjected to 72h of treatment followed by immunostaining for YAP and nuclear staining

using DAPI. During the 72h treatment period, MSCs were divided into three groups: control

samples, SMG samples which were subjected to the 72h SMG alone, and SMG+LIV$_{DT}$ samples

which were subjected to both the 72h SMG regimen and the daily LIV$_{DT}$ regimen.

Representative images for YAP and DAPI stained images are shown in **Fig.2a.** As depicted in

**Fig.2b**, quantitative analysis of confocal images revealed a 42% decrease in the nuclear YAP

intensity of the SMG group as compared to non-SMG controls (p<0.0001). Compared to the

SMG group, the LIV$_{DT}$ group increased nuclear YAP levels by 67% (p<0.0001) and there was no

significant difference between nuclear YAP levels of LIV$_{DT}$ treated MSCs and non-SMG controls.

*LIV$_{AT}$-induced YAP nuclear entry decreased by SMG was not restored by daily LIV$_{DT}$ application*

As SMG decreased basal nuclear YAP levels, we next tested whether SMG decreases LIV$_{AT}$-

induced YAP mechanosignaling (i.e. nuclear shuttling). Since LIV$_{DT}$ was able to restore nuclear

YAP levels (**Fig.2**), the SMG+LIV$_{DT}$ group was added to evaluate the effect of LIV$_{DT}$ on the SMG

response. A schematic of the experimental design is given in **Fig.3**. MSCs were divided into six

groups in which the CTRL, SMG, SMG+LIV$_{DT}$ groups were treated with ±LIV$_{AT}$ at the end of 72h

and nuclear YAP levels were measured. As shown in **Fig.4**, SMG alone decreased basal

nuclear YAP levels by 37% (p<0.0001) which were restored back to control levels in the

SMG+LIV$_{DT}$ group. As depicted in **Fig.4**, +LIV$_{AT}$ increased nuclear YAP levels in the CTRL,

SMG and SMG+LIV$_{DT}$ groups by 50%, 69% and 22%, respectively (p<0.0001) while exhibiting

the smallest increase in the SMG+LIV$_{DT}$. As a result, final nuclear YAP levels in the

SMG+LIV$_{DT}$+LIV$_{AT}$ group was not significantly different from the SMG+LIV$_{AT}$ and 23% lower than

the LIV$_{AT}$ group (p<0.0001). Representative confocal images are presented in **Fig.S4**.

*LPA treatment increases nuclear YAP levels*

As SMG+LIV$_{DT}$ treatment did not improve the acute LIV$_{AT}$ response when compared to the SMG

group, we next considered a soluble regulator of cytoskeletal contractility, LPA [20]. To test the

effect of LPA on the acute YAP nuclear entry, two LPA concentrations (50µM and 100µM) were

compared against control samples. Shown in **Fig.5**, nuclear YAP levels were almost doubled

under a two-hour exposure to 50µM LPA and 100µM LPA treatments with 99% and 107%

increases as compared to the control samples (p<0.0001). Nuclear YAP levels for 50µM LPA

and 100µM LPA treatments were not significantly different. Therefore, we chose to use 50µM

LPA treatment in the subsequent experiments.

*LPA-induced YAP nuclear entry decreased by SMG was alleviated by daily LIV$_{DT}$ application*

In order to evaluate whether LIV$_{DT}$ can restore LPA-induced YAP nuclear entry after SMG,

50µM LPA dissolved in DMSO or DMSO as vehicle control were added to the samples at the

end of the 72h treatment of either CTRL, SMG or SMG+LIV$_{DT}$ treatments. The CTRL group,

SMG group, and SMG+LIV$_{DT}$ group were subjected to the same treatment as in the previous

experiments and displayed similar results. As depicted in **Fig.6**, +LPA increased nuclear YAP

levels in the CTRL, SMG and SMG+LIV$_{DT}$ groups by 105%, 67% and 43% respectively

(p<0.0001). While final YAP nuclear levels in SMG+LIV$_{DT}$+LPA remained 70% higher than

SMG+LPA group (P<0.0001), it remained 29% lower than the LPA group (P<0.0001).

*MSC stiffness and structure remain intact under SMG and SMG+LIV$_{DT}$ treatments*

As YAP mechanosignaling of SMG+LIV$_{DT}$ MSCs remained below control levels in response to

both LIV$_{AT}$ and LPA, we quantified the effects of SMG and SMG+LIV$_{DT}$ on the cell stiffness, F-

actin intensity, cell area and nuclear area. AFM testing was used to quantify the elastic modulus

of the nucleus by measuring load-displacement curves on top of the nucleus.  AFM tests shown

in **Fig.7a** indicated a 21% and 27% stiffness decrease in the SMG or SMG+LIV$_{DT}$ groups but

differences were not significant. Quantified from confocal images (**Fig.7b**), mean F-actin

intensities for all the cells in each imaging field were quantified by dividing the mean F-actin

intensity to the number of nuclei in each imaging field. Shown in **Fig.7c**, SMG and SMG+LIVDT

treated MSCs revealed 36% and 30% decreases in the mean F-actin intensity per cell,

respectively but the differences were not statistically significant. We have further quantified

nuclear area as a measure of cyto-mechanical forces on the nucleus [39]. Shown in **Fig.7d**,

analysis of cross-sectional area of cell nuclei using DAPI stained images revealed no significant

effect on average nuclear size by either SMG or combined SMG+LIV$_{DT}$ treatment compared to

control levels.

**Discussion**

The mechanical forces that the bone and muscle cells are subjected to on Earth and in

microgravity are complex and remain incompletely understood. At the same time, it is clear that

these forces are required for healthy tissue growth and function. The complexity of these forces

makes it difficult to design experiments that comprehensively simulate *in vivo* conditions. While

the *in vitro* experiments utilizing SMG and LIV treatments used in this study are limited in this

way and do not entirely correlate with the physiological behavior of these cells *in vivo*, the

experiments presented here remain useful for testing cell behavior under well-defined

conditions.

In this study, we focused on YAP mechanosignaling of MSCs. The first experiments

demonstrated that repeated LIV$_{AT}$ application over six hours was capable of stimulating YAP

entry into the nucleus in both MSCs (**Fig.1**) and in C2C12 cell line (**Fig.S1**). These findings

suggest that similar to high magnitude substrate strains [25] smaller mechanical signals such as

LIV can be effective at increasing nuclear YAP levels. Further, in agreement with earlier reports

utilizing uniaxial strain [28], LIV$_{AT}$-induced increase in nuclear YAP levels also required functional

LINC complexes (**Fig.S2**). Integrin related FAK signaling have been shown to promote YAP

nuclear levels in the proliferative descendants of stem cells and that FAK inhibitor PF573228

decreased nuclear YAP in these cells [40]. Similarly, inhibiting integrin engagement via blocking

FAK phosphorylation in Tyr 397 residue via FAKi also mutes the increase of GTP-bound RhoA

levels in LIV treated MSCs [19]. While we confirmed the loss of phosphorylation in Tyr 397 at both

basal level and in response to $LIV_{AT}$ (**Fig.S3b**), FAKi treatment changed neither basal levels nor

the $LIV_{AT}$-induced increase in nuclear YAP (**Fig.S3c**), suggesting a FAK independent

mechanism. In these experiments, we did not compare $LIV_{AT}$ with strain because the application

of 5 to 15% stretch onto sealed culture flasks was not technically possible without significantly

altering experimental conditions. Instead, LPA addition served as the best option for applying a

simple mechanical stimulation in order to evaluate the YAP mechanotransduction. LPA is a

phospholipid derivative signaling molecule which is capable of causing the simulation of static

transient stretch of a cell by increasing the contractility of the cytoskeleton [20,41]. The first

experiments with LPA served to verify that the simulation of stretch via increased cytoskeleton

contractility was capable of triggering YAP entry into the nucleus and the analysis methods

utilized here were capable of detecting this response (**Fig.5**).

The first SMG experiments confirmed a clear decrease of basal nuclear YAP levels.

Interestingly, SMG treated cells remained responsive, as both $LIV_{AT}$ and LPA treatments were

able to increase the nuclear YAP levels at the end of acute stimulation period (< 6h). However,

final nuclear YAP levels in SMG treated MSCs remained significantly lower when compared to

non-SMG groups (**Fig.2, 4 & 6**). These findings suggested that the YAP mechanosignaling

apparatus of MSCs, to some extent, was intact under SMG. When applied in parallel to SMG,

daily $LIV_{DT}$ treatment was able to restore basal YAP levels in the cell nucleus (**Fig.2, 4 & 6**)

measured 24h after the final $LIV_{DT}$ treatment. This increase in nuclear levels supported our

earlier report that showed sustained recovery of MSC proliferation by $LIV_{DT}$ [37].

Interestingly, this increase in basal nuclear YAP levels under $LIV_{DT}$ was accompanied by a

reduced MSC response to $LIV_{AT}$ treatment (**Fig.4**). When SMG+$LIV_{DT}$ treated MSCs were

subjected to $LIV_{AT}$, the increase in nuclear YAP from the non-$LIV_{AT}$ control was only 22%

(p<0.0001), which was small compared to the 77% increase seen in the SMG groups in

response to $LIV_{AT}$. As a result of this smaller increase in the SMG+$LIV_{DT}$ group, there was no

measurable difference between SMG and SMG+$LIV_{DT}$, samples that were subjected to $LIV_{AT}$. It

has been previously reported that an application of multiple LIV bouts separated by a refractory

period is more effective at activating mechano-signaling pathways such as βcatenin [42]. It is

possible that long term application of $LIV_{DT}$ results in cell structural adaptations that serve to

reduce MSC responsiveness to $LIV_{AT}$ treatment. To test this possibility, we replaced $LIV_{AT}$ with

an LPA treatment. When $LIV_{AT}$ was replaced by LPA treatment (**Fig.6**), responsiveness of

SMG+$LIV_{DT}$ treated MSCs almost doubled to 43% (compared to 22% in response to $LIV_{AT}$) and

was significantly higher than the SMG+LPA group (p<0.0001), suggesting that $LIV_{DT}$ increases

the YAP-mechanosignaling in response to LPA.

Absolute nuclear YAP intensity in the SMG+$LIV_{DT}$+$LIV_{AT}$ group, however, remained below the

$LIV_{AT}$ group (p<0.0001). Previously published findings using the same treatment protocols

suggested that the total cellular YAP levels decreased by SMG were restored to control levels

by daily LIV [37]. This indicates that total availability of YAP protein was not responsible for this

difference between the SMG+$LIV_{DT}$+$LIV_{AT}$ and the $LIV_{AT}$ groups. In regards to other potential

effects of SMG on the components of the mechanosignaling mechanism, one current prevailing

hypothesis suggests a role for nuclear pore opening in response to cyto-mechanical forces [38]

which may be affected by changes in the nuclear stiffness. To test this possibility, we performed

additional AFM and imaging experiments. While the AFM measured nuclear stiffness was 24%

lower in the SMG and SMG+$LIV_{DT}$ groups on average, we were unable to identify any

statistically significant effects of SMG or $LIV_{DT}$ treatment on nuclear stiffness. There was also

slight F-actin intensity decreases in both the SMG and SMG+LIV$_{DT}$ groups which were also not

significant (**Fig.7**). Similarly, cell and nuclear area were not affected. While our results were not

able to detect any changes in nuclear stiffness, considering the significant role that the nuclear

membrane plays as a mechanical structural component in the cell's interpretation of mechanical

stimulus [43,44], more detailed future studies are needed to study the effects of SMG on the

nuclear envelope and nuclear structure.

In summary, while the restoration of basal nuclear levels and improvement of LPA induced YAP

nuclear entry under daily LIV$_{DT}$ treatment identify LIV as a possible countermeasure to improve

MSC response under the detrimental effects of simulated microgravity, future studies are

required to understand why acute YAP nuclear entry in response to mechanical and soluble

factors remain less responsive.

**Methods and Materials**

_Cell Culture_

Primary mouse bone marrow derived MSC's were extracted as previously described [37,45].

C2C12 mouse myoblasts were derived from muscle satellite cells. MSCs were subcultured and

plated in Iscove modified Dulbecco's cell culture medium (IMDM, 12440053, Gibco) with 10%

fetal calf serum (FCS, S11950H, Atlanta Biologicals) and 1% pen/strep. C2C12s were

subcultured and plated in Dulbecco's modified Eagle's medium (DMEM, DML09, Caisson

Laboratories) with 10% fetal calf serum (FCS, S11950H, Atlanta Biologicals) and 1% pen/strep.

MSCs were subcultured in 9cm$^2$ culture dishes at a density of 5,200cells/cm$^2$ for 1day

experiments and 1700cells/cm$^2$ for 3day experiments, while C2C12s were plated at a density of

10,000cells/cm$^2$. Experimental cells were plated and given 24h to attach to the mounting surface

prior to experiments. Cell passages for both MSCs and C2C12s used for experiments were

limited to P7-P15.

<u>*Low Intensity Vibrations Treatment*</u>

SlideFlasks with plated MSCs were filled completely with culture medium and placed in LIV

device designed and used in previous research (**Fig.3**) [37]. LIV device subjected cells to low

intensity 90 Hz lateral vibrations at 0.7g at room temperature. MSCs were vibrated for 20min

intervals separated over time. $LIV_{AT}$ regimen was applied after 72h treatment period and

consisted of 5x 20min LIV with an hour in between each. Daily $LIV_{DT}$ regimen consisted of 3x

treatments in parallel with SMG treatment each consisting of 2x 20min LIV with 2h in between.

<u>*Simulated Microgravity Treatment*</u>

SlideFlasks (Nunc, #170920) with plated MSCs were filled completely with culture medium

(**Fig.3**) and placed in a clinostat SMG device. The clinostat shown is a redesign of a custom-

made clinostat described in previous research [37] with a new flask holder casing capable of

holding SlideFlasks and is also autoclavable. The clinostat was used to subject the MSCs to

constant 15 RPM SMG for 72h.

<u>*Immunofluorescence Staining and Image Analysis*</u>

Immediately after mechanical treatment, MSCs plated in Slideflasks were removed from

treatment, and the SlideFlasks were disassembled in order to stain the MSCs on the slides

(**Fig.3**). The MSCs were fixed with 4% paraformaldehyde, then washed and permeabilized with

0.05% Triton X-100 in PBS, followed by immunostaining with YAP specific antibody (YAP

(D8H1X) Rabbit mAb, Cell Signaling Technologies) and Alexa Fluor red secondary antibodies

(Donkey anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor Plus 594 for

all experiments prior to usage of LPA. Subsequently, Donkey anti-Rabbit IgG (H+L) Cross-

Adsorbed Secondary Antibody, Alexa Fluor Plus 633 was used). Nuclear DNA was labeled via

DAPI (Vectashield Mounting Medium, Vector Laboratories). Stained samples were imaged with

a Leica TCS SP8 confocal microscope (40x, HC PL APO CS2 Oil Immersion) prior to usage of

LPA, after this Zeiss LSM 510 Meta Confocal Microscope (40x, HC PL APO CS2 Oil Immersion)

. Exported images were used to quantify relative YAP levels within each nuclei (nuclear regions

traced by DAPI stained nucleus) via custom-made MATLAB program (The MathWorks, Natick,

MA). DAPI images were analyzed using an edge-detection algorithm in order to determine the

nuclear area for each cell. The nuclear outline was then used as a mask to quantify the average

pixel intensity of the YAP stain within the nuclei of each individual cell. (n=50-100

nuclei/sample).

*Atomic Force Microscopy*

Bruker Dimension FastScan AFM was used for collection of the atomic force measurements.

Tipless MLCT-D probes with a 0.03 N/m spring constant were functionalized with 10 µm

diameter borosilicate glass beads for force collection. The AFM's optical microscope was used

to locate individual live MSCs plated on the SlideFlask slides with the flask section removed for

access to the cells. The nucleus of each cell was tested with at least 3 seconds of rest between

each test. In each test, three force-displacement curves were obtained (ramping rate: 2 µm/sec

over 2 µm total travel, 1 µm approach, 1 µm retract), which were analyzed using Nanoscope

software with the implementation of a best-fit curve to a Hertzian (spherical) model (optimized

such that $R^2$ value was greater than 0.95, or $p<0.05$) to obtain elastic moduli of nuclear

membrane of individual nuclei.

*Western Blotting*

Western blotting was performed as previously described.[23,26,27,64]  20µg of lysed cell protein from

each sample was run on a 10% polyacrylamide gels, transferred onto a polyvinylidene difluoride

(PVDF) membranes, blocked with 5%( w/v) milk for 1h. After washing, primary antibodies were

incubated overnight at 4˚C with.  Protein bands were visualized via horseradish peroxidase-

conjugated secondary antibodies (1: 5,000, Cell Signaling) and ECL plus chemiluminescence kit

(Amersham Biosciences, Piscataway, NJ) and scanned using C-DiGit blot scanner (Licor,

Lincoln, NE).  All blots derive from the same experiment and were processed in parallel.

*Statistical analysis*

All data analysis results were displayed graphically based on the mean value with standard

error bars. Differences between treatments were not assumed to follow a Gaussian distribution.

Therefore, group differences were identified via either non-parametric two-tailed Mann-Whitney

U-test (**Fig.1a**) or Kruskal-Wallis test followed by Tukey multiple comparison (**Fig.2b, 4, 5, 6, 7,**

**&S1, S2, S3**). P-values of less than 0.05 were considered significant.

*Data availability*

The datasets generated and/or analyzed during the current study are available from the

corresponding author on reasonable request.

*Ethics*

All methods were carried out in accordance with relevant guidelines and regulations of Boise

Institutional Animal Care and Use Committee and Institutional Biosafety Committee. All

procedures were approved by Boise State University Institutional Animal Care and Use

Committee, and Institutional Biosafety Committee.

*Acknowledgements*

This study was supported by NASA ISGC NNX15AI04H, NIH R01AG059923, and

5P2CHD086843-03, P20GM109095, P20GM103408 and NSF 1929188.

*Competing interests*

The author(s) declare no competing interests, financial or otherwise.

*Contributions*

**Thompson, M** experimental methods, data analysis/interpretation, manuscript writing, final

approval of manuscript.

**Woods, K** data analysis/interpretation, final approval of manuscript.

**Newberg, J** experimental methods, final approval of manuscript

**Oxford JT**, financial support, final approval of manuscript

**Uzer, G** concept/design, financial support, data analysis/interpretation, manuscript writing, final

approval of manuscript

*References*

1    Smith, S. M. *et al.* Calcium metabolism before, during, and after a 3-mo spaceflight:

kinetic and biochemical changes. *Am J Physiol* **277**, R1-10 (1999).

2    Greenleaf, J. E., Bulbulian, R., Bernauer, E. M., Haskell, W. L. & Moore, T. Exercise-

training protocols for astronauts in microgravity. *Journal of applied physiology*

*(Bethesda, Md. : 1985)* **67**, 2191-2204, doi:10.1152/jappl.1989.67.6.2191 (1989).

3    Vico, L. *et al.* Effects of long-term microgravity exposure on cancellous and cortical

weight-bearing bones of cosmonauts. *Lancet (London, England)* **355**, 1607-1611 (2000).

4    Thompson, W. R., Rubin, C. T. & Rubin, J. Mechanical regulation of signaling pathways

in bone. *Gene* **503**, 179-193, doi:10.1016/j.gene.2012.04.076 (2012).

5    Ozcivici, E. *et al.* Mechanical signals as anabolic agents in bone. *Nature reviews.*

*Rheumatology* **6**, 50-59, doi:10.1038/nrrheum.2009.239 (2010).

6    Rando, T. A. & Ambrosio, F. Regenerative Rehabilitation: Applied Biophysics Meets

Stem Cell Therapeutics. *Cell Stem Cell* **22**, 306-309, doi:10.1016/j.stem.2018.02.003

(2018).

7      Chan, M. E., Uzer, G. & Rubin, C. The Potential Benefits and Inherent Risks of Vibration

as a Non-Drug Therapy for the Prevention and Treatment of Osteoporosis. *Current*

*osteoporosis reports*, 1-9, doi:10.1007/s11914-012-0132-1 (2013).

8      Judex, S., Gross, T. S. & Zernicke, R. F. Strain Gradients Correlate with Sites of

Exercise-Induced Bone-Forming Surfaces in the Adult Skeleton. *Journal of Bone and*

*Mineral Research* **12**, 1737-1745, doi:10.1359/jbmr.1997.12.10.1737 (1997).

9      Rubin, C. T. & Lanyon, L. E. Dynamic strain similarity in vertebrates; an alternative to

allometric limb bone scaling. *Journal of Theoretical Biology* **107**, 321-327,

doi:10.1016/s0022-5193(84)80031-4 (1984).

10     Price, C., Zhou, X. Z., Li, W. & Wang, L. Y. Real-Time Measurement of Solute Transport

Within the Lacunar-Canalicular System of Mechanically Loaded Bone: Direct Evidence

for Load-Induced Fluid Flow. *Journal of Bone and Mineral Research* **26**, 277-285,

doi:10.1002/jbmr.211 (2011).

11     Gurkan, U. A. & Akkus, O. The Mechanical Environment of Bone Marrow: A Review.

*Annals of biomedical engineering* **36**, 1978-1991, doi:10.1007/s10439-008-9577-x

(2008).

12     Vainionpaa, A. *et al.* Intensity of exercise is associated with bone density change in

premenopausal women. *Osteoporosis international : a journal established as result of*

*cooperation between the European Foundation for Osteoporosis and the National*

*Osteoporosis Foundation of the USA* **17**, 455-463, doi:10.1007/s00198-005-0005-x

(2006).

13     Dickerson, D. A., Sander, E. A. & Nauman, E. A. Modeling the mechanical

consequences of vibratory loading in the vertebral body: microscale effects.

*Biomechanics and Modeling in Mechanobiology* **7**, 191-202, doi:10.1007/s10237-007-

0085-y (2008).

14    Coughlin, T. R. & Niebur, G. L. Fluid shear stress in trabecular bone marrow due to low-

magnitude high-frequency vibration. *Journal of biomechanics* **45**, 2222-2229,

doi:10.1016/j.jbiomech.2012.06.020 (2012).

15    Riddle, R. C. & Donahue, H. J. From Streaming Potentials to Shear Stress: 25 Years of

Bone Cell Mechanotransduction. *Journal of Orthopaedic Research* **27**, 143-149,

doi:10.1002/jor.20723 (2009).

16    Fritton, S. P., McLeod, K. J. & Rubin, C. T. Quantifying the strain history of bone: spatial

uniformity and self-similarity of low-magnitude strains. *Journal of biomechanics* **33**, 317-

325, doi:10.1016/s0021-9290(99)00210-9 (2000).

17    Pagnotti, G. M. *et al.* Combating osteoporosis and obesity with exercise: leveraging cell

mechanosensitivity. *Nature Reviews Endocrinology*, doi:10.1038/s41574-019-0170-1

(2019).

18    Pongkitwitoon, S., Uzer, G., Rubin, J. & Judex, S. Cytoskeletal Configuration Modulates

Mechanically Induced Changes in Mesenchymal Stem Cell Osteogenesis, Morphology,

and Stiffness. *Scientific reports* **6**, 34791, doi:10.1038/srep34791 (2016).

19    Uzer, G. *et al.* Cell Mechanosensitivity to Extremely Low-Magnitude Signals Is Enabled

by a LINCed Nucleus. *STEM CELLS* **33**, 2063-2076, doi:10.1002/stem.2004 (2015).

20    Uzer, G., Pongkitwitoon, S., Ete Chan, M. & Judex, S. Vibration induced osteogenic

commitment of mesenchymal stem cells is enhanced by cytoskeletal remodeling but not

fluid shear. *Journal of Biomechanics* **46**, 2296-2302, doi:10.1016/j.jbiomech.2013.06.008

(2013).

21    Pan, J. X. *et al.* YAP promotes osteogenesis and suppresses adipogenic differentiation

by regulating beta-catenin signaling. *Bone research* **6**, 18, doi:10.1038/s41413-018-

0018-7 (2018).

22    Dupont, S. *et al.* Role of YAP/TAZ in mechanotransduction. *Nature* **474**, 179-183,

466            doi:http://www.nature.com/nature/journal/v474/n7350/abs/10.1038-nature10137-

467            unlocked.html#supplementary-information (2011).

468    23    Kegelman, C. D. *et al.* Skeletal cell YAP and TAZ combinatorially promote bone

469            development. *FASEB journal : official publication of the Federation of American*

470            *Societies for Experimental Biology* **32**, 2706-2721, doi:10.1096/fj.201700872R (2018).

471    24    Zhao, B. *et al.* TEAD mediates YAP-dependent gene induction and growth control.

472            *Genes & development* **22**, 1962-1971, doi:10.1101/gad.1664408 (2008).

473    25    Benham-Pyle, B. W., Pruitt, B. L. & Nelson, W. J. Cell adhesion. Mechanical strain

474            induces E-cadherin-dependent Yap1 and beta-catenin activation to drive cell cycle entry.

475            *Science (New York, N.Y.)* **348**, 1024-1027, doi:10.1126/science.aaa4559 (2015).

476    26    Ho, L. T. Y., Skiba, N., Ullmer, C. & Rao, P. V. Lysophosphatidic Acid Induces ECM

477            Production via Activation of the Mechanosensitive YAP/TAZ Transcriptional Pathway in

478            Trabecular Meshwork Cells. *Investigative Ophthalmology & Visual Science* **59**, 1969-

479            1984, doi:10.1167/iovs.17-23702 (2018).

480    27    Cai, H. & Xu, Y. The role of LPA and YAP signaling in long-term migration of human

481            ovarian cancer cells. *Cell Communication and Signaling* **11**, 31, doi:10.1186/1478-811X-

482            11-31 (2013).

483    28    Driscoll, Tristan P., Cosgrove, Brian D., Heo, S.-J., Shurden, Zach E. & Mauck,

484            Robert L. Cytoskeletal to Nuclear Strain Transfer Regulates YAP Signaling in

485            Mesenchymal Stem Cells. *Biophysical journal* **108**, 2783-2793,

486            doi:10.1016/j.bpj.2015.05.010 (2015).

487    29    Janmaleki, M., Pachenari, M., Seyedpour, S. M., Shahghadami, R. & Sanati-Nezhad, A.

488            Impact of Simulated Microgravity on Cytoskeleton and Viscoelastic Properties of

489            Endothelial Cell. *Scientific reports* **6**, 32418, doi:10.1038/srep32418 (2016).

30   Qian, A. R. *et al.* Fractal Dimension as a Measure of Altered Actin Cytoskeleton in

MC3T3-E1 Cells Under Simulated Microgravity Using 3-D/2-D Clinostats. *Biomedical*

*Engineering, IEEE Transactions on* **59**, 1374-1380, doi:10.1109/TBME.2012.2187785

(2012).

31   Pardo, S. J. *et al.* Simulated microgravity using the Random Positioning Machine inhibits

differentiation and alters gene expression profiles of 2T3 preosteoblasts. *Am J Physiol*

*Cell Physiol* **288**, doi:10.1152/ajpcell.00222.2004 (2005).

32   Dai, Z. Q., Wang, R., Ling, S. K., Wan, Y. M. & Li, Y. H. Simulated microgravity inhibits

the proliferation and osteogenesis of rat bone marrow mesenchymal stem cells. *Cell*

*proliferation* **40**, 671-684, doi:10.1111/j.1365-2184.2007.00461.x (2007).

33   Shi, F. *et al.* Simulated Microgravity Promotes Angiogenesis through RhoA-Dependent

Rearrangement of the Actin Cytoskeleton. *Cellular physiology and biochemistry :*

*international journal of experimental cellular physiology, biochemistry, and pharmacology*

**41**, 227-238, doi:10.1159/000456060 (2017).

34   Corydon, T. J. *et al.* Reduced Expression of Cytoskeletal and Extracellular Matrix Genes

in Human Adult Retinal Pigment Epithelium Cells Exposed to Simulated Microgravity.

*Cellular physiology and biochemistry : international journal of experimental cellular*

*physiology, biochemistry, and pharmacology* **40**, 1-17, doi:10.1159/000452520 (2016).

35   Chen, Z., Luo, Q., Lin, C., Kuang, D. & Song, G. Simulated microgravity inhibits

osteogenic differentiation of mesenchymal stem cells via depolymerizing F-actin to

impede TAZ nuclear translocation. *Scientific reports* **6**, 30322, doi:10.1038/srep30322

(2016).

36   Uddin, S. M. & Qin, Y. X. Enhancement of osteogenic differentiation and proliferation in

human mesenchymal stem cells by a modified low intensity ultrasound stimulation under

simulated microgravity. *PLoS One* **8**, e73914, doi:10.1371/journal.pone.0073914 (2013).

37    Touchstone, H. *et al.* Recovery of stem cell proliferation by low intensity vibration under

simulated microgravity requires intact LINC complex  *npj. Microgravity* **5**,

doi:doi.org/10.1038/s41526-019-0072-5 (2019).

38    Shiu, J.-Y., Aires, L., Lin, Z. & Vogel, V. Nanopillar force measurements reveal actin-

cap-mediated YAP mechanotransduction. *Nature cell biology* **20**, 262-271,

doi:10.1038/s41556-017-0030-y (2018).

39    Saeed, M. & Weihs, D. Finite element analysis reveals an important role for cell

morphology in response to mechanical compression. *Biomechanics and Modeling in*

*Mechanobiology*, doi:10.1007/s10237-019-01276-5 (2019).

40    Hu, J. K.-H. *et al.* An FAK-YAP-mTOR Signaling Axis Regulates Stem Cell-Based

Tissue Renewal in Mice. *Cell Stem Cell* **21**, 91-106.e106,

doi:10.1016/j.stem.2017.03.023 (2017).

41    Riddick, N., Ohtani, K. & Surks, H. K. Targeting by myosin phosphatase-RhoA

interacting protein mediates RhoA/ROCK regulation of myosin phosphatase. *Journal of*

*Cellular Biochemistry* **103**, 1158-1170, doi:10.1002/jcb.21488 (2008).

42    Sen, B. *et al.* Mechanical signal influence on mesenchymal stem cell fate is enhanced by

incorporation of refractory periods into the loading regimen. *Journal of biomechanics* **44**,

593-599, doi:10.1016/j.jbiomech.2010.11.022 (2011).

43    De Magistris, P. & Antonin, W. The Dynamic Nature of the Nuclear Envelope. *Current*

*biology : CB* **28**, R487-r497, doi:10.1016/j.cub.2018.01.073 (2018).

44    Czapiewski, R., Robson, M. I. & Schirmer, E. C. Anchoring a Leviathan: How the

Nuclear Membrane Tethers the Genome. *Frontiers in genetics* **7**, 82,

doi:10.3389/fgene.2016.00082 (2016).

45    Uzer, G. *et al.* Sun-mediated mechanical LINC between nucleus and cytoskeleton

regulates betacatenin nuclear access. *J Biomech* **74**, 32-40,

doi:10.1016/j.jbiomech.2018.04.013 (2018).

**Figure 1. Acute $LIV_{AT}$ application increases nuclear YAP levels.** (a) MSCs were subjected to $LIV_{AT}$ and stained with DAPI (blue) and YAP (red). Confocal images indicated an increased nuclear YAP levels following acute $LIV_{AT}$ applied as five 20min vibration periods separated by 1hr. (b) Quantitative analysis of confocal images showed a 32% of increase of nuclear YAP in $LIV_{AT}$ samples compared to controls. n>400/grp, group comparison was made a Mann-Whitney U-test, $*p<0.05$, $**p<0.01$, $***p<0.01$, $****p<0.0001$.

**Figure 2. Basal nuclear YAP levels decreased by SMG were rescued by $LIV_{DT}$.** (a) MSCs were subjected to SMG, and SMG+$LIV_{DT}$ over 72h period and stained with DAPI (blue) and YAP (red). (b) Quantitative analysis showed a 42% decrease of nuclear YAP levels in the SMG group compared to control levels. The SMG+$LIV_{DT}$ group showed a 67% increase of nuclear YAP when compared to the SMG group. There was no statistically significant difference between CTRL and SMG+$LIV_{DT}$ groups. n>100/grp. Group comparisons were made via Kruskal-Wallis test followed by Tukey multiple comparison, $*p<0.05$, $**p<0.01$, $***p<0.01$, $****p<0.0001$.

**Figure 3. Experimental design of combined SMG, $LIV_{AT}$ and $LIV_{DT}$ application.** MSCs were subcultured and plated in SlideFlasks and allowed to attach for 24h before SlideFlasks were filled with culture medium and sealed. Treatment regimen for MSC's involved 72h of SMG (blue). $LIV_{DT}$ regimen consisted of one treatment cycle every 24hr during SMG treatment with each cycle consisting of 2x 20min LIV with an hour in between (yellow). $LIV_{AT}$ regimen was applied after 72h SMG treatment period and consisted of 5x 20min LIV with an hour in between each (red). For LIV application, MSCs plated in SlideFlasks were placed in LIV device constructed in the lab previous to this research. Vibrations were applied at peak magnitudes of 0.7 g at 90 Hz at room temperature. Control samples were treated the same but were not vibrated. For SMG application, MSCs plated in SlideFlasks were secured in lab custom-built

clinostat inside incubator. The clinostat subjected the MSCs to constant 15 RPM rotation

simulated microgravity. After treatment, flasks were removed for immunofluorescence staining.

**Figure 4. LIV$_{AT}$-induced YAP nuclear entry decreased by SMG was not restored by daily**

**LIV$_{DT}$ application.** MSCs were subjected to either CTRL, SMG, SMG+LIV$_{DT}$ over 72h period

were subsequently treated with LIV$_{AT}$. Quantitative analysis of confocal images showed that

SMG alone decreased basal nuclear YAP levels by 37% which were increased back to control

levels in the SMG+LIV$_{DT}$ group. +LIV$_{AT}$ increased nuclear YAP levels in the CTRL, SMG and

SMG+LIV$_{DT}$ groups by 50%, 69% and 22%, respectively. Nuclear YAP intensity in the

SMG+LIV$_{DT}$+LIV$_{AT}$ group remained not significantly different from the SMG+LIV$_{AT}$ and 23%

smaller than the LIV$_{AT}$ group. n>200/grp. Group comparisons were made via Kruskal-Wallis test

followed by Tukey multiple comparison, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001.

**Figure 5. LPA treatment increases nuclear YAP levels.** (a) Representative confocal images

of DAPI (blue) and YAP (red) stained MSCs with or without LPA treatment. MSCs were

subjected to LPA addition at 50μM and 100μM concentrations. (b) Quantitative analysis of

confocal images revealed a 99% and a 107% increase in the 50μM LPA and 100μM LPA

treatments compared to DMSO treated controls, respectively. Nuclear YAP levels for 50μM LPA

and 100μM LPA treatments were not significantly different.  n>30/grp. Group comparisons were

made via Kruskal-Wallis test followed by Tukey multiple comparison, *p<0.05, **p<0.01,

***p<0.01, ****p<0.0001.

**Figure 6. LPA-induced YAP nuclear entry decreased by SMG was alleviated by daily LIV$_{DT}$**

**application.** MSCs were subjected to SMG, and parallel SMG+LIV$_{DT}$ over 72h period at the end

of 72h, samples were treated with either LPA (50μM) or DMSO. Quantitative analysis of

confocal images revealed that LPA addition increased nuclear YAP levels by 105%, 67% and

43% in the CTRL, SMG and SMG+LIV$_{DT}$ when compared to DMSO controls. When compared to

nuclear YAP intensity of the LPA treatment alone, SMG+LPA and SMG+LIV$_{DT}$+LPA samples

were 55% and 29% lower, respectively. YAP nuclear levels in SMG+LIV$_{DT}$+LPA remained 70%

higher than SMG+LPA group. n>100/grp. Group comparisons were made via Kruskal-Wallis test

followed by Tukey multiple comparison, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001.

**Figure 7. MSC stiffness and structure remain intact under SMG and SMG+LIV$_{DT}$**

**treatments.** MSCs were subjected to SMG and parallel SMG+LIV$_{DT}$ over a 72h period. (a)

Compared to CTRL samples, AFM measurement of the elastic moduli of SMG and SMG+LIV$_{DT}$

treated MSCs revealed apparent decreases in elastic modules that were 21% and 27% below

control levels, measured differences were not statistically significant. n=10/grp. (b)

Quantification of confocal images show that, (c) mean F-actin intensity of SMG and SMG+LIV$_{DT}$

treated MSCs revealed decrease of 36% and 30% below control levels, measured differences

were not statistically significant. n=15/grp. (d) No significant effects of either SMG or LIV$_{DT}$

treatment on the average nucleus size were found. n>100/grp. Group comparisons were made

via Kruskal-Wallis test followed by Tukey multiple comparison, *p<0.05, **p<0.01, ***p<0.01,

****p<0.0001.

**Figure S1. LIV$_{AT}$ treatment increases nuclear YAP in C2C12 cells.** (a) C2C12 cells were

subjected to LIV$_{AT}$ and stained with DAPI (blue) and YAP (red). Confocal images displayed

increased nuclear YAP levels following LIV$_{AT}$ treatment. (b) Quantitative analysis of confocal

images showed a 40% increase of nuclear YAP in LIV$_{AT}$ samples compared to controls.

n>900/grp, group comparison was made using a Mann-Whitney U-test, *p<0.05, **p<0.01,

***p<0.01, ****p<0.0001.

**Figure S2. LINC complex disruption decreases nuclear YAP levels and reduces LIV$_{AT}$-induced YAP nuclear entry.** (a) Plasmids harboring either a dominant negative KASH domain of Nesprin (DNK) to disable LINC complex function or empty mCherry control (MC) were overexpressed in MSCs. Following puromycin selection, MC or DNK expressing MSCs were subjected to LIV$_{AT}$ and stained against DAPI (blue) and YAP (red). (b) Quantitative analysis of confocal images revealed a 49% increase of nuclear YAP following LIV$_{AT}$ in MC expressing control MSCs. Basal YAP levels of the DNK-CTRL group were 34% lower compared to MC-CTRL and LIV$_{AT}$ treatment failed to significantly increase nuclear YAP over DNK-CTRL (18%, NS). n>30/grp, group comparisons were made via Kruskal-Wallis test followed by Tukey multiple comparison, *$p < 0.05$, **$p < 0.01$, ***$p < 0.01$, ****$p < 0.0001$.

**Figure S3. Blocking FAK phosphorylation at Tyr 397 does not limit LIV$_{AT}$ induced YAP nuclear entry.** Dimethyl sulphoxide (DMSO) or Tyr 397 specific FAK inhibitor (FAKi) PF573228 (3µM) was added to MSCs in culture medium for 1h prior to LIV$_{AT}$ or control treatments. (a) Confocal images of YAP showed more intense nuclear YAP staining of LIV$_{AT}$ treated MSCs but no apparent effect of FAKi when compared to DMSO (b) FAKi application 1hr prior to LIV$_{AT}$ treatment inhibited the LIV$_{AT}$ induced FAK phosphorylation at Tyr 397 and decreased the basal levels (c) Quantitative analysis of confocal images revealed a 61% increase of nuclear YAP in both the DMSO-LIV group and a 60% increase in the FAKi-LIV$_{AT}$ group compared to the DMSO-CTRL group. Differences between the DMSO-CTRL and FAKi-CTRL groups and between the DMSO-LIV and FAKi-LIV$_{AT}$ groups were not significant. n>50/grp. Group comparisons were made via Kruskal-Wallis test followed by Tukey multiple comparison, *$p < 0.05$, **$p < 0.01$, ***$p < 0.01$, ****$p < 0.0001$.

**Figure S4.** Confocal images for SMG/LIV$_{DT}$/LIV$_{AT}$ treatments in Figure 4.

644 **Figure S5.** Confocal images for SMG/LIV$_{DT}$/LPA treatment in Figure 6.

646 **Figure S6.** Unprocessed blots used in Figure S3 as obtained by LiCor C-DiGit blot scanner.

648 **Table S1:** Cell culture and pharmacological reagents and their final concentrations.

| Cell Culture and Pharmacological Reagents | | Final Concentration |
|---|---|---|
| IMDM | GIBCO | - |
| DMEM | Caisson Laboratories | - |
| FCS | Atlanta Biologicals | 10% v/v |
| Penicillin/streptomycin | GIBCO | 1% v/v |

650 **Table S2:** Antibodies used and their final concentrations for western blots.

| Antibodies | | Final Concentration |
|---|---|---|
| p-FAK Tyr397 (3283) | Cell Signaling | 1/1000 |
| FAK (sc-558) | Santa Cruz Biotechnology | 1/500 |
| LDHA (2012S) | Cell Signaling Technology | 1/1000 |

652 **Table S3:** Immunostaining antibodies and reagents and their final concentrations.

| Immunostaining antibodies and Reagents | | Final Concentration |
|---|---|---|
| DAPI (H-1500-10) | Vector Laboratories | 1 µg/mL |
| Alexa Fluor 488 Phalloidin | Life Technologies | 0.1µM |
| YAP (14074S) | Cell Signaling Technology | 1/100 |

# Figure 1.

**Figure 1. Acute LIV_AT application increases nuclear YAP levels.** (a) MSCs were subjected to LIV_AT and stained with DAPI (blue) and YAP (red). Confocal images indicated an increased nuclear YAP levels following acute LIV_AT applied as five 20min vibration periods separated by 1hr. (b) Quantitative analysis of confocal images showed a 32% of increase of nuclear YAP in LIV_AT samples compared to controls. n>400/grp, group comparison was made a Mann-Whitney U-test, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001

**Figure 2.**

**Figure 2. Basal nuclear YAP levels decreased by SMG were rescued by LIV_{DT}.**
(a) MSCs were subjected to SMG, and SMG+LIV_{DT} over 72h period and stained
with DAPI (blue) and YAP (red). (b) Quantitative analysis showed a 42% decrease
of nuclear YAP levels in the SMG group compared to control levels. The
SMG+LIV_{DT} group showed a 67% increase of nuclear YAP when compared to the
SMG group. There was no statistically significant difference between CTRL and
SMG+LIV_{DT} groups. n>100/grp. Group comparisons were made via Kruskal-Wallis
test followed by Tukey multiple comparison, *p<0.05, **p<0.01, ***p<0.01,
****p<0.0001.

# Figure 3.

**Figure 3. Experimental design of combined SMG, LIV$_{AT}$ and LIV$_{DT}$ application.**
MSCs were subcultured and plated in SlideFlasks and allowed to attach for 24h
before SlideFlasks were filled with culture medium and sealed. Treatment regimen
for MSC's involved 72h of SMG (blue). LIV$_{DT}$ regimen consisted of one treatment
cycle every 24hr during SMG treatment with each cycle consisting of 2x 20min LIV
with an hour in between (yellow). LIV$_{AT}$ regimen was applied after 72h SMG
treatment period and consisted of 5x 20min LIV with an hour in between each
(red). For LIV application, MSCs plated in SlideFlasks were placed in LIV device
constructed in the lab previous to this research. Vibrations were applied at peak
magnitudes of 0.7 g at 90 Hz at room temperature. Control samples were treated
the same but were not vibrated. For SMG application, MSCs plated in SlideFlasks
were secured in lab custom-built clinostat inside incubator. The clinostat subjected
the MSCs to constant 15 RPM rotation simulated microgravity. After treatment,
flasks were removed for immunofluorescence staining.

**Figure 4.**

**Figure 4. LIV$_{AT}$-induced YAP nuclear entry decreased by SMG was not restored by daily LIV$_{DT}$ application.** MSCs were subjected to either CTRL, SMG, SMG+LIV$_{DT}$ over 72h period were subsequently treated with LIV$_{AT}$. Quantitative analysis of confocal images showed that SMG alone decreased basal nuclear YAP levels by 37% which were increased back to control levels in the SMG+LIV$_{DT}$ group. +LIV$_{AT}$ increased nuclear YAP levels in the CTRL, SMG and SMG+LIV$_{DT}$ groups by 50%, 69% and 22%, respectively. Nuclear YAP intensity in the SMG+LIV$_{DT}$+LIV$_{AT}$ group remained not significantly different from the SMG+LIV$_{AT}$ and 23% smaller than the LIV$_{AT}$ group. n>200/grp. Group comparisons were made via Kruskal-Wallis test followed by Tukey multiple comparison, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001.

# Figure 5.

**Figure 5. LPA treatment increases nuclear YAP levels.** (a) Representative confocal images of DAPI (blue) and YAP (red) stained MSCs with or without LPA treatment. MSCs were subjected to LPA addition at 50µM and 100µM concentrations. (b) Quantitative analysis of confocal images revealed a 99% and a 107% increase in the 50µM LPA and 100µM LPA treatments compared to DMSO treated controls, respectively. Nuclear YAP levels for 50µM LPA and 100µM LPA treatments were not significantly different.  n>30/grp. Group comparisons were made via Kruskal-Wallis test followed by Tukey multiple comparison, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001.

**Figure 6.**

**Figure 6. LPA-induced YAP nuclear entry decreased by SMG was alleviated by daily LIV$_{DT}$ application.** MSCs were subjected to SMG, and parallel SMG+LIV$_{DT}$ over 72h period at the end of 72h, samples were treated with either LPA (50µM) or DMSO. Quantitative analysis of confocal images revealed that LPA addition increased nuclear YAP levels by 105%, 67% and 43% in the CTRL, SMG and SMG+LIV$_{DT}$ when compared to DMSO controls. When compared to nuclear YAP intensity of the LPA treatment alone, SMG+LPA and SMG+LIV$_{DT}$+LPA samples were 55% and 29% lower, respectively. YAP nuclear levels in SMG+LIV$_{DT}$+LPA remained 70% higher than SMG+LPA group. n>100/grp. Group comparisons were made via Kruskal-Wallis test followed by Tukey multiple comparison, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001.

# Figure 7.

**Figure 7. MSC stiffness and structure remain intact under SMG and SMG+LIV$_{DT}$ treatments.** MSCs were subjected to SMG and parallel SMG+LIV$_{DT}$ over a 72h period. (a) Compared to CTRL samples, AFM measurement of the elastic moduli of SMG and SMG+LIV$_{DT}$ treated MSCs revealed apparent decreases in elastic modules that were 21% and 27% below control levels, measured differences were not statistically significant. n=10/grp. (b) Quantification of confocal images show that, (c) mean F-actin intensity of SMG and SMG+LIV$_{DT}$ treated MSCs revealed decrease of 36% and 30% below control levels, measured differences were not statistically significant. n=15/grp. (d) No significant effects of either SMG or LIV$_{DT}$ treatment on the average nucleus size were found. n>100/grp. Group comparisons were made via Kruskal-Wallis test followed by Tukey multiple comparison, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001.

# Supplementary Information

## Low Intensity Vibrations Restore Nuclear YAP Levels and Acute YAP Nuclear Shuttling in Mesenchymal Stem Cells Subjected to Simulated Microgravity

Thompson M[1], Woods K[2], Newberg, J[1], Oxford JT, Uzer G[1†]

[1]Mechanical and Biomedical Engineering, Boise State University
[2]Biomolecular Sciences Graduate Program, Boise State University
**[†] Corresponding Author**

**Funding support:**
NASA ISGC NNX15AI04H, NIH R01AG059923, and 5P2CHD086843-03, P20GM109095, P20GM103408 and NSF 1929188.

**Figure S1. LIV$_{AT}$ treatment increases nuclear YAP in C2C12 cells.** (a) C2C12 cells were subjected to LIV$_{AT}$ and stained with DAPI (blue) and YAP (red). Confocal images displayed increased nuclear YAP levels following LIV$_{AT}$ treatment. (b) Quantitative analysis of confocal images showed a 40% increase of nuclear YAP in LIV$_{AT}$ samples compared to controls. n>900/grp, group comparison was made using a Mann-Whitney U-test, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001.

# Figure S2.

**Figure S2. LINC complex disruption decreases nuclear YAP levels and reduces LIV$_{AT}$-induced YAP nuclear entry.** (a) Plasmids harboring either a dominant negative KASH domain of Nesprin (DNK) to disable LINC complex function or empty mCherry control (MC) were overexpressed in MSCs. Following puromycin selection, MC or DNK expressing MSCs were subjected to LIV$_{AT}$ and stained against DAPI (blue) and YAP (red). (b) Quantitative analysis of confocal images revealed a 49% increase of nuclear YAP following LIV$_{AT}$ in MC expressing control MSCs. Basal YAP levels of the DNK-CTRL group were 34% lower compared to MC-CTRL and LIV$_{AT}$ treatment failed to significantly increase nuclear YAP over DNK-CTRL (18%, NS). n>30/grp, group comparisons were made via Kruskal-Wallis test followed by Tukey multiple comparison, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001.

# Figure S3.

**Figure S3. Blocking FAK phosphorylation at Tyr 397 does not limit LIV$_{AT}$ induced YAP nuclear entry.** Dimethyl sulphoxide (DMSO) or Tyr 397 specific FAK inhibitor (FAKi) PF573228 (3µM) was added to MSCs in culture medium for 1h prior to LIV$_{AT}$ or control treatments. (a) Confocal images of YAP showed more intense nuclear YAP staining of LIV$_{AT}$ treated MSCs but no apparent effect of FAKi when compared to DMSO (b) FAKi application 1hr prior to LIV$_{AT}$ treatment inhibited the LIV$_{AT}$ induced FAK phosphorylation at Tyr 397 and decreased the basal levels (c) Quantitative analysis of confocal images revealed a 61% increase of nuclear YAP in both the DMSO-LIV group and a 60% increase in the FAKi-LIV$_{AT}$ group compared to the DMSO-CTRL group. Differences between the DMSO-CTRL and FAKi-CTRL groups and between the DMSO-LIV and FAKi-LIV$_{AT}$ groups were not significant. n>50/grp. Group comparisons were made via Kruskal-Wallis test followed by Tukey multiple comparison, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001.

# Figure S4.

**Figure S4.** Confocal images for SMG/LIV$_{DT}$/LIV$_{AT}$ treatments in Figure 4.

**Figure S5.**

**Figure S5.** Confocal images for SMG/LIV$_{DT}$/LPA treatment in Figure 6.

**Table S1:** Cell culture and pharmacological reagents and their final concentrations.

| Cell culture and pharmacological reagents | | Final Concentration |
|---|---|---|
| IMDM | GIBCO | - |
| DMEM | Caisson Laboratories | - |
| FCS | Atlanta Biologicals | 10% v/v |
| Penicillin/streptomycin | GIBCO | 1% v/v |

**Table S2:** Antibodies used and their final concentrations for western blots.

| Antibodies | | Final Concentration |
|---|---|---|
| p-FAK Tyr397 (3283) | Cell Signaling Technology | 1/1000 |
| FAK (sc-558) | Santa Cruz Biotechnology | 1/500 |
| LDHA (2012S) | Cell Signaling Technology | 1/1000 |

**Table S3:** Immunostaining antibodies and reagents and their final concentrations.

| Immunostaining antibodies and reagents | | Final Concentration |
|---|---|---|
| DAPI (H-1500-10) | Vector Laboratories | 1 µg/mL |
| Alexa Fluor 488 Phalloidin | Life Technologies | 0.1µM |
| YAP (14074S) | Cell Signaling Technology | 1/100 |

**Figure S6**

**Figure S6.** Unprocessed blots used in Figure S3 as obtained by LiCor C-DiGit blot scanner.