Low Intensity Vibration Restores Nuclear YAP Levels and Acute YAP Nuclear Shuttling in

Mesenchymal Stem Cells Subjected to Simulated Microgravity

Thompson M[1], Woods K[2], Newberg, J[1], Oxford JT[2], Uzer G[1†]

[1] Mechanical and Biomedical Engineering, Boise State University

[2] Biomolecular Sciences Graduate Program, Boise State University

† Corresponding Author

Running title: LIV rescues SMG-inhibited YAP nuclear entry

Funding support: NASA ISGC NNX15AI04H, NIH R01AG059923, and 5P2CHD086843-03,

P20GM109095, P20GM103408 and NSF 1929188.

† Corresponding author:

Gunes Uzer PhD

Boise State University

Department of Mechanical and Biomedical Engineering

1910 University Drive, MSd-2085

Boise, ID 83725-2085

Ph. (208) 426-4461

Email: gunesuzer@boisestate.edu

Abstract

Reducing the bone deterioration that astronauts experience in microgravity requires

countermeasures that can improve the effectiveness of rigorous and time-expensive exercise

regimens under microgravity. The ability of low intensity vibrations (LIV) to activate force-

responsive signaling pathways in cells suggests LIV as a potential countermeasure to improve

cell responsiveness to subsequent mechanical challenge. Mechanoresponse of mesenchymal

stem cells (MSC) that maintains bone-making osteoblasts is in part controlled by the

"mechanotransducer" protein YAP (Yes-associated protein) which is shuttled into the nucleus in

response cyto-mechanical forces. Here, using YAP nuclear shuttling as a measure of MSC

mechanoresponse, we tested the effect of 72 hours of simulated microgravity (SMG) and daily

LIV application (LIV_{DT}) on the YAP nuclear entry driven by either acute LIV (LIV_{AT}) or

Lysophosphohaditic acid (LPA), applied at the end of the 72h period. We hypothesized that

SMG-induced impairment of acute YAP nuclear entry will be alleviated by daily application of

LIV_{DT}. Results showed that while both acute LIV_{AT} and LPA treatments increased nuclear YAP

entry by 50% and 87% over the basal levels in SMG-treated MSCs, nuclear YAP levels of all

SMG groups were significantly lower than non-SMG controls. Daily dosing of LIV_{DT}, applied in

parallel to SMG, restored the SMG-driven decrease in basal nuclear YAP to control levels as

well as increasing the LPA-induced but not LIV_{AT}-induced YAP nuclear entry over the non-LIV_{DT}

treated, SMG only, counterparts. These cell level observations suggest that utilizing daily LIV_{DT}

treatments is a feasible countermeasure for increasing the YAP-mediated anabolic

responsiveness of MSCs to subsequent mechanical challenge under SMG.

Key Words: Simulated Microgravity, Low Intensity Vibrations, YAP, Nucleus, Mesenchymal

Stem Cells, Mechanosignaling

Introduction

The musculoskeletal deterioration which astronauts experience on long-term space missions and the resulting increase of traumatic physical injury risk is in part due to the reduction of mechanical loading on the musculoskeleton [1]. To alleviate the detrimental effects of unloading, astronauts undergo long, intensive regimens of running and resistance training in orbit [2]. Despite these efforts, astronauts lose an average bone density of 1% for each month they spend in space [3]. This loss necessitates new non-pharmacologic therapies in addition to exercise to keep bones healthy during long-term space missions. In bone, tissue level response to mechanical challenge is in part regulated by osteoblasts and osteocytes [4]. Both osteoblasts and osteocytes in turn share a common progenitor: the mesenchymal stem cell (MSC). Therefore, the growth and differentiation of MCSs in response to mechanical stimulation is required for the maintenance and repair of bone [5]. It is for this reason that the MSCs are a potential target for mechanical therapies aiming to alleviate bone loss in astronauts, injured service personnel with long periods of bedrest, and physically inactive aged individuals [6].

To maintain healthy bone making cell populations, MSCs rely on environmental mechanical signals inside the bone marrow niches and near bone surfaces. While the exact characteristics of the mechanical environment in which MSCs exist remains to be quantified, it is known that during habitual activities, our bones are subjected to combinations of complex loads including strain, fluid shear, and acceleration, each of which is inseparable [7]. For example, during moderate running, cortical bone can experience strains up to $2000\mu\varepsilon$ [8,9], which also generates coupled fluid flow within canaliculi of up to $100\mu m/s$ [10]. The interior of bone is filled with bone marrow with viscosities in the range of 400-800cP [11]. During moderate running, tibial accelerations are within the 2-5g range [12] ($1g = 9.81$ m/s^2), creating a complex loading at the bone-marrow interface that depends on many factors including frequency, amplitude, and viscosity [13]. *In silico* studies reveal that when exposed to vibrations (0.1-2g), marrow-filled

trabecular compartments generate fluid shear stresses up to 2Pa [13,14], capable of driving bone

cell functions [15]. Interestingly, while these high magnitude forces are only experienced a few

times during the day, bones are bombarded by smaller mechanical signals arising from muscle

contractions that generate bone strains ranging between 2 to 10με [16]. Exogenous application of

small magnitude mechanical regimes in the form of low intensity vibrations (LIV) ranging

between 0.1-2g acceleration magnitudes and 20-200Hz frequencies were shown to be effective

in improving bone and muscle indices in clinical and preclinical studies [17]. At the cellular level,

our group has reported that application of LIV increases MSC contractility [18], activates RhoA

signaling [19], and results in increased osteogenic differentiation and proliferation of MSCs [20].

One of the most actively investigated signaling pathways that regulate the MSC

mechanoresponse is the Yes-associated protein (YAP) signaling pathway. YAP depletion in

stem cells results in reduced proliferation and osteogenesis [21,22]. Similarly, depleting YAP from

osteoblast progenitors decreases both bone quality and quantity in mice [23]. Functionally, in

response to cytomechanical forces and substrate stiffness, YAP moves from the cytoplasm to

the nucleus where it interacts with its co-transcriptional activators such as TEAD to regulate

gene expression related to proliferation [24]. For example, application of substrate strain induces

YAP nuclear entry and YAP transcriptional activity which is required to activate proliferation [25].

While it has been shown that YAP nuclear entry is triggered by soluble factors that increase F-

actin contractility such as Lysophosphatidic acid (LPA)[26,27], large changes in substrate stiffness

[22], or substrate stretch ranging from 3% to 15% [25,28], it is not known if low magnitude signals like

LIV also trigger acute YAP nuclear entry.

Research aimed at studying the effects of microgravity at the cellular level often relies on

simulated microgravity (SMG) devices designed to alter the gravitational conditions that cells

experience by rotating on one or multiple axes at low speed [29-31]. SMG decreases MSC

proliferation [32] and cytoskeletal contractility [29,33,34]. In this way, application of physical or soluble

factors that induce cytoskeletal contractility are commonly used as countermeasures for SMG

[35,36]. SMG also alters nuclear structure. Research from our group has shown that SMG results in

reduced levels of integral nuclear proteins such as Lamin A/C and LINC (Linker of

Nucleoskeleton and Cytoskeleton) complex elements Sun-2 [37]. As mechanically induced

nuclear shuttling of YAP and its paralog TAZ have been associated with LINC complex function

[38], SMG also results in decreased nuclear levels of the YAP paralog protein TAZ [35].

Interestingly, twice daily application of LIV for 20 minutes during SMG recovers both MSC

proliferation and levels of nuclear envelope proteins LaminA/C and Sun-2, suggesting LIV as a

potential countermeasure to improve YAP-mediated mechanosignaling in MSCs under SMG.

In this study, using YAP nuclear shuttling as a measure of MSC mechanoresponse, we tested

the effect of 72 hours of simulated microgravity (SMG) and daily LIV application (LIV_{DT}) on the

YAP nuclear entry driven by either acute LIV (LIV_{AT}) or Lysophosphohaditic acid (LPA), applied

at the end of the 72h period. We hypothesized that SMG-induced impairment of YAP nuclear

entry in response to mechanical and soluble factors will be alleviated by daily application of LIV.

Results

Acute LIV_{AT} application increases nuclear YAP levels

To quantify the acute YAP nuclear entry in response to LIV, MSCs were plated at density of

1,700 cells/cm^2 and were allowed to attach for 24hr. Following this, MSCs were subjected to

treatment in two groups: control and acute LIV treatment regimen (LIV_{AT}). The LIV_{AT} regimen

consisted of 5x 20min vibration periods separated by 1hr in between each repetition at room

temperature while control samples were treated identically (also taken out of the incubator) but

were not vibrated. Immediately after LIV_{AT}, samples were immunostained for YAP and DAPI

was used to visualize the nucleus. MATLAB was used to quantify the changes in the nuclear

YAP levels. As shown in **Fig.1a**, confocal images showed an increase in YAP staining in the cell

nuclei. Analysis of confocal images to quantify nuclear YAP intensity shown in **Fig.1b** revealed

a 32% increase in the nuclear YAP levels in the LIV_{AT} samples as compared to the control

samples (p<0.0001). As both LIV-induced focal adhesion signaling [19] and YAP nuclear entry in

response to substrate strain [28] requires intact LINC function, disabling LINC function via a

dominant negative overexpression of Nesprin KASH (Klarsicht, ANC-1, Syne homology)

fragment both decreased basal nuclear YAP levels by 34% (p<0.0001) and impeded the LIV-

induced YAP nuclear entry when compared to empty plasmid (**Fig.S1**).

Basal nuclear YAP levels decreased by SMG were rescued by daily application of LIV_{DT}

We next tested whether SMG decreases basal YAP levels and whether a daily LIV treatment

regimen (LIV_{DT}), applied in parallel with SMG, could alleviate decreased YAP in the nucleus. As

we reported previously, LIV_{DT} consisting of 2x 20min vibrations applied every 24 hours during

the 72h period of SMG application was effective at restoring MSC proliferation and whole cell

YAP levels, when applied in conjunction with SMG [37]. MSCs were plated at a density of 1,700

cells/cm^2 in 25 cm^2 tissue culture SlideFlasks (Nunc, #170920) and were allowed to attach for

24h, after which point, the flasks were filled completely with growth medium, sealed, and

subjected to 72h of treatment followed by immunostaining for YAP and nuclear staining using

DAPI. During the 72h treatment period, MSCs were divided into three groups: control samples,

SMG samples which were subjected to the standard regimen of 72h of continuous SMG, and

SMG+LIV_{DT} samples which were subjected to both the 72h SMG regimen and the daily LIV_{DT}

regimen. Representative images for YAP and DAPI stained images are shown in **Fig.2a.** As

depicted in **Fig.2b**, quantitative analysis of confocal images revealed a 42% decrease in the

nuclear YAP intensity in the SMG-treated samples as compared to the control samples

(p<0.0001). Compared to the SMG group, nuclear YAP levels of the LIV_{DT} treated MSCs

showed an increase of 67%, respectively. There was no significant nuclear YAP level difference

between of LIV_{DT} treated MSCs and non-SMG controls.

LIV_{AT}-induced YAP nuclear entry decreased by SMG was not restored by daily LIV_{DT} application

As SMG decreased basal nuclear YAP levels, we next tested whether SMG decreases LIV_{AT}-

induced YAP mechanosignaling (i.e. nuclear shuttling). Since LIV_{DT} was able to restore nuclear

YAP levels (**Fig.2**), the SMG+LIV_{DT} group was added to evaluate the effect of LIV_{DT} on the SMG

response. A schematic of the experimental design is given in **Fig.3**. MSCs were divided into six

groups in which the CTRL, SMG, SMG+LIV_{DT} groups were treated with ±LIV_{AT} at the end of 72h

and nuclear YAP levels were measured As shown in **Fig.4**, SMG alone decreased basal nuclear

YAP levels by 37% ($p<0.0001$) which were restored back to control levels in the SMG+LIV_{DT}

group. As depicted in **Fig.4**, +LIV_{AT} increased nuclear YAP levels in the CTRL, SMG and

SMG+LIV_{DT} groups by 50%, 69% and 22%, respectively ($p<0.0001$) while exhibiting the smallest

increase in the SMG+LIV_{DT}. As a result, final nuclear YAP levels in the SMG+LIV_{DT}+LIV_{AT} group

was not significantly different from the SMG+LIV_{AT} and 23% lower than the LIV_{AT} group

($p<0.0001$).

LPA treatment increases nuclear YAP levels

As SMG+LIV_{DT} treatment did not improve the acute LIV_{AT} response when compared to the SMG

group, we next considered a soluble regulator of cytoskeletal tension, LPA. To test the effect of

LPA on the acute YAP nuclear entry, two LPA concentrations (50µM and 100µM) were

compared against control samples. Shown in **Fig.5**, nuclear YAP levels were almost doubled

under a two-hour exposure to 50µM LPA and 100µM LPA treatments with 99% and 107%

increases as compared to the control samples ($p<0.0001$). Nuclear YAP levels for 50µM LPA

and 100µM LPA treatments were not significantly different. Therefore, we chose to use 50µM

LPA treatment in the subsequent experiments.

LPA-induced YAP nuclear entry decreased by SMG was alleviated by daily LIV$_{DT}$ application

In order to evaluate whether LIV$_{DT}$ can restore LPA-induced YAP mechanosignaling after SMG,

50µM LPA dissolved in DMSO or DMSO as vehicle control were added to the samples at the

end of the 72h treatment of either CTRL, SMG or SMG+LIV$_{DT}$ treatments. The CTRL group,

SMG group, and SMG+LIV$_{DT}$ group were subjected to the same treatment as in the previous

experiments and displayed similar results. As depicted in **Fig.6**, +LPA increased nuclear YAP

levels in the CTRL, SMG and SMG+LIV$_{DT}$ groups by 105%, 67% and 43% respectively

(p<0.0001). While final YAP nuclear levels in SMG+LIV$_{DT}$+LPA remained 70% higher than

SMG+LPA group (P<0.0001), it remained 29% lower than the LPA group (P<0.0001).

Nuclear stiffness and area were not affected by SMG and SMG+LIV$_{DT}$ treatments.

As YAP mechanosignaling of SMG+LIV$_{DT}$ MSCs remained below control levels in response to

both LIV$_{AT}$ and LPA, we quantified the nuclear stiffness in CTRL, SMG and SMG+LIV$_{DT}$

samples. AFM testing was used to quantify the elastic modulus of the nucleus by measuring

load-displacement curves on top of the nucleus. As shown in **Fig.7a**, analysis revealed no

statistically significant effect on nuclear elastic modulus in the SMG or SMG+LIV$_{DT}$ groups

compared to control levels. We have further quantified nuclear area as a measure of cyto-

mechanical forces on the nucleus [39]. Shown in **Fig.7b**, analysis of cross-sectional area of cell

nuclei using DAPI stained images revealed no significant effect on average nuclear size by

either SMG or combined SMG+LIV$_{DT}$ treatment compared to control levels.

Discussion

The mechanical forces to which the bone and muscle tissue cells of the human body are

subjected both on Earth and in microgravity environments are complex and remain incompletely

understood. At the same time, it is clear that these forces are required for healthy tissue growth

on Earth. The complexity of these forces makes it difficult to design realistic experiments that

comprehensively simulate *in vivo* conditions. While the *in vitro* experiments utilizing SMG and

LIV treatments used in this study are limited in this way and do not entirely correlate with the

physiological behavior of these cells *in vivo*, the experiments presented here remain useful for

testing cell behavior under well-defined conditions.

In this study, we focused on YAP mechanosignaling in MSCs. The first experiments

demonstrated that repeated LIV_{AT} application over six hours was capable of stimulating YAP

entry into the nucleus (**Fig.1**). These findings suggest that similar to high magnitude substrate

strains [25] smaller mechanical signals such as LIV can be effective at increasing nuclear YAP

levels. Further, in agreement with earlier reports utilizing uniaxial strain [28], LIV_{AT}-induced

increase in nuclear YAP levels also required functional LINC complexes (**Fig.S1**). In these

experiments, we did not compare LIV_{AT} with strain because the application of 5 to 15% stretch

onto sealed culture flasks was not technically possible without significantly altering experimental

conditions. Instead, LPA addition served as the best option for applying a simple mechanical

stimulation in order to evaluate the YAP mechanotransduction. LPA is a phospholipid derivative

signaling molecule which is capable of causing the simulation of static transient stretch of a cell

by increasing the contractility of the cytoskeleton [20,40]. The first experiments with LPA served to

verify that the simulation of stretch via increased cytoskeleton contractility was capable of

triggering YAP entry into the nucleus and the analysis methods utilized here were capable of

detecting this response (**Fig.5**).

The first SMG experiments confirmed a clear decrease in basal nuclear YAP levels.

Interestingly, SMG treated cells remained responsive, as both LIV_{AT} and LPA treatments were

able to increase the nuclear YAP levels at the end of acute stimulation period (< 6h). However,

final nuclear YAP levels in SMG treated MSCs remained significantly lower when compared to

non-SMG groups (**Fig.2, 4 & 6**). These findings suggested that the YAP mechanosignaling

apparatus of MSCs, to some extent, was intact under SMG. When applied in parallel to SMG,

daily LIV_{DT} treatment was able to restore basal YAP levels in the cell nucleus (**Fig.2**) measured

24h after the final LIV_{DT} treatment. This increase in nuclear levels supported our earlier report

that showed sustained recovery of MSC proliferation by LIV_{DT} [37].

Interestingly, this increase in basal nuclear YAP levels under LIV_{DT} was accompanied by a

reduced MSC response to LIV_{AT} treatment (**Fig.5**). When SMG+LIV_{DT} treated MSCs were

subjected to LIV_{AT}, the increase in nuclear YAP from the non-LIV_{AT} control was only 22%

($p<0.0001$), which was small compared to the 77% increase seen in the SMG groups in

response to LIV_{AT}. As a result of this smaller increase in the SMG+LIV_{DT} group, there was no

measurable difference between SMG and SMG+LIV_{DT}, samples that were subjected to LIV_{AT}. It

has been previously reported that an application of multiple LIV bouts separated by a refractory

period is more effective at activating mechano-signaling pathways such as βcatenin [41]. It is

possible that long term application of LIV_{DT} results in cell structural adaptations that serve to

reduce MSC responsiveness to LIV_{AT} treatment. To test this possibility, we have replaced LIV_{AT}

with an LPA treatment. When LIV_{AT} was replaced by LPA treatment (**Fig.6**), responsiveness of

SMG+LIV_{DT} treated MSCs almost doubled to 43% (compared to 22% in response to LIV_{AT}) and

was significantly higher than the SMG+LPA group ($p<0.0001$), suggesting that LIV_{DT} increases

the YAP-mechanosignaling in response to LPA.

Absolute nuclear YAP intensity in the SMG+LIV_{DT}+LIV_{AT} group however, remained below the

LIV_{AT} group ($p<0.0001$). Previously published findings using the same treatment protocols

suggested that the total cellular YAP levels decreased by SMG were restored to control levels

by daily LIV [37]. This would indicate that total availability of YAP protein was not responsible for

this difference between the SMG+LIV_{DT}+LIV_{AT} and the LIV_{AT} groups. In regards to other

potential effects of SMG on the components of the mechanosignaling mechanism, one current

prevailing hypothesis suggests a role for nuclear pore opening in response to cyto-mechanical

forces [38] which may be affected by changes in the nuclear stiffness. To test this possibility, we

have performed additional AFM and imaging experiments. While the AFM measured nuclear

stiffness was 24% lower in SMG and SMG+LIV$_{DT}$ groups on average, we were unable to identify

any statistically significant effects of SMG or LIV$_{DT}$ treatment on nuclear stiffness or on nuclear

area (**Fig.7**). While our results were not able to detect any changes in nuclear stiffness,

considering the significant role that the nuclear membrane plays as a mechanical structural

component in the cell's interpretation of mechanical stimulus [42,43], more detailed future studies

are needed to study the effects of sMG on the nuclear envelope and nuclear structure.

In summary, while the restoration of basal nuclear levels and improvement of LPA induced YAP

nuclear entry under daily LIV$_{DT}$ treatment identify LIV as a possible countermeasure to improve

MSC response under the detrimental effects of simulated microgravity, future studies are

required to understand why acute YAP nuclear entry in response to mechanical and soluble

factors remain less responsive.

Methods and Materials

Mesenchymal Stem Cell Culture

Primary mouse bone marrow derived MSC's were extracted as previously described[37,44]. MSCs

were then subcultured and plated in Iscove modified Dulbecco's cell culture medium (IMDM,

12440053, Gibco) with 10% fetal calf serum (FCS, S11950H, Atlanta Biologicals) and 1%

pen/strep. MSCs were subcultured in 55cm^2 culture dishes at a density of 1,300cells/cm^2.

Experimental cells were plated in SlideFlasks at density of 1,700cells/cm^2 and given 24h to

attach to the slide prior to experiments. Cell passages for MSCs used for experiments were

limited to P7-P15.

<u>*Low Intensity Vibrations Treatment*</u>

SlideFlasks with plated MSCs were filled completely with culture medium and placed in LIV

device designed and used in previous research (**Fig.3**) (Touchstone 2019). LIV device

subjected cells to low intensity 90 Hz lateral vibrations at 0.7g at room temperature. MSCs were

vibrated for 20min intervals separated over time. LIV_{AT} regimen was applied after 72h treatment

period and consisted of 5x 20min LIV with an hour in between each. Daily LIV_{DT} regimen

consisted of 3x treatments in parallel with SMG treatment each consisting of 2x 20min LIV with

2h in between.

<u>*Simulated Microgravity Treatment*</u>

SlideFlasks (Nunc, #170920) with plated MSCs were filled completely with culture medium

(**Fig.3**) and placed in a clinostat SMG device. The clinostat shown is a redesign of a custom-

made clinostat described in previous research (Touchstone, 2019) with a new flask holder

casing capable of holding SlideFlasks and is also autoclavable. The clinostat was used to

subject the MSCs to constant 15 RPM SMG for 72h.

<u>*Immunofluorescence Staining and Image Analysis*</u>

Immediately after mechanical treatment, MSCs plated in Slideflasks were removed from

treatment, and the SlideFlasks were disassembled in order to stain the MSCs on the slides

(**Fig.3**). The MSCs were fixed with 4% paraformaldehyde, then washed and permeabilized with

0.05% Triton X-100 in PBS, followed by immunostaining with YAP specific antibody (YAP

(D8H1X) Rabbit mAb, Cell Signaling Technologies) and Alexa Fluor red secondary antibodies

(Donkey anti-Rabbit IgG (H+L) Cross-Adsorbed Secondary Antibody, Alexa Fluor Plus 594 for

all experiments prior to usage of LPA. Subsequently, Donkey anti-Rabbit IgG (H+L) Cross-

Adsorbed Secondary Antibody, Alexa Fluor Plus 633 was used). Nuclear DNA was labeled via

DAPI (Vectashield Mounting Medium, Vector Laboratories). Stained samples were imaged with

a Leica TCS SP8 confocal microscope (40x, HC PL APO CS2 Oil Immersion) prior to usage of

LPA, after this Zeiss LSM 510 Meta Confocal Microscope (40x, HC PL APO CS2 Oil

Immersion) . Exported images were used to quantify relative YAP levels within each nuclei

(nuclear regions traced by DAPI stained nucleus) via custom-made MATLAB program (The

MathWorks, Natick, MA). DAPI images were analyzed using an edge-detection algorithm in

order to determine the nuclear area for each cell. The nuclear outline was then used as a mask

to quantify the average pixel intensity of the YAP stain within the nuclei of each individual cell.

(n=50-100 nuclei/sample).

Atomic Force Microscopy

Bruker Dimension FastScan AFM was used for collection of the atomic force measurements.

Tipless MLCT-D probes with a 0.03 N/m spring constant were functionalized with 10 µm

diameter borosilicate glass beads for force collection. The AFM's optical microscope was used

to locate individual live MSCs plated on the SlideFlask slides with the flask section removed for

access to the cells. The nucleus of each cell was tested with at least 3 seconds of rest between

each test. In each test, three force-displacement curves were obtained (ramping rate: 2 µm/sec

over 2 µm total travel, 1 µm approach, 1 µm retract), which were analyzed using Nanoscope

software with the implementation of a best-fit curve to a Hertzian (spherical) model (optimized

such that R^2 value was greater than 0.95, or $p < 0.05$) to obtain elastic moduli of nuclear

membrane of individual nuclei.

Statistical analysis

All data analysis results were displayed graphically based on the mean value with standard

error bars. Differences between treatments were not assumed to follow a Gaussian distribution.

Therefore, group differences were identified via either non-parametric two-tailed Mann-Whitney

U-test (**Fig.1a**) or Kruskal-Wallis test followed by Tukey multiple comparison (**Fig.2b, 4, 5, 6, 7,**

&S1). P-values of less than 0.05 were considered significant.

Data availability

The datasets generated and/or analyzed during the current study are available from the

corresponding author on reasonable request.

Acknowledgements

This study was supported by NASA ISGC NNX15AI04H, NIH R01AG059923, and

5P2CHD086843-03, P20GM109095, P20GM103408 and NSF 1929188.

Competing interests

The author(s) declare no competing interests, financial or otherwise.

Contributions

Thompson, M experimental methods, data analysis/interpretation, manuscript writing, final

approval of manuscript.

Woods, K data analysis/interpretation, final approval of manuscript.

Newberg, J experimental methods, final approval of manuscript

Oxford JT, financial support, final approval of manuscript

Uzer, G concept/design, financial support, data analysis/interpretation, manuscript writing, final

approval of manuscript

References

1   Smith, S. M. *et al.* Calcium metabolism before, during, and after a 3-mo spaceflight:
kinetic and biochemical changes. *Am J Physiol* **277**, R1-10 (1999).

364	2	Greenleaf, J. E., Bulbulian, R., Bernauer, E. M., Haskell, W. L. & Moore, T. Exercise-training protocols for astronauts in microgravity. *Journal of applied physiology (Bethesda, Md. : 1985)* **67**, 2191-2204, doi:10.1152/jappl.1989.67.6.2191 (1989).
367	3	Vico, L. *et al.* Effects of long-term microgravity exposure on cancellous and cortical weight-bearing bones of cosmonauts. *Lancet (London, England)* **355**, 1607-1611 (2000).
369	4	Thompson, W. R., Rubin, C. T. & Rubin, J. Mechanical regulation of signaling pathways in bone. *Gene* **503**, 179-193, doi:10.1016/j.gene.2012.04.076 (2012).
371	5	Ozcivici, E. *et al.* Mechanical signals as anabolic agents in bone. *Nature reviews. Rheumatology* **6**, 50-59, doi:10.1038/nrrheum.2009.239 (2010).
373	6	Rando, T. A. & Ambrosio, F. Regenerative Rehabilitation: Applied Biophysics Meets Stem Cell Therapeutics. *Cell Stem Cell* **22**, 306-309, doi:10.1016/j.stem.2018.02.003 (2018).
376	7	Chan, M. E., Uzer, G. & Rubin, C. The Potential Benefits and Inherent Risks of Vibration as a Non-Drug Therapy for the Prevention and Treatment of Osteoporosis. *Current osteoporosis reports*, 1-9, doi:10.1007/s11914-012-0132-1 (2013).
379	8	Judex, S., Gross, T. S. & Zernicke, R. F. Strain Gradients Correlate with Sites of Exercise-Induced Bone-Forming Surfaces in the Adult Skeleton. *Journal of Bone and Mineral Research* **12**, 1737-1745, doi:10.1359/jbmr.1997.12.10.1737 (1997).
382	9	Rubin, C. T. & Lanyon, L. E. Dynamic strain similarity in vertebrates; an alternative to allometric limb bone scaling. *Journal of Theoretical Biology* **107**, 321-327, doi:10.1016/s0022-5193(84)80031-4 (1984).
385	10	Price, C., Zhou, X. Z., Li, W. & Wang, L. Y. Real-Time Measurement of Solute Transport Within the Lacunar-Canalicular System of Mechanically Loaded Bone: Direct Evidence for Load-Induced Fluid Flow. *Journal of Bone and Mineral Research* **26**, 277-285, doi:10.1002/jbmr.211 (2011).
389	11	Gurkan, U. A. & Akkus, O. The Mechanical Environment of Bone Marrow: A Review. *Annals of biomedical engineering* **36**, 1978-1991, doi:10.1007/s10439-008-9577-x (2008).
392	12	Vainionpaa, A. *et al.* Intensity of exercise is associated with bone density change in premenopausal women. *Osteoporosis international : a journal established as result of cooperation between the European Foundation for Osteoporosis and the National Osteoporosis Foundation of the USA* **17**, 455-463, doi:10.1007/s00198-005-0005-x (2006).
397	13	Dickerson, D. A., Sander, E. A. & Nauman, E. A. Modeling the mechanical consequences of vibratory loading in the vertebral body: microscale effects. *Biomechanics and Modeling in Mechanobiology* **7**, 191-202, doi:10.1007/s10237-007-0085-y (2008).
401	14	Coughlin, T. R. & Niebur, G. L. Fluid shear stress in trabecular bone marrow due to low-magnitude high-frequency vibration. *Journal of biomechanics* **45**, 2222-2229, doi:10.1016/j.jbiomech.2012.06.020 (2012).
404	15	Riddle, R. C. & Donahue, H. J. From Streaming Potentials to Shear Stress: 25 Years of Bone Cell Mechanotransduction. *Journal of Orthopaedic Research* **27**, 143-149, doi:10.1002/jor.20723 (2009).
407	16	Fritton, S. P., McLeod, K. J. & Rubin, C. T. Quantifying the strain history of bone: spatial uniformity and self-similarity of low-magnitude strains. *Journal of biomechanics* **33**, 317-325, doi:10.1016/s0021-9290(99)00210-9 (2000).
410	17	Pagnotti, G. M. *et al.* Combating osteoporosis and obesity with exercise: leveraging cell mechanosensitivity. *Nature Reviews Endocrinology*, doi:10.1038/s41574-019-0170-1 (2019).

18   Pongkitwitoon, S., Uzer, G., Rubin, J. & Judex, S. Cytoskeletal Configuration Modulates
Mechanically Induced Changes in Mesenchymal Stem Cell Osteogenesis, Morphology,
and Stiffness. *Scientific reports* **6**, 34791, doi:10.1038/srep34791 (2016).
19   Uzer, G. *et al.* Cell Mechanosensitivity to Extremely Low-Magnitude Signals Is Enabled
by a LINCed Nucleus. *STEM CELLS* **33**, 2063-2076, doi:10.1002/stem.2004 (2015).
20   Uzer, G., Pongkitwitoon, S., Ete Chan, M. & Judex, S. Vibration induced osteogenic
commitment of mesenchymal stem cells is enhanced by cytoskeletal remodeling but not
fluid shear. *Journal of biomechanics* **46**, 2296-2302, doi:10.1016/j.jbiomech.2013.06.008
(2013).
21   Pan, J. X. *et al.* YAP promotes osteogenesis and suppresses adipogenic differentiation
by regulating beta-catenin signaling. *Bone research* **6**, 18, doi:10.1038/s41413-018-
0018-7 (2018).
22   Dupont, S. *et al.* Role of YAP/TAZ in mechanotransduction. *Nature* **474**, 179-183,
doi:http://www.nature.com/nature/journal/v474/n7350/abs/10.1038-nature10137-
unlocked.html#supplementary-information (2011).
23   Kegelman, C. D. *et al.* Skeletal cell YAP and TAZ combinatorially promote bone
development. *FASEB journal : official publication of the Federation of American
Societies for Experimental Biology* **32**, 2706-2721, doi:10.1096/fj.201700872R (2018).
24   Zhao, B. *et al.* TEAD mediates YAP-dependent gene induction and growth control.
Genes & development **22**, 1962-1971, doi:10.1101/gad.1664408 (2008).
25   Benham-Pyle, B. W., Pruitt, B. L. & Nelson, W. J. Cell adhesion. Mechanical strain
induces E-cadherin-dependent Yap1 and beta-catenin activation to drive cell cycle entry.
Science (New York, N.Y.) **348**, 1024-1027, doi:10.1126/science.aaa4559 (2015).
26   Ho, L. T. Y., Skiba, N., Ullmer, C. & Rao, P. V. Lysophosphatidic Acid Induces ECM
Production via Activation of the Mechanosensitive YAP/TAZ Transcriptional Pathway in
Trabecular Meshwork Cells. *Investigative Ophthalmology & Visual Science* **59**, 1969-
1984, doi:10.1167/iovs.17-23702 (2018).
27   Cai, H. & Xu, Y. The role of LPA and YAP signaling in long-term migration of human
ovarian cancer cells. *Cell Communication and Signaling* **11**, 31, doi:10.1186/1478-811X-
11-31 (2013).
28   Driscoll, Tristan P., Cosgrove, Brian D., Heo, S.-J., Shurden, Zach E. & Mauck,
Robert L. Cytoskeletal to Nuclear Strain Transfer Regulates YAP Signaling in
Mesenchymal Stem Cells. *Biophysical journal* **108**, 2783-2793,
doi:10.1016/j.bpj.2015.05.010 (2015).
29   Janmaleki, M., Pachenari, M., Seyedpour, S. M., Shahghadami, R. & Sanati-Nezhad, A.
Impact of Simulated Microgravity on Cytoskeleton and Viscoelastic Properties of
Endothelial Cell. *Sci Rep* **6**, 32418, doi:10.1038/srep32418 (2016).
30   Qian, A. R. *et al.* Fractal Dimension as a Measure of Altered Actin Cytoskeleton in
MC3T3-E1 Cells Under Simulated Microgravity Using 3-D/2-D Clinostats. *Biomedical
Engineering, IEEE Transactions on* **59**, 1374-1380, doi:10.1109/TBME.2012.2187785
(2012).
31   Pardo, S. J. *et al.* Simulated microgravity using the Random Positioning Machine inhibits
differentiation and alters gene expression profiles of 2T3 preosteoblasts. *Am J Physiol
Cell Physiol* **288**, doi:10.1152/ajpcell.00222.2004 (2005).
32   Dai, Z. Q., Wang, R., Ling, S. K., Wan, Y. M. & Li, Y. H. Simulated microgravity inhibits
the proliferation and osteogenesis of rat bone marrow mesenchymal stem cells. *Cell
proliferation* **40**, 671-684, doi:10.1111/j.1365-2184.2007.00461.x (2007).
33   Shi, F. *et al.* Simulated Microgravity Promotes Angiogenesis through RhoA-Dependent
Rearrangement of the Actin Cytoskeleton. *Cellular physiology and biochemistry :
international journal of experimental cellular physiology, biochemistry, and pharmacology*
41, 227-238, doi:10.1159/000456060 (2017).

34 Corydon, T. J. *et al.* Reduced Expression of Cytoskeletal and Extracellular Matrix Genes
in Human Adult Retinal Pigment Epithelium Cells Exposed to Simulated Microgravity.
Cellular physiology and biochemistry : international journal of experimental cellular
physiology, biochemistry, and pharmacology **40**, 1-17, doi:10.1159/000452520 (2016).
35 Chen, Z., Luo, Q., Lin, C., Kuang, D. & Song, G. Simulated microgravity inhibits
osteogenic differentiation of mesenchymal stem cells via depolymerizing F-actin to
impede TAZ nuclear translocation. *Scientific reports* **6**, 30322, doi:10.1038/srep30322
(2016).
36 Uddin, S. M. & Qin, Y. X. Enhancement of osteogenic differentiation and proliferation in
human mesenchymal stem cells by a modified low intensity ultrasound stimulation under
simulated microgravity. *PLoS One* **8**, e73914, doi:10.1371/journal.pone.0073914 (2013).
37 Touchstone, H. *et al.* Recovery of stem cell proliferation by low intensity vibration under
simulated microgravity requires intact LINC complex *npj. Microgravity* **5**,
doi:doi.org/10.1038/s41526-019-0072-5 (2019).
38 Shiu, J.-Y., Aires, L., Lin, Z. & Vogel, V. Nanopillar force measurements reveal actin-
cap-mediated YAP mechanotransduction. *Nature cell biology* **20**, 262-271,
doi:10.1038/s41556-017-0030-y (2018).
39 Saeed, M. & Weihs, D. Finite element analysis reveals an important role for cell
morphology in response to mechanical compression. *Biomechanics and Modeling in*
Mechanobiology, doi:10.1007/s10237-019-01276-5 (2019).
40 Riddick, N., Ohtani, K. & Surks, H. K. Targeting by myosin phosphatase-RhoA
interacting protein mediates RhoA/ROCK regulation of myosin phosphatase. *Journal of*
Cellular Biochemistry **103**, 1158-1170, doi:10.1002/jcb.21488 (2008).
41 Sen, B. *et al.* Mechanical signal influence on mesenchymal stem cell fate is enhanced by
incorporation of refractory periods into the loading regimen. *Journal of biomechanics* **44**,
593-599, doi:10.1016/j.jbiomech.2010.11.022 (2011).
42 De Magistris, P. & Antonin, W. The Dynamic Nature of the Nuclear Envelope. *Current*
biology : CB **28**, R487-r497, doi:10.1016/j.cub.2018.01.073 (2018).
43 Czapiewski, R., Robson, M. I. & Schirmer, E. C. Anchoring a Leviathan: How the
Nuclear Membrane Tethers the Genome. *Frontiers in genetics* **7**, 82,
doi:10.3389/fgene.2016.00082 (2016).
44 Uzer, G. *et al.* Sun-mediated mechanical LINC between nucleus and cytoskeleton
regulates betacatenin nuclear access. *Journal of biomechanics* **74**, 32-40,
doi:10.1016/j.jbiomech.2018.04.013 (2018).

Figure 1. Acute LIV$_{AT}$ application increases nuclear YAP levels. (a) MSCs were subjected to

LIV$_{AT}$ and stained with DAPI (blue) and YAP (red). Confocal images indicated an increased

nuclear YAP levels following acute LIV$_{AT}$ applied as five 20min vibration periods separated by

1hr. (b) Quantitative analysis of confocal images showed a 32% of increase of nuclear YAP in

LIV$_{AT}$ samples compared to controls. n>400/grp, group comparison was made a Mann-Whitney

U-test, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001.

Figure 2. Basal nuclear YAP levels decreased by SMG were rescued by LIV$_{DT}$. (a) MSCs were subjected to SMG, and SMG+LIV$_{DT}$ over 72h period and stained with DAPI (blue) and YAP (red). (b) Quantitative analysis showed a 42% decrease of nuclear YAP levels in the SMG group compared to control levels. The SMG+LIV$_{DT}$ group showed a 67% increase of nuclear YAP when compared to the SMG group. There was no statistically significant difference between CTRL and SMG+LIV$_{DT}$ groups. n>100/grp. Group comparisons were made via Kruskal-Wallis test followed by Tukey multiple comparison, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001.

Figure 3. Experimental design of combined SMG, LIV$_{AT}$ and LIV$_{DT}$ application. MSCs were subcultured and plated in SlideFlasks and allowed to attach for 24h before SlideFlasks were filled with culture medium and sealed. Treatment regimen for MSC's involved 72h of SMG (blue). LIV$_{DT}$ regimen consisted of one treatment cycle every 24hr during SMG treatment with each cycle consisting of 2x 20min LIV with an hour in between (yellow). LIV$_{AT}$ regimen was applied after 72h SMG treatment period and consisted of 5x 20min LIV with an hour in between each (red). For LIV application, MSCs plated in SlideFlasks were placed in LIV device constructed in the lab previous to this research. Vibrations were applied at peak magnitudes of 0.7 g at 90 Hz at room temperature. For SMG application, MSCs plated in SlideFlasks were secured in lab custom-built clinostat inside incubator. The clinostat subjected the MSCs to constant 15 RPM rotation simulated microgravity. After treatment, flask was removed for immunofluorescence staining.

Figure 4. LIV$_{AT}$-induced YAP nuclear entry decreased by SMG was not restored by daily LIV$_{DT}$ application. MSCs were subjected to either CTRL, SMG, SMG+LIV$_{DT}$ over 72h period were subsequently treated with LIV$_{AT}$. Quantitative analysis of confocal images showed that SMG alone decreased basal nuclear YAP levels by 37% which were increased back to control levels in the SMG+LIV$_{DT}$ group. +LIV$_{AT}$ increased nuclear YAP levels in the CTRL, SMG and

SMG+LIV_DT groups by 50%, 69% and 22%, respectively. Nuclear YAP intensity in the

SMG+LIV_DT+LIV_AT group remained not significantly different from the SMG+LIV_AT and 23%

smaller than the LIV_AT group. n>200/grp. Group comparisons were made via Kruskal-Wallis test

followed by Tukey multiple comparison, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001.

Figure 5. LPA treatment increases nuclear YAP levels. MSCs were subjected to LPA

addition at 50µM and 100µM concentrations. Quantitative analysis of confocal images revealed

a 99% and a 107% increase in the 50µM LPA and 100µM LPA treatments compared to DMSO

treated controls, respectively. Nuclear YAP levels for 50µM LPA and 100µM LPA treatments

were not significantly different. n>30/grp. Group comparisons were made via Kruskal-Wallis

test followed by Tukey multiple comparison, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001.

Figure 6. LPA-induced YAP nuclear entry decreased by SMG was alleviated by daily LIV_DT

application. MSCs were subjected to SMG, and parallel SMG+LIV_DT over 72h period at the end

of 72h, samples were treated with either LPA (50µM) or DMSO. Quantitative analysis of

confocal images revealed that LPA addition increased nuclear YAP levels by 105%, 67% and

43% in the CTRL, SMG and SMG+LIV_DT when compared to DMSO controls. When compared to

nuclear YAP intensity of the LPA treatment alone, SMG+LPA and SMG+LIV_DT+LPA samples

were 55% and 29% lower, respectively. YAP nuclear levels in SMG+LIV_DT+LPA remained 70%

higher than SMG+LPA group. n>100/grp. Group comparisons were made via Kruskal-Wallis test

followed by Tukey multiple comparison, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001.

Figure 7. Nuclear stiffness and area were not affected by SMG and SMG+LIV_DT

treatments. MSCs were subjected to SMG, and parallel SMG+LIV_DT over 72h period. (a)

Compared to CTRL samples, AFM measured elastic modulus SMG and SMG+LIV treated

MSCs revealed an apparent decrease in elastic modules that were 21% and 27% below control

levels, measured differences were not statistically significant. n=10/grp (b) Quantitative analysis

of confocal images from the image groups revealed no significant effects of either SMG or LIV_{DT}

treatment on the average nucleus size. n=100/grp. Group comparisons were made via Kruskal-

Wallis test followed by Tukey multiple comparison, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001.

Figure S1. LINC complex disruption decreases nuclear YAP levels and reduces LIV_{AT}-

induced YAP nuclear entry. Plasmids harboring either a dominant negative KASH domain of

Nesprin (DNK) to disable LINC complex function or empty mCherry control (MC) were

overexpressed in MSCs. Following puromycin selection, MC or DNK expressing MSCs were

subjected to LIV_{AT}. Quantitative analysis of confocal images revealed a 49% increase of nuclear

YAP following LIV_{AT} in MC expressing control MSCs. Basal YAP levels of the DNK-CTRL group

was 34% lower compared to MC-CTRL and LIV_{AT} treatment failed to significantly increase

nuclear YAP over DNK-CTRL (18%, NS). n=30/grp, group comparisons were made via

Kruskal-Wallis test followed by Tukey multiple comparison, *p<0.05, **p<0.01, ***p<0.01,

****p<0.0001.

Figure 1.

Figure 1. Acute LIV_{AT} application increases nuclear YAP levels. (a) MSCs were subjected to LIV_{AT} and stained with DAPI (blue) and YAP (red). Confocal images indicated an increased nuclear YAP levels following acute LIV_{AT} applied as five 20min vibration periods separated by 1hr. (b) Quantitative analysis of confocal images showed a 32% of increase of nuclear YAP in LIV_{AT} samples compared to controls. n>400/grp, group comparison was made a Mann-Whitney U-test, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001.

Figure 2. Basal nuclear YAP levels decreased by SMG were rescued by LIV$_{DT}$.
(a) MSCs were subjected to SMG, and SMG+LIV$_{DT}$ over 72h period and stained with DAPI (blue) and YAP (red). (b) Quantitative analysis showed a 42% decrease of nuclear YAP levels in the SMG group compared to control levels. The SMG+LIV$_{DT}$ group showed a 67% increase of nuclear YAP when compared to the SMG group. There was no statistically significant difference between CTRL and SMG+LIV$_{DT}$ groups. n>100/grp. Group comparisons were made via Kruskal-Wallis test followed by Tukey multiple comparison, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001.

Figure 3.

Figure 3. Experimental design of combined SMG, LIV$_{AT}$ and LIV$_{DT}$ application. MSCs were subcultured and plated in SlideFlasks and allowed to attach for 24h before SlideFlasks were filled with culture medium and sealed. Treatment regimen for MSC's involved 72h of SMG (blue). LIV$_{DT}$ regimen consisted of one treatment cycle every 24hr during SMG treatment with each cycle consisting of 2x 20min LIV with an hour in between (yellow). LIV$_{AT}$ regimen was applied after 72h SMG treatment period and consisted of 5x 20min LIV with an hour in between each (red). For LIV application, MSCs plated in SlideFlasks were placed in LIV device constructed in the lab previous to this research. Vibrations were applied at peak magnitudes of 0.7 g at 90 Hz at room temperature. For SMG application, MSCs plated in SlideFlasks were secured in lab custom-built clinostat inside incubator. The clinostat subjected the MSCs to constant 15 RPM rotation simulated microgravity. After treatment, flask was removed for immunofluorescence staining.

Figure 4.

Figure 4. LIV$_{AT}$-induced YAP nuclear entry decreased by SMG was not restored by daily LIV$_{DT}$ application. MSCs were subjected to either CTRL, SMG, SMG+LIV$_{DT}$ over 72h period were subsequently treated with LIV$_{AT}$. Quantitative analysis of confocal images showed that SMG alone decreased basal nuclear YAP levels by 37% which were increased back to control levels in the SMG+LIV$_{DT}$ group. +LIV$_{AT}$ increased nuclear YAP levels in the CTRL, SMG and SMG+LIV$_{DT}$ groups by 50%, 69% and 22%, respectively. Nuclear YAP intensity in the SMG+LIV$_{DT}$+LIV$_{AT}$ group remained not significantly different from the SMG+LIV$_{AT}$ and 23% smaller than the LIV$_{AT}$ group. n>200/grp. Group comparisons were made via Kruskal-Wallis test followed by Tukey multiple comparison, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001.

Figure 5.

Figure 5. LPA treatment increases nuclear YAP levels. MSCs were subjected to LPA addition at 50µM and 100µM concentrations. Quantitative analysis of confocal images revealed a 99% and a 107% increase in the 50µM LPA and 100µM LPA treatments compared to DMSO treated controls, respectively. Nuclear YAP levels for 50µM LPA and 100µM LPA treatments were not significantly different. n>30/grp. Group comparisons were made via Kruskal-Wallis test followed by Tukey multiple comparison, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001.

Figure 6.

Figure 6. LPA-induced YAP nuclear entry decreased by SMG was alleviated by daily LIV$_{DT}$ application. MSCs were subjected to SMG, and parallel SMG+LIV$_{DT}$ over 72h period at the end of 72h, samples were treated with either LPA (50µM) or DMSO. Quantitative analysis of confocal images revealed that LPA addition increased nuclear YAP levels by 105%, 67% and 43% in the CTRL, SMG and SMG+LIV$_{DT}$ when compared to DMSO controls. When compared to nuclear YAP intensity of the LPA treatment alone, SMG+LPA and SMG+LIV$_{DT}$+LPA samples were 55% and 29% lower, respectively. YAP nuclear levels in SMG+LIV$_{DT}$+LPA remained 70% higher than SMG+LPA group. n>100/grp. Group comparisons were made via Kruskal-Wallis test followed by Tukey multiple comparison, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001.

(a)

(b)

Figure S1. LINC complex disruption decreases nuclear YAP levels and reduces LIV$_{AT}$-induced YAP nuclear entry. Plasmids harboring either a dominant negative KASH domain of Nesprin (DNK) to disable LINC complex function or empty mCherry control (MC) were overexpressed in MSCs. Following puromycin selection, MC or DNK expressing MSCs were subjected to LIV$_{AT}$. Quantitative analysis of confocal images revealed a 49% increase of nuclear YAP following LIV$_{AT}$ in MC expressing control MSCs. Basal YAP levels of the DNK-CTRL group was 34% lower compared to MC-CTRL and LIV$_{AT}$ treatment failed to significantly increase nuclear YAP over DNK-CTRL (18%, NS). n=30/grp, group comparisons were made via Kruskal-Wallis test followed by Tukey multiple comparison, *p<0.05, **p<0.01, ***p<0.01, ****p<0.0001.

Supplementary Information

Low Intensity Vibration Restores Nuclear YAP Levels and Acute YAP Nuclear Shuttling in Mesenchymal Stem Cells Subjected to Simulated Microgravity

Thompson M[1], Woods K[2], Newberg, J[1], Oxford JT, Uzer G[1†]

[1]Mechanical and Biomedical Engineering, Boise State University
[2]Biomolecular Sciences Graduate Program, Boise State University
† Corresponding Author

Funding support:
NASA ISGC NNX15AI04H, NIH R01AG059923, and 5P2CHD086843-03, P20GM109095, P20GM103408 and NSF 1929188.

Figure S1.

Figure S1. LINC complex disruption decreases nuclear YAP levels and reduces LIV$_{AT}$-induced YAP nuclear entry. Plasmids harboring either a dominant negative KASH domain of Nesprin (DNK) to disable LINC complex function or empty mCherry control (MC) were overexpressed in MSCs. Following puromycin selection, MC or DNK expressing MSCs were subjected to LIV$_{AT}$. Quantitative analysis of confocal images revealed a 49.27% increase of nuclear YAP following LIV$_{AT}$ in MC expressing control MSCs. Basal YAP levels of the DNK-CTRL group was 33.91% lower compared to MC-CTRL and LIV$_{AT}$ treatment failed to significantly increase nuclear YAP over DNK-CTRL (18.33%, NS). n=30/grp, group comparisons were made via Kruskal-Wallis test followed by Tukey multiple comparison, *$p < 0.05$, **$p < 0.01$, ***$p < 0.01$, ****$p < 0.0001$.