**Pro-inflammatory response of non-inflammatory Classical monocytes stimulated**
**with LPS *in vitro*.**

Claudio Karsulovic[1,2], Fabian Tempio[3,4] , Mercedes Lopez[3,4], Julia Guerrero[1], Annelise Goecke[1,2]

[1] Laboratorio de Inmunomodulación Neuroendocrina, Instituto de Ciencias Biomédicas, Facultad de
Medicina, Universidad de Chile, Santiago, Chile.

[2] Sección de Reumatología, Hospital Clínico Universidad de Chile, Universidad de Chile, Santiago,
Chile.

[3] Laboratorio de Regulación e Inmunología del Cáncer, Facultad de Medicina, Universidad de Chile,
Santiago, Chile.

[4] Instituto Milenio de Inmunologia e Inmunoterapia, Facultad de Medicina, Universidad de Chile

**\* Correspondence:**
Correspondence has to be addressed to Annelise Goecke at igoecke@hcuch.cl

**Keywords: Monocytes, Phenotype, Classical, Intermediate, Non-Classical, LPS, flow**
**cytometry.**

**Abstract**

BACKGROUND: CD14 (Monocyte identifying Toll-Like Receptor) and CD16 (FcyRIII co-receptor, marker of
inflammatory monocytes) were used to define 3 subpopulations of circulating monocytes with different attributes in terms
of inflammatory and phagocytic capabilities. There are contradictory reports regarding response of circulating monocytes
to pro-inflammatory or non-inflammatory stimuli *in vitro*. Here we aimed to analyze the phenotypic changes in
circulating monocytes when stimulated with pro and non-inflammatory stimuli.

METHODS: Whole blood from 9 healthy donors was extracted and studied. Monocyte subpopulations were directly
measured using flow cytometry with PBMC Ficoll extraction method. Pro-inflammatory interleukin IL-1β was measured
by intracellular cytometry. Whole blood-extracted monocytes were stimulated using LPS and IL-4 as previously
described. Changes against non-stimulated (N-S) populations were statistically analyzed.

RESULTS: Compared to N-S, LPS-stimulated monocytes display a singular milieu of markers, with higher levels of
intracellular IL-1β in parallel raise of CD14+CD163-/CD14+CD163+ ratio. CD163 shows positive correlation with levels
of IL-1β. In t-SNE (T-distributed Stochastic Neighbor Embedding) analysis, after LPS stimulation, subpopulation
CD14+CD16-CD163-, containing mainly classical monocytes, show a higher number of IL-1β+ cells.

CONCLUSION: Classical monocytes, the non-inflammatory subset, show higher levels of IL-1β in response to LPS
narrowing down to a new subpopulation of monocytes CD14+CD16-CD163-, which correlates better with this interleukin
response than widely used monocytes classification. Using CD163 in addition to CD16, we were able to show that
classical monocytes display the most intense response to LPS. Additionally, CD163 appears to be a suitable addition to
CD14-CD16 classification to improve its performance.

**Introduction**

In 2010, an international meeting sponsored by the IUIS (International Union of Immunological
Societies) and the WHO (World Health Organization), proposed a subset classification for circulating
monocytes (1). CD14 (Monocyte identifying Toll-Like Receptor) and CD16 (FcyRIII co-receptor,
marker of inflammatory monocyte) were used to define 3 subpopulations of circulating monocytes
with different attributes in terms of inflammatory protein expression and phagocytic capabilities (2).
Classical monocytes (CD14++CD16-) are 80 to 90% percent of the entire monocyte's population,

express low levels of pro-inflammatory proteins and high phagocytic characteristics. Intermediate
monocytes (CD14+CD16+) are 5 to 15% of the total monocytes. Express variable levels of pro-
inflammatory cytokines and intermediate phagocytic capacities. Non-classical monocytes
(CD14+CD16++) (3) are approximately 5% of all monocytes, express higher levels of pro-
inflammatory cytokines and minor phagocytic characteristics. The last two subsets, since have shown
to express higher levels of inflammatory cytokines when stimulated with pro inflammatory stimuli,
have been repeatedly related to inflammatory pathologies. (4).

During the last years, multiple studies have been published associating expansion or contraction of
inflammatory or non-inflammatory monocytes subpopulations in diseases like lupus, rheumatoid
arthritis, atherosclerosis or membranous glomerulonephritis (5-7), among others.  Intermediate
monocytes has shown to be expanded in infectious and non-infectious inflammatory conditions,
however many studies presents inconsistent and contradictory data.(8).

In cardiovascular diseases, intermediate monocytes have been reported both expanded and
contracted(8, 9). In the other hand, Classical monocytes show higher frequencies in atherosclerotic
lesions despite of inflammatory status, evidencing conflicting data in this regard(9, 10). In 2017,
Williams et al. classified circulating monocytes from post-stroke patients, using CD86 (co-
stimulatory T signal, typically described M1 Monocyte/Macrophage phenotype) and CD163 (High-
affinity scavenger receptor typically related with M2 Monocyte/Macrophage phenotype) to group
monocytes as M1 and M2 trying to emulate macrophage classification. They showed that M1/M2
ratio (CD86/CD163 ratio) was higher in post-stroke patients(11, 12). In other clinical entities like
membranous glomerulopathy, similar results have been found using broader group of markers
defining M1 and M2 phenotypes (5).

Until now, many changes have been introduced looking for accuracy to define monocytes
subpopulations. Thomas et al. using multiparametric analysis, tSNE (t-distributed stochastic neighbor
embedding), showed that CCR2 (CC Chemokine Receptor Type 2), HLA-DR (Human Leukocyte
Antigen – DR isotype) and CD163 were good and enough markers to a proper M1 and M2
monocytes classification (10). Interestingly CD163 and CD16, protein which defines current
monocyte classification, does not show correlation in clinical setting (13).

In vitro, when monocytes are stimulated with pro-inflammatory and anti-inflammatory stimuli, there
are also contradictory reports in terms of interleukin production response. Mukerjhee et al. showed
that monocytes from healthy donors do not display cell number changes when stimulated with LPS.
Also, intermediate and non-classical monocytes express higher levels of IL-1β and TNF-α, but
inflammatory intermediate subpopulation, responds with higher levels of regulatory IL-10 to the
same stimulus (7). Subsequent studies have shown that M1 monocytes express higher levels of IL-6
when stimulated with LPS exposing again contradictory reports(11).

Given the absence of a solid classification strategy and in vitro monocyte response to inflammatory
stimuli, we analyzed monocytes from nine healthy donors and expose it to LPS and IL-4 showing a
singular cytokine production and phenotypic pattern. We found that less-reactive-non-inflammatory
classical monocyte subpopulation has the most robust response *in vitro* to LPS stimulus. This
subpopulation also could be better classified using other membrane markers as CD163. Classical
monocytes show also a more "flexible" M1/M2 ratio, compared to intermediate and non-classical
monocytes, becoming the subpopulation with the highest functional plasticity.

Pro-inflammatory response of non-inflammatory Classical monocytes stimulated with LPS *in vitro*.

## Material and Methods

*2.1.Healthy donors.* We recruited 9 healthy donors ranging from 26 to 40 years, 5 man and 4 women, from our facility. A medical anamnesis was performed by an internal medicine specialist before whole blood extraction. No medication uses nor relevant past medical history were reported. The ethics committee of Facultad de Medicina de la Universidad de Chile approved the study protocol.

*2.2.Whole blood lysis protocol and flow cytometry.* Venous blood (30ml) was obtained by cubital venopunction from all participants. PBMC Ficoll extraction was performed as previously dercribed(REF). PBMC were stained in duplicate with the following antibodies: FITC-A-anti-CD14, APC-H7-anti CD16, PERCP-Cy5.5-anti-CD86, APC-A-anti-CD16, V450-A-anti-IL-1β (Biolegend, US) at room temperature for 30 minutes. 1ml of BD FACS Lysing Solution (BD Biosciences) at 1x concentration was used for erythrocyte lysis. Finally cells we fixed and permeabilized using a fixation/permeabilization BD Kit (BD Biosciences).

2.3.*LPS and IL-4 Stimulation and flow cytometry.* To phenotype monocyte subsets, $1 \times 10^6$/cells per tube were stimulated in duplicate with lipopolysaccharide, (1 µg/mL) (BD Biosciences, NY) and IL-4 (200 U/ml) (BD Biosciences, NY) in RPMI 1640 (BD Biosciences, NY) medium mixed with 10% fetal bovine serum (BD Biosciences, NY) for 2 h and 4 h respectively at 37°C in 5% CO2 and exposed to brefeldin A (BD Biosciences) as described in previous studies. Then, the cells were washed, followed by staining with V710-anti-CD3 (Biolegend, US), V650-anti-CD19, (Biolegend, US), FITC-A-anti-CD14, APC-H7-anti CD16, PERCP-Cy5.5-anti-CD86, APC-A-anti-CD16, (Biolegend, US). Subsequently, they were fixed and permeabilized and stained with V450-A-anti-IL-1β (Biolegend, US). The frequencies monocyte subsets were assessed on a FACS Verse instrument (BD, Franklin Lakes, US), and the data was analyzed with FlowJo software (v7.6.1; TreeStar; Ashland, US).

2.4.*Statistical Analysis*. Variables were shown as median and range values. We employed the Mann–Whitney U nonparametric test to evaluate the differences among groups. The relationship between variables was analyzed by the Spearman rank correlation test. All of the data were carried out with the GraphPad Prism version 6.01 software. P value of <0.05 represented statistical significance.

## Results

*Classical monocytes are the most responsive subpopulation to LPS stimuli measured as IL-1β MFI.* Monocytes where gated using CD14 and CD16 for usual three-subsets classification (Fig1, A, B, C). Classical monocytes are significantly higher than intermediate and non-classical monocytes in any condition and percentages are equivalent to previously described studies(1) (Fig1, D). In each subpopulation, levels of IL-1β measured as Mean Fluorescence Intensity (MFI) and count of IL-1β+ monocytes, are higher when stimulated with LPS, being significantly higher in Classical versus Non-classical monocytes (Fig1 E, F).

*CD163- monocytes rise in response to LPS and shows higher levels of intracellular IL-1β.* Monocytes were firstly gated using CD14 and then, using CD163, two different subpopulations where found with evenly split frequencies (Fig2 A, B, C).When expose with LPS, CD163- cells

count raises to a maximum reciprocally to CD163+ cells which shows lowest levels. IL-4
monocytes maintain 50%-50% proportion seen on basal conditions (Fig2, D). CD163- cells
contains significantly higher levels of IL-1β than CD163+ monocytes (Fig2, E). When
CD14+CD163-/CD14+CD163+ Ratio is analyzed, the abrupt rise in CD163- cells seen when
expose to LPS, reflects as a higher ratio only in Classical monocytes, suggesting higher plasticity
on this subpopulation (Fig2, F).

136
137    *Only CD163 and no CD16, show correlation with IL-1β after LPS stimulation*. When we
138    analyzed separately positivity to either CD163 or CD16 and levels of IL-1β after LPS
139    stimulation, only CD163 negativity pairs with IL-1β peaks in all subsets (Fig3, A). After
140    correlation tests, again only CD163 show positive and significant correlation with MFI IL-1β
141    after LPS stimulation (Fig3, B, C). No correlation is seen between CD16 and IL-1β (Fig3, D, E).

142
*CD14+CD163- inflammatory subset rather CD14+CD16+ subset, respond to LPS stimulation*
*raising intracellular IL-1β*. Using t-SNE multiparametric reduction (of CD3, CD19, CD14,
CD16, CD163, CD86 markers), we were able to reduce all monocytes to two cell clouds,
CD14+CD16+ and CD14+ CD163- which interestingly, where mutually exclusive (Fig4, A).
When stimulated with LPS, IL-1β+ monocytes invade CD163- cells without significantly grow
on CD16+ cloud, showing that CD163- monocytes are especially responsive shaping a new
inflammatory subpopulation CD14+CD16-CD163-IL-1β+ (Fig4, C).
150

## Discussion

During the last years, numerous studies have tried to find an association between certain monocyte
subpopulations and prevalent inflammatory and metabolic diseases. Autoimmune entities and
atherosclerosis as a metabolic dysregulation are the most usually studied, nevertheless, their
correlation with expansion or contraction of these subpopulations has been difficult to study and not
completely reproducible. Many reasons for the latter have been proposed. One of the most frequently
argued explanation is how difficult is to separate three "different" subpopulations using just a small
group of markers. Many times, depending on the inflammatory status of the patient, the first selection
criteria in the gating process, as is the FSC (granularity) in flow cytometry, could represent a
challenge trying to gate only monocytes out of a "universe" of neutrophils.

Thomas et al. were able to demonstrate using tSNE with more than 20 different parameters, that a
group of at least 12 markers commonly available, must be used to perform positive and negative
selection in order to obtain a proper separation (14). Even when these markers help to better classify
circulating monocytes, it is yet expensive and time consuming to process it.

When these monocyte subpopulations are analyzed *in vitro*, its response to stimuli turns more erratic.
Most of the studies available, are prone to point to intermediate monocyte subpopulation as the most
"plastic" and responsive to LPS. Many times, intermediate monocytes are presented as transition
subpopulation with oscillations on percentages attributed to inflammatory nature on each pathology.
Despite all this, until today, there is no clear correlation between any subpopulation rate changes and
clinical course in inflammatory or non-inflammatory conditions.

In our experience, we were able to show that non-inflammatory classical monocytes constitute the
most responsive subpopulation. Classical monocytes uniquely raise levels of intracellular IL-1β.
Also, when these monocytes are reclassified using CD163, CD163- cells are the most responsive.
When compared CD16 marker and CD163 marker, the last is the one who better correlate with

intracellular levels of IL-1β. Classical monocytes subset, a priori classified as less-responsive/non-
inflammatory subset, contains almost 100% of high responsive CD163- cells.

We were also able to show that, in the way of a Russian *matryoshka*, classical monocytes display an
intern and variable plasticity, with different grades of responsiveness depending at least in part, on
CD163- cells number. These findings rise the question about if we are properly classifying
monocytes subsets using a bidimensional analysis. More studies are needed to understand if the most
responsive subpopulation CD14+CD163- and a new subpopulation CD14+CD16-CD163-IL-1β+
displays different sub-phenotypes with also different grades of response to LPS and other stimuli
using more markers or multiparametric analysis. How these sub-phenotypes behave in clinical
scenarios remain a question to be investigated.

**Figures**

187
FIGURE 1: Gating strategy, flow cytometry analysis and intracellular IL-1β in CD14-CD16 defined
subpopulations under different stimulation conditions. Whole blood from healthy donors was stained
with anti-CD14, anti-CD16, anti-CD86 and anti-CD163. The cells were gated firstly on mononuclear
cells (A) and secondly on CD14+ monocytes (B). Afterwards, the count of CD14++CD16-,
CD14+CD16+ and CD14+CD16++ were determined by flow cytometry (C). Classical monocytes are
significantly higher than intermediate and non-classical monocytes in any condition
(D)(****p≤0,0001). In each subpopulation, levels of IL-1β are higher when stimulated with LPS,
being significantly higher in Classical versus Non-classical monocytes (E) (****p≤0,0001;
**p≤0,007). When IL-1β+ cells are analyzed, LPS stimulated monocytes, show significantly higher
cell count in Classical versus Non-Classical monocytes (F) (**p≤0,009).

198
199
200

Pro-inflammatory response of non-inflammatory Classical monocytes stimulated with LPS *in vitro*.

201
FIGURE 2: Gating strategy, flow cytometry analysis, intracellular IL-1β in CD14-CD163 defined
subpopulations under different stimulation conditions. Whole blood stained with anti-CD14, anti-
CD16, anti-CD86 and anti-CD163. The cells were gated firstly on mononuclear cells (A) and
secondly on CD14+ monocytes (B). Afterwards, the count of CD14+CD163- and CD14+CD163+
were determined by flow cytometry (C). CD163- cells count raises only when stimulated with LPS,
reciprocally to CD163+ cells (D)(****p≤0,0001). CD163- cells contains significantly higher levels
of IL-1β than CD163+ monocytes (E)(**p≤0,009). Only in Classical monocytes stimulated with
LPS, CD163-/CD163+ ratio raises (F)(****p≤0,0001).
.
211

Pro-inflammatory response of non-inflammatory Classical monocytes stimulated with LPS *in vitro*.

212
FIGURE 3 : Intracellular IL-1β correlation with both CD16 and CD163 markers. Association
between levels of IL-1β and CD16-CD163 positivity/negativity in each stimulation condition. In A,
positivity for CD16 marker, did not show relation with levels of IL-1β. On the contrary, negativity
for CD163 is present every time IL-1β raises, being significantly higher in N-S condition (*p≤0,02).
Correlation between CD163 and CD16 cells and levels of intracellular IL-1β. CD14+CD163- cells
correlate positive and significantly with levels of IL-1β, on the contrary CD163+ have negative
correlation (B, C). CD16+ o CD16- cells did not correlate significantly with levels of intracellular IL-
1β (D, E).
221
222

Pro-inflammatory response of non-inflammatory Classical monocytes stimulated with LPS *in vitro*.

223
FIGURE 4: t-SNE analysis for CD14+ monocytes using multiple markers. Orange colored cloud of
cells represents CD16+ cells, light blue cloud CD163- cells and green cloud IL-1β+ cells, being the
first two, "inflammatory" subpopulation according most used classifications. As is easily seen,
CD16+ and CD163- cells are completely different subpopulations (A). In IL-4 stimulation condition,
no changes are seen in green cloud, which is comparable to N-S condition (B). In C, after LPS
stimulation, green cloud "invades" CD16-CD163- forming a new subpopulation CD14+CD16-
CD163-IL-1β+ (C).
231

**Conflict of Interest**

*The authors declare that the research was conducted in the absence of any commercial or financial*
*relationships that could be construed as a potential conflict of interest.*

**Funding**

**References**

1.      Ziegler-Heitbrock L, Ancuta P, Crowe S, Dalod M, Grau V, Hart DN, et al. Nomenclature of
monocytes and dendritic cells in blood. Blood. 2010 Oct 21;116(16):e74-80.

2.      Ziegler-Heitbrock L. Blood Monocytes and Their Subsets: Established Features and Open
Questions. Front Immunol. 2015;6:423.

3.      Burbano C, Vasquez G, Rojas M. Modulatory effects of CD14+CD16++ monocytes on
CD14++CD16- monocytes: a possible explanation of monocyte alterations in systemic lupus
erythematosus. Arthritis Rheumatol. 2014 Dec;66(12):3371-81.

4.        Kunnathully V, Gomez-Lira M, Bassi G, Poli F, Zoratti E, La Verde V, et al. CD14(++)
CD16(-) monocytes are the main source of 11beta-HSD type 1 after IL-4 stimulation. Int
Immunopharmacol. 2017 Feb;43:156-63.

5.        Hou J, Zhang M, Ding Y, Wang X, Li T, Gao P, et al. Circulating
CD14(+)CD163(+)CD206(+) M2 Monocytes Are Increased in Patients with Early Stage of
Idiopathic Membranous Nephropathy. Mediators Inflamm. 2018;2018:5270657.

6.        Rossol M, Kraus S, Pierer M, Baerwald C, Wagner U. The CD14(bright) CD16+ monocyte
subset is expanded in rheumatoid arthritis and promotes expansion of the Th17 cell population.
Arthritis Rheum. 2012 Mar;64(3):671-7.

7.        Mukherjee R, Kanti Barman P, Kumar Thatoi P, Tripathy R, Kumar Das B, Ravindran B.
Non-Classical monocytes display inflammatory features: Validation in Sepsis and Systemic Lupus
Erythematous. Sci Rep. 2015 Sep 11;5:13886.

8.        Wildgruber M, Czubba M, Aschenbrenner T, Wendorff H, Hapfelmeier A, Glinzer A, et al.
Increased intermediate CD14(++)CD16(++) monocyte subset levels associate with restenosis after
peripheral percutaneous transluminal angioplasty. Atherosclerosis. 2016 Oct;253:128-34.

9.        Ammirati E, Moroni F, Magnoni M, Di Terlizzi S, Villa C, Sizzano F, et al. Circulating
CD14+ and CD14(high)CD16- classical monocytes are reduced in patients with signs of plaque
neovascularization in the carotid artery. Atherosclerosis. 2016 Dec;255:171-8.

10.       Thomas GD, Hamers AAJ, Nakao C, Marcovecchio P, Taylor AM, McSkimming C, et al.
Human Blood Monocyte Subsets: A New Gating Strategy Defined Using Cell Surface Markers
Identified by Mass Cytometry. Arterioscler Thromb Vasc Biol. 2017 Aug;37(8):1548-58.

11.       Fukui S, Iwamoto N, Takatani A, Igawa T, Shimizu T, Umeda M, et al. M1 and M2
Monocytes in Rheumatoid Arthritis: A Contribution of Imbalance of M1/M2 Monocytes to
Osteoclastogenesis. Front Immunol. 2017;8:1958.

12.       Williams H, Cassorla G, Pertsoulis N, Patel V, Vicaretti M, Marmash N, et al. Human
classical monocytes display unbalanced M1/M2 phenotype with increased atherosclerotic risk and
presence of disease. Int Angiol. 2017 Apr;36(2):145-55.

13.       Rana AK, Li Y, Dang Q, Yang F. Monocytes in rheumatoid arthritis: Circulating precursors
of macrophages and osteoclasts and, their heterogeneity and plasticity role in RA pathogenesis. Int
Immunopharmacol. 2018 Dec;65:348-59.

14.       Zhou L, Somasundaram R, Nederhof RF, Dijkstra G, Faber KN, Peppelenbosch MP, et al.
Impact of human granulocyte and monocyte isolation procedures on functional studies. Clin Vaccine
Immunol. 2012 Jul;19(7):1065-74.

279