Retrograde labeling illuminates distinct topographical organization of D1 and D2 receptor-positive neurons in the prefrontal cortex of mice.

Sara M. Green, Sanya Nathani, Joseph Zimmerman, David Fireman, Nikhil M. Urs

Department of Pharmacology, College of Medicine, University of Florida.

Correspondence:

Nikhil M. Urs

1200 Newell Dr

ARB-R5-140

Gainesville FL 32610

nikhilurs@ufl.edu

ABSTRACT

The cortex plays an important role in regulating motivation and cognition, and does so by regulating multiple subcortical brain circuits. Glutamatergic pyramidal neurons in the prefrontal cortex (PFC) are topographically organized in different subregions such as the prelimbic, infralimbic and orbitofrontal, and project to topographically-organized subcortical target regions. Dopamine D1 and D2 receptors are expressed on glutamatergic pyramidal neurons in the PFC. However, it is unclear whether D1 and D2 receptor-expressing pyramidal neurons in the PFC are also topographically organized. We used a retrograde adeno-associated virus (AAVRG)-based approach to illuminate the topographical organization of D1 and D2 receptor-expressing neurons, projecting to distinct striatal and midbrain subregions. Our experiments reveal that AAVRG injection in the nucleus accumbens (NAcc) or dorsal striatum (dSTR) of D1Cre mice labeled distinct neuronal subpopulations in medial orbitofrontal or prelimbic PFC, respectively. However, AAVRG injection in NAcc or dSTR of D2Cre mice labeled medial orbitofrontal, but not medial prelimbic PFC, respectively. Additionally, D2R+ but not D1R+ PFC neurons were labeled upon injection of AAVRG in substantia nigra pars compacta (SNpc). Thus, our data are the first to highlight a unique dopamine receptor-specific topographical pattern in the PFC, which could have profound implications for corticostriatal signaling in the basal ganglia.

INTRODUCTION

Dopamine regulates normal processes such as motivation, reinforcement-based learning, reward and movement (Palmiter, 2008; Schultz, 2002), and its dysfunction is implicated in many psychiatric and neurological disorders such as schizophrenia, Parkinson's disease, obsessive compulsive disorder, attention deficit hyperactivity disorder (ADHD), and addiction (Abi-Dargham, 2018; Robinson et al., 2006; Sulzer, 2011; Yager et al., 2015; Zhai et al., 2018).

The striatum is the main input center of the basal ganglia, receiving input from cortical, thalamic, limbic and dopaminergic nuclei (Ikemoto and Bonci, 2014). Glutamatergic signals from cortical areas, and dopamine signals from midbrain dopaminergic nuclei act upon spiny projection neurons (SPNs) in the striatum. The SPNs integrate both glutamate and dopamine signals and coordinate various aspects of learning and behavior (Bamford et al., 2018; Horvitz, 2009; Shiflett and Balleine, 2011).

Afferent cortical glutamatergic inputs into the striatum originate within various medial or lateral subregions of the PFC such as the prelimbic, infralimbic, orbitofrontal, and motor cortex. The PFC plays a critical role in motivation and cognition (Balleine and O'Doherty, 2010; Smith and Graybiel, 2014). Within the medial and lateral PFC, topographically-organized regions such as the dorsally located prelimbic corticostriatal neurons, and the ventral infralimbic or orbitofrontal PFC (OFC) neurons have dissociable effects on motivated behavior and cognitive flexibility (Ahmari et al., 2013; Barker et al., 2017; Hart et al., 2018; Killcross and Coutureau, 2003). (Burguiere et al., 2013; Gremel and Costa, 2013; Rudebeck and Murray, 2011). Although, these studies elegantly outline the role of sub-regions in the PFC or striatum, few studies have explored the neuronal and molecular diversity of PFC pyramidal neurons involved in regulating motivation and cognition. Dopamine and its receptors in the PFC also regulate motivated behavior and cognitive flexibility (Barker et al., 2013; Goldman-Rakic, 1998; Hitchcott et al., 2007; Natsheh and Shiflett, 2018; Ott and Nieder, 2019)

Dopamine activates D1 (stimulatory Gαs-coupled) and D2 (inhibitory Gαi-coupled) class of receptors in the PFC, which modulate the activity of both pyramidal neurons and interneurons (Ferguson and Gao, 2018; Santana et al., 2009; Tomasella et al., 2018; Tseng and O'Donnell, 2004, 2007). Moreover, pharmacological targeting of D1Rs or D2Rs, or genetic deletion of D2Rs in the PFC can regulate dopamine-dependent behaviors such as locomotion and goal-directed behavior (Barker et al., 2013; Del Arco et al., 2007; Hitchcott et al., 2007; Natsheh and Shiflett, 2018; Tomasella et al., 2018; Urs et al., 2016). Although PFC dopamine receptors play an important role in motivation and cognition, the topographical distribution of D1R+ or D2R+ neurons in specific subregions of the PFC is not known. Here, we use a retrograde AAV (AAVRG)-based approach to identify distinct topographically-organized subpopulations of D1R+ and D2R+ neurons in the PFC based on their target projection areas. Given the role of various subregions of the PFC in motivated behavior and cognition, and the heterogeneity of these regions, the effects of dopamine and its receptors within these subregions will expand our understanding of the molecular and neuronal mechanisms regulating motivated behavior and cognition.

RESULTS

For this study we focused primarily on dopamine-related subcortical target regions such as the striatum and midbrain dopamine nuclei i.e. substantia nigra pars compacta (SNpc). The striatum itself can be topographically divided along the dorsoventral or mediolateral axes, into the dorsal and ventral striatum, or dorsomedial (DMS) and dorsolateral (DLS), respectively.

Dorsoventral Topographical distribution of D1R+ and D2R+ neurons in the PFC.

We first injected AAVRGs in topographically distinct target regions along the dorsoventral axis in the dorsal striatum (dSTR) and the nucleus accumbens (NAcc) core within the ventral striatum, respectively, of either D1Cre or D2Cre mice. 100nl of AAVRG-CAG-Flex-TdTomato and AAVRG-

hSyn-DIO-GFP were stereotaxically injected in the dSTR or NACC core (see methods for coordinates), respectively. As shown in Figure 1, we observed distinct topographically-organized patterns of GFP (dSTR) and TdT (NAcc core) positive cells in the PFC of D1Cre and D2Cre mice. For D1Cre mice, AAVRG injection in the dSTR (GFP+) and NAcc (TdT+) labeled distinct minimally overlapping subpopulations of dorsally located layer 5, and ventrally located layer 2/3 neuronal cell bodies in the PFC, respectively (Figure 1, ai). The dSTR projecting neurons were primarily localized to dorsally located prelimbic/cingulate (Cg1/PrL) and motor cortex (M1/M2), and mediolaterally located ventral/lateral OFC (VO/LO), whereas the NAcc projecting neurons were primarily localized to the ventrally located medial OFC and infralimbic (MO/IL) regions. Quantifying the number of cell bodies in these regions from rostral, rostrocaudal and caudal PFC sections (see methods) revealed minimal overlap (~5%) between the PrL and MO/IL subpopulations (Figure 1 c, ** p< 0.01, Two-way ANOVA). "D1-dSTR" - Cg1/PrL: 145 ± 11.3; MO/IL: 22.5 ± 3.9; VO/LO: 121.7 ± 16; M1/M2: 122 ± 11.2 neurons, and "D1-NAcc" - Cg1/PrL: 20.8 ± 9.1; MO/IL: 93.4 ± 10.6; VO/LO: 10.1 ± 2.2; M1/M2: 6.7 ± 6.6 neurons. In contrast, for D2Cre mice, AAVRG injection in the dSTR and NAcc labeled predominantly ventrally located TdT+ layer 5 neuronal cell bodies in MO/IL, but few dorsally located GFP+ neurons in Cg1/PrL (Figure 1, bi). Quantifying the number of cell bodies in these regions revealed minimal overlap of TdT+ neurons (~15%) between the PrL and MO/IL subpopulations (Figure 1 d, ** p< 0.01, Two-way ANOVA). The NAcc projecting neurons were primarily localized to the ventrally located medial OFC and infralimbic (MO/IL) regions, whereas the few dSTR projecting neurons were primarily localized to M1/M2 and VO/LO regions. "D2-dSTR" - Cg1/PrL: 7 ± 1.8; MO/IL: 3.7 ± 1; VO/LO: 24.4 ± 5.9; M1/M2: 45.7 ± 11.8 neurons, and "D2-NAcc" - Cg1/PrL: 55.7 ± 9.7; MO/IL: 162.9 ± 25.5; VO/LO: 32.8 ± 6.1; M1/M2: 5.6 ± 1.7 neurons. Thus, in the medial PFC of D2Cre mice, MO/IL neurons were predominantly labeled, but not Cg1/PrL neurons.

Figure 1. <u>**Unique topographically organized subpopulations of D1R+ and D2R+ PFC neurons project to dSTR and NAcc.**</u> PFC sections from D1Cre (**ai**) and D2Cre (**bi**) mice showing cell bodies retrograde labeled with both TdTomato and GFP. Left panels are zoomed insets of medial PrL and MO showing layer-specific localization of cell bodies. PrL-prelimbic, MO-medial orbitofrontal, VO- ventral orbitofrontal, LO-lateral orbitofrontal, M1-primary motor, M2- secondary motor cortex. **aii, bii.** Injection sites are shown in striatal (STR) sections along with midbrain SNpc/VTA (SN VTA) sections. D1Cre and D2Cre mice were injected with Cre-dependent retrograde AAV, AAVRG-CAG-Flex-TdTomato and AAVRG-hSYN-DIO-GFP in dSTR (100nl) and NAcc (100nl). Representative images (n=5). **c,d.** – quantification of # of cells. Data are represented as mean ± SEM. n.s – not significant. ** $p < 0.01$ Two-way ANOVA.

In both D1Cre and D2Cre mice, distinct projection fibers were observed in midbrain regions i.e substantia nigra pars compacta (SNpc) and reticulata (SNpr). In the D1Cre mice, GFP+ projection fibers were observed in the SNpr, whereas, TdT+ projection fibers were observed in the SNpc (Figure 1, aii). In contrast, for the D2Cre mice only TdT+ projection fibers were observed in the SNpc, and no labeling in the SNpr (Figure 1, bii).

PFC pyramidal neurons also send projections to midbrain regions, specifically to dopamine nuclei (Murugan et al., 2017; Watabe-Uchida et al., 2012). However, whether D1R+ and D2R+ PFC neurons send projections to midbrain dopamine neurons is not known. We injected AAVRGs in dSTR (GFP) as reference, and in SNpc (TdT) of D1Cre or D2Cre mice. For D1Cre mice, similar to Figure 1, injection in the dSTR labeled predominantly distinct Cg1/PrL, VO/LO and M1/M2 localized populations of PFC (Figure 2, ai). However, for SNpc injections we saw minimal labeling of PFC neurons (Figure 2, ai). Quantification of neurons from these regions show a similar predominant labeling of GFP+ neurons in Cg1/PrL as in Figure 1 (Figure 2 b, **$p<0.01$, Two-way ANOVA), but minimal TdT+ (SNpc) labeling. In contrast, for the D2Cre mice we observed minimal GFP+ labeling similar to Figure 1, but we observed robust TdT+ (SNpc) labeling of PFC neurons (Figure 2, bi). TdT+ SNpc labeling was presumably from dopamine neurons since dense TdT+ axonal projections were seen in the dSTR (Figure 2, bii). Quantification of labeled neurons from these regions revealed however, that there was no topographical pattern for the midbrain projecting D2R+ neurons (Figure 2 d, **$p<0.01$, Two-way ANOVA). "D2-SNpc" - Cg1/PrL: 22.7 ± 4.3; MO/IL: 17.0 ± 3.4 VO/LO: 39.6 ± 7.1; M1/M2: 29.4 ± 4.5 neurons. Similar patterns were observed for VTA injections for both D1Cre and D2Cre mice as well (data not shown). Together these data suggest that predominantly D2R+ and not D1R+ PFC neurons project to midbrain dopamine neurons.

Figure 2. <u>**Unique topographically organized subpopulations of D1R+ and D2R+ PFC neurons project to dSTR and SNpc.**</u> PFC sections from D1Cre (**ai**) and D2Cre (**bi**) mice showing cell bodies retrograde labeled with either GFP or TdTomato, respectively. Left panels are zoomed insets of medial PrL and MO showing layer-specific localization of cell bodies. PrL-prelimbic, MO-medial orbitofrontal, VO- ventral orbitofrontal, LO-lateral orbitofrontal, M1-primary motor, M2-secondary motor cortex. **aii, bii.** Injection sites are shown in striatal (STR) and midbrain SNpc/VTA (SN VTA) sections. D1Cre and D2Cre mice were injected with Cre-dependent retrograde AAV, AAVRG-CAG-Flex-TdTomato and AAVRG-hSYN-DIO-GFP in dSTR (100nl) and SNpc (40nl). Representative images (n=5). **c,d.** – quantification of # of cells. Data are represented as mean ± SEM. * p<0.05, ** p<0.01, Two-way ANOVA.

Figure 3. <u>**Comparison of topographical distribution of D1R+ and D2R+ PFC neurons along the rostrocaudal axis**</u>. **a** – Quantification of GFP+ (dSTR) cells in PFC sections along the rostro-caudal axis of D1Cre and D2Cre mice. Right panel shows representative images and stereotaxic coordinates of rostral, rostrocaudal and caudal PFC sections. **b** - Quantification of TdT+ (NAcc core) cells in PFC sections along the rostro-caudal axis of D1Cre and D2Cre mice. **c** - Quantification of TdT+ (SNpc) cells in PFC sections along the rostro-caudal axis of D1Cre and D2Cre mice. PrL-prelimbic, MO-medial orbitofrontal, VO- ventral orbitofrontal, LO-lateral orbitofrontal, AI-agranular insular, M1-primary motor, M2- secondary motor cortex. n=5 for each group. n.s – not significant. Data are represented as mean ± SEM. * $p<0.05$, ** $p<0.01$, Two-way ANOVA.

Figure 3 shows a more detailed comparison of PFC topographical patterns of D1Cre and D2Cre mice for each injection site i.e dSTR, NAcc and SNpc, along the rostro-caudal axis, for the same mice used in Figures 1 and 2. For dSTR (Figure 3a), D1R+ neurons were the predominantly labeled subpopulation across the rostro-caudal axis, in all regions (Cg1/PrL, VO/LO, M1/M2) except in the MO/IL region in rostrocaudal and caudal regions. D2R+ neuron labeling ranged from 1-28% compared to D1R+ neurons for all regions except rostrocaudal and caudal MO/IL. For NAcc (Figure 3b), both D1R+ and D2R+ neurons showed robust labeling across the rostro-caudal axis, with more D2R+ neuron labeling than D1R+ in only the rostral and caudal sections. For SNpc (Figure 3c), D2R+ neurons were predominantly labeled, showing an incremental gradient along the rostro-caudal axis, whereas only 16-20% were D1R+.

Mediolateral Topographical distribution of D1R+ and D2R+ neurons in the PFC.

In the previous experiments the injection sites were along the dorsoventral axis in the dorsocentral striatum and NAcc core. However, within the dorsal striatum, both medial and lateral subregions have specific roles in motivated behaviors. The dorsomedial striatum (DMS) is involved in the acquisition of goal-directed actions, whereas the dorsolateral striatum (DLS) regulates acquisition of habitual behaviors (Yin et al., 2006; Yin et al., 2005). Previous studies however show that mPFC pyramidal neurons primarily project to the DMS whereas more posterior lateral sensorimotor cortex neurons project to the DLS (Kupferschmidt et al., 2017; Shiflett and Balleine, 2011). We next asked the question whether PFC D1R+ and D2R+ neurons have specific projection pattern to the DMS or DLS. Similar to Figures 1 and 2, we injected 50nl of Cre-dependent AAVRG GFP and AAVRG TdT in the DMS and DLS, respectively, of D1Cre and D2Cre mice. As seen in Figure 4, we observe distinct topographical patterns for D1R+ and D2R+ neurons projecting to DMS and DLS. For DMS injection in D1Cre mice, we observed robust GFP+ labeling predominantly in the PrL (67 ± 12.5 neurons), whereas for DLS injection we observed robust TdT+

Figure 4. Topographical organization D1R+ and D2R+ neurons projecting to DMS and DLS. PFC sections from D1Cre (**ai**) and D2Cre (**bi**) mice showing cell bodies retrograde labeled with either GFP or TdTomato, respectively. Left panels are zoomed insets of medial PrL and MO showing layer-specific localization of cell bodies. PrL-prelimbic, MO-medial orbitofrontal, AI – agranular insular, M1-primary motor, M2- secondary motor cortex. **aii, bii.** Injection sites are shown in DMS and DLS subregions in striatal sections. D1Cre and D2Cre mice were injected with Cre-dependent retrograde AAV, AAVRG-CAG-Flex-TdTomato and AAVRG-hSYN-DIO-GFP in DMS (50nl) and DLS (50nl). Representative images (n=5). **c,d.** – quantification of # of cells from D1Cre and D2Cre mice. **e, f** – Comparison of # of labeled cells in PFC of D1 and D2Cre mice projecting to DMS and DLS subregions. Data are represented as mean ± SEM. * p<0.05, ** p<0.01, Two-way ANOVA.

labeling predominantly in M1/M2 (118.3 ± 12 neurons) and AI (146 ± 24.1 neurons) (Figure 4, a and c). Overall, very few D2R+ neurons project to either DMS or DLS (Figure 4, b and d). Comparison of patterns of D1Cre and D2Cre mice for DMS show that only Cg1/PrL has significantly greater labeling of D1R+ neurons (Figure 4e, **p<0.01, Two-way ANOVA). For DLS however, Cg1/PrL, AI and M1/M2 show significantly greater labeling for D1Cre compared to D2Cre mice (Figure 4f, **p<0.01, Two-way ANOVA).

DISCUSSION

In this study, we show a previously unappreciated distinct topographical organization of D1R+ and D2R+ neurons in the PFC of mice. As summarized in Figure 5, we observed distinct topographical organization patterns of D1R+ and D2R+ neurons along the dorsoventral and mediolateral axes. Along the dorsoventral axis, PFC D1R+ neurons are topographically organized such that D1R+ neurons in prelimbic regions primarily project to the dSTR, whereas D1R+ neurons in medial OFC and infralimbic regions primarily project to the NAcc core. In contrast, PFC D2R+ neurons have a distinct pattern of organization, such that very few prelimbic D2R+ neurons project to dSTR, but medial OFC and infralimbic D2R+ neurons project to NAcc core, similar to D1R+ neurons. Along the mediolateral axis, D1R+ neurons in the medial prelimbic region primarily project to dorsomedial striatum, whereas D1R+ neurons in M1/M2 motor and agranular insular cortex primarily project to dorsolateral striatum. In contrast very few D2R+ neurons project to either DMS or DLS. However, medial PFC D2R+ but not D1R+ neurons project to midbrain dopamine nuclei.

Thus, our data provide, for the first time a detailed insight into the organization of D1R+ and D2R+ neurons in the PFC. Our data are consistent with previous findings that D1R+ neurons are primarily corticostriatal, whereas D2R+ neurons are corticostriatal and also project to more

Figure 5. **Summary of topographical organization of D1R+ and D2R+ pyramidal neurons in mouse PFC.** D1R+ neurons represented by black circles and D2R+ neurons are represented by white circles. Circle borders represent projection targets. Size of the circle represent abundance of projections to target regions.

caudal regions such as the thalamus (Gee et al., 2012). Other groups have also shown that PFC D1R+ and D2R+ neurons also project to other limbic areas such as the basolateral amygdala (BLA) (Jenni et al., 2017). Prelimbic neurons projecting to BLA, NAcc or VTA are not only distinct subpopulations but also have distinct laminar distribution (Murugan et al., 2017). In this study we see distinct laminar distribution of both D1R+ and D2R+ PFC neurons. "D1-dSTR" prelimbic neurons are predominantly localized to layer 5, whereas "D1-NAcc" MO/IL neurons are predominantly localized to layer 2/3. Interestingly, "D2-NAcc" MO/IL neurons are predominantly localized to layer 2/3 and 5.

Our data also suggests that distinct predominantly D1R+ neuron subpopulations project to the DMS and DLS. Interestingly, a similar topographical pattern is maintained within the SNpc, where medial and lateral dopamine neurons project to DMS and DLS, respectively (Lerner et al., 2015). dSTR-NAcc injection reveals distinct projection fiber patterns in the SNpr and SNpc. For injections in dSTR, we observed projection fibers in the SNpr only in D1Cre and not D2Cre mice, consistent with AAVRG-GFP labeling striatal D1R+ direct pathway SPNs. However, for NAcc core injections, we observe projection fibers in the medial SNpc and not the SNpr in both D1Cre and D2Cre mice, consistent with previous observations that D1R+ and D2R+ NAcc core SPNs do not follow the traditional direct/indirect dichotomy as dSTR SPNs, and instead send projections to ventral pallidum and SNpc (Kupchik et al., 2015). Thus, D1R+ and D2R+ positive pyramidal neurons in the MO/IL projecting to NAcc core, can not only release glutamate and regulate excitability of GABAergic SPNs, but also modulate dopamine release in the DMS by indirectly acting on medial SNpc dopamine neurons.

The dorsal striatum is topographically divided into the DMS and DLS, which have been implicated in action-outcome learning and stimulus-response learning, respectively, whereas, the NAcc, has been implicated in reward perception (Shiflett and Balleine, 2011). D1R+ and D2R+ neurons in topographically organized regions in the PFC can thus have various effects on behavior depending on their projection target and modulation by dopamine.

Illuminating this unique pattern of organization of D1R+ and D2R+ neurons in the PFC will help us better understand the regional and global effects of cortical dopamine and dopamine receptors in the regulation of motivation and cognition.

METHODS

Animals. All mouse studies were performed according to NIH guidelines for animal care and use,/ and were approved through the University of Florida Animal Care and Use Committee. All mice were housed in a 12h light-dark cycle at a maximum of five per cage, provided with food and water *ad libitum* and tested at 8-12 weeks of age. Mice were age and gender matched, and all experiments were performed in naïve animals. Dopamine D1 receptor Cre (D1Cre) and Dopamine D2 receptor Cre (D2Cre) transgenic mice were obtained from MMRRC. Cre+ hemizygous transgenics were used for all experiments.

Viral Surgeries. D1Cre or D2Cre mice were stereotaxically injected unilaterally with 40-100nl retrograde adeno-associated virus (AAVRG) (Addgene). Stereotaxic coordinates are as follows: dSTR (AP +1.0, ML +1.8, DV -3.25), NAcc core (AP +1.0, ML +1.0, DV -4.75), DMS (AP +1.0, ML +1.2, DV -2750), DLS (AP +1.0, ML +2.2, DV -3.25) and SNpc (AP +3.5, ML +1.25, DV -4.1). Mice were allowed to recover for 2 weeks to allow for viral expression of GFP or TdT before imaging and counting of cells.

Imaging and Quantification. 40 μm thick vibratome cut sections of formalin-fixed mouse brains were processed for imaging. Sections from rostral, rostrocaudal and caudal PFC, and striatal and midbrain sections were imaged using a Nikon AZ100 Zoom microscope, using the same exposure across genotypes for an injection pair. Captured images were used for quantifying number of fluorescent cells for each channel (GFP and Tdt) in different subregions of rostral, rostrocaudal and caudal PFC using ImageJ (NIH), and threshold was kept the same between genotypes and

injection pairs. At least three sections from rostral, rostrocaudal and caudal PFC were analyzed for each mouse, with an n=5 mice per injection pair.

Statistical analyses. Data were analyzed by a standard two-way ANOVA test for comparison between genotypes, and injection pairs. Individual genotypes, or injection pairs were compared using a *post hoc* Tukey's test. Data are presented as mean±SEM. $p < 0.05$ is considered as significant.

REFERENCES

Abi-Dargham, A. (2018). From "bedside" to "bench" and back: A translational approach to studying dopamine dysfunction in schizophrenia. Neurosci Biobehav Rev.

Ahmari, S.E., Spellman, T., Douglass, N.L., Kheirbek, M.A., Simpson, H.B., Deisseroth, K., Gordon, J.A., and Hen, R. (2013). Repeated cortico-striatal stimulation generates persistent OCD-like behavior. Science (New York, NY *340*, 1234-1239.

Balleine, B.W., and O'Doherty, J.P. (2010). Human and rodent homologies in action control: corticostriatal determinants of goal-directed and habitual action. Neuropsychopharmacology *35*, 48-69.

Bamford, N.S., Wightman, R.M., and Sulzer, D. (2018). Dopamine's Effects on Corticostriatal Synapses during Reward-Based Behaviors. Neuron *97*, 494-510.

Barker, J.M., Glen, W.B., Linsenbardt, D.N., Lapish, C.C., and Chandler, L.J. (2017). Habitual Behavior Is Mediated by a Shift in Response-Outcome Encoding by Infralimbic Cortex. eNeuro *4*.

Barker, J.M., Torregrossa, M.M., and Taylor, J.R. (2013). Bidirectional modulation of infralimbic dopamine D1 and D2 receptor activity regulates flexible reward seeking. Front Neurosci *7*, 126.

Burguiere, E., Monteiro, P., Feng, G., and Graybiel, A.M. (2013). Optogenetic stimulation of lateral orbitofronto-striatal pathway suppresses compulsive behaviors. Science (New York, NY *340*, 1243-1246.

Del Arco, A., Mora, F., Mohammed, A.H., and Fuxe, K. (2007). Stimulation of D2 receptors in the prefrontal cortex reduces PCP-induced hyperactivity, acetylcholine release and dopamine metabolism in the nucleus accumbens. J Neural Transm *114*, 185-193.

Ferguson, B.R., and Gao, W.J. (2018). PV Interneurons: Critical Regulators of E/I Balance for Prefrontal Cortex-Dependent Behavior and Psychiatric Disorders. Front Neural Circuits *12*, 37.

Gee, S., Ellwood, I., Patel, T., Luongo, F., Deisseroth, K., and Sohal, V.S. (2012). Synaptic activity unmasks dopamine D2 receptor modulation of a specific class of layer V pyramidal neurons in prefrontal cortex. J Neurosci *32*, 4959-4971.

Goldman-Rakic, P.S. (1998). The cortical dopamine system: role in memory and cognition. Adv Pharmacol *42*, 707-711.

Gremel, C.M., and Costa, R.M. (2013). Orbitofrontal and striatal circuits dynamically encode the shift between goal-directed and habitual actions. Nat Commun *4*, 2264.

Hart, G., Bradfield, L.A., and Balleine, B.W. (2018). Prefrontal Corticostriatal Disconnection Blocks the Acquisition of Goal-Directed Action. J Neurosci *38*, 1311-1322.

Hitchcott, P.K., Quinn, J.J., and Taylor, J.R. (2007). Bidirectional modulation of goal-directed actions by prefrontal cortical dopamine. Cereb Cortex *17*, 2820-2827.

Horvitz, J.C. (2009). Stimulus-response and response-outcome learning mechanisms in the striatum. Behav Brain Res *199*, 129-140.

Ikemoto, S., and Bonci, A. (2014). Neurocircuitry of drug reward. Neuropharmacology *76 Pt B*, 329-341.

Jenni, N.L., Larkin, J.D., and Floresco, S.B. (2017). Prefrontal Dopamine D1 and D2 Receptors Regulate Dissociable Aspects of Decision Making via Distinct Ventral Striatal and Amygdalar Circuits. J Neurosci *37*, 6200-6213.

Killcross, S., and Coutureau, E. (2003). Coordination of actions and habits in the medial prefrontal cortex of rats. Cereb Cortex *13*, 400-408.

Kupchik, Y.M., Brown, R.M., Heinsbroek, J.A., Lobo, M.K., Schwartz, D.J., and Kalivas, P.W. (2015). Coding the direct/indirect pathways by D1 and D2 receptors is not valid for accumbens projections. Nat Neurosci *18*, 1230-1232.

Kupferschmidt, D.A., Juczewski, K., Cui, G., Johnson, K.A., and Lovinger, D.M. (2017). Parallel, but Dissociable, Processing in Discrete Corticostriatal Inputs Encodes Skill Learning. Neuron *96*, 476-489 e475.

Lerner, T.N., Shilyansky, C., Davidson, T.J., Evans, K.E., Beier, K.T., Zalocusky, K.A., Crow, A.K., Malenka, R.C., Luo, L., Tomer, R.*, et al.* (2015). Intact-Brain Analyses Reveal Distinct Information Carried by SNc Dopamine Subcircuits. Cell *162*, 635-647.

Murugan, M., Jang, H.J., Park, M., Miller, E.M., Cox, J., Taliaferro, J.P., Parker, N.F., Bhave, V., Hur, H., Liang, Y.*, et al.* (2017). Combined Social and Spatial Coding in a Descending Projection from the Prefrontal Cortex. Cell *171*, 1663-1677 e1616.

Natsheh, J.Y., and Shiflett, M.W. (2018). Dopaminergic Modulation of Goal-Directed Behavior in a Rodent Model of Attention-Deficit/Hyperactivity Disorder. Front Integr Neurosci *12*, 45.

Ott, T., and Nieder, A. (2019). Dopamine and Cognitive Control in Prefrontal Cortex. Trends Cogn Sci *23*, 213-234.

Palmiter, R.D. (2008). Dopamine signaling in the dorsal striatum is essential for motivated behaviors: lessons from dopamine-deficient mice. Ann N Y Acad Sci *1129*, 35-46.

Robinson, S., Sotak, B.N., During, M.J., and Palmiter, R.D. (2006). Local dopamine production in the dorsal striatum restores goal-directed behavior in dopamine-deficient mice. Behav Neurosci *120*, 196-200.

Rudebeck, P.H., and Murray, E.A. (2011). Dissociable effects of subtotal lesions within the macaque orbital prefrontal cortex on reward-guided behavior. J Neurosci *31*, 10569-10578.

Santana, N., Mengod, G., and Artigas, F. (2009). Quantitative analysis of the expression of dopamine D1 and D2 receptors in pyramidal and GABAergic neurons of the rat prefrontal cortex. Cereb Cortex *19*, 849-860.

Schultz, W. (2002). Getting formal with dopamine and reward. Neuron *36*, 241-263.

Shiflett, M.W., and Balleine, B.W. (2011). Molecular substrates of action control in cortico-striatal circuits. Prog Neurobiol *95*, 1-13.

Smith, K.S., and Graybiel, A.M. (2014). Investigating habits: strategies, technologies and models. Front Behav Neurosci *8*, 39.

Sulzer, D. (2011). How addictive drugs disrupt presynaptic dopamine neurotransmission. Neuron *69*, 628-649.

Tomasella, E., Bechelli, L., Ogando, M.B., Mininni, C., Di Guilmi, M.N., De Fino, F., Zanutto, S., Elgoyhen, A.B., Marin-Burgin, A., and Gelman, D.M. (2018). Deletion of dopamine D2 receptors from parvalbumin interneurons in mouse causes schizophrenia-like phenotypes. Proceedings of the National Academy of Sciences of the United States of America *115*, 3476-3481.

Tseng, K.Y., and O'Donnell, P. (2004). Dopamine-glutamate interactions controlling prefrontal cortical pyramidal cell excitability involve multiple signaling mechanisms. J Neurosci *24*, 5131-5139.

Tseng, K.Y., and O'Donnell, P. (2007). Dopamine modulation of prefrontal cortical interneurons changes during adolescence. Cereb Cortex *17*, 1235-1240.

Urs, N.M., Gee, S.M., Pack, T.F., McCorvy, J.D., Evron, T., Snyder, J.C., Yang, X., Rodriguiz, R.M., Borrelli, E., Wetsel, W.C.*, et al.* (2016). Distinct cortical and striatal actions of a beta-arrestin-biased dopamine D2 receptor ligand reveal unique antipsychotic-like properties. Proceedings of the National Academy of Sciences of the United States of America *113*, E8178-E8186.

Watabe-Uchida, M., Zhu, L., Ogawa, S.K., Vamanrao, A., and Uchida, N. (2012). Whole-brain mapping of direct inputs to midbrain dopamine neurons. Neuron *74*, 858-873.

Yager, L.M., Garcia, A.F., Wunsch, A.M., and Ferguson, S.M. (2015). The ins and outs of the striatum: role in drug addiction. Neuroscience *301*, 529-541.

Yin, H.H., Knowlton, B.J., and Balleine, B.W. (2006). Inactivation of dorsolateral striatum enhances sensitivity to changes in the action-outcome contingency in instrumental conditioning. Behav Brain Res *166*, 189-196.

Yin, H.H., Ostlund, S.B., Knowlton, B.J., and Balleine, B.W. (2005). The role of the dorsomedial striatum in instrumental conditioning. Eur J Neurosci *22*, 513-523.

Zhai, S., Tanimura, A., Graves, S.M., Shen, W., and Surmeier, D.J. (2018). Striatal synapses, circuits, and Parkinson's disease. Curr Opin Neurobiol *48*, 9-16.

a D1Cre

PFC — Injection site — STR — SN VTA

PrL / MO / 2/3 5 6

Cg1, PrL, M2, VO, MO, LO, M1, AI (i)

dSTR, NAcc (STR panel)

VTA, SNpc, SNpr (SN VTA panel) (ii)

b D2Cre

PrL / MO (i)

dSTR, NAcc (STR panel) (ii)

c
D1Cre
dSTR- retro GFP
NAcc- retro TdT

no. of retrograde labeled cells

■ dSTR ■ NAcc

Cg1/PrL, MO/IL, VO/LO, M1/M2

** markers

d
D2Cre
dSTR- retro GFP
NAcc- retro TdT

no. of retrograde labeled cells

■ dSTR ■ NAcc

Cg1/PrL (n.s), MO/IL (**), VO/LO, M1/M2