1

Improved growth and morphological plasticity of *Haloferax volcanii*

Roshali T. de Silva[1], Mohd F. Abdul-Halim[2], Dorothea A. Pittrich[1], Hannah J.

Brown[1], Mechthild Pohlschroder[2] and Iain G. Duggin[1]*

[1]The ithree institute, University of Technology Sydney, New South Wales, 2007,

Australia.

[2]Department of Biology, University of Pennsylvania, Philadelphia, PA 19104, USA.

*Address correspondence to: Iain.Duggin@uts.edu.au

Running title: Morphological plasticity of *H. volcanii*

Abstract

Some microbes display pleomorphism, showing variable cell shapes in a single culture, whereas others differentiate to adapt to changed environmental conditions. The pleomorphic archaeon *Haloferax volcanii* commonly forms discoid-shaped ('plate') cells in culture, but may also be present as rods, and can develop into motile rods in soft agar, or longer filaments in certain biofilms. Here we report improvement of *H. volcanii* growth in both semi-defined and complex media by supplementing with eight trace-element micronutrients. With these supplemented media, transient development of plate cells into uniformly-shaped rods was clearly observed during the early log phase of growth; cells then reverted to plates for the late log and stationary phases. In media prepared with high-purity reagents, without supplemental trace elements, rods and other complex elongated morphologies ('pleomorphic rods') were observed at all growth stages of the culture; the highly-elongated cells sometimes displayed a substantial tubule at one or less frequently both poles, as well as unusual tapered and highly-curved forms. Polar tubules were observed forming by initial mid-cell narrowing or tubulation, causing a dumbbell-like shape, followed by cell division towards one end. Formation of the uniform early-log rods, as well as the pleomorphic rods and tubules were dependent on the function of the tubulin-like cytoskeletal protein, CetZ1. Our results have revealed the remarkable morphological plasticity of *H. volcanii*, and shown that its changes in cell shape occur in response to multiple signals.

Importance

Microbes can show morphological responses to changed environmental conditions, which are important for their survival in a wide variety of environments. These conditions and the specific role of such morphological changes are poorly defined. Here we describe improved growth media for the model archaeon *Haloferax volcanii*, and the identification and characterization of environmental conditions— micronutrient starvation and culture dilution—that induce striking morphological responses. We show that the morphological changes require the tubulin-like cytoskeletal protein, CetZ1. We expect the discovery of these responses and the application of these conditions to facilitate the use of this species in ongoing studies

49   of archaeal cell biology.

Introduction

Archaea exhibit some of the most diverse and unusual microbial cell morphologies. Cell shapes range from rods and cocci to striking triangles and squares. However, little is known about the mechanisms and environmental cues that dictate archaeal cell morphology, or the regulation of transitions between cell morphologies. The model haloarchaeon, *Haloferax volcanii*, when first isolated, was described as mainly disk-shaped cells, with cell shape and size varying significantly (1). In routine liquid cultures of *H. volcanii*, both pleomorphic discoid (plate) and rod cell morphologies may be observed, occasionally with more unusual shapes (2-4).

The conditions that influence *H. volcanii* cell shapes and the relative abundance of the distinct types are not well understood, and specific signals have not been identified. In *H. volcanii* biofilms, substantial elongation (filamentation) has been observed in a subpopulation of cells (5). Furthermore, *H. volcanii* forms rods in swimming-motility soft-agar, such that rods are observed at the forefront of expanding colonies of swimming cells (6). The tubulin-like cytoskeletal protein CetZ1 was required for rod-formation, suggesting a connection between morphology and motility that may be expected based on improved hydrodynamics or directional movement of rods (4, 6). Conversely, the peptidase archaeosortase A (ArtA) and phosphatidylethanolamine biosynthesis enzymes PssA and PssD, which are required for the C-terminal processing and covalent lipid attachment of several *H. volcanii* surface proteins including the surface (S-layer) glycoprotein, are also required for effective and stable plate-shaped cell formation (7, 8).

To better understand the regulation of cell shape switching and the importance of the distinct shapes, we aimed to define conditions that can be robustly used to study these processes. We determined that inclusion of a trace elements (TE) solution in both complex and semi-defined culture media substantially improves growth and culture reproducibility for this species. In the new media, early-log rod formation was also more reproducible, while the pleomorphic rods and other shapes seen in cultures lacking TE were absent. We further defined culture conditions and characterized these CetZ1-dependent morphological changes, which may serve as experimental models for studies of morphological development in archaea.

Results

It was previously shown that, in soft agar assays of cellular swimming motility, *H. volcanii* cells change from a discoid (plate) shape into a rod shape in a CetZ1-dependent manner (6). However, the soft-agar motility assay is not a robust method for ongoing studies of cell-shape differentiation in *H. volcanii* as the soft agar is difficult to separate from the cells, cell yields are low, and the distribution of shapes observed is sensitive to the site of sampling from the soft agar (6). We and others have noticed that *H. volcanii* can form regularly-shaped rods in the early stages of culture, from around the time the culture transitions from lag to log phase, and then cells revert to the irregular discoidal 'plate' cells later in log phase (9). As shown in Fig. 1, cells obtained from colonies grown on agar medium are initially plate-shaped upon resuspension in fresh Hv-Ca liquid medium, but after overnight shaking incubation (to an OD_{600} of 0.05) cells were predominantly rod-shaped. With further incubation, until the culture reached OD_{600} of 0.3, the rod-shaped cells had started to disappear and by 0.9 the culture predominantly contained the plate-shaped cells (Fig. 1). Early culture cells had similar morphology to motile cells withdrawn from motility agar (6), and were recently shown to be motile and suitable for direct observation of swimming (9).

The OD_{600} where rods could be observed and then returned to plates varied between cultures, possibly due to differences in the size of starting inoculum and the times cultures take to reach equal culture density. We have also noticed widely varying growth yields of *H. volcanii* in Hv-Ca medium prepared in different laboratories or with different sources of reagents, and the extent and timing of early-culture rod formation is generally sensitive to different media sources (e.g. we observed that one lower-grade source of NaCl inhibited the early-culture rod development that was seen in all other media tested). The supply of trace elements in the medium was also observed to have dramatic effects on growth and morphology, as described further below. The results shown in Fig. 1 were obtained with Hv-Ca medium without supplemental trace elements, but growth in this culture was satisfactory as cells must have had sufficient supply of essential trace elements in this medium. However, such growth was not always observed with other reagent sources. We therefore sought to

112 understand and standardize the conditions required for optimal growth and rod
113 development; this led to the identification of previously uncharacterized
114 morphological responses of *H. volcanii.*

H. volcanii growth and morphology in nutrient-limited batch culture

The commonly used media for *H. volcanii*, Hv-Ca and Hv-YPC*,* consist of
macronutrients for heterotrophic growth in a base of 18% (w/v) buffered primary sea-
water salts (10). Hv-Ca is a semi-defined medium with Casamino acids as the carbon
and energy source, and typically provides a slower growth rate and yield than the
complex Hv-YPC medium, which contains Yeast extract, Peptone and Casamino
acids added to the 18% salts base. Both media are known to produce different
morphotypes in the same culture—primarily a mix of rods and plate cells, with some
other more unusual shapes also sometimes seen.

Since these media are normally not supplemented with micronutrients, beyond what
may be present as contaminants in the media reagents (10), and a range of metal ions
or other elements are typically required by cells at low or moderate concentrations to
act as essential cofactors in enzymatic reactions, we added a solution of eight
compounds as a source of micronutrients to the Hv-Ca medium. This was based on
common requirements for microorganisms, and contained the elements Fe, Zn, Cu,
Co, Mn, Ni, Mo and B ('trace elements', TE; see Methods). A vitamins solution (10)
was also added. The new medium (named "Hv-Cab", for the 'b' variant of Hv-Ca)
substantially improved the growth rate and yield, and their reproducibility, in *H.*
volcanii cultures, which was due primarily to the inclusion of the TE solution, as
described below.

To identify any growth or morphological changes associated with omission of the
primary nutrient sources in Hv-Cab, mid-log *H. volcanii* cells were transferred to
specific nutrient-omitted media, which were prepared with high-purity reagents to
minimize contaminating sources of trace elements or substances that might inhibit
growth or development. Growth and cell morphology were monitored over three
successive passages (by dilution into fresh media) under each of these nutrient-
omitted conditions. Transfer to media without the carbon and energy source,
Casamino acids (CAA), caused cells to rapidly cease growth, as expected (Fig. 2A,

maroon data-points). Exclusion of the vitamins had a minor effect on the maximum OD_{600} that was consistent throughout the three successive culture passages, but had no effect on cell shape (Fig. 2A-C). However, cultures without supplementary TE showed successively greater reductions in growth rate and yield with each passage, such that by the third passage growth was very poor, and the culture barely reached a maximal OD_{600} of ~0.3 (Fig. 2C).

Cells were observed by phase-contrast microscopy at day 5 (stationary phase) after all three of the culture passages, and the cell shapes were visualized and analyzed by automated measurement of the circularity of cell outlines; these data were then plotted as histograms of cell circularity, expressed as a percentage compared to a perfect circle (Fig. 2D-F). The stationary phase Hv-Cab cultures contained almost exclusively plate-shaped cells that had near-circular cell outlines, similar to late log cultures. However, the TE-depleted cultures showed striking cell elongation, which became more extensive in each successive passage (Fig. 2H). Cells depleted of TE showed complex features accompanying the extensively elongated rod shapes, including rods that showed variable widths along their lengths, and some cells exhibiting substantial polar tubules of variable length and width (Fig. 2H).

Improved growth of H. volcanii in TE-supplemented complex and semi-defined media

Previous studies have shown that in complex medium, such as Hv-YPC (10), elongated and irregular cell morphologies can exist together with the plate-shaped cells (1, 3). Trace elements are expected to be present in Hv-YPC, originating primarily from the Yeast extract. We next investigated the possibility that trace elements may yet be somewhat limiting for *H. volcanii* in Hv-YPC, giving rise to the elongated forms seen in this medium, by supplementing Hv-YPC with the TE solution. By analogy to the Hv-Cab media terminology introduced above, we refer to the TE-supplemented version of Hv-YPC medium as Hv-YPCab.

We initially took mid-log and stationary-phase cells grown in Hv-Cab medium, and then washed and resuspended cells from both starters in Hv-YPC and Hv-YPCab, and then monitored growth in microtiter-plates, to observe how well the cultures grew under these new conditions. The growth rate and yields were noticeably greater in Hv-

YPCab medium compared to Hv-YPC (Fig. 3A). The Hv-YPCab culture grew well

even when started by inoculation with a stationary phase culture, compared to the Hv-

YPC culture inoculated with a mid-log starter, suggesting that the maintenance of

supplemental trace elements in the media is useful for minimizing the lag phase and

maximizing growth rates and yields of *H. volcanii*.

We also monitored growth and morphology of cells in standard tube-based batch-

cultures. Consistent with the results of Fig. 3A, cultures recommenced growth

substantially faster with Hv-YPCab medium, compared to the Hv-YPC medium (Fig.

3B). Similar trends were obtained with experiments based on the Hv-Ca medium (Fig.

3C), although growth was very poor, which was attributed to the lack of trace

elements in this medium made with high-purity reagents. Microscopy of late-stage

culture samples (at 40 h) revealed that cells were almost uniformly the plate

morphotype in Hv-YPCab and Hv-Cab (Fig 3D, '+TE' panels), and showed none of

the elongated rods or filaments with tubules previously noted in non-TE-

supplemented medium (Fig. 3D, -TE panels). The cell size also appeared substantially

smaller in both TE-supplemented media compared to the original media (Fig. 3D). An

additional control culture, supplemented with the EDTA component only showed that

culture growth, cell size and morphology were unaffected compared to Hv-YPC

alone, demonstrating that the cells were reacting specifically to the concentration of

the trace-element compounds that differ between the original and the supplemented

media (Fig 3D). Finally, we also observed a substantial difference in the morphology

when comparing cells in Hv-YPC and Hv-Ca (-TE cultures, Fig. 3D); the more

elongated cells observed in the Hv-Ca culture were attributed to the more severe TE

starvation in this medium (see Fig. 2).

Development of H. volcanii rods during an early stage of growth in culture

Based on the above results, the new supplemented media were expected to improve

consistency and reproducibility of the motile-rod development that occurs during

early-culture. It was expected that the elongated/tubulated rod cell types specifically

seen during TE-limitation would be absent, thus improving uniformity of the rods

seen in early-culture growth. In order to thoroughly characterize the cell morphology

development of *H. volcanii* during the batch-culture growth cycle with the new media,

we inoculated cultures to a fixed cell density from consistently-prepared starter

cultures, and then monitored culture growth rate and cell morphology at regular

timepoints. *H. volcanii* was initially cultured for four days on the appropriate solid

media to allow colony development, and then sufficient colonies were picked and

resuspended in the equivalent liquid medium, and the OD_{600} was adjusted to 0.05 to

start the culture at a measurable and reproducible density. Culture growth (OD_{600} and

cell number), cell volumes (Coulter cytometry) and morphology (microscopy) were

monitored over time, both with or without supplemental TE (Fig. 4).

In both Hv-Cab and Hv-YPCab media, cell volume and culture growth (OD_{600})

increased from the start of the experiments (Fig. 4B, 4D) and the cell number began to

noticeably increase after ~6 h delay (Fig. 4C). This corresponded approximately to the

onset of rod development, as seen in the representative images of the cells (Fig. 4E),

and the quantitative analysis of cell elongation (Fig. 4A); in both media (+TE), a peak

of rod development was observed commencing ~7-8 h after inoculation, reaching a

maximum mean cell elongation after ~10 h, and then from 13 h onwards, plate cells

were again almost exclusively seen (Fig. 4a, +TE). Similar results were obtained with

cultures of the H26 (+ pTA962) strain inoculated from either agar-based colonies or

late-stage liquid cultures (Fig. S1). We also found that early-culture rod development

required inclusion of the plasmid, which is based on the natural pHV2 plasmid from

H. volcanii and complements the uracil-auxotrophy of H26, indicating the importance

of plasmid inclusion or prototrophy in these assays (Fig. S1). Comparison of results

obtained with the different media showed that the peak of rod development was

greater in Hv-YPCab than in Hv-Cab, indicating a higher proportion or longer rods in

Hv-YPCab (compare the left-hand two panels in Fig. 4E, 11 h). The timing of rod

development during the first few generations and subsequent conversion back to

plates approximately corresponded to the timing of an increase (growth) and then a

decrease (division) in the mean cell volume, respectively (Fig. 4B).

The effects of TE-starvation during the above colony-resuspension assay were also

investigated. Cultures were first grown into mid-log phase in either Hv-Cab or Hv-

YPCab medium (*i.e.* +TE), and then they were separated from the medium by

centrifugation, washed (media -TE) and plated onto the respective -TE agar medium

(Hv-Ca and Hv-YPC). After growth on the agar, cells were resuspended in -TE liquid

medium. Cells resuspended from colonies were initially plate-shaped (Fig. 4A, 4E -
TE), even though there was a clear limitation of trace elements in the agar, detected
by very poor growth of colonies on the Hv-Ca agar. As expected, the Hv-Ca (-TE)
liquid cultures showed very poor overall growth, whereas Hv-YPC (-TE) cultures
grew quite well (Fig. 4C, 4D). Cells became noticeably elongated by 5 h (Hv-YPC, -
TE) or 8 h (Hv-Ca, -TE) of incubation, and they continued to show substantial
elongated pleomorphology and increased cell volume for at least 20 h, particularly in
the severe starvation conditions, (Hv-Ca, -TE) (Fig. 4A, 4B, 4E). At longer timepoints

Thus, as may be expected based on the poor growth and marked elongation responses
seen with severe TE starvation (Fig. 2), the transient appearance of the regularly-
shaped rod cell type seen in the optimally-growing cultures (+TE) was not clear
during these -TE assays. This emphasizes the importance of using complete, TE-
supplemented media for obtaining consistent results in the regular rod development
assays, and offers a standardized way of performing TE-starvation when those cell
types are required.

Reversibility and specificity of the cell elongation response to TE-starvation

To investigate the reversibility of the TE-starvation phenotype in the Hv-Ca cultures,
TEs were added at 15 h incubation after colony resuspension either by direct addition
of the TE stock solution, or by dilution of the Hv-Ca culture (1:100) into Hv-Cab, Hv-
Cab with 10x concentrated TE, or Hv-Ca (-TE) as a control. In all three media
conditions containing additional TE, the cultures reverted to almost exclusively the
plate cell morphotype at 12 h (Fig. 5). This is a similar duration required for the
appearance of elongated cells upon colony resuspension in Hv-Ca (Fig. 4A, -TE), and
shows that the effect of TE-starvation on morphology is readily reversible by resupply
of the trace element nutrients. It is concluded that *H. volcanii* undergoes substantial
cell elongation (and increased tubule development) in response to trace-element
starvation, but not during starvation associated with stationary phase (+ TE) (Fig. 2-5,
S2).

The potential effects of individual components of the TE solution on *H. volcanii* cell
shape were screened by testing a series of five TE solutions omitting either the Co,
Cu, Fe, Mn or Zn. None of these individual 'drop-out' TE media matched the degree

of cell elongation observed when all TE were omitted (Fig. S3), suggesting that the

combined effect of multiple depleted elements causes the large change in cell shape

seen in the -TE conditions. The Co and Fe drop-out media appeared to cause the most

noticeable (although moderate) cell elongation response compared to the other metal

drop-out media (Fig. S3).

Microfluidic flow-chamber visualization of H. volcanii cellular development

We next sought to directly visualize morphological development during the early log

growth phase in liquid culture. Cells resuspended from colonies were loaded into

incubated microfluidic chambers for time-lapse microscopy. Time-series of rod-cell

development in the media (+TE) are shown in Fig. 6A and 6B, and the complete time-

series are available in Movies S1 and S4. These data show that plate cells can initially

grow and divide as plates, and then most cells transition into rods as they grow and

divide over several generations. We observed that rods persisted during incubation in

the microfluidic culture system (particularly evident in Hv-YPCab), compared to the

temporally quite narrow peak of rods seen in batch culture (Fig. 4A). The continual

flow of fresh medium while cells are attached to the flow channels may therefore

create signals that maintain rod morphology.

Microfluidic cultures with media lacking the trace elements revealed striking

morphological responses (Fig. 5C, 5D, 5E, and Movies S2, S3 and S5), including

some extensively elongated cells exhibiting highly asymmetric division (Fig. 5C, 5E),

and the generation of some bizarre irregular shapes (Fig 5D). This supported the batch

culture results suggesting that formation of regularly shaped rods in early log phase is

promoted in optimal growth conditions (in the presence of TE), whereas irregularly

shaped and elongated shapes are associated with growth in the absence of TE and

hence suboptimal growth conditions. Interestingly, tubules appeared to be generated

by a combination of localized narrowing and elongation (tubulation), often in the mid-

cell region of a cell, creating an approximate dumbbell-like appearance, followed by

division near one end of the narrowing. We also saw that occasionally the narrow

zone can re-widen, in a peristaltic-like wave along the cell (Movie S5).

Requirement of tubulin-like protein CetZ1 for elongated shapes and tubule
formation

CetZ is a family of cytoskeletal proteins from the tubulin superfamily found in the
Euryarchaeota. In *H. volcanii*, the CetZ1 protein was found to be required for rod-cell
development when cells become motile (4). The robust growth accompanying the
morphological changes observed above suggests that the shape changes are biological
responses aimed at helping *H. volcanii* adapt to the conditions. In order to confirm
that the morphological changes are of biological origins and not direct abiotic effects,
we investigated the role of CetZ1 in the cell shape changes. A CetZ1-knock-out strain
was subjected to the assays defined in Fig. 4, to determine if rod-shape or tubule
development were affected under these conditions. In early-culture rod development
assays, the Δ*cetZ1* strain failed to form rods at 10 h post-resuspension in Hv-Cab or
Hv-YPCab, compared to clear rod formation in the wild-type control (Fig. 7A).
Furthermore, both the elongated rods and the tubules observed in wild-type cells in
the -TE conditions (Fig. 4) were not evident in Δ*cetZ1* cells (Fig. 7A). We also used
Coulter cytometry to assess cell volumes of wild-type and CetZ1 knock-out strains.
They showed very similar cell-size distributions in both media (15 h post-
resuspension; Fig. 7C), despite the different shapes (Fig. 7A). However, significantly,
both strains showed larger cells on average in Hv-Ca (-TE) than in Hv-Cab,
demonstrating that cell size (regulation of cell division) also responds trace-element
starvation.

We confirmed the expected absence of CetZ1 protein in the knock-out strain by
Western blotting (Fig. 7B) and observed no substantial difference in the CetZ1
detected in the presence or absence of supplementary TE. To verify that CetZ1
function is necessary for the rod and tubule formation, we expressed a dominant-
inhibitory mutant (CetZ1.E218A) from a plasmid in a wild-type genetic background
and then observed cell shapes. The E218A mutation in the GTPase active-site
strongly inhibits CetZ1 function via hyper-stabilization of localized polymers, and
was used previously for verification of CetZ1's involvement in rod development in
motility soft-agar (4). We found that expression of *cetZ1*.E218A blocked rod cell and
tubule formation in the wild-type strain background (Fig. 7A), confirming its
dominant influence on cell shape and the requirement for functional CetZ1 in trace-
element-induced rod and tubule generation.

Discussion

We report conditions for improved growth of *H. volcanii* and the identification and characterization of cell morphological responses, with liquid culture time-course methods for observing them. The rod cells that develop during the early stages of growth in liquid culture (Fig. 1) are regularly-shaped and appear similar to the motile rods taken from soft agar (6). Indeed, early culture cells were very recently shown to be motile and suitable for direct observation of swimming (9). Secondly, we observed highly elongated, pleomorphic rod cells forming upon starvation of trace-element nutrients in liquid culture, but not in stationary phase starvation when trace elements were added (Fig. 2, S). Many of the cells lacking trace elements displayed a striking tubular extension at one or both cell poles, in many cases formed by a mid-cell narrowing or tubulation followed by division at one end (Fig. 6). We further showed that commonly-used semi-defined and complex media, which do not include supplementary trace elements, can limit growth, and that trace-element limitation is a significant cause of morphological variation in such cultures (Fig. 3-5). Finally, we showed that rod and other elongated cell morphologies were dependent on the function of the CetZ1 cytoskeletal protein under all conditions investigated (Fig. 7). The variety of dynamic cell shapes observed in these conditions underscores the remarkable morphological plasticity of this archaeon.

Our preliminary observations had shown that the sources of the reagents used for preparing *H. volcanii* media substantially and unpredictably influenced growth and cell morphology. The commonly-used *H. volcanii* media, Hv-Ca and Hv-YPC, were found to be limited in their supply of trace-elements (Fig. 2-4). When high-purity reagents are used, as in Fig. 2-7 of this study, Hv-Ca medium in particular gives very poor growth due to the very low concentration of essential trace metals that must come from contaminants in the media reagents or water (Fig. 2). We had also noticed that the use of one lower-grade source of NaCl inhibited the early-culture rod development seen in all other media tested. These issues make reproducing certain experiments difficult between laboratories, when using different reagents. Such considerations are pertinent when nutrition, metabolism, stress-response or whole-organism ('omic') projects are planned.

We introduced two media, named Hv-Cab (semi-defined) and Hv-YPCab (complex),

which are supplemented with a solution containing eight trace elements that enhance

the reproducibility and growth performance (rate and yield) of *H. volcanii* cultivation

(Fig. 2 and Fig. 3). The new media also produce much more uniform cell

morphologies in routine culture, in contrast to the variable, mixed morphotype

cultures obtained with the original Hv-Ca and Hv-YPC media (3, 10). The mixed

morphotypes of *H. volcanii* that have been commonly seen in the late log and

stationary phases of routine laboratory cultures are therefore primarily a response to

trace-nutrient limitation.

Our surprising observation that TE-supplementation can improve growth and shape

uniformity even in the complex medium (based on Hv-YPC, which contains yeast

extract) made with high-purity salt, buffer and water reagents suggests that *H.*

volcanii requires relatively high levels of certain 'trace' metals for optimal growth.

Previous *H. volcanii* media have been prepared with a trace elements solution

containing Mn, Zn, Cu and Fe (11), but these are unlikely to represent all the minor

elements utilized by *H. volcanii* (Fig. 3 and Fig. S3), and the four-element solution

has only been widely adopted for use in minimal medium (10). More recent reports

using trace-elements supplementation in *H. volcanii* media have used six (12) of the

elements used in the current study, or all eight with synthetic medium (13), although

in differing proportions, and these studies did not report their influence on growth and

morphology. Use of the eight-elements solution should avoid or minimize metal-

starvation responses in *H. volcanii* cultivation in other media, including rich media,

and it greatly improves culture reproducibility.

Relatively high metal ion concentrations might be expected in hypersaline lakes,

including the Dead Sea where *H. volcanii* DS2 was originally isolated, although some

metals such as Fe can still be present at only moderate concentrations in relation to

biological requirements (1, 14). The TE-solution used here is expected to provide

similar availabilities of important metal nutrients compared to relevant natural

environments, although it should be noted that the ion ratios vary significantly

amongst different salt-lake habitats of *H. volcanii*. For example, Mn is in especially

high abundance (~100-150 μM) in the Dead Sea (14). The TE-solution used here has

a similar Mn concentration (130 μM). We observed that individually omitting

elements from the TE-solution had only moderate effects on cell shape (Fig. S2),
suggesting that the large effect of omission of the whole TE-solution would be the
cumulative effect of limiting several of these elements.

Why would trace-element limitation trigger the development of *H. volcanii* elongation
or tubules? Some of the factors that are related to morphological changes in bacteria
are nutrients, extracellular signalling, cell attachment and biofilm formation,
motility/chemotaxis, predation and polar differentiation (15, 16). For example,
starvation of multiple nutrients causes cell shape changes in a marine *Vibrio* species,
resulting in pleomorphic cell populations that stayed viable for extended periods,
suggesting that the morphological heterogeneity might offer a survival advantage
(17). In *Caulobacter crescentus*, phosphate starvation triggers stalk extension, which
is thought to improve surface area for nutrient uptake (18, 19). The reason for *H.*
volcanii morphological differentiation in response to trace element starvation is not
yet clear. The observation that cells also get larger (Fig. 4B) indicates a change in the
regulation of cell size or division too, which argues against solely a nutrient-uptake
function based on improved surface area-to-volume ratio.

The present study also characterized the timing and conditions for *H. volcanii* rod
development that occurs during the early stages of growth in dilute liquid cultures
under nutritionally-replete culture conditions (*i.e.* including supplemental TE, Fig. 4).
In bacteria, gene expression profiles in the lag- and early-log phases show that these
are dynamic stages of culture, sensitive to the history of the starting culture (20).
Consistent with this, we have found that it is important that *H. volcanii* are pre-
cultured in a consistent way when performing the time-course assays for rod
development. *H. volcanii* cell shape changes can be triggered in response to moderate
changes in the intracellular or extracellular environment. For example, we found that
early-culture rod formation in the high-purity reagent medium, requires the presence
of an *H. volcanii* plasmid, pTA962 (Fig. S1), which contains the endogenous pHV2
replication functions, and marker genes (*pyrE2*, *hdrB*) that complement auxotrophic
strains. A previous study noted that the presence of pTA963 increased the appearance
of rods (7), and we have also observed a slight decrease in cell size (unpublished
data). While this plasmid does not appear to significantly affect growth rate (21), the
results suggest that *H. volcanii* cell size and shape is influenced by auxotrophic status,

plasmid replication or other functions, or their impacts on metabolism. Other
responses, including the development of motility in soft agar, appear to be less
sensitive to these effects (6).

The rod development that occurs after resuspension or dilution of *H. volcanii* in fresh
liquid medium is consistent with the sensing of nutrients or diluted biotic signals such
as secreted quorum-sensing molecules. Consistent with this, we also observed cells
retaining their rod shapes for extended periods when in the microfluidic culture
system (Fig. 6 and Movie S4), in which cells are exposed to a continuous flow of
fresh medium. This would continuously remove biotic signals from the culture,
however cells might also sense surface attachment and shear forces that differ
between the flow chambers and liquid batch cultures. We also observed that cell
shapes in agar surface colonies differed greatly from liquid cultures made with
otherwise the same medium, both without trace elements (Fig. 4). This suggested that
the morphological effect of growth on agar (producing the plate morphotype) is
dominant over the influence of trace element limitation, and could reflect sensing of
different chemical or physical signals in colonies. *H. volcanii* is capable of quorum
sensing (22) and mechanosensing (23), so further studies could ascertain how these
processes might trigger changes in cell shape.

Materials and Methods

Archaeal strains

Haloferax volcanii H26 (Δ*pyrE2*), H53 (Δ*pyrE2* Δ*trpA*) or H98 (Δ*pyrE2* Δ*hdrB*) (7), containing the pHV2-based plasmids pTA962 or pTA963 (21) as indicated, were used as wild-type strains. *H. volcanii* ID59 (6), transformed with pTA962, was the *cetZ1* knockout strain used (based on H98). Attempts to verify the roles of CetZ1 by plasmid-based expression of wild-type *cetZ1* in the Δ*cetZ1* strain (ID59) were unsuccessful, likely due to polar effects on a downstream gene, so we expressed *cetZ1.E218A* (a dominant-inhibitory mutant) to confirm the involvement of *cetZ1*, by using H98 + pTA962-*cetZ1.E218A* (6).

Culture media and general growth conditions

Media were based on the Hv-Ca and Hv-YPC media commonly used for *H. volcanii* (10). In this study, we used defined sources of all media reagents, which we found to be important for reproducibility. However, the data reporting the initial observation of early-culture rod development (Fig. 1) were obtained with the original recipe for Hv-Ca (10). For solid media, Bacteriological Agar (Oxoid LP0011) (10 g/L), was dissolved in the volume of water required for the medium by heating in a microwave oven, before mixing with pre-heated concentrated stock solutions of macronutrients and buffered salt water solutions (listed below) and autoclaving, followed by addition of trace elements or heat-sensitive reagents. Water was obtained from an ultrapure (18.2 MOhm.cm) water purification system (Sartorius).

Hv-Ca medium was prepared similarly to that described previously (10), however, in all experiments except Fig. 1, Hv-Ca medium also contained an additional vitamins solution. Hv-Ca contained: 144 g/L NaCl (Sigma S6191), 21 g/L $MgSO_4.7H_2O$ (Sigma M1880), 18 g/L $MgCl_2.6H_2O$ (Sigma M2393), 4.2 g/L KCl (Sigma P5405), 12 mM Tris-HCl (pH 7.4) (Sigma RDD008, buffered with AR grade HCl), 3 mM $CaCl_2$ (Sigma C5670), and 5 g/L Casamino Acids (Oxoid LP0041, prepared initially as a 10x stock solution). The medium was then autoclaved. Once cool, a filter-sterilized vitamins concentrated stock solution (1000x), containing 1 g/L thiamine (Sigma T1270) and 0.1 g/L biotin (Sigma B4639), was added (1 mL per L of medium). Buffered salt water (BSW) was prepared as the buffered salts solution

described above, i.e. Hv-Ca medium (18 % w/v total salts), omitting the Casamino acids and vitamins. BSW was prepared as a concentrated (30% w/v salts) stock solution.

To prepare Hv-Cab medium, a trace-elements concentrated stock solution (100x) was added (10 mL per L) to the Hv-Ca medium after autoclaving and cooling (including vitamins). The 100x trace-elements stock solution contained: 5 g/L $Na_2.EDTA.2H_2O$ (Sigma E1644), 0.8 g/L $FeCl_3$ (Sigma 157740), 0.05 g/L $ZnCl_2$ (Sigma 793523), 0.01 g/L $CuCl_2$ (Sigma 751944), 0.01 g/L $CoCl_2$ (Sigma 232696), 0.01 g/L H_3BO_3 (Sigma B6768), 1.6 g/L $MnCl_2$ (Sigma 328146), 0.01 g/L $NiSO_4$ (Sigma 656895), and 0.01 g/L $Na_2MoO_4.2H_2O$ (Sigma M1003); the pH of the solution was adjusted to 7.0 with NaOH (AR grade, Sigma), and then the obvious brown precipitate that formed on neutralization was removed by 0.2 μm sterile filtration; stock solutions were stored at room temperature in sterile aliquots.

Hv-YPC medium (10) was prepared as per the Hv-Ca medium above, except the supplementary vitamins were omitted, the Casamino acids was used at a concentration of 1 g/L, and 5 g/L Yeast Extract (Oxoid LP0021) and 1 g/L Peptone (Oxoid LP0037) were included (initially dissolved as a 10x stock solution). To prepare "Hv-YPCab" medium, the 100x trace-elements stock solution was added (10 mL per L of medium) to Hv-YPC after autoclaving and cooling.

Most cultures (5 mL) were grown in sterile 50 mL Falcon tubes, incubated at 45°C with shaking at 200 rpm in a GFL 1092 rotary-shaking water bath (the tube's lid was maintained loosened) unless otherwise indicated. *H. volcanii* was initially grown on solid medium followed by continuous growth in liquid medium (diluted once per day), to maintain steady logarithmic growth prior to sampling, which was only done after the culture had been growing steadily in continuous log-phase for at least 2 days, unless otherwise indicated.

Time-course studies of rod development

In order to obtain a standardized way of observing rod cells, and analyzing rod-development, time-course studies of the early stages of liquid culture growth were carried out (Fig. 4). The *H. volcanii* strain was streaked onto an agar plate made from the same medium as chosen for the assay (*e.g.* Hv-YPCab), and then the plate was

incubated in sealed humidified bag at 45°C for 4 days. Colonies were collected from

the surface with a microbiological loop, and resuspended in 5 mL of liquid medium,

sufficient to give an $OD_{600} > 0.05$. The OD_{600} was immediately adjusted to 0.05 by

dilution with fresh pre-warmed medium, marking the start of the time-course.

Samples were withdrawn as required for microscopy or other analyses (Fig. 4). In

some experiments, cells were obtained directly from a mid-log liquid culture (Fig.

S1B).

To observe trace element limitation over a time-course (Fig. 4), cultures were first

grown into steady log phase in Hv-Cab medium at 45°C, with shaking at 200 rpm (5

mL culture). A sample of the culture was withdrawn and centrifuged at 5,000 rpm for

5 min in an Eppendorf microcentrifuge. The supernatant was discarded and the cells

were then washed by re-suspending the pellet in 1 mL of 18% BSW. The wash was

repeated three times and then the cells were finally re-suspended in 0.1 mL 18% BSW

and then spread onto Hv-Ca agar plate(s). The plate(s) were placed into a plastic bag

and incubated at 45°C for 4 days. 5 mL of liquid Hv-Ca was added to the plate

surface and the cells were re-suspended, avoiding disturbing the agar. The OD_{600} was

adjusted to 0.05 with Hv-Ca medium, to start the time course (incubation was

continued at 45°C at 200 rpm).

Microscopy

Samples were prepared by mounting a 2 μl volume of culture (concentrated where

necessary by centrifugation and gentle resuspension) onto a slide prepared with a

~170 μm thick, 1% agarose pad containing 18% BSW (6). Most phase-contrast

images were acquired with a Zeiss AxioPlan2, Nikon Ti, or a V3 Delta Vision Elite

(Applied Precision Inc., a GE Healthcare Company, Issaquah, USA) microscope, with

1.4 NA phase-contrast objectives. Images were analyzed using the MicrobeJ plug-in

for ImageJ (24). The cell circularity ($4\pi \times$ area / perimeter^2) was measured on the

interpolated contour generated using the particle medial axis and the specified value

of the width measured along the medial axis. Circularity values range between 0 and

1. Cell elongation was calculated as the inverse of the circularity.

For live cell imaging, a CellASIC ONIX microfluidics system (with B04A-03 plates),

was used with a Nikon Ti inverted microscope equipped with an incubated stage

(42°C). The four flow chambers were initially washed with 1 mg/mL Bovine Serum

Albumin (BSA) in phosphate-buffered saline followed by rinsing with 18% BSW at 5

psi for 5 min. Cells of *H. volcanii* (H98 + pTA962) cultured freshly cultured on the

agar media indicated with each movie were resuspended in the respective liquid

medium (pre-warmed to 42°C), and then loaded into the chambers as per the

manufacturer's instructions, and the chambers were perfused with medium at 2 psi for

20 h, with time-lapse images recorded at 10 min intervals.

Coulter cytometry

Culture samples were diluted (1:1000) with 18% BSW, and then analyzed with a

Multisizer 4 Coulter cytometer (Beckman-Coulter) as described previously (6).

Western blotting

Western blotting was carried out as described previously (6).

Author contribution statement

Designed research: RTdS, MP, IGD. Contributed data: MFAH, MP (Fig. 1), RTdS

(Fig. 2-7, S2-3, Movies S1-S5), DAP, HJB (Fig. S1). Interpreted data: RTdS, MP,

DAP, IGD. Wrote the manuscript: RTdS, IGD.

Acknowledgements

This work was supported by the Australian Research Council (FT160100010 and

DP160101076). M.P. and M.F.A.-H. were supported by the National Science

Foundation grant 1817518. We are grateful for use of the equipment and technical

support from the UTS Microbial Imaging Facility, and Solenne Ithurbide and Stephen

Vadia for comments on the manuscript.

References

1.   Mullakhanbhai MF, Larsen H. 1975. *Halobacterium volcanii spec. nov.*, a Dead Sea halobacterium with a moderate salt requirement. Archives of microbiology 104:207-14.

2.   Bisson-Filho AW, Zheng J, Garner E. 2018. Archaeal imaging: leading the hunt for new discoveries. Mol Biol Cell 29:1675-1681.

3.   Delmas S, Duggin IG, Allers T. 2013. DNA damage induces nucleoid compaction via the Mre11-Rad50 complex in the archaeon *Haloferax volcanii*. Molecular Microbiology 87:168-79.

4.   Liao Y, Ithurbide S, de Silva RT, Erdmann S, Duggin IG. 2018. Archaeal cell biology: diverse functions of tubulin-like cytoskeletal proteins at the cell envelope. Emerging Topics in Life Sciences 2:547-559.

5.   Chimileski S, Franklin MJ, Papke RT. 2014. Biofilms formed by the archaeon Haloferax volcanii exhibit cellular differentiation and social motility, and facilitate horizontal gene transfer. BMC Biol 12:65.

6.   Duggin IG, Aylett CH, Walsh JC, Michie KA, Wang Q, Turnbull L, Dawson EM, Harry EJ, Whitchurch CB, Amos LA, Lowe J. 2015. CetZ tubulin-like proteins control archaeal cell shape. Nature 519:362-5.

7.   Abdul Halim MF, Karch KR, Zhou Y, Haft DH, Garcia BA, Pohlschroder M. 2015. Permuting the PGF Signature Motif Blocks both Archaeosortase-Dependent C-Terminal Cleavage and Prenyl Lipid Attachment for the *Haloferax volcanii* S-Layer Glycoprotein. J Bacteriol 198:808-15.

8.   Abdul-Halim MF, Schulze S, DiLucido A, Pfeiffer F, Bisson Filho AW, Pohlschroder M. 2020. Lipid Anchoring of Archaeosortase Substrates and Midcell Growth in Haloarchaea. mBio 11.

9.   Li Z, Kinosita Y, Rodriguez-Franco M, Nußbaum P, Braun F, Delpech F, Quax TEF, Albers S-V. 2019. Positioning of the Motility Machinery in Halophilic Archaea. mBio 10:e00377-19.

10.   Allers T, Ngo HP, Mevarech M, Lloyd RG. 2004. Development of additional selectable markers for the halophilic archaeon *Haloferax volcanii* based on the *leuB* and *trpA* genes. Applied and environmental microbiology 70:943-53.

11.   Cline SW, Schalkwyk LC, Doolittle WF. 1989. Transformation of the archaebacterium *Halobacterium volcanii* with genomic DNA. J Bacteriol 171:4987-91.

12.   Hattori T, Shiba H, Ashiki K, Araki T, Nagashima YK, Yoshimatsu K, Fujiwara T. 2016. Anaerobic Growth of Haloarchaeon *Haloferax volcanii* by Denitrification Is Controlled by the Transcription Regulator NarO. J Bacteriol 198:1077-86.

13.   Maurer S, Ludt K, Soppa J. 2018. Characterization of Copy Number Control of Two Haloferax volcanii Replication Origins Using Deletion Mutants and Haloarchaeal Artificial Chromosomes. J Bacteriol 200.

14.   Nissenbaum A. 1977. Minor and trace elements in Dead sea water. Chemical Geology 19:99-111.

15.   Young KD. 2007. Bacterial morphology: why have different shapes? Current opinion in microbiology 10:596-600.

16.   Justice SS, Hunstad DA, Cegelski L, Hultgren SJ. 2008. Morphological
606   plasticity as a bacterial survival strategy. Nature reviews Microbiology
607   6:162-8.

608   17.   Stretton S, Danon SJ, Kjelleberg S, Goodman AE. 1997. Changes in cell
609   morphology and motility in the marine *Vibrio sp.* strain S14 during
610   conditions of starvation and recovery. FEMS Microbiology Letters 146:23-
611   29.

612   18.   Klein EA, Schlimpert S, Hughes V, Brun YV, Thanbichler M, Gitai Z. 2013.
613   Physiological role of stalk lengthening in *Caulobacter crescentus*. Commun
614   Integr Biol 6:e24561.

615   19.   Wagner JK, Setayeshgar S, Sharon LA, Reilly JP, Brun YV. 2006. A nutrient
616   uptake role for bacterial cell envelope extensions. Proc Natl Acad Sci U S A
617   103:11772-7.

618   20.   Pin C, Rolfe MD, Munoz-Cuevas M, Hinton JC, Peck MW, Walton NJ,
619   Baranyi J. 2009. Network analysis of the transcriptional pattern of young
620   and old cells of *Escherichia coli* during lag phase. BMC Syst Biol 3:108.

621   21.   Allers T, Barak S, Liddell S, Wardell K, Mevarech M. 2010. Improved
622   strains and plasmid vectors for conditional overexpression of His-tagged
623   proteins in *Haloferax volcanii*. Applied and environmental microbiology
624   76:1759-69.

625   22.   Megaw J, Gilmore BF. 2017. Archaeal Persisters: Persister Cell Formation
626   as a Stress Response in *Haloferax volcanii*. Front Microbiol 8:1589.

627   23.   Le Dain AC, Saint N, Kloda A, Ghazi A, Martinac B. 1998. Mechanosensitive
628   ion channels of the archaeon *Haloferax volcanii*. J Biol Chem 273:12116-9.

629   24.   Ducret A, Quardokus EM, Brun YV. 2016. MicrobeJ, a tool for high
630   throughput bacterial cell detection and quantitative analysis. Nat
631   Microbiol 1:16077.

 Figure Legends

Figure 1. *H. volcanii* forms rods during the early stages of growth at low culture
density in liquid batch culture. A fresh colony of *H. volcanii* (H53 + pTA963) was
used to inoculate Hv-Ca liquid medium (without additional vitamins or trace
elements, but prepared with standard-purity reagents as the only possible source of
trace elements). Cells from early-, mid- and late-log phase cultures (OD_{600} 0.05, 0.3,
and 0.9, respectively) were observed by phase-contrast microscopy. Scale bars are 10
μm (inset = 2.5 μm).

Figure 2. Effects of nutrient depletion on *H. volcanii* growth and morphology in
liquid culture. Strain H98 + pTA962 was initially grown to mid-log phase in Hv-Cab
(OD_{600} = 0.5), and then was washed 3 times by centrifugation at 5,000 rpm in a
microcentrifuge for 5 min, followed by resuspension of the cell pellet in same volume
of 18% BSW, and diluted (1:100) into fresh Hv-Cab or in media excluded for one of
the major nutrient components, *i.e.* trace elements (TE, red data-points), vitamins
(biotin and thiamine, purple triangles), or Casamino acids (CAA, maroon triangles).
(A) Culture growth was monitored by measuring the OD_{600} over 5-days. Cells from
day-5 of the first dilution were then diluted (1:100) into the respective nutrient-
depleted media and incubated and monitored the same way for a further 5 days **(B)**.
Cells from the second dilution (day-5) were again diluted (1:100) into the respective
nutrient-depleted media and then incubated for five more days **(C)**. Cells imaged by
phase-contrast microscopy at day-5 (final data point) in each of the cultures (A-C) by
determining the circularity of cell outlines (N = 100, randomly selected cells), shown
as a corresponding histogram (D-F). G and H shows the phase-contrast images of
cells in (G) Hv-Cab (+TE) and (H) Hv-Ca (-TE) on day-5 of the third passage (a
representative image of the cells analysed in histogram F). TE depletion also resulted
in formation of polar tubules in some cells. The scale bars are 5 μm.

Figure 3. Media supplementation with a trace elements (TE) solution improves
growth and cell shape uniformity in *H. volcanii*. (A) Growth curves in microtiter
plate format. Two *H. volcanii* (H98 + pTA962) Hv-Cab batch cultures (5 mL) were
started from separate colonies. The cells were taken at the mid-log phase (for 6
technical replicates) of one culture and at the stationary phase (for 3 technical

replicates) of the other. Both were incubated at 45$^\text{o}$C and then washed with 18% BSW

three times before inoculating to an OD$_{600}$ of 0.1 the indicated Hv-YPC based media

with or without TEs as indicated. Growth at 42°C with 200 rpm shaking was then

monitored in a Tecan Spark microtiter plate spectrophotometer (30 min reading time

intervals). The error bars indicate the standard deviation. **(B)** Growth curves in tube-

culture format, started from steady mid-log cultures in the respective media,

comparing Hv-YPC medium with the same media supplemented with the complete

TE solution (*i.e.* Hv-YPCab), or with only the EDTA component of the solution

(control). **(C)** Tube-culture experiments, as per panel (B) except based on Hv-Ca

medium. The Hv-Ca culture was started at ~10-fold lower cell density than the other

two, owing to the relatively poor growth of the starter culture. **(D)** Cells sampled after

40 h (stationary phase) were imaged by phase-contrast microscopy. All scale bars

represent 5 μm.

Figure 4. Assays for *H. volcanii* reversible morphological transitions during the

growth cycle. Samples of *H. volcanii* (H98 + pTA962), grown in the indicated liquid

media (after inoculation from colonies obtained from 4 days of growth on the

equivalent solid agar medium), were withdrawn at time-points for microscopy (A)(E),

Coulter cytometry determination of cell volumes (B) and counts (C), and OD$_{600}$

measurements (D). The microscopy analysis in panel (A) shows mean cell elongation

(expressed as inverse circularity; a value of 1 is a perfect circle and becomes greater

as cells become more elongated) of 250 randomly selected cells at each time point.

Datapoints shown represent the mean from two independent replicates, and error bars

indicate standard deviation of this mean (some are within the datapoints).

Figure 5. Reversal and specificity TE starvation. *H. volcanii* (H98 + pTA962)

colonies from Hv-Ca agar were resuspended in Hv-Ca liquid medium. After 15 h of

incubation, the culture was treated as indicated in the key, and then each new culture

was monitored by sampling at 3 h intervals for microscopy over 12 hours. (A) The

mean cell circularity (sample size N = 150 at each time point) versus time. (B)

Representative phase-contrast image of the cells from the Hv-Ca (-TE) culture at 15 h,

immediately prior to introducing TE. (C) Images containing cells after 4 h, 8 h and 12

h incubation under the conditions given in the key. Scale bars represent 5 microns.

Figure 6. Time-lapse microscopy of *H. volcanii* (H98 + pTA962) with and without
trace elements. Microfluidic chambers were loaded with cells and phase-contrast
images were collected at 10 min intervals during perfusion (2 psi for 20 h). (A)
Selected time frames showing initial growth after resuspension in Hv-Cab (+TE). (B)
Selected time frames during microfluidic culture in Hv-YPCab (+TE). (C) Selected
time frames showing the substantial cell elongation observed Hv-Ca (-TE) and a
highly asymmetric division event. (D) Another field of cells cultured in Hv-Ca (-TE)
showing slow growth and striking morphologies, including the generation of a cells
that shows a distinctive highly curved narrowing or tubule. (E) Selected images
during cell elongation in Hv-YPC (-TE).

Figure 7. CetZ1 is essential for rod development during early-growth after
colony resuspension and during TE-depletion. (A) Phase-contrast images of wild-
type (H98 + pTA962), the CetZ1 knock-out (H98.*ΔcetZ1* (ID59) + pTA962), and the
point-mutant (H98 + pTA962-*cetZ1*.E218A) were recorded 10 h post-resuspension of
colonies in the indicated media (including 1 mM L-tryptophan, as the inducer for the
P.tna promoter in pTA962). In *ΔcetZ1*, or during expression of CetZ1.E218A (-TE
conditions), cells failed to form rods or substantially elongate, and sometimes
exhibited a ruffled cell shape, showing small protrusions and grooves. (B) Western
blot probing CetZ1 in the wild-type and CetZ1 knock-out strains at 15 h post-
resuspension in the indicated media. (C) Coulter cytometry cell-volume distributions
of the wild-type and CetZ1 knock-out strains at 15 h post-resuspension in Hv-Cab and
Hv-Ca (-TE).

Fig 1

Colony - resuspended	Liquid culture		
	OD = 0.05	OD = 0.3	OD = 0.9

2.5 µm

10 µm

Fig 2

(A) Passage 1

OD$_{600}$

- HvCab
- HvCab (-TE)
- HvCab (-Vitamin)
- HvCab (-CAA)

(B) Passage 2

OD$_{600}$

(C) Passage 3

OD$_{600}$

Time (days)

(D) Passage 1

Frequency (%)

(E) Passage 2

Frequency (%)

(F) Passage 3

Frequency (%)

Cell Circularity (%)

(G) Hv-Cab (+TE)

(H) Hv-Ca (-TE)

Tubulated cells (-TE)

Fig 3

Fig 4

Fig 5

Key (panels A and C):

(a) TEs were added directly to culture
(b) Culture diluted into Hv-Cab (+TE)
(c) Culture diluted into Hv-Cab (+10xTE)
(d) Culture diluted into Hv-Ca (-TE)

(B) 0 h

(C) 4 h   8 h   12 h

(A)

Fig 6

(A) Hv-Cab (+TE)

(B) Hv-YPCab (+TE)

(C) Hv-Ca (-TE)

(D) Hv-Ca (-TE)

(E) Hv-YPC (-TE)

Fig 7

(A)

| | Hv-Cab (+TE) | Hv-YPCab (+TE) | Hv-Ca (-TE) | Hv-YPC (-TE) |

Wild-type

ΔcetZ1

+CetZ1(E218A)

(B)

1 2 3 4 5

1) Pure CetZ1
2) Wild-type – Hv-Cab (+TE)
3) ΔcetZ1 – Hv-Cab (+TE)
4) Wild-type – Hv-Ca (-TE)
5) ΔcetZ1 – Hv-Ca (-TE)

(C)

- WT in Hv-Cab
- ΔcetZ1 in Hv-Cab
- WT in Hv-Ca(-TE)
- ΔcetZ1 in Hv-Ca(-TE)

Frequency (%)

Cell Volume (µm³)