A first perturbome of *Pseudomonas aeruginosa*: Identification of core genes

related to multiple perturbations by a machine learning approach

Jose Arturo Molina Mora (***corresponding author***)

Research Center in Tropical Diseases (CIET), University of Costa Rica, Costa Rica

Email: jose.molinamora@ucr.ac.cr

Pablo Montero-Manso

Department of Mathematics, University of A Coruña, Spain

Email: pmonteromanso@udc.es

Raquel García Batán

Research Center in Tropical Diseases (CIET), University of Costa Rica, Costa Rica

Email: raquel.garcia@ucr.ac.cr

Rebeca Campos Sánchez

Research Center in Cellular and Molecular Biology, (CIBCM), University of Costa Rica, Costa Rica

Email: rebeca.campos@ucr.ac.cr

Jose Vilar Fernández

Department of Mathematics, University of A Coruña, Spain

Email: josevilarf@udc.es

Fernando García Santamaría

Research Center in Tropical Diseases (CIET), University of Costa Rica, Costa Rica

Email: fernando.garcia@ucr.ac.cr

Abstract

Tolerance to stress conditions is vital for organismal survival, including bacteria under specific environmental

conditions, antibiotics, and other perturbations. Some studies have described common modulation and

shared genes during stress response to different types of disturbances (termed as perturbome), leading to the

idea of central control at the molecular level. We implemented a robust machine learning approach to identify

and describe genes associated with multiple perturbations or perturbome in a *Pseudomonas aeruginosa* PAO1

model.

Using microarray datasets from the Gene Expression Omnibus (GEO), we evaluated six approaches to rank

and select genes: using two methodologies, data single partition (SP method) or multiple partitions (MP

method) for training and testing datasets, we evaluated three classification algorithms (SVM Support Vector

Machine, KNN K-Nearest neighbor and RF Random Forest). Gene expression patterns and topological features

at the systems level were included to describe the perturbome elements.

We were able to select and describe 46 core response genes associated with multiple perturbations in *P.*

aeruginosa PAO1 and it can be considered a first report of the *P. aeruginosa* perturbome. Molecular

annotations, patterns in expression levels, and topological features in molecular networks revealed biological

functions of biosynthesis, binding, and metabolism, many of them related to DNA damage repair and aerobic

respiration in the context of tolerance to stress. We also discuss different issues related to implemented and

assessed algorithms, including data partitioning, classification approaches, and metrics. Altogether, this work

offers a different and robust framework to select genes using a machine learning approach.

Keywords: Perturbations, *Pseudomonas aeruginosa*, machine learning, gene selection, perturbome.

1. Introduction

Cell stress can be defined as a wide range of molecular changes that cells undergo in response to environmental, physical, chemical, or biological stressors; sensing and responding to stress is critical for survival [1]. These biological functions and metabolic activities are executed through complex physical and regulatory interactions of genes that resemble networks [2]. Additionally, tolerance to stress conditions (i.e. stressors, perturbations, or disturbances) is vital for organismal survival, including bacteria under diverse environmental conditions, including antibiotics [1].

Several studies have revealed diverse molecular levels that can explain the general response to disturbances in many organisms. However, detailed mechanisms related to responses to perturbations remain poorly understood [3]. Available reports usually focus on specific stressors and relatively few studies have focused on common, central, and universal determinants affected by multiple perturbations [3]. This concept has been recently referred to as the perturbome [4,5]. For example, in eukaryotic organisms, including plants and human cancer models, some studies have described diverse stress-response genes as common modulators for different types of disturbances, suggesting a central control mechanism [2,4–6]. In prokaryotic models, similar findings have been reported for *Escherichia coli* [3,7].

Additionally, a comprehensive study of gene-interactions allows the identification of functional relationships among genes [8], their products, and the underlying biological phenomena that are critical to understand phenotypes under different biological conditions [9,10]. In the context of cell stress, the response to different environmental or experimental stimuli can be recognized by distinct gene expression patterns. This can be inferred from transcriptomic profiling data and functional associations using high throughput molecular technologies such as microarrays or RNASeq [2]. However, a challenge with these technologies is the large amount of high complexity data generated. Specialized bioinformatics analyses are required to select relevant information and to reduce noise that distinguishes the molecular determinants for particular biological conditions. Thus, a primary objective of transcriptomic profiling is to find an optimal subset of genes that could be used to characterize and classify unknown samples [11]. This gene selection is not obvious and complex due to the thousands of genes to select from [12].

To study the central response determinants to different perturbations in a living organism, we used the

model *Pseudomonas aeruginosa* PAO1 (the reference strain of Pseudomonals). *P. aeruginosa* is a Gram-

negative gamma-proteobacterium with a noteworthy metabolic versatility and adaptability enabling the

colonization of diverse niches infecting plants, animals, and humans alike [8,13]. This bacterium is an

opportunistic pathogen that causes a variety of infections in immunocompromised hosts, and it is associated

with complications in cystic fibrosis and is a major cause of death among patients with this disease [14]. For

Pseudomonals, the molecular mechanisms associated with most biological processes remain unclear causing

limited action to modulate responses, including the susceptibility to stressors, environmental factors, and

experimental conditions.

In *P. aeruginosa*, the response to disturbances has been reported to be influenced by <10% of coding

genes [15,16], including complex molecular networks with pleiotropic patterns [17], i.e., a single perturbation

induces the modulation of hundreds of genes [15,16]. These networks can be modeled using system biology

approaches to recognized emergent patterns at the molecular level, as previously reported [18–21]. Large-

scale molecular networks are regularly built using databases and can be used not only to understand biological

functions, regulatory mechanisms, pathways, or gene groups associated with the molecular response [22] but

also to eventually manipulate it [23].

In the response to stress conditions, gene expression profiling using transcriptomic analysis and

topological metrics can be used to provide a global view of transcriptional regulation [24]. The identification

of hub genes and pathways associated with the network can be used to gain insights into the biological

meaning of the complex interactions [24,25]. This is a key point to translate molecular patterns into the

phenotypes, in particular for the response to perturbations [26].

Several studies have used machine learning algorithms at the transcriptomic level to recognize gene

expression patterns [2,27–29]. However, for many common biological contexts, the applicability and utility of

these machine learning approaches have not been fully explored and utilized [30], for example in the exposure

to multiple stressors and the molecular response. To our knowledge, only a few studies have used feature

selection methods on biological data to describe the effects of multiple perturbations in complex biological

systems [4,6] and so far none in *P. aeruginosa*. A related work in *P. aeruginosa* used a machine learning

approach to identify sets of genes that correlate with multiple culture media [31]. Due to differences in the

experimental design, this study and our work are not comparable.

The use of microarray and other high throughput technologies data involve challenges for machine

learning approaches. These include the curse of dimensionality [32,33], normalization of raw values to

compare samples [34,35], data partitions for training and testing models [36,37], and evaluation of

performance [34,38]. Since comparison between the machine learning algorithms is completely variable

[11,30,33,39,40], Support Vector Machine (SVM) [41], K-Nearest neighbor (KNN) [40], and Random Forest (RF)

[42] have been successfully used with microarray gene expression data allowing the recognition of emerging

patterns [39].

Here, we hypothesize that perturbations on living cells will trigger global reprogramming of multiple

molecular determinants that can be sensed at the transcriptional level. The initial central response after acute

stress will then expand producing the global molecular rearrangement (Figure 1-A). Then, pleiotropic and

specific effects on gene networks will be reflected as changes in gene expression profiles and the complexity

of molecular regulation at other levels. Therefore, by using a machine learning approach, common molecular

features (for all stressors) could be identified as a central or core determinant.

Thus, the aims for this study were (i) to implement a machine learning approach to select genes from

gene expression data, and (ii) to identify and describe a subset of genes that can be associated with multiple

perturbations or the core perturbome in *P. aeruginosa*.

2. Materials and Methods

The overall methodology of the study is presented in Figure 1-B. In brief, after a data selection,

normalization and gene selection to define the perturbome were run. To achieve this, we considered two

different procedures to split the data (training and testing datasets for the machine learning approaches: SVM,

KNN, and RF): a single partition (SP method) and another with multiple partitions (MP method). Relations

between genes were represented using both large scale and small-world networks, and a final comparison

between conditions, an analysis of differential expression, and gene annotation were performed.

2.1 Datasets

To compare gene expression profiles of strain *P. aeruginosa* PAO1 when exposed to multiple

perturbations, the GEO database (https://www.ncbi.nlm.nih.gov/geo/) was used for a systematic selection of

datasets (Figure 2-A-B). Initial evaluation by the organism and mRNA profiles by GPL84 platform (Affymetrix

Pseudomonas aeruginosa PAO1 Array, with all 5549 protein-coding sequences) identified 156 series of

datasets with 1310 samples (Date of Access: January 25th, 2018). In a second step, data were selected

according to experimental design if they included perturbations, leaving only 47 series. Finally, to make

datasets comparable by experimental conditions, evaluation and selection were done for series with similar

culture conditions (Luria Bertani LB medium and exponential phase when measuring mRNA profile) and if a

control condition was available (without any perturbation or treatment). The final dataset was composed of

10 series with 71 samples (Series GSE2430, GSE3090, GSE4152, GSE5443, GSE7402, GSE10605, GSE12738,

GSE13252, GSE14253 and GSE36753).

Some series included temporal measurements (see Experiment 3 of the Figure 2-B) which we considered

as separate perturbations, resulting in replicates of 10 controls and 14 perturbations (Figure 3-A):

azithromycin with 2 series (AZM-a and AZM-b) [43,44], Hydrogen peroxide (H_2O_2) [45], copper (Cu) [46],

sodium hypoclorite (NaClO) [47], ortho-phenylphenol (OPP) at 20 and 60 minutes [48], colistin (COL) [49],

chlorhexidine diacetate (CDA) at 10 and 60 minutes [50], E-4-bromo-5-bromomethylene-3-methylfuran-2-5H-

one (BF8) [51] and ciprofloxacin CIP at 0, 30 and 120 minutes [19]. Details of each study (experimental design

and main results) are provided in Table S1.

2.2 Pre-processing and comparison of global transcriptomic profiles

To compare all the 71 samples, a first analysis was the pre-processing step using Bioconductor 3.8

(https://www.bioconductor.org/) in the R software (Version 3.5) with classical functions for microarrays.

Robust MultiArray Average algorithm (RMA) was performed in the Affy package to correct background,

normalization, and summarization.

Subsequently, a clustering algorithm of Principal Component Analysis (PCA) was implemented to

compare global transcriptomic profiles between perturbations and controls. PCA was done with default

parameters of the Caret Package (caret.r-forge.r-project.org/) in R software.

In order to robustly select genes that could separate experimental conditions (controls and

perturbations) and to identify the core response of *P. aeruginosa*, two approaches of feature selection

protocols were implemented, as detailed below (Figure 2-B-C).

2.3 Gene ranking and selection by Single Partition SP method

To identify genes which expression values were commonly related to multiple perturbations, a first

approach was implemented considering a particular partition of the dataset for training and testing sets

(Figure 2-B-C). Single partition was established using the last replicate of each experiment, in both control and

perturbation. Because there were 14 perturbations and 10 controls (71 samples including replicates or

individual samples), a total of 24 samples were included in the testing dataset and the remaining 47 samples

were included for the training dataset (66% for training and 34% for the testing set).

Using this partition, the ranking of genes was done by a machine learning approach using three

classification algorithms: SVM, KNN, and RF. A homemade script in R included these functions of the Caret

package. For all three algorithms, default parameters were used for training and 10-fold cross-validation was

included. After this, the variable importance metric was calculated for all genes using the *varImp* function,

associating a specific value for each gene. In the case of SVM and KNN, the same importance is calculated

because the function is model-free for these cases (as detailed in Caret Package), resulting in the same list of

genes but metrics are specifically calculated for each algorithm. For RF, the accuracy and model parameters

were used to calculate the metric.

As similarly reported in [11], to evaluate the number of genes that should be selected in the top group

(the first K ranked genes, as candidates for the core response by each algorithm), multiple classification

models were systematically run, starting only with the highest-score ranked gene. In brief, after the training

with one gene, model performance was evaluated by calculating metrics using the testing dataset. Then,

training was run again when the next ranked gene was added, and new metrics were calculated. This process

was repeated up to complete all the ranked genes. Metrics included accuracy (correct classification

percentage), kappa value (inter-rater classification agreement), sensitivity, specificity, precision, recall,

prevalence, and F1 score (harmonic average of the precision and recall).

Selection of the K value of top genes was based on the following criteria: (i) the stability of the metrics

(priority for accuracy, kappa, and F1) when the increment of ranked genes was done, as suggested in [11], (ii)

consensus value suitable for all the three algorithms (including a 10% of tolerance), and (iii) minimum number

of genes as possible. After the selection of the K value, ROC (Receiver-operating characteristic) curve and AUC

(Area under the curve) value were calculated for each algorithm. Finally, the selection of top K genes between

algorithms was compared by metrics and- the list of genes.

2.4 Gene ranking and selection by Multiple Partitions (MP) method

In order to identify genes related to multiple perturbations independently of a single/specific partition,

a second method using multiple random partitions was implemented (Figure 2-B-C). To address this, a random

data selection for training and testing sets was done using the *createDataPartition* function. Partition was set

to 80% (57 samples) for the training set and the remaining for the testing set (14 samples) with experimental

conditions equally distributed. Then, protocols with SVM, KNN, and RF algorithms (same conditions as

previously described in SP method, with 10-fold cross-validation) resumed the analysis with a final ranking of

genes using the *varImp* function. Using only the top K of ranked genes (K value determined using the criteria

described in the SP method), a new set of training/testing sets were used for evaluating the performance of

the models and each metric was stored with the list of the K ranked genes. This full process was automatically

repeated 100 times using the *replicate* function, starting with a new random partition and finishing with the

list of the K genes and the metrics associated with that partition. Finally, for each algorithm, full data of all the

runs were analyzed for determining the frequency of the appearance of genes (*table* function) and calculating

the average and dispersion of metrics across all the 100 runs. The definitive list of the K more frequent genes

was established for each algorithm after this comparison by frequency.

2.5 Identification of core response genes

After selection of top K genes (candidates of the perturbome) in each algorithm by SP and MP methods,

a comparison of genes was done using UpSets plots (https://github.com/hms-dbmi/UpSetR) in R software.

We called the "support" of a gene the number of algorithms (out of six approaches) in which a gene was

selected as a candidate. Genes with a support ≥4 (i.e., it was identified by at least four algorithms) were

considered as part of the final list of genes of the core perturbome. This consideration guaranteed that a gene

was identified by the two methods and at least by two different classification algorithms.

Gene relationships were represented using molecular networks using a large scale model (using a top-

down systems biology approach). The network was built using protein-protein interaction (PPI) graphs in

PseudomonasNet database (www.inetbio.org/pseudomonasnet/). Visualization of the network and

measurement of the topological features were performed using Cytoscape software [52].

2.6 Description and comparison of core response genes

In order to describe and compare the genes associated with the core response of *P. aeruginosa,* by

experimental conditions, four analyzes were established. First, PCA was evaluated again but now only

considering genes of the core response. Based on distribution in the case of PCA, representation of centroids

was done using the Kmeans algorithm.

Second, using the PseudomonasNet database, a small-world network was built and then exported into

Cytoscape software with the genes of the core response. The information of topological features (including

connectivity) and expression levels of kmedoids were incorporated into different versions of the network. To

identify central or hub genes in the network in Cytoscape, the cytoHubba app was used [53] with the top 5

elements according to the metrics of degree, bottleneck, and betweenness. All identified genes in any out of

the three approaches were assigned as hub genes.

Third, a classic analysis of differentially expressed genes (DEGs, *p<0.05*) was implemented in R with

Limma package (https://www.rdocumentation.org/packages/limma/versions/3.28.14) using empirical Bayes

moderated t-statistic (eBayes) with Benjamini and Hochberg method for *p*-value correction [54]. This was

done to compare our results with a classical approach for gene expression.

Finally, to give biological interpretation to the selected genes, to determine levels of expression reported

in databases, and to study metabolic pathways involved under each perturbation, an exhaustive annotation

was made using the databases PseudomonasDB (gene ontology), GEO database, KEGG annotation, and

particular literature. This information was integrated with the results obtained by all the analysis and the DEGs

to fully describe the genes that make up the core response or perturbome of *P. aeruginosa* PAO1.

Availability of data and material

The datasets generated and/or analyzed during the current study are available in:

• Public raw data used in this study: GEO database (https://www.ncbi.nlm.nih.gov/geo/), data Series

GSE2430, GSE3090, GSE4152, GSE5443, GSE7402, GSE10605, GSE12738, GSE13252, GSE14253 and

GSE36753.

• Normalized data and R Scripts: https://github.com/josemolina6/CoreResponsePae

3. Results

3.1 Perturbome genes of P. aeruginosa *can be identified by a machine learning approach using SP and MP*

methods

A total of 71 samples of 10 controls and 14 perturbations were considered for the study, with comparable

expression levels (Figure 3-A). Global transcriptomic profiles (all 5549 genes) were compared by a PCA,

revealing a mixed pattern (no separation) between perturbations and controls (Figure 3-B). Two samples (BF8

and Control 8) resulted in "extreme" global profiles. Two methods using machine learning (SP and MP) were

implemented in order to robustly rank and select genes associated with multiple perturbations in *P.*

aeruginosa (Figure 2). In each method, three classification algorithms were evaluated: SVM, KNN, and RF.

The first method (SP) implemented a gene ranking based on variable importance using a single/specific

data partition. Metrics associated with RF are shown in Figure 4, and in supplementary Figure S1 for SVM and

KNN. After the ranking was established, multiple classification models were run with the ranked genes (Figures

4-A and supplementary Figure S1-A-C). For each classification model, the stability of the three metrics was

evaluated to select the suitable K value of genes that could be applied to all algorithms at the same time. For

SVM and RF, stable values of metrics are given with at least the first 51 ranked genes, meanwhile it is 45 genes

for KNN. Considering a 10% of tolerance with the highest of these values, K=56 was selected as the number

of top genes that were included as preliminary candidates of the core response according to each algorithm.

With this value, metrics of each algorithm were compared (Table S2). For example, accuracy was 0.79, 0.71,

and 0.75 for SVM, KNN, and RF respectively in the SP method. SVM obtained a better performance according

to kappa, sensitivity, recall, and F1 scores, but higher values of specificity and precision resulted for RF. Also,

the ROC curve and AUC value were calculated (Figures 4-B and supplementary Figure S1-B-D). The best

performance was obtained for RF with AUC = 0.92, then 0.82 for SVM, and finally 0.76 for KNN. Since the

ranking by *importance* for SVM and KNN is the same (model-free strategy, see details in

https://topepo.github.io/caret/variable-importance.html), they shared the same list of top 56 genes.

Comparison between implementations showed that 21 genes were identified by the three algorithms at the

same time, 35 exclusively by RF, and the same number for SVM/KNN. In total 91 genes were identified by any

of the algorithms. The list of genes and the importance value of the top 56 ranked genes for each approach is

presented in Figures 4-C for RF and Figure S1-E for SVM/KNN.

In a second approach, the MP method was implemented using multiple random partitions. Same SVM,

KNN, and RF algorithms were evaluated by running 100 iterations with different partitions and top-56 more

frequent genes for each method were selected. Details of ranking and frequency are shown in Figure 5-A

(SVM) and supplementary Figure S2 (KNN and RF). Metrics for all 100 iterations are presented in Table S2,

Figure 5-B, and supplementary Figure S2. Accuracies for all the models were 0.66, 0.69, and 0.70 for SVM,

KNN, and RF respectively. Specific values for kappa, precision, recall, and F1 score were found for each

algorithm. When a comparison of the list of genes was done, 29 genes were identified at the same time by all

implementations, and 23 by both SVM and KNN.

To identify preliminary core response genes in *P. aeruginosa*, the lists of top 56 genes selected by each

algorithm and method were jointly represented using UpSet plots (Figure 6-A). A total of 118 different genes

were identified and 15 genes were simultaneously identified by all the algorithms. The KEGG annotation of all

the genes revealed functions associated with energy and DNA metabolisms, biosynthesis, ribosomal activity,

and other pathways (Figure 6-B). See below for more details of the annotation.

The distribution of all 118 genes on a large-scale molecular network is presented in Figure 7-A. Results

show that selected genes are associated with different subgroups of highly connected genes.

The topological metrics of the network showed a non-random arrangement with 90 edges (more than

expected by chance, enrichment p-value: < 1.0e-16), 3.9 average number of neighbors, a clustering coefficient

of 0.51, and others. Besides, 10 hub genes were identified among the core perturbome, which are

representative elements of the network. More details in Figure 7-A and Table 1.

The final version of the core perturbome was established by selecting genes with a support ≥4 (i.e found

by at least four algorithms). In total 46 genes were finally associated with core response. Notwithstanding six

individual classifiers were implemented (each one with specific datasets), the consensus strategy to select the

final genes (by the support) had no effects on the resolution of the classes (control or perturbation), as shown

with the PCA with the transcriptomic profile using the 46 genes.

In the representation as a small-world network, relationships between the 46 genes revealed different

topological patterns of connectivity between molecules, being *nuoC* and *nuoF* the ones with higher

connectivity (connection degree). In addition, only six genes had no connections between them (Figure 7-B).

3.2 Comparisons of core response genes show the separation of global transcriptomic profiles according to

experimental conditions and biological functions related to tolerance to stress

After the selection of the core response genes, comparisons between controls and perturbations were

done to characterize these genes by global profiles. First, the PCA was implemented to compare the global

profiles using the 46 genes of the perturbome, in which the distribution shows a well-defined separation

between controls and perturbations (Figure 7-C). A Kmedoid algorithm (k = 2) was able to identify two clusters

enriched by samples of each condition: one consisting entirely of perturbation samples (11 samples, blue

color) and the other mostly by control samples (10 controls and 3 disturbances, red color). In the case of the

blue cluster, the kmedoid was sample CIP-120min, meanwhile control 6 was selected for another cluster.

Supplementary analysis of gene expression was included by comparing levels for all core response genes

(Figure S3-A) and comparing expression levels of the kmedoids on the small-world network (Figure S3-B-C).

On the other hand, gene annotation revealed that biological processes related to most of the genes are

metabolism, molecule binding, and biosynthesis (Figure 8-A). At the level of pathways, genes of the core

perturbome have participation in processes of DNA damage response, DNA repairing, and response to

general/specific stimuli (Figure 6-B and Table 1). Other studies show distinct expression patterns for each

gene, depending on the disturbance as shown in Table 1. Finally, in order to compare the results of the

machine learning strategy with another approach, differential expression analysis was run. A total of 101 DEGs

were identified, of which 33 were shared by the core response (Figure 8-B). This means that 72% of core

response genes were also identified by another single and independent method. Annotation of DEGs for

biological processes (Figure 8-C) showed a similar profile to our machine learning approach.

4. Discussion

Living organisms face external and internal conditions that compromise cellular functions at molecular,

metabolic, and structural levels, disrupting their homeostasis [55,56]. The cell stress response is crucial for

organismal survival and complex networks are usually involved in the molecular mechanism related to

tolerance [1]. However, few studies have identified central and possible universal regulation of the response

to multiple disturbances, a concept termed as perturbome [4,5]. The common molecular response was

previously reported as a network of a common set of genes and pathways that can be generically associated

with multiple perturbations in plants [7], pathogenic bacteria [3], or cell lines models [4], and others.

In our approach using *P. aeruginosa,* but also applicable to other organisms, we hypothesized the

existence of a set of core genes regulating the response to stressors in a generic sense of different pathways.

This bacterium is a ubiquitous organism that causes infections in immunocompromised hosts [14]. It has a

high proportion (about 5%) of its genome dedicated to regulatory mechanisms, probably explaining its

adaptability to such a broad range of growth conditions [38]. Since the strain *P. aeruginosa* PAO1 is a clinical

isolate with a profile of multiresistance to many antibiotics [57], characterization of molecular mechanisms

involved in the tolerance to stressors in this strain could eventually help to modulate sensitivity and overcome

resistance. Exhaustive integration of -omics data and network analysis are required to clarify the molecular

mechanisms related to stress conditions and eventually use them for modulating cell response.

For example, we have a particular interest in *P. aeruginosa* due to we have been studying a high-risk

strain (called *P. aeruginosa* AG1) which was isolated in 2010 from a Costa Rican Hospital. We reported this

isolated in 2015 [58] and it is a Priority 1 organism [59] because it is resistant to carbapenems, a last-resort

antibiotic. We assembled the genome using Illumina and Nanopore sequencing [13], revealing the genomic

features including two integrons, which were studied in detail in another work [60]. Proteomic profiles after

exposure to antibiotics have been also studied [61]. Finally, we studied the transcriptome in response to

ciprofloxacin using RNA-Seq and systems biology [16]. We expect to integrate our results of the perturbome

(in the reference strain) with the Costa Rican strain (AG1) to identify the "expected general response" to

multiple perturbations (as a Pseudomonal), and more importantly, to identify the "exclusive response" to

antibiotics, to perform new assays. In the response to ciprofloxacin, we already demonstrated that energy

metabolism, RNA degradation, and stress response are induced, as found in other perturbations (that here

we call "the perturbome"), but also other more exclusive responses such as phage induction (very exclusive

and not part of the perturbome) are modulated with ciprofloxacin.

On the other hand, although nowadays RNA-Seq is one of the most common high-throughput molecular

strategies to analyze transcriptomic data, the selection of microarray data to analyze the perturbome of *P.*

aeruginosa was based on the availability of data in databases. First, unlike other microorganisms (for example

intra-cellular bacteria), *P. aeruginosa* has a very variable genome size between 6 and 7.5 Mb, which is

explained in part due to the great repertory of the accessory genome [13]. Because of this, it is difficult to

combine molecular data of multiple strains. Second, the PAO1 strain is the reference strain (model for the

group during more than 20 years), and no other strain has the completeness of gene annotation as found for

this model. Finally, transcriptomic data of PAO1 is mainly found with microarray strategies, which is

significantly more abundant in databases than RNA-Seq.

4.1 Insights of algorithms to identify core response genes or perturbome

In our study, initial global transcriptomic profiles showed mixed patterns between samples according to

their experimental condition. Because of its complexity, raw microarray data showed noise and redundant

information that can explain the poor resolution of the initial PCA to identify classes [33]. Thus, a first step

was the normalization, which was particularly critical for multi-origin samples as in our case. Comparative

analysis needs to consider variation that depends not only on the experimental design but also parameters

including laboratory/culture conditions and mRNA extraction methods [62]. Subsequently, a feature (gene)

selection analysis was implemented for not only identifying genes associated with multiple perturbations in

P. aeruginosa, but also to improve the performance of predictive/descriptive models capable of separate

control and perturbation categories. In three experiments, two or three time points were considered as

perturbations (Figure 3-A and the example of Experiment 3 in Figure 2-B). Although an overrepresentation of

that perturbation cannot be discarded, we assume they are independent assays due to the global profile is

known to be different over time.

The potential patterns of the core response were investigated using a robust machine learning approach

by implementing six protocols, using SP and MP methods, and SVM, KNN, and RF algorithms in each case. This

was a critical step because gene selection from microarray data is complex *per se*. Feature reduction remains

a challenging task in transcriptomic studies because thousands of genes to select from, and it introduces an

additional layer of complexity in the modeling task [12,63]. To avoid bias and overfitting, implementations of

diverse strategies of data partitioning such as bootstrapping, random partitions, and cross-validation are

recommended. In general, these methodologies can robustly minorate the influence from noise, outliers,

absence of ground truth sets, and reduce variance [2,37,64]. In the specific case of implementations for the

study of molecular networks, partitions are suggested not only because to counteracts the curse of

dimensionality but also ground truth of candidate genes or regulatory networks are rarely available to allow

for an unbiased assessment of these methods [37].

The single partition SP method consisted of particular and invariable data for training (with internal cross-

validation) and another to test, and it is probably the most common approach used in machine learning. In

the case of the multiple partitions MP method, 100 random partitions of the dataset were run. MP method

had a dual consideration when splitting data (multiple partitions and the internal cross-validation). This

method can be considered as an *ensemble based on different data partitioning*, as it had been previously

proposed [36]. Datasets were divided using multiple random partitioning procedures and then genes were

ranked. After all runs, a final feature subset is determined by calculating the frequency of features in all the

runs [36]. An equivalent approach was implemented by Pai and collaborators to classify gene expression data

in a cancer model [65].

However, one possible problem with the MP approach is that cross-validation results may depend on

the similarity of testing and training sets. For example, all the samples of the experiment can be assigned as

part of the same training set and none in the testing set (or vice versa, for example, the red/blue/dark-green

replicates of the last arrangement of the MP method in Figure 2-B). A classification prediction method is only

expected to learn how to predict on unseen samples that are drawn from the same distribution as training

samples [37,64], and the MP method could violate this assumption. SP method guarantees this because it was

built using the last replicate for each perturbation and control (Figure 2-B, SP method).

In contrast, because of the presence of several biological replicates of the same perturbation in both

datasets of the SP method (one in the testing and the remaining in the training set), it can be considered that

the conformation of the testing data set is not completely independent, i.e. every sample in the training

dataset has a biological "copy" (replicate) itself in the training set. This scenario can introduce a bias in the

performance of the classifiers and robust analyses, such as multiple partitions and/or cross-validation, are

required to avoid a false impression of high predictive ability of the model and hence of the reliability of the

inferred influences on the gene [37].

Thus, both SP and MP methods have drawbacks but they can be compensated using the best features of

each strategy to robustly select features (genes).

On the other hand, many algorithms for dimension reduction have been proposed [6,32,33] but no

standard machine learning algorithm can be selected due to multiple evaluation results on completely variable

metrics associated with the performance [39]. Many studies have shown that SVM, KNN, and RF generally

outperform other traditional supervised classifiers [30,39,66]. In our case, a variable pattern was found for

different metrics in all evaluated implementations. For example, based on accuracy SVM for SP method and

RF for MP method resulted in higher scores, which agree with other studies using machine learning and

biological data [39,65].

In our subsequent analysis, we were interested in identifying a consensus list of candidate genes for the

core response in *P. aeruginosa*, resulting in 46 genes of the perturbome. When global profiles were compared

using these genes (PCA), control and perturbation classes were clearly separated (Figure 7-C). This gene

number seems to be a modest number of elements (less than 1% of all the available datasets with 5549 genes)

but it agrees with other studies, including machine learning methods [11,30] or other approaches

[11,35,67,68]. Besides, 72% of core response genes were also identified as DEGs with a similar annotation

profile. Differences can be explained by significant fluctuations in the differential expression results as

previously reported, mainly because it is not a consensus strategy (only based on *p*-value) and it does not

incorporate the estimates of the test performance (true positive/negative rates and other metrics) on the

results [2].

4.2 Biological insights of the core response genes in P. aeruginosa*: the perturbome*

In the study of complex systems, the lack of specific biological information at the single gene level and

the interactions make the extraction of the relevant information a complicated task in this field. This situation

remains as an unsolved issue of molecular high-throughput technologies, in which molecular data is generated

at a higher rate than our ability to find patterns or extract relevant information [69]. Thus, large-scale/top-

down system biology-based approaches have been used as a strategy to exploit and extract relevant

information of ample amount of -omics data [21], letting to predict interactions between genes [23].

With the representation of gene-to-gene interactions, system biology allows studying not only specific

genes or gene clusters but also the emergent properties found in the molecular networks [17]. Thus,

pleiotropic patterns can be recognized towards the response to perturbations, in which large-scale networks

can be built [69]. In the case of the core perturbome, the representation of genes showed a complex network

with topological properties that defines a non-random arrangement, including hub genes and pathways

associated with the response to stress and stimulus, metabolism, and ribosome activity. The network can be

inferred as a coordinated central response to the perturbations, independently of a particular stressor.

Genes of the core response or perturbome can be related to a central regulation network, and as the

convergent point of signal transduction, transcriptional regulation, and stress-related pathways, as it has been

suggested [2,4,5,56]. Annotation of the 46 genes shows that most of them are functionally related to

biosynthesis, molecule binding, and metabolism (including an important number of hits for lipids), including

additional functions for the regulation of DNA damage repair, response to stimuli, and aerobic respiration.

Interestingly, hub genes represent these processes with high connectivity in the small-world network. For

example, the main functions associated with *fabA* are lipid metabolism, fatty acid, and lipid biosynthesis,

meanwhile for *nuoC* are aerobic respiration, electron/proton transport, and DNA damage. Finally, *nuoF* is

associated with aerobic respiration, electron/proton transport, and Krebs cycle.

In this sense, cells are equipped with systems and mechanisms to recover from the environmental stress

and stimuli to maintain all necessary physiological functions [70]. Other common stimuli such as low ATP, slow

growth, and ROS production can also occur before cells express stress-specific factors, but mediating a

common effect. For example, response to stress includes modulation of energy production and aerobic

electron transfer chain components. As it has been reported in *E. coli*, aerobic electron transport chain

components are down-regulated in response to growth arrest [71]. This corresponds with the global profiles

of expression of 46 core response genes. Also, regulation of lipid metabolism is relevant for survival in the

wide range of environmental conditions where bacteria thrive [72], even for biofilm-living forms [3] as *P.*

aeruginosa. Core response genes *fabG*, *lpxA*, and PA5174 could be implied in this process.

In the case of DNA damage repair (including the case of *cycB* and *gltP* genes), responses mediated by

SOS and *rpoS* help to maintain genome integrity, colonization, and virulence [19,73]. These responses are

activated under multiple disturbances and modulating a low energy production and shutdown of the

metabolism, promotes the formation of antibiotic resistance and biofilms [3]. Other related pathways for

some specific genes included regulation of the transcription during stress by RNA-binding proteins to

reprogram or shut down translation and to rescue the ribosome stalled by a variety of mechanisms induced

[74]. Three core response genes (*rpmH*, *tsf,* and PA2735) were annotated with these functions.

Jointly, the relatively few diversity of metabolic functions and pathways makes sense in order to ensure

redundancy and robustness in the response to stimuli. Similar results, regarding enriched pathways, have

been obtained in other studies with eukaryotic models, including disturbed human cell lines [4,5,56],

Arabidopsis thaliana under physical and genotoxic stresses [2], or a genome-wide association study of a

generic response to stress conditions [6]. In the case of prokaryotic organisms, two studies have used

Escherichia coli as a model to identify differentially expressed genes after exposure to stress conditions [3]

and to create networks associated with the response to fluctuating environments [7]. Differences with other

organisms and disturbances can suggest that response cell stress can be organismal specific, although

heterogeneity has also been suggested as a reasonable explanation because of differences in the response in

a homogeneous cell population [56,75].

As in our case, the response to stimuli and stressors is orchestrated by a pleiotropic modulation [20]

which can be associated with central regulation. Alternative mechanisms such as cross stress protection (the

ability of a stress condition to protect against other stressors) [7], the role of sigma factors, and specific two-

component systems [3] can contribute to explaining this phenomenon. The molecular response can lead to

regulate multiple biological activities including metabolism, replication, transcription and translation, changes

in membrane composition, motility, modification gene expression, expression of virulence factors, multi-drug

resistant phenotypes and biofilm formation, and others [56].

For further works, we are running experiments to validate our findings here, in which *P. aeruginosa* is

exposed to different perturbations and gene expression is quantified by RT-qPCR. This will also allow us to

build small-world networks with the representation of the core perturbome as a directed graph. Using a

bottom-up system biology approach using differential equations, sensitivity analysis, and other machine

learning algorithms, we expect to identify and manipulate the points of control the network, as we have

previously done in other models [76].

Taking all together, results of our study suggest that identification of core response genes associated

with multiple perturbations or perturbome in *P. aeruginosa* can define a central network available to

modulate a basic response that includes biological functions such as biosynthesis, binding, and metabolism,

many of them related to DNA damage repair and aerobic respiration. To our knowledge, this study can be

considered a first report of the *P. aeruginosa* perturbome.

Further analyses are required to explore the potential use of perturbome network to modulate

(positively or negatively) the response to disturbances, to model molecular circuits, to identify possible

biomarker genes of stressed states, and to experimentally validate our findings. Besides, this approach can be

used to model the perturbome in other *P. aeruginosa* strains, as we hope to run soon with a genome we

recently described [13], and other organisms.

5. Conclusions

A robust machine learning approach was implemented to identify and describe core response genes to

multiple perturbations in *P. aeruginosa*. Using public microarray data, two independent partition strategies

(single and multiple with SP and MP methods respectively), and three classification algorithms, we were able

to identify 46 perturbome elements. Both network analysis and functional annotations of these genes showed

coordinated modulation of biological processes in response to multiple perturbations, including metabolism,

biosynthesis, and molecule binding, associated with DNA damage repairing, and aerobic respiration, all

probably related to tolerance to stressors, growth arrest, and molecular regulation. We also discussed

different issues related to implemented and assess algorithms of normalization analysis, data partitioning,

classification approaches, and metrics.

Abbreviations

AUC: Area under the curve

AZM: Azithromycin

B8F: E-4-bromo-5-bromomethylene-3-methylfuran-2-5H-one

CDA: Chlorhexidine diacetate

CIP: Ciprofloxacin

COL: Colistin

Cu: Copper

DEGs: Differentially expressed genes

GEO: Gene Expression Omnibus

H2O2: Hydrogen peroxide

HC: Hierarchical Clustering

KNN: K-Nearest Neighbor

LB: Luria Bertani

mRNA: Messenger RNA

NaClO: Sodium hypochlorite

OPP: Ortho-phenylphenol

PCA: Principal Component Analysis

PPI: Protein-protein interaction

RF: Random Forest

ROC: Receiver-operating characteristic

SVM: Support Vector Machine

Declarations

Ethics approval and consent to participate

Not Applicable

Consent for Publication

Not Applicable

Availability of data and material

The datasets generated and/or analyzed during the current study are available in:

Public raw data used in this study: GEO database (https://www.ncbi.nlm.nih.gov/geo/), data Series GSE2430,

GSE3090, GSE4152, GSE5443, GSE7402, GSE10605, GSE12738, GSE13252, GSE14253 and GSE36753.

Normalized data and R Scripts: https://github.com/josemolina6/CoreResponsePae

Competing interests

No competing interests to declare.

Funding

This work was funded by projects B8114 Definición de la red transcripcional y de las alteraciones genómicas

inducidas por la ciprofloxacina en *Pseudomonas aeruginosa* AG1 and B8152 proNGS 1.0: Implementación y

evaluación de protocolos de análisis de datos de tecnologías NGS y afines para el estudio de sistemas

biológicos complejos, Universidad de Costa Rica (period 2017-2020), Costa Rica.

Authors' contributions

JMM and FGS participated in the conception, design of the study, and data acquisition. JMM, PMM, RGB, and

JVF participated in data analysis and interpretation. JMM, RCS, JVF, and FGS participated in the interpretation

of the data analysis. JMM drafted the manuscript and all authors were involved in its revision. All authors read

and approved the final manuscript.

Acknowledgments

We thank students Iosif Forero Trelles, Daniel Solano Alvarado, and Daniela Aguilar Orozco for their

collaborations in different activities of the project.

References

1. DeLong, E.F. *Prokaryotes : prokaryotic physiology and biochemistry*; Springer, 2012; ISBN 9783642301407.

2. Ma, C.; Xin, M.; Feldmann, K.A.; Wang, X. Machine Learning-Based Differential Network Analysis: A Study of

Stress-Responsive Transcriptomes in Arabidopsis. *Plant Cell* **2014**, *26*, 520–537, doi:10.1105/tpc.113.121913.

3. Nagar, S.D.; Aggarwal, B.; Joon, S.; Bhatnagar, R.; Bhatnagar, S. A Network Biology Approach to Decipher Stress

Response in Bacteria Using *Escherichia coli* As a Model. *Omi. A J. Integr. Biol.* **2016**, *20*, 310–324,

doi:10.1089/omi.2016.0028.

4. Caldera, M.; Müller, F.; Kaltenbrunner, I.; Licciardello, M.P.; Lardeau, C.H.; Kubicek, S.; Menche, J. Mapping the

perturbome network of cellular perturbations. *Nat. Commun.* **2019**, *10*, doi:10.1038/s41467-019-13058-9.

5. Sadeh, S.; Clopath, C. Theory of Neuronal Perturbome: Linking Connectivity to Coding via Perturbations. *bioRxiv*

2020, 2020.02.20.954222, doi:10.1101/2020.02.20.954222.

6. Bermingham, M.L.; Pong-Wong, R.; Spiliopoulou, A.; Hayward, C.; Rudan, I.; Campbell, H.; Wright, A.F.; Wilson,

J.F.; Agakov, F.; Navarro, P.; et al. Application of high-dimensional feature selection: Evaluation for genomic

prediction in man. *Sci. Rep.* **2015**, *5*, 1–12, doi:10.1038/srep10312.

7. Dragosits, M.; Mozhayskiy, V.; Quinones-Soto, S.; Park, J.; Tagkopoulos, I. Evolutionary potential, cross-stress

behavior and the genetic basis of acquired stress resistance in Escherichia coli. *Mol. Syst. Biol.* **2014**, *9*, 643–643,

doi:10.1038/msb.2012.76.

8.   Nogales, J.; Guðmundsson, S.; Duque, E.; Ramos, J.L.; Palsson, B.Ø. Expanding the computable reactome in

Pseudomonas putida reveals metabolic cycles providing robustness. *bioRxiv* **2017**, 139121, doi:10.1101/139121.

9.   Kc, K.; Li, R.; Cui, F.; Haake, A.R. GNE: A deep learning framework for gene network inference by aggregating

biological information. *Bioinformatics* **2018**, 1–9.

10.   Szklarczyk, D.; Gable, A.L.; Lyon, D.; Junge, A.; Wyder, S.; Huerta-Cepas, J.; Simonovic, M.; Doncheva, N.T.;

Morris, J.H.; Bork, P.; et al. STRING v11: protein–protein association networks with increased coverage,

supporting functional discovery in genome-wide experimental datasets. *Nucleic Acids Res.* **2019**, *47*, D607–

D613, doi:10.1093/nar/gky1131.

11.   Li, Y.; Wang, N.; Perkins, E.J.; Zhang, C.; Gong, P. Identification and optimization of classifier genes from multi-

class earthworm microarray dataset. *PLoS One* **2010**, *5*, 1–9, doi:10.1371/journal.pone.0013715.

12.   Saeys, Y.; Inza, I.; Larranaga, P. A review of feature selection techniques in bioinformatics. *Bioinformatics* **2007**,

23, 2507–2517, doi:10.1093/bioinformatics/btm344.

13.   Molina-Mora, J.-A.; Campos-Sánchez, R.; Rodríguez, C.; Shi, L.; García, F. High quality 3C de novo assembly and

annotation of a multidrug resistant ST-111 Pseudomonas aeruginosa genome: Benchmark of hybrid and non-

hybrid assemblers. *Sci. Rep.* **2020**, *10*, 1392, doi:10.1038/s41598-020-58319-6.

14.   Kumar, S.S.; Penesyan, A.; Elbourne, L.D.H.; Gillings, M.R.; Paulsen, I.T. Catabolism of Nucleic Acids by a Cystic

Fibrosis Pseudomonas aeruginosa Isolate: An adaptive pathway to cystic fibrosis sputum environment. *Front.*

Microbiol. **2019**, *10*, 1–14, doi:10.3389/fmicb.2019.01199.

15.   Wu, W.; Jin, S. PtrB of Pseudomonas aeruginosa suppresses the type III secretion system under the stress of

DNA damage. *J. Bacteriol.* **2005**, *187*, 6058–6068, doi:10.1128/JB.187.17.6058-6068.2005.

16.   Molina-Mora, J.A.; Chinchilla, D.; Chavarría, M.; Ulloa, A.; Campos-Sanchez, R.; Mora-Rodríguez, R.A.; Shi, L.;

García, F. Transcriptomic determinants of the response of ST-111 Pseudomonas aeruginosa AG1 to ciprofloxacin

identified by a top-down systems biology approach. *Sci. Rep.* **2020**, *10*, 1–23, doi:10.1038/s41598-020-70581-2.

17.   Schulz, S.; Eckweiler, D.; Bielecka, A.; Nicolai, T.; Franke, R.; Dötsch, A.; Hornischer, K.; Bruchmann, S.; Düvel, J.;

Häussler, S. Elucidation of Sigma Factor-Associated Networks in Pseudomonas aeruginosa Reveals a Modular

Architecture with Limited and Function-Specific Crosstalk. *PLoS Pathog.* **2015**, *11*, 1–21,

doi:10.1371/journal.ppat.1004744.

18.   Brazas, M.D.; Brazas, M.D.; Hancock, R.E.W.; Hancock, R.E.W. Ciprofloxacin Induction of a Susceptibility

Determinant in Pseudomonas aeruginosa. *Antimicrob. Agents Chemother.* **2005**, *49*, 3222–3227,

doi:10.1128/AAC.49.8.3222.

19. Cirz, R.T.; O'Neill, B.M.; Hammond, J.A.; Head, S.R.; Romesberg, F.E. Defining the Pseudomonas aeruginosa SOS

response and its role in the global response to the antibiotic ciprofloxacin. *J. Bacteriol.* **2006**, *188*, 7101–7110,

doi:10.1128/JB.00807-06.

20. Molina-Mora, J.A.; Campos-Sanchez, R.; Garcia, F. Gene Expression Dynamics Induced by Ciprofloxacin and Loss

of Lexa Function in Pseudomonas aeruginosa PAO1 Using Data Mining and Network Analysis. In Proceedings of

the 2018 IEEE International Work Conference on Bioinspired Intelligence (IWOBI); IEEE, 2018; pp. 1–7.

21. Anupama, R.; Sajitha Lulu, S.; Mukherjee, A.; Babu, S. Cross-regulatory network in Pseudomonas aeruginosa

biofilm genes and TiO 2 anatase induced molecular perturbations in key proteins unraveled by a systems

biology approach. *Gene* **2018**, *647*, 289–296, doi:10.1016/j.gene.2018.01.042.

22. Fang, G.; Passalacqua, K.D.; Hocking, J.; Llopis, P.M.; Gerstein, M.; Bergman, N.H.; Jacobs-Wagner, C.

Transcriptomic and phylogenetic analysis of a bacterial cell cycle reveals strong associations between gene co-

expression and evolution. *BMC Genomics* **2013**, *14*, doi:10.1186/1471-2164-14-450.

23. Linde, J.; Schulze, S.; Henkel, S.G.; Guthke, R. Data- and knowledge-based modeling of gene regulatory

networks: An update. *EXCLI J.* **2015**, *14*, 346–378, doi:10.17179/excli2015-168.

24. Langfelder, P.; Horvath, S. WGCNA: An R package for weighted correlation network analysis. *BMC*

Bioinformatics **2008**, *9*, doi:10.1186/1471-2105-9-559.

25. Dai, H.; Zhou, J.; Zhu, B. Gene co-expression network analysis identifies the hub genes associated with immune

functions for nocturnal hemodialysis in patients with end-stage renal disease. *Med. (United States)* **2018**, *97*, 1–

8, doi:10.1097/MD.0000000000012018.

26. Lovelace, A.H.; Smith, A.; Kvitko, B.H. Pattern-triggered immunity alters the transcriptional regulation of

virulence-associated genes and induces the sulfur starvation response in pseudomonas syringae pv. tomato

DC3000. *Mol. Plant-Microbe Interact.* **2018**, *31*, 750–765, doi:10.1094/MPMI-01-18-0008-R.

27. Zhao, W.; Chen, J.J.; Perkins, R.; Wang, Y.; Liu, Z.; Hong, H.; Tong, W.; Zou, W.; Metzker, M.; Didelot, X.; et al. A

novel procedure on next generation sequencing data analysis using text mining algorithm. *BMC Bioinformatics*

2016, *17*, 213, doi:10.1186/s12859-016-1075-9.

28. Cornforth, D.M.; Dees, J.L.; Ibberson, C.B.; Huse, H.K.; Mathiesen, I.H.; Kirketerp-Møller, K.; Wolcott, R.D.;

Rumbaugh, K.P.; Bjarnsholt, T.; Whiteley, M. Pseudomonas aeruginosa transcriptome during human infection.

Proc. Natl. Acad. Sci. U. S. A. **2018**, *115*, doi:10.1073/pnas.1717525115.

29.   Glaab, E.; Bacardit, J.; Garibaldi, J.M.; Krasnogor, N. Using rule-based machine learning for candidate disease

645   gene prioritization and sample classification of cancer gene expression data. *PLoS One* **2012**, *7*,

646   doi:10.1371/journal.pone.0039932.

647   30.   Leung, R.K.K.; Wang, Y.; Ma, R.C.W.; Luk, A.O.Y.; Lam, V.; Ng, M.; So, W.Y.; Tsui, S.K.W.; Chan, J.C.N. Using a

648   multi-staged strategy based on machine learning and mathematical modeling to predict genotype-phenotype

649   risk patterns in diabetic kidney disease: A prospective case-control cohort analysis. *BMC Nephrol.* **2013**, *14*, 1,

650   doi:10.1186/1471-2369-14-162.

651   31.   Tan, J.; Doing, G.; Lewis, K.A.; Price, C.E.; Chen, K.M.; Cady, K.C.; Perchuk, B.; Laub, M.T.; Hogan, D.A.; Greene,

652   C.S. Unsupervised Extraction of Stable Expression Signatures from Public Compendia with an Ensemble of

653   Neural Networks. *Cell Syst.* **2017**, *5*, 63-71.e6, doi:10.1016/j.cels.2017.06.003.

654   32.   Kar, S.C. Comparing Prediction Accuracy for Supervised Techniques in Gene Expression Data. *Math. Theory*

655   *Model.* **2014**, *4*, 108–116.

656   33.   Raza, K.; Hasan, A. A Comprehensive Evaluation of Machine Learning Techniques for Cancer Class Prediction

657   Based on Microarray Data. *Int. J. Bioinform. Res. Appl.* **2015**, *11*, 397–416, doi:10.1504/IJBRA.2015.071940.

658   34.   Savli, H.; Karadenizli, A.; Kolayli, F.; Gundes, S.; Ozbek, U.; Vahaboglu, H. Expression stability of six housekeeping

659   genes: a proposal for resistance gene quantification studies of Pseudomonas aeruginosa by real-time

660   quantitative RT-PCR. *J. Med. Microbiol.* **2003**, *52*, 403–408, doi:10.1099/jmm.0.05132-0.

661   35.   Casares, F.M. A Simple Method for Optimization of Reference Gene Identification and Normalization in DNA

662   Microarray Analysis. *Med. Sci. Monit. Basic Res.* **2016**, *22*, 45–52, doi:10.12659/MSMBR.897644.

663   36.   Yang, P.; Zhou, B.B.; Yang, J.Y.-H.; Zomaya, A.Y. Stability of feature selection algorithms and ensemble feature

664   selection methods in bioinformatics. *Biol. Knowl. Discov. Handb. Preprocessing, mining, postprocessing Biol.*

665   *data* **2013**, 333–352, doi:10.1002/9781118617151.ch14.

666   37.   Tabe-Bordbar, S.; Emad, A.; Zhao, S.D.; Sinha, S. A closer look at cross-validation for assessing the accuracy of

667   gene regulatory networks and models. *Sci. Rep.* **2018**, *8*, 1–11, doi:10.1038/s41598-018-24937-4.

668   38.   Alqarni, B.; Colley, B.; Klebensberger, J.; McDougald, D.; Rice, S.A. Expression stability of 13 housekeeping genes

669   during carbon starvation of Pseudomonas aeruginosa. *J. Microbiol. Methods* **2016**, *127*, 182–187,

670   doi:10.1016/j.mimet.2016.06.008.

671   39.   Noi, P.T.; Kappas, M. Comparison of random forest, k-nearest neighbor, and support vector machine classifiers

672   for land cover classification using sentinel-2 imagery. *Sensors (Switzerland)* **2018**, *18*, doi:10.3390/s18010018.

673   40.   Li, L.; Weinberg, C.R.; Darden, T.A.; Pedersen, L.G. Gene selection for sample classification based on gene

674 expression data: study of sensitivity to choice of parameters of the GA/KNN method. *Bioinformatics* **2001**, *17*,

675 1131–42.

676 41. Vapnik, V.N.; Vladimir *Estimation of dependences based on empirical data*; Springer-Verlag, 1982; ISBN

677 0387907335.

678 42. Breiman, L. Random Forests. *Mach. Learn.* **2001**, *45*, 5–32, doi:10.1023/A:1010933404324.

679 43. Yusuf Nalca; Lothar Jänsch; Florian Bredenbruch; Robert Geffers, J.B.; Susanne Häussler Quorum-sensing

680 antagonistic activities of azithromycin in Pseudomonas aeruginosa PAO1: a global approach. *Antimicrob Agents*

681 *Chemother* **2008**, *50*, 1680–1688, doi:10.1128/AAC.50.5.1680.

682 44. Kai, T.; Tateda, K.; Kimura, S.; Ishii, Y.; Ito, H.; Yoshida, H.; Kimura, T.; Yamaguchi, K. A low concentration of

683 azithromycin inhibits the mRNA expression of N-acyl homoserine lactone synthesis enzymes, upstream of lasI or

684 rhlI, in Pseudomonas aeruginosa. *Pulm. Pharmacol. Ther.* **2009**, *22*, 483–486, doi:10.1016/j.pupt.2009.04.004.

685 45. Chang, W.; Small, D.A.; Toghrol, F.; Bentley, W.E. Microarray analysis of Pseudomonas aeruginosa reveals

686 induction of pyocin genes in response to hydrogen peroxide. *BMC Genomics* **2005**, *6*, 1–14, doi:10.1186/1471-

687 2164-6-115.

688 46. Teitzel, G.M.; Geddie, A.; De Long, S.K.; Kirisits, M.J.; Whiteley, M.; Parsek, M.R. Survival and Growth in the

689 Presence of Elevated Copper: Transcriptional Profiling of Copper-Stressed Pseudomonas aeruginosa. *J.*

690 *Bacteriol.* **2006**, *188*, 7242–7256, doi:10.1128/JB.00837-06.

691 47. Small, D.A.; Chang, W.; Toghrol, F.; Bentley, W.E. Comparative global transcription analysis of sodium

692 hypochlorite, peracetic acid, and hydrogen peroxide on Pseudomonas aeruginosa. *Appl. Microbiol. Biotechnol.*

693 **2007**, *76*, 1093–1105, doi:10.1007/s00253-007-1072-z.

694 48. Nde, C.W.; Jang, H.-J.; Toghrol, F.; Bentley, W.E. Toxicogenomic response of Pseudomonas aeruginosa to ortho-

695 phenylphenol. *BMC Genomics* **2008**, *9*, 473, doi:10.1186/1471-2164-9-473.

696 49. Cummins, J.; Reen, F.J.; Baysse, C.; Mooij, M.J.; O'Gara, F. Subinhibitory concentrations of the cationic

697 antimicrobial peptide colistin induce the pseudomonas quinolone signal in Pseudomonas aeruginosa.

698 *Microbiology* **2009**, *155*, 2826–2837, doi:10.1099/mic.0.025643-0.

699 50. Nde, C.W.; Jang, H.J.; Toghrol, F.; Bentley, W.E. Global transcriptomic response of Pseudomonas aeruginosa to

700 chlorhexidine diacetate. *Environ. Sci. Technol.* **2009**, *43*, 8406–8415, doi:10.1021/es9015475.

701 51. Pan, J.; Bahar, A.A.; Syed, H.; Ren, D. Reverting Antibiotic Tolerance of Pseudomonas aeruginosa PAO1 Persister

702 Cells by (Z)-4-bromo-5-(bromomethylene)-3-methylfuran-2(5H)-one. *PLoS One* **2012**, *7*,

703 doi:10.1371/journal.pone.0045778.

52. Shannon, P.; Markiel, A.; Ozier, O.; Baliga, N.S.; Wang, J.T.; Ramage, D.; Amin, N.; Schwikowski, B.; Ideker, T.

Cytoscape: A Software Environment for Integrated Models of Biomolecular Interaction Networks. *Genome Res.*

2003, *13*, 2498–2504, doi:10.1101/gr.1239303.

53. Chin, C.-H.; Chen, S.-H.; Wu, H.-H.; Ho, C.-W.; Ko, M.-T.; Lin, C.-Y. cytoHubba: identifying hub objects and sub-

networks from complex interactome. *BMC Syst. Biol.* **2014**, *8 Suppl 4*, S11, doi:10.1186/1752-0509-8-S4-S11.

54. Smyth, G.K. Linear Models and Empirical Bayes Methods for Assessing Differential Expression in Microarray

Experiments. *Stat. Appl. Genet. Mol. Biol.* **2004**, *3*, 1–25, doi:10.2202/1544-6115.1027.

55. Richter, K.; Haslbeck, M.; Buchner, J. The Heat Shock Response: Life on the Verge of Death. *Mol. Cell* **2010**, *40*,

253–266, doi:10.1016/j.molcel.2010.10.006.

56. Vihervaara, A.; Duarte, F.M.; Lis, J.T. Molecular mechanisms driving transcriptional stress responses. *Nat. Rev.*

Genet. **2018**, *19*, 385–397, doi:10.1038/s41576-018-0001-6.

57. Holloway, B.W. Genetic Recombination in Pseudomonas aeruginosa. *Microbiology* **1955**, *13*, 572–581,

doi:10.1099/00221287-13-3-572.

58. Toval, F.; Guzmán-Marte, A.; Madriz, V.; Somogyi, T.; Rodríguez, C.; García, F. Predominance of carbapenem-

resistant Pseudomonas aeruginosa isolates carrying blaIMP and blaVIM metallo-β-lactamases in a major

hospital in Costa Rica. *J. Med. Microbiol.* **2015**, *64*, 37–43, doi:10.1099/jmm.0.081802-0.

59. World Health Organization *Guidelines for the prevention and control of carbapenem-resistant*

Enterobacteriaceae, Acinetobacter baumannii and Pseudomonas aeruginosa in health care facilities; Geneva,

2017;

60. Molina-Mora, J.-A.; Garcia-Batan, R.; Garcia, F. Genomic context of the two integrons of ST-111 Pseudomonas

aeruginosa AG1: A VIM-2-carrying old-acquaintance and a novel IMP-18-carrying integron. *Res. Sq.* **2020**,

doi:10.21203/RS.3.RS-41474/V1.

61. Molina-Mora, J.A.; Chinchilla-Montero, D.; Castro-Peña, C.; García, F. Two-dimensional gel electrophoresis (2D-

GE) image analysis based on CellProfiler. *Medicine (Baltimore).* **2020**, *99*, e23373,

doi:10.1097/MD.0000000000023373.

62. Bustin, S.A.; Benes, V.; Nolan, T.; Pfaffl, M.W. Quantitative real-time RT-PCR – a perspective. *J. Mol. Endocrinol.*

2005, *34*, 597–601, doi:10.1677/jme.1.01755.

63. Piao, J.; Sun, J.; Yang, Y.; Jin, T.; Chen, L.; Lin, Z. Target gene screening and evaluation of prognostic values in

non-small cell lung cancers by bioinformatics analysis. *Gene* **2018**, *647*, 306–311,

doi:10.1016/j.gene.2018.01.003.

64.   Touw, W.G.; Bayjanov, J.R.; Overmars, L.; Backus, L.; Boekhorst, J.; Wels, M.; van Hijum, S.A.F.T. Data mining in

735   the Life Sciences with Random Forest: a walk in the park or lost in the jungle? *Brief. Bioinform.* **2013**, *14*, 315–

736   326, doi:10.1093/bib/bbs034.

737   65.   Pai, S.; Hui, S.; Isserlin, R.; Shah, M.A.; Kaka, H.; Bader, G.D. netDx: interpretable patient classification using

738   integrated patient similarity networks. *Mol. Syst. Biol.* **2019**, *15*, e8497, doi:10.15252/msb.20188497.

739   66.   Park, H.; Shimamura, T.; Imoto, S.; Miyano, S. Adaptive NetworkProfiler for Identifying Cancer Characteristic-

740   Specific Gene Regulatory Networks. *J. Comput. Biol.* **2017**, *25*, cmb.2017.0120, doi:10.1089/cmb.2017.0120.

741   67.   Falciani, F.; Diab, A.; Sabine, V.; Williams, T.; Ortega, F.; George, S.; Chipman, J. Hepatic transcriptomic profiles

742   of European flounder (Platichthys flesus) from field sites and computational approaches to predict site from

743   stress gene responses following exposure to model toxicants. *Aquat. Toxicol.* **2008**, *90*, 92–101,

744   doi:10.1016/j.aquatox.2008.07.020.

745   68.   Nota, B.; Verweij, R.A.; Molenaar, D.; Ylstra, B.; van Straalen, N.M.; Roelofs, D. Gene Expression Analysis Reveals

746   a Gene Set Discriminatory to Different Metals in Soil. *Toxicol. Sci.* **2010**, *115*, 34–40, doi:10.1093/toxsci/kfq043.

747   69.   Liu, W.; Li, L.; Long, X.; You, W.; Zhong, Y.; Wang, M.; Tao, H.; Lin, S.; He, H. Construction and Analysis of Gene

748   Co-Expression Networks in Escherichia coli. *Cells* **2018**, *7*, 19, doi:10.3390/cells7030019.

749   70.   Krämer, R. Bacterial stimulus perception and signal transduction: Response to osmotic stress. *Chem. Rec.* **2010**,

750   *10*, 217–229, doi:10.1002/tcr.201000005.

751   71.   Schurig-Briccio, L.A.; Farías, R.N.; Rodríguez-Montelongo, L.; Rintoul, M.R.; Rapisarda, V.A. Protection against

752   oxidative stress in Escherichia coli stationary phase by a phosphate concentration-dependent genes expression.

753   *Arch. Biochem. Biophys.* **2009**, *483*, 106–110, doi:10.1016/j.abb.2008.12.009.

754   72.   Parsons, J.B.; Rock, C.O. Bacterial lipids: metabolism and membrane homeostasis. *Prog. Lipid Res.* **2013**, *52*,

755   249–76, doi:10.1016/j.plipres.2013.02.002.

756   73.   Storvik, K.A.M.; Foster, P.L. RpoS, the stress response sigma factor, plays a dual role in the regulation of

757   Escherichia coli's error-prone DNA polymerase IV. *J. Bacteriol.* **2010**, *192*, 3639–44, doi:10.1128/JB.00358-10.

758   74.   Starosta, A.L.; Lassak, J.; Jung, K.; Wilson, D.N. The bacterial translation stress response. *FEMS Microbiol. Rev.*

759   **2014**, *38*, 1172–1201, doi:10.1111/1574-6976.12083.

760   75.   Adamson, B.; Norman, T.M.; Jost, M.; Cho, M.Y.; Nuñez, J.K.; Chen, Y.; Villalta, J.E.; Gilbert, L.A.; Horlbeck, M.A.;

761   Hein, M.Y.; et al. A Multiplexed Single-Cell CRISPR Screening Platform Enables Systematic Dissection of the

762   Unfolded Protein Response. *Cell* **2016**, *167*, 1867-1882.e21, doi:10.1016/j.cell.2016.11.048.

76.   Molina-Mora, J.A.; Kop-Montero, M.; Quirós-Fernández, I.; Quiros, S.; Crespo-Mariño, J.L.; Mora-Rodríguez, R.A.

A hybrid mathematical modeling approach of the metabolic fate of a fluorescent sphingolipid analogue to

predict cancer chemosensitivity. *Comput. Biol. Med.* **2018**, *97*, 8–20, doi:10.1016/j.compbiomed.2018.04.008.

LEGENDS OF FIGURES AND LEGENDS

Figure 1. General pipeline for identifying core response genes in *P. aeruginosa* by a machine learning
approach. (A) Schematic representation of hypothesis for identifying core response determinants when
bacteria are exposed to multiple perturbations. (B) The workflow of the machine learning approach using
microarray data and model fitting by SP and MP methods for identifying and describing core response genes.

Figure 2. Data selection and partitioning. (A) Microarray-data selection in GEO-NCBI database according to
experimental design. (B) Samples, including replicates in each experiment -perturbation and control-, were
separated into the training or the testing data sets using the SP (the last replicate of each condition was moved
to the testing dataset) or MP (random separation with multiple arrangements) methods. (C) Representation
of classification procedures for the SP and MP methods, including internal 10-fold cross-validation for the
training data set.

Figure 3. Comparison of samples by global profiles with transcriptomic data of 5549 genes of *P. aeruginosa*.
(A) Dispersion of intensities of samples, showing similar distribution between samples after normalization. (B)
Global profiles of samples were compared by PCA, using all the 5549 genes in the microarray.

Figure 4. **Evaluation of SP method for gene ranking by importance, the case for RF algorithm.** (A) Accuracy,
F1, and kappa values after iterations of classification with the first top 200 genes (adding genes 1-to-1). (B)
ROC plot using selected top 56 genes for evaluation of the performance of the algorithm. (C) Ranking and
importance value of top 56 genes. Similar results are shown for SVM and KNN algorithms in supplementary
Figure S1.

Figure 5. **Evaluation of MP method for gene ranking by frequency, case for SVM algorithm.** (A) Ranking of
top 56 genes by frequency after iterations of 100 data partitions and classification model fitting. (B) Dispersion
of metrics across 100 iterations. Similar results are shown for KNN and RF algorithms in supplementary Figure
S2.

Figure 6. Comparison of 118 candidate genes of the core perturbome. (A) The top 56 genes of each
algorithm were selected and compared using upset plots, showing the intersection of the lists
among classifiers. This comparison resulted in 118 distinct candidate genes. A gene was selected in
the final list if it was found at least by four different algorithms, i.e. the support was ≥ 4. (B) Gene
annotation using KEGG shows the enriched pathways found for the core perturbome genes.

Figure 7. Network analysis and comparison of global transcriptomic profiles using the 46 core perturbome genes. (A) Distribution and support (number of algorithms) of preliminary 118 selected genes on a basal large-scale network of functional associations in *P. aeruginosa.* Topological metrics of the network and identity of hub genes (circles with light-blue border) are included. (B) Small-world network showing relationships between the 46 core response genes and connectivity metric for each gene. (C) The global profile of samples by PCA, showing the separation of conditions. For reference, centroids were plotted as triangles in each cluster. More details are shown in Supplementary Figure S3.

Figure 8. Annotation of core response genes and comparison with a differential expression analysis. (A) General annotation profile of identified genes by core response genes showing associated biological processes. (B) Comparison of identified genes by our machine learning approach and DEGs lists. (C) General annotation of DEGs showing a similar profile to our approach.

Table 1. Annotation of genes of the core perturbome *Pseudomonas aeruginosa*

Supplementary Figures and Tables

Figure S1. **Evaluation of SP method for gene ranking by importance, the case for SVM and KNN algorithms.** (A-C) Accuracy, F1, and kappa values after iterations of classification with the first top 200 genes (adding genes 1-to-1). (B-D) ROC plot using selected top 56 genes for evaluation of the performance of the algorithm. (E) Ranking and importance value of top 56 genes (same importance for both algorithms).

Figure S2. **Evaluation of MP method for gene ranking by frequency, the case for KNN and RF algorithms.** (A) KNN evaluation of ranking of top 56 genes by frequency after iterations of 100 data partitions and classification model fitting and dispersion of metrics across 100 iterations. (B) Same evaluation, the case for RF algorithm.

Figure S3. Systems-level description and gene expression comparison by core response genes. (A) Gene expression and clustering of the 46 core response determinants of *P. aeruginosa*. (B-C) Comparison of gene expression (logR) of kmedoids for each cluster using small-world networks.

Table S1. Studies of datasets included in this work to identify the core perturbome of *Pseudomonas aeruginosa*

Table S2. Metrics associated with the performance of each method and algorithm

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

A

B

Figure 6

A

Topological properties of the core perturbome network:

¬ 46 genes (nodes) and 10 hub genes
¬ Highly connected network
¬ Edges: 90
¬ PPI enrichment p-value: < 1.0e-16
¬ Avg. number of neighbors: 3.9
¬ Clustering coefficient: 0.51
¬ Network centralization: 0.165
¬ Network heterogeinity: 0.738

Support
0 1 2 3 4 5 6

B

Connectivity
11
0

C

Class
Control
Perturbation

Figure 7

A *Core response genes*

Translation 3%
Others 4%
Transport 8%
Metabolism 40%
Biosynthesis 19%
Molecule binding 26%

B

Core response genes

13   33   68

DEGs

C *DEGs*

3% 4%
7%
14%
49%
23%

Figure 8

Figure S1

Figure S2

Figure S3

Table 1. Annotation of genes of the core perturbome of *Pseudomonas aeruginosa*

Support*	Code**	Name	Top annotated functions	DNA damage	DNA repair	Stimulus response	Other stimulus	Differential expression*** (↑ upregulated, ↓downregulated and ↑↓ variable regulation)
6	PA0500	bioB	Biotin biosynthesis, catabolic process, diaminopimelate process	x	x	x	heat	↓AZM and ↑↓COL
	PA1812	mltD	Microcin transport, cytokinesis, peptidoglycan biosynthesis	-	-	x	drug,heat	↓AZM and ↓COL
	PA2647	nuoL	Aerobic respiration, electron/proton transport	x	-	x	response to superoxide, antibiotic	↓NaClO and ↓CIP
	PA2735	-	DNA metabolism, translation, DNA damage	x	x	-		↑Temperature, ↑↓AZM and ↓Cu
	PA2851	efp	Peptidyl-prolyl isomerization, DNA damage, ATP (proton)synthesis	x		x	starvation, heat, antibiotic	↑AZM, ↓↑COL and ↓↑CIP
	PA2967	fabG	Lipid metabolism, fatty acid and lipid biosynthesis	x	-	x	cold, osmotic stress	↓↑AZM, ↓↑COL and ↓Cu
	PA3644	lpxA	Lipid metabolism, fatty acid and peptidoglycan biosynthesis	-	-	x	drug, antibiotic	↓COL, ↓AZM and ↓Cu
	PA4563	rpsT	Cold response, ribosomal assembly, transcription antitermination	-	-	x	stress, antibiotic, heat	↓AZM, ↓Cu and ↓↑COL
	PA4569	ispB	Ubiquinone biosynthesis, transport, aerobic respiration	x	-	x	stress, antibiotic	↓AZM, ↓↑COL and ↓Cu
	PA4645	-	Enzyme-directed rRNA pseudouridine synthesis, L-serin catabolism, glycine biosynthesis	x	x	-		↑AZM and ↓Cu
	PA5016	aceF	Aerobic respiration, electron transport, Krebs cycle	x	-	x	heat, stress	↓AZM, ↓↑CIP, ↓↑COL and ↓Cu
	PA5051	argS	DNA damage, antibiotic response, peptidoglycan biosynthesis	x	-	x	heat	↑Cu, ↓COL and ↓AZM
	PA5300	cycB	DNA damagage recognition and repair	x	x	-		↓AZM, ↓Cu and ↓COL
	PA5479	gltP	DNA damage, transport, fumarate transport	x	-	x	stimulus	↓CIP, ↓Cu and ↓↑COL
	PA5570	rpmH	Ribosomal assembly, translation, transcription DNA dependent	x	-	x	drug, cold, antibiotic	↓CIP, ↓Cu and ↓↑AZM, ↓↑COL
5	PA0003	recF	Response to DNA damage, SOS response, DNA metabolic process	x	x	x	drug, heat, UV	↓Cu and ↓AZM, ↓↑COL
	PA0429	-	Uncharacterized protein	-	-	-		↓Cu and ↓AZM, ↓↑COL
	PA1609	fabB	Fatty acid biosynthesis, lipid metabolism and biosynthesis	x	-	x	drug	↓Cu, ↓CIP, ↓↑AZM and ↓↑COL
	PA2537	-	Phospholipid biosynthesis, wall biogenesis, regulation of endopeptidase	-	-	-		↓Cu and ↑AZM
	PA2641	nuoF	Aerobic respiration, electron/proton transport, Krebs cycle	x	x	x	acidity, oxidative stress,	↓AZM, ↓Cu and ↓COL
	PA2733	-	Movement, pilus assembly, histidine biosynthesis	-	-	-		↓↑COL, ↓Cu and ↑AZM
	PA2950	-	Lipid metabolism, fatty acid and lipid biosynthesis	-	x	x	magnesium starvation	↓↑COL, ↓↑Cu and ↓AZM
	PA4265	tufA	Heat response, transcription antitermination, DNA damage	x	-	x	drug, antibiotic, cold	↑light, ↓↑CIP and ↓AZM
	PA4403	secA	Protein transport, DNA damage, transport	x	-	x	heat, cold, antibiotic,	↓↑COL, ↓Cu, ↓↑CIP and ↓AZM
	PA4428	sspA	DNA transcription, outer membrane assembly, lipopolysaccharide transport	-	-	x	drug, oxidative stress,	↓↑COL, ↓Cu, ↓CIP and ↓↑AZM
	PA4430	-	Response to DNA damage, catabolism, phenylacetate catabolism	x	-	x	heat, oxidative stress	↓Cu, ↓↑AZM and ↓NaClO
	PA5040	pilQ	Movement, pilus assembly, secretion	x	-	x	antibiotic	↓Cu and ↓AZM
	PA5110	fbp	Precursor metabolites and energy, gluconeogenesis, glycolisis	x	-	-		↓↑COL and ↓↑AZM
	PA5174	-	Lipid metabolism, fatty acid and lipid biosynthesis	-	-	-		↓↑COL, ↓AZM and ↓↑Cu
	PA5339	-	DNA damage, nucleobase interconversion, guanosine tetraphosphate metabolism	x	-	-	starvation, oxidative estress	↓↑COL, ↓AZM and ↓Cu
4	PA0002	dnaN	DNA damage response, SOS response, DNA metabolic process	x	x	x	heat, stress, UV, drugs	↓↑COL, ↓AZM, ↓↑CIP and ↓Cu
	PA1610	fabA	Lipid metabolism, fatty acid and lipid biosynthesis	x	-	x	drug	↓↑COL, ↓AZM and ↓Cu
	PA2639	nuoD nuoC	Aerobic respiration , electron/proton transport, DNA damage	x	-	x	heat	↓COL, ↓CIP, ↓AZM, and ↓Cu
	PA2644	nuoI	Aerobic respiration, electron/proton transport	x	-	-		↑↓COL, ↓AZM and ↓Cu
	PA2645	nuoJ	Aerobic respiration, electron/proton transport			x	oxidative stress	↓↑COL and ↓NaClO
	PA2646	nuoK	Aerobic respiration, electron/proton transport	x	x	x	antibiotic, oxidative stress	↓Cu, ↓CIP and ↓AZM and ↓NaClO
	PA2739	pheT	Protein transport, ARNt aminocylation for protein transport	-	-	x	cold, heat	↑↓Cu and ↓AZM
	PA2744	thrS	Heat response, ATP catabolism, cold response	-	-	x	cold, heat, antibiotic	↑COL, ↓Cu and ↓AZM
	PA2965	fabF1	Lipid metabolism, fatty acid and lipid biosynthesis	-	-	x	drug, osmotic stress, Temp,	↓CIP, ↓AZM, and ↓Cu
	PA4461	-	Lipopolysacaridchard transport, outer memebrane assembly, cold response	-	-	-		↓CIP, ↓AZM, ↑↓COL and ↓Cu
	PA4740	pnp	RNA catabolism, transcription antitermination cold response	-	-	x	cold, osmotic stress, heat	↓AZM, ↑↓COL and ↑Cu

PA4847	*accB*	Fatty acid biosynthesis, lipid metabolism and biosynthesis	x	-	x	drug, heat	↓CIP, ↓↑COL, ↓↑AZM and ↓ low oxigen
PA5063	*ubiE*	Ubiquinone biosynthesis, aerobic respiration, DNA-dependent transcription	x	x	x	light, oxidative stress	↑↓AZM, ↑↓COL, ↑↓CIP and ↓Cu
PA5163	*rmlA*	Lipopolysaccharide biosynthesis, DNA damage, antibiotics response	x	x	x	antibiotic	↓AZM, ↓COL, ↓CIP and ↓Cu
PA5337	***rpoZ***	Regulation of transcription, transcription antitermination	x	-	x	stress, heat, cold, antibiotic	↑↓AZM, ↑↓COL, ↑↓CIP and ↓Cu
PA5554	*atpD*	Response to DNA damage, aerobic respiration, amino acid metabolism	x	x	x	heat, cold, drug	↑↓AZM, ↓COL, ↑↓CIP and ↓Cu

* Support: number of algorithms (out of six). A gene was included in the core perturbome if the support was ≥4.

** Bold names refer to hub genes of the core perturbome.

*** Expression patterns found in significant works. See Methods for details of the references.

Table S1. Studies of datasets included in this work to identify the core perturbome of *Pseudomonas aeruginosa*

Perturbation (name, acronym and GEO series ID)	Main results
Azithromycin AZM-a GSE2430	In this study it was demonstrated that Azithromycin affected the metabolic pathways related to quorum sensing, as an antagonist. In addition, the production of virulence factors decreased, the response to oxidative stress also decreased, the mobility of the bacteria was reduced and the ability to form biofilm was impaired. Additionally, a very strong induction of type II secretion systems was observed (Yusuf Nalca, Lothar Jänsch, Florian Bredenbruch, Robert Geffers, Jan Buer, & Susanne Häussler, 2008).
Hydrogen peroxide H2O2 GSE3090	The results obtained in the study reinforce the conclusion that DNA repair pathways and proteins with catalase activity could be within the antioxidant defense systems that play a more relevant role *in P. aeruginosa*. In the experiment, a sublethal dose of H2O2 was used and a reduction in active and facilitated transport was found. Finally, an increase in the transcription of all the F, R and S type pyocins and at the same time a reduction in the transcription of a cellular immunity protein (pyocin S2) was shown (Chang, Small, Toghrol, & Bentley, 2005).
Copper Cu GSE4152	In the study it was observed that bacteria that were exposed to a copper shock showed a response very similar to that of oxidative stress that was not observed in a population adapted to high concentrations of iron. It is suggested that tolerance to iron could be given by two efflux systems: a P-type ATPase and a two-component regulatory system (Teitzel et al., 2006).
Ciprofloxacin CIP GSE5443 (0, 30 and 120 min)	In this study it was found that the genes involved in virtually all metabolic pathways present a negative regulation in response to ciprofloxacin by knocking down lexA gene, regulator of SOS response. The LexA-controlled SOS regulon identified by microarray analysis includes only 15 genes but does include several genes that encode proteins involved in recombination and replication, including two inducible polymerases known to play a role in mutation and the evolution of antibiotic resistance in other organisms (Cirz, O'Neill, Hammond, Head, & Romesberg, 2006).
Sodium hypoclorite NaClO GSE7402	Exposition to sodium hypochlorite showed a negative regulation in the genes related to metabolism and energy. This regulation suggests that the production of energy in the cell is limited. In addition, another strong negative regulation was shown in the genes linked to small molecule transport proteins. Finally, an increase in the transcription of hypothetical enzymes and membrane proteins was shown (Small, Chang, Toghrol, & Bentley, 2007).
Ortho-Phenylphenol OPP GSE10605 (20 and 60 min)	The results of the study show that after the exposure of Ortho-Phenylphenol, bacteria had a negative regulation in the synthesis of multiple amino acids, in particular the synthesis of enzymes linked to lysine biosynthesis was reduced. Among the genes that showed a positive regulation are genes coding for ribosomal proteins and genes coding for virulence factors. Genes related to phospholipid metabolism and transport and acquisition of iron were inhibited (Nde, Jang, Toghrol, & Bentley, 2008).
Azithromycin AZM-b GSE12738	In this study it was found that in the case of concentrations below the minimum inhibitory concentration, no change in the expression of the genes related to the quorum sensing was observed, however a negative change in the genetic expression was observed on N-acyl homoserine lactones (Kai et al., 2009).
Colistin COL GSE13252	*Pseudomonas* sp faced with a suboptimal dose of colistin, an alteration was obtained in the routes associated with virulence factors and bacterial colonization. Differential expression was obtained for genes related to osmotic stress, biofilm formation, mobility and quorum sensing (Cummins, Reen, Baysse, Mooij, & O'Gara, 2009).
Chlorhexidine diacetate CDA GSE14253 (10 and 60 min)	In this study it is shown that in both times 10 and 60 minutes the genes related to membrane transport, oxidative phosphorylation and electron transport presented a negative regulation. After 10 minutes, inhibition was found in the mechanisms of DNA repair and a stimulation of the oprH gene that causes self-stimulation of antimicrobial absorption. At minute 60, inhibition was found in genes related to flagella, pylons, oxidative phosphorylation, electron transport and outer membrane proteins (Nde, Jang, Toghrol, & Bentley, 2009).
E-4-bromo-5-bromomethylene-3-methylfuran-2-5H-one BF8 GSE36753	In the study it is indicated that the use of BF8 can restore the susceptibility to antibiotics of the *Pseudomonas* strains under study. Interestingly, although BF8 is an inhibitor of quorum sensing in gram-negative bacteria, the data in this study suggest that the activities of BF8 to revert antibiotic tolerance of *P. aeruginosa* PAO1 persister cells is not through QS inhibition and may involve other targets (Pan, Bahar, Syed, & Ren, 2012).

Table S2. Metrics associated with the performance of each method and algorithm

Metrics	SP method*			MP method**		
	SVM	KNN	RF	SVM	KNN	RF
Accuracy	0.79	0.71	0.75	0.66	0.69	0.70
Kappa	0.58	0.39	0.45	0.26	0.37	0.35
McnemarPValue	1.00	1.00	0.22	0.67	0.75	0.78
Sensitivity	0.80	0.60	0.50	0.50	0.69	0.56
Specificity	0.79	0.79	0.93	0.76	0.69	0.80
Positive Predictive Value	0.73	0.67	0.83	0.62	0.59	0.64
Negative Predictive Value	0.85	0.73	0.72	0.72	0.80	0.76
Precision	0.73	0.67	0.83	0.62	0.59	0.64
Recall	0.80	0.60	0.50	0.50	0.69	0.56
F1 score	0.76	0.63	0.63	0.52	0.63	0.59
Prevalence	0.42	0.42	0.42	0.38	0.38	0.38
Balanced Accuracy	0.79	0.69	0.71	0.63	0.69	0.68

* Metrics for top 56 genes, averaged after internal cross validation

** Average after 100 iterations of random partitions and internal cross validation