A first *Pseudomonas aeruginosa* perturbome: Identification of core genes

related to multiple perturbations by a machine learning approach

Jose Arturo Molina-Mora[1*], Pablo Montero-Manso[2], Raquel García Batán[1], Rebeca Campos Sánchez[3], Jose

Vilar Fernández[2], Fernando García[1].

[1] Research Center in Tropical Diseases (CIET), University of Costa Rica, Costa Rica

[2] Department of Mathematics, University of A Coruña, Spain

[3] Research Center in Cellular and Molecular Biology, (CIBCM), University of Costa Rica, Costa Rica

* Corresponding author

E-mail: jose.molinamora@ucr.ac.cr

Abstract

Tolerance to stress conditions is vital for organismal survival, including bacteria under specific environmental conditions, antibiotics and other perturbations. Some studies have described common modulation and shared genes during stress response to different types of disturbances (termed as perturbome), leading to the idea of a central control at the molecular level. We implemented a robust machine learning approach to identify and describe genes associated with multiple perturbations or perturbome in a *Pseudomonas aeruginosa* PAO1 model.

Using public transcriptomic data, we evaluated six approaches to rank and select genes: using two methodologies, data single partition (SP method) or multiple partitions (MP method) for training and testing datasets, we evaluated three classification algorithms (SVM Support Vector Machine, KNN K-Nearest neighbor and RF Random Forest). Gene expression patterns and topological features at systems level were include to describe the perturbome elements.

We were able to select and describe 46 core response genes associated to multiple perturbations in *Pseudomonas aeruginosa* PAO1 and it can be considered a first report of the *P. aeruginosa* perturbome. Molecular annotations, patterns in expression levels and topological features in molecular networks revealed biological functions of biosynthesis, binding and metabolism, many of them related to DNA damage repair and aerobic respiration in the context of tolerance to stress. We also discuss different issues related to implemented and assessed algorithms, including normalization analysis, data partitioning, classification approaches and metrics. Altogether, this work offers a different and robust framework to select genes using a machine learning approach.

Keywords: Perturbations, *Pseudomonas aeruginosa*, machine learning, gene selection, perturbome.

Introduction

Cell stress can be defined as a wide range of molecular changes that cells undergo in response to environmental, physical, chemical or biological stressors; sensing and responding to stress is critical for survival (1). These biological functions and metabolic activities are executed through complex physical and regulatory interactions of genes that resemble networks (2). Additionally, tolerance to stress conditions (i.e. stressors, perturbations or disturbances) is vital for organismal survival, including bacteria under diverse environmental conditions, including antibiotics (1).

Several studies have revealed diverse molecular levels that can explain the general response to disturbances in many organisms. However, detailed mechanisms related to responses to perturbations remain poorly understood (3). Available reports usually focus on specific stressors and relatively few studies have focused on common, central and universal determinants affected by multiple perturbations (3). This concept has been recently referred as the perturbome (4,5). For example, in eukaryotic organisms, including plants and human cancer models, some studies have described diverse stress-response genes as common modulators for different types of disturbances, suggesting a central control mechanism (2,4–6). In prokaryotic models, similar findings have been reported for *Escherichia coli* (3,7).

Additionally, comprehensive study of gene-interactions allows the identification of functional relationships among genes (8), their products and the underlying biological phenomena that are critical to understand phenotypes under different biological conditions (9,10). In the context of cell stress, the response to different environmental or experimental stimuli can be recognized by distinct gene expression patterns. This can be inferred from transcriptomic profiling data and functional associations using high throughput molecular technologies such as microarrays or RNASeq (2). However, a challenge with these technologies is the large amount of high complexity data generated. Specialized bioinformatics analysis are required to select relevant information and to reduce noise that distinguishes the molecular determinants for particular biological conditions. Thus, a primary objective of transcriptomic profiling is to find an optimal subset of genes that could be used to characterize and classify unknown samples (11). This gene selection is not obvious and complex due to the thousands of genes to select from (12).

To study the central response determinants to different perturbations in a living organism, we used the model *Pseudomonas aeruginosa* PAO1 (reference strain). *P. aeruginosa* is a Gram negative gamma-proteobacterium with a noteworthy metabolic versatility and adaptability enabling colonization of diverse niches infecting plants, animals and humans alike (8,13). In this group, the molecular mechanisms associated to most biological processes remain unclear causing limited action to modulate responses, including the susceptibility to stressors, environmental factors and experimental conditions.

Several studies have used machine learning algorithms at the transcriptomic level to recognize gene expression patterns (2,14–16). However, for many common biological contexts, applicability and utility of these machine learning approaches have not been fully explored and utilized (17), for example in the exposure to multiple stressors and the molecular response. To our knowledge, only a few studies have used feature selection methods on biological data to describe the effects of multiple perturbations in complex biological systems (4,6) and so far none in *P. aeruginosa*. A related work in *P. aeruginosa* used a machine learning approach to identify sets of genes that correlate with multiple culture media, but without other conditions (18).

The use of microarray and other high throughput technologies data involve challenges for machine learning approaches. These include the curse of dimensionality (19,20), normalization of raw values to compare samples (21,22), data partitions for training and testing models (23,24), and evaluation of performance (21,25). Since comparison between the machine learning algorithms are completely variable (11,17,20,26,27), Support Vector Machine (SVM) (28), K-Nearest neighbor (KNN) (27) and Random Forest (RF) (29) have been successfully used with microarray gene expression data allowing the recognition of emerging patterns (26).

Here, we hypothesize that perturbations on living cells will trigger global reprogramming of multiple molecular determinants that can be sensed at the transcriptional level. The initial response after an acute stress will then expand producing the global molecular rearrangement. Then, pleotropic and specific effects on gene networks will be reflected as changes in gene expression profiles and the complexity of molecular regulation at other levels. Therefore, by using a machine learning approach, common molecular features (for all stressors) could be identified as a central or core determinant (see Fig. 1-A-B).

Fig. 1. General pipeline for identifying core response genes in *P. aeruginosa* by a machine learning approach.
(A) Schematic representation of hypothesis for identifying core response determinants when bacteria are
exposed to multiple perturbations. (B) Workflow of the machine learning approach using microarray data and
model fitting by SP and MP methods for identifying and describing core response genes. (C) Representation
of data partition methods, SP and MP, including subsamples for testing and an internal 10-fold cross validation
for training data set.

Thus, the aims of this study were (i) to implement a machine learning approach to select genes from
microarray expression data, and (ii) to identify and describe a subset of genes than can be associated with
multiple perturbations or perturbome in *P. aeruginosa,* i.e. the core response components.

Materials and methods

Overall methodology of the study is presented in Fig. 1-B. In brief, after a data selection, normalization
and gene selection to define the perturbome were run. To achieve this, we considered two different
procedures to split the data (training and testing datasets for the machine learning approaches: SVM, KNN
and RF): a single partition (SP method) and another with multiple partitions (MP method). Relations between
genes were represented using both large scale and small world networks, and a final comparison between
conditions, an analysis of differential expression and gene annotation were performed.

Datasets

In order to compare gene expression profiles of strain *P. aeruginosa* PAO1 when exposed to multiple
perturbations, GEO database (https://www.ncbi.nlm.nih.gov/geo/) was used for a systematic selection of
datasets (details in supplementary Fig. S1-A). Initial evaluation by organism and mRNA profiles by GPL84
platform (Affymetrix *Pseudomonas aeruginosa* PAO1 Array, with all 5549 protein-coding sequences) identified
156 series of datasets with 1310 samples (Date of Access: January 25th 2018). In a second step, data were
selected according to experimental design if they included perturbations, leaving only 47 series. Finally, to
make datasets comparable by experimental conditions, evaluation and selection were done for series with

similar culture conditions (Luria Bertani LB medium and exponential phase when measuring mRNA profile)

and if a control condition was available (without any perturbation or treatment). The final dataset was

composed of 10 series with 71 samples (Series GSE2430, GSE3090, GSE4152, GSE5443, GSE7402, GSE10605,

GSE12738, GSE13252, GSE14253 and GSE36753).

Some series included temporal measurements which we considered as separate perturbations, resulting

in replicates of 10 controls and 14 perturbations: azithromycin with 2 series (AZM-a and AZM-b) (30,31),

Hydrogen peroxide (H_2O_2) (32), copper (Cu) (33), sodium hypoclorite (NaClO) (34), ortho-phenylphenol (OPP)

at 20 and 60 minutes (35), colistin (COL) (36), chlorhexidine diacetate (CDA) at 10 and 60 minutes (37), E-4-

bromo-5-bromomethylene-3-methylfuran-2-5H-one (BF8) (38) and ciprofloxacin CIP at 0, 30 and 120 minutes

(39).

Normalization and comparison of global transcriptomic profiles

To compare all the 71 samples, a first normalization step using housekeeping genes was performed, as

recommended previously for multiple origin samples (22,40,41). Since *ampC, fabD, proC, pbpA (PBP2), rpoD*

and *rpoS* genes have been reported as stable housekeeping genes for *P. aeruginosa* (21)*,* normalization by

each one of these genes were compared according to effect on the other. After normalization was done using

a specific gene, the other five genes were evaluated by intensity dispersion of controls vrs perturbations and

p values were calculated using T-test ($p < 0.001$). Then, comparison was done by calculating variation

coefficient for control conditions. These metrics were used for selecting the best normalizer gene (assuming

no differences between other housekeeping genes when exposed to stressors).

After normalization, clustering algorithms were implemented in order to compare global transcriptomic

profiles between perturbations and controls. Principal Component Analysis (PCA) and Hierarchical Clustering

(HC) were run with default parameters using the Caret Package (caret.r-forge.r-project.org/) in R software

(Version 3.4.3).

In order to robustly select a number of genes that could separate experimental conditions (controls and

perturbations), reduce noise in microarray data and for finally identifying the core response of *P. aeruginosa*,

two approaches of feature selection protocols were implemented, as detailed below.

Gene ranking and selection by Single Partition SP method

With the aim of identify genes which expression values were commonly related to multiple

perturbations, a first approach was implemented considering a particular partition of dataset for training and

testing sets (Fig. 1-C). Single partition was established using the last replicate of each experiment, in both

control and perturbation. Because there were 14 perturbations and 10 controls (71 samples including

replicates), a total of 24 samples were included in the testing dataset and the remaining 47 samples were

included for the training dataset (66% for training and 34% for testing set).

Using this partition, ranking of genes was done by a machine learning approach using three classification

algorithms: SVM, KNN and RF. A homemade script in R included these functions of the Caret package. For all

three algorithms, default parameters were used for training and 10-fold cross validation (Fig. 1-B) was

included. After this, variable importance metric was calculated for all genes using the *varImp* function,

associating a specific value for each gene. In the case of SVM and KNN, same importance is calculated because

function is model-free for these cases (as detailed in Caret Package), resulting in the same list of genes but

metrics are specifically calculated for each algorithm.

As similarly reported (11), to evaluate the number of genes that should be selected in the top group (the

first K ranked genes, as candidates for the core response by each algorithm), multiple classification models

were systematically run, starting only with the highest-score ranked gene. In brief, after the training with one

gene, model performance was evaluated by calculating metrics using the testing dataset. Then, training was

run again when the next ranked gene was added, and new metrics were calculated. This process was repeated

up to complete all the ranked genes. Metrics included accuracy (correct classification percentage), kappa

value (inter-rater classification agreement), sensitivity, specificity, precision, recall, prevalence and F1 score

(harmonic average of the precision and recall).

Selection of the K value of top genes was based on the following criteria: (i) the stability of the metrics

(priority for accuracy, kappa and F1) when increment of ranked genes was done, as suggested in (11), (ii)

consensus value suitable for all the three algorithms (including a 10% of tolerance), and (iii) minimum number

of genes as possible. After the selection of the K value, ROC (Receiver-operating characteristic) curve and AUC

(Area under the curve) value were calculated for each algorithm. Finally, selection of top K genes between

algorithms were compared by metrics and list of genes (see below for details).

Gene ranking and selection by Multiple Partitions (MP) method

In order to identify genes related to multiple perturbations independently of a single/specific partition,

a second method using multiple random partitions was implemented (Fig. 1-C). To address this, a random

data selection for training and testing sets was done using the *createDataPartition* function. Partition was set

to 80% (57 samples) for training set and remaining for testing set (14 samples) with experimental conditions

equally distributed. Then, protocols with SVM, KNN and RF algorithms (same conditions as previously

described in SP method, with 10-fold cross validation) resumed the analysis with a final ranking of genes using

the *varImp* function. Using only top K of ranked genes (K value determined using the criteria described in the

SP method), new set of training/testing sets were used for evaluating performance of the models and each

metric was stored with the list of the K ranked genes. This full process was automatically repeated 100 times

using *replicate* function, starting with a new random partition and finishing with the list of the K genes and

the metrics associated to that partition. Finally, for each algorithm, full data of all the runs were analyzed for

determining frequency of the appearance of genes (*table* function) and calculating average and dispersion of

metrics across all the 100 runs. Definitive list of the K more frequent genes was established for each algorithm

after this comparison by frequency.

Identification of core response genes

After selection of top K genes in each algorithm by SP and MP methods, comparison of genes was done

using   Venn   diagrams   in   order   to   identify   all   the   candidate   genes   using   Venn-tool

(http://bioinformatics.psb.ugent.be/webtools/Venn/). Genes identified by at least four algorithms were

considered part of the perturbome (this guarantees that a gene was identified by the two methods and at

least by two different classification algorithms).

Gene relationships were represented using molecular networks using a large scale model (using a top-

down systems biology approach). The network was built using protein-protein interaction (PPI) graphs in

PseudomonasNet database (www.inetbio.org/pseudomonasnet/). Network was downloaded and visualized

using the Cytoscape software (https://cytoscape.org/).

Description and comparison of core response genes

In order to describe and compare the genes associated with the core response of *P. aeruginosa,* by

experimental conditions, four analyzes were established. First, clustering analysis by PCA and HC algorithms

were evaluated again but now only considering genes of the core response. Based on distribution in the case

of PCA, an additional analysis to cluster the samples into two groups was done using the Kmedoids algorithm

in R with fastkmed package (https://rdrr.io/cran/kmed/man/fastkmed.html, default parameters). This

allowed not only evaluating the capacity of the core response genes to separate the experimental conditions,

but also to calculate the kmedoids for the groups. Additional representation of centroids was done using

Kmeans algorithm.

Second, using the PseudomonasNet database, a small world network was built and then exported into

Cytoscape software with the genes of the core response. The information of topological features (including

connectivity) and expression levels of kmedoids were incorporated into different versions of the network.

Third, a classic analysis of differentially expressed genes (DEGs, *p<0.05*) was implemented in R with

Limma package (https://www.rdocumentation.org/packages/limma/versions/3.28.14) using empirical Bayes

moderated t-statistic (eBayes) with Benjamini and Hochberg method for *p* value correction (42). This led us

to compare our results with a classical approach for gene expression.

Finally, in order to give biological interpretation to the selected genes, to determine levels of expression

reported in databases and to study metabolic pathways involved under each perturbation, an exhaustive

annotation was made using the databases PseudomonasDB (gene ontology), GEO database and particular

literature. This information was integrated with the results obtained by all the analysis and the DEGs in order

to fully describe the genes that make up the core response or perturbome of *P. aeruginosa* PAO1.

Results

Normalization and comparison of global transcriptomic profiles

In order to compare multi-study microarray data, after data selection (supplementary Fig. S1-A),

normalization was performed using six classical housekeeping genes for *P. aeruginosa*, which were evaluated

according to two metrics (supplementary Table S1). First, after each normalization by a specific housekeeping

gene, intensity dispersion of controls vrs perturbations were compared for other genes and a *p* value was

obtained by T-test. For all genes, comparisons between conditions revealed no statistical differences

(*p*>0.001). A second criterion was the comparison of variation coefficient for control conditions after each

normalization. The lower values of the coefficient were obtained after the normalization with *proC* and with

pbpA. However, *proC* was selected due it has been reported as a more stable housekeeping gene in *P.*

aeruginosa (21). Details of the expression levels (between conditions) of the other housekeeping genes after

proC normalization are shown in supplementary Fig. S1-B. Similarly, average intensity between conditions was

compared for each perturbation after *proC* normalization (Fig. 2-A), showing differences in the expression of

housekeeping genes for some specific disturbances. Comparison of final dispersion of mean intensities

(expression level) of all genes showed similar distribution as expected (Fig. S1-C, case for first replicate of each

condition), making samples comparable.

Global transcriptomic profiles (all 5549 genes) were compared by both PCA and HC clustering algorithms,

revealing a mixed pattern (no separation) between perturbations and controls. Two samples (BF8 and Control

8) resulted with extreme global profiles (Fig. 2 B-C).

Fig. 2. Normalization and comparison of samples by global profiles using all genes. (A) Evaluation of

housekeeping genes after normalization by *proC* gene, comparing intensity between conditions for each

perturbation. Details of gene selection for normalization are shown in Table S1 and Fig. S1. (B-C) Global profiles

were compared by both PCA and HC clustering algorithms, showing mixed patterns between classes. More

details in supplementary Fig. S1.

Perturbome genes of P. aeruginosa *can be identified by a machine learning approach using SP and MP methods*

Two methods using machine learning algorithms (by SP and MP methods) were implemented in order to

robustly rank and select genes associated to multiple perturbations in *P. aeruginosa*. In each method, three

classification algorithms were evaluated: SVM, KNN and RF. Metric results associated to RF are shown in Fig.

3, and supplementary Fig. S2 for SVM and KNN.

Fig. 3. **Evaluation of SP method for gene ranking by importance, case for RF algorithm.** (A) Accuracy, F1 and

kappa values after iterations of classification with the first top 200 genes (adding genes 1-to-1). (B) ROC plot

using selected top 56 genes for evaluation of performance of the algorithm. (C) Ranking and importance value

of top 56 genes. Similar results are shown for SVM and KNN algorithms in supplementary Fig. S2.

The first method (SP) implemented a gene ranking based on variable importance using a single/specific

data partition. After the ranking was established, multiple classification models were run with the ranked

genes (Fig. 3-A and supplementary Fig. S2-A-C). For each classification model, stability of the three metrics

were evaluated to select the suitable K value of genes that could be applied to all algorithms at the same time,

as detailed in criteria of Material and Methods section. For SVM and RF, stable values of metrics are given

with at least the first 51 ranked genes, meanwhile it is 45 genes for KNN. Considering a 10% of tolerance with

the highest of these values, K=56 was selected as the number of top genes that were included as preliminary

candidates of the core response according to each algorithm. With this value, metrics of each algorithm were

compared (Table 1). For example, accuracy was 0.79, 0.71 and 0.75 for SVM, KNN and RF respectively in the

SP method. SVM obtained a better performance according to kappa, sensitivity, recall and F1 scores, but

higher values of specificity and precision resulted for RF. Also, ROC curve and AUC value were calculated (Fig.

3-B and supplementary Fig. S2-B-D). Best performance was obtained for RF with AUC = 0.92, then 0.82 for

SVM and finally 0.76 for KNN. Since *importance* metric for SVM and KNN is the same, they shared same list of

top 56 genes. Comparison between implementations showed that 21 genes were identified by the three

algorithms at same time, 35 exclusively by RF and same number for SVM/KNN. In total 91 genes were identify

by any of the algorithms. List of genes and importance value of top 56 ranked genes for each approach is

presented in Fig. 3-C for RF and Fig. S2-E for SVM/KNN.

Table 1. Performance of six different machine learning approaches to identify perturbation-associated
genes

Metrics	SP method*			MP method**		
	SVM	KNN	RF	SVM	KNN	RF
Accuracy	0.79	0.71	0.75	0.66	0.69	0.70
Kappa	0.58	0.39	0.45	0.26	0.37	0.35
Sensitivity	0.80	0.60	0.50	0.50	0.69	0.56
Specificity	0.79	0.79	0.93	0.76	0.69	0.80
Positive Predictive Value	0.73	0.67	0.83	0.62	0.59	0.64
Negative Predictive Value	0.85	0.73	0.72	0.72	0.80	0.76
Precision	0.73	0.67	0.83	0.62	0.59	0.64
Recall	0.80	0.60	0.50	0.50	0.69	0.56
F1 score	0.76	0.63	0.63	0.52	0.63	0.59
Prevalence	0.42	0.42	0.42	0.38	0.38	0.38
Balanced Accuracy	0.79	0.69	0.71	0.63	0.69	0.68

283 * Metrics for top 56 genes, averaged after internal cross validation
284 ** Average after 100 iterations of random partitions and internal cross validation
285

In a second approach, the MP method was implemented using multiple random partitions. Same SVM,

KNN and RF algorithms were evaluated by running 100 iterations with different partitions and top-56 more

frequent genes for each method were selected. Details of ranking and frequency is shown in Fig. 4-A (SVM)

and supplementary Fig. S3 (KNN and RF) and metrics for all 100 iterations are presented in Table 1, Fig. 4-B

and supplementary Fig. S3. Accuracies for all the models were 0.66, 0.69 and 0.70 for SVM, KNN and RF

respectively. Specific values for kappa, precision, recall and F1 score were found for each algorithm. When

comparison of list of genes was done, 29 genes were identified at same time by all implementations, and 23

by both SVM and KNN.

Fig. 4. **Evaluation of MP method for gene ranking by frequency, case for SVM algorithm.** (A) Ranking of top
56 genes by frequency after iterations of 100 data partitions and classification model fitting. (B) Dispersion of
metrics across 100 iterations. Similar results are shown for KNN and RF algorithms in supplementary Fig. S3.

With the aim of identifying preliminary core response genes in *P. aeruginosa*, the lists of top 56 genes

selected by each algorithm and method were jointly represented using a Venn diagram (Fig. 5-B). Note that

the same set was used for SVM and KNN since both procedures lead to the same genes. A total of 118 different

genes were identified and 15 genes were simultaneously identified by all the algorithms. Distribution of all

118 genes on large scale molecular networks is presented in Fig. 5-A. Results show that selected genes are

connected but they do not establish a defined cluster. These genes seem to be associated to different

subgroups of highly connected genes.

Fig. 5. Identification, systems level description and global profiles comparison given by core response genes.
(A) Distribution and number of algorithms of preliminary 118 selected genes on a basal large scale network of
functional associations in *P. aeruginosa.* (B) Set comparison of genes given by classification algorithms of SP
and MP methods, with a final selection of 46 genes which were identified by at least 4 algorithms. (C) Small
world network showing relationships between the 46 core response genes and connectivity metric for each
gene. (D-E) Global profile comparison of samples by both PCA and HC clustering algorithms, showing
separation of conditions. For reference, centroids of each cluster were plotted as triangles in each cluster.
More details are shown in supplementary Fig. S4.

Final version of the core perturbome components was established by selecting genes recognized for at

least four algorithms, according to the criteria detailed in Materials and Methods. A total of 46 genes were

finally associated to core response. In the representation as small world network, relationships between the

46 genes revealed different topological patterns of connectivity between molecules, being *nuoC* and *nuoF* the

ones with higher connectivity (connection degree). In addition, only six genes had no connections between

them (Fig. 5-C).

Comparisons of core response genes show separation of global transcriptomic profiles according to

experimental conditions and biological functions related to tolerance to stress

After the selection of the core response genes, comparisons between controls and perturbations were

done to characterize these genes by global profiles. First, clustering algorithms to compare global profiles

were run using the 46 genes of the perturbome. In the case of PCA (Fig. 5-D), samples distribution let to

differentiate controls and perturbations. A Kmedoid algorithm (k = 2) was able to identify two clusters

enriched by samples of each condition: one consisting entirely of perturbation samples (11 samples, blue

color) and the other mostly by control samples (10 controls and 3 disturbances, red color). In the case of the

blue cluster, the kmedoid was sample CIP-120min, meanwhile control 6 was selected for the another cluster.

For the case of HC (Fig. 5-E), the same distribution of samples was obtained. Supplementary analysis of

gene expression was included by comparing levels for all core response genes (Fig. S4-A) and comparing

expression levels of the kmedoids on the small world network (Fig. S4-B-C).

Gene annotation revealed that biological processes related to most of the genes are metabolism,

molecule binding and biosynthesis (Fig. 6-A). Specific information about participation in processes of DNA

damage response, DNA repairing and response to general/specific stimuli was also searched for each gene,

showing that most of genes includes participation in such processes. Literature support shows variable

patterns of expression, depending on the disturbance as shown in supplementary Table S2.

Fig. 6. Annotation of core response genes and comparison with a differential expression analysis. (A)
General annotation profile of identified genes by core response genes showing associated biological
processes. (B) Comparison of identified genes by our machine learning approach and DEGs lists. (C) General
annotation of DEGs showing similar profile than our approach. Specific annotation per gene is shown in
supplementary Table S2.

Finally, in order to compare the results of the machine learning strategy with another approach, a

differential expression analysis was run. A total of 101 DEGs were identified, which 33 were shared by the

core response (Fig. 6-B). This means that 72% of core response genes were also identified by another single

and independent method. Annotation profile of DEGs (Fig. 6-C) showed a similar pattern as our machine

learning approach.

Discussion

358   Living organisms face external and internal conditions that compromise cellular functions at molecular,

359  metabolic and structural levels, disrupting their homeostasis (43,44). Cell stress response is crucial for

360  organismal survival and complex networks are usually involved in the molecular mechanism related to

361  tolerance (1). However, few studies have identified central and possible universal regulation of the response

362  to multiple disturbances, a concept termed as perturbome (4,5). Common molecular response was previously

363  reported as a network of common set of genes and pathways that can be generically associated to multiple

364  perturbations in plants (7), pathogenic bacteria (3) or cell lines models (4), and others.

365   In our approach using *P. aeruginosa,* but applicable to other organisms, we hypothesized the existence

366  of a set of core genes regulating the response to stressors in a generic sense of different pathways. *P.*

367  *aeruginosa* has a high proportion (about 5%) of its genome dedicated to regulatory mechanisms, probably

368  explaining its adaptability to such a broad range of growth conditions (25). Since strain *P. aeruginosa* PAO1 is

369  a clinical isolate with a profile of multiresistance to many antibiotics (45), characterization of molecular

370  mechanisms involved in the tolerance to stressors in this strain could eventually help to modulate sensitivity

371  and overcome resistance. Exhaustive integration of multi-omic data and network analysis are required in

372  order to clarify the molecular mechanisms related to stress conditions and eventually use them for modulating

373  cell response.

375  *Insights of algorithms to identify core response genes or perturbome*

376   In our study, a first analysis to compare data was done by normalizing intensities using *proC*

377  housekeeping gene. This approach is a prerequisite in qRT-PCR (22), but as reported and exhaustively

378  evaluated in (22), it is also suitable for microarray data, in particular to compare multiple origin samples with

379  divergent sample processing steps (21,25). After normalization, global transcriptomic profiles showed mixed

380  patterns between samples according to their experimental condition. Because its complexity, raw microarray

381  data showed noise and redundant information that can explain the poor resolution of this clustering

382  algorithms to identify classes (20). Thus, a feature (gene) selection analysis was implemented for not only

identifying genes associated to multiple perturbations in *P. aeruginosa,* but also to improve performance of

predictive/descriptive models capable of separate control and perturbation categories.

In our case, these potential patterns were investigated using a robust machine learning approach by

implementing six protocols, using SP and MP methods and SVM, KNN and RF algorithms in each case. This was

a critical step because gene selection from microarray data is complex *per se*. Feature reduction remains as a

challenging task in transcriptomic studies because thousands of genes to select from, and it introduces an

additional layer of complexity in the modelling task (12,46). To avoid bias and overfitting, implementations

of diverse strategies of data partitioning such as bootstrapping, random partitions and cross validation are

recommended. In general, these methodologies can robustly minorate the influence from noise, outliers,

absence of ground truth sets, and to reduce variance (2,24,47).

The single partition SP method consisted of a particular and invariable data for training (with internal

cross validation) and another to test, and it is probably the most common approach used in machine learning.

In the case of the multiple partitions MP method, 100 random partitions of dataset were run. MP method had

a dual consideration when splitting data (multiple partitions and the internal cross validation). This method

can be considered as an *ensemble based on different data partitioning*, as it had been previously proposed

(23). Datasets were divided using multiple random partitioning procedures and then genes were ranked. After

all runs, a final feature subset is determined by calculating the frequency of features in all the runs (23). A

equivalent approach was implemented by Pai and collaborators to classify gene expression data in a cancer

model (48).

However, one possible problem with MP approach is that cross validation results may depend on

similarity of testing and training sets. A classification prediction method is only expected to learn how to

predict on unseen samples that are drawn from the same distribution as training samples (24,47), and MP

method could violate this assumption. SP method guarantees this because was built always using a replicate

for each perturbation and control. Thus, both methods SP and MP are required to robustly select features.

On the other hand, many algorithms for dimension reduction have been proposed (6,19,20) but no

standard machine learning algorithm can be selected due multiple evaluations results on completely variable

metrics associated to performance (26). Many studies have shown that SVM, KNN, and RF generally

outperform other traditional supervised classifiers (17,26,49). In our case, a variable pattern was found for

different metrics in all evaluated implementations. For example, based on accuracy SVM for SP method and

RF for MP method resulted in higher scores, which agree with other studies using machine learning and

biological data (26,48).

In our subsequent analysis, we were interested in identifying a consensus list of candidate genes for the

core response in *P. aeruginosa*, resulting in 46 genes of the perturbome. When global profiles were compared

using these genes (PCA and HC), control and perturbation classes (Fig. 5-D) were clearly separated. This gene

number seems to be a modest number of elements (less than 1% of all the available dataset with 5549 genes)

but it agrees completely with other studies, including machine learning methods (11,17) or other approaches

(11,22,50,51). In addition, 72% of core response genes were also identified as DEGs with similar annotation

profiles; differences can be explained by significant fluctuations in the differential expression results as

previously reported, mainly because it is not a consensus strategy (only based on *p*-value) and it does not

incorporate the estimates of the test performance (true positive/negative rates and other metrics) on the

results (2).

Biological insights of the core response genes in P. aeruginosa: the perturbome

Core response genes or perturbome can be related to a central regulation network, and as convergent

point of signal transduction, transcriptional regulation and stress-related pathways, as it has been suggested

(2,4,5,44). Annotation of the 46 genes shows that most of them are functionally related to biosynthesis,

molecule binding and metabolism (including an important number of hits for lipids), including additional

functions for regulation of DNA damage repair, response to stimuli and aerobic respiration. Interestingly,

these processes are represented by genes with high connectivity in the small world network. For example,

main functions associated to *fabA* are lipid metabolism, fatty acid and lipid biosynthesis, meanwhile for *nuoC*

are aerobic respiration, electron/proton transport and DNA damage. Finally, *nuoF* is associated to aerobic

respiration, electron/proton transport and Krebs cycle.

In this sense, cells are equipped with systems and mechanisms to recover from the environmental stress

and stimuli to maintain all necessary physiological functions (52). Other common stimuli such as low ATP, slow

growth, and ROS production can also occur before cells express stress specific factors, but mediating a

common effect. For example, response to stress includes modulation of energy production and aerobic

electron transfer chain components. As it has been reported in *E. coli*, aerobic electron transport chain

components are down-regulated in response to growth arrest (53). This corresponds with the global profiles

of expression of 46 core response genes. Also, regulation of lipid metabolism is relevant for survival in the

wide range of environmental conditions where bacteria thrive (54), even for biofilm-living forms (3) as *P.*

aeruginosa. Core response genes *fabG*, *lpxA*, and PA5174 could be implied in this process.

In the case of DNA damage repair (including the case of *cycB* and *gltP* genes), responses mediated by

SOS and *rpoS* help to maintain genome integrity, colonization, and virulence (39,55). These responses are

activated under multiple disturbances and modulating a low energy production and shutdown of the

metabolism, promotes formation of antibiotic resistance and biofilms (3). Other related pathways for some

specific genes included regulation of the transcription during stress by RNA-binding proteins in order to

reprogram or shut down translation and to rescue the ribosome stalled by a variety of mechanisms induced

(56). Three core response genes (*rpmH*, *tsf* and PA2735) were annotated with these functions.

Jointly, the relatively few diversity of metabolic functions and pathways has sense in order to ensure

redundancy and robustness in the response to stimuli. Similar results, regarding enriched pathways, have

been obtained in other studies with eukaryotic models, including disturbed human cell lines (4,5,44),

Arabidopsis thaliana under physical and genotoxic stresses (2) or a genome-wide association study of a generic

response to stress conditions (6). In the case of prokaryotic organisms, two studies have used *Escherichia coli*

as model to identify differentially expressed genes after exposure to stress conditions (3) and to create

networks associated with the response to fluctuating environments (7). Differences with other organisms and

disturbances can suggest that response cell stress can be organismal specific, although heterogeneity has also

been suggested a reasonable explanation because differences in the response in a apparently homogeneous

cell population (44,57).

As in our case, the response to stimuli and stressors is orchestrated by a pleotropic modulation (58)

which can be associated to a central regulation. Alternative mechanisms such as cross stress protection (ability

of a stress condition to provide protection against other stressors) (7), role of sigma factors and specific two

component systems (3) can contribute to explain this phenomenon. The molecular response can lead to

regulate multiple biological activities including metabolism, replication, transcription and translation, changes

in membrane composition, motility, modification gene expression, expression of virulence factors, multi-drug

resistant phenotypes and biofilm formation, and others (44).

Taking all together, results of our study suggest that identification of core response genes associated to

multiple perturbations or perturbome in *P. aeruginosa* can define a central network available to modulate a

basic response that includes biological functions such as biosynthesis, binding and metabolism, many of them

related to DNA damage repair and aerobic respiration. To our knowledge, this study can be considered a first

report of the *P. aeruginosa* perturbome.

Further analyses are required to explore potential use of perturbome network to modulate (positively

or negatively) the response to disturbances, to model molecular circuits, to identify possible biomarker genes

of stressed states, and to experimentally validate our findings. In addition, this approach can be used to model

the perturbome in other *P. aeruginosa* strains, as we hope to run soon with a genome we recently described

(13), and other organisms.

Conclusions

A robust machine learning approach was implemented in order to identify and describe core response

genes to multiple perturbations in *P. aeruginosa*. Using public microarray data, two independent partition

strategies (single and multiple with SP and MP methods respectively) and three classification algorithms, we

were able to identify 46 perturbome elements. Both network analysis and functional annotations of these

genes showed coordinated modulation of biological processes in response to multiple perturbations, including

metabolism, biosynthesis and molecule binding, associated to DNA damage repairing, and aerobic respiration,

all probably related to tolerance to stressors, growth arrest and molecular regulation. We also discussed

different issues related to implemented and assess algorithms of normalization analysis, data partitioning,

classification approaches and metrics.

Abbreviations

491 AUC: Area under the curve

492 AZM: Azithromycin

493 B8F: E-4-bromo-5-bromomethylene-3-methylfuran-2-5H-one

494 CDA: Chlorhexidine diacetate

495 CIP: Ciprofloxacin

496 COL: Colistin

497 Cu: Copper

498 DEGs: Differentially expressed genes

499 DNA: Deoxyribonucleic acid

500 GEO: Gene Expression Omnibus

501 H2O2: Hydrogen peroxide

502 HC: Hierarchical Clustering

503 KNN: K-Nearest Neighbor

504 LB: Luria Bertani

505 mRNA: Messenger RNA

506 NaClO: Sodium hypoclorite

507 OPP: Ortho-phenylphenol

508 PCA: Principal Component Analysis

509 PPI: Protein-protein interaction

510 RF: Random Forest

511 RNA: Ribonucleic acid

512 ROC: Receiver-operating characteristic

513 SVM: Support Vector Machine

Declarations

Ethics approval and consent to participate

Not Applicable

Availability of data and material

The datasets generated and/or analysed during the current study are available in:

Public raw data used in this study: GEO database (https://www.ncbi.nlm.nih.gov/geo/), data Series GSE2430,

GSE3090, GSE4152, GSE5443, GSE7402, GSE10605, GSE12738, GSE13252, GSE14253 and GSE36753.

Normalized data and R Scripts: https://github.com/josemolina6/CoreResponsePae

Competing interests

No competing interests to declare.

Funding

This work was funded by projects B8114 Definición de la red transcripcional y de las alteraciones genómicas

inducidas por la ciprofloxacina en *Pseudomonas aeruginosa* AG1 and B8152 proNGS 1.0: Implementación y

evaluación de protocolos de análisis de datos de tecnologías NGS y afines para el estudio de sistemas

biológicos complejos, Universidad de Costa Rica (period 2017-2020), Costa Rica.

Authors' contributions

JMM and FGS participated in the conception, design of the study and data acquisition. JMM, PMM, RGB and

JVF participated in data analysis and interpretation. JMM, RCS, JVF and FGS participated in the interpretation

of the data analysis. JMM drafted the manuscript and all authors were involved in its revision. All authors read

and approved the final manuscript.

Acknowledgements

We thank students Iosif Forero Trelles, Daniel Solano Alvarado and Daniela Aguilar Orozco for their collaborations in different activities of the project.

References

1. DeLong EF. Prokaryotes : prokaryotic physiology and biochemistry. Springer; 2012.

2. Ma C, Xin M, Feldmann KA, Wang X. Machine Learning-Based Differential Network Analysis: A Study of Stress-Responsive Transcriptomes in Arabidopsis. Plant Cell [Internet]. 2014;26(2):520–37. Available from: http://www.plantcell.org/cgi/doi/10.1105/tpc.113.121913

3. Nagar SD, Aggarwal B, Joon S, Bhatnagar R, Bhatnagar S. A Network Biology Approach to Decipher Stress Response in Bacteria Using *Escherichia coli* As a Model. Omi A J Integr Biol [Internet]. 2016 May [cited 2018 Nov 16];20(5):310–24. Available from: http://www.ncbi.nlm.nih.gov/pubmed/27195968

4. Caldera M, Müller F, Kaltenbrunner I, Licciardello MP, Lardeau CH, Kubicek S, et al. Mapping the perturbome network of cellular perturbations. Nat Commun [Internet]. 2019;10(1). Available from: http://dx.doi.org/10.1038/s41467-019-13058-9

5. Sadeh S, Clopath C. Theory of Neuronal Perturbome: Linking Connectivity to Coding via Perturbations. bioRxiv [Internet]. 2020 Feb 20 [cited 2020 Apr 29];2020.02.20.954222. Available from: https://www.biorxiv.org/content/10.1101/2020.02.20.954222v1.full

6. Bermingham ML, Pong-Wong R, Spiliopoulou A, Hayward C, Rudan I, Campbell H, et al. Application of high-dimensional feature selection: Evaluation for genomic prediction in man. Sci Rep [Internet]. 2015;5:1–12. Available from: http://dx.doi.org/10.1038/srep10312

7. Dragosits M, Mozhayskiy V, Quinones-Soto S, Park J, Tagkopoulos I. Evolutionary potential, cross-stress behavior and the genetic basis of acquired stress resistance in Escherichia coli. Mol Syst Biol [Internet]. 2014 Apr 28 [cited 2018 Nov 17];9(1):643–643. Available from: http://www.ncbi.nlm.nih.gov/pubmed/23385483

8. Nogales J, Guðmundsson S, Duque E, Ramos JL, Palsson B∅. Expanding the computable reactome in Pseudomonas putida reveals metabolic cycles providing robustness. bioRxiv [Internet]. 2017;139121. Available from: https://www.biorxiv.org/content/early/2017/05/19/139121

9.   Kc K, Li R, Cui F, Haake AR. GNE: A deep learning framework for gene network inference by aggregating

biological information. Bioinformatics [Internet]. 2018;1–9. Available from: https://github.com/kckishan/GNE

10.   Szklarczyk D, Gable AL, Lyon D, Junge A, Wyder S, Huerta-Cepas J, et al. STRING v11: protein–protein association

networks with increased coverage, supporting functional discovery in genome-wide experimental datasets.

Nucleic Acids Res [Internet]. 2019 Jan 8 [cited 2019 Nov 13];47(D1):D607–13. Available from:

https://academic.oup.com/nar/article/47/D1/D607/5198476

11.   Li Y, Wang N, Perkins EJ, Zhang C, Gong P. Identification and optimization of classifier genes from multi-class

earthworm microarray dataset. PLoS One. 2010;5(10):1–9.

12.   Saeys Y, Inza I, Larranaga P. A review of feature selection techniques in bioinformatics. Bioinformatics

[Internet]. 2007 Oct 1 [cited 2018 Nov 16];23(19):2507–17. Available from:

https://academic.oup.com/bioinformatics/article-lookup/doi/10.1093/bioinformatics/btm344

13.   Molina-Mora J-A, Campos-Sánchez R, Rodríguez C, Shi L, García F. High quality 3C de novo assembly and

annotation of a multidrug resistant ST-111 Pseudomonas aeruginosa genome: Benchmark of hybrid and non-

hybrid assemblers. Sci Rep [Internet]. 2020 Dec 29 [cited 2020 Jan 29];10(1):1392. Available from:

http://www.nature.com/articles/s41598-020-58319-6

14.   Zhao W, Chen JJ, Perkins R, Wang Y, Liu Z, Hong H, et al. A novel procedure on next generation sequencing data

analysis using text mining algorithm. BMC Bioinformatics [Internet]. 2016;17(1):213. Available from:

http://bmcbioinformatics.biomedcentral.com/articles/10.1186/s12859-016-1075-9

15.   Cornforth DM, Dees JL, Ibberson CB, Huse HK, Mathiesen IH, Kirketerp-Møller K, et al. Pseudomonas aeruginosa

transcriptome during human infection. Proc Natl Acad Sci U S A. 2018;115(22).

16.   Glaab E, Bacardit J, Garibaldi JM, Krasnogor N. Using rule-based machine learning for candidate disease gene

prioritization and sample classification of cancer gene expression data. PLoS One. 2012;7(7).

17.   Leung RKK, Wang Y, Ma RCW, Luk AOY, Lam V, Ng M, et al. Using a multi-staged strategy based on machine

learning and mathematical modeling to predict genotype-phenotype risk patterns in diabetic kidney disease: A

prospective case-control cohort analysis. BMC Nephrol [Internet]. 2013;14(1):1. Available from: BMC

Nephrology

18.   Tan J, Doing G, Lewis KA, Price CE, Chen KM, Cady KC, et al. Unsupervised Extraction of Stable Expression

Signatures from Public Compendia with an Ensemble of Neural Networks. Cell Syst [Internet]. 2017 Jul [cited

2020 Apr 30];5(1):63–71.e6. Available from: https://linkinghub.elsevier.com/retrieve/pii/S2405471217302314

19.   Kar SC. Comparing Prediction Accuracy for Supervised Techniques in Gene Expression Data. Math Theory

Model. 2014;4(6):108–16.

20.   Raza K, Hasan A. A Comprehensive Evaluation of Machine Learning Techniques for Cancer Class Prediction

Based on Microarray Data. Int J Bioinform Res Appl [Internet]. 2015;11(5):397–416. Available from:

http://arxiv.org/abs/1307.7050%0Ahttp://arxiv.org/abs/1307.7050%5Cnhttp://www.inderscienceonline.com/d

oi/abs/10.1504/IJBRA.2015.071940

21.   Savli H, Karadenizli A, Kolayli F, Gundes S, Ozbek U, Vahaboglu H. Expression stability of six housekeeping genes:

a proposal for resistance gene quantification studies of Pseudomonas aeruginosa by real-time quantitative RT-

PCR. J Med Microbiol [Internet]. 2003 May 1 [cited 2018 Jan 31];52(5):403–8. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/12721316

22.   Casares FM. A Simple Method for Optimization of Reference Gene Identification and Normalization in DNA

Microarray Analysis. Med Sci Monit Basic Res [Internet]. 2016 Apr 28 [cited 2018 Nov 10];22:45–52. Available

from: http://www.ncbi.nlm.nih.gov/pubmed/27122237

23.   Yang P, Zhou BB, Yang JY-H, Zomaya AY. Stability of feature selection algorithms and ensemble feature selection

methods in bioinformatics. Biol Knowl Discov Handb Preprocessing, mining, postprocessing Biol data.

2013;(January 2014):333–52.

24.   Tabe-Bordbar S, Emad A, Zhao SD, Sinha S. A closer look at cross-validation for assessing the accuracy of gene

regulatory networks and models. Sci Rep [Internet]. 2018;8(1):1–11. Available from:

http://dx.doi.org/10.1038/s41598-018-24937-4

25.   Alqarni B, Colley B, Klebensberger J, McDougald D, Rice SA. Expression stability of 13 housekeeping genes

during carbon starvation of Pseudomonas aeruginosa. J Microbiol Methods [Internet]. 2016;127:182–7.

Available from: http://dx.doi.org/10.1016/j.mimet.2016.06.008

26.   Noi PT, Kappas M. Comparison of random forest, k-nearest neighbor, and support vector machine classifiers for

land cover classification using sentinel-2 imagery. Sensors (Switzerland). 2018;18(1).

27.   Li L, Weinberg CR, Darden TA, Pedersen LG. Gene selection for sample classification based on gene expression

data: study of sensitivity to choice of parameters of the GA/KNN method. Bioinformatics [Internet]. 2001 Dec

[cited 2018 Nov 17];17(12):1131–42. Available from: http://www.ncbi.nlm.nih.gov/pubmed/11751221

28.   Vapnik VN, Vladimir. Estimation of dependences based on empirical data [Internet]. Springer-Verlag; 1982

[cited 2018 Nov 17]. 399 p. Available from: https://dl.acm.org/citation.cfm?id=1098680

29.   Breiman L. Random Forests. Mach Learn [Internet]. 2001 [cited 2018 Nov 17];45(1):5–32. Available from:

http://link.springer.com/10.1023/A:1010933404324

30.   Yusuf Nalca, Lothar Jänsch, Florian Bredenbruch, Robert Geffers JB, Susanne Häussler. Quorum-sensing

antagonistic activities of azithromycin in Pseudomonas aeruginosa PAO1: a global approach. Antimicrob Agents

Chemother. 2008;50(5):1680–8.

31.   Kai T, Tateda K, Kimura S, Ishii Y, Ito H, Yoshida H, et al. A low concentration of azithromycin inhibits the mRNA

expression of N-acyl homoserine lactone synthesis enzymes, upstream of lasI or rhlI, in Pseudomonas

aeruginosa. Pulm Pharmacol Ther [Internet]. 2009 Dec [cited 2018 Nov 13];22(6):483–6. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/19393329

32.   Chang W, Small DA, Toghrol F, Bentley WE. Microarray analysis of Pseudomonas aeruginosa reveals induction of

pyocin genes in response to hydrogen peroxide. BMC Genomics. 2005;6:1–14.

33.   Teitzel GM, Geddie A, De Long SK, Kirisits MJ, Whiteley M, Parsek MR. Survival and Growth in the Presence of

Elevated Copper: Transcriptional Profiling of Copper-Stressed Pseudomonas aeruginosa. J Bacteriol [Internet].

2006 Oct 15 [cited 2018 Nov 13];188(20):7242–56. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/17015663

34.   Small DA, Chang W, Toghrol F, Bentley WE. Comparative global transcription analysis of sodium hypochlorite,

peracetic acid, and hydrogen peroxide on Pseudomonas aeruginosa. Appl Microbiol Biotechnol.

2007;76(5):1093–105.

35.   Nde CW, Jang H-J, Toghrol F, Bentley WE. Toxicogenomic response of Pseudomonas aeruginosa to ortho-

phenylphenol. BMC Genomics. 2008 Oct;9(1):473.

36.   Cummins J, Reen FJ, Baysse C, Mooij MJ, O'Gara F. Subinhibitory concentrations of the cationic antimicrobial

peptide colistin induce the pseudomonas quinolone signal in Pseudomonas aeruginosa. Microbiology. 2009

Sep;155(9):2826–37.

37.   Nde CW, Jang HJ, Toghrol F, Bentley WE. Global transcriptomic response of Pseudomonas aeruginosa to

chlorhexidine diacetate. Environ Sci Technol. 2009;43(21):8406–15.

38.   Pan J, Bahar AA, Syed H, Ren D. Reverting Antibiotic Tolerance of Pseudomonas aeruginosa PAO1 Persister Cells

by (Z)-4-bromo-5-(bromomethylene)-3-methylfuran-2(5H)-one. PLoS One. 2012;7(9).

39.   Cirz RT, O'Neill BM, Hammond JA, Head SR, Romesberg FE. Defining the Pseudomonas aeruginosa SOS response

and its role in the global response to the antibiotic ciprofloxacin. J Bacteriol. 2006 Oct;188(20):7101–10.

40.   Tseng GC. Issues in cDNA microarray analysis: quality filtering, channel normalization, models of variations and

assessment of gene effects. Nucleic Acids Res [Internet]. 2001;29(12):2549–57. Available from:

https://academic.oup.com/nar/article-lookup/doi/10.1093/nar/29.12.2549

41.   Mar J. Applying Data-Driven Normalization Strategies for qPCR Data Using Bioconductor Preprocessing Raw

qPCR Data Normalizing qPCR Data. 2010;1–8.

42.   Smyth GK. Linear Models and Empirical Bayes Methods for Assessing Differential Expression in Microarray

Experiments. Stat Appl Genet Mol Biol [Internet]. 2004 Jan 12 [cited 2019 Apr 29];3(1):1–25. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/16646809

43.   Richter K, Haslbeck M, Buchner J. The Heat Shock Response: Life on the Verge of Death. Mol Cell [Internet].

2010 Oct 22 [cited 2018 Nov 16];40(2):253–66. Available from:

http://www.ncbi.nlm.nih.gov/pubmed/20965420

44.   Vihervaara A, Duarte FM, Lis JT. Molecular mechanisms driving transcriptional stress responses. Nat Rev Genet

[Internet]. 2018;19(6):385–97. Available from: http://dx.doi.org/10.1038/s41576-018-0001-6

45.   Holloway BW. Genetic Recombination in Pseudomonas aeruginosa. Microbiology [Internet]. 1955 Dec 1 [cited

2018 Nov 14];13(3):572–81. Available from: http://www.ncbi.nlm.nih.gov/pubmed/13278508

46.   Piao J, Sun J, Yang Y, Jin T, Chen L, Lin Z. Target gene screening and evaluation of prognostic values in non-small

cell lung cancers by bioinformatics analysis. Gene [Internet]. 2018 Mar 20 [cited 2018 Nov 6];647(1):306–11.

Available from: http://www.ncbi.nlm.nih.gov/pubmed/29305979

47.   Touw WG, Bayjanov JR, Overmars L, Backus L, Boekhorst J, Wels M, et al. Data mining in the Life Sciences with

Random Forest: a walk in the park or lost in the jungle? Brief Bioinform [Internet]. 2013 May 1 [cited 2018 Nov

18];14(3):315–26. Available from: http://www.ncbi.nlm.nih.gov/pubmed/22786785

48.   Pai S, Hui S, Isserlin R, Shah MA, Kaka H, Bader GD. netDx: interpretable patient classification using integrated

patient similarity networks. Mol Syst Biol [Internet]. 2019 Mar 14 [cited 2019 May 1];15(3):e8497. Available

from: http://www.ncbi.nlm.nih.gov/pubmed/30872331

49.   Park H, Shimamura T, Imoto S, Miyano S. Adaptive NetworkProfiler for Identifying Cancer Characteristic-Specific

Gene Regulatory Networks. J Comput Biol [Internet]. 2017;25(2):cmb.2017.0120. Available from:

http://online.liebertpub.com/doi/10.1089/cmb.2017.0120

50.   Falciani F, Diab A, Sabine V, Williams T, Ortega F, George S, et al. Hepatic transcriptomic profiles of European

flounder (Platichthys flesus) from field sites and computational approaches to predict site from stress gene

responses following exposure to model toxicants. Aquat Toxicol [Internet]. 2008 Nov 11 [cited 2018 Nov

18];90(2):92–101. Available from: http://www.ncbi.nlm.nih.gov/pubmed/18823667

51.   Nota B, Verweij RA, Molenaar D, Ylstra B, van Straalen NM, Roelofs D. Gene Expression Analysis Reveals a Gene

Set Discriminatory to Different Metals in Soil. Toxicol Sci [Internet]. 2010 May [cited 2018 Nov 18];115(1):34–

40. Available from: http://www.ncbi.nlm.nih.gov/pubmed/20133373

52. Krämer R. Bacterial stimulus perception and signal transduction: Response to osmotic stress. Chem Rec [Internet]. 2010 Aug 16 [cited 2018 Nov 23];10(4):217–29. Available from: http://www.ncbi.nlm.nih.gov/pubmed/20607761

53. Schurig-Briccio LA, Farías RN, Rodríguez-Montelongo L, Rintoul MR, Rapisarda VA. Protection against oxidative stress in Escherichia coli stationary phase by a phosphate concentration-dependent genes expression. Arch Biochem Biophys [Internet]. 2009 Mar 1 [cited 2018 Nov 22];483(1):106–10. Available from: http://www.ncbi.nlm.nih.gov/pubmed/19138658

54. Parsons JB, Rock CO. Bacterial lipids: metabolism and membrane homeostasis. Prog Lipid Res [Internet]. 2013 Jul [cited 2018 Nov 22];52(3):249–76. Available from: http://www.ncbi.nlm.nih.gov/pubmed/23500459

55. Storvik KAM, Foster PL. RpoS, the stress response sigma factor, plays a dual role in the regulation of Escherichia coli's error-prone DNA polymerase IV. J Bacteriol [Internet]. 2010 Jul [cited 2018 Nov 22];192(14):3639–44. Available from: http://www.ncbi.nlm.nih.gov/pubmed/20472798

56. Starosta AL, Lassak J, Jung K, Wilson DN. The bacterial translation stress response. FEMS Microbiol Rev [Internet]. 2014 Nov [cited 2018 Nov 22];38(6):1172–201. Available from: http://www.ncbi.nlm.nih.gov/pubmed/25135187

57. Adamson B, Norman TM, Jost M, Cho MY, Nuñez JK, Chen Y, et al. A Multiplexed Single-Cell CRISPR Screening Platform Enables Systematic Dissection of the Unfolded Protein Response. Cell [Internet]. 2016 Dec 15 [cited 2018 Nov 16];167(7):1867–1882.e21. Available from: http://www.ncbi.nlm.nih.gov/pubmed/27984733

58. Molina-Mora JA, Campos-Sanchez R, Garcia F. Gene Expression Dynamics Induced by Ciprofloxacin and Loss of Lexa Function in Pseudomonas aeruginosa PAO1 Using Data Mining and Network Analysis. In: 2018 IEEE International Work Conference on Bioinspired Intelligence (IWOBI) [Internet]. IEEE; 2018 [cited 2018 Nov 16]. p. 1–7. Available from: https://ieeexplore.ieee.org/document/8464130/

Supporting information

Supplementary Figures

Fig. S1. Evaluation of normalization by *proC* gene. (A) Systematic microarray data selection in GEO-NCBI database according to experimental design; (B) Comparison of conditions (control vrs perturbations) for other housekeeping genes after normalization by *proC* gene; (C) Dispersion of intensities of samples, showing similar distribution between samples.

Fig. S2. **Evaluation of SP method for gene ranking by importance, case for SVM and KNN algorithms.** (A-C) Accuracy, F1 and kappa values after iterations of classification with the first top 200 genes (adding genes 1-to-1). (B-D) ROC plot using selected top 56 genes for evaluation of performance of the algorithm. (E) Ranking and importance value of top 56 genes (same importance metric for both algorithms).

Fig. S3. **Evaluation of MP method for gene ranking by frequency, case for KNN and RF algorithms.** (A) KNN evaluation of ranking of top 56 genes by frequency after iterations of 100 data partitions and classification model fitting and dispersion of metrics across 100 iterations. (B) Same evaluation, case for RF algorithm.

Fig. S4. Systems level description and gene expression comparison given by core response genes. (A) Gene expression and clustering of the 46 core response determinants of *P. aeruginosa*. (B-C) Comparison of gene expression (logR) of kmedoids for each cluster using small world networks.

Supplementary tables

Table S1. Evaluation of housekeeping genes (columns) after data normalization with another specific housekeeping gene (rows).

Table S2. Annotation of genes of core response genes of *P. aeruginosa* PAO1

Figure1

A

Figure2

Figure3

Figure4

Figure5

A Core response genes

Translation 3%
Others 4%
Transport 8%
Biosynthesis 19%
Metabolism 40%
Molecule binding 26%

B

Core response genes

13 33 68

DEGs

C DEGs

3% 4%
7%
14%
49%
23%

Figure6