A comprehensive portrait of antimicrobial resistance in the zoonotic

pathogen *Streptococcus suis*

Nazreen F. Hadjirin, Eric L. Miller, Gemma G. R. Murray, Phung L. K. Yen, Ho

D. Phuc, Thomas M. Wileman, Juan Hernandez-Garcia, Susanna M.

Williamson, Julian Parkhill, Duncan J. Maskell, Rui Zhou, Nahuel Fittipaldi,

Marcelo Gottschalk, A.W. (Dan) Tucker, Ngo Thi Hoa, John J. Welch, Lucy A.

Weinert

Abstract

Antimicrobial resistance (AMR) is among the gravest threats to human health

and food security worldwide. Pigs receive more antimicrobials than most other

livestock, and are a known source of zoonotic disease. We studied AMR in

Streptococcus suis, a commensal found in most pigs, but which can also

cause serious disease in both pigs and humans. We obtained replicated

measures of Minimum Inhibitory Concentration (MIC) for 16 antibiotics, across

a panel of 678 isolates, from the major pig-producing regions of the world. For

several drugs, there was no natural separation into "resistant" and

"susceptible", highlighting the need to treat MIC as a quantitative trait. We

found differences in MICs between countries, consistent with their patterns of

antimicrobial usage. AMR levels were high even for drugs not used to treat *S.*

suis, with many multi-drug resistant isolates. And similar levels of resistance

were found in pigs and humans from zoonotic regions. We next used whole

genome sequences for each isolate to identify 43 candidate resistance

determinants, 22 of which were novel in *S. suis*. The presence of these

determinants explained most of the variation in MIC. But there were also

complications, including epistatic interactions, where known resistance alleles

had no effect in some genetic backgrounds. Beta-lactam resistance involved

many variants of small effect, appearing in a characteristic order. Our results

confirm the potential for genomic data to aid in the fight against AMR, but also

demonstrate that it cannot be tackled one species or one drug at a time.

Introduction

The ability of bacterial pathogens to evolve resistance to antimicrobials is one of the gravest threats to human health and food security worldwide. Antimicrobial resistance (AMR) to a given drug can be quantified via the minimum inhibitory concentration (MIC), i.e., the concentration of the drug that is sufficient to inhibit growth of a bacterial culture. While, for practical use, bacterial isolates are often categorised as either susceptible or resistant, MIC is a continuously varying trait. In this study, we investigate how AMR genes, variants and ecology explain antibiotic phenotype in the bacterium *Streptococcus suis*, treating MIC phenotype as a quantitative trait.

S. suis primarily exists as a commensal in pigs, colonising the nasopharynx, gut and vagina, but it also causes systemic and respiratory disease, particularly in young pigs [1]. *S. suis* is also a serious zoonotic disease, being the leading cause of adult bacterial meningitis in Vietnam [2]. While some autogenous vaccines are used in pig production, they are serotype-specific and give inconsistent cross protection against heterogeneous *S. suis* [3]. Antimicrobials therefore remain the standard treatment for *S. suis*, and as such, *S. suis* is a leading driver of antimicrobial usage in pig farms [4].

As well as being a serious problem in itself, *S. suis* also has unique benefits as a model for studying AMR. By weight, more pork is consumed globally than any other meat [5, 6], and *S. suis* is found in most, if not all pigs [7]. Furthermore, antibiotic consumption is higher in pigs (172 mg per population corrected unit) than any other livestock (e.g. cattle (45 mg) and chicken (148

mg)) [8]. As a result, most *S. suis* lineages will experience antibiotics. These

include not only antibiotics administered directly against *S. suis*, whether as a

therapeutic, prophylaxis or metaphylaxis, but also, and perhaps more

commonly, in response to many other bacterial infections.

The strong selection pressure caused by widespread use of antimicrobials in

pig farming is expected to give rise to AMR in *S. suis*. Consistent with this,

several phenotypic studies show high MICs for each of the major classes of

antibiotics in one or more *S. suis* collections [9-13]. In addition, there have

been demonstrations of individual resistance determinants affecting MIC in *S.*

suis [14-16], and also some mining of *S. suis* genome collections for known

resistance determinants [17-19]. However, to our knowledge, no study has

combined complete genomic and phenotypic information in large numbers of

isolates, collected from a diverse range of populations.

To this end, we obtained replicated measures of MIC for 16 antibiotics, each

widely used in pigs, for six different *S. suis* collections, comprising 678

isolates. These collections were chosen to include the three main pig-

producing regions of the globe - namely the Americas, South East Asia and

Europe - and targeted both human and pig hosts, as well as a range of years,

serotypes and clinical phenotypes [20-22]. We obtained whole genome

sequences for each of our isolates, allowing us to compare phenotype and

genotype on an unprecedented scale.

Results

1. Measurement of MIC for 16 antibiotics in *S. suis*

We scored 678 isolates of *Streptococcus suis* for replicated measures of

Minimum Inhibitory Concentration (MIC) for 16 antibiotics (Table S1). Four of

these are beta-lactams (amoxicillin, cefquinome, ceftiofur and penicillin),

which are typically used to treat *S. suis* infection in pigs. The other antibiotics

are all widely used in the pig industry, but found in different drug classes.

They comprise macrolide-lincosamide-streptogramin B (MLS$_B$: erythromycin,

lincomycin, tilmicosin and tylosin), tetracyclines (doxycycline and tetracycline),

and fluoroquinolones (enrofloxacin and marbofloxacin), plus one each of an

aminoglycoside (spectinomycin), a pleuromutilin (tiamulin), trimethoprim

(TMP), and a phenicol (florfenicol). Results, shown in the left-hand plot of

Figure 1, reveal wide variation among our isolates in MIC for most of the

antibiotics. For a few antibiotics, especially the MLS$_B$ and tetracyclines, the

MIC values are clearly bimodal, suggesting a meaningful division between

"resistant" and "wild-type" isolates. However, for most antibiotics, the

distributions are either roughly lognormal (fluoroquinolones and phenicols) or

positively skewed (spectinomycin, pleuromutilin, TMP and beta-lactams). It is

striking that the distributions for the beta-lactams are not clearly dissimilar

from any of the other antibiotic classes, despite their being the most common

treatment against *S. suis*. Given these distributions, and the general lack of

defined clinical breakpoints for these antibiotics in *S. suis* [23], we treated MIC

as a quantitative trait in our subsequent analyses.

2. Ecological and genomic predictors of MIC

Our *S. suis* isolates were collected at different times, and from different

countries, hosts and body sites, including sites associated with respiratory

and systemic disease. Isolates were also genetically heterogeneous, and

differed in their serotype; including serotypes with a known disease

association (see methods and Table S1). We first used linear models, to ask

whether MIC levels varied systematically with these ecological and genomic

factors.

S. suis is highly recombining, such that no single genealogy describes its

diversity [20]. As such, to characterise genomic diversity in our sample, we

identified 30 clusters of similar strains (see methods). The resulting clusters

were a highly significant predictor of MIC (p<10^{-15}; Table 1a) showing that

genomically similar isolates tend to have similar MIC values. To account for

this genetic structure, "cluster" was included as a random effect in subsequent

analyses.

A model including all potential predictors showed that country of isolation was

also a highly significant predictor of MIC. In addition, there were weaker

effects of year of collection (with a slight trend for increasing MIC over time);

for serotype (with non-disease-associated serotypes tending to have higher

MIC); and for host disease status (with non-clinical isolates having higher

MIC) (Table 1b).

Separate analyses for each antibiotic, showed that the effect of country was

driven by consistently higher MIC in the samples from Canadian pigs. As

shown in Figure 2a (left-hand panel), this applied to 15/16 antibiotics, and also

applied to various models (Table 1c), and subsets of the data (Figure S1). For

example, the effect is seen consistently in systemic pathogens, respiratory

pathogens and non-clinical isolates (Figure S1). By contrast, the other

predictors (year, serotype and clinical status) had effects that were less

consistent or robust (Table 1d-f). The difference in the MIC values between

Canada and the UK is consistent with higher antimicrobial usage in Canada

[24, 25].

Predictions for Vietnam are more difficult, as there is a lack of official records

of antibiotic usage, although it is likely that usage is both higher than in the EU

and focussed on different drugs [26]. However, as shown in Figure 2b (left-

hand panel), we saw no such signal in the MIC data, even when we restricted

the comparison to the genetic cluster BAPS4 that includes all of our

Vietnamese sample (see methods). The clearest pattern is for lower MIC for

the MLS_B class in the Vietnamese isolates (pink points in Figure 2b left-hand

panel). As shown in Figure 2c (left-hand panel), we also found no robust

differences between Vietnamese isolates sampled from humans and pigs.

This is consistent with pigs being a reservoir for this zoonotic disease, and the

lack of evidence of genomic adaptation to human hosts [20].

3. Identification of candidate AMR determinants within *S. suis* genomes

Using a candidate locus approach, we next identified known and putative

determinants of AMR that were present in our sampled genomes. We

considered both the presence/absence of whole genes, and sequence

variants within genes. In total, we detected 43 determinants: 10 completely

novel variants of known resistance genes, 12 known resistance genes

detected in *S. suis* for the first time, and 21 previously reported in *S. suis*

(Table 2 and Table S1). Our candidates come from three sources: the

antimicrobial resistance database CARD [27], previously published *S. suis*

variants not present in the CARD database [28-30] and novel variants

detected using a subset of our data (see methods).

Of the 10 novel determinants, six are variants in genes previously associated

with resistance in other bacteria (including a promoter variant). In particular,

we discovered four novel haplotypes within the loop region of the central

transpeptidase domains of the *pbp2B* and *pbp2X* genes and within the signal

peptide region of the *mraY* gene (encoding the enzyme, acetylmuramoyl-

pentapeptide-transferase, essential to cell wall biosynthesis) that were

associated with variation in beta-lactam MIC (Table 2). In particular, we noted

a PBP2X mutation at the conserved location T551, similar to that found in *S.*

pneumoniae (T550), group B streptococci (T555) and *S. pyogenes* (T553)

PBP2X proteins [31] indicative of a shared overall mechanism mediating beta-

lactam resistance across the genus. Polymorphisms in *mraY* have also been

identified in a genome-wide association study of beta-lactam resistance in *S.*

pneumoniae [32]. In addition to these four beta-lactam haplotypes, two

variants – a variant of the chromosomal dihydrofolate reductase gene *dhfr*

and its promoter – were associated with reduced susceptibility against

trimethoprim (Table 2). Furthermore, all of these six variants were

independently associated with changes in MIC in at least seven different

genetic clusters (Table S2) implying that the association between these

variants and MIC is either directly causal or compensatory to the causal

variant [33].

The remaining four novel variants were whole genes with homologies to

known resistance genes: three located on mobile genetic elements (MGEs)

and one chromosomal (Table 2). Three novel aminoglycoside genes, carried

by isolates with high spectinomycin MIC, were identified based on

aminoglycoside resistance determinant protein homologies. Although

previously undescribed, we found homologues in other bacteria using a *blastn*

search of the non-redundant database in *Genbank* (Table 2). We also

characterised a chromosomally-encoded *vgaC* homologue (37% protein

homology to *vgaC*), encoding an ABC-F ATP-binding cassette ribosomal

protection protein, that we designate *vgaF.* This gene has arisen in 13

different BAPS clusters, each time associated with reduced susceptibility to

tiamulin (Table S2).

The 12 previously-known AMR genes detected in *S. suis* for the first time are

mobile genetic element (MGE)-linked genes that confer resistance to

aminoglycosides (2/12), MLS$_B$ (2/12), tetracyclines (5/12), TMP (2/12) and

phenicol (1/12*).* Based on the CARD and NCBI databases, these genes are

associated with other gram-positive and gram-negative bacteria (Table 2).

Overall, most of our candidate determinants (28/43) were found in fewer than

5% of the isolates, but *ermB* (which confers resistance to the MLS$_B$ class) and

tetO (which confers resistance to tetracyclines) were present in a majority of the isolates (Table 2). Only about a quarter of isolates carry a candidate determinant for the beta-lactams highlighting the continuing susceptibility to these first line treatment drugs in *S. suis*.

Nevertheless, as shown in Figure 3, many isolates carried multiple determinants, and multi-drug resistance was widespread. Around 40% of isolates carried resistance determinants to three or more classes of drug (275/678), and over 10% carried resistance determinants to five classes (81/678); this is more than carried no determinants at all (59/678).

4. Candidate AMR determinants explain most of the variation in MIC

We next investigated how well our candidate variants explain the variation we observe in MIC (see Figures S2-S8 for detailed plots). The right-hand panel of Figure 1 compares the distribution of MIC values for isolates carrying one or more of the candidate AMR determinants for that antibiotic class (denoted "1+"), to the MIC of the remaining isolates, which carry no such determinant (denoted "0").

The proportion of the variation in MIC explained by our variants varied between antibiotics. Explanatory power was strongest ($r^2 > 0.75$) for antibiotics where the distributions of MIC were most clearly bimodal (MLS$_B$ and tetracyclines). By contrast, we found no association between our variants and MIC for the fluoroquinolones and phenicol, where the distribution of MIC values was roughly lognormal, consistent with our isolates representing a "wild-type" population. For the remaining antibiotics, our determinants had

intermediate explanatory power (Figure 1), but there were very few isolates

with high MIC that did not carry a candidate determinant. This suggests that

we have detected most of the causal variants in our dataset, removing the

need to perform additional genome-wide associations.

We next asked whether the variable presence of our candidate determinants

in different ecological settings would explain the significant ecological

predictors of MIC in our linear models (Table 1). Consistent with this

hypothesis, the typical MIC level in each genetic cluster was highly correlated

with the frequency of determinants in that cluster (Figure S9).

Differences between countries were explained in the same way. The higher

MIC in Canada than the UK was due to consistently higher frequencies of the

same determinants (Figure 2a right-hand panel) – with the exception of

trimethoprim (see below). By contrast, tetracycline resistance throughout the

BAPS4 cluster, though at similar levels in all three countries, was conferred by

different determinants in Vietnam (Figure 2b).

5. Ineffective variants and epistasis

While our data contained few isolates with high MIC that did not carry a

candidate determinant, there were many isolates with low MIC despite

carrying a determinant (Figure 1).

In some cases, this was due to previously identified candidate genes that had

no appreciable effect on MIC in our data. For example, spectinomycin MIC

was unchanged by some determinants identified from the CARD database

(e.g. *ant(6')-Ib* and *aph(3')-IIIa* in Figure S6). If we allow for the presence of

these non-functional genes, the r^2 value for spectinomycin increases from

0.37 (Figure 1) to 0.69 (Figure S10a).

In other cases, as is well known, variants act against only some of the

antibiotics in a class. In MLS_B, for example, modification of the ribosomal

target confers cross resistance to macrolides and lincosamides, while

mechanisms such as efflux and enzymatic inactivation do not [34]; so *msrD*

acts against erythromycin, but not against lincomycin, while *linB* inactivates

lincomycin, but not erythromycin (Figure S3). Again, taking this into account

further increases predictive power (Figure S10a).

As well as these simple cases, there were some clear examples of genetic

interactions. For trimethoprim, two candidate variants – in the protein

DHFR102 and its promoter (Table 2) – were found at higher frequency in the

UK than Canada (Figure 2a right-hand-panel), but in Canada, the two variants

were more often found together, leading to higher MIC overall (Figure 2a;

Figure S8).

Finally, we observed complex patterns for the second most common

determinant in our dataset, *ermB*. For the 334 isolates that carried only *ermB*

(and no other candidate determinant to MLS_B class drugs), MIC values were

clearly bimodal, with many carriers having very low MIC (Figures S3 and

S11). Further analysis revealed a small number of isolates with frameshift or

premature stop codons (4/326 isolates with complete sequences in our

assemblies). While the remaining isolates with low MIC carried rare amino

acid variants at one of four positions (T75X, N100S, R118H and V226I) – the

last three of which differentiate the *ermB* sequences found in *Streptococcus*

pneumoniae from *Clostridium perfringens* [35]. Nevertheless, no particular

sequence was always associated with low MIC, suggesting some unidentified

source of epistasis (see methods and Figure S11 for full details).

6. Resistance to beta-lactams

While beta-lactams are the major treatment class for *S. suis*, it is notable that

the explanatory power of our variants seems to be weaker for this drug class

than for, e.g. tetracyclines or MLS$_B$.

In Streptococci a typical route to beta-lactam resistance involves variants in

the *pbp* genes, and it is well established that the joint action of many *pbp*

variants is necessary to explain substantial changes in MIC [36, 37]. Single

point mutations in *pbp2x,* for example, cause very modest elevations in *S.*

pneumoniae [36], group B Streptococci [38], *Streptococcus dysgalactiae*

subsp. equisimilis [39] and *Streptococcus pyogenes* [31].

Like *S. pneumoniae, S. suis* has three key mosaic *pbp* genes, *pbp1A, pbp2b*

and *pbp2x*, and shares a similarly broad MIC distribution (although with

typically lower MIC values) [37, 40, 41]. Our data also show that mutations in

pbp2x alone have small effects (Figure S2), while genotypes carrying four or

more variants have the highest mean MIC (Figure S2). Our explanatory power

increases greatly when we predict beta-lactam MIC from the total number of

candidate variants carried (Figure S10b). Although official breakpoints are

lacking for many beta-lactam antibiotics in *S. suis*, for penicillin, only the

isolates with the most variants reach clinical significance (i.e. penicillin

resistance ≥ 1 or log(0) MIC).

In *S. pneumoniae* it has also been noted that mutations conferring resistance

to beta-lactams occur in a set order, with amino acid changes in PBP2B and

PBP2X often acting as the first step [36, 42, 43]. Figure 4 shows that our data

also show this characteristic "nested" pattern (see also Figure S12; [44]), with

PBP2X mutations largely occurring in backgrounds containing PBP2B

mutations and *mraY* mutations occurring in backgrounds containing both

PBP2B and PBP2X mutations.

It is notable that mutations in *mraY* – which are generally last to occur, and

never occur alone – might have a compensatory effect. Altered PBPs, while

conferring resistance, might be less active transpeptidases than their wild type

counterparts [45]. For example, the amidotransferase enzyme encoded by

murT enables cross-linking of cell wall peptidoglycans by preparing lipid II

used by pneumococcal PBPs [46, 47]. Mutations in *mraY* might therefore

compensate for reduced enzymatic activity of altered PBPs.

Studies have shown that cefotaxime, a third-generation cephalosporin like

ceftiofur, selectively inactivates PBP2X but not PBP2B [48] and that mutations

within the catalytic domain in PBP2X give rise to cefotaxime, but not to

penicillin, resistance [49]. Our results echo this pattern because our PBP2B

haplotype has a smaller change on mean MIC of ceftiofur than penicillin

(Figure S2), and the explanatory effects of our PBP haplotypes differ between

the drugs (Figure 1). Previous variants in PBP2X thought to confer cefuroxime

resistance [28], whilst validated by our data, were only found at very low

frequency in our collection (Table 2).

Discussion

We have examined antimicrobial susceptibility of *Streptococcus suis* to 16
antibiotics that are commonly used in pigs. Replicated measures of MIC
suggest that, even when some isolates have very high MIC, there is often no
natural separation into "resistant" and "susceptible" (Figure 1). This highlights
both the need for clinical breakpoints in *S. suis*, and the need to evaluate
AMR with quantitative measurements of MIC. This is particularly important for
beta-lactam resistance, given its multi-allelic nature (Figures 4, S2), and the
possibility of "MIC creep" that may eventually lead to treatment failures (e.g.
vancomycin resistance in MRSA [50]).

We have also shown that AMR levels differ systematically between countries.
For example, compared to the UK, Canada has consistently higher MICs, and
frequencies of the same candidate AMR determinants (Figures 2, S1).
Differences in antimicrobial usage are the most plausible explanation for this
trend, because usage is much higher in Canada than the UK [24, 25].
However,
our results are correlational and other factors could be explanatory e.g.
different microbiome compositions in pigs from different areas.

Next, we have identified 43 candidate resistance determinants, and shown
that they explain the majority of the observed variation in MIC for 11/16
antibiotics, and a substantial fraction of the variation for a further 2/16 (Figure
1).

While many of these variants were previously described, and most were found

at low frequency (Table 2; Figure 2), we detected three novel candidates that

are common and associated with clinically-relevant changes in MIC: *vgaF*,

DHFR102 and the *dhfr* promoter. A new mechanism for tiamulin resistance,

such as the *vgaF*, could have implications for treatment success because

tiamulin is a common drug for treating infections in livestock, including pigs

[51]. High-level trimethoprim resistance was associated with mutations in the

promoter regions of the *dhfr* gene. This is in contrast to the *folA* (I100L), *folP*

1-2 codon insertion combination conferring high level resistance to

trimethoprim/sulfamethoxazole (TMS) in *S. pneumoniae* [52]. While our MICs

were for trimethoprim, our observations suggest the evolution of divergent

mechanisms of resistance against TMS between these species. If these

resistance mechanisms are functionally verified, they should be included in

routine AMR gene testing for *S. suis*.

While our candidate variants were explanatory, we also found several

complexities in the genotype-phenotype map. These include epistatic

interactions, where the effects of some candidate genes vary with their *S. suis*

genetic background. Phenotypic reversion (i.e. mutations in transcriptional

regulators or elsewhere in the genome, counteracting a resistance

determinant) can be common if resistance has a fitness cost [53], and this

could explain why we see many isolates carrying candidate determinants but

with low MIC (Figure 1). This same pattern is also apparent in up to 10% of *S.*

aureus isolates [54]. These complexities have consequences for diagnostic

investigation of AMR of *S. suis* using whole genome sequences. While the

predictive power is quite high for many antibiotics, we should expect many

Type I errors (false positives). These results also caution against using the

presence of AMR determinants as a measure of resistance more generally.

While susceptibility to penicillin in clinical cases of *S. suis* remains high, we

see substantial variation in beta-lactam MIC (Figure 1a). Consistent with

studies in other Streptococci, our results show incremental and ordered

changes in amino acids, first within PBP2A and then PBP2X that lead to

clinically relevant elevations in MICs. Accurate predictions of genotype-

phenotype have already been developed in *S. pneumoniae* [37, 52, 55-57],

group B Streptococci [58] and *S. pyogenes* [31]. With no effective treatment

for *S. suis* other than antibiotics, developing similar models and monitoring of

these variants in *S. suis* populations should be a priority.

Two final aspects of our results highlight the fact that AMR is not a problem

that can be tackled one disease at a time.

First, we have found high levels of resistance in *S. suis* even to antibiotics that

are not typically used to treat this infection, including high rates of multi-drug

resistance (Figure 3). This trend could be partly due to the co-occurrence of

resistance genes on MGEs. However, determinants for beta lactams (found in

190/678 isolates) – which are the primary treatment – were less common than

those for tetracyclines (574/678), MLS_B (395/678), trimethoprim (246/678) and

tiamulin (193/678). This pattern probably reflects a "bystander selection

effect", common in opportunistic pathogens that are part of the healthy

microbiota and frequently exposed to antibiotics used as growth promoters or

therapeutics.

Second, many of the rarer AMR genes in our sample were not previously

known in *S. suis*, but are common in other bacteria, both gram negative and

gram positive (Table 2). Together with the presence of non-typical

Streptococcal *ermB* in some isolates, this raises the possibility of *S. suis*

acting as a reservoir for AMR determinants in other bacteria. Indeed *S. suis*

shares conserved chromosomal insertion sites with many human pathogens,

such as *S. pyogenes, S. pneumoniae*, and *S. agalactiae* [17]. Given that *S.*

suis causes human clinical disease, a direct exchange of AMR genes

between *S. suis* and other human pathogens is plausible. Our results lend

further support to a one health approach to tackling AMR.

Material and Methods

***Streptococcus suis* isolates**

We assembled a collection of 678 *S. suis* isolates, including isolates from the

UK (*n*=423), Canada (*n*=205) and Vietnam (*n*=50) (Table S1).

In the UK, isolates came from three different collections. The first collection in

2009-2011 sampled non-clinical and clinical isolates from pigs across England

and Wales (described in Weinert *et al.* 2015 [20]). The second collection in

2013-2014 sampled non-clinical isolates from five farms (described in Zou *et*

al. 2018 [22]). The third collection during 2013–2014 targeted clinical isolates

from pigs across England and Wales (described in Wileman *et al.* 2019 [21]).

In pigs that showed clinical symptoms consistent with *S. suis* infections (e.g.

meningitis, septicaemia and arthritis), the site of isolation was classified as

'systemic' if recovered from systemic sites. The site of recovery was classified

as 'respiratory' if derived from lungs with gross lesions of pneumonia. *S. suis*

isolates from the tonsils or tracheo-bronchus of healthy pigs or dead pigs

without any typical signs of *S. suis* infections were defined as 'non-clinical'.

Isolates that could not confidently be assigned to these categories (e.g. a

tonsil isolate from a pig with systemic signs) were classified as unknown.

Altogether, the UK isolates were classified by clinical status as 'systemic'

(*n*=94), 'respiratory' (*n*=50), 'non-clinical' (*n*=197) or 'unknown' (*n*=82),

respectively.

The Canadian pig *S. suis* isolates from 1983-2016 were collected to target

similar numbers of clinical and non-clinical isolates and were also classified by

clinical status as 'systemic' (*n*=81), 'respiratory' (*n*=30), 'non-clinical' (*n*=55) or

'unknown' (*n*=39).

The Vietnamese isolates were collected to sample related populations from

human and pig (described in Weinert *et al.* 2015 [20]). These comprised

'systemic' isolates (*n*=28) from human clinical cases of meningitis from

provinces in southern and central Vietnam, and 'systemic' (*n*=4) or 'non-

clinical' isolates (*n*=18) from pigs, collected between 2000 and 2010. These

isolates were exclusively serotype 2 or serotype 14 and belong to one genetic

population (Table S1).

Antimicrobial susceptibility testing

The minimum inhibitory concentrations (MIC) for a range of antibiotics were

determined by the micro-broth dilution method, which was performed and the

results interpreted in accordance with CLSI Approved Standards, M100-S25

(2015), Vet01S 3[rd] Edition (2015) and de Jong *et al.* (2014) [9, 59]. MIC

measurements for some of our isolates were previously published [9]. For the

remaining MIC measurements, the MICs were determined for sixteen different

antimicrobial compounds, representing nine antimicrobial classes at LGC,

Fordham, UK (formerly Quotient Bioresearch, Fordham, UK) for the UK and

the Canadian isolates. MIC testing of antibiotics for the Vietnamese isolates

was performed at the Oxford University Clinical Research Unit, Ho Chi Minh

City, Vietnam, in collaboration with the Department of Veterinary Medicine,

University of Cambridge, UK.

Whole genome sequencing, assembly and inference of population

structure

Genome DNA extractions and whole genome sequencing of newly sequenced

S suis in this study were as previously described by Weinert *et al.* (2015) [20].

Briefly, single colonies of strains were grown up in broth culture, DNA was

extracted using DNeasy kits (Qiagen), Illumina library preparations were

performed as described by Quail *et al.* 2008 [60] and the whole genomes

sequenced on the HiSeq2000 according to the manufacturer's instructions

(Illumina, San Diego, CA, USA) at the Welcome Trust Sanger Institute,

Cambridge, UK. Sequencing generated 125bp paired end reads, which were

assessed for quality using Sickle [61] after the removal of adapter sequences.

Sequencing reads that passed the quality threshold were put forward for *de-*

novo assembly generation using Spades v.3.10.1 [62] utilising a variety of

parameters conditions. Our 678 isolates were combined with 401 additional

genomes given in Table S3 to increase robustness of *S. suis* population

structure estimation, although these isolates were unavailable for MIC testing.

Draft genomes were annotated using Prokka v1.12-beta (v2.8.2). [63] and

bacterial species assignment was performed by a combination of MLST

assignment and FastQ Screen v.0.11.1 using a custom database [64]. We

mapped the Illumina reads back to the *de-novo* assembly to investigate

polymorphic reads in the samples (indicative of mixed cultures) using BWA

v.0.7.16a [65]. Genomes that exhibited poor sequencing quality (i.e poor

assembly as indicated by a large number of contigs, low N50 values or a high

number of polymorphic reads) or that which were inconsistent with an *S. suis*

species assignment were excluded from the analysis.

In order to group our *S. suis* isolates in to different genetic clusters, we used

the program hierBAPs in R [66]. First, we inferred core genes from our

isolates using Roary (v2.8.2) [67], aligned them using MACSE [68], and

stripped regions that could not be aligned unambiguously due to high

divergence, indels or missing data. This conserved region of the core genome

was used as input in hierBAPS.

Known AMR determinant detection

ARIBA [69] identifies AMR determinants (or any sequence of interest) directly

from paired sequencing reads using a public or custom reference database

and relies on mapping the reads to reference sequence clusters followed by

the formation of local assemblies of the mapped reads. The tool is further able

to confirm the intactness of resistance genes to identify known or novel SNPs

within a gene of interest.

The AMR determinants in *S. suis* were identified using ARIBA v2.10.0 based

on the public AMR database, CARD [27], which was further supplemented

with a custom database. The custom database contained gene sequences

not present within the CARD database at the time of testing. These were

previously published *S. suis* AMR genes, single nucleotide polymorphisms

(SNPs) in genes known to confer antibiotic resistance and AMR genes found

in other bacteria. In addition, we investigated whether there were additional

novel resistance variants in known resistance genes (described below). The

sequence identity threshold against a reference was set at 90%. Only paired-

end sequence reads that passed the quality control thresholds were used as

input for ARIBA. To identify genes falling under the 90% identity threshold, we

also performed blast searches of the draft assemblies against the non-

redundant NCBI protein or nucleotide databases to identify variants or

chimeric alleles of known resistance determinants that might not be present in

the CARD database. Examples of allelic variants identified this way include

tet, *aad*, *ant* and *cat* (Table 2).

Novel AMR candidate determinant detection

We identified novel resistance determinants by a range of methods.

Candidate determinants that might encode novel resistance mechanisms

were identified by scanning literature describing experimental studies and

others found by Genome Wide Association Studies (GWAS) in related

bacterial species, for example, variants of *pbp*s, *mraY*, *dhfr* and *folA* [36, 70-

72] along with their promoter regions. To avoid over-fitting our generalised

linear models, gene or promoters of interest in a "training" subset of the

collection (*n*=205) were then extracted, aligned and ranked from the highest

to lowest MIC values, using MUSCLE [73] in SEAVIEW [74]. Manual

sequence analysis was then performed to identify either amino acid or

nucleotide variations that associated with high MIC.

Kinetic and structural studies have previously established that beta-lactam

resistance is conferred by substitutions within PBPs in *Streptococcus*

pneumoniae [36, 75] and other streptococci [39, 76, 77]. While mutations are

present throughout the entire PBPs, we noted statistically significant

mutations (Table S2) in altered PBP2B and PBP2X proteins in strains

exhibiting high penicillin (≥ 1mg/L) and ceftiofur MICs (≥ 2mg/L), respectively.

In PBP2B, the residue variations were present in loop regions, K479T/A,

D512E/Q/K/A, K513E/D and T515S, within the *S. suis* transpeptidase domain

(residues 351-681). In PBP2X, the mutations were located at positions

M437L, S445T, T467S, Y525F and T551S, also in loop regions within its

catalytic domain (residues 265–619). No statistically significant variations

were found within PBP1A, although the variant P405T was shared by some of

the BAPs clusters. As multiple amino acid variants were identified within

PBP2X and PBP2B, many of which were in the same genomic region (Table

2), we grouped the variants in to haplotypes. We did this by scoring haplotype

presence if the isolate had all of the amino acid variants associated with high

MIC in a given gene.

In addition to allelic differences in PBP2B and PBP2X, we also detected

mutations in *mraY*. Found immediately upstream of *pbp2X*, *mraY* encodes an

acetylmuramoyl-pentapeptide-transferase enzyme which is essential to lipid

cycle reactions in the peptidoglycan cell wall biosynthesis pathway. Residue

substitutions were present at A4S/T, M6I/L and G8S within the signal peptide

regions of the protein. These substitutions were in strong linkage

disequilibrium, such that they almost always occurred in pairs. In particular,

both A4S/T and G8S occurred only in isolates also carrying M6I/L, while M6I/L

appeared alone in only a single isolate. As such, we scored each isolate as

carrying the *mraY* determinant only if it carried two of these three amino acid

substitutions (either A4S/T and M6I/L, or G8S and M6I/L; Table 2). This set of

isolates differed significantly in their MICs in both penicillin and ceftiofur

across multiple BAPs groups (Table S2).

Not all isolates with high tiamulin MICs possessed a *lsaE* gene, suggesting an

additional mechanism in *S. suis* conferring resistance to pleuromutilins. Using

the CARD RGI (resistance gene identifier) protein homology models, a *vgaC*

homologue was identified as a candidate gene associated with variation in

tiamulin MIC. The gene, designated as *vgaF* in this work, encodes an ABC-F

ATP-binding cassette ribosomal protection protein and shares 37.14%

homology to the reference. Located within the chromosome, *vgaF* is found

intact (~1386bp) in statistically significant numbers in isolates with MIC ≥

8mg/L (Table S2) but is truncated (<750bp) in others. Plasmid borne *vga*

homologues including *vgaC* are frequently detected in staphylococci and have

been shown to confer cross resistance to pleuromutilins, lincosamides and

streptogramin A antibiotics. A chromosome based *vgaA* gene variant,

encoding an ATP-binding cassette protein conferring resistance to

streptogramin A and related antibiotics in *S. aureus* has also been described

[78].

An amino acid substitution (I102-L) was detected in the dihydrofolate

reductase (DHFR) gene, *dhfr* (counterpart of *folA* in *S. pneumoniae*), in the

majority of isolates with trimethoprim MICs of 0.12mg/L or greater. Similar

substitutions of isoleucine to leucine at position 100 in *S. pneumoniae* [71, 79]

and *Streptococcus pyogenes* [80] is known to cause resistance to TMP.

Additionally, we also identified polymorphisms within the promoter region (0-

30bp upstream) of the *dhfr* gene; an A5G substitution and insertions, in

isolates exhibiting MICs ≥ 1mg/L.

Mutations such as 1-2 codon insertions within *folP*, another core metabolic

gene that is documented to confer resistance to

trimethoprim/sulfamethoxazole in *S. pneumoniae*, were also examined. Whilst

insertions and or deletions were absent, amino acid residue variations were

observed at position 198 (A198G) either alone (9/678) or in combination with

dhfr I102L (26/678 isolates) in isolates exhibiting a wide range of trimethoprim

MICs; 0.03mg/L-32mg/mL. However, this data was not statistically significant

in the Binomial sign tests, and hence was excluded from further analysis.

All of these variants might be causal, compensatory or linked to high MIC

because of population structure. We therefore examined our variants and

showed that susceptible and resistance alleles were found repeatedly in

different BAPs clusters, which were significant using binomial sign tests when

defining a cut-off (Table S2).

The *pbp*, *mraY* and *vgaF* genes were unable to be included into the custom

database in ARIBA because of the sequence divergence. Therefore, we

manually aligned the genes in all 678 isolates using MUSCLE [73] in

SEAVIEW [74].

Similarly, analysis for known variants in *gyrA* and *parC* were also performed

by manually aligning the genes. The *tet(O/W/32/O)* gene was scored by using

blastn because ARIBA was unable to differentiate mosaic sequence patterns

626
627

Serotype inference from whole genomes

Serotypes were determined *in silico* using the Athey *et al.* (2016) [81]

serotype database, which we implemented in ARIBA. Failed ARIBA runs and

sequence non-matches were designated as not available (NA).

Statistical analyses

All general linear models were fit in R v. 3.3 [82] using the built-in function *lm*

for models with solely fixed effects, or via Reduced Maximum Likelihood,

using the function *lme* from the package nlme v. 3.1-141 [83], when BAPs

cluster was included as a random effect. The response MIC values were log

transformed. Our data for host and country is confounded because all of our

human samples came from Vietnam (human *S. suis* has a higher prevalence

in South East Asia [84]). Therefore, we coded "Country" as separate

populations – Canada, UK, Vietnam-pig and Vietnam-human. We classified

"serotypes" into either disease-associated or non-disease associated

according to Wileman *et al.* (2019) [21].

Data availability

Whole genome sequence assemblies are available from NCBI under Project

ID PRJNA628943. The gene *cat* from isolate FX419 has accession

MT367165. The gene *aadE1* from isolate CF2D3-1A has

accession MT383663. The gene *aadE2* from isolate BH3D7-4E has

accession MT383664. The gene *ant1* from isolate CF2D3-1C has accession

MT383665. The gene *tetM* from isolate 1129711 has accession MT383666.

The gene *tetM* from isolate LSS85 has accession MT383667. The gene *tetM*

from isolate SS981 has accession MT383668. The gene tetO from isolate

D16-010262 has accession MT383669. The gene *vgaF* from isolate 1230091

has accession MT431628.

Acknowledgements

This work was primarily supported by a Challenge grant from the Royal

Society (reference: CH160114) and an Isaac Newton Trust Research Grant

(reference: 17.24(u)). In addition, LAW, ELM, GGRM were supported by a Sir

Henry Dale Fellowship jointly funded by the Wellcome Trust and the Royal

Society (109385/Z/15/Z). We also acknowledge a BBSRC grant to AWT

(BBSRC grant BB/L003902/1), a LoLa grant to DJM and AWT

(BB/G019274/1) and a China MoST grant to RZ (2013DFG32360) for sample

collections. In addition, we wish to thank Abiyad Baig, Andrew Balmer and

Olivier Restif for helpful discussion on antimicrobial resistance in *S. suis*.

Contributions

LAW and NTH conceived the study. AWT, SMW, RZ, MG, NF, TMW, JP, DJM

contributed isolates. NFH, ELM, PLKY, HDP, NTH collected new data. NFH,

ELM, GGRM, JJW and LAW analysed data. JJW produced all figures. LAW,

NFH and JJW wrote the manuscript.

Table 1. Ecological and genomic predictors of MIC

Analysis	#strains	Fixed effects	Df	F	p
(a)	678	Antibiotic	15	1404.125	$<10^{-15}$
		genetic cluster	29	40.644	$<10^{-15}$
(b)	450	Antibiotic	15	972.3107	$<10^{-4}$
		Year	1	6.6078	0.0102
		Serotype	1	6.6261	0.0101
		Disease Status	2	2.9760	0.0511
		Country	3	92.5155	$<10^{-4}$
(c)	678	Antibiotic	15	1457.4043	$<10^{-4}$
		Country	3	139.5902	$<10^{-4}$
(d)	557	Antibiotic	15	1144.3846	$<10^{-4}$
		Disease Status	2	9.9619	$<10^{-4}$
(e)	542	Antibiotic	15	1131.6347	$<10^{-4}$
		Serotype	1	8.7216	0.0032
(f)	652	Antibiotic	15	1350.8781	$<10^{-4}$
		Year	1	0.1591	0.69

Table 2. AMR determinants of *Streptococcus suis*

AMR gene	Name in figures	Resistance conferred (Antibiotic Class)	No. of isolates	Reported in *S. suis*	Reported in other species (CARD and NCBI)	Reported genetic location in *S. suis* or other species
ant(4')-Ib	ant4Ib	Aminoglycoside	1	This report	*Staphylococcus aureus, Staphylococcus epidermidis, Enterococcus spp.*	Plasmid
ant(6')-Ia/aadE	ant6Ia	Aminoglycoside	56	[19]	*Staphylococcus epidermidis, Streptococcus spp., Enterococcus faecalis*	Plasmid, ICE
ant(6')-Ib	ant6Ib	Aminoglycoside	12	This report	*Campylobacter spp., Bacillus subtilis, Clostridium spp.*	Pathogenicity Island
ant(9')-Ia/spw/aad9	ant9Ia	Aminoglycoside	45	[19]	*Staphylococcus aureus, Enterocoocus spp., Staphylococcus spp.*	Plasmid, ICE
aph(3')-IIIa	aph3IIIa	Aminoglycoside	46	[85]	*Campylobacter spp., Staphylococcus spp., Streptococcus spp., Enterococcus spp.*	Plasmid, transposon
ant1*	ant1	Aminoglycoside	12	This report	*Campylobacter spp., Clostridium difficile, Lactobacillus amylovorus*	Plasmid, ICE
aadE1*	aadE1	Aminoglycoside	2	This report	*Enterococcus faecium, Staphylococcus aureus*	Plasmid
aadE2*	aadE2	Aminoglycoside	5	This report	*Clostridium spp., Campylobacter fetus*	ICE
ermB	ermB	MLS$_B$ (Macrolide/ Lincosamide	370	[86]	*Campylobacter spp., Clostridium spp., Enterococcus spp., Escherichia coli, Klebsiella pneumoniae, Proteus mirabilis, Shigella spp., Staphylococcus spp., Streptococcus spp., Vibrio vulnificus.*	Plasmid, ICE
ermG	ermG	MLS$_B$ (Macrolide/ Lincosamide	1	This report	*Clostridium difficile, Enterococcus faecium*	Conjugative transposon
ermT	ermT	MLS$_B$ (Macrolide/ Lincosamide)	3	This report	*Enterococcus faecium, Staphylococcus aureus, Streptococcus spp.*	Plasmid
mefA	mefA	Macrolide	11	[86]	*Streptococcus spp., Enterococcus faecium, Haemophilus parainfluenzae, Proteus mirabilis*	Transposon, ICE
mel/msrD	mel	Macrolide	9	[86]	*Streptococcus spp., Enterococcus faecium, Haemophilus parainfluenzae, Proteus mirabilis.*	Transposon, ICE
lnuB	lnuB	Lincosamide	3	[17]	*Enterococcus spp., Staphylococcus spp., Streptococcus agalactiae*	Plasmid
linB	linB	Lincosamide	42	[87]		
lnuC	lnuC	Lincosamide	9	[88]	*Campylobacter spp., Clostridium difficile, Streptococcus agalactiae.*	Plasmid
lsaE	lsaE	Pleuromutilin	43	[17]	*Enterococcus spp., Staphylococcus spp., Streptococcus agalactiae.*	Plasmid, ICE
tet44	tet44	Tetracycline	4	This report	*Clostridium spp.*	ICE
tetM	tetM	Tetracycline	81	[19]	*Streptococcus spp., Escherichia coli, Staphylococcus spp., Enterococcus spp., Klebsiella pneumonia, Neisseria meningitides, Clostridium difficile, Salmonella enterica*	Plasmid, ICE
tetM1*	tetM1	Tetracycline	34	This report	*Staphylococcus spp., Streptococcus spp., Enterococcus spp., Bacillus spp., Clostridium spp.*	Plasmid, transposon
tetM2*	tetM2	Tetracycline	43	This report	*Enterococcus spp., Clostridium difficile.*	
tetM3*	tetM3	Tetracycline	13	This report	*Streptococcus pneumoniae, Streptococcus pyogenes, Enterococcus spp., Clostridium spp.*	Transposon, ICE
tetO	tetO	Tetracycline	465	[19]	*Campylobacter spp.*	ICE, plasmid
tetO1*	tetO1	Tetracycline	2	This report	*Dorea spp.,* uncultured bacterium	Conjugative transposon
tetW	tetW	Tetracycline	8	[16]	*Streptococcus agalactiae, Clostridium difficile, Chlamydia trachomatis.*	Plasmid

tet40	tet40	Tetracycline	7	[17, 29]	*Clostridium difficile*		Composite mobile genetic element, ICE
tetL	tetL	Tetracycline	12	[19]	*Streptococcus spp., Escherichia coli, Staphylococcus spp., Enterococcus spp., Klebsiella pneumoniae.*		Plasmid
tet(O/W/32/O) †	tetO:W:32:O	Tetracycline	20	[84]			ICE
dfrF	dfrF	TMP	11	This report	*Clostridium difficile, Enterococcus spp., Staphylococcus aureus, Streptococcus agalactiae*		plasmid
dfrK	dfrK	TMP	7	This report	*Staphylococcus aureus, Staphylococcus epidermidis.*		plasmid
*vgaF**	vgaF	Pleuromutilin	173	This report	Not known		Chromosome. Found as an intact gene (~1386bp) or truncated (<750bp)
*cat**	cat	Florfenicol	3	This report	*Enterococcus faecalis, Streptococcus iniae, Listeria monocytogenes, Staphylococcus aureus*		Plasmid, Transposon
*dhfr**	DHFR102	TMP	235	This report	Mutations in I100L in *Streptococcus spp.* and *Staphylococcus aureus*		DHFR: I102L
*dhfr promoter**	DHFRPromoter	TMP	86	This report	None reported		DHFR : A5G upstream/indels 0-30bp upstream
*pbp2B/penA**	PBP2B:hap	Penicillin	181	This report	Mutations in other locations in *Streptococcus pneumonia* and Group B *Streptococcus spp*		PBP2B : K479T/A, D512E/Q/K/A, K513E/D,T515S
*pbp2X**	PBP2X:hap	Ceftiofur	93	This report	None reported		PBP2X : M437L, S445T, T467S Y525F
pbp2X	PBP2X551	Penicillin	153	This report	Mutation in PBP2X in *S. pneumoniae* (T550), group B streptococci (T555) and *S. pyogenes* (T553)		PBP2X : T551S
pbp2X	PBP2X594:596	Penicillin	15	[28]	*Streptococcus suis.* Mutations in other locations in *Streptococcus pneumonia*, Group B *Streptococcus sp.* and *Streptococcus uberis*		PBP2X : L594Y/F, V596G
pbp2X	PBP2X569	Penicillin	10	[28]			PBP2X : N569Q
*mraY**	MraY:hap	Penicillin	62	This report	None reported		*MraY :* M6I/L and either A4S/T or G8S
parC†	ParC79	Fluoroquinolone	4	[30]	*Escherichia coli, Pseudomonas aeruginosa, Streptococcus spp., Staphylococcus aureus*		*ParC* : S79
gyrA†	GyrA81	Fluoroquinolone	13	[30]	*Escherichia coli, Klebsiella spp., Streptococcus spp., Staphylococcus aureus*		GyrA : S81
gyrA†	GyrA85	Fluoroquinolone	1	[30]	*Escherichia coli, Klebsiella spp., Streptococcus spp., Staphylococcus aureus*		GyrA : E85

*novel determinants reported in this study
† Not documented in the CARD database
MLS$_B$ Macrolide Linocsomide Streptogramin B
TMP Trimethoprim

 Figure legends

Figure 1. Candidate AMR determinants explain most of the variation in

MIC. Histograms of log transformed MIC measures for each of our 16 different

antibiotics, across our panel of 678 *S. suis* isolates. Antibiotics are coloured

by their class (Beta: beta-lactams; MLS_B: macrolide-lincosamide-

streptogramin B; Tetra: tetracyclines; Fluoro: fluoroquinolones; Amino:

aminoglycoside; Pleuro: pleuromutilin; TMP: trimethoprim and Phen:

phenicol). In the 16 square panels, the left-hand histograms (labelled 0) show

the MIC values for isolates that carry no determinant for that antibiotic class,

while the right-hand histograms (labelled 1+) show the MIC values for isolates

carrying one or more such determinant. If all resistance determinants perfectly

explain MIC, then we expect to see histogram distributions on the bottom left

(low MIC, no determinant) and the top right (high MIC, presence of candidate

determinant(s)). For the first antimicrobial in each class, we show the number

of candidate AMR determinants in square brackets, along with the number of

isolates where candidate determinants are absent or present. r^2 values show

the proportion of the variance explained in a standard ANOVA by the

presence of one or more candidate determinant.

Figure 2. Differences in MICs between subsets of the data. Each row

compares two subsets of the isolates: (a) the 423 isolates from UK pigs, and

the 205 isolates from Canadian pigs. (b) the 50 isolates from Vietnam (all of

which are from the genetic cluster "BAPS4"), and the 112 BAPS4 isolates

from the UK and Canada. (c) the 22 Vietnamese isolates from pigs, and the

28 Vietnamese isolates from people. Left-hand panels compare the mean log

MICs for each antibiotic. Consistent deviations from the dotted 1:1 line

suggest consistently higher or lower MICs in that subset of the data (so the

rightward shift in panel (a) shows that MICs are consistently higher in

Canada). Right-hand panels show the proportion of isolates that carry each of

the 43 candidate AMR determinants. Each point or bar is coloured according

to its drug class, according to the colour scheme in Figure 1.

Figure 3. High levels of multidrug resistance in *S. suis*. The upper panel

shows the number of our 678 isolates that carry a given number of candidate

AMR determinants. The low panel shows the number of isolates that carry

one or more AMR determinant for a given number of drug classes. Results

show that more isolates carry determinants against 5 drug classes than carry

no determinant at all.

Figure 4. Beta-lactam resistance determinants have additive effects, and

appear in a consistent order. The left-hand panel shows that the candidate

determinants against the beta-lactam drug class often appear in a consistent

order. For example, the mutation at site 551 of PBP2X tends to appear in

backgrounds that already carry mutations in PBP2B. The plot was generated

according to the method and plotting convention of [44], where each row

represents an isolate, and those isolates that fit the nested pattern are shown

in pink. Results show that 93.5% (634/678) of our isolates fit the nested

pattern. The right-hand plots show the log MIC values for these 634 isolates,

comparing isolates carrying different numbers of candidate determinants.

1.   de Moor CE: **Septicaemic infections in pigs, caused by haemolytic**
streptococci of new Lancefield groups designated R, S and T. *Antonie*
van Leeuwenhoek 1963, **29**(1):272-280.
2.   Wertheim HFL, Nguyen HN, Taylor W, Lien TTM, Ngo HT, Nguyen TQ,
Nguyen BNT, Nguyen HH, Nguyen HM, Nguyen CT *et al*: **Streptococcus**
suis, an Important Cause of Adult Bacterial Meningitis in Northern
Vietnam. *PloS one* 2009, **4**(6):e5973.
3.   Baums CG, Kock C, Beineke A, Bennecke K, Goethe R, Schröder C,
Waldmann K-H, Valentin-Weigand P: **Streptococcus suis bacterin and**
subunit vaccine immunogenicities and protective efficacies against
serotypes 2 and 9. *Clinical and vaccine immunology* 2009, **16**(2):200-
208.
4.   van Rennings L, von Münchhausen C, Ottilie H, Hartmann M, Merle R,
Honscha W, Käsbohrer A, Kreienbrock L: **Cross-sectional study on**
antibiotic usage in pigs in Germany. *PloS one* 2015, **10**(3):e0119114.
5.   McGlone JJ: **The Future of Pork Production in the World: Towards**
Sustainable, Welfare-Positive Systems. *Animals* 2013, **3**(2):401-415.
6.   Steinfeld H, Gerber PJ, Wassenaar T, Castel V, Rosales M, De haan C:
Livestock's Long Shadow: Environmental Issues and Options, vol. 24;
2006.
7.   Goyette-Desjardins G, Auger J-P, Xu J, Segura M, Gottschalk M:
Streptococcus suis, an important pig pathogen and emerging
zoonotic agent—an update on the worldwide distribution based on
serotyping and sequence typing. *Emerging Microbes and Infections*
2014, **3**(1):1-20.
8.   Van Boeckel TP, Brower C, Gilbert M, Grenfell BT, Levin SA, Robinson TP,
Teillant A, Laxminarayan R: **Global trends in antimicrobial use in food**
animals. *Proceedings of the National Academy of Sciences* 2015,
112(18):5649-5654.
9.   Hernandez-Garcia J, Wang J, Restif O, Holmes MA, Mather AE, Weinert LA,
Wileman TM, Thomson JR, Langford PR, Wren BW *et al*: **Patterns of**
antimicrobial resistance in Streptococcus suis isolates from pigs with
or without streptococcal disease in England between 2009 and 2014.
Vet Microbiol 2017, **207**:117-124.
10. Yongkiettrakul S, Maneerat K, Arechanajan B, Malila Y, Srimanote P,
Gottschalk M, Visessanguan W: **Antimicrobial susceptibility of**
Streptococcus suis isolated from diseased pigs, asymptomatic pigs,
and human patients in Thailand. *BMC Veterinary Research* 2019,
15(1):5.
11. Marie J, Morvan H, Berthelot-Hérault F, Sanders P, Kempf I, Gautier-
Bouchardon AV, Jouy E, Kobisch M: **Antimicrobial susceptibility of**
Streptococcus suis isolated from swine in France and from humans in
different countries between 1996 and 2000. *Journal of Antimicrobial*
Chemotherapy 2002, **50**(2):201-209.
12. Šeol B, Kelnerić Ž, Hajsig D, Madić J, Naglić T: **Susceptibility to**
Antimicrobial Agents of Streptococcus suis Capsular Type 2 Strains
Isolated from Pigs. *Zentralblatt für Bakteriologie* 1996, **283**(3):328-331.

13.   Turgeon PL, Higgins R, Gottschalk M, Beaudoin M: **Antimicrobial**
susceptibility of *Streptococcus suis* **isolates**. *British Veterinary Journal*
1994, **150**(3):263-269.
14.   Gurung M, Tamang MD, Moon DC, Kim S-R, Jeong J-H, Jang G-C, Jung S-C,
Park Y-H, Lim S-K: **Molecular Basis of Resistance to Selected**
Antimicrobial Agents in the Emerging Zoonotic Pathogen
Streptococcus suis. *Journal of Clinical Microbiology* 2015, **53**(7):2332-
2336.
15.   Yao J, Shang K, Huang J, Ran W, Kashif J, Wang L: **Overexpression of an**
ABC transporter and mutations of GyrA, GyrB, and ParC in
contributing to high-level ciprofloxacin resistance in *Streptococcus*
suis **type 2**. *BioScience Trends* 2014, **8**(2):84-92.
16.   Hoa NT, Chieu TT, Nghia HD, Mai NT, Anh PH, Wolbers M, Baker S,
Campbell JI, Chau NV, Hien TT *et al*: **The antimicrobial resistance**
patterns and associated determinants in *Streptococcus suis* **isolated**
from humans in southern Vietnam, 1997-2008. *BMC infectious diseases*
2011, **11**:6.
17.   Huang J, Ma J, Shang K, Hu X, Liang Y, Li D, Wu Z, Dai L, Chen L, Wang L:
Evolution and Diversity of the Antimicrobial Resistance Associated
Mobilome in *Streptococcus suis*: **A Probable Mobile Genetic Elements**
Reservoir for Other Streptococci. *Frontiers in Cellular and Infection*
Microbiology 2016, **6**(118).
18.   Hu P, Yang M, Zhang A, Wu J, Chen B, Hua Y, Yu J, Chen H, Xiao J, Jin M:
Comparative genomics study of multi-drug-resistance mechanisms
in the antibiotic-resistant *Streptococcus suis* **R61 strain**. *PloS one*
2011, **6**(9):e24988.
19.   Holden MT, Hauser H, Sanders M, Ngo TH, Cherevach I, Cronin A,
Goodhead I, Mungall K, Quail MA, Price C *et al*: **Rapid evolution of**
virulence and drug resistance in the emerging zoonotic pathogen
Streptococcus suis. *PloS one* 2009, **4**(7):e6072.
20.   Weinert LA, Chaudhuri RR, Wang J, Peters SE, Corander J, Jombart T, Baig
A, Howell KJ, Vehkala M, Välimäki N *et al*: **Genomic signatures of human**
and animal disease in the zoonotic pathogen *Streptococcus suis*.
Nature Communications 2015, **6**(1):6740.
21.   Wileman TM, Weinert LA, Howell KJ, Wang J, Peters SE, Williamson SM,
Wells JM, Langford PR, Rycroft AN, Wren BW *et al*: **Pathotyping the**
Zoonotic Pathogen *Streptococcus suis*: **Novel Genetic Markers To**
Differentiate Invasive Disease-Associated Isolates from Non-Disease-
Associated Isolates from England and Wales. *Journal of Clinical*
Microbiology 2019, **57**(7):e01712-01718.
22.   Zou G, Zhou J, Xiao R, Zhang L, Cheng Y, Jin H, Li L, Zhang L, Wu B, Qian P
et al: **Effects of Environmental and Management-Associated Factors**
on Prevalence and Diversity of *Streptococcus suis* **in Clinically**
Healthy Pig Herds in China and the United Kingdom. *Applied and*
Environmental Microbiology 2018, **84**(8):e02590-02517.
23.   van Hout J, Heuvelink A, Gonggrijp M: **Monitoring of antimicrobial**
susceptibility of *Streptococcus suis* **in the Netherlands, 2013–2015**.
Vet Microbiol 2016, **194**:5-10.

24. Dunlop RH, McEwen SA, Meek AH, Black WD, Clarke RC, Friendship RM:
Individual and group antimicrobial usage rates on 34 farrow-to-
finish swine farms in Ontario, Canada. *Preventive Veterinary Medicine*
1998, **34**(4):247-264.
25. Directorate VM: **UK veterinary antibiotic resistance and sales**
surveillance report. In.; 2018.
26. Carrique-Mas JJ, Choisy M, Van Cuong N, Thwaites G, Baker S: **An**
estimation of total antimicrobial usage in humans and animals in
Vietnam. *Antimicrobial Resistance and Infection Control* 2020, **9**(1):16.
27. Jia B, Raphenya AR, Alcock B, Waglechner N, Guo P, Tsang KK, Lago BA,
Dave BM, Pereira S, Sharma AN *et al*: **CARD 2017: expansion and**
model-centric curation of the comprehensive antibiotic resistance
database. *Nucleic Acids Res* 2017, **45**(D1):D566-D573.
28. Ge Y, Wu J, Xia Y, Yang M, Xiao J, Yu J: **Molecular dynamics simulation of**
the complex PBP-2x with drug cefuroxime to explore the drug
resistance mechanism of *Streptococcus suis* **R61**. *PloS one* 2012,
7(4):e35941.
29. Palmieri C, Magi G, Mingoia M, Bagnarelli P, Ripa S, Varaldo PE, Facinelli B:
Characterization of a *Streptococcus suis* **tet(O/W/32/O)-carrying**
element transferable to major streptococcal pathogens. *Antimicrobial*
agents and chemotherapy 2012, **56**(9):4697-4702.
30. Escudero JA, San Millan A, Catalan A, de la Campa AG, Rivero E, Lopez G,
Dominguez L, Moreno MA, Gonzalez-Zorn B: **First characterization of**
fluoroquinolone resistance in *Streptococcus suis*. *Antimicrobial agents*
and chemotherapy 2007, **51**(2):777-782.
31. Vannice KS, Ricaldi J, Nanduri S, Fang FC, Lynch JB, Bryson-Cahn C, Wright
T, Duchin J, Kay M, Chochua S *et al*: *Streptococcus pyogenes* **pbp2x**
Mutation Confers Reduced Susceptibility to β-Lactam Antibiotics.
Clinical infectious diseases : an official publication of the Infectious Diseases
Society of America 2020, **71**(1):201-204.
32. Chewapreecha C, Marttinen P, Croucher NJ, Salter SJ, Harris SR, Mather
AE, Hanage WP, Goldblatt D, Nosten FH, Turner C *et al*: **Comprehensive**
identification of single nucleotide polymorphisms associated with
beta-lactam resistance within pneumococcal mosaic genes. *PLoS*
genetics 2014, **10**(8):e1004547.
33. Albarracin Orio AG, Pinas GE, Cortes PR, Cian MB, Echenique J:
Compensatory evolution of pbp mutations restores the fitness cost
imposed by beta-lactam resistance in *Streptococcus pneumoniae*.
PLoS pathogens 2011, **7**(2):e1002000.
34. Leclercq R: **Mechanisms of Resistance to Macrolides and**
Lincosamides: Nature of the Resistance Elements and Their Clinical
Implications. *Clinical Infectious Diseases* 2002, **34**(4):482-492.
35. Farrow KA, Lyras D, Polekhina G, Koutsis K, Parker MW, Rood JI:
Identification of Essential Residues in the Erm(B) rRNA
Methyltransferase of *Clostridium perfringens*. *Antimicrobial agents and*
chemotherapy 2002, **46**(5):1253.
36. Grebe T, Hakenbeck R: **Penicillin-binding proteins 2b and 2x of**
Streptococcus pneumoniae **are primary resistance determinants for**

different classes of beta-lactam antibiotics. *Antimicrobial agents and*
chemotherapy 1996, **40**(4):829-834.

37.   Li Y, Metcalf BJ, Chochua S, Li Z, Gertz RE, Walker H, Hawkins PA, Tran T,
Whitney CG, McGee L *et al*: **Penicillin-Binding Protein Transpeptidase**
Signatures for Tracking and Predicting β-Lactam Resistance Levels
in ***Streptococcus pneumoniae***. *mBio* 2016, **7**(3):e00756-00716.

38.   Dahesh S, Hensler ME, Van Sorge NM, Gertz RE, Jr., Schrag S, Nizet V, Beall
BW: **Point mutation in the group B streptococcal pbp2x gene**
conferring decreased susceptibility to beta-lactam antibiotics.
Antimicrobial agents and chemotherapy 2008, **52**(8):2915-2918.

39.   Fuursted K, Stegger M, Hoffmann S, Lambertsen L, Andersen PS, Deleuran
M, Thomsen MK: **Description and characterization of a penicillin-**
resistant ***Streptococcus dysgalactiae*** **subsp.** ***equisimilis*** **clone isolated**
from blood in three epidemiologically linked patients. *Journal of*
Antimicrobial Chemotherapy 2016, **71**(12):3376-3380.

40.   Holmer I, Salomonsen CM, Jorsal SE, Astrup LB, Jensen VF, Høg BB,
Pedersen K: **Antibiotic resistance in porcine pathogenic bacteria and**
relation to antibiotic usage. *BMC Veterinary Research* 2019, **15**(1):449.

41.   Liñares J, Ardanuy C, Pallares R, Fenoll A: **Changes in antimicrobial**
resistance, serotypes and genotypes in ***Streptococcus pneumoniae***
over a 30-year period. *Clinical Microbiology and Infection* 2010,
16(5):402-410.

42.   Mouz N, Di Guilmi AM, Gordon E, Hakenbeck R, Dideberg O, Vernet T:
Mutations in the active site of penicillin-binding protein PBP2x from
Streptococcus pneumoniae. **Role in the specificity for beta-lactam**
antibiotics. *The Journal of biological chemistry* 1999, **274**(27):19175-
19180.

43.   Nagai K, Davies TA, Jacobs MR, Appelbaum PC: **Effects of amino acid**
alterations in penicillin-binding proteins (PBPs) 1a, 2b, and 2x on
PBP affinities of penicillin, ampicillin, amoxicillin, cefditoren,
cefuroxime, cefprozil, and cefaclor in 18 clinical isolates of
penicillin-susceptible, -intermediate, and -resistant pneumococci.
Antimicrobial agents and chemotherapy 2002, **46**(5):1273-1280.

44.   Lehtinen S, Blanquart F, Lipsitch M, Fraser C, with the Maela
Pneumococcal Collaboration: **On the evolutionary ecology of multidrug**
resistance in bacteria. *PLoS pathogens* 2019, **15**(5):e1007763.

45.   Calvez P, Breukink E, Roper DI, Dib M, Contreras-Martel C, Zapun A:
Substitutions in PBP2b from beta-Lactam-resistant ***Streptococcus***
pneumoniae **Have Different Effects on Enzymatic Activity and Drug**
Reactivity. *Journal of Biological Chemistry* 2017, **292**(7):2854-2865.

46.   Figueiredo TA, Sobral RG, Ludovice AM, Almeida JM, Bui NK, Vollmer W,
de Lencastre H, Tomasz A: **Identification of genetic determinants and**
enzymes involved with the amidation of glutamic acid residues in
the peptidoglycan of ***Staphylococcus aureus***. *PLoS pathogens* 2012,
8(1):e1002508.

47.   Zapun A, Philippe J, Abrahams KA, Signor L, Roper DI, Breukink E, Vernet
T: **In vitro reconstitution of peptidoglycan assembly from the Gram-**
positive pathogen ***Streptococcus pneumoniae***. *ACS chemical biology*
2013, **8**(12):2688-2696.

48. Kocaoglu O, Tsui HC, Winkler ME, Carlson EE: **Profiling of beta-lactam**
selectivity for penicillin-binding proteins in *Streptococcus*
***pneumoniae* D39**. *Antimicrobial agents and chemotherapy* 2015,
59(6):3548-3555.
49. Coffey TJ, Daniels M, McDougal LK, Dowson CG, Tenover FC, Spratt BG:
Genetic analysis of clinical isolates of *Streptococcus pneumoniae* with
high-level resistance to expanded-spectrum cephalosporins.
Antimicrobial agents and chemotherapy 1995, **39**(6):1306-1313.
50. Sakoulas G, Moellering RC, Jr.: **Increasing Antibiotic Resistance among**
Methicillin-Resistant *Staphylococcus aureus* Strains. *Clinical Infectious*
Diseases 2008, **46**(Supplement_5):S360-S367.
51. van Duijkeren E, Greko C, Pringle M, Baptiste KE, Catry B, Jukes H, Moreno
MA, Pomba MCMF, Pyörälä S, Rantala M *et al*: **Pleuromutilins: use in**
food-producing animals in the European Union, development of
resistance and impact on human and animal health. *Journal of*
Antimicrobial Chemotherapy 2014, **69**(8):2022-2031.
52. Metcalf BJ, Chochua S, Gertz RE, Jr., Li Z, Walker H, Tran T, Hawkins PA,
Glennen A, Lynfield R, Li Y *et al*: **Using whole genome sequencing to**
identify resistance determinants and predict antimicrobial
resistance phenotypes for year 2015 invasive pneumococcal disease
isolates recovered in the United States. *Clinical microbiology and*
infection : the official publication of the European Society of Clinical
Microbiology and Infectious Diseases 2016, **22**(12):1002.e1001-
1002.e1008.
53. Dunai A, Spohn R, Farkas Z, Lázár V, Györkei Á, Apjok G, Boross G,
Szappanos B, Grézal G, Faragó A *et al*: **Rapid decline of bacterial drug-**
resistance in an antibiotic-free environment through phenotypic
reversion. *Elife* 2019, **8**:e47088.
54. Kime L, Randall CP, Banda FI, Coll F, Wright J, Richardson J, Empel J,
Parkhill J, O'Neill AJ: **Transient Silencing of Antibiotic Resistance by**
Mutation Represents a Significant Potential Source of Unanticipated
Therapeutic Failure. *mBio* 2019, **10**(5):e01755-01719.
55. Li Y, Metcalf BJ, Chochua S, Li Z, Gertz RE, Jr., Walker H, Hawkins PA, Tran
T, McGee L, Beall BW: **Validation of β-lactam minimum inhibitory**
concentration predictions for pneumococcal isolates with newly
encountered penicillin binding protein (PBP) sequences. *BMC*
genomics 2017, **18**(1):621.
56. Beall B, Chochua S, Gertz RE, Jr., Li Y, Li Z, McGee L, Metcalf BJ, Ricaldi J,
Tran T, Walker H *et al*: **A Population-Based Descriptive Atlas of**
Invasive Pneumococcal Strains Recovered Within the U.S. During
2015-2016. *Frontiers in microbiology* 2018, **9**:2670.
57. Varghese J, Chochua S, Tran T, Walker H, Li Z, Snippes Vagnone PM,
Lynfield R, McGee L, Li Y, Metcalf BJ *et al*: **Multistate population and**
whole genome sequence-based strain surveillance of invasive
pneumococci recovered in the USA during 2017. *Clinical microbiology*
and infection : the official publication of the European Society of Clinical
Microbiology and Infectious Diseases 2020, **26**(4):512.e511-512.e510.
58. Metcalf BJ, Chochua S, Gertz RE, Jr., Hawkins PA, Ricaldi J, Li Z, Walker H,
Tran T, Rivers J, Mathis S *et al*: **Short-read whole genome sequencing**

		for determination of antimicrobial resistance mechanisms and capsular serotypes of current invasive *Streptococcus agalactiae* recovered in the USA. *Clinical microbiology and infection : the official publication of the European Society of Clinical Microbiology and Infectious Diseases* 2017, **23**(8):574.e577-574.e514.
1008	59.	de Jong A, Thomas V, Simjee S, Moyaert H, El Garch F, Maher K, Morrissey I, Butty P, Klein U, Marion H *et al*: **Antimicrobial susceptibility monitoring of respiratory tract pathogens isolated from diseased cattle and pigs across Europe: The VetPath study**. *Vet Microbiol* 2014, **172**(1):202-215.
1013	60.	Quail MA, Kozarewa I, Smith F, Scally A, Stephens PJ, Durbin R, Swerdlow H, Turner DJ: **A large genome center's improvements to the Illumina sequencing system**. *Nature Methods* 2008, **5**(12):1005-1010.
1016	61.	Joshi N, Fass J: **Sickle: A sliding-window, adaptive, quality-based trimming tool for FastQ files**. In.; 2011.
1018	62.	Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, Lesin VM, Nikolenko SI, Pham S, Prjibelski AD *et al*: **SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing**. *Journal of Computational Biology* 2012, **19**(5):455-477.
1022	63.	Seemann T: **Prokka: rapid prokaryotic genome annotation**. *Bioinformatics* 2014, **30**(14):2068-2069.
1024	64.	Wingett SW, Andrews S: **FastQ Screen: A tool for multi-genome mapping and quality control**. *F1000Res* 2018, **7**:1338-1338.
1026	65.	Li H, Durbin R: **Fast and accurate short read alignment with Burrows–Wheeler transform**. *Bioinformatics* 2009, **25**(14):1754-1760.
1028	66.	Cheng L, Connor TR, Sirén J, Aanensen DM, Corander J: **Hierarchical and Spatially Explicit Clustering of DNA Sequences with BAPS Software**. *Molecular biology and evolution* 2013, **30**(5):1224-1228.
1031	67.	Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MTG, Fookes M, Falush D, Keane JA, Parkhill J: **Roary: rapid large-scale prokaryote pan genome analysis**. *Bioinformatics* 2015, **31**(22):3691-3693.
1034	68.	Ranwez V, Harispe S, Delsuc F, Douzery EJP: **MACSE: Multiple Alignment of Coding SEquences Accounting for Frameshifts and Stop Codons**. *PloS one* 2011, **6**(9):e22594.
1037	69.	Hunt M, Mather AE, Sánchez-Busó L, Page AJ, Parkhill J, Keane JA, Harris SR: **ARIBA: rapid antimicrobial resistance genotyping directly from sequencing reads**. *Microb Genom* 2017, **3**(10):e000131.
1040	70.	Lopez P, Espinosa M, Greenberg B, Lacks St: **Sulfonamide resistance in *Streptococcus pneumoniae*: DNA sequence of the gene encoding dihydropteroate synthase and characterization of the enzyme**. *Journal of Bacteriology* 1987, **169**(9):4320-4326.
1044	71.	Adrian PV, Klugman KP: **Mutations in the dihydrofolate reductase gene of trimethoprim-resistant isolates of *Streptococcus pneumoniae***. *Antimicrobial agents and chemotherapy* 1997, **41**(11):2406-2413.
1048	72.	Hakenbeck R, Grebe T, Zähner D, Stock JB: **β-Lactam resistance in *Streptococcus pneumoniae*: penicillin-binding proteins and non-penicillin-binding proteins**. *Molecular Microbiology* 1999, **33**(4):673-678.

73.   Edgar RC: **MUSCLE: multiple sequence alignment with high accuracy**
and high throughput. *Nucleic Acids Res* 2004, **32**(5):1792-1797.
74.   Gouy M, Guindon S, Gascuel O: **SeaView version 4: A multiplatform**
graphical user interface for sequence alignment and phylogenetic
tree building. *Molecular biology and evolution* 2010, **27**(2):221-224.
75.   Contreras-Martel C, Dahout-Gonzalez C, Martins Ados S, Kotnik M, Dessen
A: **PBP active site flexibility as the key mechanism for beta-lactam**
resistance in pneumococci. *Journal of molecular biology* 2009,
387(4):899-909.
76.   Haenni M, Galofaro L, Ythier M, Giddey M, Majcherczyk P, Moreillon P,
Madec JY: **Penicillin-binding protein gene alterations in *Streptococcus***
***uberis* isolates presenting decreased susceptibility to penicillin**.
Antimicrobial agents and chemotherapy 2010, **54**(3):1140-1145.
77.   Piccinelli G, Carlentini G, Gargiulo F, Caruso A, De Francesco MA: **Analysis**
of Point Mutations in the pbp2x, pbp2b, and pbp1a Genes of
***Streptococcus agalactiae* and Their Relation with a Reduced**
Susceptibility to Cephalosporins. *Microbial Drug Resistance* 2017,
23(8):1019-1024.
78.   Haroche J, Allignet J, Buchrieser C, El Solh N: **Characterization of a**
Variant of vga (A) Conferring Resistance to Streptogramin A and
Related Compounds. *Antimicrobial agents and chemotherapy* 2000,
44(9):2271.
79.   Pikis A, Donkersloot JA, Rodriguez WJ, Keith JM: **A conservative amino**
acid mutation in the chromosome-encoded dihydrofolate reductase
confers trimethoprim resistance in *Streptococcus pneumoniae*.
Journal of Infectious Diseases 1998, **178**(3):700-706.
80.   Bergmann R, van der Linden M, Chhatwal GS, Nitsche-Schmitz DP:
Factors that cause trimethoprim resistance in *Streptococcus*
pyogenes. *Antimicrobial agents and chemotherapy* 2014, **58**(4):2281-
2288.
81.   Athey TBT, Teatero S, Lacouture S, Takamatsu D, Gottschalk M, Fittipaldi
N: **Determining *Streptococcus suis* serotype from short-read whole-**
genome sequencing data. *BMC Microbiol* 2016, **16**(1):162-162.
82.   Team RC: **R: A language and environment for statistical computing**.
2013.
83.   Pinheiro J, Bates D, DebRoy S, Sarkar D, Heisterkamp S, Van Willigen B,
Maintainer R: **Package 'nlme'**. 2019.
84.   Palmieri C, Varaldo PE, Facinelli B: ***Streptococcus suis*, an Emerging**
Drug-Resistant Animal and Human Pathogen. *Frontiers in microbiology*
2011, **2**:235.
85.   Huang K, Zhang Q, Song Y, Zhang Z, Zhang A, Xiao J, Jin M:
Characterization of Spectinomycin Resistance in *Streptococcus suis*
Leads to Two Novel Insights into Drug Resistance Formation and
Dissemination Mechanism. *Antimicrobial agents and chemotherapy*
2016, **60**(10):6390-6392.
86.   Chang B, Wada A, Ikebe T, Ohnishi M, Mita K, Endo M, Matsuo H, Asatuma
Y, Kuramoto S, Sekiguchi H *et al*: **Characteristics of *Streptococcus suis***
isolated from patients in Japan. *Japanese journal of infectious diseases*
2006, **59**(6):397-399.

87. Bojarska A, Molska E, Janas K, Skoczynska A, Stefaniuk E, Hryniewicz W,
Sadowy E: *Streptococcus suis* **in invasive human infections in Poland:**
clonality and determinants of virulence and antimicrobial
resistance. *European Journal of Clinical Microbiology and Infectious*
Diseases 2016, **35**(6):917-925.
88. Athey TB, Auger JP, Teatero S, Dumesnil A, Takamatsu D, Wasserscheid J,
Dewar K, Gottschalk M, Fittipaldi N: **Complex Population Structure and**
Virulence Differences among Serotype 2 *Streptococcus suis* **Strains**
Belonging to Sequence Type 28. *PloS one* 2015, **10**(9):e0137760.
1110

log(MIC)

Amoxicillin r^2=0.143
Cefquinome r^2=0.284
Ceftiofur r^2=0.516
Penicillin r^2=0.513
Erythromy. r^2=0.802
Lincomycin r^2=0.78
Tilmicosin r^2=0.751
Tylosin r^2=0.789
Doxycycline r^2=0.875
Tetracycline r^2=0.837
Enrofloxa. r^2=0.002
Marbofloxa. r^2=0.003
Spectinomy. r^2=0.371
Tiamulin r^2=0.642
Trimethop. r^2=0.538
Florifenicol r^2=0

[6] (468−/210+)
[8] (283−/395+)
[11] (104−/574+)
[3] (662−/16+)
[8] (550−/128+)
[2] (485−/193+)
[4] (432−/246+)
[1] (675−/3+)