A novel platform to accelerate antimicrobial susceptibility testing in *Neisseria gonorrhoeae*

using RNA signatures

Marjan M.Hashemi[1], Nikhil Ram-Mohan[1], Xi Yang[1], Nadya Andini[1], Nicholas R. Gessner[1],

Karen C. Carroll[2], Tza-Huei Wang[3], Samuel Yang[1]*

[1]Department of Emergency Medicine, Stanford University, Stanford, CA, USA

[2]Department of Pathology, Johns Hopkins University, Baltimore, MD, USA

[3]Department of Mechanical Engineering, Johns Hopkins University, Baltimore, MD, USA

*Corresponding author Email: syang5@stanford.edu, Phone: (650) 725-9492, Fax: (650) 723-

0121

Running title: RNA markers for AST in *Neisseria gonorrhoeae*

Abstract

The rise of antimicrobial-resistant pathogens can be attributed to the lack of a rapid pathogen

identification (ID) or antimicrobial susceptibility testing (AST), resulting in delayed therapeutic

decisions at the point of care. Gonorrhea is usually empirically treated with no AST results

available before treatment, thus contributing to the rapid rise in drug resistance. Herein we

present a rapid AST platform using RNA signatures for *Neisseria gonorrhoeae* (NG). RNA-seq

followed by bioinformatic tools were applied to explore potential markers in the transcriptome

profile of NG upon minutes of azithromycin exposure. Validation of candidate markers using

PCR showed that two markers (*arsR* (NGO1562) and *rpsO*) can deliver accurate AST results

across 14 tested isolates. Further validation of our cutoff in comparison to MIC across 64 more

isolates confirmed the reliability of our platform. Our RNA markers combined with emerging

molecular point-of-care systems has the potential to greatly accelerate both ID and AST to

inform treatment.

Introduction

The use of conventional clinical methods for pathogen identification (ID) and antimicrobial

susceptibility testing (AST) is a time-consuming process that contributes to the rise of

antimicrobial resistance and may in turn increase mortality rates (1). Thus, expedition of

diagnostic methods for timely and directed therapy is an urgent clinical need that is to be

addressed.

Neisseria gonorrhoeae (NG), the causative agent of gonorrhea, is an increasingly common

sexually transmitted infection with more than 550, 000 cases reported annually in the United

States (2). Gonorrhea is usually seen in female cervicitis and male urethritis (3). Untreated

gonorrhea infections can lead to major complications such as heart and nervous system

infections, infertility, pelvic inflammatory disease, and newborn blindness (4, 5). Gonorrhea is

often empirically treated based on a clinical syndromic diagnosis or after laboratory molecular

detection with nucleic acid amplification testing (NAAT). NAAT have largely replaced

diagnostic culture methods because they are faster, automated, with higher sensitivity to allow

for cost-effective diagnosis (6). Treatments often proceed without AST results. NG AST requires

laborious and time-consuming (at least 1-2 days) culture methods and is only undertaken in cases

of treatment failure (7, 8).

NG can rapidly develop resistance to antimicrobial agents due to innate mechanisms for

acquiring resistance genes (9). Treatment is more challenging due to rapidly increasing

resistance to all of the most commonly used antimicrobials including sulfonamides, penicillin,

tetracyclines, and second-generation fluoroquinolones such as ciprofloxacin (10). Azithromycin

is a macrolide antimicrobial that binds to the 23S rRNA component of the 50S ribosome, thereby

inhibiting protein synthesis (11, 12). Azithromycin has been shown an effective treatment against

gonococci with prolonged level in tissues and cells. Following oral administration, azithromycin

concentrates in tissues including genital sites (12). However high levels of azithromycin

resistance have been progressively reported worldwide (13, 14). Some of the known mechanisms

of resistance to azithromycin include overexpression of the efflux pump (*mtrR*), 23S rRNA

mutation in azithromycin binding sites (C2611T and A2059G) and ribosomal target modification

by methylase (*ermC* and *ermB*) (3).

CDC reported antimicrobial resistant NG as one of the three most urgent threats to public health

and currently recommends dual therapy with ceftriaxone and azithromycin for the treatment of

uncomplicated gonorrhea (15). Despite recommended dual therapy, azithromycin monotherapy

could be still used to treat uncomplicated gonorrhea in cephalosporin-allergic patients (16). In

this study azithromycin has been selected for our phenotypic AST development for gonorrhea as

a proof of concept; however our approach has potential to be extended to other antimicrobials.

Given the time-consuming conventional AST, increasing level of resistance, and the challenge of

slow growth rate (doubling time 60min) of NG (17), development of a novel AST that can guide

initial treatment at the point of care is critically needed. Recently, RNA-seq has been effectively

applied in discovery of reliable genomic biomarkers to develop clinical diagnosis (18). Previous

work in our lab described a novel and accelerated AST workflow based on RNA-seq in

Klebsiella pneumoniae upon exposure to ciprofloxacin (19). This method is independent of cell

division unlike other molecular phenotypic AST that measure growth kinetics of bacteria based

on DNA copies (20, 21). As proof-of-concept, in this study we focused on developing an ultra-

rapid molecular phenotypic AST for NG. RNA-seq followed by bioinformatic analysis was used

to identify candidate diagnostic RNA markers to determine susceptibility upon a short exposure

to azithromycin. Further validation of selected markers was performed through qRT-PCR using

14 isolates followed by validation of our cutoff in comparison to reported MICs using 64 more

isolates to determine susceptibility.

Material and methods

Microorganisms and culturing

Reference strains of azithromycin susceptible and resistant NG, SPL-4 and SPJ-15, were

obtained from CDC. Clinical isolates of NG were obtained from John Hopkins Medical

Institutions, Department of Pathology, Division of Medical Microbiology (Table S1). Isolates

were cultured from glycerol stocks on GC agar (#BD 228950, Becton Dickinson, Cockeysville,

MD) supplemented with 1% IsoVitaleX (#BD 211876, Becton Dickinson, Cockeysville, MD).

Plates were incubated at 35 °C, in 5% CO_2 for 24 h. A single colony of each isolate was re-

suspended in GW broth (22) and incubated overnight at 37 °C, 5% CO_2, 200 rpm to an optical

density equal to that of a 0.5 MacFarland standard.

Antimicrobial susceptibility testing

The E-test, bioMérieux (Durham, NC, USA), or agar dilution method, was used to determine the

minimum inhibitory concentration (MIC) of azithromycin (Astatech, Inc, Bristol, PA) (23). Agar

dilution method was performed using GC agar supplemented with 1% IsoVitaleX described by

the Clinical and Laboratory Standards Institute (CLSI). Briefly, two-fold dilutions of

azithromycin (range 0.03-256 µg/mL) was added to CG medium (24). Each plate was inoculated

with a 0.5 McFarland standard of bacterial suspension. Inoculated plates were incubated at 35

°C, in 5% CO_2 for 24 h. Measurements were performed in triplicate and positive and negative

controls were included for each MIC test. Following incubation, MICs were determined as the

lowest concentrations of azithromycin that inhibited visible growth of bacteria. According to the

guidelines set forth by the CLSI, isolates with MIC ≤2 µg/mL are classified as azithromycin

susceptible (25).

RNA extraction and sequencing

Bacterial culture (2mL) was prepared as described previously. Samples were exposed to

azithromycin (2µg/mL) and incubated at 37 °C, in 5% CO_2 for 10 minutes and 60 minutes (see

workflow in Fig. 1). Bacteria without exposure to azithromycin were also incubated in the same

condition as a control (water was added to the controls). Samples were collected and preserved in

4mL RNA Protect Reagent (Qiagen, Valencia, CA, USA) at each time point. RNA extraction

was conducted using RNeasy Mini Kit (Qiagen,74524) according to the manufacturer's

instruction. Extracted RNA was subjected to DNase treatment (Turbo DNase complete kit Life

Technologies,   AM1907).   Concentration of RNA was measured using QIA

expert spectrophotometer (Qiagen, USA). To consider biological variability triplicate RNA

samples were prepared from susceptible and resistant isolates.

Sample preparation for sequencing was performed using ScriptSeq complete kit for bacteria with

rRNA depleted (Illumina Inc, BB1224, RSBC10948 and SSIP1202). Samples were bio-analyzed

at Stanford Functional Genomics Facility (SFGF) and libraries were sequenced at 75 pair-end

reads on Illumina MiSeq. Quality control on raw reads from the sequenced libraries was

conducted to remove the low-quality reads (using FastQC) (26), remove PCR duplicates (using

SuperDeduper) (27) and trim the adaptor sequences (using Trim Galore, version 0.4.5) (28).

Finally, reads were aligned to NG FA 1090 (NCBI Reference Sequence: NC_002946.2RNA-seq

data are archived in the GenBank SRA (BioProject ID: PRJNA627416).

Differential expression analysis

Quantification of transcript abundance and differential expression analyses were carried out

using Rockhopper (29). Spearman correlation between samples/replicates from the normalized

transcript counts obtained from Rockhopper were calculated using the cor function in R (30).

The multidimensional scaling (MDS) that show the dissimilarity of the replicates/samples by

projecting them into two dimensions was also done using the cmdscale function.

To further analyze the differentially expressed genes under azithromycin treatment, the list was

cut down to 568 differentially expressed genes with a p-value<0.05 in susceptible compared to

control (SvC) and resistant (SvR) strains. Heatmaps were generated using the heatmap.2 function

to represent the logFC between these comparisons.

Marker selection and validation

Candidate markers were selected from the RNA-seq dataset based on the following criteria: 1)

significant fold change in susceptible but not in resistant after 10min drug exposure; 2) high level

of expression after 10min drug exposure; 3) same trend of expression in 10&60 min drug

exposure. Selected candidate markers were further evaluated using qPCR.

Validation of final markers in clinical isolates was performed using quantitative real-time PCR

(qRT-PCR). Primer-blast was used to design primers for conserved regions of selected genes

(Table S2). Quantification of gene expression level was conducted with Rotor-Gene SYBR

Green RT-PCR kit (Qiagen, cat# 204174) on the Rotor-Gene-Q PCR cycler (Qiagen). The

concentration of component used in final 25μL PCR reaction mix were as follows: 1x SYBR

Green Master mix, 1mM forward primer, 1mM reverse primer, 0.05U/μL Rotor Gene RT mix

and 0.2U/μL SUPERase-in RNase inhibitor and 8% RNA template.

Expression of each candidate marker was quantified in the control and treated samples.

Threshold of 0.1 was set to obtain accurate CT values for each isolate. This threshold is a point at

which amplification plot reaches a fluorescent intensity above background level (31). To

calculate ΔCT value, a housekeeping gene was used to normalize potential differences between

the control and treated samples. From our RNA-seq results, *atpA* gene was selected as an internal

control as its expression level did not affect significantly upon antimicrobial treatment for the

tested isolates in this study. Finally, fold change (FC) in gene expression between the control and

treated samples was calculated as follows: $FC=2^{(-\Delta\Delta CT)}$ (18).

Results

Shift in transcriptome response following azithromycin exposure

To identify markers that significantly differentiate susceptible and resistant isolates of *N.*

gonorrhoeae, RNA-seq was used to compare their transcriptome response after 10 and 60 min of

azithromycin exposure. A summary of workflow is shown in Fig. 1.

Multidimensional scaling analysis of the expression profile of the biological replicates, treatment

conditions, and the two strains revealed strong reproducibility amongst replicates and

dissimilarity between the conditions and strains (Fig. 2). MDS plot was also used to provide

insights into the association between transcriptional profile and exposure time (10 and 60 min) to

azithromycin. Biological replicates of control and treated samples clustered very closely

indicating high correlation among replicates and high reproducibility of library preparation.

MDS plot demonstrated four distinct transcriptional clusters within the first two dimensions

(dim1 and dim2), an indication of remarkable transcriptional changes upon antimicrobial

exposure. Well separation of the control and treated samples in the first dimension showed that

the sequencing data were qualified for identification of differentially expressed genes.

Additionally, a significant diversity between the gene expression of 10min and 60 min treated

samples was observed suggesting that distinct gene expression profiles are triggered by

azithromycin in a time-dependent manner.

Subsequently, statistical analysis of differentially expressed genes following exposure to

azithromycin was calculated using logFC (gene expression ratio of treated compared to the

control) against their p-values <0.05 (Fig. 3A). Consistent with MDS results, a global shift in

SvC and SvR was observed as early as 10 min of azithromycin exposure, although distribution of

gene expression profile illustrated a higher magnitude of fold change at 60 min.

Marker selection and validation of candidate markers in *NG* clinical isolates

Differential expression analyses of the susceptible strain revealed more down than up regulated

genes at 10 minutes (185:129). At 60 minutes however, there are 386 and 212 genes up and

down regulated, respectively (Table S3). Since we are interested in markers that will successfully

determine susceptibility, we further screened for genes that were differentially expressed

between SvC and SvR. A p-value cutoff of 0.05 resulted in 568 genes (Fig. S1) which narrows

down to 48 when screening for abs(logFC) ≥1 under both conditions (Fig. 3B).

Among the differentially expressed genes with significant logFC, two more criteria including

high level of expression and same trend of expression in 10&60 min (see methods) were applied

to select the candidate markers. The initially selected markers include NGO0373, NGO1920,

NGO1562, NGO0191, NGO0405 and NGO1078, hereafter referred to as *ABC*, *bolA*, *arsR*, *rpsO*,

dinD and *acoT*, respectively (Table S2). These were tested in one susceptible and one resistant

strain to confirm the diagnostic potential of markers to characterize azithromycin susceptibility

using RT-qPCR. All six candidate RNA markers are highlighted in Table S4, among which *ABC*,

bolA, *arsR*, and *rpsO* are upregulated while *dinD* and *acoT* are downregulated. Further validation

of markers was conducted in eleven susceptible and one resistant clinical isolates of NG to

confirm that RNA-seq nominated markers can be applied in different strains (Fig. 4). We

calculated -△△CT values for samples following 10 min and 60 min exposure to azithromycin

and compared it between resistant and susceptible isolates for each marker. Overall -△△CT of all

six selected candidate markers was significantly different across resistant and susceptible isolates

indicating qualification of these candidate markers to be applied in our new AST platform.

Among six tested markers *arsR* and *rpsO* demonstrated consistent upregulation trends across all

tested isolates. Although *ABC* and *bolA* were also validated as significantly upregulated RNA

markers in most tested isolates, down regulation of these two markers was observed in a few

tested susceptible isolates. On the other hand, *dinD* and *acoT* were validated as significantly

downregulated RNA markers, however inconsistency of results from a few isolates was also

found. According to the validation results of six selected markers, further analysis was conducted

using *arsR* and *rpsO* as our final markers.

We examined the reliability of our method compared to the gold standard using relationship

between $2^{-\Delta\Delta CT}$ values of final markers and MIC in susceptible and resistant strains after 10min

exposure to azithromycin (Fig. 5A). For both *arsR* and *rpsO*, $2^{-\Delta\Delta CT}$ values significantly changed

between susceptible and resistant and notably no overlap in $2^{-\Delta\Delta CT}$ values was observed between

two groups of susceptible and resistant strains. Additionally, a $2^{-\Delta\Delta CT} = 2$ was defined as

threshold susceptibility for susceptible and non-susceptible isolates ($2^{-\Delta\Delta CT} \geq 2$ susceptible, 2^{-}

$^{\Delta\Delta CT} < 2$ non-susceptible). We also used a linear fitting to show how transcriptional response of

the final markers are associated to the MIC in susceptible and resistant strains. A strong negative

relationship was observed between $\log_2 2^{-\Delta\Delta CT}$ and $\log_2$ MIC values, (R^2 values 0.64 and 0.63 for

arsR and *rpsO* respectively), meaning $2^{-\Delta\Delta CT}$ value in susceptible strains is significantly higher

than that in resistant strains (Fig. 5B).

Finally, to validate our susceptibility threshold, we used MIC information of two panels of NG

including 64 clinical isolates from CDC (Antibiotic Resistance Isolate Bank) on the equation for

the line of best fit for our markers. Notably, *arsR* supported the delineated $2^{-\Delta\Delta CT}$ value by

correctly classifying susceptibility of all 64 CDC isolates based on our $2^{-\Delta\Delta CT}$ threshold as 51

susceptible strains and 13 resistant strains that was in agreement with CLSI susceptibility

category (MIC breakpoint of ≥2μg/ml) (Table S5). *rpsO* also accurately identified susceptibility

of most tested isolates with a broad range of MIC, although susceptible isolates with MIC at

1μg/ml were classified as non-susceptible isolates; however, this category is in agreement with

EUCAST breakpoint values (MIC breakpoint of ≥1μg/ml).

In summary, these observations confirmed the accurate application of $2^{-\Delta\Delta CT}$ values to categorize

susceptibility of isolates as a replacement of MIC in traditional AST.

Discussion

To stem the tide of multi-drug resistant NG, a rapid NAAT-based diagnostic with compatible

molecular phenotypic AST capability is critically needed to inform initial treatment decisions. In

this study, we have discovered azithromycin-susceptibility RNA markers and demonstrated their

early promise for ultra-rapid AST in NG.

Transcriptome profiling of susceptible and resistant NG upon exposure to azithromycin revealed

a significantly altered response as early as 10 min in SvC and SvR (Figs. 2 & 3). While

significantly different transcriptional response among NG was previously reported after 60 min

azithromycin exposure (32), our main objective was to discover the earliest susceptibility RNA

markers to minimize AST time. Therefore, two time points of 10min and 60min were included in

azithromycin treatment to be able to identify consistent and reliable RNA markers as part of

early cellular stress responses to antimicrobials. Unlike the previous reported studies at which

sublethal concentration of drugs was used (8, 32), we studied bacterial transcriptomic profile

following a high concentration of azithromycin (2μg/mL) to enhance the discovery of

differentially expressed genes (19). Six candidate markers were initially identified through RNA-

seq and subsequently tested in 14 isolates with qRT-PCR. *arsR* and *rpsO* were chosen as our two

final markers as they were able to consistently determine susceptibility based on CT value only

after 10min antimicrobial exposure (Fig. 4).

arsR gene (NGO1562) encodes a transcriptional regulatory protein that has been shown to

respond to environmental stimuli, such as iron. Upregulation of *arsR* may also play a role in

anaerobic growth of NG (33-35). Our study is the first to associate *arsR* with antimicrobial

susceptibility. Transcript *rpsO*, encoding 30S ribosomal protein S15, also forms a bridge to the

50S subunit to contact the 23S rRNA (36). Upregulation of ribosomal protein-encoding

transcripts such as *rpsO* upon antimicrobial exposure has been shown previously (32). Notably,

previously reported RNA markers for NG upon antimicrobial exposure included a different panel

of markers (except *rpsO* which is in common in both studies) suggesting that drug concentration,

antimicrobial exposure time, path of marker selection and bioinformatic pipelines could result in

different outcomes and should be considered in further application of this approach.

In order to use any novel AST platform in clinical decision making, it is necessary to compare it

to the gold standard AST method, MIC, which has been in used for decades (37). However,

current phenotypic assays still mistakenly classify some resistant strains as susceptible resulting

in failed clinical antimicrobial therapy (38). In this study, the measured changes in transcription

level, $2^{-\Delta\Delta CT}$, was correlated to MIC and translated to the susceptibility category resulted in a

threshold of a $2^{-\Delta\Delta CT} \geq 2$ for susceptibility (Fig. 5). This threshold is further supported by the

clustering of 64 susceptible and resistant CDC isolates with the 14 experimentally tested strains

using their known MICs and calculated $\Delta\Delta CT$ values (Table S5). However, when isolates with

MIC=1 μg/mL, categorized as susceptible based on CLSI breakpoint, were tested using *rpsO*

linear fitting, it was categorized as non-susceptible based on our $2^{-\Delta\Delta CT}$ threshold. Of note, this

classification is in agreement with EUCAST susceptibility breakpoint where an isolate with

MIC≥1μg/ml is called resistant. This result highlights the difficulties underlying azithromycin

susceptibility across NG isolates due to lack of a universal breakpoint interpretation in

conventional AST confirming the critical need for a new AST with a reliable susceptibility

breakpoint (24).

Overall, we have demonstrated the potential for antimicrobial susceptibility RNA marker

discovery through combined application of NGS, bioinformatics, and validation assays. A larger

scale experimental validation of these putative RNA markers by testing additional NG strains

with different resistance mechanisms to azithromycin is still needed to better assess test accuracy

and reproducibility. We envision translation of our ultra-rapid susceptibility markers into an

NAAT-based diagnostic with combined ID and AST capability for point-of-care use (Fig. 6). We

have previously developed a palm-sized magnetofluidic platform with combined bacterial ID and

AST directly from swab samples in less than 2.5 hours, of which AST required 2 hours of

antimicrobial incubation to assess for bacterial doubling (39). Integrating our growth-

independent, ultra-rapid AST markers on this platform would compress total assay time down to

40 minutes, drastically shifting the current paradigm for diagnosing and treating NG infections.

Acknowledgments

The authors would like to thank the funding support from the National Institute of Health (NIH

R01AI137272, R01AI138978)

307 Fig. 1

319   Fig. 2

330   Fig. 3

331   A

339   B

350   Fig.4

354 Fig. 5

355   A

365   B

Fig. 6

Figure Legends:

Fig. 1: Summarized workflow for generation of candidate RNA markers of the proposed

molecular AST. NG was cultured and exposed to azithromycin for 10 and 60 min along with a

control without azithromycin. Samples were collected for RNA-seq library and sequencing. Data

analysis was conducted to find differentially expressed genes followed by marker selection steps.

Selected markers were validated by qRT-PCR and ΔCT calculation.

This Fig. is created with BioRender.com and PPT.

Fig. 2: MDS plot to display differences in the azithromycin-induced gene expression profile at

10 and 60 min. Dissimilarity in the expression profiles between the replicates, strains, and

conditions were calculated and plotted. CS: control susceptible strain; AziS: azithromycin-

treated susceptible strain; CR: control resistant strain; AziR: azithromycin-treated resistant strain.

Fig. 3: (A) Volcano plot to show the gene expression profile induced by azithromycin varied

significantly at 10 and 60 min between the treated and untreated susceptible strains. Colored

points indicate genes that are up or down regulated in the SvC as well as SvR comparisons at

each time point. (B) Heatmap for differentially expressed genes (p-value<0.05 with a FC≥ 1) in

SvC and SvR induced by 10min azithromycin exposure.

Fig. 4. Validation of 6 RNA markers across 14 tested NG isolates using qRT-PCR. Black:

susceptible isolates; grey: resistant isolates. *ABC*, *bolA*, *arsR* and *rpsO* upregulated and *dinD* and

acoT downregulated RNA markers.

Fig. 5. (A) Determination of susceptibility threshold by correlation of MIC and our AST

platform using *arsR* and *rpsO* across14 tested NG isolates. (B) Linear fitting for *arsR* and *rpsO*

to show how transcriptional response is associated with the MIC across14 tested NG isolates (12

susceptible and 2 resistant).

Fig. 6. Proposed workflow for combined ID and AST using novel molecular point-of-care

systems such as magnetofluidic device (39). Clinical samples are collected and exposed to

antimicrobial for 10min. Control and antimicrobial treated samples are loaded into

magnetofluidic device for further ID and AST.

This Fig. is created with BioRender.com and PPT.

Supplementary data:

Table S1: Susceptibility profile of tested reference and clinical isolates of NG.

Table S2: Candidate markers information including gene description, locus tag and primer

sequences used for qPCR validation.

Table S3: List of differentially expressed genes in NG upon 10min and 60min exposure to

azithromycin.

Table S4: List of 568 differentially expressed genes (p-value<0.05) in S versus C and S versus R

induced by 10min azithromycin exposure.

Table S5: Validation of our susceptibility threshold ($2^{-\Delta\Delta CT} = 2$) across two peanels including 64

clinical isolates from CDC (Antibiotic Resistance Isolate Bank) with a wide range of MIC

values.

Fig. S1: Heatmap for differentially expressed genes (p-value<0.05) in S/C and S/R

induced by 10min azithromycin exposure

References

1. Shin DJ, Andini N, Hsieh K, Yang S, Wang T-H. 2019. Emerging analytical

techniques for rapid pathogen identification and susceptibility testing. Annu Rev Anal

Chem 12:41-67.

2. Voter AF, Callaghan MM, Tippana R, Myong S, Dillard JP, Keck JL. 2020.

Antigenic variation in *Neisseria gonorrhoeae* occurs independently of RecQ-

mediated unwinding of the *pilE* G quadruplex. J Bacteriol 202:e00607-19.

3. Shigemura K, Osawa K, Miura M, Tanaka K, Arakawa S, Shirakawa T, Fujisawa M.

2015. Azithromycin resistance and its mechanism in *Neisseria gonorrhoeae* strains in

Hyogo, Japan. Antimicrob Agents Chemother 59:2695-9.

4.   Burnett AM, Anderson CP, Zwank MD. Laboratory-confirmed gonorrhea and/or

chlamydia rates in clinically diagnosed pelvic inflammatory disease and cervicitis.

2012. Am J Emerg Med 30:1114-7.

5.   Centers for Disease Control and Prevention (CDC). Gonorrhea--United States, 1998.

2000. MMWR 49:538.

6.   Pond MJ, Hall CL, Miari VF, Cole M, Laing KG, Jagatia H, Harding-Esch E,

Monahan IM, Planche T, Hinds J, Ison CA, Chisholm S, Butcher PD, Sadiq ST. 2016.

Accurate detection of *Neisseria gonorrhoeae* ciprofloxacin susceptibility directly

from genital and extragenital clinical samples: towards genotype-guided antimicrobial

therapy. J Antimicrob Chemother 71:897-902.

7.   Chen L, Shin DJ, Zheng S, Melendez JH, Gaydos CA, Wang T-H. 2018. Direct-

qPCR assay for coupled identification and antimicrobial susceptibility testing of

Neisseria gonorrhoeae. ACS Infect Dis 4:1377-84.

8.   Khazaei T, Barlow JT, Schoepp NG, Ismagilov RF. 2018. RNA markers enable

phenotypic test of antibiotic susceptibility in *Neisseria gonorrhoeae* after 10 minutes

of ciprofloxacin exposure. Sci Rep 8:1-10.

9.   Zarantonelli L, Borthagaray G, Lee EH, Veal W, Shafer WM. 2001. Decreased

susceptibility to azithromycin and erythromycin mediated by a novel *mtrR* promoter

mutation in *Neisseria gonorrhoeae*. J Antimicrob Chemother 47:651-4.

10.   Ma KC, Mortimer TD, Hicks AL, Wheeler NE, Sánchez-Busó L, Golparian D,

Taiaroa G, Rubin DH, Wang Y, Williamson DA, Unemo M. 2020. Increased

antibiotic susceptibility in *Neisseria gonorrhoeae* through adaptation to the cervical

environment. bioRxiv 896696.

11.   Dinos GP, Michelinaki M, Kalpaxis DL. 2001. Insights into the mechanism of
459   azithromycin interaction with an *Escherichia coli* functional ribosomal complex. Mol
460   Pharmacol 59:1441-5.

461   12.   Zarantonelli L, Borthagaray G, Lee E-H, Shafer WM. 1999. Decreased azithromycin
462   susceptibility of *Neisseria gonorrhoeae* due to *mtrR* mutations. Antimicrob Agents
463   Chemother 43:2468-72.

464   13.   Eyre DW, Sanderson ND, Lord E, Regisford-Reimmer N, Chau K, Barker L, Morgan
465   M, Newnham R, Golparian D, Unemo. 2018. Gonorrhoea treatment failure caused by
466   a *Neisseria gonorrhoeae* strain with combined ceftriaxone and high-level
467   azithromycin resistance, England. Euro Surveill 23(27).

468   14.   Wan C, Li Y, Le W-J, Liu Y-R, Li S, Wang B-X, Rice PA, Su X-H. 2018. Increasing
469   resistance to azithromycin in *Neisseria gonorrhoeae* in eastern Chinese cities:
470   Resistance mechanisms and genetic diversity among isolates from Nanjing.
471   Antimicrob Agents Chemother 62:e02499-17.

472   15.   Workowski KA, Bolan GA, 2015. Centers for Disease Control and Prevention
473   (CDC). Sexually transmitted diseases treatment guidelines. MMWR 64:1-137.

474   16.   Centers for Disease Control and Prevention (CDC). 2011. *Neisseria gonorrhoeae*
475   with reduced susceptibility to azithromycin--San Diego County, California, 2009.
476   MMWR 60(18):579.

477   17.   Tobiason DM, Seifert HS. 2006. The obligate human pathogen, *Neisseria*
478   *gonorrhoeae*, is polyploid. PLoS Biol 4(6).

479   18.   Zhao YH, Qin XL, Yang JY, Liao YW, Wu XZ, Zheng HP. 2019. Identification and
480   expression analysis of ceftriaxone resistance-related genes in *Neisseria gonorrhoeae*

integrating RNA-Seq data and qRT-PCR validation. J Glob Antimicrob Resist

16:202-9.

19.   Yang X, Hashemi MM, Andini N, Li MM, Kuang S, Carroll KC, Wang T-H, Yang S.

2020. RNA markers for ultra-rapid molecular antimicrobial susceptibility testing in

fluoroquinolone-treated *Klebsiella pneumoniae*. J Antimicrob Chemother dkaa078.

20.   Diallo K, Coulibaly MD, Rebbetts LS, Harrison OB, Lucidarme J, Gamougam K,

Tekletsion YK, Bugri A, Toure A, Issaka B. 2018. Development of a PCR algorithm

to detect and characterize *Neisseria meningitidis* carriage isolates in the African

meningitis belt. PLoS One 13(12).

21.   Andini N, Hu A, Zhou L, Cogill S, Wang T-H, Wittwer CT, Yang S.. 2018. A

"culture" shift: Broad bacterial detection, identification, and antimicrobial

susceptibility testing directly from whole blood. Clin Chem 64:1453-62.

22.   Wade JJ, Graver MA. 2007. A fully defined, clear and protein-free liquid medium

permitting dense growth of *Neisseria gonorrhoeae* from very low inocula. FEMS

Microbiol Lett 273:35-7.

23.   Gose S, Kong CJ, Lee Y, Samuel MC, Bauer HM, Dixon P, Soge OO, Lei J, Pandori

M. 2013. Comparison of *Neisseria gonorrhoeae* MICs obtained by Etest and agar

dilution for ceftriaxone, cefpodoxime, cefixime and azithromycin. J Microbiol

Methods 95:379-80.

24.   McAuliffe GN, Smith M, Cooper G, Forster RF, Roberts SA. 2019. Variability in

azithromycin susceptibility results for *Neisseria gonorrhoeae* obtained using gradient

MIC strip and agar dilution techniques. J Clin Microbiol 57(12).

25.   Clinical and Laboratory Standards Institute. 2020. M100 Performance Standards for
505   Antimicrobial Susceptibility Testing, 30th ed. Clinical and Laboratory Standards
506   Institute, Wayne, PA.

507   26.   Andrew S. 2010. A quality control tool for high throughput sequence data. Fast QC
508   2010.

509   27.   Petersen KR, Streett DA, Gerritsen AT, Hunter SS, Settles ML. 2015. Super deduper,
510   fast PCR duplicate detection in fastq files. Proceedings of the 6th ACM Conference
511   on Bioinformatics, Computational Biology and Health Informatics 491-2.

512   28.   https://github.com/FelixKrueger/TrimGalore.

513   29.   Tjaden B.2015. De novo assembly of bacterial transcriptomes from RNA-seq data.
514   Genome Biol 16:1.

515   30.   Team RC. 2017. R: a language and environment for statistical computing. R
516   Foundation for Statistical Computing, Vienna, Austria. URL https://www. R-project.
517   org. 2017.

518   31.   Biosystems A. 2004. Guide to performing relative quantitation of gene expression
519   using real-time quantitative PCR. Applied Biosystems.

520   32.   Wadsworth CB, Sater MR, Bhattacharyya RP, Grad YH. 2019. Impact of species
521   diversity on the design of RNA-based diagnostics for antibiotic resistance in
522   *Neisseria gonorrhoeae*. Antimicrob Agents Chemother 63:e00549-19.

523   33.   Isabella VM, Clark VL. 2011. Deep sequencing-based analysis of the anaerobic
524   stimulon in *Neisseria gonorrhoeae*. BMC Genomics 12(1):51.

525   34.   Yu C, McClure R, Nudel K, Daou N, Genco CA. 2016. Characterization of the
526   *Neisseria gonorrhoeae* iron and fur regulatory network. J Bacteriol 198:2180-91.

35.   Isabella V, Wright LF, Barth K, Spence JM, Grogan S, Genco CA, Clark VL. 2008.

Cis- and *trans*-acting elements involved in regulation of *norB* (*norZ*), the gene

encoding nitric oxide reductase in *Neisseria gonorrhoeae*. Microbiology 154:226-39.

36.   Marzi S, Myasnikov AG, Serganov A, Ehresmann C, Romby P, Yusupov M, Klaholz

BP. 2007. Structured mRNAs regulate translation initiation by binding to the platform

of the ribosome. Cell 130:1019-31.

37.   Liu H, Taylor TH, Pettus K, Trees D. 2014. Assessment of Etest as an alternative to

agar dilution for antimicrobial susceptibility testing of *Neisseria gonorrhoeae*. J Clin

Microbiol 52:1435-40.

38.   Lutgring JD, Limbago BM. 2016. The problem of carbapenemase-producing-

carbapenem-resistant-*Enterobacteriaceae* detection. J Clin Microbiol 54:529-34.

39.   Lee PW, Chen L, Trick AY, Athamanolap P, Chen FE, Wang TH. 2019. Palm-sized

magnetofluidic platform for bacterial identification and antimicrobial susceptibility

testing of infected wounds. 23rd International Conference on Miniaturized Systems

for Chemistry and Life Sciences; Basel, Switzerland.