Article Summary Line: In this study, we present a rapid antimicrobial susceptibility testing

platform using RNA signatures for *Neisseria gonorrhoeae* upon minutes of azithromycin

exposure.

Running title: RNA markers for AST in *Neisseria gonorrhoeae*

Keywords: antimicrobial susceptibility testing, RNA markers, *Neisseria gonorrhoeae,*

azithromycin

A novel platform to accelerate antimicrobial susceptibility testing in *Neisseria gonorrhoeae*

using RNA signatures

Marjan M. Hashemi, Nikhil Ram-Mohan, Nicholas R. Gessner, Karen C. Carroll, Tza-Huei

Wang, Samuel Yang

Stanford University, Stanford, CA, USA (M.M. Hashemi, N, Ram-Mohan, N.R. Gessner, S.

Yang); and Johns Hopkins University, Baltimore, MD, USA (K.C. Carroll, T-H. Wang)

Address for correspondence: Samuel Yang, Stanford Emergency Medicine Dept, 900 Welch

Road, Suite 350, Palo Alto, CA 94304; Email: syang5@stanford.edu; Phone: (650) 725-9492;

Fax: (650) 723-0121

Abstract

The rise of antimicrobial resistant pathogens can be attributed to the lack of a rapid pathogen

identification (ID) or antimicrobial susceptibility testing (AST), resulting in delayed therapeutic

decision at the point of care. Gonorrhea is usually empirically treated with no AST results

available before treatment, thus contributing to rapid rise in drug resistance. Herein we present a

rapid AST platform using RNA signatures for *Neisseria gonorrhoeae* (NG). RNA-seq followed

by bioinformatic tools were applied to explore potential markers in transcriptome profile of NG

upon minutes of azithromycin exposure. Validation of candidate markers using PCR showed that

two markers (*arsR* (NGO1562) and *rpsO*) can deliver accurate AST results across 14 tested

isolates. Further validation of our cutoff in comparison to MIC across 64 more isolates

confirmed reliability of our platform. Our RNA markers combined with emerging molecular

point-of-care systems has the potential to greatly accelerate both ID and AST to inform

treatment.

Introduction

The use of conventional clinical methods for pathogen identification (ID) and antimicrobial

susceptibility testing (AST) is a time-consuming process that contributes to the rise of

antimicrobial resistance and may in turn increase mortality rates (*1*). Thus, expedition of

diagnostic methods for timely and directed therapy is an urgent clinical need that is to be

addressed.

Neisseria gonorrhoeae (NG), the causative agent of gonorrhea, is an increasingly common

sexually transmitted infection with more than 550, 000 cases reported annually in the United

States (*2*). Gonorrhea is usually seen in female cervicitis and male urethritis (*3*). Untreated

gonorrhea infections can lead to major complications such as heart and nervous system

infections, infertility, pelvic inflammatory disease, and newborn blindness (*4*, *5*). Gonorrhea is

often empirically treated based on a clinical syndromic diagnosis or after laboratory molecular

detection with nucleic acid amplification testing (NAAT). NAAT have largely replaced

diagnostic culture methods because they are faster, automated, with higher sensitivity to allow

for cost-effective diagnosis (*6*). Treatments often proceed without AST results. NG AST requires

laborious and time-consuming (at least 1-2 days) culture methods and is only undertaken in cases

of treatment failure (*7*, *8*).

NG can rapidly develop resistance to antimicrobial agents due to innate mechanisms for

acquiring resistance genes (*9*). Treatment is more challenging due to rapidly increasing

resistance to all of the most commonly used antimicrobials including sulfonamides, penicillin,

tetracyclines, and second-generation fluoroquinolones such as ciprofloxacin (*10*). Azithromycin

is a macrolide antimicrobial that binds to the 23S rRNA component of the 50S ribosome, thereby

inhibiting protein synthesis (*11*, *12*). Azithromycin has been shown an effective treatment against

gonococci with prolonged level in tissues and cells. Following oral administration, azithromycin

concentrates in tissues including genital sites (*12*). However high levels of azithromycin

resistance have been progressively reported worldwide (*13*, *14*). Some of the known mechanisms

of resistance to azithromycin include overexpression of the efflux pump (*mtrR*), 23S rRNA

mutation in azithromycin binding sites (C2611T and A2059G) and ribosomal target modification

by methylase (*ermC* and *ermB*) (*3*).

CDC reported antimicrobial resistant NG as one of the three most urgent threats to public health

and currently recommends dual therapy with ceftriaxone and azithromycin for the treatment of

uncomplicated gonorrhea (*15*). Despite recommended dual therapy, azithromycin monotherapy

could be still used to treat uncomplicated gonorrhea in cephalosporin-allergic patients (*16*). In

this study azithromycin has been selected for our phenotypic AST development for gonorrhea as

a proof of concept; however our approach has potential to be extended to other antimicrobials.

77
Given the time-consuming conventional AST, increasing level of resistance, and the challenge of

slow growth rate (doubling time 60min) of NG (*17*), development of a novel AST that can guide

initial treatment at the point of care is critically needed. Recently, RNA-seq has been effectively

applied in discovery of reliable genomic biomarkers to develop clinical diagnosis (*18*). Previous

work in our lab described a novel and accelerated AST workflow based on RNA-seq in

Klebsiella pneumoniae upon exposure to ciprofloxacin (*19*). This method is independent of cell

division unlike other molecular phenotypic AST that measure growth kinetics of bacteria based

on DNA copies (*20*, *21*). As proof-of-concept, in this study we focused on developing an ultra-

rapid molecular phenotypic AST for NG. RNA-seq followed by bioinformatic analysis was used

to identify candidate diagnostic RNA markers to determine susceptibility upon a short exposure

to azithromycin. Further validation of selected markers was performed through qRT-PCR using

14 isolates followed by validation of our cutoff in comparison to reported MICs using 64 more

isolates to determine susceptibility.

91
92

93   Material and methods
94
Microorganisms and culturing

Reference strains of azithromycin susceptible and resistant NG, SPL-4 and SPJ-15, were

obtained from CDC. Clinical isolates of NG were obtained from John Hopkins Medical

Institutions, Department of Pathology, Division of Medical Microbiology (Appendix Table 1).

Isolates were cultured from glycerol stocks on GC agar (#BD 228950, Becton Dickinson,

Cockeysville, MD) supplemented with 1% IsoVitaleX (#BD 211876, Becton Dickinson,

Cockeysville, MD). Plates were incubated at 35 °C, in 5% CO_2 for 24 h. A single colony of each

isolate was re-suspended in GW broth (22) and incubated overnight at 37 °C, 5% CO_2, 200 rpm

to an optical density equal to that of a 0.5 MacFarland standard.

Antimicrobial susceptibility testing

The E-test, bioMérieux (Durham, NC, USA), or agar dilution method, was used to determine the

minimum inhibitory concentration (MIC) of azithromycin (Astatech, Inc, Bristol, PA) (23). Agar

dilution method was performed using GC agar supplemented with 1% IsoVitaleX described by

the Clinical and Laboratory Standards Institute (CLSI). Briefly, two-fold dilutions of

azithromycin (range 0.03-256 µg/mL) was added to CG medium (24). Each plate was inoculated

with a 0.5 McFarland standard of bacterial suspension. Inoculated plates were incubated at 35

°C, in 5% CO_2 for 24 h. Measurements were performed in triplicate and positive and negative

controls were included for each MIC test. Following incubation, MICs were determined as the

lowest concentrations of azithromycin that inhibited visible growth of bacteria. According to the

guidelines set forth by the CLSI, isolates with MIC ≤2 µg/mL are classified as azithromycin

susceptible (25).

RNA extraction and sequencing

Bacterial culture (2mL) was prepared as described previously. Samples were exposed to

azithromycin (2μg/mL) and incubated at 37 °C, in 5% CO_2 for 10 minutes and 60 minutes (see

workflow in Figure 1). Bacteria without exposure to azithromycin were also incubated in the

same condition as a control (water was added to the controls). Samples were collected and

preserved in 4mL RNA Protect Reagent (Qiagen, Valencia, CA, USA) at each time point. RNA

extraction was conducted using RNeasy Mini Kit (Qiagen,74524) according to the

manufacturer's instruction. Extracted RNA was subjected to DNase treatment (Turbo DNase

complete kit Life Technologies, AM1907). Concentration of RNA was measured using QIA

expert spectrophotometer (Qiagen, USA). To consider biological variability triplicate RNA

samples were prepared from susceptible and resistant isolates.

Sample preparation for sequencing was performed using ScriptSeq complete kit for bacteria with

rRNA depleted (Illumina Inc, BB1224, RSBC10948 and SSIP1202). Samples were bio-analyzed

at Stanford Functional Genomics Facility (SFGF) and libraries were sequenced at 75 pair-end

reads on Illumina MiSeq. Quality control on raw reads from the sequenced libraries was

conducted to remove the low-quality reads (using FastQC) (26), remove PCR duplicates (using

SuperDeduper) (27) and trim the adaptor sequences (using Trim Galore, version 0.4.5) (28).

Finally, reads were aligned to NG FA 1090 (NCBI Reference Sequence: NC_002946.2RNA-seq

data are archived in the GenBank SRA (BioProject ID: PRJNA627416).

Differential expression analysis

Quantification of transcript abundance and differential expression analyses were carried out

using Rockhopper (*29*). Spearman correlation between samples/replicates from the normalized

transcript counts obtained from Rockhopper were calculated using the cor function in R (*30*).

The multidimensional scaling (MDS) that show the dissimilarity of the replicates/samples by

projecting them into two dimensions was also done using the cmdscale function.

To further analyze the differentially expressed genes under azithromycin treatment, the list was

cut down to 568 differentially expressed genes with a p-value<0.05 in susceptible compared to

control (SvC) and resistant (SvR) strains. Heatmaps were generated using the heatmap.2 function

to represent the logFC between these comparisons.

149
150
151   Marker selection and validation

152   Candidate markers were selected from the RNA-seq dataset based on the following criteria: 1)

153   significant fold change in susceptible but not in resistant after 10min drug exposure; 2) high level

154   of expression after 10min drug exposure; 3) same trend of expression in 10&60 min drug

155   exposure. Selected candidate markers were further evaluated using qPCR.

156
Validation of final markers in clinical isolates was performed using quantitative real-time PCR

(qRT-PCR). Primer-blast was used to design primers for conserved regions of selected genes

(Appendix Table 2). Quantification of gene expression level was conducted with Rotor-Gene

SYBR Green RT-PCR kit (Qiagen, cat# 204174) on the Rotor-Gene-Q PCR cycler (Qiagen).

The concentration of component used in final 25µL PCR reaction mix were as follows: 1x SYBR

Green Master mix, 1mM forward primer, 1mM reverse primer, 0.05U/µL Rotor Gene RT mix

and 0.2U/µL SUPERase-in RNase inhibitor and 8% RNA template.

Expression of each candidate marker was quantified in the control and treated samples.

Threshold of 0.1 was set to obtain accurate CT values for each isolate. This threshold is a point at

which amplification plot reaches a fluorescent intensity above background level (*31*). To

calculate ΔCT value, a housekeeping gene was used to normalize potential differences between

the control and treated samples. From our RNA-seq results, *atpA* gene was selected as an internal

control as its expression level did not affect significantly upon antimicrobial treatment for the

tested isolates in this study. Finally, fold change (FC) in gene expression between the control and

treated samples was calculated as follows: $FC=2^{(-\Delta\Delta CT)}$ (*18*).

172
Results

Shift in transcriptome response following azithromycin exposure

To identify markers that significantly differentiate susceptible and resistant isolates of *N.*

gonorrhoeae, RNA-seq was used to compare their transcriptome response after 10 and 60 min of

azithromycin exposure. A summary of workflow is shown in Figure 1.

Multidimensional scaling analysis of the expression profile of the biological replicates, treatment

conditions, and the two strains revealed strong reproducibility amongst replicates and

dissimilarity between the conditions and strains (Figure 2). MDS plot was also used to provide

insights into the association between transcriptional profile and exposure time (10 and 60 min) to

azithromycin. Biological replicates of control and treated samples clustered very closely

indicating high correlation among replicates and high reproducibility of library preparation.

MDS plot demonstrated four distinct transcriptional clusters within the first two dimensions

(dim1 and dim2), an indication of remarkable transcriptional changes upon antimicrobial

exposure. Well separation of the control and treated samples in the first dimension showed that

the sequencing data were qualified for identification of differentially expressed genes.

Additionally, a significant diversity between the gene expression of 10min and 60 min treated

samples was observed suggesting that distinct gene expression profiles are triggered by

azithromycin in a time-dependent manner.

Subsequently, statistical analysis of differentially expressed genes following exposure to

azithromycin was calculated using logFC (gene expression ratio of treated compared to the

control) against their p-values <0.05 (Figure 3A). Consistent with MDS results, a global shift in

SvC and SvR was observed as early as 10 min of azithromycin exposure, although distribution of

gene expression profile illustrated a higher magnitude of fold change at 60 min.

Marker selection and validation of candidate markers in *NG* clinical isolates

Differential expression analyses of the susceptible strain revealed more down than up regulated

genes at 10 minutes (185:129). At 60 minutes however, there are 386 and 212 genes up and

down regulated, respectively (Appendix Table 3). Since we are interested in markers that will

successfully determine susceptibility, we further screened for genes that were differentially

expressed between SvC and SvR. A p-value cutoff of 0.05 resulted in 568 genes (Appendix

Figure 1) which narrows down to 48 when screening for abs(logFC) ≥1 under both conditions

(Figure 3B).

Among the differentially expressed genes with significant logFC, two more criteria including

high level of expression and same trend of expression in 10&60 min (see methods) were applied

to select the candidate markers. The initially selected markers include NGO0373, NGO1920,

NGO1562, NGO0191, NGO0405 and NGO1078, hereafter referred to as *ABC*, *bolA*, *arsR*, *rpsO*,

dinD and *acoT*, respectively (Appendix Table 2). These were tested in one susceptible and one

resistant strain to confirm the diagnostic potential of markers to characterize azithromycin

susceptibility using RT-qPCR. All six candidate RNA markers are highlighted in Appendix

Table 4, among which *ABC*, *bolA*, *arsR*, and *rpsO* are upregulated while *dinD* and *acoT* are

downregulated. Further validation of markers was conducted in eleven susceptible and one

resistant clinical isolates of NG to confirm that RNA-seq nominated markers can be applied in

different strains (Figure 4). We calculated -$\Delta\Delta$CT values for samples following 10 min and 60

min exposure to azithromycin and compared it between resistant and susceptible isolates for each

marker. Overall -$\Delta\Delta$CT of all six selected candidate markers was significantly different across

resistant and susceptible isolates indicating qualification of these candidate markers to be applied

in our new AST platform. Among six tested markers *arsR* and *rpsO* demonstrated consistent

upregulation trends across all tested isolates. Although *ABC* and *bolA* were also validated as

significantly upregulated RNA markers in most tested isolates, down regulation of these two

markers was observed in a few tested susceptible isolates. On the other hand, *dinD* and *acoT*

were validated as significantly downregulated RNA markers, however inconsistency of results

from a few isolates was also found. According to the validation results of six selected markers,

further analysis was conducted using *arsR* and *rpsO* as our final markers.

We examined the reliability of our method compared to the gold standard using relationship

between $2^{-\Delta\Delta CT}$ values of final markers and MIC in susceptible and resistant strains after 10min

exposure to azithromycin (Figure 5A). For both *arsR* and *rpsO*, $2^{-\Delta\Delta CT}$ values significantly

changed between susceptible and resistant and notably no overlap in $2^{-\Delta\Delta CT}$ values was observed

between two groups of susceptible and resistant strains. Additionally, a $2^{-\Delta\Delta CT} = 2$ was defined as

threshold susceptibility for susceptible and non-susceptible isolates ($2^{-\Delta\Delta CT} \geq 2$ susceptible, 2^{-}

$^{\Delta\Delta CT}$< 2 non-susceptible). We also used a linear fitting to show how transcriptional response of

the final markers are associated to the MIC in susceptible and resistant strains. A strong negative

relationship was observed between $\log_2 2^{-\Delta\Delta CT}$ and $\log_2$ MIC values, (R^2 values 0.64 and 0.63 for

arsR and *rpsO* respectively), meaning $2^{-\Delta\Delta CT}$ value in susceptible strains is significantly higher

than that in resistant strains (Figure 5B).

Finally, to validate our susceptibility threshold, we used MIC information of two panels of NG

including 64 clinical isolates from CDC (Antibiotic Resistance Isolate Bank) on the equation for

the line of best fit for our markers. Notably, *arsR* supported the delineated $2^{-\Delta\Delta CT}$ value by

correctly classifying susceptibility of all 64 CDC isolates based on our $2^{-\Delta\Delta CT}$ threshold as 51

susceptible strains and 13 resistant strains that was in agreement with CLSI susceptibility

category (MIC breakpoint of ≥2μg/ml) (Appendix Table 5). *rpsO* also accurately identified

susceptibility of most tested isolates with a broad range of MIC, although susceptible isolates

with MIC at 1μg/ml were classified as non-susceptible isolates; however, this category is in

agreement with EUCAST breakpoint values (MIC breakpoint of ≥1μg/ml).

In summary, these observations confirmed the accurate application of $2^{-\Delta\Delta CT}$ values to categorize

susceptibility of isolates as a replacement of MIC in traditional AST.

Discussion

To stem the tide of multi-drug resistant NG, a rapid NAAT-based diagnostic with compatible

molecular phenotypic AST capability is critically needed to inform initial treatment decisions. In

this study, we have discovered azithromycin-susceptibility RNA markers and demonstrated their

early promise for ultra-rapid AST in NG.

Transcriptome profiling of susceptible and resistant NG upon exposure to azithromycin revealed

a significantly altered response as early as 10 min in SvC and SvR (Figures 2 & 3). While

significantly different transcriptional response among NG was previously reported after 60 min

azithromycin exposure (*32*), our main objective was to discover the earliest susceptibility RNA

markers to minimize AST time. Therefore, two time points of 10min and 60min were included in

azithromycin treatment to be able to identify consistent and reliable RNA markers as part of

early cellular stress responses to antimicrobials. Unlike the previous reported studies at which

sublethal concentration of drugs was used (*8, 32*), we studied bacterial transcriptomic profile

following a high concentration of azithromycin (2µg/mL) to enhance the discovery of

differentially expressed genes (*19*). Six candidate markers were initially identified through RNA-

seq and subsequently tested in 14 isolates with qRT-PCR. *arsR* and *rpsO* were chosen as our two

final markers as they were able to consistently determine susceptibility based on CT value only

after 10min antimicrobial exposure (Figure 4).

arsR gene (NGO1562) encodes a transcriptional regulatory protein that has been shown to

respond to environmental stimuli, such as iron. Upregulation of *arsR* may also play a role in

anaerobic growth of NG (*33-35*). Our study is the first to associate *arsR* with antimicrobial

susceptibility. Transcript *rpsO*, encoding 30S ribosomal protein S15, also forms a bridge to the

50S subunit to contact the 23S rRNA (*36*). Upregulation of ribosomal protein-encoding

transcripts such as *rpsO* upon antimicrobial exposure has been shown previously (*32*). Notably,

previously reported RNA markers for NG upon antimicrobial exposure included a different panel

of markers (except *rpsO* which is in common in both studies) suggesting that drug concentration,

antimicrobial exposure time, path of marker selection and bioinformatic pipelines could result in

different outcomes and should be considered in further application of this approach.

283
284
In order to use any novel AST platform in clinical decision making, it is necessary to compare it

to the gold standard AST method, MIC, which has been in used for decades (*37*). However,

current phenotypic assays still mistakenly classify some resistant strains as susceptible resulting

in failed clinical antimicrobial therapy (*38*). In this study, the measured changes in transcription

level, $2^{-\Delta\Delta CT}$, was correlated to MIC and translated to the susceptibility category resulted in a

threshold of a $2^{-\Delta\Delta CT} \geq 2$ for susceptibility (Figure 5). This threshold is further supported by the

clustering of 64 susceptible and resistant CDC isolates with the 14 experimentally tested strains

using their known MICs and calculated $\Delta\Delta CT$ values (Appendix Table 5). However, when

isolates with MIC=1 μg/mL, categorized as susceptible based on CLSI breakpoint, were tested

using *rpsO* linear fitting, it was categorized as non-susceptible based on our $2^{-\Delta\Delta CT}$ threshold. Of

note, this classification is in agreement with EUCAST susceptibility breakpoint where an isolate

with MIC≥1μg/ml is called resistant. This result highlights the difficulties underlying

azithromycin susceptibility across NG isolates due to lack of a universal breakpoint

interpretation in conventional AST confirming the critical need for a new AST with a reliable

susceptibility breakpoint (*24*).

Overall, we have demonstrated the potential for antimicrobial susceptibility RNA marker

discovery through combined application of NGS, bioinformatics, and validation assays. A larger

scale experimental validation of these putative RNA markers by testing additional NG strains

with different resistance mechanisms to azithromycin is still needed to better assess test accuracy

and reproducibility. We envision translation of our ultra-rapid susceptibility markers into an

NAAT-based diagnostic with combined ID and AST capability for point-of-care use (Figure 6).

We have previously developed a palm-sized magnetofluidic platform with combined bacterial ID

and AST directly from swab samples in less than 2.5 hours, of which AST required 2 hours of

antimicrobial incubation to assess for bacterial doubling (*39*). Integrating our growth-

independent, ultra-rapid AST markers on this platform would compress total assay time down to

40 minutes, drastically shifting the current paradigm for diagnosing and treating NG infections.

Funding

The authors would like to thank the funding support from the National Institute of Health (NIH

R01AI137272, R01AI138978)

Author Bio

Dr. Marjan M.Hashemi is a postdoctoral fellow in the Department of Emergency Medicine at

Stanford University, Palo Alto. Her primary research interests include developing molecular

diagnostic assays for pathogen identification and antimicrobial susceptibility testing.

321
322

325
326
327
328

329
Figure 1: Summarized workflow for generation of candidate RNA markers for the proposed
molecular AST. NG was cultured and exposed to azithromycin for 10 min and 60 min along with
a control without azithromycin. Samples were collected for RNA-seq library and sequencing.
Data analysis was conducted to find differentially expressed genes followed by marker selection
steps. Selected markers were validated by qRT-PCR and ΔCT calculation.
This figure is created with BioRender.com and PPT.

355
356
357 Figure 2: MDS plot to display differences in the azithromycin-induced gene expression profile at
358 10 and 60 min. Dissimilarity in the expression profiles between the replicates, strains, and
359 conditions were calculated and plotted.

377 **A**

378
379 **B**

380
381 Figure 3: (A) Volcano plot to show the gene expression profile induced by azithromycin varied
382 significantly at 10 and 60 min between the treated and untreated susceptible strains. Colored
383 points indicate genes that are up or down regulated in the SvC as well as SvR comparisons at

384 each time point. (B) Heatmap for differentially expressed genes (p-value<0.05 with a FC≥ 1) in
385 SvC and SvR induced by 10min azithromycin exposure.
386

387
388
389 Figure 4. Validation of 6 RNA markers across 14 tested NG isolates using qRT-PCR. Black:
390 susceptible isolates; grey: resistant isolates. *ABC*, *bolA*, *arsR* and *rpsO* upregulated and *dinD* and
391 *acoT* downregulated RNA markers.

397
398
399   **A**   **B**
400

401
402
403   Figure 5. (A) Determination of susceptibility threshold by correlation of MIC and our AST
404   platform using *arsR* and *rpsO* across14 tested NG isolates.(B) Linear fitting for *arsR* and *rpsO* to
405   show how transcriptional response is associated with the MIC across14 tested NG isolates (12
406   susceptible and 2 resistant).

419
420
421

422
423 Figure 6. Proposed workflow for combined ID and AST using a novel molecular point-of-care
424 systems such as a magnetofluidic device (*39*). Clinical samples are collected and exposed to
425 antimicrobial for 10min. Control and antimicrobial treated samples are loaded into
426 magnetofluidic device for further ID and AST.
427 This figure is created with BioRender.com and PPT.
428
429

Supplementary data:
Appendix Table1: Susceptibility profile of tested reference and clinical isolates of NG.
Appendix Table2: Candidate markers information including gene description, locus tag and
primer sequences used for qPCR validation.
Appendix Table3: List of differentially expressed genes in NG upon 10min and 60min exposure
to azithromycin.
Appendix Table 4: List of 568 differentially expressed genes (p-value<0.05) in S/C and S/R
induced by 10min azithromycin exposure.
Appendix Table5: Validation of our susceptibility threshold ($2^{-\Delta\Delta CT} = 2$) across 64 clinical
isolates from CDC (Antibiotic Resistance Isolate Bank) with a wide range of MIC values.
Appendix Figure1: Heatmap for differentially expressed genes (p-value<0.05) in S/C and S/R
induced by 10min azithromycin exposure

453
454 References
455
1. Shin DJ, Andini N, Hsieh K, Yang S, Wang T-H. Emerging analytical techniques for
rapid pathogen identification and susceptibility testing. Annu Rev Anal Chem.
2019;12(1):41-67.
2. Voter AF, Callaghan MM, Tippana R, Myong S, Dillard JP, Keck JL. Antigenic
variation in *Neisseria gonorrhoeae* occurs independently of RecQ-mediated
unwinding of the *pilE* G quadruplex. J Bacteriol. 2020;202(3):e00607-19.
3. Shigemura K, Osawa K, Miura M, Tanaka K, Arakawa S, Shirakawa T, et al.
Azithromycin resistance and its mechanism in *Neisseria gonorrhoeae* strains in
Hyogo, Japan. Antimicrob Agents Chemother. 2015;59(5):2695-9.
4. Burnett AM, Anderson CP, Zwank MD. Laboratory-confirmed gonorrhea and/or
chlamydia rates in clinically diagnosed pelvic inflammatory disease and cervicitis.
Am J Emerg Med. 2012;30(7):1114-7.
5. Centers for Disease Control and Prevention (CDC). Gonorrhea--United States, 1998.
MMWR. 2000;49(24):538.
6. Pond MJ, Hall CL, Miari VF, Cole M, Laing KG, Jagatia H, et al. Accurate detection
of *Neisseria gonorrhoeae* ciprofloxacin susceptibility directly from genital and
extragenital clinical samples: towards genotype-guided antimicrobial therapy. J
Antimicrob Chemother. 2016;71(4):897-902.
7. Chen L, Shin DJ, Zheng S, Melendez JH, Gaydos CA, Wang T-H. Direct-qPCR assay
for coupled identification and antimicrobial susceptibility testing of *Neisseria
gonorrhoeae*. ACS Infect Dis. 2018;4(9):1377-84.
8. Khazaei T, Barlow JT, Schoepp NG, Ismagilov RF. RNA markers enable phenotypic
test of antibiotic susceptibility in *Neisseria gonorrhoeae* after 10 minutes of
ciprofloxacin exposure. Sci Rep. 2018;8(1):1-10.
9. Zarantonelli L, Borthagaray G, Lee EH, Veal W, Shafer WM. Decreased
susceptibility to azithromycin and erythromycin mediated by a novel *mtrR* promoter
mutation in *Neisseria gonorrhoeae*. J Antimicrob Chemother. 2001;47(5):651-4.
10. Ma KC, Mortimer TD, Hicks AL, Wheeler NE, Sánchez-Busó L, Golparian D, et al.
Increased antibiotic susceptibility in *Neisseria gonorrhoeae* through adaptation to the
cervical environment. bioRxiv. 2020;896696.
11. Dinos GP, Michelinaki M, Kalpaxis DL. Insights into the mechanism of azithromycin
interaction with an *Escherichia coli* functional ribosomal complex. Mol Pharmacol.
2001;59(6):1441-5.
12. Zarantonelli L, Borthagaray G, Lee E-H, Shafer WM. Decreased azithromycin
susceptibility of *Neisseria gonorrhoeae* due to *mtrR* mutations. Antimicrob Agents
Chemother. 1999;43(10):2468-72.
13. Eyre DW, Sanderson ND, Lord E, Regisford-Reimmer N, Chau K, Barker L, et al.
Gonorrhoea treatment failure caused by a *Neisseria gonorrhoeae* strain with
combined ceftriaxone and high-level azithromycin resistance, England, February
2018. Euro Surveill. 2018;23(27).
14. Wan C, Li Y, Le W-J, Liu Y-R, Li S, Wang B-X, et al. Increasing resistance to
azithromycin in *Neisseria gonorrhoeae* in eastern Chinese cities: Resistance

mechanisms and genetic diversity among isolates from Nanjing. Antimicrob Agents
Chemother. 2018;62(5):e02499-17.

15. Workowski KA, Bolan GA, Centers for Disease Control and Prevention (CDC).
     Sexually transmitted diseases treatment guidelines, 2015. MMWR. 2015;64:1-137.

16. Centers for Disease Control and Prevention (CDC). *Neisseria gonorrhoeae* with
     reduced susceptibility to azithromycin--San Diego County, California, 2009.
     MMWR. 2011;60(18):579.

17. Tobiason DM, Seifert HS. The obligate human pathogen, *Neisseria gonorrhoeae*, is
     polyploid. PLoS Biol. 2006;4(6).

18. Zhao YH, Qin XL, Yang JY, Liao YW, Wu XZ, Zheng HP. Identification and
     expression analysis of ceftriaxone resistance-related genes in *Neisseria gonorrhoeae*
     integrating RNA-Seq data and qRT-PCR validation. J Glob Antimicrob Resist.
     2019;16(1):202-9.

19. Yang X, Hashemi MM, Andini N, Li MM, Kuang S, Carroll KC, et al. RNA markers
     for ultra-rapid molecular antimicrobial susceptibility testing in fluoroquinolone-
     treated *Klebsiella pneumoniae*. J Antimicrob Chemother. 2020.

20. Diallo K, Coulibaly MD, Rebbetts LS, Harrison OB, Lucidarme J, Gamougam K, et
     al. Development of a PCR algorithm to detect and characterize *Neisseria meningitidis*
     carriage isolates in the African meningitis belt. PLoS One. 2018;13(12).

21. Andini N, Hu A, Zhou L, Cogill S, Wang T-H, Wittwer CT, et al. A "culture" shift:
     Broad bacterial detection, identification, and antimicrobial susceptibility testing
     directly from whole blood. Clin Chem. 2018;64(10):1453-62.

22. Wade JJ, Graver MA. A fully defined, clear and protein-free liquid medium
     permitting dense growth of *Neisseria gonorrhoeae* from very low inocula. FEMS
     Microbiol Lett. 2007;273(1):35-7.

23. Gose S, Kong CJ, Lee Y, Samuel MC, Bauer HM, Dixon P, et al. Comparison of
     Neisseria gonorrhoeae MICs obtained by Etest and agar dilution for ceftriaxone,
     cefpodoxime, cefixime and azithromycin. J Microbiol Methods. 2013;95(3):379-80.

24. McAuliffe GN, Smith M, Cooper G, Forster RF, Roberts SA. Variability in
     azithromycin susceptibility results for *Neisseria gonorrhoeae* obtained using gradient
     MIC strip and agar dilution techniques. J Clin Microbiol. 2019;57(12).

25. Clinical and Laboratory Standards Institute. Performance Standards for Antimicrobial
     Susceptibility Testing, 30 ed. CLSI supplement M100, Table 2F. Clinical and
     Laboratory Standards Institute Suite 2500, Wayne Pennsylvania, 19087, USA, 2020.

26. Andrew S. A quality control tool for high throughput sequence data. Fast QC. 2010.

27. Petersen KR, Streett DA, Gerritsen AT, Hunter SS, Settles ML. Super deduper, fast
     PCR duplicate detection in fastq files. Proceedings of the 6th ACM Conference on
     Bioinformatics, Computational Biology and Health Informatics; 2015. p. 491-2.

28. https://github.com/FelixKrueger/TrimGalore.

29. Tjaden B. De novo assembly of bacterial transcriptomes from RNA-seq data.
     Genome Biol. 2015;16(1):1.

30. Team RC. R: a language and environment for statistical computing. R Foundation for
     Statistical Computing, Vienna, Austria. URL https://www. R-project. org. 2017.

31. Biosystems A. Guide to performing relative quantitation of gene expression using
     real-time quantitative PCR. Applied Biosystems. 2004.

543   32.   Wadsworth CB, Sater MR, Bhattacharyya RP, Grad YH. Impact of species diversity
544   on the design of RNA-based diagnostics for antibiotic resistance in *Neisseria*
545   *gonorrhoeae*. Antimicrob Agents Chemother. 2019;63(8):e00549-19.
546   33.   Isabella VM, Clark VL. Deep sequencing-based analysis of the anaerobic stimulon in
547   *Neisseria gonorrhoeae*. BMC Genomics. 2011;12(1):51.
548   34.   Yu C, McClure R, Nudel K, Daou N, Genco CA. Characterization of the *Neisseria*
549   *gonorrhoeae* iron and fur regulatory network. J Bacteriol. 2016;198(16):2180-91.
550   35.   Isabella V, Wright LF, Barth K, Spence JM, Grogan S, Genco CA, et al. *Cis-* and
551   *trans*-acting elements involved in regulation of *norB* (*norZ*), the gene encoding nitric
552   oxide reductase in *Neisseria gonorrhoeae*. Microbiology. 2008;154(1):226-39.
553   36.   Marzi S, Myasnikov AG, Serganov A, Ehresmann C, Romby P, Yusupov M, et al.
554   Structured mRNAs regulate translation initiation by binding to the platform of the
555   ribosome. Cell. 2007;130(6):1019-31.
556   37.   Liu H, Taylor TH, Pettus K, Trees D. Assessment of Etest as an alternative to agar
557   dilution for antimicrobial susceptibility testing of *Neisseria gonorrhoeae*. J Clin
558   Microbiol. 2014;52(5):1435-40.
559   38.   Lutgring JD, Limbago BM. The problem of carbapenemase-producing-carbapenem-
560   resistant-*Enterobacteriaceae* detection. J Clin Microbiol. 2016;54(3):529-34.
561   39.   Lee PW, Chen L, Trick AY, Athamanolap P, Chen FE, Wang TH. Palm-sized
562   magnetofluidic platform for bacterial identification and antimicrobial susceptibility
563   testing of infected wounds. 23rd International Conference on Miniaturized Systems
564   for Chemistry and Life Sciences; Basel, Switzerland, 27 - 31 October 2019.
565
566
567