Optimizing experimental design for genome sequencing and

assembly with Oxford Nanopore Technologies

John M. Sutton[1], Joshua D. Millwood[1], A. Case McCormack[1], and Janna L. Fierst[1]*

[1] Department of Biological Sciences, The University of Alabama, Tuscaloosa, AL 35487-0344

* Author for correspondence, janna.l.fierst@ua.edu

Abstract (250 words)

Background

High quality reference genome sequences are the core of modern genomics. Oxford Nanopore

Technologies (ONT) produces inexpensive DNA sequences in excess of 100,000 nucleotides but

high error rates make sequence assembly and analysis a non-trivial problem as genome size and

complexity increases. To date there has been no comprehensive attempt to generate robust

experimental design for ONT genome sequencing and assembly. In this study, we simulate ONT

and Illumina DNA sequence reads for the model organisms *Escherichia coli*, *Caenorhabditis*

elegans, *Arabidopsis thaliana,* and *Drosophila melanogaster* and assemble with Canu, Flye, and

MaSuRCA software to quantify the influence of sequencing coverage and assembly approach.

Heterozygosity in outbred eukaryotes is a common problem for genome assembly. We show

broad applicability of our methods using real ONT data generated for four strains of the highly

heterozygous nematode *Caenorhabditis remanei* and *C. latens*.

Results

- ONT libraries have a unique error structure and high sequence depth is necessary to

assemble contiguous genome sequences.

- As sequence depth increases errors accumulate and assembly statistics plateau.

• High-quality assembled sequences require a combination of experimental techniques that

increase sequence read length and computational protocols that reduce error through

correction, read selection and 'polishing' with higher accuracy short sequence reads.

• Our robust experimental design results in highly contiguous and accurate genome

assemblies for the four strains of *C. remanei* and *C. latens*.

Conclusions

ONT sequencing is inexpensive and accessible but the technology's error structure requires

robust experimental design. Our quantitative results will be helpful for a broad array of

researchers seeking guidance for *de novo* assembly projects.

Background

Many factors affect the quality of a *de novo* genome assembly. Genome size increases the size of

the "puzzle" to put together, while the size of the pieces (sequence reads) remains the same.

Repetitive regions and mobile genetic elements create unique challenges as they are present in

more than one location in the genome and without contextual information it is difficult to

identify how many copies exist in the complete genome. For example, *Alu* repeat elements reach

>1 million copies in the human genome [1].

Third generation sequencing technologies can theoretically solve these problems. The

long sequence reads span repetitive regions, potentially allowing for the identification of the

exact size and location of repeats on a chromosome. Long sequence reads increase the puzzle

piece size for assembly and require less sequencing effort to span the entire genome. Pacific

Biosciences (PacBio) and Oxford Nanopore Technologies (ONT) are the current front-runners in

long-read sequencing platforms; both are capable of average read lengths in the tens of thousands

of base pairs and, theoretically, entire chromosomes can be sequenced in a single read [2, 3].

Both are also capable of producing quality genome assemblies with reasonable amounts of data

[4].

ONT offers several advantages over PacBio. Nanopore sequencing relies on running

molecular fragments through engineered nanopores and recording the resulting alterations in

electrical current. The technology is versatile and can be used for DNA sequencing, mRNA

sequencing, amplification-free mRNA quantification [5], and measuring DNA base

modifications like methylation [6, 7]. ONT libraries can be readily prepared with low amounts of

input DNA, which is an important consideration when studying organisms that are small or

difficult to sample. ONT platforms are inexpensive and designed to be used in individual

research laboratories and classrooms [8]. For these reasons we have chosen to study ONT and

quantify how this inexpensive, accessible technology may be best utilized to produce high-

quality assembled reference genome sequences. Similar projects analyzed experimental design

for PacBio [9, 10] but the few benchmarking studies for ONT sequencing and assembly are

carried out on a narrow scope of organisms [11-13]

ONT sequence reads present unique challenges for genome sequence assembly. DNA

molecules do not move through protein nanopores at a constant rate and changes in current are a

composite signature reflecting 3-5 nucleotides occupying the nanopore (for R9.4.1 flow cells).

The signal processing has trouble detecting changes in current with homopolymers (single

nucleotide repeats, for example AAAAAA), short tandem repeats and heavily methylated sites

[14, 15]. As a result, ONT sequence reads contain small nucleotide stretches that have been

incorrectly identified, inserted and deleted (Fig. 1A, C). This error structure results in relatively

few large, contiguous stretches of correctly identified nucleotides (Fig. 1B, D) and is uniquely

challenging for assembly algorithms.

To achieve our results we simulate ONT DNA libraries for the model organisms

Escherichia coli, *Caenorhabditis elegans*, *Arabidopsis thaliana,* and *Drosophila melanogaster*.

For each organism we assemble sequence read sets at different depths (i.e., the average times a

nucleotide in the genome is sequenced). We measure the contiguity, completeness, and accuracy

of each assembled sequence relative to the current reference genome. Many *de novo* assembly

projects target organisms without reference genomes and we also measure the identification of a

set of genes thought to be found in single copy in each organism. Pure ONT datasets result in

superior assembled genome sequences but can be cost-prohibitive for large genomes where

multiple flow cells may be necessary to gather enough data. We also analyze assembled

sequences from 'hybrid' DNA read datasets composed of ONT and Illumina sequence reads.

For eukaryotes both ploidy and breeding system can influence the assembly of genome

sequence. Polyploidy can create scenarios where many sites in a genome look highly similar to

one another, making it tough to place these regions within an assembled genome [16, 17]. For

non-haploid organisms, there is a potential for heterozygosity and at these sites the effective

sequencing coverage is cut in half. For small, non-clonal metazoans sufficient DNA for

sequencing cannot be acquired from a single individual. DNA must be extracted from pools of

individuals, and for outbreeding organisms there may be substantial individual diversity within

the pool. Here, we use results from our simulated sequences to guide experimental ONT

sequencing and assembly for four strains of the highly heterozygous *Caenorhabditis remanei* and

C. latens. We present the experimental and computational protocols that result in high quality

assembled genome sequences.

We find that ONT approaches can produce contiguous assembled sequences for the

organisms we study with relatively high sequencing coverage of >60x, well above current

recommendations [18]. Pure ONT sequencing and assembly outperforms our tested hybrid

approach. We find that contiguous assemblies cannot be achieved by solely increasing ONT

sequencing depth as errors accumulate and assembly statistics plateau. Pre-assembly filtering and

read correction improve contiguity, and post-assembly polishing using short Illumina DNA

sequence reads increases accuracy. We also find that the use of Illumina data, even at very low

sequencing depths, increases accuracy through iterative polishing. Our results demonstrate that

rigorous experimental design can improve our ability to infer a genome sequence and reduce the

effort and cost required.

Data Description

Simulated sequence libraries

We obtained ONT sequences (R9 chemistry) from the National Center for Biotechnology

Information (NCBI) Sequence Read Archive (SRA) on February 2, 2019 [19]. The *E. coli*

dataset contained 11,652,330 sequenced bases in 120,151 reads (~2.5x coverage of the 4.64Mb

reference sequence), the *C. elegans* dataset contained 8,860,671,330 sequenced bases in 583,466

reads (~87.9x coverage of the 100.8Mb reference sequence), the *A. thaliana* dataset contained

3,421,779,258 sequenced bases in 300,071 reads (~25.22x coverage of the 135.67Mb reference

sequence), and the *D. melanogaster* dataset contained 4,617,842,308 sequenced bases in 663,784

reads (~32.39x coverage of the 142.57Mb reference sequence). We obtained the reference

genome sequence for *E. coli* strain K12_MG1655 from NCBI; all other reference genome

sequences were obtained from Ensembl (Release 95). Accession numbers are provided under

'Resource Availability.'

We simulated 150bp paired-end Illumina DNA libraries with the software ART [20] and

ONT DNA libraries with the software NanoSim (version 2.0.0; [21]). Both software programs

utilize an assembled sequence to generate simulated libraries with read profiles similar to an

empirically generated DNA library. NanoSim additionally requires real ONT flowcell data to

simulate unique, organism-specific mismatch, insertion and deletion rates. ART libraries were

simulated to a coverage of 300x with a fragment standard deviation of 50. NanoSim libraries

mirror the characteristics of the real ONT flowcells including read length distributions. We used

NanoSim to simulate 500,000 *E. coli* reads (943x coverage; read N50 8,158), 2,000,000 *C.*

elegans reads (336x coverage; N50 21,559), 5,000,000 *A. thaliana* reads (420x coverage; N50

19,577) and 10,000,000 *D. melanogaster* reads (219x coverage; N50 11,955). The sequence read

N50 indicates 50% of the total sequenced nucleotides are in reads that long or longer.

C. remanei *sequence libraries*

Output for each flow cell used for each strain is summarized in STable 1. Based on an estimated

genome size of 125Mb [22] we generated 76x-156x coverage of each strain with sequence read

N50 8,203-22,891bp.

Genome assembly

Assembly of Simulated Data

Genome sequences were assembled using three approaches. We first assembled the simulated

ONT read sets with Canu v1.9 [18]. Each genome was assembled at decreasing coverage depths

until the assembler failed to produce an assembly with the given data. In order to minimize the

influence of individual reads and stochastic assembly artifacts, each read set was generated by

selecting a random subset of the full simulated dataset. The second approach used the same

subsets of ONT data, but with Flye 2.8 [23] as the assembly software.

The long-read datasets that performed the best were paired with 2x150 simulated paired-

end Illumina data and assembled using MaSuRCA version 3.3.9 [24]; coverage depths were

adjusted for both datasets to better understand the effects of increasing or decreasing coverage on

the final assembly. Here, we retained the ONT dataset to maximize our ability to draw parallels

between assembly approaches. For example, the minimum *C. elegans* ONT dataset that

assembled with Canu [18] was an average of ~60x coverage across the genome and this read set

was used in the MaSuRCA trials with 50x and 100x Illumina coverage, respectively.

We also used Flye [23] to assemble read sets that had been corrected using the 'canu –

correct' function. Canu correction uses all-versus-all overlap information to correct each

individual read [18]. In addition to the consensus information, Canu employs two filters to avoid

biases due to sequence quality or repeats. From read-length estimates, Canu uses the longest

available reads up to the user-specified coverage depth for correction [18]. By default, this is set

to 40x coverage. Assembling these datasets with both Canu and Flye allowed us to compare the

effectiveness of the error correction modules within each software package. We also assembled

libraries with reads selected by the Canu correction module but not subject to correction to test

the influence of error correction vs. read length on assembly.

The most contiguous assemblies from the long-read only and hybrid categories were error

corrected using Pilon version 1.23 [25] to determine the effect of short-read polishing on the

accuracy of the draft assemblies. Each simulated assembled sequence was polished with the

entire simulated paired-end data set. Four rounds of polishing were completed for each assembly

with statistics measured after each round with QUAST v5.0.2 [26] and BUSCO v4.0.3 [27].

Assembly of ONT Nematode Data

Preliminary assemblies were created for each strain by first correcting the complete

library with canu –correct (Canu v2.0) [18]. Corrected reads from Canu were then assembled

using Flye 2.8 [23]. The draft assembly for each strain was polished using Pilon v1.23 [28] and

Illumina paired-end reads. All data sets contained a small sub-set of contaminants from the

nematode microbiome that were removed from the final assemblies [29, 30].

Analyses

Assessing the quality of assembled simulated sequences

We measured genome statistics relative to the reference sequence of each organism with QUAST

[26]. We assessed contiguity and accuracy of the assembled sequence through eight statistics:

(1) **NG50** is a size median statistic and indicates that 50% of the expected assembled genome

sequence (where the expectation is based on a known reference) is contained in contiguous

sequences that large or larger.

(2) **NGA50** is a similar size median but indicates that 50% of the expected assembled genome

sequence that aligns to the reference genome is contained in contiguous sequences that large or

larger.

(3) **LG50** is the number of linkage groups or contiguous assembled sequences containing 50% of

the expected assembled genome sequence.

(4) **LGA50** is similar but again measured in the portion of the assembled sequence that aligns to

the reference genome.

(5) **Genome fraction (%)** is the fraction of reference genome captured in the assembled

sequence.

(6) **Duplication** measures the fraction of reference genome found multiple times in the

assembled sequence.

(7) **Mismatches** is the number of mismatched nucleotides per 100,000 nucleotides or base pairs

(bp).

(8) **Indels** are the number of insertions and/or deletions per 100,000 nucleotides.

For our metazoan organisms we also used the software package BUSCO (version 4.0.3;

released February 2020) to search for a set of unique genes expected to be conserved in single

copy in an evolutionarily related group of organisms [27]. Below, we describe the analyses of

each organismal dataset.

Escherichia coli

The *E. coli* genome is relatively small at 4.64Mbp and contains few repeats (87.8% of the

genome codes for proteins; [31]). Assembly of 50,000 ONT sequences (~62x coverage; N50

8,350bp) with Canu [18] and Flye [23] resulted in circular contigs with high contiguity and few

mismatches, insertions, and deletions (Fig. 2A-D). At low ONT coverage (~19x) the assemblies

produced by Canu and Flye were contained in 6 and 9 contigs, respectively. Both assembled

sequences had high rates of mismatches, insertions and deletions when compared with sequences

assembled from high coverage libraries (Fig. 2C-D). Increasing ONT coverage beyond 62x

resulted in poor contiguity for Canu assembled sequences (Fig. 2A) and increased mismatches

for Flye assembled sequences (Fig. 2C). Assembly of hybrid Illumina-ONT libraries with

MaSuRCA [24] resulted in sequences with variable contiguity and accuracy. Two of the tested

hybrid sets were able to assemble the genome into a single contig but the library with the highest

Illumina coverage (300x) and highest ONT coverage (227x) assembled into two contigs.

MaSuRCA was unable to perform as well as either Canu or Flye when given the same long-read

dataset. For instance, the top performing Canu and Flye runs used 50,000 ONT reads (~62x

coverage); the same reads passed through MaSuRCA produced 2 contigs, regardless of Illumina

coverage (Fig. 2B).

After polishing the assemblies with Pilon [25], the Canu-assembled sequence had fewer

mismatches and indels per 100 Kbp than the Flye assembled sequence or an assembly produced

by Flye using reads that had first been corrected with Canu (SFig. 1A-C). Despite its dependence

on higher accuracy Illumina sequences the MaSuRCA assembled sequence had higher numbers

of mismatches, insertions and deletions both pre- and post-polishing (SFig 1B). We used

MaSuRCA to assemble a set of paired end Illumina sequences to test the influence of adding

ONT data to the assembly process. The resulting sequence was 98.76% of the *E. coli* genome

and was fragmented into 74 contiguous pieces. The accuracy, 1.7 mismatches and 0.04 indels per

100 Kbp, was improved when compared with the hybrid assembled sequences. This indicates

that the inclusion of ONT data can introduce errors in hybrid approaches that cannot be corrected

with post-assembly polishing.

Caenorhabditis elegans

The *C. elegans* genome is 100.8 Mbp [32] contained in a single X chromosome and five

autosomes [33]. *C. elegans* is a diploid self-fertile hermaphrodite with low levels of genetic

diversity [34] and approximately 16% of the genome is repetitive [35]. Canu, Flye, and

MaSuRCA assembled sequences plateaued in contiguity and accuracy above ~70x coverage

(Fig. 3A-D). The read N50 for these datasets ranged from 20,742 to 22,840. Both Canu and Flye

assembled chromosome-scale contigs with low rates of mismatches, insertions and deletions.

The MaSuRCA [24] hybrid assembly approach did not perform as well, even with high ONT

coverage (Fig. 3A,B).

To improve these assemblies we experimented with polishing the Canu and Flye

assembled sequences, Additionally, we used Flye to assemble the Canu corrected dataset

("Corrected" in Fig. 4) and used Flye to assemble the reads selected by the Canu correction

module but not corrected ("Selected" in Fig. 4). The dataset corrected with Canu and assembled

with Flye had a read N50 of 28,641bp and produced 6 chromosome-scale contigs with low

numbers of mismatches, insertions and deletions that further decreased after polishing (Fig. 4A-

D). Flye assembly of the same dataset prior to correction resulted in an assembly with higher

fragmentation (Fig. 4B) and lower accuracy (Fig. 4C-D). This dataset had a read N50 of

29,404bp, indicating that both read length and correction contribute to assembly quality. The

assembly graph for the Canu corrected, Flye assembled dataset (Fig. 5) shows 6 chromosome-

scale contigs with smaller disjunct assembled sequences [36].

We used QUAST [26] to identify annotated features in our assembled sequences to

compare each software's ability to assemble different classes of genomic elements. We used the

WS277 annotations provided by WormBase Parasite and identified exons, introns, pseudogenes,

repeats, small RNAs, transposable elements (TEs) and untranslated regions (UTRs). The worst

performing assembly had >99.67% representation of exons, introns, small RNAs and UTRs and

>98.85% representation of repetitive sequences like pseudogenes, repeats and TEs (Table 1). The

Flye assembled sequences had the best representation of exons, introns, small RNAs and UTRs

while the Canu assembled sequences had the best representation of repetitive pseudogenes,

repeats and TEs.

Drosophila melanogaster

The 180Mb genome of the diploid fruitfly *D. melanogaster* presents multiple challenges for

sequencing and assembly including approximately 60Mb of repetitive heterochromatin [37]. The

genome sequence is contained in a large X chromosome, a small ("dot") Y chromosome and 3

autosomes. Assemblies of the simulated ONT *D. melanogaster* libraries repeated patterns seen

with the *C. elegans* dataset; the most contiguous Canu assembly was produced with ~113x ONT

coverage and a read N50 of 11,679 (~16Gbp of data; Fig. 6A,B). This assembly produced 145

contiguous pieces but many of these were small and the LGA50 was 4 contigs (Fig. 6B). The

Flye assembly of the same dataset was much less contiguous, producing 482 contigs. When the

dataset was corrected with Canu then assembled with Flye the assembled sequences had high

accuracy (SFig. 2).

265   The top performing Canu assembly contained 91% of the metazoan genes expected to be

conserved in single copy with BUSCO [27] prior to polishing and 94% after four rounds of Pilon

with paired-end reads (SFig. 3). In comparison, the *D. melanogaster* reference sequence contains

94.1% of the expected single copy metazoan genes. Additional data decreased contiguity with a

slight increase in accuracy and NGA50 (Fig. 6 A,D). Hybrid MaSuRCA [24] assemblies for *D.*

melanogaster performed markedly worse than Canu assemblies of ONT data (Fig. 6B). The top

performing MaSuRCA assembly produced 220 contigs with 93.9% of the expected metazoan

genes identified [27].

Arabidopsis thaliana

The 135Mbp genome sequence of the self-fertile plant *A. thaliana* is contained in 5 autosomes

[38]. *A. thaliana* has undergone at least two rounds of whole genome duplication [39] and

contains large tracts of highly similar genomic regions. Assembly of the ONT libraries with

Canu [18] and Flye [23] resulted in sequences with lower contiguity and higher fragmentation

when compared with the other model organisms, even when the library contained ~420x

coverage and a read N50 of 19,577bp (56.7Gbp of sequenced nucleotides). Canu produced 30

contiguous pieces (LGA50 5 chromosomes) with 98.8% of the expected Viridiplantae genes

(Supplementary Data) identified prior to polishing [27] and Flye produced 26 contigs and 98.8%

of the expected Viridiplantae genes. The Flye assembly contained many more mismatches and

indels than its Canu counterpart, but this discrepancy was alleviated with Pilon polishing (SFig.

5). Following polishing with Pilon [25], the Flye assembled sequences contained 99.1% of the

expected Viridiplantae genes [27], matching that of the TAIR10 reference genome for *A.*

thaliana. Interestingly, our combined approach of long-read methods did not work with *A.*

thaliana datasets and neither Flye nor Canu was able to assemble a draft genome with <40x

coverage.

The hybrid MaSuRCA assemblies for *A. thaliana* performed the best of the three

eukaryotes. The top performing MaSuRCA [24] assembly produced 45 contiguous pieces with

98.8% of the expected Viridiplantae genes identified in the sequence [27]. This was also brought

up to 99.1% after polishing with Pilon (Supplementary Data).

Caenorhabditis remanei and *Caenorhabditis latens*

We used our simulated data to create a protocol for *de novo* sequencing and assembly for 3

strains of the nematode *C. remanei* [40] and one of the closely related nematode *C. latens* [41].

Both *C. remanei* and *C. latens* are obligate outcrossing species with high levels of nucleotide

diversity [42] that have hobbled previous assembly attempts [35, 43]. The worm's small size

means that pools of individuals harboring both individual and population-level diversity must be

sacrificed for DNA sequencing and assembly. We prepared high-molecular-weight DNA extracts

and generated ONT libraries for an outbred *C. remanei* strain EM464, two inbred laboratory

strains PX356 and PX439 and the inbred laboratory *C. latens* strain PX534. We tested library

preparation and computational protocols with the inbred *C. remanei* strain PX356 and used these

to generate assembled sequences for the other *C. remanei* and *C. latens* strains.

The most contiguous *C. remanei* PX356 assembly was achieved by using Canu [18] to

correct the entire ~156x coverage ONT dataset prior to assembly with Flye [23]. After

correction, the ~40x coverage dataset had a read N50 of 21,340bp and resulted in an assembled

sequence contained in 163 contigs with 89.1% of the expected conserved nematode genes [27].

Decontamination with Blobtools2 [29] and SIDR [30] resulted in a final assembly for PX356

with 124,499,812bp contained in 70 contigs with an N50 of 6.2Mbp and an N90 of 1.93Mbp

contained in just 18 large contigs, approximately 3 per chromosome. We compared our results

(Fig. 7A-D) with a recently published sequence for the related *C. remanei* strain PX506 [22]

generated with Pacific Biosystems sequencing and chromatin conformation capture to produce

chromosome-scale sequences [44].

Canu-only and Flye-only assemblies contained more contigs and a smaller N50 than the

combined approach described above (Fig. 7A-D). The assembly graph (Fig. 8) shows several

large contiguous fragments but also multiple regions difficult to disentangle and smaller disjunct

fragments. Analyses with GenomeScope [45] suggested ~0.7% of the genome remains

heterozygous (SFigure 6). Assemblies with subsamples of the long-read data were increasingly

fragmented with decreasing coverage. Flye assemblies with initial coverages of 102x, 70x, and

52x produced 183, 242, and 267 contigs respectively. A MaSuRCA-hybrid approach [24], using

102x ONT coverage and 450x paired-end coverage yielded 336 contigs and 96.6% of BUSCO

single-copy genes.

BUSCO identified 97.7% (increased from 89.1%) of the expected conserved single copy

genes following 4 rounds of polishing using Pilon [28] with Illumina paired end reads at ~225x

average depth (Fig. 6). We tested the influence of Illumina coverage on polishing and found that

97.6% of the expected conserved nematode genes could be identified after polishing with just

~20x Illumina coverage (STable 3), indicating that a large amount of data is not necessary to

correct the majority of errors in an assembly. However, this was achieved after three successive

rounds of error correction with the Pilon software package [28] utilizing the same Illumina DNA

sequence reads. This assembled sequence is less fragmented and contains a higher percentage of

conserved nematode genes (Fig. 6) when compared with the previously published assembly for

C. remanei PX356 produced with Illumina libraries and a genetic linkage map [35].

We generated assembled genome sequences for *C. remanei* PX439, *C. remanei* EM464,

and *C. latens* PX534 using the 'best practices' protocols developed through simulations and our

C. remanei PX356 experimentation. Our *C. remanei* PX439 dataset had an ONT coverage depth

of ~120x with a read N50 of 18,233bp. After Canu correction the read N50 increased to

38,434bp and we assembled a 132,046,643bp genome sequence contained in 53 contigs (N50

12.6Mbp). The N90 for this assembled sequence was 2.27Mbp contained in 12 large contigs,

approximately 2 per chromosome.

C. remanei EM464 is outbred and retains high levels of heterozygosity. We did not

expect to assemble a cohesive genome sequence but generated ONT libraries and an assembled

sequence to identify the distribution of allelic variants in the population. The ONT library had a

coverage depth of ~76x and a read N50 of 22,891bp. The Canu corrected dataset used for Flye

assembly had an N50 of 34,951bp. Flye produced a 138,626,018bp genome sequence contained

in 320 contigs (N50 3.4Mbp). The flow-cytometry estimated genome size of *C. remanei* is 124-

131Mbp and we estimate that 5-12% of the assembled sequences are allelic variants. The *C.*

latens PX534 dataset had a coverage depth of ~119x and a read N50 of 14,759bp. After Canu

correction the read N50 of the dataset 34,591bp and we assembled a 120,367,464bp genome

sequence in 97 contigs with an N50 of 13.6Mbp.

Discussion

We have found that Canu [18] and Flye software packages [23] produce high-quality contiguous

draft assemblies from ONT libraries for a broad range of organisms. In many cases, Flye slightly

outperforms Canu while being much faster and using fewer computational resources [13, 46],

however there are many instances in our results where the best assembly is produced by first

error correcting the data with the Canu correction modules, then assembling the corrected reads

with Flye. Given the same set of corrected reads, the Flye assembler regularly creates a better

assembly than Canu. This finding highlights the need for performing multiple assembly

strategies to identify the top performing strategy for each dataset.

We found the hybrid assemblies produced by MaSuRCA [24] contained a higher

proportion of expected conserved genes when compared with the unpolished long-read only

assemblies and had fewer mismatches and insertion/deletion errors. However, given the same

amount of ONT data, both Canu [18] and Flye [23] assembled more contiguous genome

sequences. The long-read only assemblies also produced larger NG50 values and contained a

larger fraction of the expected genome when compared to hybrid MaSuRCA assemblies. The

shortfalls of long-read only assembly can be overcome by 'polishing' with low coverage (~20x)

Illumina DNA sequences using a software such as Pilon [28] to error correct the draft

assemblies. We found that polishing played a larger role in increasing accuracy of the real-world

data compared to the results seen with simulated datasets. These methods improved the accuracy

of Canu or Flye assemblies to be on par with, or better than, those produced by MaSuRCA.

These highly contiguous assemblies were achieved with relatively high sequencing depth,

at least 100x coverage across the genome. Although ONT can theoretically produce megabase-

sized reads, in reality, many of the sequence reads in 'real' projects are shorter due to

fragmentation that occurs during DNA purification and library preparation. ONT libraries may

have many more small reads and >100x 'real' coverage may be necessary to achieve >40x

coverage with reads >25 Kbp and highly contiguous assemblies. Increasing read N50 in a library

through improvement of DNA isolation and library preparation methods reduces the necessary

coverage to produce a high-quality assembly [47]. Therefore, researchers should adjust their

coverage goals based on the quality of libraries they are able to reliably produce.

Our findings with simulated datasets were supported by all real-world datasets tested.

Caenorhabditis remanei PX356, PX439 and *C. latens* PX534 all have previously published

genome assemblies generated with paired-end Illumina sequences [35]. Our assemblies proved to

be at least one order of magnitude more contiguous than any of the previous assemblies,

highlighting the ability of ONT long reads to drastically improve contiguity. Outbreeding

Caenorhabditis are resistant to inbreeding [48] and previous assembly attempts resulted in over-

assembly due to residual heterozygosity [49] or under-assembly. Despite this, we were able to

finalize a highly contiguous and complete assembly for these nematodes. The strains we targeted

are closely related and interfertile [50] yet their assembled genome sequences vary up to 10% in

length. This level of structural variation between closely related species highlights the need for

de novo assembled sequences in molecular evolution research.

Our *de novo* assembled *Caenorhabditis* sequences are contiguous and achieved BUSCO

scores greater than those of Illumina-only assembled sequences (STable 4). The findings we

present here clearly demonstrate existing genome sequences can be improved by adding in ONT

datasets. As sequencing and assembly technologies continue to improve and drop in cost, many

researchers will find published datasets can be updated to further improve downstream analyses.

Potential Implications

In light of our findings, we suggest that long-read data be prioritized when undertaking *de novo*

genome assembly projects. Our results indicate that an assembly with sufficient ONT long read

coverage will be highly contiguous and polishing with Illumina data can achieve high levels of

accuracy. For situations where high-coverage ONT libraries are not feasible, MaSuRCA-

assembled [24] Illumina and ONT read sets can produce reliable draft sequences. However, the

quality and contiguity of the assembled sequence is determined by Illumina read depth and effort

should be made to increase Illumina read depth, even if it is at the expense of ONT sequences.

MaSuRCA-assembled Illumina sequences have fewer mismatches and insertion/deletion errors

when compared with MaSuRCA-assembled ONT and Illumina hybrid read sets, indicating that

the inclusion of ONT sequences introduces errors. We suggest that error correction with Illumina

DNA sequences and the Pilon software package [25] is a necessary finishing step in any

assembly project utilizing ONT data.

Our results demonstrate that near-chromosome-level genome sequences are achievable

with sufficient ONT data. However, chromosome-level genome assemblies are often not

necessary to address many research questions, particularly those focused on small numbers of

genes or phylogenomic information. Researchers should approach genome sequencing by first

determining what genome-completion level will be sufficient for their research goals. To aid in

this approach, we hope our study will help researchers determine the amount of sequencing

effort, and the sequencing approaches, that will best suit their needs.

Resource Availability

Lead Contact:

Janna L. Fierst: janna.l.fierst@ua.edu

Materials Availability:

This study did not generate new unique reagents.

Data and Code Availability:

E. coli genome sequence GCF_000005845.2; ONT read set SRR8154670

C. elegans genome sequence GCA_000002985.3; ONT read set ERR2092776

A. thaliana genome sequence GCA_000001735.1; ONT read set ERR2173373

D. melanogaster genome sequence GCA_000001215.4; ONT read set SRR6702603

C. remanei PX356 BioProject PRJNA248909

C. remanei PX439 BioProject PRJNA248911

C. remanei EM464 PRJNA562722

C. latens PX534 BioProject PRJNA248912

Full datasets are available at https://doi.org/10.5061/dryad.3r2280gd2

Bioinformatic scripts and workflows are available at

https://github.com/jmsutton2

Methods

DNA Extraction and Sequencing

Nematodes used for genomic sequencing were grown on two 100mm NGM plates [51]

seeded with *E. coli* OP50. Worms were harvested by washing plates with M9 minimal media

into 15mL conical tubes. Samples were rocked on a tabletop rocker for 1 hour before being

centrifuged to pellet worms. The supernatant was removed and tubes refilled with sterile M9,

mixed and pelleted by centrifugation again. This process was repeated five times or until the

supernatant was clear after centrifugation. The pellet was moved to 2mL tubes and frozen at –

20°C until extraction. Worm pellets were allowed to thaw to room-temperature then flash frozen

in liquid nitrogen and repeated three times. Worm pellets were placed in 1.2mL of lysis buffer

solution (100mM EDTA, 50mM Tris, and 1%SDS) and 20μL of Proteinase K (100mg/mL).

Tubes were then placed on a 56°C heat block for 30 minutes with shaking. Genomic DNA was

then isolated using a modified phenol-chloroform extraction[52]. DNA concentrations and purity

were measured with a Qubit 4 (Life Technologies, Carlsbad, CA USA) and NanoDrop® 1000

spectrophotometer, respectively (Thermo Fischer Scientific, Waltham, MA USA). Extracts were

visualized on a 0.8% agarose gel to verify high-molecular weight gDNA. DNA size selection

was carried out using the Short Read Eliminator Kit from Circulomics Inc. according to

manufacturer guidelines (Baltimore, MD USA).

DNA libraries were prepared for each sample using the SQK-LSK109 ligation

sequencing kit and loaded on to R9.4.1 RevD flow cells. The recommended protocol from ONT

was modified by replacing the first AmpureXP bead clean step with an additional treatment with

the Short Read Eliminator Kit. Approximately 700ng of gDNA from each library was loaded on

to a flow cell sequenced for 48 hours on a GridION X5 platform with basecalling performed by

Guppy v.4.0.11. ONT reads for each strain were trimmed of adapters using Porechop

(https://github.com/rrwick/Porechop) with the –discard_middle flag to remove chimeric reads.

Assembly and Decontamination of Real Datasets

We evaluated contaminants in the *C. remanei* PX356, *C. remanei* PX439, *C. remanei*

EM464 and *C. latens* PX534 draft assemblies with Blobtools2 [29] and SIDR [30]. Blobtools2

uses phylum-level taxonomic assignment, read coverage depth and GC content to identify non-

target contigs within the assembly. To prepare data for visualization with BlobToolKit, the ONT

reads were aligned to the initial assemblies using minimap2 v2.17 [53] and sorted using

SAMtools v1.8 [54]. A reference database for taxonomic identification of scaffolds was created

using blastn 2.2.31 [55] using the nt database from the National Center for Biotechnology

Information. An additional database was generated using diamond tblastx [56]. Contigs that were

>10,000bp and taxonomically identified as *Nematoda* were retained in the assembly.

SIDR [30] utilizes ensemble based machine learning to train a model capable of

discriminating target and contaminant contigs based on measured predictor variables. To prepare

data for SIDR decontamination Illumina DNA and RNA sequence reads were aligned to the draft

assembly with bwa [57] using the BWA-MEM algorithm. We used bbtools/bbmap software [58]

to calculate the average sequence coverage, length, GC, bases covered in alignment, RNA

average sequence coverage and bases covered in RNA alignment for each contig. We used these

data to train a bootstrap aggregated (bagged) decision tree model based on the blast identification

of contig origin and used this model to predict the origin, target or contaminant of each contig.

SIDR allowed us to assign probable *Nematoda* origin to contigs that lacked blast identification.

We discarded contaminants common to *Nematoda* genome assemblies including E. coli,

Pseudomonas, Serratia and Stenotrophomonas [59]. We also identified one large 4.57Mb *E. coli*

contig in the *C. latens* PX534 that had been labeled *Nematoda* by BlobToolKit and removed it

from the *C. latens* PX534 assembled sequences.

Heterozygosity analyses

We assessed heterozygosity using the software Jellyfish version 2.2.4 [Marcais 2011] and

GenomeScope [Ranallo-Benavidex 2020]. Jellyfish is a fast, memory efficient kmer counting

tool that searches for occurrences of unique kmers in raw sequence reads based on user selected

size. GenomeScope estimates genome heterozygosity, genome size, and repeat content using a k-

mer based statistical approach. Jellyfish was run using the default settings and a kmer size of 21.

This kmer size was selected based on recommendation from the maker of Jellyfish and should be

sufficient to obtain all unique kmers present. A histogram file of all kmer occurrences was

created by Jellyfish and then uploaded into the online interface of GenomeScope. GenomeScope

was run using default settings with a kmer length of 21 and a read length consistent with each

sample's sequencing parameters.

Evaluation

We used the software BUSCO version 4.0.1 [27] to identify conserved gene sets. Briefly,

BUSCO searches assembled DNA sequences for a set of unique genes that are expected to be

conserved in single copy in an evolutionarily related group of organisms. We used BUSCO

version 4.0.1 [27] and the datasets Nematoda_odb10 for *C. elegans*, *C. remanei,* and *C. latens*,

Metazoan_odb10 for *D. melanogaster*, and Viridiplantae_odb10 for *A. thaliana*. We also

measured BUSCO completeness with the Diptera_odb10 for the *D. melanogaster* assemblies but

found that in some instances our assembled sequence contained a greater proportion of

conserved genes than the reference sequence. We chose to focus on the Metazoan_odb10 for *D.*

melanogaster and present both sets of statistics in the Supplementary Data. BUSCO genes in

each assembly were classified as single copy, duplicated, fragmented or missing. For single-copy

and duplicated classifications, the target gene must be present and >95% identical to the

expected size and sequence of the reference in the database [60].

Acknowledgements

We thank Paula Adams, Denise Akob, Louis Bubrig, Rebecca Varney and Kevin Kocot for

helpful discussions. Patrick C. Phillips generously provided the PX356, PX439 and PX534

inbred strains and the Caenorhabditis Genetics Center provided the EM464 outbred strain. This

work is funded by NSF uRoL 1921585 and NSF DEB 1941854.

519
520   **Author Contributions**

521   Conceptualization, J.M.S. and J.L.F.; Methodology, J.M.S. and J.L.F.; Formal Analysis, J.M.S.,

522   J.D.M., A.C.M, and J.L.F.; Resources, J.L.F.; Writing – Original Draft, J.M.S., J.D.M. and

523   J.L.F.; Writing – Review & Editing, J.M.S., J.D.M and J.L.F.; Visualization, J.L.F.; Supervision,

524   J.L.F.

Declaration of Interests

527 The authors declare no competing interests.

References

1. Consortium IHGS. Initial sequencing and analysis of the human genome. Nature. 2001;409 6822:860. doi:doi:10.1038/35057062.
2. Schneider GF and Dekker C. DNA sequencing with nanopores. Nature Biotechnology. 2012;30:326-8.
3. Quail MA, Smith M, Coupland P, Otto TD, Harris SR, Connor TR, et al. A tale of three next generation sequencing platforms: comparison of Ion torrent, pacific biosciences and Illumina MiSeq sequencers. BMC Genomics. 2012;13 1:341.
4. Murigneux V, Rai SK, Furtado A, Bruxner TJC, Tian W, Harliwong I, et al. Comparison of long-read methods for sequencing and assembly of a plant genome. GigaScience. 2020;9 12 doi:10.1093/gigascience/giaa146.
5. Byrne A, Beaudin AE, Olsen HE, Jain M, Cole C, Palmer T, et al. Nanopore long-read RNAseq reveals widespread transcriptional variation among the surface receptors of individual B cells. Nature communications. 2017;8:16027-. doi:10.1038/ncomms16027.
6. Jain M, Olsen HE, Paten B and Akeson M. The Oxford Nanopore MinION: delivery of nanopore sequencing to the genomics community. Genome biology. 2016;17 1:239. doi:10.1186/s13059-016-1103-0.
7. Simpson JT, Workman RE, Zuzarte PC, David M, Dursi LJ and Timp W. Detecting DNA cytosine methylation using nanopore sequencing. Nature Methods. 2017;14 4:407-10. doi:10.1038/nmeth.4184.
8. Salazar A, Nobrega F, Anyansi C, Aparicio-Maldonado C, Costa A, Haagsma A, et al. An educational guide for nanopore sequencing in the classroom. Plos Computational Biology. 2020;16 1 doi:10.1371/journal.pcbi.1007314.
9. Chakraborty M, Baldwin-Brown JG, Long AD and Emerson JJ. Contiguous and accurate de novo assembly of metazoan genomes with modest long read coverage. Nucleic Acids Research. 2016;44 19:e147-e. doi:10.1093/nar/gkw654.
10. Ou S, Liu J, Chougule K, Fungtammasan A, Seetharam A, Stein J, et al. Effect of sequence depth and length in long-read assembly of the maize inbred NC358. Nature Communications. 2020;11 1 doi:10.1038/s41467-020-16037-7.
11. Sun J, Li R, Chen C, Sigwart JD and Kocot KM. Benchmarking Oxford Nanopore read assemblers for high-quality molluscan genomes. bioRxiv. 2021:2020.12.31.424979. doi:10.1101/2020.12.31.424979.
12. Chen Z, Erickson DL and Meng J. Benchmarking hybrid assembly approaches for genomic analyses of bacterial pathogens using Illumina and Oxford Nanopore sequencing. BMC Genomics. 2020;21 1:631. doi:10.1186/s12864-020-07041-8.
13. Wick R and Holt K. Benchmarking of long-read assemblers for prokaryote whole genome sequencing [version 3; peer review: 4 approved]. F1000Research. 2020;8 2138 doi:10.12688/f1000research.21782.3.
14. Mitsuhashi S, Frith M, Mizuguchi T, Miyatake S, Toyota T, Adachi H, et al. Tandem-genotypes: robust detection of tandem repeat expansions from long DNA reads. Genome Biology. 2019;20 doi:10.1186/s13059-019-1667-6.
15. Giesselmann P, Brandl B, Raimondeau E, Bowen R, Rohrandt C, Tandon R, et al. Analysis of short tandem repeat expansions and their methylation state with nanopore sequencing. Nature Biotechnology. 2019;37 12:1478-+. doi:10.1038/s41587-019-0293-x.
16. Kyriakidou M, Tai HH, Anglin NL, Ellis D and Strömvik MV. Current Strategies of Polyploid Plant Genome Sequence Assembly. Front Plant Sci. 2018;9:1660-. doi:10.3389/fpls.2018.01660.
17. Claros MG, Bautista R, Guerrero-Fernández D, Benzerki H, Seoane P and Fernández-Pozo N. Why assembling plant genome sequences is so challenging. Biology (Basel). 2012;1 2:439-59. doi:10.3390/biology1020439.
18. Koren S, Walenz BP, Berlin K, Miller JR, Bergman NH and Phillippy AM. Canu: scalable and accurate long-read assembly via adaptive k-mer weighting and repeat separation. Genome Research. 2017;27 5:722-36. doi:10.1101/gr.215087.116.
19. Ncbi Resource Coordinators. Database Resources of the National Center for Biotechnology Information. Nucleic acids research. 2017;45 D1:D12-D7. doi:10.1093/nar/gkw1071.
20. Huang W, Li L, Myers JR and Marth GT. ART: a next-generation sequencing read simulator. Bioinformatics (Oxford, England). 2012;28 4:593-4. doi:10.1093/bioinformatics/btr708.
21. Yang C, Chu J, Warren RL and Birol I. NanoSim: nanopore sequence read simulator based on statistical characterization. GigaScience. 2017;6 4:1-6. doi:10.1093/gigascience/gix010.

22.   Teterina A, Willis J and Phillips P. Chromosome-Level Assembly of the Caenorhabditis remanei Genome Reveals
Conserved Patterns of Nematode Genome Organization. Genetics. 2020;214 4:769-80.
doi:10.1534/genetics.119.303018.
23.   Kolmogorov M, Yuan J, Lin Y and Pevzner P. Assembly of long, error-prone reads using repeat graphs. Nature
Biotechnology. 2019;37 5:540-+. doi:10.1038/s41587-019-0072-8.
24.   Zimin AV, Puiu D, Luo MC, Zhu T, Koren S, Marcais G, et al. Hybrid assembly of the large and highly repetitive
genome of Aegilops tauschii, a progenitor of bread wheat, with the MaSuRCA mega-reads algorithm. Genome
research. 2017;27 5:787-92. doi:10.1101/gr.213405.116.
25.   Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, et al. Pilon: an integrated tool for comprehensive
microbial variant detection and genome assembly improvement. PloS one. 2014;9 11:e112963.
doi:10.1371/journal.pone.0112963.
26.   Gurevich A, Saveliev V, Vyahhi N and Tesler G. QUAST: quality assessment tool for genome assemblies.
Bioinformatics (Oxford, England). 2013;29 8:1072-5. doi:10.1093/bioinformatics/btt086.
27.   Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV and Zdobnov EM. BUSCO: assessing genome assembly and
annotation completeness with single-copy orthologs. Bioinformatics. 2018;31 19:3210-2.
doi:10.1093/bioinformatics/btv351.
28.   Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, et al. Pilon: An Integrated Tool for
Comprehensive Microbial Variant Detection and Genome Assembly Improvement. Plos One. 2014;9 11 doi:ARTN
e112963

10.1371/journal.pone.0112963.
29.   Challis R, Richards E, Rajan J, Cochrane G and Blaxter M. BlobToolKit - Interactive Quality Assessment of Genome
Assemblies. G3-Genes Genomes Genetics. 2020;10 4:1361-74. doi:10.1534/g3.119.400908.
30.   Fierst JL and Murdock DA. Decontaminating eukaryotic genome assemblies with machine learning. BMC
Bioinformatics. 2017;18 1:533. doi:10.1186/s12859-017-1941-0.
31.   Blattner F, Plunkett G, Bloch C, Perna N, Burland V, Riley M, et al. The complete genome sequence of Escherichia
coli K-12. Science. 1997;277 5331:1453-+. doi:10.1126/science.277.5331.1453.
32.   Consortium CeS. Genome sequence of the nematode C. elegans. Science. 1998;282 5396:2012-8.
33.   Nigon V and Dougherty EC. Reproductive patterns and attempts at reciprocal crossing of Rhabditis elegans Maupas,
1900, and Rhabditis briggsae Dougherty and Nigon, 1949 (Nematoda: Rhabditidae). Journal of Experimental Zoology.
1949;112 3:485-503.
34.   Andersen E, Gerke J, Shapiro J, Crissman J, Ghosh R, Bloom J, et al. Chromosome-scale selective sweeps shape
Caenorhabditis elegans genomic diversity. Nature Genetics. 2012;44 3:285-U83. doi:10.1038/ng.1050.
35.   Fierst JL, Willis JH, Thomas CG, Wang W, Reynolds RM, Ahearne TE, et al. Reproductive Mode and the Evolution of
Genome Size and Structure in Caenorhabditis Nematodes. PLoS Genet. 2015.
36.   Wick RR, Schultz MB, Zobel J and Holt KE. Bandage: interactive visualization of *de novo* genome assemblies. 2015,
p. 3350-2.
37.   Adams M, Celniker S, Holt R, Evans C, Gocayne J, Amanatides P, et al. The genome sequence of Drosophila
melanogaster. Science. 2000;287 5461:2185-95. doi:10.1126/science.287.5461.2185.
38.   Kaul S, Koo H, Jenkins J, Rizzo M, Rooney T, Tallon L, et al. Analysis of the genome sequence of the flowering plant
Arabidopsis thaliana. Nature. 2000;408 6814:796-815. doi:10.1038/35048692.
39.   Bomblies K and Madlung A. Polyploidy in the Arabidopsis genus. Chromosome Research. 2014;22 2:117-34.
doi:10.1007/s10577-014-9416-x.
40.   Baird SE, Fitch DHA and Emmons SW. Caenorhabditis vulgaris sp.n. (Nematoda: Rhabditidae): A necromenic
associate of pillbugs and snails. Nematologica. 1994;40:1-11.
41.   Dey A, Jeon Y, Wang GX and Cutter AD. Global Population Genetic Structure of Caenorhabditis remanei Reveals
Incipient Speciation. Genetics. 2012;191 4:1257-U314. doi:10.1534/genetics.112.140418.
42.   Cutter AD, Baird SE and Charlesworth D. High nucleotide polymorphism and rapid decay of linkage disequilibrium in
wild populations of Caenorhabditis remanei. Genetics. 2006;174 2:901-13. doi:10.1534/genetics.106.061879.
43.   Barriere A, Yang SP, Pekarek E, Thomas CG, Haag ES and Ruvinsky I. Detecting heterozygosity in shotgun genome
assemblies: Lessons from obligately outcrossing nematodes. Genome Research 2009;19 3:470-80.
doi:10.1101/gr.081851.108.
44.   Fierst JL. Using linkage maps to correct and scaffold de novo genome assemblies: methods, challenges and
computational tools. Frontiers in Genetics. 2015;6:220.
45.   Ranallo-Benavidez TR, Jaron KS and Schatz MC. GenomeScope 2.0 and Smudgeplot for reference-free profiling of
polyploid genomes. Nature Communications. 2020;11 1:1432. doi:10.1038/s41467-020-14998-3.
46.   Shafin K, Pesout T, Lorig-Roach R, Haukness M, Olsen HE, Bosworth C, et al. Nanopore sequencing and the Shasta
toolkit enable efficient de novo assembly of eleven human genomes. Nature Biotechnology. 2020;38 9:1044-53.
doi:10.1038/s41587-020-0503-6.
47.   Ou S, Liu J, Chougule KM, Fungtammasan A, Seetharam AS, Stein JC, et al. Effect of sequence depth and length in
long-read assembly of the maize inbred NC358. Nature Communications. 2020;11 1:2288. doi:10.1038/s41467-020-
16037-7.

48. Dolgin ES, Charlesworth B, Baird SE and Cutter AD. Inbreeding and outbreeding depression in caenorhabditis
nematodes. Evolution. 2007;61 6:1339-52. doi:10.1111/j.1558-5646.2007.00118.x.
49. Barriere A, Yang S-P, Pekarek E, Thomas CG, Haag ES and Ruvinsky I. Detecting heterozygosity in shotgun genome
assemblies: Lessons from obligately outcrossing nematodes. Genome Research. 2009;19:470-80.
50. Dey A, Jin Q, Chen Y-C and Cutter AD. Gonad morphogenesis defects drive hybrid male sterility in asymmetric
hybrid breakdown of Caenorhabditis nematodes. Evolution and Development. 2014;16 6:362-72.
51. Stiernagle T. Maintenance of *C. elegans*. Wormbook. 2006.
52. Sambrook J. Molecular cloning : a laboratory manual. Third edition. Cold Spring Harbor, N.Y. : Cold Spring Harbor
Laboratory Press, [2001] ©2001; 2001.
53. Haft DH, DiCuccio M, Badretdin A, Brover V, Chetvernin V, O'Neill K, et al. RefSeq: an update on prokaryotic
genome annotation and curation. Nucleic Acids Res. 2018;46 D1:D851-d60. doi:10.1093/nar/gkx1068.
54. Genome Project Data Processing S, Wysoker A, Handsaker B, Marth G, Abecasis G, Li H, et al. The Sequence
Alignment/Map format and SAMtools. Bioinformatics. 2009;25 16:2078-9. doi:10.1093/bioinformatics/btp352.
55. Petrov VM, Nolan JM, Bertrand C, Levy D, Desplats C, Krisch HM, et al. Plasticity of the Gene Functions for DNA
Replication in the T4-like Phages. Journal of Molecular Biology. 2006;361 1:46-68.
doi:https://doi.org/10.1016/j.jmb.2006.05.071.
56. Buchfink B, Xie C and Huson D. Fast and sensitive protein alignment using DIAMOND. Nature Methods. 2015;12
1:59-60. doi:10.1038/nmeth.3176.
57. Li H and Durbin R. Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics. 2009;25
14:1754-60. doi:10.1093/bioinformatics/btp324.
58. B. B. BBMap short-read aligner, and other bioinformatics tools. http://sourceforgenet/projects/bbmap/. 2016.
59. Fierst JL, Murdock DA, Thanthiriwatte C, Willis JH and Phillips PC. Metagenome-assembled draft genome sequence
of a novel microbial Stenotrophomonas maltophilia strain isolated from Caenorhabditis remanei tissue. Genome
Announcements. 2017;5 7:e01646-16.
60. Simao FA, Waterhouse RM, Ioannidis P, Kriventseva EV and Zdobnov EM. BUSCO: assessing genome assembly and
annotation completeness with single-copy orthologs. Bioinformatics. 2015;31 19:3210-2.

Table 1. The proportion of genomic features that QUAST identified in the assembled genome sequence. The best performing software and dataset combination is highlighted in bold for each genomic feature. Flye reliably assembled exons, introns, small RNAs and UTRs while Canu performed best when assembling pseudogenes, repeats and transposable elements. Polishing the assembled genome sequence with Illumina short reads did not affect the inference of genomic features.

Assembly Software	Dataset	Exons	Introns	Pseudo-genes	Repeats	Small RNA	TEs	UTRs
Flye	Selected	**0.9988**	**0.9993**	0.9954	0.9967	**0.9985**	0.9941	0.9996
Flye	Corrected	**0.9988**	**0.9993**	0.9954	0.9971	0.9984	0.9942	**0.9997**
Flye	Corrected, Polished	**0.9988**	**0.9993**	0.9954	0.9971	0.9984	0.9942	**0.9997**
Canu	Corrected, Polished	0.9986	0.9986	**0.9986**	**0.9981**	0.9981	**0.9978**	0.9988
MaSuRCA		0.9967	0.9973	0.9885	0.9897	0.9972	0.9887	0.9975

Figure 1. ONT sequence reads contain mixtures of errors including miscalled nucleotides, deletions, insertions and truncated homopolymers. When aligned to the reference genome this results in a large number of single and multi-base deletions, insertions and mismatches for (A) *E. coli* and (B) relatively few stretches of contiguous matching sequence that extend beyond a few nucleotides. For (C) *D. melanogaster* the error profile is similar but the large, repeat-rich genome results in multi-nucleotide deletions and insertions and again (D) few stretches of long, contiguous matching sequence.

725
726
Figure 2. The *E. coli* genome is relatively small at 4.64Mbp and less complex when compared
with metazoan genome sequences. Assembly of ONT libraries at relatively high coverage (>60x
average sequence depth) with Canu and Flye results in assembled sequences with (A) high
contiguity; (B) low contig number converging on a single chromosome; (C) few mismatches;
and (D) few insertion/deletion errors (indels) when compared with the reference sequence. The
influence of sequencing coverage on contiguity or fragmentation was not monotonic for Canu or
MaSuRCA assembled sequences. The relationship between sequencing coverage and accuracy
was not monotonic for Flye assembled sequences.
735
736
737

738
739
740 **Figure 3.** Both Canu and Flye assembled libraries with >70x ONT coverage into genome
741 sequences with A) high contiguity; B) low fragmentation; C) few mismatches; and D) few
742 insertions and deletions when compared with the *C. elegans* reference sequence. In comparison
743 MaSuRCA performed poorly across a range of ONT sequencing coverages.
744
745

Figure 4. The relative performance of Canu and Flye assembly software with raw datasets, datasets with reads selected for length ('Selected'), reads selected for length and pre-corrected ('Corrected' in the figure legend) and sequences post-assembly polished with Illumina short reads ('Polished' in the figure legend) as measured by A) genome contiguity; B) mismatches and C) insertions and deletions relative to the *C. elegans* reference sequence. Canu assembly of *C. elegans* ONT libraries produces assembled sequences with high contiguity, low fragmentation, few mismatches and relatively more insertions and deletions that are eliminated with post-assembly polishing. The performance of Flye was increased through read selection, correction and post-assembly polishing with the highest-quality *C. elegans* sequence produced through Canu correction, Flye assembly and post-assembly polishing. The dataset with reads selected for length did not assemble as well as the dataset with reads selected and corrected, indicating that correction is an important step in assembly.

759
760
761 **Figure 5.** The assembly graph for the Flye assembled Canu selected and corrected *C. elegans*
762 dataset shows 6 large sequences corresponding to the 6 chromosomes of *C. elegans* with 5 small
763 unconnected sequences. Two of the large sequences contain unresolved 'bubbles' in the
764 assembly graph.
765

Figure 6. The *D. melanogaster* genome presents multiple challenges for genome sequencing and assembly including approximately 60Mb of repetitive heterochromatin. Canu assembly of ONT libraries with >60x sequencing depth produced a sequence with (A) high contiguity; (B) low contig number; (C) few mismatches and (D) few insertions and deletions (indels) when compared with the reference. The accuracy of the Canu assembled sequence increased with ONT library coverage. Flye assembly of the same ONT libraries produced sequences with relatively less contiguity, more fragmentation, more mismatches and fewer insertions and deletions. MaSuRCA assembly of the same ONT sequences resulted in fragmented sequences with similar error levels with compared with the reference sequence.

C. remanei

783
784 **Figure 7.** Both Canu and Flye produced assembled sequences with A) contiguity; B)
785 fragmentation; C) the largest fragment and D) BUSCO gene completeness comparable to Pacific
786 BioSystems sequencing after read correction and post-assembly polishing.

787
788 **Figure 8.** The *C. remanei* PX356 assembly graph shows multiple unresolved regions, bacterial
789 contaminants (visible as circular chromosomes) and small fragments.

SFigure 1. The relative performance of Canu, Flye and MaSuRCA assembly software with raw datasets, datasets with reads pre-corrected ('Corrected' in the figure legend) and post-assembly polishing with Illumina short reads ('Polished' in the figure legend) as measured by A) genome contiguity; B) mismatches and C) insertions and deletions relative to the *E. coli* reference sequence. Both Canu and Flye assembled highly contiguous sequences for all datasets and the MaSuRCA assembled sequence was highly contiguous post-polishing. The Canu assembled sequences had few mismatches and insertion/deletion errors were eliminated through polishing. The MaSuRCA assembled sequence had numerous mismatches that were not addressed through polishing and few insertions and deletions. The accuracy of all Flye assembled sequences was improved with polishing.

SFigure 2. The relative performance of Canu and Flye assembly software with raw datasets, datasets with reads selected for length ('Selected'), reads selected for length and pre-corrected ('Corrected' in the figure legend) and sequences post-assembly polished with Illumina short reads ('Polished' in the figure legend) as measured by A) genome contiguity; B) mismatches and C) insertions and deletions relative to the *D. melanogaster* reference sequence. Canu assembly of *D. melanogaster* ONT libraries produces assembled sequences with high contiguity, low fragmentation, few mismatches and relatively more insertions and deletions that are eliminated with post-assembly polishing. The performance of Flye can be increased by selecting, correcting and polishing.

SFigure 3. Polishing the assembled *D. melanogaster* sequence with Illumina libraries and the software package Pilon (REF) increases (A) the number of conserved genes found in single copy; and reduces (B) the number of mismatches and indels compared with the reference sequence. The bulk of this improvement occurs after 1-2 rounds of polishing.

SFigure 4. Canu, Flye and MasuRCA assembled sequences across a range of ONT sequencing coverages had A) low contiguity; B) high fragmentation; C) mismatches and D) insertions and deletions when compared with the *A. thaliana* reference sequence.

SFigure 5. Post-assembly polishing increased the number of mismatches for Canu assembled sequences but decreased the number of insertions and deletions. Post-assembly polishing decreased mismatches, insertions and deletions for Flye assembled sequences.

SFigure 6. Analysis with the GenomeScope software revealed residual heterozygosity in the C. remanei PX356 sequences (estimated at 0.712%). Here, the residual heterozygosity is visible as a shoulder at ~50x sequencing coverage.

SFigure 7. Canu and MaSuRCA assembled *C. remanei* sequences with A) low contiguity and B) high fragmentation compared with the true chromosome size and number but the sequences were of comparable contiguity and fragmentation when compared with Pacific BioSystems sequences. The C) largest fragment was also comparable to that achieved by Pacific BioSystems and the BUSCO genome completeness was high with assembly of a high coverage ONT library.

SFigure 8. Polishing the *C. remanei* PX356 empirical assembled sequence with Illumina libraries and the Pilon software package (REF) increases the BUSCO completeness with a single round. The assembled sequence has a greater number of conserved genes found in single copy when compared with the published sequence (REF).

Organism	Flow Cell ID	Total # Reads (K)	Total bp passed(Gbp)
C. remanei	464.1	391	4.11
PX464	464.2	457	5.39
	534.1	545	0.579
	534.2	504	0.784
C. latens	534.3	492	2.76
PX534	534.4	824	2.35
	534.5	719	4.35
	534.6	1,020	5.41
	439.1	2,080	6.1
C. remanei	439.2	940	5.3
PX439	439.3	398	5.59
	356.1*	2,240	4.31
	356.2	940	5.01
C. remanei	356.3	1,048	4.19
PX356	356.4	750	0.723
	356.5	1,710	4.16

*Sequenced with ONT SQK-LSK108 Library Prep Kit

STable 2. Genome statistics for MaSuRCA-assembled Illumina paired sequences and error corrected Canu-assembled sequences. Although the Illumina paired end sequence is highly fragmented and incomplete it has fewer mismatches, insertions and deletions when compared with hybrid ONT and Illumina read sets.

Sample	Read Coverage		BUSCO				Mismatches per 100Kpb	Indels per 100Kpb
	ONT	Illumina	Single (%)	Duplicated	Fragmented	Missing		
A. thaliana	0	100	98.1	0.9	0.5	0.5	10.99	1.03
	420	polished	99.1	0.5	0	0.4	11.36	7.3
C. elegans	0	200	95.8	0.8	1.6	1.8	4.78	1.02
	336	polished	97.7	0.6	0.4	1.3	0.73	0.88
D. melanogaster	0	100	93.5	1.5	0.4	4.6	10.01	1.7
	108	polished	94	1.2	0.3	4.5	8.84	12.87
E. coli	0	100	-	-	-	-	1.7	0.04
	62	polished	-	-	-	-	0.02	0.02

STable 3: Polishing the *C. remanei* assembled sequence with an Illumina library at 20x depth increases the percentage of conserved genes identified by BUSCO (Simão et al., 2018) after 3 rounds of polishing with Pilon (Walker et al., 2014). Polishing with a higher depth Illumina library (114x average sequencing coverage) produces similar results after 2 rounds of polishing with Pilon. The highest percentage of conserved genes found in single copy is achieved after 3 rounds of polishing with Pilon and a high depth Illumina library at 227x average coverage.

| Sample | Polishing | | BUSCO | | | |
	Rounds	Coverage	Single (%)	Duplicated	Fragmented	Missing
Canu	0	20x	80.3	0.9	7.7	11.1
Assembled	1	20x	97.2	1	0.5	1.3
C. remanei	2	20x	97.3	1	0.4	1.3
	3	**20x**	**97.6**	**1**	**0.2**	**1.2**
	4	20x	97.6	1	0.2	1.2
	1	114x	97.6	1.1	0.3	1
	2	**114x**	**97.6**	**1**	**0.3**	**1.1**
	3	114x	97.5	1	0.3	1.2
	4	114x	97.5	1	0.3	1.2
	1	227x	97.6	1	0.3	1.1
	2	227x	97.5	1	0.3	1.1
	3	**227x**	**97.7**	**0.9**	**0.3**	**1.1**
	4	227x	97.7	0.9	0.3	1.1

STable 4

Organism	Assembly Version	ONT Coverage	Read N50	Post-correction N50	Assembly Size (Mbp)	Contigs	BUSCO Single Copy	Duplicate	Fragmented	Missing
	Published PX356	-	-	-	145.44	1,591				
C. remanei	Draft PX356	156	8,203	21,340	154.64	163				
PX356	Final PX356				124.5	70	97.7	1	0.2	1.1
	Published PX439	-	-	-	124.54	912				
C. remanei	Draft PX439	95	18,233	38,434	144.15	85				
PX439	Final PX439				131.75	53	97.6	0.9	0.3	1.2
	Published EM464	-	-	-	N/A	N/A				
C. remanei	Draft EM464	76	22,891	33,688	154.81	400				
EM464	Final EM464				138.62	320	96.2	2.6	0.3	0.9
	Published PX534	-	-	-	117.12	1,858				
C. latens	Draft PX534	119	14,759	34,951	125.13	132				
PX534	Final PX534				124.94	98	97.6	1.1	0.3	1

Accessions
PX356   GCA_001643735.2
PX439   GCA_002259225.1
EM464   N/A
PX534   GCA_002259235.1