1   Tuba1a is uniquely important for axon guidance through midline commissural structures

2

3   Georgia Buscaglia[1], Jayne Aiken[1,2], Katelyn J. Hoff[2], Kyle R. Northington[1], Emily A. Bates[1]

4

5   1. Department of Pediatrics, University of Colorado Anschutz Medical Campus

6   2. Department of Cell and Developmental Biology, University of Colorado Anschutz Medical

7   Campus

8

9   Corresponding Author:

10   Emily Anne Bates,

11   12800 E 19th Avenue, RC1 North Mail stop 8313, Aurora, CO 80045

12   Email: Emily.Bates@CUAnschutz.edu

13

14   <u>Abstract</u>

15   Developing neurons undergo dramatic morphological changes to appropriately migrate

16   and extend axons to make synaptic connections. The microtubule cytoskeleton, made of α/β-

17   tubulin dimers, drives neurite outgrowth, promotes neuronal growth cone responses, and

18   facilitates intracellular transport of critical cargoes during neurodevelopment. *TUBA1A*

19   constitutes the majority of α-tubulin in the developing brain and mutations to *TUBA1A* in

20   humans cause severe brain malformations accompanied by varying neurological defects,

21   collectively termed tubulinopathies. Studies of *TUBA1A* function *in vivo* have been limited by

22   the presence of multiple genes encoding highly similar tubulin proteins, which prevents

23   TUBA1A-specific antibody generation and makes genetic manipulation challenging. Here we

present a novel tagging method for studying and manipulating *TUBA1A* in cells without

impairing tubulin function. Using this tool, we show that a *TUBA1A* loss-of-function mutation

$TUBA1A^{N102D}$ ($TUBA1A^{ND}$), reduced the amount of TUBA1A protein and prevented

incorporation of TUBA1A into microtubule polymers. Reduced Tuba1a α-tubulin in

heterozygous $Tuba1a^{ND/+}$ mice significantly impacted axon extension and impaired formation of

forebrain commissures. Neurons with reduced Tuba1a caused by $Tuba1a^{ND}$ had altered

microtubule dynamics and slower neuron outgrowth compared to controls. Neurons deficient in

Tuba1a failed to localize microtubule associated protein-1b (Map1b) to the developing growth

cone, likely impacting reception of developmental guidance cues. Overall, we show that reduced

Tuba1a is sufficient to support neuronal migration, but not axon guidance, and provide

mechanistic insight as to how *TUBA1A* tunes microtubule function to support neurodevelopment.

Introduction

Mammalian brain development is a complex process that requires precise coordination of

multiple cell types and extracellular cues to form a fully specified tissue. Despite many advances

in understanding the cellular and molecular players involved in brain development, there is still

much that remains unknown. Insights into the molecular pathways governing neurodevelopment

can be gained from studying genetic mutations that disrupt specific aspects of brain

development. Severe cortical and neurodevelopmental phenotypes associated with mutations that

disrupt tubulin genes, termed tubulinopathies, have recently been described in humans [1-4].

Tubulinopathy mutations cause a spectrum of neurodevelopmental phenotypes, but frequently

involve cortical malformations such as lissencephaly, agenesis or hypoplasia of the corpus

callosum, and cerebellar hypoplasia [1, 2, 4, 5]. Recent studies of human tubulinopathy

mutations have revealed that each variant may impact different aspects of microtubule function,

such as protein folding, polymerization competency, and microtubule-associated protein (MAP)-

binding, among others [6-9]. Tubulin mutations can therefore be used to interrogate the

requirement for specific aspects of microtubule function throughout neurodevelopment.

Developing neurons must migrate to the correct location, extend axons to meet

sometimes distant synaptic partners and form functional connections. Throughout this process,

neurons undergo dramatic morphological changes that require coordinated interaction between

the cytoskeleton and the extracellular environment. In post-mitotic neurons, microtubule

polymers made of α/β-tubulin dimers facilitate nucleokinesis and cellular migration, support

growth cone navigation, promote axon formation and form the tracks upon which intracellular

trafficking occurs [10-13]. The microtubule network needs to be precisely controlled to fulfill

diverse functions in neurons. Microtubule properties can be modulated through post-translational

modifications (PTMs), association with MAPs and through the particular tubulin genes, or

isotypes, that a cell expresses [14]. The human genome contains at least nine unique α- and ten

unique β-tubulin genes [15, 16]. The α-tubulin isotype encoded by the gene *TUBA1A* is abundant

in the brain and is the most highly expressed α-tubulin in post-mitotic, developing neurons [17-

19]. *TUBA1A* mutations are highly represented in cases of human tubulinopathies [20],

suggesting that *TUBA1A* plays an important role in neurodevelopment. However, due to the high

degree of sequence conservation between α-tubulin genes, it has been historically challenging to

study *TUBA1A* function *in vivo*, due to the limited availability of tools.

Mouse models harboring mutations to *Tuba1a* can be used as tools to interrogate the

function of Tuba1a in the context of the neuronal milieu. As tubulin genes are often required for

life and the nucleotide sequence between isotypes is conserved, generation of mutant mouse lines

70 to study *Tuba1a* function *in vivo* has been challenging. To date, only a handful of *Tuba1a* mutant

71 mouse lines have been generated, three by ENU-induced forward genetic screens and one by

72 site-directed CRISPR gene editing [21-23]. We previously showed that the ENU-induced

73 *Tuba1a^{N102D}* allele (*Tuba1aND*) impaired microtubule function in both *S. cerevisiae* and mice

74 [22]. Homozygous *Tuba1aND* mice exhibit severely impaired brain development and are neonatal

75 lethal, similar to phenotypes seen in the *Tuba1a^{null}* and *Tuba1a*-R215* mutant mice [20-22]. In

76 homozygous *Tuba1aND*, *Tuba1a^{Null}* and *Tuba1a*-R215* mice, as well as many *TUBA1A*

77 tubulinopathy patients, cortical migration and commissural formation are severely disrupted,

78 making it difficult to infer whether axon pathfinding is a direct consequence of altered Tuba1a

79 function or if it is secondary to abnormal cortical layering and migration. *Tuba1a$^{ND/+}$*

80 heterozygous mutant mice have reduced Tuba1a function during brain development, which was

81 sufficient to support survival and cortical migration resulting in normal cortical layers [22, 24],

82 but does not support formation of commissures. Therefore, *Tuba1a$^{ND/+}$* heterozygous animals can

83 provide insight into how Tuba1a contributes specifically to axon pathfinding.

84   Here, we show that a reduction in developmental Tuba1a protein causes specific impairments

85 in axon guidance through large brain commissures. Using a novel tubulin visualization

86 technique, we demonstrate that the *TUBA1A^{N102D}* mutation prevents incorporation of *TUBA1A*

87 into microtubule polymers in cells. In mice, heterozygous *Tuba1a$^{ND/+}$* brains fail to form the

88 corpus callosum, anterior and hippocampal commissures due to apparent deficits in axon

89 pathfinding through the midline. Cultured neurons from *Tuba1a$^{ND/+}$* and wild-type cortices

90 revealed that *Tuba1a$^{ND/+}$* neurons have shorter neurites than wild-type and grow slower overall,

91 potentially due to alterations in microtubule polymerization dynamics. Further, we demonstrate

92 that *Tuba1a$^{ND/+}$* neurons fail to localize at least one critical developmental MAP to the

developing growth cone. Collectively, our data present evidence for precise mechanisms by

which loss of Tuba1a causes axonal pathfinding deficits during development.

Materials and Methods

Animals

All animal research was performed in accordance with the Institutional Animal Care and Use

Committee at the University of Colorado School of Medicine. All mice used were maintained on

a 129S1/C57Bl6 genetic background. Mice were kept on a 12:12 light:dark cycle with *ad libitum*

access to food and water. *Tuba1aND* and wild-type littermate mice were maintained on water

supplemented with 0.2g/L MgSO$_4$ to promote *Tuba1a$^{ND/+}$* survival and ability to reproduce. For

timed mating male and female mice were introduced overnight and separated upon confirmation

of mating, which was then considered embryonic day 0.5. Male and female mice were

represented in all studies. All mice were genotyped by PCR amplification of tail DNA followed

by Sanger sequencing to differentiate homozygous or heterozygous *Tuba1a$^{ND/+}$* mice from wild-

type. Primers used to amplify mouse DNA for genotyping were:

forward:TGGATGGTACGCTTGGTCTT; reverse: CTTTGCAGATGAAGTTCGCA; and

sequencing: GTCGAGGTTTCTACGACAGATATC.

Histology

Mice were anesthetized and trans-cardially perfused with 0.1M NaCl and 4% paraformaldehyde

(PFA) for histology. Tissues of interest were dissected and post-fixed in 4% PFA. Tissue

sectioning was performed on a CM1520 cryostat (Leica, Wetzlar, Germany) and 30μm

cryosections were obtained for all histology. For luxol fast blue staining, sections from brain

were stained using a 0.1% luxol fast blue solution. For immunofluorescence studies PFA-fixed

tissues or cells were blocked in phosphate-buffered saline (PBS) containing 5% goat serum or

bovine serum albumin (BSA) with 0.3% Triton-X 100. Primary and secondary antibodies were

diluted in PBS containing 1% BSA with 0.1% Triton-X 100.

Electron Microscopy

Mice used for electron microscopy were perfused with 0.1M NaCl and 2.5% glutaraldehyde 4%

PFA, after which the brain was dissected and post-fixed in 2.5% glutaraldehyde 4% PFA

overnight at 4°C. Following post-fixation, brains were sent for sectioning and imaging by the CU

School of Medicine Electron Microscopy Core facility.

Plasmids and Reagents

The hexahisitidine (His6) epitope tag was inserted in the α-tubulin internal loop region [25-27].

Codon optimization for *rattus norvegicus* (https://www.idtdna.com/codonopt) was used to

generate the His6 sequence CATCACCACCATCATCAC, which was inserted into the coding

region of human *TUBA1A* from the Mammalian Genome Collection (clone

ID:LIFESEQ6302156) between residues I42 and G43. Gibson cloning was used to insert the

gBlock of *TUBA1A* internally tagged with His6 (*TUBA1A*-His6) into the pCIG2 plasmid (shared

by Dr. Matthew Kennedy, University of Colorado) linearized with NruI and HindIII. *TUBA1A*-

His6 incorporation was confirmed by sequencing across the complete *TUBA1A* coding region.

The $TUBA1A^{T349E}$ ($TUBA1A^{TE}$) polymerization incompetent, and $TUBA1A^{E255A}$ ($TUBA1A^{EA}$)

highly polymer-stable α-tubulin mutants were identified and described in prior publications [28-

30].The GFP-MACF43 vector was shared by Dr. Laura Anne Lowery (Boston College) and Dr.

Casper Hoogenraad (Utrecht University). Myr-TdTomato plasmid DNA was shared from Mark

Gutierrez and Dr. Santos Franco (University of Colorado).

Cell Culture and Nucleofection

Cos-7 cells (Thermo Fisher Scientific, Waltham, MA; ATCC® CRL-1651™) were cultured in a

37°C humidified incubator with 5% CO_2 in DMEM (Gibco) supplemented with 10% fetal bovine

serum (Gibco), 1mM sodium pyruvate (Thermo), and penn/strep (1,000 IU/ 1,000 µg/mL;

Thermo). Cos-7 cells were transfected with 2.5µg of hexahistidine (His6) tagged *TUBA1A*

plasmid DNA using Lipofectamine 3000 (Invitrogen). For Cos-7 proteasome inhibition assays,

5µm Lactacystin A [31, 32] was added to normal culture medium for 24 hours, the day following

transfection with *TUBA1A*-His6 constructs. Dissociated neurons were cultured from male and

female P0-P2 mouse or rat cortices. Brains were removed and placed into HBSS (Life

Technologies) supplemented with 1M HEPES (Life Technologies) and 1mM kynurenic acid

(Tocris Bioscience, Bristol, UK). Meninges were removed and cortices were dissected and cut

into approximately 1mm pieces. Cortical pieces were triturated to a single-cell suspension using

glass Pasteur pipettes. Cortical neurons were plated onto 35mm Poly-D-Lysine coated glass-

bottom culture dishes at a density of 350,000 cells (Willco Wells, HBSt-3522). For nucelofected

mouse and rat neurons, 4 µg of plasmid DNA was introduced to $4x10^6$ neurons using an

AMAXA nucleofection kit (VPG-1001, VPG-1003; Lonza). AMAXA-nucleofected cells were

plated in 35mm glass bottom imaging dishes. Neurons were maintained in a 37°C humidified

incubator with 5% CO_2 in phenol-free Neurobasal-A medium (Life Technologies) supplemented

with B-27 (Thermo Fisher Scientific, Waltham, MA), Penn/strep (Thermo), GlutaMax (Thermo),

5ng/mL β-FGF (Gibco), and Sodium Pyruvate (Thermo).

RNA isolation + RTPCR

RNA was isolated from Cos-7 cells, 48-hours post-transfection using TRIzol Reagent (Thermo;

15596026). RNA concentration and purity were determined using a spectrophotometer, then

cDNA was synthesized using the RT2 First Strand Kit (Qiagen, Hilden, Germany; 330401).

qRT-PCR reactions were prepared with SYBR Green RT-PCR Master mix (Thermo; S9194) and

run with a CFX Connect Real-Time System (Bio-Rad). Samples were run in triplicate, results

were analyzed in Excel. All qPCR data presented in this manuscript was normalized to

expression of GFP, which was present on the same plasmid as *TUBA1A*-His6 constructs. Wild-

type *TUBA1A* mRNA quantity was set to = 1 and *TUBA1A*ND relative mRNA quantity was

presented relative to wild-type. For all qRT-PCR experiments 3 biological replicates were used

per genotype.

Neuron Immunocytochemistry

DIV 3 primary cortical neurons were washed with PBS and fixed with a fixation solution of 4%

PFA and 0.2% glutaraldehyde in PBS for 15 minutes at room temperature. For tubulin

extraction, cells were washed with PBS followed by PHEM buffer, then soluble tubulin dimers

were extracted using 0.1% triton with 10μM taxol and 0.1% DMSO in PHEM buffer. Extracted

cells were fixed with 2% PFA and 0.05% glutaraldehyde in PBS for 10 minutes, washed with

PBS and then reduced in 0.1% NaBH4 in PBS for 7 minutes at room temperature. Cells were

then washed with PBS and blocked in 3% BSA and 0.2% Triton in PBS for 20 minutes at room

temperature, with agitation. Immunostaining was performed using primary antibodies directed

against: 6X-Histidine (Invitrogen, 4A12E4 37-2900; 1:500), total α-tubulin (Sigma, DM1A

T6199; 1:5,000), Acetylated Tubulin (Sigma, T7451; 1:1,000), Tyrosinated Tubulin (Chemicon,

MAB1864; 1:1,000), Map1b (Santa Cruz Biotech, sc-135978; 1:500), Map2 (Novus Biologicals,

NB300-213; 1:2,000). Primary antibodies were diluted in blocking buffer and incubated

overnight at 4ºC in a humidified chamber. After primary antibody staining, cells were washed

three times with PBS. Fluorescently-conjugated secondary antibodies were diluted 1:500 in

blocking buffer and incubated for 2 hours at room temperature, protected from light. Secondary

185 antibodies were from Life Technologies (Carlsbad, CA) all used at 1:500. Alexa Fluor 568-

186 conjugated Phalloidin (Thermo, A12380; 1:20) was added during secondary antibody incubation

187 for labeling of filamentous actin. Tissues or cells of interest were mounted onto glass microscope

188 slides and sealed with glass coverslips and aqueous mounting media containing DAPI (Vector

189 Laboratories, #H-1200) and imaged on a Zeiss 780 or 880 confocal microscope with a 40X or

190 63X oil objective.

191 *Microtubule Dynamics and Neuron Growth Rates*

192 Primary neurons from wild-type and *Tuba1a*$^{ND/+}$ neonatal mouse cortices were cultured as

193 described above. Prior to plating, mouse cortical neurons were nucleofected with 4μg each of

194 GFP-MACF43 and Myr-TdTomato plasmid DNA. Following 1 day in culture, neurons were

195 transferred to a 37°C imaging chamber and time-lapse images of GFP-MACF43 comets and

196 Myr-TdTomato membrane label were acquired using a Zeiss 780 confocal microscope with 40X

197 oil objective. Following acquisition of baseline images, images were acquired every 2 seconds

198 for 2 minutes. A total of four time-lapse videos were acquired per neuron, with a 10-minute

199 waiting period in between imaging sessions. Data from all four GFP-MACF43 time points were

200 pooled by cell to generate a cell average and then grouped by genotype for further analysis.

201 Membrane-bound Myr-TdTomato images were acquired at the beginning and end of the imaging

202 period and were used to track neuronal growth. Kymographs of GFP-MACF43 comets in single

203 neurites were generated from GFP-MACF43 videos in ImageJ/FIJI (National Institutes of

204 Health) and were used to assess microtubule polymerization velocity, along with the duration

205 and distance of individual plus-end polymerization events. In brief, polymerization velocity was

206 determined by measuring the change in position (X_2-X_1; μm) divided by the change in time (Y_2-

207 Y_1; min) for each GFP-MACF43 comet, duration assessed the total time (Y_2-Y_1) each GFP-

MACF43 comet spent moving in seconds, and distance assessed the total distance (X_2-X_1)

covered by each polymerization event in μm.

Western Blotting

Protein was isolated from brains of P0-P2 mice by dounce homogenization and ultra-

centrifugation. Tubulin-enriched protein fractions with MAPs were isolated according to a

previously established protocol [33]. Cos-7 cell protein was extracted using a Tris-Triton lysis

buffer with protease inhibitor cocktail (Sigma). Protein concentrations were assessed using a

BCA assay (Thermo), and relative concentration was determined using a Synergy H1 microplate

reader (BioTek Instruments, Winooski, VT). 5μg of either whole brain lysate or tubulin-enriched

protein fraction was loaded per lane and run on 4-20% Mini-Protean TGX Stain-Free precast

gels (4568093; Bio-Rad Laboratories, Hercules, CA) at 150mV for 1hr. Prior to protein transfer,

Stain-Free gels were activated with UV light for 1 minute and imaged for total protein on gel

using a ChemiDoc MP imager (Bio-Rad). Proteins were transferred to PVDF blotting

membranes (Bio-Rad) in standard 25mM Tris-base, 192mM glycine, 15% methanol transfer

buffer, or transfer buffer optimized for high molecular-weight proteins (>200kDa) by the

addition of 0.025% SDS. Blots were transferred at 4°C and 75V for either 1 hour for standard

molecular-weight proteins, or 3 hours for high molecular-weight proteins. Immediately following

transfer, PVDF membranes were rinsed in TBST and imaged to quantify the total protein on blot

using UV-activated Stain-Free blots. Gels were also imaged post-transfer to assess transfer

efficiency for each blot. Membranes were blocked in Tris-buffered Saline containing 0.1%

Tween-20 (TBST) with 5% BSA for 1 hour and incubated in primary antibody diluted in TBST

containing 1% BSA overnight at 4°C. Primary antibodies were: mouse anti 6X-Histidine

(Invitrogen, 4A12E4; 1:500), chicken anti-GFP (Invitrogen, A10262; 1:1,000), and mouse anti-

Map1b (Santa Cruz, sc-135978; 1:500). Blots were incubated in HRP-conjugated Goat-anti-

mouse (1:5,000; Santa Cruz) secondary antibody diluted in TBST containing 0.5% BSA with

streptavidin-HRP (Bio-Rad, 1:10,000) for 1 hour at room temperature. Blots were developed in

ECL solution for 5 minutes at room temperature (Bio-Rad) and imaged.

Experimental design and statistical analyses.

Band volume of all Western blots was analyzed using Image Lab software (Bio-Rad).

Fluorescence intensity measurements, area and morphological assessment, kymograph

generation, and quantification of EM images was performed using ImageJ/FIJI software (NIH)

and Excel (Microsoft). Statistical analyses were performed, and graphs were created using Prism

version 8.0 (GraphPad). Most graphs display all data points to accurately represent the variability

in each dataset, except in cases where such presentation obscured visibility. For all statistical

analyses, statistical significance was considered to be $p < 0.05$. Statistical analyses used in each

experiment are indicated in their respective figure legends. For all graphs mean ± SEM was

reported unless otherwise noted. Normality of each dataset was assessed using a Shapiro-Wilk

test. In datasets with two groups, parametric data was analyzed using a Student's t-test, while

non-parametric data was assessed by Mann-Whitney *U* analysis of medians. Multiple groups

were compared by one-way or two-way ANOVA and analyzed *post hoc* by either a Bonferroni

or Kruskal-Wallis test for parametric and non-parametric data, respectively.

Results

***Tuba1a* is required for formation of midline commissural structures**

TUBA1A is a major component of developing neuronal microtubules, and is critical for

proper brain development [20]. Human *TUBA1A*-tubulinopathy patients and homozygous

Tuba1a mutant mouse models both exhibit severe brain malformations. *Tuba1a*$^{ND/+}$ heterozygous

mutant mice undergo normal cortical migration, display comparable brain weight to wild-type

littermates at birth, and survive to adulthood [22, 24]. However, *Tuba1a*$^{ND/+}$ brains exhibit

agenesis of the corpus callosum and abnormal formation of the anterior and hippocampal

commissures (Fig. 1A). In wild-type mice, nascent callosal 'pioneer' axons extend through

midline at E15.5, and early 'follower' axons begin extending at E17 in mice [34]. Evidence of

abnormal callosal projections were apparent as early as P0 in *Tuba1a*$^{ND/+}$ brains, as seen by the

early formation of aberrant axon bundles adjacent to the callosum, known as Probst bundles (Fig.

1A)[35]. In addition to agenesis of the corpus callosum at midline, lateral regions of adult

Tuba1a$^{ND/+}$ corpus callosum were found to be significantly thinner than wild-type (Fig. 1B).

Similarly, adult *Tuba1a*$^{ND/+}$ anterior commissures are smaller than that of wild-type littermates

(Fig. 1C). In wild-type mice, corpus callosum thickness and anterior commissure area both

increased significantly between P0 and adulthood; however, normal postnatal expansion of these

tracts was not evident in *Tuba1a*$^{ND/+}$ mice (Fig. 1B, C). *Tuba1a*$^{ND/+}$ axons fail to organize into

typical midline commissural structures, indicating that half of the normal complement of Tuba1a

during brain development is not sufficient for commissural axon guidance.

Figure 1 *Tuba1a is required for formation of midline commissural structures.*
A. Luxol fast blue-stained coronal brain sections from postnatal day 0 (P0; top) and Adult (middle-bottom) wild-type and *Tuba1a^{ND/+}*
mice. Images portray abnormal midline commissural formation in *Tuba1a^{ND/+}* mouse brains, with labels highlighting the corpus callosum
(CC), anterior commissures (AC), and hippocampal commissure (HC). Scale bars are 500μm. ***B.*** Scatter plot representing corpus
callosum width at P0, 3-months, and 10-months-old. Arrows indicate Probst bundles. Asterisk in *A.* shows where measurements for *B.*
were obtained. ***C.*** Scatter plot displaying anterior commissure area in P0, 3-month, and 10-month-old wild-type and *Tuba1a^{ND/+}* brains.
Wild-type and *Tuba1a^{ND/+}* measurements compared by two-way ANOVA. * $p<0.05$; ** $p<0.01$; *** $p<0.001$; and **** $p<0.0001$.

 Examination of sagittal brain sections taken directly at midline revealed dramatic

 disorganization of corpus callosum axons in *Tuba1a^{ND/+}* brains (Fig. 2A). Compared to wild-

 type, *Tuba1a^{ND/+}* midline commissural axons were largely absent, and the existing axons failed to

 organize into a tract with uniform orientation (Fig. 2A). Despite dramatic differences between

 wild-type and *Tuba1a^{ND/+}* callosal axon organization, *Tuba1a^{ND/+}* axons were highly colocalized

 with immunolabeled myelin sheaths (Fig. 2A). To further assess the impact of *Tuba1a^{ND/+}*

substitution on callosal axon morphology and myelination, we performed electron microscopy

(EM) in both wild-type and *Tuba1a*$^{ND/+}$ corpus callosi. Due to the scarcity of axons present

directly at midline in the *Tuba1a*$^{ND/+}$ corpus callosum, we sampled a region of corpus callosum

2mm lateral to midline for both wild-type and *Tuba1a*$^{ND/+}$ animals (Fig. 2B). EM images

revealed a striking difference in axon density between wild-type and *Tuba1a*$^{ND/+}$ corpus callosi

(Fig. 2B, D; p=0.03). Myelin thickness, measured by G-ratio, was similar between wild-type and

Tuba1a$^{ND/+}$ brains (Fig. 2C; p=0.34), as was axon diameter (Fig. 2E; p=0.14). There was a trend

towards decreased myelination in *Tuba1a*$^{ND/+}$ animals (p=0.07), but this difference was not

statistically significant (Fig. 2F). These data provide evidence that *Tuba1a*$^{ND/+}$ callosal axons do

not correctly organize to form a commissure. Previously published data indicated that reduced

developmental Tuba1a function is tolerable for cortical neuron migration [22]; however, our

results indicate that reduced Tuba1a is not sufficient to support axon pathfinding.

Figure 2 *Axon density is reduced by Tuba1a^{ND/+} in midline corpus callosum.*
A. Sagittal brain sections at midline from wild-type (top) and *Tuba1a^{ND/+}* (bottom) brains stained with neurofilament-heavy (NFT-H; magenta) to label axons, and proteolipid protein (PLP; green) to label myelin. Images were taken at 4X (left) and 20X (right) magnification with 500μm and 50μm scale bars, respectively. **B.** Electron microscopy (EM) images of sagittal wild-type (top) and *Tuba1a^{ND/+}* (bottom) sections of corpus callosum 2mm adjacent to midline. Enlarged regions on right are denoted by boxes, scale bar is 1μm. **C.** Scatter plot of G-ratio measurements for wild-type and *Tuba1a^{ND/+}* axons. Data points represent individual myelinated axons and are clustered by animal (N=3 mice, n=100 axons; p=0.34). **D.** Scatter plot representing axon density of wild-type and *Tuba1a^{ND/+}* axons in analyzed EM images. Data points represent average values for each animal (n=6 regions containing same total area; p=0.03). **E.** Scatter plot of axon diameters in wild-type and *Tuba1a^{ND/+}* corpus callosum by EM. Only those axons captured in cross section were assessed for diameter, as skewed axons provide inaccurate measurements. Data points represent individual axons and are clustered by animal (n=100; p=0.14). **F.** Scatter plot representing the average percent of myelinated axons per animal in wild-type and *Tuba1a^{ND/+}* corpus callosum. 6 regions containing the same total area were assessed (n=6; p=0.07). Statistical differences between means of wild-type and *Tuba1a^{ND/+}* datasets were assessed by t test, with * p<0.05.

291   *TUBA1A^{ND}* α-tubulin does not incorporate into neuronal microtubules

292   Reduced developmental Tuba1a is not adequate to support long-distance axon targeting,

293   but the molecular and cellular mechanisms by which Tuba1a alters neuronal microtubule

294   function are unknown. The *Tuba1a^{ND}* allele was previously demonstrated to cause partial loss of

295 microtubule function in yeast and mice [22, 24]. However, the mechanism by which *Tuba1a*ND

296 substitution leads to loss-of-function (LOF) is unclear. The neuronal microtubule network is

297 complex, containing many different tubulin isotype proteins, PTMs, and MAPs decorating the

298 lattice, all of which can modify functional microtubule properties. Further, the high degree of

299 sequence similarity between α-tubulin isotypes has precluded study of individual tubulin

300 isotypes at the protein level *in vivo*. No α-tubulin isotype or species-specific antibodies exist for

301 TUBA1A, and prior attempts to tag TUBA1A neuronal microtubules with N- or C-terminal

302 fusion proteins have had detrimental effects on protein function [36]. These challenges have

303 made the direct visualization of specific tubulin isotypes or mutant tubulin proteins in neurons

304 historically difficult. Thus, the ways in which TUBA1A specifically contributes to neuronal

305 microtubule protein function have been difficult to ascertain. To address the need for better tools

306 to study TUBA1A protein, we generated a functional hexahistidine (His6)-tagged TUBA1A

307 construct based on a previously identified internal loop in the α-tubulin protein for *in vivo*

308 visualization and manipulation of microtubules [26] (Fig. 3A). We inserted a His6 tag into an

309 internal loop of TUBA1A between residues I42 and G43. This region of α-tubulin was

310 previously shown to tolerate addition of amino acids without disrupting tubulin function [26],

311 and previous groups added a His6 tag in this loop to affinity purify tubulin subunits for

312 reconstituted systems [25, 27]. However, to our knowledge this internal His6 tag on α-tubulin

313 has never been used for immunohistochemical assays to visualize the microtubule network *in*

314 *vivo*. Ectopically expressed wild-type *TUBA1A*-His6 is incorporated into Cos-7 cell microtubules

315 (Fig. 3B). Compared to traditional immunolabeling of cellular microtubules (DM1A), *TUBA1A*-

316 His6 provides excellent labeling of microtubule polymers (Fig. 3B).

Figure 3 *TUBA1A^{ND} impairs incorporation into cellular microtubules and reduces α-tubulin protein abundance.*
A. Schematic of *TUBA1A*-His6 tag and experimental design. His6 epitope tag was added to an internal loop on TUBA1A (top), and TUBA1A-His6 plasmid DNA was transfected into Cos-7 cells to label cellular microtubules in the presence or absence of proteasome inhibitor treatment (bottom). ***B.*** Images showing Cos-7 cells transfected with wild-type, *TUBA1A^{ND}*, *TUBA1A^{TE}* (polymerization-incompetent mutant), and *TUBA1A^{EA}* (highly polymer stable mutant) *TUBA1A*-His6 plasmid DNA. Cells were immunolabeled with α-His6 antibodies to reveal microtubule incorporation of wild-type and mutant TUBA1A-His6 protein. TUBA1A-His6 images on the left are compared to traditional immunolabeling of α-tubulin using DM1A antibody in untransfected Cos-7 cells on right. ***C.*** Western blot for His6 in TUBA1A-His6 transfected Cos-7 cell lysates. A subset of transfected cells were treated with 5μM Lactacystin A (LactA) for 24 hours to block proteasomal degradation. His6 signal was normalized to GFP, which was expressed from the same plasmid. ***D.*** Quantification of band density for His6 western blot shown in *C*. His6 band density was normalized to GFP-expressing cells in control-treated (left), and cells treated with 5μM LactA for 24-hours (right). Data were analyzed by t test. N=3 transfections; n=3 technical replicates; p=0.04 for all comparisons marked with asterisks; p=0.83 for control vs LactA-treated *TUBA1A^{ND}*. ***E.*** Bar graph representing *TUBA1A* mRNA expression in Cos-7 cells transfected with TUBA1A-His6. *TUBA1A* mRNA expression was normalized to GFP mRNA expression. Data were normalized to *TUBA1A* expression in wild-type (WT) TUBA1A-His6-transfected cells and represent 3 separate transfection experiments and 3 qRT-PCR replicates. Differences between groups were assessed by t test (p=0.97). All images were taken at 40X magnification, scale bars are 10μm. All graphs show mean of data ± SEM, *p<0.05.

   His6-tagged mutant TUBA1A can be ectopically expressed in cells to evaluate the abundance

   and polymerization capability of mutant TUBA1A proteins. We ectopically expressed three

   TUBA1A variants in Cos-7 cells: *TUBA1A^{ND}* , the mutation analogous to the *Tuba1a^{ND}* mice;

   TUBA1A^{T349E} (*TUBA1A^{TE}*) an α-tubulin mutation shown to prevent polymerization in yeast [30];

   and *TUBA1A^{E255A}* (*TUBA1A^{EA}*) an α-tubulin mutation which inhibits depolymerization and thus

   becomes locked in a polymer-bound state [28, 29](Fig. 3B). As predicted, *TUBA1A^{TE}*-His6

   protein was not highly detected in Cos-7 cell microtubule polymers, while *TUBA1A^{EA}*-His6

   protein was abundantly incorporated into cellular microtubules (Fig. 3B). In contrast to wild-

   type, but similar to the polymerization-incompetent *TUBA1A^{TE}*-His6 protein, ectopic

   TUBA1A^{ND}-His6 protein was not detected in Cos-7 cells (Fig. 3B). Western blotting of lysates

329 from Cos-7 cells, 48-hours post-transfection revealed that $TUBA1A^{ND}$-His6 protein was

330 significantly reduced compared to wild-type $TUBA1A$-His6 (Fig. 3C, D; p=0.04), despite similar

331 $TUBA1A$ mRNA levels between wild-type and $TUBA1A^{ND}$ transfected Cos-7 cells (Fig. 3E;

332 p=0.97). To evaluate if $TUBA1A^{ND}$ mutant protein is targeted for proteasomal degradation, we

333 treated $TUBA1A$-His6 transfected Cos-7 cells with 5μM proteasome inhibitor, Lactacystin A

334 (LactA [31, 32]), for 24-hours and probed for His6 abundance by western blot (Fig. 3C).

335 Treatment with LactA significantly increased wild-type $TUBA1A$-His6 protein compared to

336 control-treated cells, but had no effect on $TUBA1A^{ND}$-His6 protein abundance (Fig. 3D; p=0.83).

337 These results indicate that $TUBA1A^{ND}$ mutant protein is likely not targeted for degradation by the

338 proteasome, but may reduce cellular TUBA1A by other mechanisms.

339 We next investigated if $TUBA1A^{ND}$ substitution impairs incorporation of TUBA1A

340 protein into neuronal microtubule polymers (Fig. 4). Wild-type rat cortical neurons were

341 nucleofected with $TUBA1A$-GFP, wild-type $TUBA1A$-His6, $TUBA1A^{ND}$-His6 and $TUBA1A^{TE}$-

342 His6 DNA at day *in vitro* 0 (DIV 0; Fig. 4A). Following 2-days in culture (DIV 2), cells were

343 fixed and a subset of neurons were permeabilized to remove soluble tubulin dimers ("tubulin

344 extraction"). Extraction of soluble tubulin leaves behind only polymer-bound tubulin, enabling

345 visualization of ectopic tubulin protein polymerization competence [37]. Neurons expressing

346 $TUBA1A$-GFP exhibited abundant GFP signal in intact cells, but tubulin extraction revealed that

347 GFP fusion impaired incorporation of TUBA1A into neuronal microtubule polymers (Fig. 4B).

348 In contrast, wild-type $TUBA1A$-His6 protein was highly present in both intact and tubulin

349 extracted neurons, demonstrating that His6-tagging does not impair polymerization ability of

350 TUBA1A in neurons

Figure 4 *TUBA1A^{ND} α-tubulin does not incorporate into neuronal microtubule polymers.*
A. Schematic of cortical neuron isolation and transfection. ***B.*** Cortical rat neurons at DIV 2 transfected with *TUBA1A*-GFP.
Top panel shows neurons with intracellular environment intact (unextracted), containing soluble tubulin dimers and
polymerized microtubules transfected to express a TUBA1A-GFP fusion protein. Bottom panel shows neurons with soluble
tubulin dimers extracted, showing only GFP-labeled TUBA1A that is incorporated into microtubule polymer. ***C.*** Rat cortical
neurons at DIV 2 transfected with wild-type (WT) *TUBA1A*-His6, *TUBA1A^{ND}*-His6 LOF mutation, and *TUBA1A^{TE}*-His6
polymerization-incompetent mutant as a negative control. Top panels show composite image containing membrane-bound GFP
(green) for confirmation of transfection, α-His6 (Magenta) and DAPI (blue) immunolabeling. Middle panels show unextracted
and bottom panels show tubulin extracted DIV 2 neurons (described in *B.*), labeled with α-His6 antibodies to visualize ectopic
TUBA1A-His6 proteins. All images were taken at 63X magnification, scale bar is 25μm.

(Fig. 4C). As predicted, polymerization incompetent *TUBA1A^{TE}*-His6 mutant protein was

diffusely visible in intact neurons, but was absent from tubulin extracted neurons (Fig. 4C).

TUBA1A^{ND}-His6 protein was detectable at very low levels in unextracted neurons, but was not

visible following tubulin extraction, indicating that *TUBA1A^{ND}* impairs incorporation into

neuronal microtubules (Fig. 4C). These experiments illustrate that *TUBA1A^{ND}* reduces

abundance of TUBA1A protein upstream of proteasomal degradation and prevents incorporation

of mutant TUBA1A into cellular microtubules.

Figure 5 *Reduced Tuba1a alters neuronal microtubule polymerization at DIV 1.*
A. Schematic of nucleofection and imaging experimental setup. Neurons were nucleofected with GFP-MACF43 plasmid DNA and imaged at DIV 1 for microtubule polymerization events in a neurite region of interest. ***B.*** Representative images showing GFP-MACF43 expressing wild-type (WT) and *Tuba1a^(ND/+)* neurons at 40X magnification (left) and representative kymographs that were generated from GFP-MACF43 images (right). Scale bars are 10µm and 5µm on neuron images and kymographs, respectively. ***C.*** Bar graph representing the average polymerization velocity for anterograde microtubule polymerization events in WT and *Tuba1a^(ND/+)* cortical neurons at DIV 1 (n=688 events; p=0.0008 by Mann-Whitney test). ***D.*** Bar graph representing polymerization distance in DIV 1 WT and *Tuba1a^(ND/+)* cortical neurons (n=688 events, p<0.0001). ***E.*** Bar graph representing polymerization duration in DIV 1 WT and *Tuba1a^(ND/+)* cortical neurons (n=688 events, p<0.0001). Bars represent the mean of each group and error bars represent SEM. Differences between groups were assessed by t test, unless otherwise noted.

 Reduced Tuba1a alters microtubule dynamic properties in neurons

   Dynamic instability is a defining feature of microtubule polymers that is heavily

 influenced by a number of factors to tune microtubule network function in cells [14]. MAP-

 binding, PTMs, and incorporation of different tubulin isotypes can all influence microtubule

365 dynamics [38-41]. *TUBA1AND* tubulin is less abundant than wild-type and does not incorporate

366 into neuronal microtubule polymers (Figs. 3-4). We next assessed microtubule dynamics in

367 developing *Tuba1a$^{ND/+}$* neurons. Wild-type and *Tuba1a$^{ND/+}$ cortical* neurons were transfected

368 with GFP-MACF43, a fluorescently tagged protein containing the SxIP motif bound by

369 microtubule plus-end binding proteins (EBs), allowing for visualization of microtubule plus-end

370 polymerization in live cells (Fig. 5A)[42]. Kymograph plots generated from time-lapse GFP-

371 MACF43 images were analyzed for velocity, duration, and distance of GFP-MACF43

372 polymerization events (Fig. 5B). Previous analysis of *Tuba1a$^{ND/+}$* GFP-MACF43 data revealed a

373 significant reduction in the number of microtubule plus-ends per micron of neurite, which is

374 evident in the example kymographs (Fig. 5B; [24]). Kymograph analysis showed that

375 *Tuba1a$^{ND/+}$* microtubules had accelerated polymerization velocity (19.67±1.1µm/min) compared

376 to wild-type at DIV 1 (17.36±0.43 µm/min; Fig. 5C; n=688 events per genotype; p=0.0008 by

377 Mann-Whitney test), consistent with prior data from *S. cerevisiae* [22]. Additionally, *Tuba1a$^{ND/+}$*

378 polymerization events covered a larger distance (Fig. 5D; n=688 events, p<0.0001) and lasted a

379 longer amount of time (Fig. 5E; n=688 events, p<0.0001) than polymerization events in wild-

380 type neurons. The increased polymerization rates of *Tuba1a$^{ND/+}$* neuronal microtubules was

381 surprising, but collectively with our previous results these data indicate that *Tuba1aND* protein is

382 not functional, and that the diminished α-tubulin pool leads to altered microtubule dynamics and

383 changes to microtubule organization in developing neurons.

384

Figure 6 *Tuba1a^{ND} impairs neurite outgrowth and alters growth cone cytoskeleton in cortical neurons.*
 A. Schematic of experimental design and time-lapse images of membrane-labeled Myr-TdTomato neurons at DIV 1. Myr-TdTomato images were aquired 60 minutes apart to assess neuronal growth rate in wild-type (WT; top) and *Tuba1a^{ND/+}* (bottom) cortical neurons. **B.** Scatter plot of growth rates in live cortical neurons from WT and *Tuba1a^{ND/+}* mice at DIV 1. Data points represent individual neurons (N=3 mice, n=13 neurons; p=0.03 by Mann-Whitney test). **C.** Scatter plot of neurite length at DIV 3 in fixed WT and *Tuba1a^{ND/+}* primary cortical neurons. For each cell, the longest neurite, designated a putative 'axon', was measured from the cell soma to the distal neurite tip using an acetylated tubulin marker. Data points represent individual neurons (N=3 mice, n=124 neurons; p=0.02). **D.** Images of DIV 3 WT (top) and *Tuba1a^{ND/+}* (bottom) cortical neurons stained with antibodies directed against acetylated tubulin (green) and rhodamine phalloidin (F-actin, magenta). Composite images are shown (left) with single channel grayscale images for acetylated tubulin (middle) and F-actin (right). Scale bars are 10μm. **E.** Scatter plot of growth cone area at DIV 3 in WT and *Tuba1a^{ND/+}* cortical neurons. Data points represent individual growth cones (N=3 mice, n=52 growth cones; p=0.26). **F.** Scatter plot of acetylated tubulin intensity within the growth cone of WT and *Tuba1a^{ND/+}* cortical neurons at DIV 3. Data points represent individual growth cones (n=49 growth cones; p=0.89). **G.** F-actin intensity within growth cone of WT and *Tuba1a^{ND/+}* cortical neurons at DIV 3. Data points represent individual growth cones (N=3 mice, n=49 growth cones; p=0.0014). **H.** Representative images of WT and *Tuba1a^{ND/+}* cortical neuron growth cones showing distribution of acetylated tubulin (green) and F-actin (magenta). **I.** Line plot showing the average acetylated tubulin fluorescence intensity across a 20μm line scan through the central domain of the growth cone in DIV 3 WT and *Tuba1a^{ND/+}* neurons. Differences between WT and *Tuba1a^{ND/+}* growth cones were assessed by two-way ANOVA which showed a significant interaction between genotype and fluorescence intensity by position (n=20 growth cones; p<0.0001). **J.** Scatter plot representing the ratio of acetylated tubulin to F-actin intensity within the growth cones of DIV 3 WT and *Tuba1a^{ND/+}* cortical neurons. Data points represent individual growth cones (N=3 mice, n=49 growth cones; p=0.0003 by Mann-Whitney test). For all plots, lines represent mean and error bars report SEM. Differences between WT and *Tuba1a^{ND/+}* datasets were assessed by t test unless otherwise noted. *p<0.05; **p<0.01;

385 ***Tuba1a* is necessary for neurite extension and cytoskeletal organization in growth cone**

386   To assess potential mechanisms by which reduced Tuba1a prevents commissural neurons

387 from reaching their contralateral targets, we measured neurite growth rates in cultured primary

388 cortical neurons from P0 wild-type and *Tuba1a^{ND/+}* mice (Fig. 6A). We labeled primary wild-

389 type and *Tuba1a^{ND/+}* cortical neurons with a membrane-bound Myr-TdTomato to visualize

390 growth rates in live neurons (Fig. 6A). Analysis of membrane bound Myr-TdTomato images

391 taken one hour apart revealed that *Tuba1a^{ND/+}* neurons grew at a significantly slower rate than

392   wild-type neurons (Fig. 6B; N=3 mice, n=13 neurons; p=0.03 by Mann-Whitney test) with

393   *Tuba1a$^{ND/+}$* neurons growing an average of 4.67±1.15μm/hr compared to 12.60±3.12μm/hr in

394   wild-type neurons. Additionally, measurements of the longest neurite, designated a putative

395   'axon', at DIV 3 revealed that *Tuba1a$^{ND/+}$* neurites were significantly shorter than that of wild-

396   type (Fig. 6C; N=3 mice, n=124 neurons; p=0.02).Together, these data show that developing

397   neurons with reduced Tuba1a have shorter neurites and grow at a slower rate than wild-type.

398   To explore the precise mechanisms by which reduced Tuba1a contributes to slowed

399   outgrowth in *Tuba1a$^{ND/+}$* neurons, we assessed the abundance of acetylated microtubules and

400   filamentous actin (F-actin) in developing growth cones of wild-type and *Tuba1a$^{ND/+}$* cortical

401   neurons at DIV 3 (Fig. 6D). The growth cone is a dynamic developmental structure that uses the

402   coordinated action of the actin and microtubule cytoskeleton to drive neuronal outgrowth in

403   response to internal and external cues [43, 44]. Growth cone area was not changed in *Tuba1a$^{ND/+}$*

404   neurons (317.4±31.3μm^2) compared to wild-type (380.9±30.4μm^2; Fig. 6E; n=49 growth cones;

405   p=0.15). We examined the amount of acetylated tubulin, a PTM associated with stable

406   microtubules (Fig. 6F) and found there to be no difference in the overall fluorescence intensity of

407   acetylated tubulin in *Tuba1a$^{ND/+}$* neurons compared to wild-type (n=49 growth cones; p=0.89). In

408   contrast, we observed a significant increase in F-actin intensity within the growth cones of

409   *Tuba1a$^{ND/+}$* neurons compared to wild-type (Fig. 6G; n=49 growth cones; p=0.0014). Neuronal

410   microtubules splay out in the central, actin-dominated regions of the growth cone, but are

411   bundled towards the peripheral domains of the growth cone [45]. To assess the degree of growth

412   cone microtubule bundling, we next performed line scans across the widest point of DIV 3

413   growth cones ≥10μm (Fig. 6H). Line scans of acetylated tubulin through the growth cone

414   revealed differences in microtubule organization between wild-type and *Tuba1a$^{ND/+}$* neurons

415 (Fig. 6I; n=39 growth cones; p<0.0001 between genotypes by two-way ANOVA). Specifically,

416 we observed peaks in fluorescence, indicating bundled microtubules, at the edges of the growth

417 cone in wild-type neurons, where acetylated tubulin was more diffuse and lacked obvious

418 organization in *Tuba1a*^{ND/+} growth cones (Fig. 6I). Intriguingly, the ratio of acetylated

419 microtubules to F-actin in the growth cone was significantly reduced in *Tuba1a*^{ND/+} neurons

Figure 7 *Tuba1a*ND *neurons do not correctly localize Map1b to the developing growth cone.*
A. Western blots showing Map1b protein associate with a tubulin-enriched fraction from brain (top panel) and total Map1b protein in whole brain lysate (bottom panel) from wild-type (WT) and *Tuba1a*^{ND/+} mice. Due to the amount of protein that was loaded for Map1b western blots, antibody-stained bands for α-tubulin were oversaturated and could not be quantified, thus Map1b was normalized to the 50kDa band (presumed to be primarily tubulin) on a UV-activated stain-free blot. **B.** Scatter plot quantifying Map1b associated with the tubulin-enriched brain lysate, normalized to the 50kDa presumed tubulin band using stain-free western blotting. p=0.03 **C.** Scatter plot representing total Map1b protein in brain lysate by western blot, normalized to the total protein on a stain-free blot. p=0.98 **D.** Scatter plot showing the subcellular distribution of Map1b protein in WT and *Tuba1a*^{ND/+} cortical neurons at DIV 3. Data are represented as Map1b fluorescent signal in growth cone region divided by a region proximal to the cell body. p=0.009. **E.** Representative images showing altered subcellular distribution of Map1b in *Tuba1a*^{ND/+} (bottom) cortical neurons compared to WT (top) at DIV 3. Composite and individual channel grayscale images of MAP2 and Map1b immunocytochemistry are shown, *i* and *ii* indicate enlarged regions shown in insets. Scale bars are 10μm. Differences between groups were evaluated by t test. * p<0.05; ** p<0.01

compared to wild-type, indicating changes to the overall growth cone cytoskeletal environment

in $Tuba1a^{ND/+}$ neurons (Fig. 6J; n=49 growth cones per genotype; p=0.0003). Together, these

data indicate that neurite growth rate is particularly sensitive to the amount of Tuba1a tubulin

available, and impaired Tuba1a function leads to abnormal actin and microtubule architecture in

the developing growth cone.

$Tuba1a^{ND}$ neurons fail to localize critical developmental proteins to growth cone

MAPs play a crucial role in regulating neuronal microtubule function to support proper

neurodevelopment. Microtubule-associated protein 1b (Map1b) promotes axon extension and is

required for formation of the corpus callosum in mice [46, 47]. There was no significant deficit

in the association of Map1b with microtubules from $Tuba1a^{ND/+}$ brain lysates compared to wild-

type; in fact, $Tuba1a^{ND/+}$ lysates bound slightly more Map1b than wild-type (Fig. 7A, B; p=0.03).

Western blot analysis of whole brain lysates from wild-type and $Tuba1a^{ND/+}$ mouse brains

showed no difference in the total amount of Map1b protein (Fig. 7A,C; p=0.98). These data

indicate $Tuba1a^{ND}$ does not impair Map1b's interaction with neuronal microtubules. In

developing wild-type neurons, Map1b localizes strongly to the growth cone to promote axon

growth and facilitate microtubule response to guidance cues [47-51]. $Tuba1a^{ND/+}$ neurons

contained Map1b protein, but exhibited very little Map1b fluorescence in the growth cone

compared to wild-type neurons (Fig. 7D, E; n=31 growth cones; p=0.009). These data provide

evidence that while the abundance of Map1b protein is unchanged by $Tuba1a^{ND}$, reduced

Tuba1a does not allow for correct subcellular localization of Map1b to the growth cone. Failure

of $Tuba1a^{ND/+}$ neurons to localize Map1b to the developing growth cone provides a putative

442 mechanism by which developing axons may fail to respond to critical guidance cues, leading to

443 the commissural deficits observed in $Tuba1a^{ND/+}$ mice.

444

Figure 8 *Mechanisms of Tuba1a-induced axonal pathfinding deficits.*
A. Schematic illustrating how reduced Tuba1a may impair ability of $Tuba1a^{ND/+}$ axons to reach key signaling points at the correct developmental time to support proper brain formation. Wild-type axons (WT; top) reaching the midline crossing point of corpus callosum at embryonic day (E) 18.5 are compared to the potentially stunted axonal growth in $Tuba1a^{ND/+}$ brains (bottom). The reduced density of microtubules and altered polymerization dynamics likely contribute to slower growth of developing $Tuba1a^{ND/+}$ commissural neurons. ***B.*** Schematics illustrating potential molecular mechanisms by which Tuba1a supports navigating axons. ***i.*** Inset from *(A)*, showing WT growth cones navigating through midline corpus callosum. WT growth cones are rich with Map1b, which binds growth cone microtubules to guide cellular response to guidance molecules like Netrin. ***ii.*** Inset from *(A)* showing $Tuba1a^{ND/+}$ growth cones, which fail to localize Map1b and may be rendered unable to mount a cytoskeletal response to extracellular cues.

445   **Multi-faceted functions of TUBA1A during neurodevelopment**

446   Our results support two potential models by which TUBA1A regulates neuronal

447   microtubules to promote commissural axon pathfinding. Using our *TUBA1A*-His6 tool, we

448   illustrated that the *TUBA1A^{ND}* allele diminishes TUBA1A protein abundance in cells (Fig. 3).

449   Here, we demonstrate that diminished Tuba1a alters microtubule dynamics and impairs neurite

450   outgrowth *in vitro* (Figs. 5-6). Thus, reduced Tuba1a could render neurons incapable of

451   supplying the microtubule bulk needed to drive axon outgrowth forward at a specific rate (Fig.

452   8A). The timing of neurodevelopment is precisely regulated, and neurons which fail to reach

453   targets at the correct time can miss crucial developmental signaling events. We additionally show

454   that neurons with reduced Tuba1a fail to localize a critical developmental MAP, Map1b, to the

455   growth cone (Fig. 7). Interactions between MAPs and microtubules play a major role in adapting

456   microtubule function in response to a changing intra- and extra-cellular developmental

457   environment [46, 51, 52]. Therefore, neurons with reduced Tuba1a may be rendered unable to

458   respond to extracellular guidance cues, such as Netrin1, due to failed sub-cellular localization of

459   MAPs and other critical cargoes (Fig. 8B). Our evidence supports a multi-faceted role for

460   TUBA1A during neurodevelopment, where it tunes microtubule dynamics and density to fuel

461   growth and also provides stable tracks for rapid intracellular transport.

462

463   Discussion

464   **Studying α-tubulin isotypes *in vivo***

465   *TUBA1A* has long been associated with neurodevelopment due to its spatial and temporal

466   expression as well as its role in tubulinopathies [2-4, 17-20, 53-55], but it has been historically

467   difficult to study the contribution of a single α-tubulin isotype to microtubule network function

in vivo due to the limited availability of isotype-specific tools. Here, we present a novel tool for

studying TUBA1A protein *in vivo* without impacting native microtubule properties by

introducing a hexahistidine (His6) tag into a previously identified internal loop of TUBA1A (Fig.

3) [25-27]. In this study, we provide the first evidence that TUBA1A is essential for regulating

neuronal microtubule function to support long-distance targeting of central nervous system

axons. Ectopic expression of *TUBA1A*-His6 protein in cells revealed that *TUBA1AND* protein is

approximately half as abundant as wild-type (Fig. 3D), despite similar amounts of mRNA

expression in transfected cells (Fig. 3E). Based on this evidence, *TUBA1AND* substitution causes

targeted depletion of the mutant TUBA1A protein, but our results suggest that degradation by the

proteasome is unlikely (Fig. 3D). Newly synthesized α- and β-tubulin proteins enter a complex

tubulin folding pathway, where they interact with cytosolic chaperonins and tubulin-binding

cofactors to become folded and assembled into tubulin heterodimers [56]. As *TUBA1AND* protein

is diminished compared to wild-type, but is not proteasomally degraded, we predict that mutant

TUBA1AND protein may ineffectively cycle through the tubulin folding pathway. Introduction of

the ND substitution into the primary yeast α-tubulin, *Tub1* (*Tub1ND*) was previously shown to be

lethal when combined with tubulin folding pathway mutants, providing evidence that ND

substitution impairs tubulin assembly [22]. Further testing will be needed to evaluate the precise

molecular mechanisms responsible for reduced *TUBA1AND* protein in mammalian neurons, but it

is evident that Tuba1a function is lost in the *Tuba1aND* rodent model.

Neurons with reduced Tuba1a function exhibited accelerated microtubule polymerization

compared to wild-type, but also demonstrated deficits in neurite extension likely due to the

decreased axonal microtubule density (Figs. 5, 6; [24]). Additionally, Tuba1a-deficient

microtubules were not adequate to support growth cone localization of at least one critical

developmental MAP associated with commissural axon pathfinding, Map1b (Fig. 7). We show

that Tuba1a-rich microtubules promote axon outgrowth and pathfinding, and reduced Tuba1a

was not sufficient to form forebrain commissures in $Tuba1a^{ND/+}$ mice (Figs. 1-2). Collectively,

these data support the conclusion that reduced Tuba1a during neurodevelopment is adequate for

cortical neuron migration [22], but does not allow for sufficient microtubule function to properly

localize proteins to the growth cone for axon guidance. Thus, long range axon guidance may be

exquisitely sensitive to α-tubulin levels and microtubule structure.

498   Studying individual α-tubulin isotypes in neurons has been historically arduous as the

high degree of amino acid sequence similarity between α-tubulin isotypes has prevented

generation of a TUBA1A-specific antibody and has made genetically targeting a single α-tubulin

gene challenging. The abundance of clinically identified mutations to *TUBA1A* provide strong

evidence that *TUBA1A* is a major player in both tubulinopathy and typical neurodevelopment;

however, the lack of available tools to study TUBA1A *in vivo* has prevented researchers from

understanding precisely how *TUBA1A* contributes to neurodevelopment. As such, previous

studies of tubulinopathy mutations have relied heavily on mRNA analysis and indirect methods

of evaluating TUBA1A function. Here we introduce an important advancement in the study of

TUBA1A protein *in vivo*, by harnessing a previously-identified internal loop within TUBA1A

that tolerates addition of small epitope tags without impacting TUBA1A incorporation or

dynamics [26]. Our internal *TUBA1A-His6* construct marks an important advancement for the

study of tubulinopathies and neuronal α-tubulin as a whole. *TUBA1A-His6* was readily expressed

in both Cos-7 cells and neurons and was able to incorporate into microtubule polymers, unlike

TUBA1A containing a GFP fusion that prohibited incorporation into neuronal microtubules (Fig.

4). Additionally, we successfully used this epitope-tagged TUBA1A to model mutant tubulin

514 behavior *in vitro* (Figs. 3-4). Overall, this tool provides an important advancement in the study of

515 α-tubulin protein *in vivo*, and makes interrogating the function of specific α-tubulin isotypes

516 accessible to more researchers.

517

518 **Tuba1a influences microtubule density, dynamics, and function in cells**

519   The microtubule cytoskeleton supports a wide range of different cellular functions in

520 different cell types, ranging from facilitating chromosome segregation during mitosis to forming

521 dynamic and motile structures like the neuronal growth cone. Understanding how different cells

522 use the same basic building blocks to create vastly different microtubule-based structures is a

523 major question in microtubule biology. Many different mechanisms have been identified through

524 which cells can regulate microtubule network properties and overall function. We show that a

525 mutation which reduced Tuba1a incorporation into cellular microtubules accelerates microtubule

526 polymerization rates (Fig. 5). Alterations to tubulin isotype composition were previously shown

527 to change microtubule polymerization dynamics when the analogous mutation was made in

528 yeast, $Tub1^{ND}$. A similar acceleration of microtubule polymerization was observed in $Tub1^{ND}$

529 yeast mutants and was likely caused by a shift in the α-tubulin isotype ratio at the protein level

530 [22]. We propose that this polymerization rate increase is not due to a reduction of available

531 cellular tubulin, but instead is due to a change in the ratio of tubulin isotypes available for

532 microtubule growth. This result is also supported by recent evidence showing that increased

533 incorporation of Tuba1a tubulin, with subsequent decreased incorporation of alternative tubulin

534 isotypes, slowed microtubule polymerization rates in *in vitro* reconstituted microtubules [40].

535 These results are fitting with the "tubulin code" model, which proposes that incorporation of

536 different tubulin isotypes can modify microtubule network behavior [14, 57]. Importantly,

previous work has shown that local changes to growth cone microtubule dynamics facilitate

growth cone turning in response to extracellular cues [11, 13, 43, 45, 58, 59]. Thus,

dysregulation of growth cone microtubule dynamics, as was observed in $Tuba1a^{ND/+}$ cortical

neurons, could diminish the ability of developing neurons to appropriately interact with their

environment. Together, these data support the conclusion that incorporation of Tuba1a α-tubulin

tunes neuronal microtubule polymerization rates to support neurodevelopmental processes.

While the abundance of acetylated tubulin was not significantly different between

$Tuba1a^{ND/+}$ and wild-type growth cones, the distribution of acetylated microtubules was different

by genotype (Fig. 6). Differences in distribution of acetylated tubulin within the growth cone

likely reflects altered microtubule organization (Fig. 6). Acetylation is a microtubule PTM that is

associated with stable microtubule populations and as such is sparse in dynamic structures like

growth cones [60-65]. However, microtubule acetylation can be induced in growth cones

following contact with extracellular matrix proteins and was shown to promote cortical neuron

migration *in vivo* and suppress axon branching *in vitro*, demonstrating a clear role for this PTM

in development [66-68]. Tubulin PTMs, like acetylation, have been shown to impact MAP-

binding affinity and function, providing a clear mechanism by which changing the PTM

landscape of microtubules could alter neuronal microtubule function [41, 57, 65, 69-72]. Thus,

any changes to the organization or distribution of acetylated microtubules in the growth cone

could impact the ability of developing neurons to appropriately navigate their environment and

establish correct synaptic targets.

$Tuba1a^{ND/+}$ growth cones showed a significant increase in F-actin signal compared to

wild-type, causing an overall shift in the growth cone microtubule-actin balance (Fig. 6). It is

well established that interplay between the actin and microtubule cytoskeleton drives growth

cone movements in developing neurons [43, 44, 73, 74]. Growth cone microtubule

polymerization has been shown to induce F-actin assembly, and coordination of actin and

microtubules is regulated by interactions with MAPs to drive appropriate growth cone response

[75-78]. As actin and microtubules are tightly regulated within the growth cone, it is reasonable

to assume that mutations which disrupt microtubule function, like $Tuba1a^{ND}$, likely also impact

the actin cytoskeleton of developing neurons. In $Tuba1a^{ND/+}$ neurons, the actin cytoskeleton may

occupy increased growth cone territory as the result of microtubule deficiencies, but additional

testing of actin-response in developing $Tuba1a^{ND/+}$ neurons is needed to assess whether the

increase in growth cone actin has any functional consequences. Further, we showed that

$Tuba1a^{ND/+}$ neurons do not effectively localize at least one developmental MAP, Map1b, to the

growth cone (Fig. 7). Map1b acts downstream of several important developmental signaling

pathways to regulate function of both actin and microtubules within the growth cone, and

dysregulation of this or other MAPs could therefore impact multiple cytoskeletal components

[50, 51, 79]. The mechanisms by which $Tuba1a^{ND}$ induces changes to the growth cone actin

cytoskeleton remain to be explored, but could reveal important insights on how microtubules and

actin are coordinately regulated to support growth cone navigation.

Models of Tuba1a-dysfunction reveal potential roles in developmental signaling

$Tuba1a^{ND/+}$ neuronal microtubules were not sufficient to support growth cone localization

of Map1b (Fig. 7). Microtubules are the tracks upon which intracellular cargo transport occurs in

neurons. Here we present evidence that impaired Tuba1a function in neurons causes aberrant

localization of Map1b (Fig. 7). Map1b mRNA is a known target of the mRNA transport protein

FMRP and is locally translated within developing neurons [80, 81]. As we previously

583 demonstrated that intracellular transport is impaired in developing $Tuba1a^{ND/+}$ neurons [24], this

584 provides a putative model by which reduced Tuba1a could lead to altered localization of

585 developmental MAPs. Intracellular transport is a crucial function of neuronal microtubules

586 throughout life; however, microtubule-based transport has been shown to be essential during

587 neurodevelopment [12, 82-87]. Correct localization of developmental MAPs, mRNAs and

588 organelles are crucial for cytoskeletal response to extracellular guidance cues [84, 86, 88-90]. In

589 particular, Map1b is required for neuronal response to the guidance cue, Netrin1, a key player in

590 commissural formation [91-96]. We showed that neuronal microtubules with reduced Tuba1a do

591 not support neuronal growth to the same degree as wild-type microtubules, causing shorter

592 neurite length and slower growth rates *in vitro* (Fig. 6). The timing of developmental processes is

593 crucial for effective signal transduction and supports the formation of appropriate synaptic

594 contacts [97, 98]. Collectively, the data presented in this study support two potential models by

595 which $Tuba1a^{ND}$ neuronal microtubules fail to support proper neurodevelopment. The first model

596 proposes that the timing of axon extension during development is crucial for effective axon

597 guidance, as neurons with reduced Tuba1a exhibit impaired neurite extension (Fig. 6). If neurons

598 lacking TUBA1A are not reaching the correct location at the correct time, it is possible that

599 neurons will fail to receive key developmental signals (Fig. 8).The second model posits that

600 neurons lacking functional TUBA1A do not have adequate intracellular transport to support

601 localization of critical developmental proteins (Fig. 8). Inappropriate protein localization during

602 critical points in axon extension and guidance could render neurons unable to respond to

603 incoming guidance cues, as the machinery required to induce microtubule response to

604 extracellular cues is absent. Importantly, these models are not mutually exclusive, as we

605 demonstrated that TUBA1A is crucial for both developmental protein localization and neuron

606   outgrowth. The extent to which these processes contribute to the overall deficit in commissural

607   axon guidance remains to be explored in future studies.

608   Understanding the mechanisms by which microtubules contribute to discrete aspects of

neurodevelopment is an active area of research. Human neurodevelopmental disorders that

impact microtubule function, such as tubulinopathies, demonstrate that microtubules are critical

for proper neurodevelopment to occur. Tubulinopathy patients exhibit severe, sometimes lethal,

brain malformations that frequently impact multiple neurodevelopmental processes, including

neuronal survival, migration and axon extension [3, 6-8, 53, 99-101]. The range of phenotypes

exhibited by tubulinopathy patients have made it challenging for scientists to pinpoint specific

aspects of neuronal function that are reliant on TUBA1A tubulin. In this way, mutations such as

the *Tuba1aND* variant whose severity can be tuned according to gene dosage can be used as

important tools to interrogate the requirement for Tuba1a in discrete aspects of

neurodevelopment. Importantly, though tubulinopathy patients exhibit a range of brain

phenotypes, commissural abnormalities such as agenesis of the corpus callosum, are one of the

most commonly reported features of this disease [1-4, 20]. Cortical malformations and neuronal

migration errors are also common features of *TUBA1A* tubulinopathies; however, it has thus far

been unclear as to whether commissural deficits occur as a primary or secondary consequence of

TUBA1A dysfunction. In this study, we provide evidence that neurons deficient in Tuba1a fail to

properly navigate to meet contralateral binding partners. These data demonstrate that TUBA1A

is required for forebrain commissural formation, independent of its role in neuronal survival or

migration. The insights presented in this manuscript expand upon the currently known role for

TUBA1A in neurodevelopment, and advance the study of tubulinopathy by presenting specific

mechanisms by which TUBA1A supports neurodevelopment.

References:

1.   Cushion, T.D., et al., *Overlapping cortical malformations and mutations in TUBB2B and*
631   *TUBA1A*. Brain, 2013. 136(Pt 2): p. 536-48.

632   2.   Oegema, R., et al., *Recognizable cerebellar dysplasia associated with mutations in*
633   *multiple tubulin genes*. Hum Mol Genet, 2015. 24(18): p. 5313-25.

634   3.   Fallet-Bianco, C., et al., *Neuropathological phenotype of a distinct form of lissencephaly*
635   *associated with mutations in TUBA1A*. Brain, 2008. 131(Pt 9): p. 2304-20.

636   4.   Poirier, K., et al., *Large spectrum of lissencephaly and pachygyria phenotypes resulting*
637   *from de novo missense mutations in tubulin alpha 1A (TUBA1A)*. Hum Mutat, 2007.
638   28(11): p. 1055-64.

639   5.   Tischfield, M.A., et al., *Phenotypic spectrum of the tubulin-related disorders and*
640   *functional implications of disease-causing mutations*. Curr Opin Genet Dev, 2011. 21(3):
641   p. 286-94.

642   6.   Tian, G., et al., *Disease-associated mutations in TUBA1A result in a spectrum of defects*
643   *in the tubulin folding and heterodimer assembly pathway*. Hum Mol Genet, 2010. 19(18):
644   p. 3599-613.

645   7.   Tian, G., et al., *A pachygyria-causing alpha-tubulin mutation results in inefficient cycling*
646   *with CCT and a deficient interaction with TBCB*. Mol Biol Cell, 2008. 19(3): p. 1152-61.

647   8.   Aiken, J., J.K. Moore, and E.A. Bates, *TUBA1A mutations identified in lissencephaly*
648   *patients dominantly disrupt neuronal migration and impair dynein activity*. Hum Mol
649   Genet, 2019. 28(8): p. 1227-1243.

650   9.   Aiken, J., et al., *Tubulin mutations in brain development disorders: Why*
651   *haploinsufficiency does not explain TUBA1A tubulinopathies*. Cytoskeleton (Hoboken),
652   2019.

653   10.   Kapitein, L.C. and C.C. Hoogenraad, *Building the Neuronal Microtubule Cytoskeleton*.
654   Neuron, 2015. 87(3): p. 492-506.

655   11.   Liu, G. and T. Dwyer, *Microtubule dynamics in axon guidance*. Neurosci Bull, 2014.
656   30(4): p. 569-83.

12.   Miller, K.E. and D.M. Suter, *An Integrated Cytoskeletal Model of Neurite Outgrowth.*
Front Cell Neurosci, 2018. 12: p. 447.

13.   Sakakibara, A., et al., *Microtubule dynamics in neuronal morphogenesis.* Open Biol,
2013. 3(7): p. 130061.

14.   Gadadhar, S., et al., *The tubulin code at a glance.* J Cell Sci, 2017. 130(8): p. 1347-1353.

15.   Khodiyar, V.K., et al., *A revised nomenclature for the human and rodent alpha-tubulin*
gene family. Genomics, 2007. 90(2): p. 285-9.

16.   Findeisen, P., et al., *Six subgroups and extensive recent duplications characterize the*
evolution of the eukaryotic tubulin protein family. Genome Biol Evol, 2014. 6(9): p.
2274-88.

17.   Gloster, A., et al., *Early induction of Talpha1 alpha-tubulin transcription in neurons of*
the developing nervous system. J Comp Neurol, 1999. 405(1): p. 45-60.

18.   Gloster, A., et al., *The T alpha 1 alpha-tubulin promoter specifies gene expression as a*
function of neuronal growth and regeneration in transgenic mice. J Neurosci, 1994.
14(12): p. 7319-30.

19.   Miller, F.D., et al., *Isotypes of alpha-tubulin are differentially regulated during neuronal*
maturation. J Cell Biol, 1987. 105(6 Pt 2): p. 3065-73.

20.   Aiken, J., et al., *The alpha-Tubulin gene TUBA1A in Brain Development: A Key*
Ingredient in the Neuronal Isotype Blend. J Dev Biol, 2017. 5(3).

21.   Bittermann, E., et al., *Differential requirements of tubulin genes in mammalian forebrain*
development. PLoS genetics, 2019. 15(8): p. e1008243-e1008243.

22.   Gartz Hanson, M., et al., *Novel alpha-tubulin mutation disrupts neural development and*
tubulin proteostasis. Dev Biol, 2016. 409(2): p. 406-19.

23.   Keays, D.A., et al., *Mutations in alpha-tubulin cause abnormal neuronal migration in*
mice and lissencephaly in humans. Cell, 2007. 128(1): p. 45-57.

24.   Buscaglia, G., et al., *Reduced TUBA1A tubulin causes defects in trafficking and impaired*
683   *adult motor behavior.* eNeuro, 2020.

684   25.   Heilemann, M., et al., *Subdiffraction-resolution fluorescence imaging with conventional*
685   *fluorescent probes.* Angewandte Chemie (International ed. in English), 2008. 47(33): p.
686   6172-6176.

687   26.   Schatz, P.J., et al., *Insertions of up to 17 amino acids into a region of alpha-tubulin do*
688   *not disrupt function in vivo.* Molecular and cellular biology, 1987. 7(10): p. 3799-3805.

689   27.   Hotta, T., et al., *Affinity Purification and Characterization of Functional Tubulin from*
690   *Cell Suspension Cultures of Arabidopsis and Tobacco.* Plant physiology, 2016. 170(3): p.
691   1189-1205.

692   28.   Anders, K.R. and D. Botstein, *Dominant-lethal alpha-tubulin mutants defective in*
693   *microtubule depolymerization in yeast.* Mol Biol Cell, 2001. 12(12): p. 3973-86.

694   29.   Roostalu, J., et al., *The speed of GTP hydrolysis determines GTP cap size and controls*
695   *microtubule stability.* Elife, 2020. 9.

696   30.   Johnson, V., et al., *Design, overexpression, and purification of polymerization-blocked*
697   *yeast αβ-tubulin mutants.* Biochemistry, 2011. 50(40): p. 8636-44.

698   31.   Ōmura, S. and A. Crump, *Lactacystin: first-in-class proteasome inhibitor still excelling*
699   *and an exemplar for future antibiotic research.* The Journal of Antibiotics, 2019. 72(4):
700   p. 189-201.

701   32.   Tomoda, H. and S. Omura, *Lactacystin, a proteasome inhibitor: discovery and its*
702   *application in cell biology.* Yakugaku zasshi : Journal of the Pharmaceutical Society of
703   Japan, 2000. 120(10): p. 935-949.

704   33.   Vallee, R.B., *A taxol-dependent procedure for the isolation of microtubules and*
705   *microtubule-associated proteins (MAPs).* J Cell Biol, 1982. 92(2): p. 435-42.

706   34.   Nishikimi, M., K. Oishi, and K. Nakajima, *Axon guidance mechanisms for establishment*
707   *of callosal connections.* Neural Plast, 2013. 2013: p. 149060.

708   35.   Probst, M., *Uber den Bau des balkenlosen Grobhirns, sowie uber Mikrogyrie and*
709   *Heterotypie der grauer Substanz.* Arch F Psychiatr, 1901(34): p. 709-786.

710   36.   Kimble, M., et al., *Microtubule organization and the effects of GFP-tubulin expression in*
711   *dictyostelium discoideum.* Cell motility and the cytoskeleton, 2000. 47(1): p. 48-62.

712   37.   Zieve, G. and F. Solomon, *Direct isolation of neuronal microtubule skeletons.* Mol Cell
713   Biol, 1984. 4(2): p. 371-4.

714   38.   Wloga, D., E. Joachimiak, and H. Fabczak, *Tubulin Post-Translational Modifications*
715   *and Microtubule Dynamics.* Int J Mol Sci, 2017. 18(10).

716   39.   Zanic, M., *Measuring the Effects of Microtubule-Associated Proteins on Microtubule*
717   *Dynamics In Vitro.* Methods Mol Biol, 2016. 1413: p. 47-61.

718   40.   Vemu, A., et al., *Tubulin isoform composition tunes microtubule dynamics.* Mol Biol
719   Cell, 2017. 28(25): p. 3564-3572.

720   41.   Sirajuddin, M., L.M. Rice, and R.D. Vale, *Regulation of microtubule motors by tubulin*
721   *isotypes and post-translational modifications.* Nat Cell Biol, 2014. 16(4): p. 335-44.

722   42.   Honnappa, S., et al., *An EB1-Binding Motif Acts as a Microtubule Tip Localization*
723   *Signal.* Cell, 2009. 138(2): p. 366-376.

724   43.   Craig, E.M., *Model for Coordination of Microtubule and Actin Dynamics in Growth*
725   *Cone Turning.* Frontiers in cellular neuroscience, 2018. 12: p. 394-394.

726   44.   Dent, E.W., S.L. Gupton, and F.B. Gertler, *The growth cone cytoskeleton in axon*
727   *outgrowth and guidance.* Cold Spring Harb Perspect Biol, 2011. 3(3).

728   45.   Buck, K.B. and J.Q. Zheng, *Growth cone turning induced by direct local modification of*
729   *microtubule dynamics.* The Journal of neuroscience : the official journal of the Society
730   for Neuroscience, 2002. 22(21): p. 9358-9367.

731   46.   Meixner, A., et al., *MAP1B is required for axon guidance and Is involved in the*
732   *development of the central and peripheral nervous system.* J Cell Biol, 2000. 151(6): p.
733   1169-78.

47. Gonzalez-Billault, C., J. Avila, and A. Caceres, *Evidence for the role of MAP1B in axon*
formation. Mol Biol Cell, 2001. 12(7): p. 2087-98.

48. Black, M.M., T. Slaughter, and I. Fischer, *Microtubule-associated protein 1b (MAP1b) is*
concentrated in the distal region of growing axons. The Journal of neuroscience : the
official journal of the Society for Neuroscience, 1994. 14(2): p. 857-870.

49. González-Billault, C., et al., *Participation of structural microtubule-associated proteins*
(MAPs) in the development of neuronal polarity. Journal of neuroscience research, 2002.
67(6): p. 713-719.

50. Mack, T.G., M.P. Koester, and G.E. Pollerberg, *The microtubule-associated protein*
MAP1B is involved in local stabilization of turning growth cones. Molecular and cellular
neurosciences, 2000. 15(1): p. 51-65.

51. Tymanskyj, S.R., T.M. Scales, and P.R. Gordon-Weeks, *MAP1B enhances microtubule*
assembly rates and axon extension rates in developing neurons. Mol Cell Neurosci, 2012.
49(2): p. 110-9.

52. Del Rio, J.A., et al., *MAP1B is required for Netrin 1 signaling in neuronal migration and*
axonal guidance. Curr Biol, 2004. 14(10): p. 840-50.

53. Kumar, R.A., et al., *TUBA1A mutations cause wide spectrum lissencephaly (smooth*
brain) and suggest that multiple neuronal migration pathways converge on alpha
tubulins. Hum Mol Genet, 2010. 19(14): p. 2817-27.

54. Lecourtois, M., et al., *Human lissencephaly with cerebellar hypoplasia due to mutations*
in TUBA1A: expansion of the foetal neuropathological phenotype. Acta Neuropathol,
2010. 119(6): p. 779-89.

55. Bamji, S.X. and F.D. Miller, *Comparison of the expression of a T alpha 1:nlacZ*
transgene and T alpha 1 alpha-tubulin mRNA in the mature central nervous system. J
Comp Neurol, 1996. 374(1): p. 52-69.

56. Lewis, S.A., G. Tian, and N.J. Cowan, *The alpha- and beta-tubulin folding pathways.*
Trends Cell Biol, 1997. 7(12): p. 479-84.

57. Janke, C. and M. Kneussel, *Tubulin post-translational modifications: encoding functions*
on the neuronal microtubule cytoskeleton. Trends Neurosci, 2010. 33(8): p. 362-72.

763 58. Fassier, C., et al., *Motor axon navigation relies on Fidgetin-like 1-driven microtubule*
764   *plus end dynamics.* J Cell Biol, 2018. 217(5): p. 1719-1738.

765 59. Kiss, A., et al., *Neuronal Growth Cone Size-Dependent and -Independent Parameters of*
766   *Microtubule Polymerization.* Front Cell Neurosci, 2018. 12: p. 195.

767 60. Baas, P.W., et al., *Stability properties of neuronal microtubules.* Cytoskeleton
768   (Hoboken), 2016. 73(9): p. 442-60.

769 61. Eshun-Wilson, L., et al., *Effects of α-tubulin acetylation on microtubule structure and*
770   *stability.* Proceedings of the National Academy of Sciences of the United States of
771   America, 2019. 116(21): p. 10366-10371.

772 62. Palazzo, A., B. Ackerman, and G.G. Gundersen, *Cell biology: Tubulin acetylation and*
773   *cell motility.* Nature, 2003. 421(6920): p. 230.

774 63. Robson, S.J. and R.D. Burgoyne, *Differential localisation of tyrosinated, detyrosinated,*
775   *and acetylated alpha-tubulins in neurites and growth cones of dorsal root ganglion*
776   *neurons.* Cell motility and the cytoskeleton, 1989. 12(4): p. 273-282.

777 64. Schulze, E., et al., *Posttranslational modification and microtubule stability.* J Cell Biol,
778   1987. 105(5): p. 2167-77.

779 65. Mansfield, S.G. and P.R. Gordon-Weeks, *Dynamic post-translational modification of*
780   *tubulin in rat cerebral cortical neurons extending neurites in culture: effects of taxol.*
781   Journal of neurocytology, 1991. 20(8): p. 654-666.

782 66. Creppe, C., et al., *Elongator controls the migration and differentiation of cortical*
783   *neurons through acetylation of alpha-tubulin.* Cell, 2009. 136(3): p. 551-64.

784 67. Dan, W., et al., *α-Tubulin Acetylation Restricts Axon Overbranching by Dampening*
785   *Microtubule Plus-End Dynamics in Neurons.* Cerebral cortex (New York, N.Y. : 1991),
786   2018. 28(9): p. 3332-3346.

787 68. Bergstrom, R.A., R.C. Sinjoanu, and A. Ferreira, *Agrin induced morphological and*
788   *structural changes in growth cones of cultured hippocampal neurons.* Neuroscience,
789   2007. 149(3): p. 527-536.

69. Balabanian, L., C.L. Berger, and A.G. Hendricks, *Acetylated Microtubules Are*
Preferentially Bundled Leading to Enhanced Kinesin-1 Motility. Biophysical journal,
2017. 113(7): p. 1551-1560.

70. Bonnet, C., et al., *Differential binding regulation of microtubule-associated proteins*
MAP1A, MAP1B, and MAP2 by tubulin polyglutamylation. J Biol Chem, 2001. 276(16):
p. 12839-48.

71. Lacroix, B., et al., *Tubulin polyglutamylation stimulates spastin-mediated microtubule*
severing. J Cell Biol, 2010. 189(6): p. 945-54.

72. Reed, N.A., et al., *Microtubule acetylation promotes kinesin-1 binding and transport.*
Curr Biol, 2006. 16(21): p. 2166-72.

73. Dent, E.W. and K. Kalil, *Axon branching requires interactions between dynamic*
microtubules and actin filaments. J Neurosci, 2001. 21(24): p. 9757-69.

74. Piper, M., et al., *Differential requirement of F-actin and microtubule cytoskeleton in cue-*
induced local protein synthesis in axonal growth cones. Neural Dev, 2015. 10: p. 3.

75. Rochlin, M.W., M.E. Dailey, and P.C. Bridgman, *Polymerizing microtubules activate*
site-directed F-actin assembly in nerve growth cones. Molecular biology of the cell,
1999. 10(7): p. 2309-2327.

76. Biswas, S. and K. Kalil, *The Microtubule-Associated Protein Tau Mediates the*
Organization of Microtubules and Their Dynamic Exploration of Actin-Rich
Lamellipodia and Filopodia of Cortical Growth Cones. J Neurosci, 2018. 38(2): p. 291-
307.

77. Slater, P.G., et al., *XMAP215 promotes microtubule-F-actin interactions to regulate*
growth cone microtubules during axon guidance in Xenopus laevis. Journal of cell
science, 2019. 132(9): p. jcs224311.

78. Szikora, S., et al., *The formin DAAM is required for coordination of the actin and*
microtubule cytoskeleton in axonal growth cones. Journal of cell science, 2017. 130(15):
p. 2506-2519.

79. Noiges, R., et al., *Microtubule-associated protein 1A (MAP1A) and MAP1B: light chains*
determine distinct functional properties. J Neurosci, 2002. 22(6): p. 2106-14.

819 80. Lu, R., et al., *The fragile X protein controls microtubule-associated protein 1B*
820 *translation and microtubule stability in brain neuron development.* Proc Natl Acad Sci U
821 S A, 2004. 101(42): p. 15201-6.

822 81. Wang, B., et al., *FMRP-Mediated Axonal Delivery of miR-181d Regulates Axon*
823 *Elongation by Locally Targeting Map1b and Calm1.* Cell Rep, 2015. 13(12): p. 2794-
824 807.

825 82. Baraban, M., et al., *Zebrafish embryonic neurons transport messenger RNA to axons and*
826 *growth cones in vivo.* J Neurosci, 2013. 33(40): p. 15726-34.

827 83. Dent, E.W. and P.W. Baas, *Microtubules in neurons as information carriers.* J
828 Neurochem, 2014. 129(2): p. 235-9.

829 84. Feltrin, D., et al., *Growth cone MKK7 mRNA targeting regulates MAP1b-dependent*
830 *microtubule bundling to control neurite elongation.* PLoS Biol, 2012. 10(12): p.
831 e1001439.

832 85. Hoogenraad, C.C. and F. Bradke, *Control of neuronal polarity and plasticity--a*
833 *renaissance for microtubules?* Trends Cell Biol, 2009. 19(12): p. 669-76.

834 86. Leung, K.M., et al., *Cue-Polarized Transport of beta-actin mRNA Depends on 3'UTR and*
835 *Microtubules in Live Growth Cones.* Front Cell Neurosci, 2018. 12: p. 300.

836 87. Liu, J.S., et al., *Molecular basis for specific regulation of neuronal kinesin-3 motors by*
837 *doublecortin family proteins.* Mol Cell, 2012. 47(5): p. 707-21.

838 88. Pilaz, L.J., et al., *Dynamic mRNA Transport and Local Translation in Radial Glial*
839 *Progenitors of the Developing Brain.* Curr Biol, 2016. 26(24): p. 3383-3392.

840 89. Qu, C., et al., *Direct binding of TUBB3 with DCC couples netrin-1 signaling to*
841 *intracellular microtubule dynamics in axon outgrowth and guidance.* J Cell Sci, 2013.
842 126(Pt 14): p. 3070-81.

843 90. Welshhans, K. and G.J. Bassell, *Netrin-1-induced local beta-actin synthesis and growth*
844 *cone guidance requires zipcode binding protein 1.* J Neurosci, 2011. 31(27): p. 9800-13.

845   91.   Arbeille, E. and G.J. Bashaw, *Brain Tumor promotes axon growth across the midline*
846   *through interactions with the microtubule stabilizing protein Apc2.* PLoS Genet, 2018.
847   14(4): p. e1007314.

848   92.   Barallobre, M.J., et al., *Aberrant development of hippocampal circuits and altered neural*
849   *activity in netrin 1-deficient mice.* Development, 2000. 127(22): p. 4797-810.

850   93.   Finger, J.H., et al., *The netrin 1 receptors Unc5h3 and Dcc are necessary at multiple*
851   *choice points for the guidance of corticospinal tract axons.* J Neurosci, 2002. 22(23): p.
852   10346-56.

853   94.   Fothergill, T., et al., *Netrin-DCC signaling regulates corpus callosum formation through*
854   *attraction of pioneering axons and by modulating Slit2-mediated repulsion.* Cereb
855   Cortex, 2014. 24(5): p. 1138-51.

856   95.   Lindwall, C., T. Fothergill, and L.J. Richards, *Commissure formation in the mammalian*
857   *forebrain.* Curr Opin Neurobiol, 2007. 17(1): p. 3-14.

858   96.   Serafini, T., et al., *Netrin-1 is required for commissural axon guidance in the developing*
859   *vertebrate nervous system.* Cell, 1996. 87(6): p. 1001-14.

860   97.   Krol, A. and G. Feng, *Windows of opportunity: timing in neurodevelopmental disorders.*
861   Current opinion in neurobiology, 2018. 48: p. 59-63.

862   98.   Rossi, A.M., V.M. Fernandes, and C. Desplan, *Timing temporal transitions during brain*
863   *development.* Current opinion in neurobiology, 2017. 42: p. 84-92.

864   99.   Bamba, Y., et al., *In vitro characterization of neurite extension using induced pluripotent*
865   *stem cells derived from lissencephaly patients with TUBA1A missense mutations.* Mol
866   Brain, 2016. 9(1): p. 70.

867   100.   Belvindrah, R., et al., *Mutation of the alpha-tubulin Tuba1a leads to straighter*
868   *microtubules and perturbs neuronal migration.* J Cell Biol, 2017.

869   101.   Keays, D.A., et al., *The role of Tuba1a in adult hippocampal neurogenesis and the*
870   *formation of the dentate gyrus.* Dev Neurosci, 2010. 32(4): p. 268-77.

871