Title: Resolving spatial complexities of hybridization in the context of the gray zone of

speciation in North American ratsnakes (*Pantherophis obsoletus* complex)

Frank T. Burbrink[1*]

Marcelo Gehara[1,2]

Edward A. Myers[1,3]

[1]Department of Herpetology, The American Museum of Natural History, Central Park West and

79[th] Street, New York, NY 10024 USA

[2]Department of Biological Sciences, Rutgers University Newark, 195 University Ave,

Newark, NJ 07102

[3]Department of Vertebrate Zoology, National Museum of Natural History, Smithsonian

Institution, Washington, DC, USA

*Corresponding author – fburbrink@amnh.org

Abstract:

Estimating species divergence with gene flow has been crucial for characterizing the gray

zone of speciation, which is the period of time where lineages have diverged but have not yet

achieved strict reproductive isolation. However, estimates of divergence times and gene flow

often ignores spatial information, for example the formation and shape of hybrid zones. Using

population genomic data from the eastern ratsnake complex (*Pantherophis obsoletus*), we infer

phylogeographic groups, gene flow, changes in demography, the timing of divergence, and

hybrid zone widths. We examine the spatial context of diversification by linking migration and

timing of divergence to the size, shape, and types of hybridization (e.g., F1, backcrosses) in

hybrid zones. Rates of migration between lineages are associated with the width and shape of

hybrid zones. Timing of divergence is not related to migration rate across species pairs and is

therefore a poor proxy for inferring position in the gray zone. Artificial neural network

approaches are applied to understand how landscape features and past climate have influenced

population genetic structure among these lineages prior to hybridization. The Mississippi River

produced the deepest divergence in this complex, whereas Pleistocene climate and elevation

secondarily structured lineages.

Key Words: Eastern Nearctic, migration, isolation, neural networks, reproductive isolation, cline,

hybrid zone

Wide-ranging species complexes that cross numerous biogeographic barriers provide

opportunities to better understand how changing landscapes affects diversification, gene flow,

demography, and the formation of hybrid zones. Lineages within species complexes ranging

across heterogeneous landscapes are likely at different stages of the speciation process reflecting

how specific environmental and biogeographic barriers have uniquely altered changes in gene

flow and other demographic processes in a particular complex (Myers et al. 2020). Even at the

same barrier within related groups of organisms, the timing of divergence and degree of gene

flow can be species specific and dependent on when they encountered the barrier (Riddle 2016;

Myers et al. 2019). Alternatively, generalist species not as constrained to specific habitats may

not show a correlation between genetic variation and landscape features (Joseph and Wilke 2007;

Makowsky et al. 2009; Lourenço et al. 2017).

Lineage divergence can be placed in the context of the gray zone of speciation, which

defines the range of time where speciation proceeds from early population differentiation with

unfettered gene flow to nearly reproductive isolation defined by low rates of gene flow (de

Queiroz, 2007; Hewitt, 2008; Jackson et al., 2016; Roux et al., 2016). The ability to delimit

species therefore may be correlated with their position in the gray zone, which has previously

been assessed by relating measures of genetic isolation and migration (the genealogical

divergence index; GDI) with species-delimitation probabilities (Jackson et al. 2016; Leaché et al.

2019). Moreover, given a model of speciation that proceeds by Dobzhansky-Muller

incompatibilities and genetic drift alone, the degree of gene flow might be predicted by the age

of lineage formation relative to population size and thus position within the gray zone (Orr 1995;

Gavrilets 2004; Singhal and Moritz 2013). Clearly some bias exists here; enhanced gene flow

between the youngest divergences may never proceed to the later stages in the gray zone as they

may rapidly collapse into a single lineage (Garrick et al. 2019), such as in the case where

physical barriers to gene flow disappear. On the other hand, strong selection may prevent young

lineages from exchanging alleles at particular loci, shortening the time in the gray zone (Mayr

1963; Barton 2010; Feder et al. 2012; Roux et al. 2016; Edwards et al. 2020). Therefore, finding

a threshold that can determine if two species are unique given only generations since divergence

may fail in cases where rates of gene flow between lineages drastically change over time and

space (Gourbière and Mallet 2010; Nosil et al. 2017).

Conceptualizing the gray zone without a spatial demographic component therefore may

be inadequate for understanding speciation and delimiting species. Isolation-migration models

(Hey and Nielsen 2004; Hey 2010) are coalescent estimators of historical processes and can

assess gene flow throughout time, reflecting position in the gray zone of speciation. These

models, however, lack any spatial component necessary to understand the underlying geographic

and environmental processes influencing migration. Studies of hybrid zones on the other hand

can clarify the location of migration and rates of contemporary gene flow, provided selection

does not eliminate F1 hybrids or backcrossing. If species pairs rapidly adapt to unique habitats

across their distribution, then age of divergence and degree of gene flow may be unrelated

relative to neutral processes when compared across taxa (Nosil and Crespi 2006; Agrawal et al.

2011; Nosil 2012; Karrenberg et al. 2019). Therefore, divergence over landscape can show how

selection at a barrier can be discordant from the timing of origin of the lineages (Barton and

Hewitt 1985; Harrison 1993; Jiggins and Mallet 2000; Gay et al. 2008; Seehausen et al. 2014;

Stankowski et al. 2017). Combining spatial information with genetic data will better help

understand how genetic variation is structured by geographic distance, environment, and the

presence of hybrid zones. This should provide a clearer understanding of why lineages maintain

particular rates of gene flow along a biogeographic barrier that has resulted in a hybrid zone.

Moreover, investigating gene flow in space determines if lineages were generated via isolation

and migration or are simply artificially partitioned groups with continuous gene flow over the

landscape defined by IBD (Wright 1943; Frantz et al. 2010; Bradburd et al. 2018).

To understand the factors shaping reproductive isolation in a species complex, we ask

whether divergence time predicts rates of gene flow or cline width across space in eastern

ratsnakes (*Pantherophis obsoletus* complex). This complex is a wide-ranging group of four

species found throughout the forested regions of the Eastern Nearctic and nearby Chihuahuan

Desert that have diverged at three unique biogeographic barriers. The primary divergence in this

complex occurred at the Mississippi River, which has consistently been identified as one of the

main biogeographic barriers in the ENA (Robison 1986; Burbrink et al. 2000; Soltis et al. 2006;

Brandley et al. 2010; Zellmer et al. 2012; Myers et al. 2020). Further east, the Appalachian

Mountains and Apalachicola/Chattahoochee River System (AARS) have noted barrier effects in

other organisms and likely contributed to the divergence between *P. alleghaniensis* and *P.*

spiloides (Walker and Avise 1998; Burbrink et al. 2000; Soltis et al. 2006). West of the

Mississippi River, divergences occurred at the transition between temperate forests to the rocky

areas on the western edge of the Edwards Plateau into the Chihuahuan Desert and isolated the

species *P. obsoletus* and *P. bairdi* (Fig.1; Lawson and Lieb 1990; Burbrink et al. 2000; Burbrink

2001). Since the Pliocene, the ENA has experienced numerous climate change events associated

with glacial cycles that forced species into refugia and compressed populations; upon climate

amelioration populations expanded into formerly glaciated areas (Hewitt 2000; Bintanja and van

de Wal 2008; Burbrink et al. 2016). Using dense population sampling and genomic-scale data,

we combine isolation-migration estimates with spatial information to understand species

divergence in the context of geographic distance, contemporary and historical environments, and

biogeographic boundaries. We also examine how migration rates and divergence dates relate to

the width and types of hybrids (F1, backcrosses) in these hybrid zones at biogeographic

boundaries. We also investigate demographic changes through time to understand if population

expansion occurs among all lineages as predicted given glacial cycling. Addressing these

questions provides an integrated view of phylogeographic history over a physically and

historically complex landscape and yields a clearer understanding of speciation processes and

delimitation in the gray zone.

Methods

Dataset

We sampled 288 individuals liberally covering the range of all species within the

Pantherophis obsoletus complex (Fig.1; Dryad XXX). DNA was extracted from all samples

using Qiagen DNeasy Blood & Tissue Kits and samples were screened for quality using broad-

range Qubit Assays. We used services from RAPiD Genomics (https://www.rapid-

genomics.com/services/) to generate 5472 baits and to sequence 5060 conserved elements

(UCEs) loci following the protocols from (Faircloth et al. 2012) and (Sun et al. 2014). These

markers have been used to address phylogeographic/population genetic and deeper phylogenetic

questions (Branstetter and Longino 2019; Younger et al. 2019). We mapped UCE reads to a

Chromium 10x *Pantherophis spiloides* genome (in prep), removed loci containing >50% missing

samples, and removed individual specimens missing >30% of all alignments (details on the

assembly of UCE loci are available in Supporting Information Material 1). We produced both

phased locus datasets, used in PHRAPL and PipeMaster, and single SNPs/locus for all other

analyses, furthermore we filtered the unlinked SNP dataset to remove alleles with a minor allele

frequency <0.1 and used these data for population structuring analyses.

Geographic groupings

We estimated population structure by comparing Discriminant Analysis of Principal

Components (DAPC; Jombart et al., 2010) in adegenet v2.1.2 (Jombart 2008). We compared this

to estimated effective migration surfaces (EEMS v; Petkova et al., 2016), which models effective

migration rates over geography to represent regions where migration is low in cases where

genetic dissimilarity increases rapidly, thus providing a uniquely distinct view of the location of

population clusters relative to biogeographic barriers as compared to DAPC (details on these

population grouping methods and other methods for assessing structure are available in

Supporting Information Material 1). We also examined fixation indices (F_{ST}) estimated from Nei

(1973) in adegenet v2.1.2 (Jombart 2008) across all loci comparing spatially adjacent lineage

pairs from DAPC .

Isolation, migration, and historical demography

We generated a coalescent-based species tree with SNAPP v1.3.0 (Bryant et al. 2012) to

understand relationships among the four taxa identified from clustering analyses and test species

delimitation. Because generating a tree in SNAPP using all individuals was computationally

intractable, we used four individuals per taxon that were sampled from different regions of the

taxon's distribution. We estimated SNAPP trees using BEAST v2.5.2 (Bouckaert et al. 2014;

details on parameter setting for SNAPP are available in Supporting Information Material 1). We

assessed four alternative species delimitation models including four taxa, three taxa (collapsing

P. bairdi/P. obsoletus or *P. alleghaniensis/P. spiloides*)*,* and two taxa (collapsing *P. bairdi/P.*

obsoletus and *P.alleghaniensis/P. spiloides).*

Using the four inferred lineages, we examined which models best described the origins of

these groups given divergence time, historical demographic change, and migration between

spatially adjacent taxa. We tested 2,300 candidate isolation-migration models including all

possible topologies with or without migration between all spatially adjacent pairs using the

program PHRAPL (Jackson et al. 2016) implemented in R. To further test that four genetic

lineages exist, we ran pairwise comparisons between the best selected model and models of three

populations where sister species were collapsed into a single entity; four *Pantherophis* species

against one model where *P. obsoletus* and *P. bairdi* were collapsed, and another where *P.*

spiloides and *P. alleghaniensis* were collapsed (details on PHRAPL runs are available in

Supporting Information Material 1).

To estimate gene flow, timing of divergence, and demographic change, we used

PipeMaster (Gehara et al., 2017; Gehara et al. in review) to simulate genetic data and perform

approximate Bayesian computation (ABC) and supervised machine-learning. Here we used the

best model selected in the PHRAPL analysis as a template (same topology and migration

parameters) to generate three competing models: (i) an isolation migration model with constant

population size for each lineage and constant migration, (ii) demographic change along each

lineage and constant migration through time and demographic change, and (iii) population size

change with migration occurring after the Last Glacial Maximum. We simulated 100,000 data

sets of 54 summary statistics and performed ABC rejection with 0.01 tolerance level to select the

best of these three models (Table S1). We then took the selected model and performed rejection

using a 0.1 tolerance level with a neural network regression using a single layer and 20 nodes to

estimate the model parameters.

Spatial population genetics

We used artificial neural networks (ANN; Lek and Guégan 1999; Legendre and Fortin

2010; Legendre et al. 2011) to understand how contemporary and historical aspects of the

landscape and climate generated genetic diversity. We first compared niche models of current

climates between adjacent species pairs to determine if contemporary niches were significantly

different prior to using these variables for understanding their effects on genetic structure (see

Supporting Information Material 1 for details). We then used Nei's distance (Nei, 1972) to

estimate genetic distance among all individuals in the R package adegenet. For predictor

variables, we used 1) geographic distances measured as pairwise great-circle distances among all

points to account for isolation-by-distance using the R package fossil (Vavrek 2020), 2) binary

categorization of isolation east and west of the Mississippi River, 3) elevation, and 4) bioclimatic

variables at 2.5 mins representing current climate, Last Glacial Maximum (21kya), Last

Interglacial (130 kya), Pleistocene Marine Isotope Stage 19 (787 kya) and Mid-Pliocene

Warming (3.2 Ma) all obtained from the PaleoClim database (Brown et al. 2018). For the

bioclimatic data and elevation, we extracted parameter values given the latitude and longitude for

each individual sample. To reduce dimensionality of the bioclimatic variables, we transformed

the former using principal components analyses using the R package raster (Hijmans et al. 2014).

Spatial distances were transformed using principal coordinates analyses in the R package ape

(Paradis and Schliep 2019). For principal coordinate/component analyses we retained axes that

account for >95% of variance.

We used regression-based ANN to determine which variables predict genetic distances

and rank those with the highest model importance. Machine learning methods can infer non-

linear interactions among many predictor variables, regardless of the distribution or variable type

and can generate a range of models of varying complexity (Lek et al. 1996; Zhang 2010;

Libbrecht and Noble 2015; Sheehan et al. 2016). To conduct the ANN regressions we used the R

package *caret* (Kuhn 2008) and partitioned the data into the standard 70% training and 30% test

sets (Lek et al. 1996; Zhang 2010; Burbrink et al. 2020). This analysis was run using 1,000

maximum iterations to ensure convergence. We resampled the data using the default 25 bootstrap

replicates to reach convergence over the following parameters: weight decay, root mean squared

error, r^2, and mean absolute error. We assessed the power of these models by recomposing the

test and training sets 100 times and compared three test statistics (root mean squared error, r^2,

and mean absolute error) to those from randomized response variables for each of these 100 re-

estimated models. We ranked the most important variables over these replicates after accounting

for multicollinearity among variable affects using the R package Rnalytica (see Supporting

Information Material 1). We also compared results from our ANN analysis to those using

redundancy analyses (RDA, see Supporting Information Material 1 for details).

Hybrid-zone dynamics

To understand if the composition and width of hybrid-zones differed among adjacent

lineages we first estimated the composition of individuals and their locations (parentals, F1s,

backcrosses). We first used *snapclust* (Beugin et al. 2018) in adegenet to assign pure parentals

(P1 and P2), backcrosses and F1s between the following spatially adjacent pairs of taxa, herein

referred to by geographic area: 1) eastern (*P. allegahniensis/P. spiloides)*, 2) central (*P.*

spiloides/P. obsoletus), and 3) western (*P. obsoletus/P. bairdi)*. This method maximizes the

likelihood of two fixed panmictic populations using a geometric approach with an expectation-

maximization algorithm. We estimated group membership first using the "k-means" option in the

function. To check the probability that *snapclust* identified hybrids correctly with these data, we

isolated pure parental individuals (membership >0.95), used the *hybridize* function to generate

the same number of hybrids and parentals and then determined the probability that *snapclust*

finds the correct number of hybrids and parentals.

We then fit clines in the program HZAR v.2-9 (Derryberry et al. 2014) for the same

species pairs using individual genetic assignments from DAPC to determine if cline widths differ

among groups. We compared these to ancestry alternatively estimated in *snaplcust* (see

Supporting Information Material 1 for details). Using the Gaussian cline model, we estimated the

center and width of the cline and determined if these sigmoidal distributions have significant tails

by fitting the following models: 1) no tails, 2) right tail only, 3) left tail only, 4) mirrored tails,

and 5) both tails estimated independently (see Derryberry et al. 2014). We fit these models to our

data using AICc and ran the MCMC chains for 5×10^6 generations, thinned by 5×10^3 generations,

and estimated stationarity using ESS >200 in the R package CODA (Plummer et al. 2006). We

note that making transects through multiple locations along each hybrid zone would better define

the width throughout the length of the cline, though this was not possible here for all clines.

Using equations from Barton and Gale (1993) and applied in Bailey et al. (2015) we

addressed selection against hybridization for each cline. Using estimates of maximum lifetime

dispersal (σ) for ratsnakes, here 0.937-4.3 km (Weatherhead and Blouin-Demers pers comm.),

we predict the width of the cline under neutrality over a range of generation times (*T*) since the

origin of each sister pair and group using the equation: $w = 2.51\sigma\sqrt{T}$. We also predict selection

at the center of the cline, also using σ and estimates of cline width (*w*) from HZAR using the

equation:

$S = \left(\frac{2\sigma}{w}\right)^2$. We estimate these given the range of dispersal estimates and estimates for the widths

of clines.

All scripts, genetic, and spatial data are available on Dryad XX.

Results

Data and population genetic structure

After filtering loci for presence in 70% or more individuals, we generated 2,491 UCE

loci. For all downstream analyses we randomly sampled one SNP per locus yielding a total of

846 SNPs for 238 individuals for an average of 13.96% missing data.

Both EEMS and DAPC find population structure that generally matched the geographic

ranges of *P. alleghaniensis*, *P. spiloides, P. obsoletus*, and *P. bairdi* (Fig. 1 and Fig. S1;

Burbrink, 2001). All three EEMS generated similar acceptance proportions for all proposal types

(12-51%). Total DAPC assignment probabilities without apriori species groupings were 0.975

(*P. bairdi* = 1.0, *P. obsoletus* = 0.96; *P. spiloides* = 1.0, and *P. alleghaniensis* = 0.97). Similarly,

all three EEMS runs suggest low migration at the MR and the area separating the *P. bairdi* and

P. obsoletus in west Texas. East of the MR, estimated low migration occurred near the

Appalachian Mountains, though this area is a complex mixture of isolation and gene flow (Fig.

1).

These estimates of population structure all showed *P. alleghaniensis* as occurring mostly

in the Florida Peninsula and north along the east coast to Virginia/Delaware (not as far north as

in Burbrink 2001), *P. spiloides* is found from western Florida to the Mississippi River up through

the Midwest and east to the northeastern US, *P. obsoletus* is found west of the Mississippi River

in the forested regions of the Midwest, and *P. bairdi* is distributed in the Chihuahuan Desert of

west Texas (Figs.1-2).

Isolation-migration processes

Our SNAPP analyses produced a phylogeny with a root separating the taxa east and west

of the Mississippi River and then a sister relationship between *P. bairdi* and *P. obsoletus* and

then *P. spiloides* and *P. alleghaniensis* (Fig. 2), consistent with Burbrink et al., (2000). Bayes

factors (BF) showed the four-taxon model as decisively superior to the three-taxon (BF =35.40)

and two-taxon model (BF=2022.46).

Using those four lineages we filtered models of divergence using PHRAPL. Models with

all four taxa, incorporating migration between spatially adjacent lineages was preferred with the

same topology as found using SNAPP. ΔAIC between the best ranked model and next model was

> 20, suggesting high confidence in model selection (Table S2).

The demographic model estimated from PipeMaster that incorporated historical

population size change and constant migration (IMD) best fit the data (posterior probability 1.0;

see PCA plots in Fig. S2; Figs. 2 & 3). Median migration rates were highest between eastern

groups (*P. alleghaniensis* to *P. spiloides* = 2.79 and *P. spiloides* to *P. alleghaniensis* = 3.79

individuals/generation), lower in the central groups across the Mississippi River (*P. spiloides* to

P. obsoletus = 1.96 and *P. obsoletus* to *P. spiloides* = 1.28 individuals/generation), and lowest in

western groups (*P. bairdi* to *P. obsoletus*= 0.55 and *P. obsoletus* to *P. bairdi* = 0.65

individuals/generations; Table 1). Divergence occurred earliest in the central lineages at the

Mississippi River at 1.08 My (95% CI =0.553 to 1.674 My), then the eastern lineages at 0.650

My (95%CI = 0.225 to 0.983 My) and western lineages at 0.367 My (95% CI 0.367 to 0.875 My;

Table 1). Divergence dates and average rates of migration were poorly correlated (ρ=0.264,

P=0.83, n=3). For all taxa we inferred population expansion from ancestral N_e sizes ranging from

5,200 to 5,600 to modern sizes at 138,000 to 281,000. The posterior distributions for all

parameters were different from the uniform priors, though wide distributions of several

parameters suggested some uncertainty in estimates (Fig. 3).

Spatial population genetics and hybrid zones

We used ANN to determine what features of the landscape and environmental layers

through time best predict genetic structure (Fig.4). Contemporary niches between taxa were

significantly different for all taxa (see Supporting Information Material 1 for details, Fig. S3).

Estimates of accuracy using ANN were high (>90%) for all comparisons. Most genetic structure

can be predicted by the Mississippi River showing 100% variable importance (Table S3).

Distance also played a role in structuring these data (although they were partialed out in RDA

analyses yielding similar conclusions regarding other landscape variables, see Supporting

Information Material 1). For those lineages east of the Mississippi River, population structure

was predicted by climate at the Last Glacial Maxima and elevation (variable importance=100).

For those lineages west of the Mississippi River, climate during Pleistocene Marine Isotope

Stage 19 and elevation structured these lineages (variable importance = 100).

Using *snapclust,* we showed that the number of hybrids of all types decreases with

species pair comparisons from east to west (Fig. 5). The parentals, hybrids and backcrosses were

predicted with a mean accuracy of 0.99 in simulations. All hybrid types were spatially oriented

between parental types where the number of hybrids and backcrosses between the eastern groups

approached the number of parental individuals (49%), for the central groups the hybrids and

backcrosses to parental ratio was 5.9%, and for the western groups there were no hybrids

detected (Fig. 5). It is important to note, however, that sampling was more concentrated between

P. alleghaniensis/P. spiloides compared to the other pairs.

For the eastern lineages, HZAR showed strong support for a right-tailed model

(ΔAIC_{DAPC} =-31.96 $-$ -107.42), whereas central lineages at the Mississippi River (ΔAIC_{DAPC} = -

4.26 $-$ -8.869) and western lineages at the Chihuahuan Desert and Edward's Plateau (ΔAIC_{DAPC} =

-5.51 $-$ -221.35) both supported a no-tails model (Fig. 6; Table S4 & S5). However, estimating

tailed models for the latter two comparisons may have required more samples than were

available. The width of clines decreased from the eastern lineages (width=126 km, 1st and 3rd

quartiles 114-138 km, tail length =53 km, 1st and 3rd quartiles 44-61 km), to central (width = 98

km, 1st and 3rd quartile 89-107 km), to the western lineages (width = 8 km, 1st and 3rd quartile =4-

15 km; Table 1). Sharply declining hybrid zones characterized the central and western lineages,

whereas the cline for eastern lineages showed a long right tail extending north.

Hybrid zone widths under neutrality should be larger than the actual widths given that

migration between sister pairs likely occurred prior to the LGM (Fig. S4). Contact time for each

of these pairs to approach the same actual width sizes all occurred within the Holocene: 516-

10,785 ybp (eastern lineages*),* 261-5424 ybp (central lineages) and 3-27 ybp (western lineages).

We also showed that selection against hybrids at the center of hybrid zones ranged from a

median of 0.3 % (0.02-0.6%) for eastern, 0.5% (0.03 to 1%) for central, and 79% (1.5-100%) for

western lineages.

When using a single SNP per locus we showed that 30% of loci from central lineages,

10.9% of loci for western lineages, and only 1.9% of loci for eastern lineages attained an F_{st}

>0.35. Given demographic histories between comparisons, this threshold for F_{st} is generally

considered extreme population disconnection beyond which advantageous loci will be exchanged

between populations (Wright 1978; Frankham et al. 2010).

Discussion

Grey Zone Dynamics

To understand the factors shaping reproductive isolation in a young species complex, we

ask whether divergence time predicts rates of gene flow or cline width across space in eastern

ratsnakes. This species complex is composed of four divergent lineages, with *P. alleghaniensis*

and *P. spiloides* meeting in the southeastern US at the intersection of subtropical and temperate

areas, *P. spiloides* and *P. obsoletus* separated at the Mississippi River, and *P. obsoletus* and *P.*

bairdi meeting at the connection on the forested and rocky habitats on the western edge of the

Edwards Plateau. All three spatially adjacent pairs of taxa show variable migration each having

unique cline shapes. The western lineages (*P. obsoletus*, *P. bairdi*) show very low migration

rates, a cline equivalent to only 0.35% of the parental range, fixation ($F_{st} > 0.35$) in 10.9% of

loci, and likely selection against hybrids. This is in comparison with central lineages (*P.*

obsoletus, P. spiloides) showing more migration, a cline occupying 5% of the parental ranges,

but with 35% of loci showing fixation, and possibly lower selection. Finally, eastern lineages (*P.*

alleghanienis, P. spiloides) have incomplete reproductive isolation (showing numerous F1 and

backcrosses) with only 1.9% of loci showing fixation, and high migration rates with a clinal tail

(Figs. 3 & 6). Where reproductive isolation occurs given low numbers of migrants/generation,

there is increased fixation among loci and steeper clines occur. The inverse is also true; when

reproductive isolation is incomplete, higher rates of gene flow less fixation of loci, and variable

cline widths occur. Importantly, however, reproductive isolation, gene flow estimates, and hybrid

clines show no association with timing of divergence.

Hybrid zone dynamics help to understand the interactions between pairs of taxa as

characteristics of the hybrid zone reveal whether there is selection against hybrids (tension

zones), selection for hybrids (bounded superiority), or ecological gradients with reduced fitness.

They also reveal whether selection is endogenous (genomic) or exogenous (environmental;

Barton and Hewitt 1985; Barton and Gale 1993). Our models indicate that migration at the

Mississippi River and in the eastern lineages occurred through the Pleistocene suggesting some

form of a hybrid zone existed through major climate change cycles. Given this age and assuming

selective neutrality, we demonstrate that these hybrid zones should be much larger, in most cases

exceeding the ranges of parental taxa (Fig. S4). The widths of the current hybrid zones are

maintained over time, preserving the historical and geographic identity of the parental taxa.

While there is some selection against hybrids as estimated from hybrid zone width and dispersal,

depending on species-pair comparison, it is possible that other factors may be constraining these

hybrid zones like fitness changes along an ecological gradient or isolation by environment (Case

and Taper 2000; Case et al. 2005; McEntee et al. 2018). There is support for this hypothesis

given significant differences among niches for all adjacent pairs of lineages (Fig S3).

Influence of barriers and environment on population structure

These ratsnake lineages are structured by a complex of physical barriers, historical

climate profiles, and geographic distance. Machine learning approaches show that the

Mississippi River has served as a strong barrier for isolating lineages of ratsnakes, forming

groups east and west of this river during the mid-Pleistocene at 1.08 My (95% CI =0.553 to

1.674 My; Figs 1&4, Table 1). These results are consistent with previous mtDNA and

morphological conclusions (Burbrink, 2001; Burbrink et al., 2000). The Mississippi River has

served as a strong barrier to gene flow given evidence from historical estimates of increased

drainage (Cox et al. 2014) and numerous unrelated organisms show genetic disconnection across

the river (Robison 1986; Burbrink 2002; Soltis et al. 2006; Burbrink et al. 2008; Pyron and

Burbrink 2009; Brandley et al. 2010; Satler and Carstens 2017; Myers et al. 2020).

Hybridization, while low (1-2 individuals/generation, Fig.3), does occur in a zone (~100km) east

of the Mississippi River between the central lineages (*P. obsoletus* and *P. spiloides*; Fig. 6).

Furthermore, divergence in this group can be predicted by elevation, geographic distance, and

mid-Pleistocene climate change (787kya). The timing of the mid-Pleistocene climate change falls

within the timing of divergence estimated from PipeMaster (Fig.4).

The Mississippi River is responsible for the earliest divergence in this complex but other

geographic features are also important. East of the Mississippi River, divergence between *P.*

alleghaniensis and *P. spiloides* was influenced by elevation, climate at the LGM, and geographic

distance after accounting for variable inflation (Fig 4). For these taxa, divergence occurred at the

Appalachian Mountains and Apalachicola/Chattahoochee River System, which suggests that

changes in elevation at the Appalachian Mountains, in part, played a role in isolating lineages or

maintaining divergence. Finer-scale testing is necessary to tease apart the effects of these river

systems, ancient embayments, and connections to uplifted areas. Although the timing of

divergence predates the LGM, it is possible that repeated 100,000 year glacial cycles isolated

these taxa into refugia, as has been suggested previously for ratsnakes and other organisms

(Waltari et al. 2007; Noss et al. 2015). Alternatively, *Pantherophis alleghaniensis* may have

diverged in an isolated Florida, due to sea level changes during interglacials, and subsequently

expanded north and west. The concentration of the yellow, striped morphology in the range of *P.*

alleghaniensis in Florida and parts of the southeastern coast of the US indicates that this isolation

likely resulted in these morphologically unique taxa (Schultz 1996; Burbrink 2001). This area of

isolation, defined as a suture zone between the Florida peninsula and the continental US has been

found for many taxa (Remington 1968; Swenson and Howard 2005; Ruane et al. 2015). While

these two forms can be delimited with genetic data and occupy distinctly different geographic

regions, reproductive isolation is not complete. However, niches are significantly different and *P.*

alleghaniensis is typically found in the Florida peninsula and coastal plain environments (Bailey

1995; Burbrink 2001), whereas *P. spiloides* is found throughout the remainder of the forested

habitats east of the Mississippi River including the ecoregions defined as interior river valleys

and hills, interior plateau, Appalachian habitats, southeastern and southcentral plains, and

Mississippi River valley. Uncertainty in the identity of taxa using morphology where these taxa

meet is likely due to extensive gene flow in hybrid zones at barriers following postglacial range

expansion (Fig. 1, 3; Burbrink 2001; Gibbs et al. 2006).

Divergence between *P. bairdi* and *P. obsoletus* occurred in the western edge of the

Edwardas Plateau, with the former extending westward into xeric forested or Chihuahuan desert

scrub habitats always associated with rocky habitats, whereas the latter extends east into Texas,

preferring forested habitats, (Lawson and Lieb 1990; Werler and Dixon 2000). Population

genetic structure between these taxa is associated with mid-Pleistocene climate, elevation, and

geographic distance (Fig. 4). Dates of origin of these taxa are within the mid-Pleistocene,

suggesting that past climate played a strong role in promoting population divergence. During this

period, the Chihuahuan Desert was much drier relative to the late-Pleistocene (Graham and Mead

1987; Metcalfe et al. 2002; Metcalfe 2006) and this increased aridity could have driven

ecological divergence between this species pair. While divergence time is not the oldest here,

migration rates are lowest, and a hybrid zone is narrow. Previous research shows limited

hybridization with backcrosses between these taxa occurring in a narrow region of the

southwestern Edwards Plateau (Lawson and Lieb 1990; Vandewege et al. 2012).

Despite each lineage occupying uniquely different habitats, population responses to

Pleistocene climate change were similar (Fig. 3). Given the divergence time for each group, N_e

has expanded by 50 times since the Pleistocene. These estimates are consistent with other

predictions in the ENA, where 75% of the tested vertebrates show population size expansion and

75% of tested snakes show synchronous expansion of N_e, likely coinciding with the retreat of

glaciers at various times in the Pleistocene or during the Holocene (Bintanja and van de Wal

2008; Burbrink et al. 2016). However, *Pantherophis bairdi* has a range only in a small area of

the west Texas and isolated populations in northeastern Mexico, where habitats may have been

indirectly affected by glacial cycles.

Taxonomy and delimitation in the genomic age

Delimiting species using genetic data, while useful, has proved controversial (Bauer et al.

2011; Sukumaran and Knowles 2017; Leaché et al. 2019). In particular, most genetic

delimitation methods fail to account for gene flow (but see Flouris et al. 2019), do not consider

spatial information, and place individuals into predefined groups prior to testing (O'Meara

2010). These methods all assume species diverge at an evolutionary time scale where mutations

accumulate through drift or selection (Rannala 2015). And while simulations show that a

threshold above one migrating individual/10 generations will cause model-based delimitation

methods to fail at finding two species (Zhang et al. 2011), it is not clear how pulses or

inconsistent migration affect species transitioning through the gray zone. Importantly, Wright

(1931) indicated that up to 1,000 individuals migrating/generation may not prevent populations

from drifting given effective population size and spatial segregation.

Modern thresholds for species delimitation (Jackson et al. 2016; Leaché et al. 2019) use

GDI (>0.7), estimates of species divergence time over population size ($2\tau/\theta>1$), absolute

divergence time (10^4) and maximum migration rates ($M=Nm<1$), metrics that are all likely

influenced by sampling effort, location of samples sequenced, and detection of hybrids. For

example, if large hybrid zones exist but remain poorly sampled then estimates of migration will

be reduced, GDI will be high, and the probability of MSC estimators will suggest multiple

species. Conversely, if only hybrid zones are sampled then all metrics will predict only a single

species. For all pairs of ratsnakes $2\tau/\theta>1$ and divergence time is always greater than 10^4.

However, migration rates remain high between adjacent pairs in the eastern lineages. This forces

us to address how we are defining a species in the context of hybrid zones, how we infer

reproductive isolation beyond MSC methods, and how selection is maintaining hybrid zones. For

all lineages of ratsnakes, hybrid zones are smaller than predicted by neutral selection if these

hybrid zones formed in the Pleistocene. It is possible that these hybrid zones formed after the

retreat of the last glacial cycle during the Holocene. However, geographic lineage identity has

been maintained for >133,000 generations in all pairs, regardless of migration, and niches are

unique among all lineages. Moreover, where hybrid zones extend over large areas, gene flow

likely varies throughout space and may have changed over time (Barton and Hewitt 1985; Mallet

2007), particularly in response to glacial cycles. Thus, identifying and delimiting species in the

context of admixture is complex. Still, maintaining lineage and geographic identity, even with a

hybrid zone suggests that selection may have maintained unique evolutionary trajectories.

Here we have demonstrated that these four species, delimited previously using mtDNA

and morphology (Burbrink, 2001; Burbrink et al., 2000) can be recognized as distinct taxa given

methods of spatial grouping and species delimitation, though ranges in the eastern-most lineage

are restricted much further south than previously thought. However, migration rates between *P.*

spiloides and *P. alleghaniensis* (Fig. 3) are high enough to draw some concern that these taxa

should not be recognized given that they may not represent independent evolutionary trajectories

(de Queiroz, 2007; Rannala, 2015; Zhang et al., 2011). Unique niches in the subtropical areas of

the southeastern US and morphology provides some evidence for isolation between these taxa:

the striped forms (with variable ground colors of orange, yellow, green, and gray) are indicative

of *P. alleghaniensis* whereas adults with dark saddle patterns on grey, olive, brown, or black

ground colors (or completely black with no pattern) are found west of Florida and east of the

Mississippi River, throughout the Midwest and up to the northeastern US and Canada represent

P. spiloides. However, hybridization north and west of Florida is likely represented by a dulling

of this yellow ground color and mixed patterned individuals (Schultz 1996; see Supporting

Information Material 1 for more information on taxonomy).

Future Directions

To better understand the complexity of speciation processes, an even more integrated

whole-genomic approach with dense population sampling should be implemented. Even when

researchers have demonstrated that integrating spatial information along with estimates of timing

and gene-flow among groups presents a more comprehensive picture of speciation over a

complex landscape, there still remain several areas of investigation. For instance, estimating

gene flow and cline shape using detailed transects across various intersections along hybrid

zones would provide a better understanding of how migration changes in different habitats at

different latitudes and what type of hybrid zone is present. Additionally, better integrating

spatial, habitat, and migration estimates would allow researchers to test dynamic speciation

models that could be used to understand how migration changes over the landscape at various

times throughout the history of these taxa. Finally, using whole genome data along clines will

help determine if selection is occurring in the hybrid zones and determine the function of loci

showing evidence of selection and resisting gene flow.

Author contributions: FTB and EAM contributed to data collection, all authors contributed to

analyses, FTB wrote the initial draft, and all authors edited subsequent versions.

Acknowledgments: We thank R. Glor, R. Brown, and L. Welton at KU and C. Austin and D.

Dittman at LSUMN and K. Krysko and FLMNH, D. Shepard at Louisiana Tech University, and

A. McKelvy for providing tissues. We also thank L. and G. Derryberry for help with HZAR. We

also thank E. Chen for help extracting DNA. EAM was funded by the Gerstner Scholar/Theodore

Roosevelt and Peter Buck/Walter Rathbone Bacon fellowships. We thank D. Frost, A. Leaché,

A. Bauer, and K. deQuieroz for helpful discussions on species delimitation, type specimens, and

admixed type specimens.

Data Accessibility Statement: All data and unique scripts are available on Dryad XXX.

Literature Cited

Agrawal, A. F., J. L. Feder, and P. Nosil. 2011. Ecological Divergence and the Origins of

Intrinsic Postmating Isolation with Gene Flow. Int. J. Ecol. 2011:1–15.

Bailey, R. G. 1995. Description of the ecoregions of the United States. 2nd ed. USDA Forest

Service.

Bailey, R. I., M. R. Tesaker, C. N. Trier, and G.-P. Saetre. 2015. Strong selection on male

plumage in a hybrid zone between a hybrid bird species and one of its parents. J. Evol. Biol.

28:1257–1269.

Barton, H. H., and K. Gale. 1993. Genetic analysis of hybrid zones. Pp. 13–45 *in* Hybrid Zones

and the Evolutionary Process. Oxford University Press, New York.

Barton, N. H. 2010. What role does natural selection play in speciation? Philos. Trans. R. Soc.

Lond. B. Biol. Sci. 365:1825–40.

Barton, N. H., and G. M. Hewitt. 1985. Analysis of Hybrid Zones. Annu. Rev. Ecol. Syst.

16:113–148.

Bauer, A. M., J. F. Parham, R. M. Brown, B. L. Stuart, L. Grismer, T. J. Papenfuss, W. Böhme,

J. M. Savage, S. Carranza, J. L. Grismer, P. Wagner, A. Schmitz, N. B. Ananjeva, and R. F.

Inger. 2011. Availability of new Bayesian-delimited gecko names and the importance of

character-based species descriptions. Proc. R. Soc. B Biol. Sci. 278:490.

Beugin, M.-P., T. Gayet, D. Pontier, S. Devillard, T. Jombart, and T. Hansen. 2018. A fast

likelihood solution to the genetic clustering problem. Methods Ecol. Evol. 9:1006–1016.

Bintanja, R., and R. S. W. van de Wal. 2008. North American ice-sheet dynamics and the onset

of 100,000-year glacial cycles. Nature 454:869–72.

Bouckaert, R., J. Heled, D. Kühnert, T. Vaughan, C.-H. Wu, D. Xie, M. A. Suchard, A.

Rambaut, and A. J. Drummond. 2014. BEAST 2: A software platform for Bayesian

evolutionary analysis. PLoS Comput. Biol. 10:e1003537.

Bradburd, G. S., G. M. Coop, and P. L. Ralph. 2018. Inferring continuous and discrete

population genetic structure across space. Genetics 210:33–52.

Brandley, M. C., T. J. Guiher, R. A. Pyron, C. T. T. Winne, and F. T. Burbrink. 2010. Does

dispersal across an aquatic geographic barrier obscure phylogeographic structure in the

diamond-backed watersnake (*Nerodia rhombifer*)? Mol. Phylogenet. Evol. 57:552–560.

Branstetter, M. G., and J. T. Longino. 2019. Ultra-conserved element phylogenomics of New

World Ponera (Hymenoptera: Formicidae) illuminates the origin and phylogeographic

history of the endemic exotic ant Ponera exotica. Insect Syst. Divers. 3.

Brown, J. L., D. J. Hill, A. M. Dolan, A. C. Carnaval, and A. M. Haywood. 2018. PaleoClim,

high spatial resolution paleoclimate surfaces for global land areas. Sci. Data 5:180254.

Bryant, D., R. Bouckaert, J. Felsenstein, N. A. Rosenberg, and A. RoyChoudhury. 2012.

Inferring species trees directly from biallelic genetic markers: bypassing gene trees in a full

coalescent analysis. Mol. Biol. Evol. 29:1917–1932.

Burbrink, F. T. 2002. Phylogeographic analysis of the cornsnake (*Elaphe guttata*) complex as

inferred from maximum likelihood and Bayesian analyses. Mol. Phylogenet. Evol. 25:465–

476.

Burbrink, F. T. 2001. Systematics of the eastern ratsnake complex (*Elaphe obsoleta*). Herpetol.

Monogr. 1–53.

Burbrink, F. T., Y. L. Chan, E. A. Myers, S. Ruane, B. T. Smith, and M. J. Hickerson. 2016.

Asynchronous demographic responses to Pleistocene climate change in Eastern Nearctic

vertebrates. Ecol. Lett. 19:1457–1467.

Burbrink, F. T., F. Fontanella, R. A. Pyron, T. J. Guiher, C. Jimenez, and R. Alexander Pyron.

2008. Phylogeography across a continent: The evolutionary and demographic history of the

North American racer (Serpentes : Colubridae : *Coluber constrictor*). Mol. Phylogenet.

Evol. 47:274–288.

Burbrink, F. T., F. G. Grazziotin, R. A. Pyron, D. Cundall, S. Donnellan, F. Irish, J. S. Keogh, F.

Kraus, R. W. Murphy, B. Noonan, C. J. Raxworthy, S. Ruane, A. R. Lemmon, E. M.

Lemmon, and H. Zaher. 2020. Interrogating genomic-scale data for Squamata (lizards,

snakes, and amphisbaenians) shows no support for key traditional morphological

relationships. Syst. Biol. 69:502–520.

Burbrink, F. T., R. Lawson, and J. B. Slowinski. 2000. Mitochondrial DNA phylogeography of

the polytypic North American rat snake (*Elaphe obsoleta*): A critique of the subspecies

concept. Evolution (N. Y). 54:2107–2118.

Case, T. J., R. D. Holt, M. A. McPeek, and T. H. Keitt. 2005. The community context of species'

borders: ecological and evolutionary perspectives. Oikos 108:28–46.

Case, T. J., and M. L. Taper. 2000. Interspecific competition, environmental gradients, gene

flow, and the coevolution of species' borders. Am. Nat. 155:583–605.

Cox, R. T., D. N. Lumsden, and R. B. Van Arsdale. 2014. Possible relict meanders of the

Pliocene Mississippi River and their implications. J. Geol. 122:609–622.

De Queiroz, K. 2007. Species concepts and species delimitation. Syst. Biol. 56:879–886.

Derryberry, E. P., G. E. Derryberry, J. M. Maley, and R. T. Brumfield. 2014. hzar : hybrid zone

analysis using an R software package. Mol. Ecol. Resour. 14:652–663.

Edwards, S., R. Hopkins, and J. Mallet. 2020. Speciation. Pp. 296–318 *in* S. Scheiner and D.

Mindell, eds. The Theory of Evolution. University of Chicago Press, Chicago.

Faircloth, B. C., J. E. McCormack, N. G. Crawford, M. G. Harvey, R. T. Brumfield, and T. C.

Glenn. 2012. Ultraconserved Elements Anchor Thousands of Genetic Markers Spanning

Multiple Evolutionary Timescales. Syst. Biol. 61:717–726.

Feder, J. L., S. P. Egan, and P. Nosil. 2012. The genomics of speciation-with-gene-flow. Trends

Genet. 28:342–350.

Flouris, T., X. Jiao, B. Rannala, and Z. Yang. 2019. A Bayesian implementation of the

multispecies coalescent model with introgression for phylogenomic analysis. Mol. Biol.

Evol., doi: 10.1093/molbev/msz296.

Frankham, R., J. D. (Jonathan D. . Ballou, and D. A. (David A. Briscoe. 2010. Introduction to

conservation genetics. Cambridge University Press.

Frantz, A. C., L. C. Pope, T. R. Etherington, G. J. Wilson, and T. Burke. 2010. Using isolation-

by-distance-based approaches to assess the barrier effect of linear landscape elements on

badger (Meles meles) dispersal. Mol. Ecol. 19:1663–1674.

Garrick, R. C., J. D. Banusiewicz, S. Burgess, C. Hyseni, and R. E. Symula. 2019. Extending

phylogeography to account for lineage fusion. J. Biogeogr. 46:268–278.

Gavrilets, S. 2004. Fitness landscapes and the origin of species. Princeton University Press.

Gay, L., P.-A. Crochet, D. A. Bell, and T. Lenormand. 2008. Comparing clines on molecular and

phenotypic traits in hybrid zones: a window on tension zone models. Evolution (N. Y).

62:2789–2806.

Gehara, M., A. A. Garda, F. P. Werneck, E. F. Oliveira, E. M. da Fonseca, F. Camurugi, F. d. M.

Magalhães, F. M. F. M. Lanna, J. W. Sites, R. Marques, R. Silveira-Filho, V. A. São Pedro,

G. R. Colli, G. C. Costa, and F. T. Burbrink. 2017. Estimating synchronous demographic

changes across populations using hABC and its application for a herpetological community

from northeastern Brazil. Mol. Ecol. 26:4756–4771.

Gibbs, H. L., S. J. Corey, G. Blouin-Demers, K. A. Prior, and P. J. Weatherhead. 2006.

Hybridization between mtDNA-defined phylogeographic lineages of black ratsnakes

(*Pantherophis sp.*). Mol. Ecol. 15:3755–3767.

Gourbière, S., and J. Mallet. 2010. Are species real? The shape of the species boundry with

exponential failure, reinforcement, and the "missing snowball." Evolution (N. Y). 64:1–24.

Graham, R. W., and J. I. Mead. 1987. Environmental fluctuations and evolution of mammalian

faunas during the last deglaciation in North America. Pp. 371–402 *in* North America and

Adjacent Oceans During the Last Deglaciation. Geological Society of America, Boulder,

Colorado 80301.

Harrison, R. G. 1993. Hybrid zones and the evolutionary process. Oxford University Press.

Hewitt, G. 2000. The genetic legacy of the Quaternary ice ages. Nature 405:907–913.

Hewitt, G. M. 2008. Speciation, hybrid zones and phylogeography - or seeing genes in space and

time. Mol. Ecol. 10:537–549.

Hey, J. 2010. Isolation with migration models for more than two populations. Mol. Biol. Evol.

27:905–920.

Hey, J., and R. Nielsen. 2004. Multilocus methods for estimating population sizes, migration

rates and divergence time, with applications to the divergence of *Drosophila pseudoobscura*

and *D-persimilis*. Genetics 167:747–760.

Hijmans, R. J., J. van Etten, and J. Checg. 2014. Geographic Data Analysis and Modeling [R

package raster version 2.5-8]. Comprehensive R Archive Network (CRAN).

Jackson, N. D., B. C. Carstens, A. E. Morales, and B. C. O'Meara. 2016. Species delimitation

with gene flow. Syst. Biol. 66:799–812.

Jiggins, C. D., and J. Mallet. 2000. Bimodal hybrid zones and speciation. Trends Ecol. Evol.

15:250–255.

Jombart, T. 2008. adegenet: a R package for the multivariate analysis of genetic markers.

Bioinformatics 24:1403–1405.

Jombart, T., S. Devillard, and F. Balloux. 2010. Discriminant analysis of principal components: a

new method for the analysis of genetically structured populations. BMC Genet. 11:94.

Joseph, L., and T. Wilke. 2007. Lack of phylogeographic structure in three widespread

Australian birds reinforces emerging challenges in Australian historical biogeography. J.

Biogeogr. 34:612–624.

Karrenberg, S., X. Liu, E. Hallander, A. Favre, J. Herforth-Rahmé, and A. Widmer. 2019.

Ecological divergence plays an important role in strong but complex reproductive isolation

in campions (Silene). Evolution (N. Y). 73:245–261.

Kuhn, M. 2008. Caret: Classification and Regression Training Package. R package version 6.0-

77.

Lawson, R., and C. S. Lieb. 1990. Variation and hybridization in *Elaphe bairdi* (Serpentes:

Colubridae). J. Herpetol. 24:280.

Leaché, A. D., T. Zhu, B. Rannala, and Z. Yang. 2019. The spectre of too many species. Syst.

Biol. 68:168–181.

Legendre, P., and M. J. Fortin. 2010. Comparison of the Mantel test and alternative approaches

for detecting complex multivariate relationships in the spatial analysis of genetic data. Mol.

Ecol. Resour. 10:831–844.

Legendre, P., J. Oksanen, and C. J. F. ter Braak. 2011. Testing the significance of canonical axes

in redundancy analysis. Methods Ecol. Evol. 2:269–277.

Lek, S., M. Delacoste, P. Baran, I. Dimopoulos, J. Lauga, and S. Aulagnier. 1996. Application of

neural networks to modelling nonlinear relationships in ecology. Ecol. Modell. 90:39–52.

Lek, S., and J. F. Gue. 1999. Artificial neural networks as a tool in ecological modelling , an

introduction. 120:65–73.

Libbrecht, M. W., and W. S. Noble. 2015. Machine learning applications in genetics and

genomics. Nat. Rev. Genet. 16:321–332.

Lourenço, C. R., K. R. Nicastro, C. D. McQuaid, R. M. Chefaoui, J. Assis, M. Z. Taleb, and G. I.

Zardi. 2017. Evidence for rangewide panmixia despite multiple barriers to dispersal in a

marine mussel. Sci. Rep. 7:10279. Nature Publishing Group.

Makowsky, R., J. Chesser, and L. J. Rissler. 2009. A striking lack of genetic diversity across the

wide-ranging amphibian *Gastrophryne carolinensis* (Anura: Microhylidae). Genetica

135:169–183. Springer Netherlands.

Mallet, J. 2007. Hybrid speciation. Nature 446:279–283. Nature Publishing Group.

Mayr, E. 1963. Animal species and evolution. Belknap Press of Harvard University Press,

Cambridge,.

McEntee, J. P., J. G. Burleigh, and S. Singhal. 2018. Dispersal predicts hybrid zone widths

across animal diversity: Implications for species borders under incomplete reproductive

isolation. bioRxiv 472506. Cold Spring Harbor Laboratory.

Metcalfe, S. E. 2006. Late Quaternary environments of the northern deserts and central

transvolcanic belt of Mexico. Ann. Missouri Bot. Gard. 93:258–273. Missouri Botanical

Garden Press.

Metcalfe, S., A. Say, S. Black, R. McCulloch, and S. O'Hara. 2002. Wet Conditions during the

Last Glaciation in the Chihuahuan Desert, Alta Babicora Basin, Mexico. Quat. Res. 57:91–

101.

Myers, E. A., A. D. McKelvy, and F. T. Burbrink. 2020. Biogeographic barriers, Pleistocene

refugia, and climatic gradients in the southeastern Nearctic drive diversification in

cornsnakes (*Pantherophis guttatus* complex). Mol. Ecol. 29:797–811. John Wiley & Sons,

Ltd.

Myers, E. A., A. T. Xue, M. Gehara, C. L. Cox, A. R. Davis Rabosky, J. Lemos-Espinal, J. E.

Martínez-Gómez, and F. T. Burbrink. 2019. Environmental heterogeneity and not vicariant

biogeographic barriers generate community-wide population structure in desert-adapted

snakes. Mol. Ecol. 28:4535–4548.

Nei, M. 1973. Analysis of gene diversity in subdivided populations. Proc Natl Acad Sci USA

70:3321–3323.

Nosil, P. 2012. Ecological Speciation. Oxford University Press, Oxford.

Nosil, P., and B. J. Crespi. 2006. Ecological divergence promotes the evolution of cryptic

reproductive isolation. Proc. R. Soc. B-Biological Sci. 273:991–997.

Nosil, P., J. L. Feder, S. M. Flaxman, and Z. Gompert. 2017. Tipping points in the dynamics of

speciation. Nat. Ecol. Evol. 1:1. Nat Ecol Evol.

Noss, R. F., W. J. Platt, B. A. Sorrie, A. S. Weakley, D. B. Means, J. Costanza, and R. K. Peet.

2015. How global biodiversity hotspots may go unrecognized: lessons from the North

American Coastal Plain. Divers. Distrib. 21:236–244.

O'Meara, B. C. 2010. New heuristic methods for joint species delimitation and species tree

inference. Syst. Biol. 59:59–73.

Orr, H. A. 1995. The population genetics of speciation: the evolution of hybrid incompatibilities.

Genetics 139:1805–13. Genetics Society of America.

Paradis, E., and K. Schliep. 2019. ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. Bioinformatics 35:526–528. .

Petkova, D., J. Novembre, and M. Stephens. 2016. Visualizing spatial population structure with

estimated effective migration surfaces. Nat. Genet. 48:94–100. Nature Publishing Group.

Plummer, M., N. Best, K. Cowles, and K. Vines. 2006. CODA: Convergence diagnosis and

output analysis for MCMC. R News 6:7–11.

Pyron, R. A., and F. T. Burbrink. 2009. Lineage diversification in a widespread species: roles for

niche divergence and conservatism in the common kingsnake, Lampropeltis getula. Mol.

Ecol. 18:3443–3457.

Rannala, B. 2015. The art and science of species delimitation. Curr. Zool. 61:846–853.

Remington, C. L. 1968. Suture-zones of hybrid interaction between recently joined biotas. Pp.

321–428 *in* Evolutionary Biology. Springer US, Boston, MA.

Riddle, B. R. 2016. Comparative phylogeography clarifies the complexity and problems of

continental distribution that drove A. R. Wallace to favor islands. Proc. Natl. Acad. Sci. U.

S. A. 113:7970.

Robison, H. W. 1986. Zoogeographic implications of the Mississippi River Basin. Pp. 267–285

in The Zoogeography of North American Freshwater Fishes. Wiley, New York.

Roux, C., C. Fraïsse, J. Romiguier, Y. Anciaux, N. Galtier, and N. Bierne. 2016. Shedding light

on the grey zone of speciation along a continuum of genomic divergence. PLoS Biol.

14:e2000234.

Ruane, S., O. Torres-Carvajal, and F. T. Burbrink. 2015. Independent demographic responses to

climate change among temperate and tropical milksnakes (Colubridae: Genus

Lampropeltis). PLoS One 10:e0128543.

Satler, J. D., and B. C. Carstens. 2017. Do ecological communities disperse across biogeographic

barriers as a unit? Mol. Ecol. 26:3533–3545.

Schultz, K. D. 1996. A monograph of the colubrid snakes of the genus *Elaphe* Fitzinger. Koelz

Scientific Books, Havlicuv.

Seehausen, O., R. K. Butlin, I. Keller, C. E. Wagner, J. W. Boughman, P. A. Hohenlohe, C. L.

Peichel, G.-P. Saetre, C. Bank, A. Brannstrom, A. Brelsford, C. S. Clarkson, F.

Eroukhmanoff, J. L. Feder, M. C. Fischer, A. D. Foote, P. Franchini, C. D. Jiggins, F. C.

Jones, A. K. Lindholm, K. Lucek, M. E. Maan, D. A. Marques, S. H. Martin, B. Matthews,

J. I. Meier, M. Most, M. W. Nachman, E. Nonaka, D. J. Rennison, J. Schwarzer, E. T.

Watson, A. M. Westram, and A. Widmer. 2014. Genomics and the origin of species. Nat

Rev Genet 15:176–192.

Sheehan, S., Y. S. Song, E. Buzbas, D. Petrov, A. Boyko, and A. Auton. 2016. Deep learning for

population genetic inference. PLOS Comput. Biol. 12:e1004845.

Singhal, S., and C. Moritz. 2013. Reproductive isolation between phylogeographic lineages

scales with divergence. Proceedings. Biol. Sci. 280:20132246.

Soltis, D. E., A. B. Morris, J. S. McLachlan, P. S. Manos, and P. S. Soltis. 2006. Comparative

phylogeography of unglaciated eastern North America. Mol. Ecol. 15:4261–4293.

Stankowski, S., J. M. Sobel, and M. A. Streisfeld. 2017. Geographic cline analysis as a tool for

studying genome-wide variation: a case study of pollinator-mediated divergence in a

761  monkeyflower. Mol. Ecol. 26:107–122.

762 Sukumaran, J., and L. L. Knowles. 2017. Multispecies coalescent delimits structure, not species.

763  Proc. Natl. Acad. Sci. U. S. A. 114:1607–1612.

764 Sun, K., K. A. Meiklejohn, B. C. Faircloth, T. C. Glenn, E. L. Braun, and R. T. Kimball. 2014.

765  The evolution of peafowl and other taxa with ocelli (eyespots): a phylogenomic approach.

766  Proc. R. Soc. B Biol. Sci. 281:20140823.

767 Swenson, N. G., and D. J. Howard. 2005. Clustering of contact zones, hybrid zones, and

768  phylogeographic breaks in North America. Am. Nat. 166:581–591.

769 Vandewege, M. W., D. Rodriguez, J. P. Weaver, T. D. Hibbitts, M. R. J. Forstner, and L. D.

770  Densmore. 2012. Evidence of hybridization between *Elaphe bairdi* and *Elaphe obsoleta*

771  *lindheimeri* including comparative population genetics inferred from microsatellites and

772  mitochondrial DNA. J. Herpetol. 46:56–63. Society for the Study of Amphibians and

773  Reptiles.

774 Vavrek, M. J. 2020. Fossil. https://cran.r-project.org/web/packages/fossil/index.html.

775 Walker, D., and J. C. Avise. 1998. Principles of phylogeography as illustrated by freshwater and

776  terrestrial turtles in the southeastern United States. Annu. Rev. Ecol. Syst. 29:23–58.

777 Waltari, E., R. J. Hijmans, A. T. Peterson, Á. S. Nyári, S. L. Perkins, and R. P. Guralnick. 2007.

778  Locating Pleistocene refugia: comparing phylogeographic and ecological niche model

779  predictions. PLoS One 2:e563.

780 Werler, J. E., and J. R. Dixon. 2000. Texas snakes : identification, distribution, and natural

781  history. 1st ed. University of Texas Press, Austin.

782 Wright, S. 1978. Evolution and the Genetics of Populations. Vol. 4. Variability within and

783  among Natural Populations. University Chicago Press.

Wright, S. 1931. Evolution in Mendelian populations. Genetics 16:97–159.

Wright, S. 1943. Isolation by Distance. Genetics 28:114–38. Genetics Society of America.

Younger, J. L., P. Dempster, Á. S. Nyári, T. O. Helms, M. J. Raherilalao, S. M. Goodman, and S.

Reddy. 2019. Phylogeography of the Rufous vanga and the role of bioclimatic transition

zones in promoting speciation within Madagascar. Mol. Phylogenet. Evol. 139:106535.

Zellmer, A. J., M. M. Hanes, S. M. Hird, and B. C. Carstens. 2012. Deep phylogeographic

structure and environmental differentiation in the carnivorous plant Sarracenia alata. Syst.

Biol. 61:763–777.

Zhang, C., D.-X. Zhang, T. Zhu, and Z. Yang. 2011. Evaluation of a bayesian coalescent method

of species delimitation. Syst. Biol. 60:747–761.

Zhang, W. 2010. Computational Ecology: Artificial Neural Networks and Their Applications.

World Scientific.

807 Table 1. Measures of divergence dates in millions of years (My), migration rates (first number is
808 first taxon into second taxa, e.g., *Pantherophis alleganiensis* into *P.spiloides*, and second vice
809 versa), and migration rates (individuals/generation) for each contacting group in the
810 *Pantherophis obsoletus* complex.
811

Group	Age (My)	Migration Rate (ind/gen)	Cline Width (m)
Eastern (*P. alleghaniensis* x *P.spiloides*)	0.65 (0.367-0.875)	2.79/3.79	125 (114-138)
Central (*P.spiloides* x *P.obsoletus*)	1.08 (0.55-1.67)	1.96/1.28	97 (88-107)
Western (*P.obsoletus* x *P. bairdi*)	0.367 (0.37-0.88)	0.65/0.55	8 (4-15)

Figures and Legends

Fig.1. Distribution and population delimitation for the ratsnakes *Pantherophis obsoletus*

complex. (A) Map showing range and geographic features: Chihuahuan Desert Province, the

Mississippi River, and the Appalachian Mountains and Apalachicola/Chattahoochee River

System (AARS). (B) Estimated effective migration surfaces (EEMS) showing areas of both high

(blue) and low migration (brown). (C) Population structure using Discriminant Analysis of

Principal Components (DAPC) shows the location of four geographically distinct lineages. (D)

Bi-plots of PC space shows relative distances among the four taxa using DAPC.

Fig.2. The results of SNAPP species-tree estimation for the four taxa over all loci (A) and models showing Isolation Migration (IM) and Isolation Migration with Demographic Change (IMD; best-fit model) and Isolation with Recent Migration and Demographic Change (IMRD) estimated in PipeMaster. Colored arrows represent migration among spatially adjacent pairs. (B). Photographs representing some of the typical color patterns seen in ratsnakes: 1) *Pantherophis alleghaniensis* (photo F. Burbrink), 2) *P. alleghaniensis* (photo N. Claunch), 3) *P. spiloides* (photo A. Meier), 4) *P. obsoletus* (photo D. Shepard), and 5) *P. bairdi* (E. Myers).

Fig.3. Results from PipeMaster showing estimates and directionality of gene flow (A),

divergence times between groups and taxon pairs (B), and changes in population sizes over time

(C).

862   Fig.4. Artificial neural network results showing accuracy predicting genetic structure among all

863   lineages and species pairs (left) and ranked variable importance (filtered for multicollinearity in

864   inset box, right).

Fig.5. Results from estimating hybrid types between taxon pairs using *snapclust* showing the spatial distribution of parentals and hybrids along with their frequencies.

Eastern Lineages *(Pantherophis alleghaniensis x Pantherophis spiloides)*

Central Lineages (Pantherophis spiloides x Pantherophis obsoletus)

Western Lineages (Pantherophis bairdi x Pantherophis obsoletus)

Fig.6. Cline model results for all three species pairs using Gaussian Cline (HZAR) model given

locations and admixture estimates using DAPC.

Supplemental Material

DRYAD XXX

Genomic Dataset

Geographic Localities

R scripts

Supporting Information Material 1

Additional Methods

Table S1 – Priors and summary stats for PipeMaster; estimating migration, timing of divergence,

and historical demography

Table S2 – PHRAPL Results (*open with R read.table() or Excel*)

Table S3 – RDA and ANN Results

Table S4 – HZAR Results for *snapclust* assignments

Table S5 – HZAR results for DAPC assignments

Fig.S1. Delimiting populations using sparse nonnegative matrix factorization (SNMF) and

spatial Principal Component analysis (sPCA) showing the location of four geographically

distinct lineages.

Fig. S2 – PipeMaster model fit of observed data to simulations under IM, IMD, and IMD-LGM

models for PC1-PC10.

Fig. S3 – Symmetric background niche similarity tests for spatially adjacent pairs (eastern,

central, and western lineage comparisons) for the *D* and *I* statistics.

Fig. S4 – Prediction of cline width under neutrality given origin times for each species pair: A)

Pantherophis alleghaniensis/P. spiloides, B) *P. spiloides/P.obsoletus*, and C) *P.*

obsoletus/P.bairdi.

Supporting Information Material 1

Supplemental Methods and Results

Dataset

We sampled 288 individuals liberally covering the range of all species within the

Pantherophis obsoletus complex (Fig.1; Dryad XXX). DNA was extracted from all samples

using Qiagen DNeasy Blood & Tissue Kits and samples were screened for quality using broad-

range Qubit Assays (https://www.thermofisher.com/us/en/home/industrial/spectroscopy-

elemental-isotope-analysis/molecular-spectroscopy/fluorometers/qubit/qubit-assays.html). We

used services from RAPiD Genomics (https://www.rapid-genomics.com/services/) to generate

5472 baits and to sequence 5060 conserved elements (UCEs) loci following the protocols from

(Faircloth et al., 2012) and (Sun et al., 2014).

Raw sequence reads were trimmed of adapter contamination using illumiprocessor

(Faircloth, 2013), a wrapper around the trimmomatic package (Bolger et al., 2014). These

sequence capture data were then assembled by mapping reads to all UCEs found within a

Chromium 10x assembled *Pantherophis spiloides* genome (Burbrink and Myers, *in prep*) after

2,500 base pairs of flanking DNA around UCE baits were pulled from this genome assembly

using the Phyluce commands *phyluce_probe_run_multiple_lastzs_sqlite* and

phyluce_probe_slice_sequence_from_genomes (Faircloth, 2016). Sequence reads for each

sample were then mapped to the 'reference' UCE dataset using bwa v0.7.17 (Li and Durbin,

2009). Samtools was used to convert sam format to bam files and alignments were soft clipped

using the CleanSam tool in Picard (http://broadinstitute.github.io/picard/). To phase these data,

we mapped reads back to each individual sample's alignments. To do this, bam files were

converted to fastq using samtools mpileup (Li et al., 2009), fastq files were converted to fasta

format via seqtk (https://github.com/lh3/seqtk) and all missing sites were removed using seqkit

(Shen et al 2016). These newly created fasta files were then used as a reference sequence for

phasing the data following the seqcap_pop pipeline (Harvey et al., 2016). Briefly we followed

this pipeline and used GATK (McKenna et al., 2010) to mark duplicates, call, realign, and

annotate/mask indels, call and annotate SNPs via GATK, restrict SNP calling to high quality

SNPs, and then conducted read backed phasing. Finally, we created a sample specific fasta file of

all phased loci for every sample using 'add_phased_snps_to_seqs_filter.py' available via the

seqcap_pop pipeline (Harvey et al., 2016). All samples were combined into locus specific fasta

files using a perl script developed in (Myers et al., 2019). Locus specific fasta files were aligned

using muscle as implemented in the R package ape (Paradis and Schliep, 2019) after removing

individuals from alignments that fell below the 25% quantile of the distribution of sequence

lengths for each locus. All fasta files were conservatively trimmed of missing data resulting in

alignments with no missing sites (i.e., trimming to the shortest sequence in the alignment). We

filtered these alignments by removing all loci that were missing >50% sequenced samples and

removed all individuals that were missing from >30% of all alignments. Finally, we created a vcf

file from these alignments and using vcftools (Danecek et al., 2011) and filtered this for minor

allele frequency >10% retaining only 1 SNP/locus for population assignment analyses (Linck

and Battey, 2019).

Geographic groupings

We estimated population structure by comparing Discriminant Analysis of Principal Components

(DAPC; Jombart et al., 2010) and spatial Principal Component analysis (sPCA; Jombart et al.,

2008) in adegenet v2.1.2(Jombart 2008), sparse nonnegative matrix factorization (SNMF;

Frichot et al., 2014) in LEA v1.4.0 (Frichot and François 2015), and estimated effective

migration surfaces (EEMS v; Petkova et al., 2016), which accounts for IBD . Each of these

methods use different assumptions for grouping individuals and we compare congruence among

them.The model-free method DAPC sequentially estimates K-means for clustering and selects

the best fit to infer genetic clusters in the absence of prior group identification. Using the

package adegenet (Jombart 2008) in R (R Core Team 2015) we first transformed data using PCA

by choosing a large number of PC axes (n=200), then picking the number of discriminant

functions yielding large F-statistics (>2000) and optimal number of groups using BIC (Bayesian

inference criterion). Because using a large number of the PCs may yield arbitrary solutions, we

estimated the optimal *a-score*, which measures this bias by calculating the difference between

the actual cluster assignment probabilities and randomly assigned probabilities (predicting 15-23

PCs). We also used cross-validation with a 90% training and 10% test dataset to estimate the

average predicted success for each group.

Because DAPC does not account for spatial structure, such as IBD, we also used sPCA.

Similar to DAPC, this method does not require data to be in Hardy-Weinberg or linkage

equilibrium. sPCA also adds an element of space by isolating global structure, representing

disconnected groups or clines, from local structure, representing repulsion (selection against

similar genetic types co-occurring), and random noise. Using adegenet, we estimated sPCA

given SNPs and locations of each sample using up to 15 positive eigenvalues (global structures)

and 15 negative eigenvalues (local structures). To connect individual locations to build a

connection network, we used Delaunay triangulation. We also implemented Monte-Carlo tests

with 100 permutations using the function *spca_randtests* to determine if significant local or

global spatial structures exist.

We also assessed ancestral coefficients using SNMF in the package LEA (Frichot and

François 2015). This methodology is comparable to ADMIXTURE (Alexander et al. 2009) and

Structure (Pritchard et al. 2000) methods, but runs 10-30x faster using genomic-scale data. This

method produces a least-squares estimate of ancestry populations given K ancestral populations

We estimated ancestry coefficients over 1-6 K using 100 repetitions per K, 100 iterations per

algorithm, and a tolerance of 0.05. To determine the appropriate number of groups, we used a

cross-validation technique with the entropy criterion, which masks genotypes to fit a particular

model to each K.

Finally, to compare these methods of grouping individuals into populations, we estimated

migration rates over the landscape using EEMS (Petkova et al. 2016). This method models

effective migration rates over geography to represent regions where migration is low in cases

when genetic dissimilarity increases rapidly, thus providing a uniquely distinct view of the

location of population clusters relative to biogeographic barriers as compared to DAPC, sPCA,

and SNMF. We ran EEMS 3 times for 3×10^6 generations with burnin set at 1×10^6 generations,

thinned by 9,999 generations, with 1,000 demes covering the known range of the *P. obsoletus*

complex (Burbrink 2001).

Both EEMS and DAPC find population structure, generally matched the geographic

ranges of *P. alleghaniensis*, *P. spiloides, P. obsoletus*, and *P. bairdi* (Fig. 1 and Fig. S1;

Burbrink, 2001). Total DAPC assignment probabilities were 0.975 (*P. bairdi* = 1.0, *P. obsoletus*

= 0.96; *P. spiloides* = 1.0, and *P. alleghaniensis* = 0.97). Comparing these assignments to those

from SMNF, we found that only *P. alleghaniensis* was misclassified as *P. spiloides* 8.5% of the

time. Similarly, SNMF cross validation predicted four groups (cv = 0.599) as compared to two

(cv=0.627), three (cv=0.607), and five (cv=0.601). These four groups generally match the same

ranges as those found in DAPC. sPCA showed the presence of four groups, with PC1 showing

the separation of groups at the MR and the two lineages east of this river, whereas PC2-5 shows

divergence in all taxa. Importantly, global spatial structures were significant (observed spatial

structure test statistic = 75.47, random expectation = 28.14; $P < 0.009$). Similarly, all three

EEMS runs suggest low migration at the MR and the area separating the *P. bairdi* and *P.*

obsoletus in west Texas. East of the MR, estimated low migration occurred near the Appalachian

Mountains, though this area is a complex mixture of isolation and gene flow (Fig. 1). All three

EEMS generated similar acceptance proportions for all proposal types (12-51%).

1005

Running SNAPP

We set forward and reverse mutation rates to 1. We applied a gamma distribution (α = 2, β =200)

for the speciation rate (λ) prior and the *snapprior* was set to α = 1, β =250, and κ = 1. To test the

four alternative species delimitation models including four taxa, three taxa (collapsing *P.*

bairdi/P. obsoletus or *P. alleghaniensis/P. spiloides*), and two taxa (collapsing *P. bairdi/P.*

obsoletus and *P.alleghaniensis/P. spiloides* we used a stepping-stone analysis with the

PathSampleAnalyser with 72 steps and a chain length of 200000, burnin percentage at 50%, and

a preBurnin of 10,000. We checked for stationarity using Tracer v1.7.1 (Drummond and

Rambaut 2007) determining that estimated sample size (ESS) were > 200 for all parameters.

Running PHRAPL

PHRAPL uses gene-tree distances between simulated and empirical data to rank models using

approximate likelihoods and AICs. To generate the empirical observations we estimated

unrooted gene trees for each loci using IQ-Tree 1.6.12 (Nguyen et al. 2015) and testing all

substitution models per locus prior to estimation. We rooted the trees using midpoint rooting and

performed 100 subsamples of two individuals per population. To reduce model space, we

allowed only one population size parameter (all populations have equal size) and one migration

parameter (all migration rates are equal). We then ran 1000 simulations per model, searching

over a grid of four CollapseStarts values and five MigrationStarts values.

1024

Niche models

We determined if the four taxa show uniquely distinct niches. We quantified niche

similarity using methods introduced in Warren et al. (2008) using Schoener's *D* and Hellinger

distance *I* (Schoener 1968). We used the georeferenced genetic samples here as inputs,

predictions of species identity from DAPC, and 19 bioclimatic variables cropped to the eastern

half of the US (Fick and Hijmans 2017). Estimates of D and I between adjacent species pairs

were calculated and significance of niche identity was derived from 500 random permutations of

lineage identity and re-estimation of the ecological niche model (ENM) using maxent (Phillips

and Dudik 2008) in the R program ENMTools (Warren et al. 2010). Because underlying

differences between lineage-pair regions may not be different from the exact locations of

individuals estimating niche identity we used the background test to produce a null distribution

of ENM symmetrically between a lineage and randomized occurrence points within the adjacent

range of the other adjacent lineage (using a radius of 20km and 1000 points). Therefore, if D and

I from the identity tests were significantly lower than the background test null distribution then

similarity between species, provided that accessible habitat exists in the region, is rejected

(Warren et al. 2008).

Identity tests for niche overlap for both the D and I statistics were significant ($P <$

$1.6 \times 10^{-12} - 4.02 \times 10^{-214}$) for all adjacent pairs. Dissimilarity for the background tests was

significantly lower than the null distributions for all pairs of taxa (P=0.003 to 0.019; Fig. S3).

These results suggests niches overlap between pairs of taxa less than expected by chance.

1045

1046

Spatial genetics using RDA

We predicted genetic distances from environmental, elevation, river, and distance variables using

RDA in the R package Vegan (Dixon 2003), which avoided problems with spatial data using

Mantel tests (Legendre et al. 2015). This method was used to compare results from the ANN

analyses. RDA is an asymmetric canonical analysis here used to determine the significance of

predictor axes (Legendre et al. 2011) at generating predict genetic structure over a landscape

(McGaughran et al. 2014; Noguerales et al. 2016). We ran this using the *capscale* function where

dissimilarity data were ordinated with metric scaling and then passed to RDA. All variable

distances were used to predict genetic distance and significance of each variable was tested using

anova-like permutation tests to examine the joint effect of predictor variables given partial

effects from spatial distance to account for IBD. Also, given that some of these variables might

be correlated, we assessed multicollinearity and reran RDA using the reduced number of

predictor variables. To examine how multicollinearity among variables affects model inference

given linear dependence among these independent variables (De Veaux and Ungar 1994; Hastie

et al. 2001; Shan et al. 2006; Dormann et al. 2013), we used stepwise variable inflation factors in

the package Rnalytica (https://rdrr.io/github/software-analytics/Rnalytica/). The function

stepwise.vif selects non-correlated variables by excluding those with the highest variable

inflation factor above five.

Estimates of accuracy using ANN were high (>90%) for all comparisons and yielded

similar conclusions to RDA. Here, RDA demonstrated that most of the structure for all taxa can

be predicted by the Mississippi River, MIS19 (climate at 787Kya), and elevation (*P*=0.001-

0.002) and these predictors similarly showed 100% variable importance using ANN (Table S3).

Distance also played a role in structuring these data, though they were partialed out in RDA. For

those lineages east of the MR, population structure was predicted by LGM and elevation (RDA:

P = 0.001, ANN: variable importance=100). For those lineages west of the MR, we

demonstrated that MIS19 and elevation structure these populations (RDA: *P*= 0.001, ANN:

variable importance = 100).

1074

Hyrbid zone modeling in HZAR using snaplust and DAPC assignments:

We also fit clines for the same species pairs using individual genetic assignments: pure parental

(0,1), backcross (0.25, 0.75), and F1 (0.5,0.5) and, alternatively, using adegenet prediction

probabilities in the program HZAR v.2-9 (Derryberry et al. 2014). Using the Gaussian cline

model, we estimated the center and width of the cline and determined if these sigmoidal

distributions have significant tails by fitting the following models: 1) no tails, 2) right tail only,

3) left tail only, 4) mirrored tails, and 5) both tails estimated independently (see Derryberry et al.

2014). We fit these models to our data using AICc and ran the MCMC chains for 5×10^6

generations, thinned by 5×10^3 generations, and estimated stationarity using ESS >200 in the R

package CODA (Plummer et al. 2006).

We found agreement for best-fit model between *spaclust* and DAPC assignments and

parameter estimates. For the eastern lineages, HZAR showed strong support for a right-tailed

model ($\Delta AIC_{snapclust}$ =-1.91 　− 　-6.08, ΔAIC_{DAPC} =-31.96 　− 　-107.42), whereas central lineages at

the MR ($\Delta AIC_{snapclust}$ =-4.28 − -8.70, ΔAIC_{DAPC} = -4.26 − -8.869) and western lineages at the

Desert and Edward's Plateau ($\Delta AIC_{snapclust}$ =-5.51 − -221.35, ΔAIC_{DAPC} = -5.51 − -221.35) both

supported a no-tails model (Fig. 6; Table S4 & S5).

1091

1092

Genomics and the ICZN

Another problem likely to be encountered by other researchers using genomic data is that

previously named type specimens at particular type localities may be from regions that show

some admixture. For instance, the type locality of the eastern-most species, *P. alleghaniensis* is

"the summit of the Blue Ridge in Virginia and the Highlands of the Hudson" (Holbrook 1836;

Schultz 1996). Both areas are likely composed of admixed individuals and the International Code

of Zoological Nomenclature (ICZN) forbids naming species based on hybrids (Article

1.3.3;ICZN, 1999) though the intention of this ICZN article did not consider proportions of

admixture. One possible solution would be to use the next available name from Florida, where

no admixture is present (eliminating *P. quadrivittata*), which would be *P.deckerti* (type locality:

Lower Matecumber Key, FL; Brady 1932). Another solution would be to designate a Neotype

from a locality showing no admixture and retain *P. alleghaniensis*. If *P. alleghaniensis/P.*

spiloides were not considered unique, then *P. alleghaniensis* has priority by 18 years (Holbrook

1836; Duméril et al. 1854).

The type locality for the remaining taxa, *P. spiloides (*New Orleans, Louisiana; Duméril

et al. 1854), *P. obsoletus* (On the Missouri River from the Vicinity Isle au Vache to Council

Bluff; Say, 1823), and *P. bairdi* (Fort Davis, Apache Mountains, Jeff Davis Co.: Texas; Yarrow

1880) are all from regions that likely represent individuals not admixed (Fig.1 & 5, Burbrink,

2001). Species delimitation methods, estimates of fixation for loci, and observation that these

four organisms have remained geographically distinct in the face of gene flow throughout the

Pleistocene suggests that these taxa should be considered unique species though hybrid zones

exist. We acknowledge difficulties recognizing the eastern lineages as distinct and could argue

for recognizing them as a single taxon, *P. alleghaniensis*.

1116

1117   Table S1. Priors and summary stats for PipeMaster for the IM, IMD, IMD-LGM models;
1118   estimating migration, timing of divergence, and historical demography.
1119
1120   **IM**

PARAMETER	PRIOR.1	PRIOR.2	DISTRIBUTION	
1	Ne0.pop1	20000	500000	runif
2	Ne0.pop2	20000	5e+05	runif
3	Ne0.pop3	20000	5e+05	runif
4	Ne0.pop4	20000	5e+05	runif
5	mig0.1_2	2	5	runif

6	mig0.1_3	0	0	runif	
7	mig0.1_4	0	0	runif	
8	mig0.2_1	2	5	runif	
9	mig0.2_3	2	5	runif	
10	mig0.2_4	0	0	runif	
11	mig0.3_1	0	0	runif	
12	mig0.3_2	2	5	runif	
13	mig0.3_4	2	5	runif	
14	mig0.4_1	0	0	runif	
15	mig0.4_2	0	0	runif	
16	mig0.4_3	2	5	runif	
17	join1_2	30000	1000000	runif	
18	join3_4	30000	1000000	runif	
19	join2_4	500000	1700000	runif	

1121
1122

IMD

PARAMETER		**PRIOR.1**	**PRIOR.2**	**DISTRIBUTION**	
1	Ne0.pop1	10000	500000	runif	
2	Ne0.pop2	10000	500000	runif	
3	Ne0.pop3	10000	500000	runif	
4	Ne0.pop4	10000	500000	runif	
5	mig0.1_2	2	5	rtnorm	
6	mig0.1_3	0	0	rtnorm	
7	mig0.1_4	0	0	rtnorm	
8	mig0.2_1	2	5	rtnorm	
9	mig0.2_3	2	5	rtnorm	
10	mig0.2_4	0	0	rtnorm	
11	mig0.3_1	0	0	rtnorm	
12	mig0.3_2	2	5	rtnorm	
13	mig0.3_4	2	5	rtnorm	
14	mig0.4_1	0	0	rtnorm	
15	mig0.4_2	0	0	rtnorm	
16	mig0.4_3	2	5	rtnorm	
17	Ne1.pop1	1000	10000	runif	
18	Ne1.pop2	1000	10000	runif	
19	Ne1.pop3	1000	10000	runif	
20	Ne1.pop4	1000	10000	runif	
21	t.Ne1.pop1	3000	170000	runif	
22	t.Ne1.pop2	3000	170000	runif	
23	t.Ne1.pop3	3000	170000	runif	
24	t.Ne1.pop4	3000	170000	runif	
25	join1_2	30000	1000000	runif	
26	join3_4	30000	1000000	runif	
27	join2_4	500000	1700000	runif	

1123
1124   **IMD-LGM**

PARAMETER	PRIOR.1	PRIOR.2	DISTRIBUTION	
1	Ne0.pop1	10000	500000	runif
2	Ne0.pop2	10000	500000	runif
3	Ne0.pop3	10000	500000	runif
4	Ne0.pop4	10000	500000	runif
5	mig0.1_2	2	5	rtnorm
6	mig0.1_3	0	0	rtnorm
7	mig0.1_4	0	0	rtnorm
8	mig0.2_1	2	5	rtnorm
9	mig0.2_3	2	5	rtnorm
10	mig0.2_4	0	0	rtnorm
11	mig0.3_1	0	0	rtnorm
12	mig0.3_2	2	5	rtnorm
13	mig0.3_4	2	5	rtnorm
14	mig0.4_1	0	0	rtnorm
15	mig0.4_2	0	0	rtnorm
16	mig0.4_3	2	5	rtnorm
17	Ne1.pop1	1000	10000	runif
18	Ne1.pop2	1000	10000	runif
19	Ne1.pop3	1000	10000	runif
20	Ne1.pop4	1000	10000	runif
21	t.Ne1.pop1	3000	170000	runif
22	t.Ne1.pop2	3000	170000	runif
23	t.Ne1.pop3	3000	170000	runif
24	t.Ne1.pop4	3000	170000	runif
25	mig1.1_2	0	0	runif
26	mig1.1_3	0	0	runif
27	mig1.1_4	0	0	runif
28	mig1.2_1	0	0	runif
29	mig1.2_3	0	0	runif
30	mig1.2_4	0	0	runif
31	mig1.3_1	0	0	runif
32	mig1.3_2	0	0	runif
33	mig1.3_4	0	0	runif
34	mig1.4_1	0	0	runif
35	mig1.4_2	0	0	runif
36	mig1.4_3	0	0	runif
37	t.mig1.1_2	0	8700	runif
38	t.mig1.1_3	0	8700	runif
39	t.mig1.1_4	0	8700	runif
40	t.mig1.2_1	0	8700	runif
41	t.mig1.2_3	0	8700	runif
42	t.mig1.2_4	0	8700	runif
43	t.mig1.3_1	0	8700	runif
44	t.mig1.3_2	0	8700	runif
45	t.mig1.3_4	0	8700	runif

46	t.mig1.4_1	0	8700	runif
47	t.mig1.4_2	0	8700	runif
48	t.mig1.4_3	0	8700	runif
49	join1_2	30000	1000000	runif
50	join3_4	30000	1000000	runif
51	join2_4	500000	1700000	runif

1125

1126 Table S2 – PHRAPL Results. Available upon request (*open with R read.table() or Excel*).

1127

1128 Table S3. Results from RDA (F and P values) and ANN (variable importance [VAR_IMP]) for
1129 climate, distance, Mississippi River, and elevation variables . Significance of variables after
1130 variable inflation factors (VIF) are considered.
1131

VARIABLES	RDA F VALUE	RDA P VALUE	P_VALUE_VIF	NN_VAR_IMP	NN_VAR_IMP_VIF
ALL TAXA					
MISSISSIPPI	14.1683	0.0001	0.001	100	100
MPW	1.3642	0.06	NA	31	NA
LGM	1.6455	0.015	NA	91	NA
LIG	1.4935	0.024	NA	6	NA
MIS	1.7603	0.005	0.001	81	100
CURRENT	1.8522	0.009	NA	0	NA
ELEVATION	0.9627	0.458	0.002	91	100
DISTANCE	Factored	Factored	Factored	100	100
EAST_OF THE_MS					
LGM	1.7148	0.001	0.001	70	100
LIG	2.0434	0.001	NA	100	NA
MIS	3.1272	0.001	NA	41	NA
CURRENT	1.8641	0.001	NA	98	NA
ELEVATION	1.0464	0.275	0.001	0	100
DISTANCE	Factored	Factored	Factored	100	100
WEST_OF_THE_MS					
LGM	1.4705	0.013	NA	72	NA
LIG	1.1802	0.192	NA	100	NA
MIS	0.9094	0.632	0.001	79	100
CURRENT	0.9583	0.533	NA	91	NA
ELEVATION	0.626	0.984	0.31	90	100
DISTANCE	Factored	Factored	Factored	100	100

1132

1133

Table S4. Results from cline modeling using HZAR for each species pair grouping using
snapclust ancestry estimates

1136

PANTHEROPHIS_ALLEGHANIENSIS_X_P.SPILOIDES

MODEL	Center	Width	Tail_Right	Tail_Left	Tail_Mirror	AICc	ESS
NONE	729 (723-735)	141 (127-157)	-	-	-	-65.158	717-5200
RIGHT	729 (723-736)	141 (126-157)	859 (375-1599)	-	-	-67.067	530-37423
LEFT	729 (723-735)	141 (126-157)	-	1143 (593-1679)	-	-60.988	829-53146
MIRROR	729 (723-735)	141 (126-157)			880 (251-1636)	-61.575	321-53591
BOTH	730 (724-736)	141 (126-157)	769 (147-1562)	1226 (719-1723)	-	-62.885	358-23637

PANTHEROPHIS_SPILOIDES_X_P.OBSOLETUS

MODEL	Center	Width	Tail_Right	Tail_Left	Tail_Mirror	AICc	ESS
NONE	801 (795-807)	100 (90-110)	-	-	-	-535.0162	14399-95534
RIGHT	801 (795-808)	100 (90-110)	1290 (721-1816)	-	-	-530.734	676-79146
LEFT	801 (795-808)	100 (90-110)	-	1276 (709-1835)	-	-530.735	20873-100521
MIRROR	801 (795-807)	100 (90-110)	-	-	1307 (762-1848)	-530.732	10695-120834
BOTH	802 (795-808)	99 (90-110)	1299 (770-1830)	885 (323-1638)	-	-526.3181	612-32572

PANTHEROPHIS_BAIRDI_P.OBSOLETUS

MODEL	Center	Width	Tail_Right	Tail_Left	Tail_Mirror	AICc	ESS
NONE	617 (575-670)	7 (3-13)	-	-	-	-150.277	5-1e5
RIGHT	617 (575-670)	10 (4-17)	1204 (613-1803)	-	-	-144.766	14-109080
LEFT	188 (635-2094)	70 (6-123)	-	647 (440-970)	-	71.072	3-523
MIRROR	642 (590-876)	12 (5-182)	-	-	757(37-1578)	58.219	1-915
BOTH	616 (574-660)	9 (4-16)	1218 (617-1810)	1198 (609-1810)	-	-138.23	9-98456

1152

Table S5. Results from cline modeling using HZAR for each species pair grouping using adegenet ancestry estimates.

1153
1154

1155

PANTHEROPHIS_ALLEGHANIENSIS_X_P. SPILOIDES

MODEL	Center	Width	Tail_Right	Tail_Left	Tail_Mirror	AICc	ESS
NONE	938 (918-954)	552 (507-599)	-	-	-	62.806	1393-2686
RIGHT	753(747-760)	141 (126-157)	53 (45-61)	-	-	-44.61	970-3782
LEFT	912 (883-939)	610 (544-681)	-	188 (59-1433)	-	-60.988	129-514
MIRROR	760 (746-911)	155 (116-502)			91 (66-177)	64.53	6-350
BOTH	750 (740-757)	120 (64-136)	53 (47-60)	819 (278-1559)	-	-12.65	33-1717

PANTHEROPHIS_SPILOIDES_X_P.OBSOLETUS

MODEL	Center	Width	Tail_Right	Tail_Left	Tail_Mirror	AICc	ESS
NONE	797 (790-803)	97 (89-107)	-	-	-	-528.48	12066-16809
RIGHT	797 (790-803)	98 (88-107)	11335 (809-1857)	-	-	-524.22	363-29907
LEFT	797 (790-803)	98 (89-107)	-	1281 (718-1841)	-	-524.22	4278-36224
MIRROR	797 (790-803)	98 (89-107)	-	-	1302 (763-1851)	-524.22	3565-41252
BOTH	797 (791-804)	97 (88-106)	1327 (791-1864)	890 (235-1646)	-	-519.79	184-7761

PANTHEROPHIS_BAIRDI_P.OBSOLETUS

MODEL	Center	Width	Tail_Right	Tail_Left	Tail_Mirror	AICc	ESS
NONE	617 (575-660)	7 (4-13)	-	-	-	-150.277	10-34416
RIGHT	617 (575-661)	10 (5-20)	1204 (595-1793)	-	-	-144.766	6-24620
LEFT	652 (595-1874)	7 (3-81)	-	647 (465-1515)	-	71.072	7-400
MIRROR	617 (574-661)	11 (5-19)	-	-	1219(622-1807)	-144.76	9-38425
BOTH	617 (574-660)	8 (3-13)	1211 (619-1806)	1235 (635-1824)	-	-138.29	2-367770

1156
1157

1186   Fig.S1. Delimiting populations using sparse nonnegative matrix factorization (SNMF) and

1187   spatial Principal Component analysis (sPCA) showing the location of four geographically

1188   distinct lineages.

1189

1199

1200

1201 Fig. S2 – PipeMaster model fit of observed data to simulations under IM, IMD, and IMD-LGM

1202 models for PC1-PC10.

Eastern Lineages
P = 0.0039

Symmetric background test
P.alleghaniensis vs. *P. spiloides*

Central Lineages
P = 0.0039

Western Lineages
P = 0.048

Symmetric background test
P.obsletus vs. *P. bairdi*

1203   Fig. S3 – Symmetric background niche similarity tests for spatially adjacent pairs (eastern,

1204   central, and western lineage comparisons) for the *D* and *I* statistics.

1206   Fig. S4 – Prediction of cline width under neutrality given origin times for each species pair: A)

1207   *Pantherophis alleghaniensis/P. spiloides*, B) *P. spiloides/P.obsoletus*, and C) *P.*

1208   *obsoletus/P.bairdi.*