Development and simulation of fully glycosylated molecular models of ACE2-Fc fusion proteins and their interaction with the SARS-CoV-2 spike protein binding domain

Austen Bernardi [a], Yihan Huang [b], Yongao Xiong [a], Somen Nandi [a,c], Karen A. McDonald [a,c], Roland Faller [a*]

[a] Chemical Engineering, [b] Materials Science and Engineering, [c] Global HealthShare Initiative, UC Davis, Davis, CA, 95616
*corresponding author

Abstract

We develop fully glycosylated computational models of ACE2-Fc fusion proteins which are promising targets for a COVID-19 therapeutic. These models are tested in their interaction with fragments of the binding domain of the Spike Protein S1 of the SARS-CoV-2 virus. We see that some ACE2 glycans interact with S1 and glycans are influencing the conformation of the ACE2 receptor. Additionally, we optimize algorithms for protein glycosylation modelling in order to expedite future model development. All models and algorithms are openly available.

Introduction

As of May 5, 2020 more than 3.6 Million people have been confirmed to be infected with coronavirus disease 2019 (COVID-19) which is caused by the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). This zoonotic pandemic has disrupted society worldwide on a peacetime-unprecedented scale. It also spurred a wide range of scientific endeavors to attack the various aspects of this disease. As the disease spreads there is a critical need for tools that enable the strategic design of biopharmaceutical countermeasures. We are here addressing computationally a molecular approach to aid in the design of a specific class of potential COVID-19 countermeasures.

The genomic sequence of the virus responsible for COVID-19, SARS-CoV-2, was made available in January, 2020 (1), providing critical information on the primary amino acid sequences of potential targets. A particularly important target is the SARS-CoV-2 spike (S) protein that is responsible for the first step in the viral infection process, binding to human cells via the angiotensin converting enzyme 2 (hACE2) receptor. The conserved expression and interaction of ACE2 indicates a wide range of hosts (human and non-human) for SARS-CoV-2 (2). The S protein is a homotrimer with each monomer comprised of 1281 amino acids. The monomers are expected to be highly glycosylated with 22 N-linked glycosylation sequons and 4 O-linked predicted glycosylation sites (3), although only 16 N-linked glycosylation sites were observed in a cryo-EM map of S produced in HEK293F cells (4). Viral coat proteins are often glycosylated which helps pathogens evade the host immune system, modulate access of host proteases, and can enhance cellular attachment through modification of protein structure and/or direct participation at the viral coat protein/cell receptor interface. These glycans are, however, only partially resolved in the experimental structure such that a computational approach is helpful to predict their behavior.

Figure 1. Proposed strategy for SARS-CoV-2 neutralization by ACE2-Fc immunoadhesin. ACE2-Fc binds to the spike (S) protein on the virus and blocks binding to the human cellular receptor ACE2, preventing cellular entrance of SARS-CoV-2.

The human ACE2 protein is a 788 amino acid integral membrane protein with seven N-linked glycosylation sites in the extracellular domain. The binding kinetics between the SARS-CoV-2 spike protein and the hACE2 receptor will depend on the 3D structures of both molecules and their molecular interactions which can be impacted strongly by glycosylation. Knowledge of the spike protein and ACE2 amino acid sequences have led to the commercial availability of the spike protein, ACE2, and various fragments of these, with and without purification/fusion tags, produced recombinantly in various expression hosts including Human embryonic kidney cells (HEK293), insect cells, Chinese Hamster Ovary cells (CHO), and *E. coli.* While the availability of recombinant sources for S and ACE2 glycoproteins have greatly contributed to our understanding of the structure and interactions between these proteins, it is important to recognize that glycosylation of recombinant S and ACE proteins will depend on the host cell (5), the recombinant protein, as well as production (6, 7) and purification methods (8). As molecular models and molecular dynamics simulations can describe the interactions of proteins with glycans and the modulation of protein structure by glycans (9, 10) they are powerful tools to assess the significance of glycosylations on 3D structure and binding site interactions between the SARS-CoV-2 spike protein and the human ACE2 receptor, and to design novel biotherapeutics including optimizing glycosylation.

A promising strategy for the design of COVID-19 therapeutic proteins is a fusion of the extracellular domain of ACE2, the human receptor for SARS-CoV-2, with the Fc region of human immunoglobulin, IgG1, by a linker separating the two domains (11). The neutralization strategy behind ACE2-Fc is shown in Figure

1 (11). This therapeutic design is often called an immunoadhesin, a chimeric protein combining the ligand-binding region of the cell receptor with the constant domain of an immunoglobulin (12). These chimeric molecules form dimers through disulfide bonds between Fc domains; this bivalency increases the affinity for the ligand. The human ACE2 receptor has been shown to be the primary receptor that SARS-CoV-2 uses for entry into and infection of human cells (13, 14), although the binding site is distinct from the catalytic domain of ACE2. With an ACE2-Fc immunoadhesin the ACE2 portion can act as a circulating "bait or decoy" to bind the SARS-CoV-2 spike protein preventing it from entering human cells while the Fc region confers longer circulatory half-life, provides effector functions of the immune system to clear the virus, and allows simple well-established purification using Protein A affinity chromatography. Immunoadhesins are a distinct class of antivirals that can be used prophylactically as well as post-infection and differ from both vaccines and antibodies. Unlike vaccines, they are not intended to elicit an immune response to the viral infection, and unlike antibody therapies, their design is greatly simplified since once the cellular receptor for viral entry is identified the immunoadhesin can be quickly designed and produced.

This strategy also precludes the coronavirus mutating to escape binding with the ACE2-Fc protein, as it would also lose binding affinity to the native ACE2 human cell receptor resulting in a less pathogenic virus. The re-emergent SARS virus in 2003-2004 had a lower affinity for human ACE2 resulting in less severe infection and no secondary transmissions (15). In this strategy the exogenous ACE2 would compensate for decreased ACE2 levels in the lungs during infection, contributing to the treatment of acute respiratory distress, and potentially reduce inflammation and reactive oxygen species in the lung (16). Most importantly, recombinant ACE2-Fc fusion proteins, with native ACE2 catalytic activity as well as a mutant version with lower ACE2 catalytic activity, produced using transfection of HEK293, have shown high affinity binding to the SARS-CoV-2 spike protein and to potently neutralize SARS-CoV-2 *in vitro* (17). Simulations are an ideal tool to optimize such a construct and guide the experimental production of ACE2-Fc.

Glycans are branched, flexible chains of carbohydrates that explore a much wider range of conformations at equilibrium conditions than the protein chain itself as the latter is typically not dynamically changing strongly from its folded form as that would affect its functionality. The faster dynamics of glycans complicates the structural and conformational characterization of glycans in laboratory experiments (18). In atomistic molecular dynamics (MD) simulations, glycan conformations can be straight-forwardly analyzed to obtain structural information, as glycan dynamics are much closer to the computational timescale than the protein dynamics. However, neighboring glycans can interact with each other and essentially lock each other in which can lead to very slow equilibration into the correct conformation (9). Therefore, algorithms are needed to generate realistic glycan configurations as glycans are regularly not fully resolved experimentally. Consequently, only a few simulations of related fully glycosylated proteins available (19-22) among them very recently a proposed glycosylated model of the Spike protein (23) . Our group has made significant progress in the field of glycan modeling in recent years (9, 10, 24).

N-glycan structure is highly heterogeneous, and the relative abundance of glycans depends on the expression system for glycoprotein production. Plant-based transient expression system is well-suited to produce recombinant ACE2-Fc under the current COVID-19 pandemic given its high production speed. Two glycovariants of ACE2-Fc are simulated in this work: one is targeted for ER retention with high mannose glycoforms, and the second is targeted for secretion with plant complex glycoforms. These glycovariants are currently being expressed and purified at UC Davis.

In order to properly understand the interaction between the spike protein and the variant ACE2 receptors bound to its fusion partner the glycosylation of both entities needs to be taken into account. The few existing computational studies of ACE2 interaction with the spike protein we are aware of are using aglycosylated proteins (25-27). Also molecular docking studies have been performed with the older related SARS-CoV virus protein implicated in the SARS epidemic in the early 2000s (28). We develop *in silico* models to predict the 3D structure of two glycosylated ACE2-Fc variants. Additionally, we evaluate the interactions between these two ACE2-Fc variants and a glycosylated spike protein fragment (SpFr) which contains the receptor binding domain of the SARS-CoV-2 spike protein.

Materials and Methods

Sequences and Initial Structure

ACE2-Fc is a homodimer of ACE2 bound to Fc via a synthetic linker. Two sequence variants are used in this work to model ACE2-Fc. Variant 1 (Sequence Seq1 in Supporting Information) contains a C-terminal SEKDEL tag, which is used to express predominantly ER-retained proteins with high-mannose glycoforms in plant-based expression systems. Variant 2 (Sequence Seq2) does not use a C-terminal SEKDEL tag, and will express standard plant glycoforms in plant-based expression systems. Variant 2 has two mutations: H357N and H361N. These mutations are used to deactivate the standard function of ACE2, by preventing the coordination of Zn^{2+} in the active site (17). The ACE2-Fc variants contain 18 disulfide bonds, with 4 of them being interchain. Table ST1 (in Supporting Information) describes the disulfide linkages. The variants also contain 8 N-glycosylation sites per monomer. Each peptidase domain of the ACE2-Fc variants is capable of binding one SARS-CoV-2 SpFr (Sequence Seq3), which contains one glycosylation site. The ACE2-Fc/SpFr structure is depicted in Figure 2. The coordinated Zn^{2+} site is shown in Figure SF1. All 3D structures are rendered with VMD (29).

Figure 2. Infographic of the ACE2-Fc variant 1 homodimer bound to two SpFr.

Simulated systems

ACE2-Fc variant 1 will express high-mannose type glycans when synthesized in plants, while variant 2 will express standard plant glycans. Additionally, SARS-CoV-2 SpFr will exhibit its own glycosylation depending on the host cell; here we assume common mammalian-like glycosylation. Our simulations employed uniform glycosylation profiles to approximate these glycosylation profiles. ACE2-Fc variant 1 is fully glycosylated with Man8 glycans, variant 2 is fully glycosylated with GnGnXF3 glycans the latter is consistent with a very recent experimental study (30), and the SpFr is glycosylated with ANaF[6] (31). Figure 3 shows these glycans using the Consortium of Functional Glycomics nomenclature.

Figure 3. Glycans used in the simulated systems. All structures were built using GlycanBuilder (32).

Four systems containing ACE2-Fc variants were simulated in this work. The first system contains ACE2-Fc variant 1 with Man8 glycans. The second system contains ACE2-Fc variant 2 with GnGnXF[3] glycans. The third and fourth systems are the immunoadhesins of the first and second systems with the SARS-CoV-2 SpFr bound, respectively. The SpFr is always glycosylated with ANaF[6]. Table 1 summarizes the simulated systems.

Table 1. Description of simulated systems.

System ID	ACE2-Fc Sequence	ACE2-Fc Glycosylation	SpFr bound?	ACE2/SpFr ref. PDB
AF[M8]	Variant 1	Man8	no	6M17
AF[GG]	Variant 2	GnGnXF[3]	no	6M17
AF[M8]/SpFr	Variant 1	Man8	yes	6M18
AF[GG]/SpFr	Variant 2	GnGnXF[3]	yes	6M18

This work is largely made possible due to the recent cryogenic electron microscopy work that resolves the ACE2-B[0]AT1 and ACE2-B[0]AT1/SpFr structures, corresponding to PDB codes 6M18 and 6M17, respectively (33). The ACE2 and ACE2/SpFr domains were taken from these structures and fused to the Fc domain (PDB 3SGJ) (34). The Zn^{2+} and coordinating residues in 6M17 and 6M18 are poorly coordinated in these structures. The conformation of these residues along with a coordinating water were instead taken from

PDB 1R42 (35). Histidine protonation states for each system were determined using Reduce (36), and are summarized in Table ST2.

Simulation Procedure

The simulation procedure includes the following steps:

1. Fuse ACE2 with Fc to ACE2-Fc using Modeller (37)
2. Model Zn^{2+} and coordinating residues with MCPB.py (38)
3. Attach glycans using glycam.org (39)
4. Merge structures from 2. and 3. using github.com/austenb28/MCPB_Glycam_merge (40)
5. Generate topology files using AmberTools (38)
6. Convert topology files to Gromacs format using Acpype (24, 41)
7. Perform rigid energy minimization (EM) of glycans using github.com/austenb28/GlyRot (42)
8. Perform EM (emtol = 1000 kJ/mol/nm)
9. Solvate and add ions
10. Perform 10 ps constant volume (NVT) (dt = 0.2 fs, T = 310 K)
11. Perform EM (emtol = 1000 kJ/mol/nm)
12. Perform 100 ps NVT (dt = 2 fs, T = 310 K)
13. Perform 100 ps constant pressure (NPT) (dt = 2 fs, T = 310 K, P = 1 atm)
14. Perform 10 ns production NPT (dt = 2 fs, T = 310 K, P = 1 atm)

Steps 2 and 4 are only required for AF^{M8} and AF^{M8}/SpFr, since they contain the coordinated Zn^{2+} sites. Steps 4 and 7 exhibit new, publicly available software under GNU General Public Licenses. GlyRot has previously been used to model glycosylated butyrylcholinesterase and CMG2-Fc (9, 10). Forcefield topologies were generated using the AmberFF14SB (43) forcefield for protein atoms, the Glycam06-j (44) forcefield for glycan atoms, and the SPC/E water model (45). Steps 8 through 14 are performed using the Gromacs suite (46-48). Systems were solvated in rectangular boxes such the minimum distance between the solute and periodic boundary is 1.2 nm. A rectangular box (for size see Table ST3) was found to be sufficient for 10 ns; longer simulations may require a larger cubic box if the solute rotates significantly. A reduced timestep NVT in step 10 is required to relax solute-solvent contacts. Steps 10 and 11-13 used position restraints on the protein atoms. All simulations were performed at 310 K and 1 atm with the Velocity Rescale thermostat (49) and Parinello-Rahman barostat (50) using time constants of 0.1 ps and 2 ps, respectively. All water bonds are constrained with SETTLE (51); all other bonds are constrained with LINCS (52). A 1 nm cutoff was used for short-range nonbonded interactions. Particle Mesh Ewald was used to model long-range electrostatics (53). Table ST3 contains additional information on system sizes and solvation. Each system was simulated using one compute node with 16 cores. Simulations averaged 2.9 ns/day for systems without the SpFr, and 2.0 ns/day for systems with the SpFr.

Results and Discussion

Figure 3 shows the starting configurations generated as described above (left) and the configurations after MD for 10 ns (right) of all simulated systems. Systems of this size will not fully equilibrate in 10 ns, but evidence of structural stability and concerted motion can still be observed. All systems exhibit shortening of the flexible linker domain between ACE2 and Fc. This shortening can be quantified by analyzing the center of mass distance between the ACE2 and Fc ordered domains, shown in Figure SF2. AF^{M8} and AF^{M8}/SpFr exhibit more shortening of the linker domain than AF^{GG} and AF^{GG}/SpFr, possibly due to the

difference in glycosylation in the Fc domain, which is closest to the flexible linker region. In the AF^{M8}/SpFr and AF^{GG}/SpFr systems, ACE2 glycans near the ACE2/SpFr interface form contacts across the binding interface, indicating that glycosylation may significantly affect binding kinetics. Additionally, glycans on SpFr that were initially oriented away from the protein are reoriented towards ACE2 after 10 ns. The structure of the ordered domains of ACE2-Fc and SpFr appear to retain structural stability. The glycans, on the other hand, show significant reorientation, as the configurational dynamics of glycans is faster than proteins as expected (9).

Figure 3: Initial (left) and 10ns simulated (right) configurations of all systems.

To quantitatively assess structural stability, the root mean square deviation (RMSD) of the ordered domains of ACE2-Fc are shown in Figure 4. All profiles exhibit stable dynamics near or below 2.5 Å. Conformational trending occurs when the RMSD increases from the initial and decreases towards the final. Conformational trending is particularly evident in the Fc domains of AF^{GG} and AF^{GG}/SpFr, while it is particularly absent from the Fc domains of AF^{M8} and AF^{M8}/SpFr. This difference indicates that $GnGnXF^3$ glycosylation in the Fc domain promotes a conformational change in the Fc domain. Backbone RMSD profiles for the SpFr are provided in Figure SF3. The SpFr domains show stable RMSD profiles with significant conformational trending, potentially due to contacts with nearby glycans.

Figure 4. Backbone RMSD profiles of ACE2 (top) and Fc (bottom) ordered domains referenced to initial and final simulation configurations. ACE2: residues 4-707. Fc: residues 745-950. (see SI for sequences)

Conclusions

We have developed fully glycosylated models of ACE2-Fc immunoadhesins with and without interactions to glycosylated SARS-CoV-2 spike protein fragments. Atomic resolution models can be used to help guide the development of ACE2 and/or ACE2-Fc therapeutics for COVID-19 and potentially other coronavirus borne diseases.

We found that glycosylations affects protein structure, and potentially ACE2/SpFr binding. It is not yet clear how important these differences are, but they must be treated carefully when designing ACE2-Fc variants. The work exhibited here provides a direct avenue for collaborations between experimental and computational researchers.

All models developed here are freely available for researchers and future COVID-19 related simulations. The open-source workflows and tools that have been generated for glycoprotein simulations will be useful for general simulations of glycosylated systems. We hope that glycosylation becomes a standard variable in protein molecular simulations in the near future.

Acknowledgements

Simulations were performed on the HPC1 computing cluster at UC Davis. This work was partially supported by a COVID-19 Research Accelerator Funding Track project at UC Davis.

Supporting Information

Supporting information contains all sequence definitions as well as additional analysis. Also, all configurations discussed in this paper are available in pdf format both in supporting information as well as on the webpage of the Faller group https://research.engineering.ucdavis.edu/multiscale. Software tools developed for this project are available on GitHub (40, 42).

References

1.	Wu F, Zhao S, Yu B, Chen Y-M, Wang W, Song Z-G, et al. A new coronavirus associated with human respiratory disease in China. Nature. 2020;579(7798):265-9.
2.	Chen Y, Guo Y, Pan Y, Zhao ZJ. Structure analysis of the receptor binding of 2019-nCoV. Biochemical and Biophysical Research Communications. 2020;525(1):135-40.
3.	Andersen KG, Rambaut A, Lipkin WI, Holmes EC, Garry RF. The proximal origin of SARS-CoV-2. Nature Medicine. 2020;26(4):450-2.
4.	Walls AC, Park Y-J, Tortorici MA, Wall A, McGuire AT, Veesler D. Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. Cell. 2020;181(2):281-92.e6.
5.	Goh JB, Ng SK. Impact of host cell line choice on glycan profile. Critical Reviews in Biotechnology. 2018;38(6):851-67.
6.	Ehret J, Zimmermann M, Eichhorn T, Zimmer A. Impact of cell culture media additives on IgG glycosylation produced in Chinese hamster ovary cells. Biotechnol Bioeng. 2019;116(4):816-30.
7.	Hossler P, Khattak SF, Li ZJ. Optimal and consistent protein glycosylation in mammalian cell culture. Glycobiology. 2009;19(9):936-49.
8.	Zhang P, Woen S, Wang T, Liau B, Zhao S, Chen C, et al. Challenges of glycosylation analysis and control: an integrated approach to producing optimal and consistent therapeutic drugs. Drug Discovery Today. 2016;21(5):740-65.
9.	Bernardi A, Kirschner KN, Faller R. Structural analysis of human glycoprotein butyrylcholinesterase using atomistic molecular dynamics: The importance of glycosylation site ASN241. PLOS ONE. 2017;12(11):e0187994.
10.	Xiong Y, Karuppanan K, Bernardi A, Li Q, Kommineni V, Dandekar AM, et al. Effects of N-Glycosylation on the Structure, Function, and Stability of a Plant-Made Fc-Fusion Anthrax Decoy Protein. Frontiers in Plant Science. 2019;10(768).
11.	Kruse R. Therapeutic strategies in an outbreak scenario to treat the novel coronavirus originating in Wuhan, China [version 2; peer review: 2 approved]. 2020;9(72).
12.	Wycoff K, Maclean J, Belle A, Yu L, Tran Y, Roy C, et al. Anti-infective immunoadhesins from plants. Plant biotechnology journal. 2015;13(8):1078-93.
13.	Yan R, Zhang Y, Li Y, Xia L, Guo Y, Zhou Q. Structural basis for the recognition of the SARS-CoV-2 by full-length human ACE2. 2020:eabb2762.
14.	Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh C-L, Abiona O, et al. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. 2020:eabb2507.
15.	Li W, Zhang C, Sui J, Kuhn JH, Moore MJ, Luo S, et al. Receptor and viral determinants of SARS-coronavirus adaptation to human ACE2. EMBO J. 2005;24(8):1634-43.
16.	Verdecchia P, Cavallini C, Spanevello A, Angeli F. The pivotal link between ACE2 deficiency and SARS-CoV-2 infection. Eur J Intern Med. 2020:S0953-6205(20)30151-5.

17.   Lei C, Fu W, Qian K, Li T, Zhang S, Ding M, et al. Potent neutralization of 2019 novel coronavirus by recombinant ACE2-Ig. bioRxiv. 2020:2020.02.01.929976.

18.   Petrescu AJ, Petrescu SM, Dwek RA, Wormald MR. A statistical analysis of N- and O-glycan linkage conformations from crystallographic data. Glycobiology. 1999;9(4):343-52.

19.   Lu D, Yang C, Liu Z. How Hydrophobicity and the Glycosylation Site of Glycans Affect Protein Folding and Stability: A Molecular Dynamics Simulation. The Journal of Physical Chemistry B. 2012;116(1):390-400.

20.   Jo S, Qi Y, Im W. Preferred conformations of N-glycan core pentasaccharide in solution and in glycoproteins. Glycobiology. 2015;26(1):19-29.

21.   Harbison A, Fadda E. An atomistic perspective on antibody-dependent cellular cytotoxicity quenching by core-fucosylation of IgG1 Fc N-glycans from enhanced sampling molecular dynamics. Glycobiology. 2019.

22.   Yanaka S, Yogo R, Inoue R, Sugiyama M, Itoh SG, Okumura H, et al. Dynamic Views of the Fc Region of Immunoglobulin G Provided by Experimental and Computational Observations. Antibodies. 2019;8(3):39.

23.   Grant OC, Montgomery D, Ito K, Woods RJ. 3D Models of glycosylated SARS-CoV-2 spike protein suggest challenges and opportunities for vaccine development. bioRxiv. 2020:2020.04.07.030445.

24.   Bernardi A, Faller R, Reith D, Kirschner KN. ACPYPE update for nonuniform 1–4 scale factors: Conversion of the GLYCAM06 force field from AMBER to GROMACS. SoftwareX. 2019;10:100241.

25.   Cheng P, Zhengdan Z, Yulong S, Xiaoyu W, Kaijie M, Yanqing Y, et al. Exploring the Binding Mechanism and Accessible Angle of SARS-CoV-2 Spike and ACE2 by Molecular Dynamics Simulation and Free Energy Calculation. chemrxiv. 2020:11877492.v1.

26.   Brielle ES, Schneidman-Duhovny D, Linial M. The SARS-CoV-2 exerts a distinctive strategy for interacting with the ACE2 human receptor. bioRxiv. 2020:2020.03.10.986398.

27.   Nami B, Ghanaeian A, Ghanaeian K, Nami N. The Effect of ACE2 Inhibitor MLN-4760 on the Interaction of SARS-CoV-2 Spike Protein with Human ACE2: A Molecular Dynamics Study. . ChemRxiv. 2020; Preprint:https://doi.org/10.26434/chemrxiv.12159945.v1.

28.   Zhang Y, Zheng N, Hao P, Cao Y, Zhong Y. A molecular docking model of SARS-CoV S1 protein in complex with its receptor, human ACE2. Computational Biology and Chemistry. 2005;29(3):254-7.

29.   Humphrey W, Dalke A, Schulten K. VMD - Visual Molecular Dynamics. J Molec Graphics. 1996;14(1):33-8.

30.   Shajahan A, Archer-hartmann SA, Supekar NT, Gleinich AS, Heiss C, Azadi P. Comprehensive characterization of N- and O- glycosylation of SARS-CoV-2 human receptor angiotensin converting enzyme 2. bioRxiv. 2020:2020.05.01.071688.

31.   Fisher P, Spencer H, Thomas-Oates J, Wood AJ, Ungar D. Modeling Glycan Processing Reveals Golgi-Enzyme Homeostasis upon Trafficking Defects and Cellular Differentiation. Cell Reports. 2019;27(4):1231-43.e6.

32.   Ceroni A, Dell A, Haslam SM. The GlycanBuilder: a fast, intuitive and flexible software tool for building and displaying glycan structures. Source Code for Biology and Medicine. 2007;2(1):3.

33.   Yan R, Zhang Y, Li Y, Xia L, Guo Y, Zhou Q. Structural basis for the recognition of SARS-CoV-2 by full-length human ACE2. Science. 2020;367(6485):1444-8.

34.   Ferrara C, Grau S, Jäger C, Sondermann P, Brünker P, Waldhauer I, et al. Unique carbohydrate–carbohydrate interactions are required for high affinity binding between FcγRIII and antibodies lacking core fucose. Proceedings of the National Academy of Sciences. 2011;108(31):12669-74.

35.   Towler P, Staker B, Prasad SG, Menon S, Tang J, Parsons T, et al. ACE2 X-Ray Structures Reveal a Large Hinge-bending Motion Important for Inhibitor Binding and Catalysis. J Biol Chem. 2004;279(17):17996-8007.

36.   Word JM, Lovell SC, Richardson JS, Richardson DC. Asparagine and glutamine: using hydrogen atom contacts in the choice of side-chain amide orientation11Edited by J. Thornton. Journal of Molecular Biology. 1999;285(4):1735-47.

37.   Webb B, Sali A. Comparative Protein Structure Modeling Using MODELLER. Current Protocols in Bioinformatics. 2016;54(1):5.6.1-5.6.37.

38.   Case DA, Betz RM, Cerutti DS, T.E. Cheatham I, Darden TA, Duke RE, et al. AMBER 2016. University of California, San Francisco; 2016.

39.   Woods Group. GLYCAM Web Athens, GA: Complex Carbohydrate Research Center, University of Georgia; 2005-2020 [Available from: http://glycam.org.

40.   Bernardi A. MCPB_Glycam_merge: GitHub; 2020 [Available from: https://github.com/austenb28/MCPB_Glycam_merge.

41.   Sousa da Silva AW, Vranken WF. ACPYPE - AnteChamber PYthon Parser interfacE. BMC Research Notes. 2012;5(1):367.

42.   Bernardi A. GlyRot: GitHub; 2020 [Available from: https://github.com/austenb28/GlyRot.

43.   Maier JA, Martinez C, Kasavajhala K, Wickstrom L, Hauser KE, Simmerling C. ff14SB: Improving the Accuracy of Protein Side Chain and Backbone Parameters from ff99SB. Journal of Chemical Theory and Computation. 2015;11(8):3696-713.

44.   Kirschner KN, Yongye AB, Tschampel SM, González-Outeiriño J, Daniels CR, Foley BL, et al. GLYCAM06: A generalizable biomolecular force field. Carbohydrates. Journal of Computational Chemistry. 2008;29(4):622-55.

45.   Berendsen HJC, Grigera JR, Straatsma TP. The missing term in effective Pair Potentials. J Phys Chem. 1987;91(24):6269-71.

46.   Pronk S, Pall S, Schulz R, Larsson P, Bjelkmar P, Apostolov R, et al. GROMACS 4.5: a high-throughput and highly parallel open source molecular simulation toolkit. Bioinformatics. 2013;29(7):845-54.

47.   Van Der Spoel D, Lindahl E, Hess B, Groenhof G, Mark AE, Berendsen HJC. GROMACS: Fast, flexible, and free. J Comp Chem. 2005;26(16):1701-18.

48.   Abraham MJ, Murtola T, Schulz R, Páll S, Smith JC, Hess B, et al. GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. SoftwareX. 2015;1-2:19-25.

49.   Bussi G, Donadio D, Parrinello M. Canonical sampling through velocity rescaling. The Journal of Chemical Physics. 2007;126(1):014101.

50.   Parinello M, Rahman A. Strain fluctuations and elastic constants. J Chem Phys. 1982;76(5):2662-6.

51.   Miyamoto S, Kollman PA. Settle: An analytical version of the SHAKE and RATTLE algorithm for rigid water models. Journal of Computational Chemistry. 1992;13(8):952-62.

52.   Hess B, Bekker H, Berendsen HJC, Fraaije JGEM. LINCS: A Linear Constraint Solver for Molecular Simulations. J Comput Chem. 1997;18(12):1463-72.

53.   Essman U, Perela L, Berkowitz ML, Darden HLT, Pedersen LG. A smooth particle mesh Ewald method. J Chem Phys. 1995;103(19):8577 - 92.