Dispersal alters the nature and scope of sexually antagonistic variation

Ewan O Flintham[1], Vincent Savolainen[1,*], and Charles Mullon[2]

[1]Department of Life Sciences, Silwood Park Campus, Imperial College London, Ascot, SL5 7PY United Kingdom
[2]Department of Ecology and Evolution, University of Lausanne, 1015 Lausanne, Switzerland

[*]Correspondence: v.savolainen@imperial.ac.uk

Abstract

Intra-locus sexual conflict, or sexual antagonism, occurs when alleles have opposing fitness effects in the two sexes. Previous theory suggests that sexual antagonism is a driver of genetic variation by generating balancing selection. However, these studies assume that populations are well-mixed, neglecting the effects of spatial subdivision. Here we use mathematical modelling to show that limited dispersal can fundamentally change evolution at sexually antagonistic autosomal and X-linked loci due to inbreeding and sex-specific kin competition. We find that if the sexes disperse at different rates, kin competition within the philopatric sex biases intralocus conflict in favour of the more dispersive sex. Furthermore, kin competition diminishes the strength of balancing selection relative to genetic drift, reducing genetic variation in small subdivided populations. Meanwhile, by decreasing heterozygosity, inbreeding reduces the scope for sexually antagonistic polymorphism due to non-additive allelic effects, and this occurs to a greater extent on the X-chromosome than autosomes. Overall, our results demonstrate that spatial structure is an important factor in predicting where to expect sexually antagonistic alleles. We suggest that observed interspecific and intragenomic variation in sexual antagonism may be explained by sex-specific dispersal ecology and demography.

Keywords: Polymorphism, Balancing selection, Island model, Sex-specific dispersal, Genetic drift.

1   Introduction

Due to the different reproductive roles of males and females, many traits are selected in different directions in the two sexes (Parker et al., 1979). Responding to these divergent selection pressures, however, is not straightforward. Because the sexes share a large part of their genomes and traits are typically determined by the same genes, homologous traits in males and females tend to be genetically correlated (Poissant et al., 2010). Opposing selection pressures on the two sexes therefore lead to a genetic tug-of-war, whereby some alleles are favoured in one sex but disfavoured in the other (Connallon and Clark, 2014). This tug of war, also known as intralocus sexual conflict or sexual antagonism (Bonduriansky and Chenoweth, 2009), can result in balancing selection and the long-term maintenance of polymorphism, within and even between species (e.g. Chippindale et al., 2001; Brommer et al., 2007; Mainguy et al., 2009; Foerster et al., 2007; Lewis et al., 2011; Svensson et al., 2009; Ruzicka et al., 2019; Eyer et al., 2019).

Mathematical population genetics models have helped elucidate the conditions that favour the emergence and maintenance of sexually antagonistic variation (e.g. Owen, 1953; Kidwell et al., 1977; Rice, 1984; Albert and Otto, 2005; Gavrilets and Rice, 2006; Fry, 2010; Connallon and Clark, 2012; Jordan and Charlesworth, 2012; Mullon et al., 2012; Jaquiéry et al., 2013; Jordan and Connallon, 2014; Tazzyman and Abbott, 2015; de Vries and Caswell, 2019). Firstly, models have highlighted the role of dominance, i.e. non-additive effects of alleles, in

facilitating balancing selection at sexually antagonistic loci on both autosomes and sex-chromosomes (Kidwell et al., 1977; Rice, 1984; Fry, 2010; Spencer and Priest, 2016). In line with this prediction for autosomes, a number of empirical studies have found that loci underlying intralocus sexual conflict often exhibit sex differences in dominance that increase the fitness of both male and female heterozygotes (Hager et al., 2008; Barson et al., 2015; Grieshop and Arnqvist, 2018; Pearse et al., 2019). Secondly, stochastic models have emphasised the importance of genetic drift in countering the effects of sexually antagonistic selection in finite populations. In fact, because sexually antagonistic selection is relatively weak, its effects on polymorphism can be easily negated by genetic drift (Connallon and Clark, 2012; Mullon et al., 2012).

A common assumption to previous models of sexual antagonism is that populations are well-mixed (i.e. individuals compete at random, Owen, 1953; Kidwell et al., 1977; Rice, 1984; Albert and Otto, 2005; Gavrilets and Rice, 2006; Fry, 2010; Connallon and Clark, 2012; Jordan and Charlesworth, 2012; Mullon et al., 2012; Jaquiéry et al., 2013; Jordan and Connallon, 2014; Tazzyman and Abbott, 2015; de Vries and Caswell, 2019). In reality, however, most natural populations are spatially structured or subdivided, simply due to the physical constraints of movement that lead to limited dispersal (Clobert et al., 2001). Where dispersal is limited, individuals interacting with one another are more likely to share common alleles than individuals sampled at random in the population (Rousset, 2004). Such structure has important genetic and evolutionary implications as it leads to inbreeding and kin selection (Hamilton, 1964; Frank, 1998; Rousset, 2004; Charlesworth and Charlesworth, 2010). These effects may be particularly relevant for the fate of sexually antagonistic alleles as inbreeding influences the abundance of heterozygotes and hence how dominance affects selection at sexually antagonistic loci (Jordan and Connallon, 2014; Tazzyman and Abbott, 2015). In addition, limited dispersal influences the effective size of a population and thus the strength of genetic drift (Caballero, 1994; Wang and Caballero, 1999; Rousset, 2004). In spite of its potential importance, the consequences of population subdivision for sexually antagonistic variation remain poorly understood.

Here, we investigate the effects of limited dispersal on the segregation of sexually antagonistic alleles. To do this, we extend panmictic population genetics models of intralocus sexual conflict (e.g. Owen, 1953; Kidwell et al., 1977; Fry, 2010) to a subdivided population consisting of patches, or groups, interconnected by dispersal. Our model allows for sex-specific dispersal and demography (i.e. different dispersal rates for males and females, and different numbers of breeding males and females within groups) as reported in many taxa (see Trochet et al., 2016; Li and Kokko, 2019, for reviews). We examine how the interplay between selection, sex-specific ecology and genetic drift influences the segregation of sexually antagonistic alleles at autosomal and sex-linked positions.

2 Model

We consider a dioecious population that is subdivided among patches, each composed of n_{m} male and n_{f} female adults, with the following life-cycle. (i) Within each patch, adults mate at random, producing juveniles of each sex in equal proportion. After reproduction, adults die. (ii) Each juvenile either remains in its natal patch or disperses to another randomly chosen patch (so dispersal is uniform among patches as in Wright, 1931's island model). Juveniles disperse independently from one another, with a sex specific probability, m_{m} and m_{f} in males and females respectively. (iii) Finally, within each patch and each sex, juveniles compete to fill n_{m} male and n_{f} female breeding positions in the local mating pool and become the adults of the next generation.

Following previous authors (e.g. Owen, 1953; Kidwell et al., 1977; Fry, 2010), we consider a genetic locus, either on an autosome or on a sex-chromosome (X in XY species or Z in ZW species), with two segregating alleles, a and A. These alleles have opposing effects on male and female competitiveness (step (iii) of the life-cycle): allele a increases competitiveness when present in males but decreases competitiveness when present in females; conversely, allele A improves female competitiveness but comes at a cost when in males (where the cost to male and female homozygotes carrying detrimental alleles is given by c_{m} and c_{f} respectively, Fig. 1A). So the fittest males are homozygous aa while the fittest females are AA. In Aa heterozygotes, the costs of the antagonistic alleles are mediated by sex-specific dominance coefficients, h_{m} in males and h_{f} in females, allowing for a variety of dominance effects (Fig. 1B-D).

Figure 1: Competitiveness in each sex under different dominance scenarios. A shows male and female phenotypes (competitiveness) according to their genotype. **B**-**D** show how male and female competitiveness varies across genotypes depending on allele dominance effects when costs to homozygotes are equal across the sexes ($c = c_m = c_f$). **B** shows additive allele effects in both sexes (co-dominance, $h_m = h_f = 0.5$), under these conditions heterozygote competitiveness is equal in both sexes and exactly halfway between the homozygote values. **C** shows non-additive effects (incomplete dominance) where the dominant and recessive alleles are the same in both sexes ($h_m = 1 - h_f \neq 0.5$). **D** shows non-additive and sex-specific dominance effects, whereby the detrimental allele in each sex is recessive ($h_m, h_f < 0.5$).

3   Results

3.1   The effects of limited dispersal on sexually antagonistic selection

We first explore the effect of limited dispersal on sexually antagonistic selection by considering mathematically the change in average frequency (weighted by reproductive value e.g. Roze and Rousset, 2003; Van Cleve et al., 2010) of the 'male-detrimental' or 'female-beneficial' allele A. For tractability, our analysis initially assumes weak selection (i.e. small c_m and c_f), large female fecundity (large enough to ignore demographic stochasticity) and an effectively infinite number of patches (see Appendix A for derivation).

3.1.1   Limited dispersal introduces kin selection and inbreeding

We find that the change $\Delta p(p)$ in average frequency p in the population of the A allele at an autosomal locus over one generation is

$$\Delta p(p) = \frac{1}{2} p(1 - p) \left(c_f \times s_f(p) - c_m \times s_m(p) \right), \tag{1}$$

where $p(1 - p)$ is the genetic variance in the population, $c_f \times s_f(p)$ captures (positive) selection on A due to its effect in females, and $-c_m \times s_m(p)$ captures (negative) selection on A due to its effect in males (see Appendices A.2.1-A.2.5 for derivation). Selection through females and males, $s_f(p)$ and $s_m(p)$ both depend on allelic

frequency p and can be decomposed as the sum of three relevant effects

$$s_{\mathrm{f}}(p) = p + F_{\mathrm{IT}}(1-p) + (1-h_{\mathrm{f}})(1-2p)(1-F_{\mathrm{IT}}) \; - (1-m_{\mathrm{f}})^2 \left[K + 2(F_{\mathrm{ST}} - K)\left(p + (1-h_{\mathrm{f}})(1-2p) \right) \right]$$

$$s_{\mathrm{m}}(p) = \underbrace{p + F_{\mathrm{IT}}(1-p)}_{\text{direct effect in } AA} + \underbrace{h_{\mathrm{m}}\,(1-2p)(1-F_{\mathrm{IT}})}_{\text{direct effect in } Aa} - \underbrace{(1-m_{\mathrm{m}})^2 \left[K + 2(F_{\mathrm{ST}} - K)\left(p + h_{\mathrm{m}}(1-2p) \right) \right]}_{\text{kin competition}}, \qquad (2)$$

where F_{IT}, F_{ST} and K denote probabilities that different types of genes are identical-by-descent (IBD) under neutrality (when $c_{\mathrm{f}} = c_{\mathrm{m}} = 0$), capturing various genetic consequences of limited dispersal. F_{IT} is the probability that the two homologous genes at an autosomal locus of one individual are IBD (which corresponds to the absolute coefficient of inbreeding in the infinite island model, Wright, 1922; Caballero, 1994; Rousset, 2004). F_{ST}, meanwhile, is the probability that two autosomal genes in two different juveniles of the same sex and group before dispersal are IBD. This measures the degree of genetic differentiation among groups and is proportional to standard relatedness coefficients (Rousset, 2004). Note that due to random mating within groups, these coefficients are equal in male and female juveniles prior to dispersal. After sex-specific dispersal however, male and female genetic differentiation among groups diverge and become $(1-m_{\mathrm{m}})^2 F_{\mathrm{ST}}$ and $(1-m_{\mathrm{f}})^2 F_{\mathrm{ST}}$, respectively. Finally, K is the probability that the two homologous genes of a juvenile plus a third gene sampled in another juvenile of the same sex and from the same group are all IBD (also prior to dispersal). As such, $(1-m_u)^2 K$ is the probability that among two competing individuals of sex u after dispersal, one is homozygote for a given allele and the other carries at least one IBD copy of that allele.

To understand eq. (2) and how limited dispersal influences selection, note first that in a well-mixed and randomly mating population (where $F_{\mathrm{IT}} = F_{\mathrm{ST}} = K = 0$), selection reduces to: $s_{\mathrm{f}}(p) = p + (1-h_{\mathrm{f}})(1-2p)$ and $s_{\mathrm{m}}(p) = p + h_{\mathrm{m}}(1-2p)$ (as found previously, e.g. eq. 2 of Mullon et al., 2012). In these baseline expressions, the term p captures selection on A due to the effects of the allele on the fitness of its bearer in homozygotes, and $(1-h_{\mathrm{f}})(1-2p)$ or $h_{\mathrm{m}}(1-2p)$ captures selection on A through heterozygotes. Under limited dispersal, we find that the direct effects of A in homozygotes (labelled "direct effect in AA" in eq. 2) increase to $p + F_{\mathrm{IT}}(1-p)$ in both sexes. Conversely, the direct effects of A in heterozygotes (labelled "direct effect in Aa" in eq. 2) decrease to $(1-h_{\mathrm{f}})(1-2p)(1-F_{\mathrm{IT}})$ in females, and to $h_{\mathrm{m}}(1-2p)(1-F_{\mathrm{IT}})$ in males. Selection through homozygotes is therefore more important under limited dispersal. This is because mating within groups leads to inbreeding and therefore a relative excess of homozygotes and a deficit of heterozygotes (according to the inbreeding coefficient F_{IT}).

The remaining terms of eq. (2) (labelled "kin competition") capture a second effect of limited dispersal: that competing individuals are more likely to carry identical gene copies than randomly sampled individuals. Such kin competition effects increase in both sexes with the probabilities that different competing individuals carry IBD genes ($(1-m_{\mathrm{f}})^2 F_{\mathrm{ST}}$ and $(1-m_{\mathrm{f}})^2 K$ in females, and $(1-m_{\mathrm{m}})^2 F_{\mathrm{ST}}$ and $(1-m_{\mathrm{m}})^2 K$ in males). As shown by the negative sign in front of these terms in eq. (2), kin competition effects oppose direct selection effects and therefore weaken the selective force favouring the most adaptive allele in each sex. This is because kin competition results in an individual's reproductive success coming at the expense of genetic relatives, thus reducing the strength of selection on competitiveness.

3.1.2 Additive effects: selection favours the most dispersive sex

Eq. (2) shows that where dispersal differs between males and females ($m_{\mathrm{f}} \neq m_{\mathrm{m}}$), the strength of kin competition differs among the sexes. Sex-specific dispersal therefore biases the intersexual tug-of-war over allele frequency. Specifically, the sex that is the most dispersal limited, and therefore experiences the greatest level of kin competition, is under weaker selection for competitiveness. This can be seen more clearly if we consider an additive locus with equally antagonistic allele effects ($h_{\mathrm{f}} = h_{\mathrm{m}} = 1/2$ and $c_{\mathrm{f}} = c_{\mathrm{m}} = c$). In this case, eqs. (1)-(2) reduce to

$$\Delta p(p) = c F_{\mathrm{ST}}\, p(1-p) \left(1 - \frac{m_{\mathrm{m}} + m_{\mathrm{f}}}{2} \right)(m_{\mathrm{f}} - m_{\mathrm{m}}), \qquad (3)$$

whose sign is determined by the sign of $m_{\mathrm{f}} - m_{\mathrm{m}}$ (for $0 < p < 1$; by contrast $\Delta p(p) = 0$ for all p for an additive locus with equally weak antagonistic effects in well-mixed populations, Fry, 2010). This means that according to the sign of $m_{\mathrm{f}} - m_{\mathrm{m}}$, allele A or allele a fixes. Female beneficial A fixes when $m_{\mathrm{f}} > m_{\mathrm{m}}$, and male beneficial a fixes when $m_{\mathrm{f}} < m_{\mathrm{m}}$. This yields a simple rule for the fate of an additive sexually antagonistic allele: the allele

that benefits the most dispersive sex will fix, and this is because the sex that disperses the most suffers less intense kin competition than the other.

3.1.3   Non-additive effects: subdivision impedes the maintenance of sexually antagonistic polymorphism

In a well-mixed population, selection favours the maintenance of sexually antagonistic polymorphism at an autosomal locus when sex-specific dominance is such that the effect of the detrimental allele in each sex is recessive (i.e. reciprocal invasion, $\Delta p'(0) > 0$ and $\Delta p'(1) < 0$ where $'$ refers to differentiation with respect to p, is favoured when $h_f < 1/2$ and $h_m < 1/2$, Fig. 1D, Kidwell et al., 1977; Fry, 2010). To investigate the effects of population subdivision on sexually antagonistic polymorphisms with sex-specific dominance, we first calculated the relevant coalescence probabilities (F_{IT}, F_{ST}, K) in eqs. (1)-(2) in terms of demographic parameters (see Appendix A.2.6 for derivation). We find that these coalescence probabilities are complicated expressions that depend on sex-specific dispersal (m_f and m_m), as well as the number of reproducing adults in each group (n_f and n_m). When dispersal among groups is low and group size is large, they simplify to

$$F = F_{IT} = F_{ST} = \frac{1}{1 + 8\big(M_m(1-r) + M_f r\big)}$$
$$K = \frac{F}{1 + 4\big(M_m(1-r) + M_f r\big)}, \tag{4}$$

where $M_f = n_f m_f$ and $M_m = n_m m_m$ are the expected numbers of female and male immigrants in a patch at each generation, and $r = n_m/(n_m + n_f)$ is the proportion of males among adults in a patch (i.e. the adult sex ratio; eq. 4 is consistent with eq. 8 in Ramachandran et al., 2008). Unsurprisingly, eq. (4) shows that coalescence becomes more likely when there are fewer immigrants (M_m and M_f are small). But eq. (4) further reveals that coalescence probabilities increase when the adult and immigrant sex ratios are both biased in a similar way (i.e. when $M_m > M_f$ and $r > 0.5$ or when $M_m < M_f$ and $r < 0.5$, see also Fig. 2A for exact F_{ST}). This is because such local bottleneck effects on the non-dispersing sex increases the coalescence of gene lineages through this sex disproportionately.

Figure 2: Coalescence probabilities according to adult and immigrant sex ratio. Contour plots of the probability that two neutral genes carried by two randomly sampled group neighbours are IBD when these genes are: **A.** residing on autosomes; **B.** residing on the X-chromosomes of females; **C.** residing on the X-chromosomes of males (see Appendix A.2.6 and A.3.2 for calculations; other parameters: group size, $n_m + n_f = 10$; number of immigrants per generation, $M_m + M_f = 1$).

Substituting eq. (4) into eqs. (1)-(2), we find that selection favours the maintenance of sexually antagonistic (i.e. $\Delta p'(0) > 0$ and $\Delta p'(1) < 0$) when

$$\frac{h_m}{1 - h_f} + F\frac{1 - h_f - h_m}{1 - h_f} < \frac{c_f}{c_m} < \frac{1 - h_m}{h_f} - F\frac{1 - h_f - h_m}{h_f}. \tag{5}$$

Where $F = 0$, condition eq. (5) reduces to the condition for the maintenance of polymorphism in a well-mixed population (e.g. eq. 3 in Fry, 2010). But as dispersal becomes limited in both sexes (and F increases), eq. (5)

reveals that the conditions for polymorphism become more stringent (as the lower and upper bounds for c_f/c_m respectively increase and decrease with F, see green region Figure 3A Top). Our analysis thus shows that limited dispersal decreases the scope for the maintenance of sexually antagonistic variation. This is because limited dispersal leads to inbreeding so that selection through heterozygotes, which favours sexually antagonistic polymorphism, is less relevant than in well-mixed populations.

Figure 3: Parameters that favour the maintenance of sexually antagonistic variation. Combinations of parameters that lead to either balancing selection (green, when $\Delta p'(0) > 0$ and $\Delta p'(1) < 0$), positive selection for A (red, when $\Delta p'(p) > 0$ for all p), and positive selection for a (blue, when $\Delta p'(p) < 0$ for all p) at **A.** an autosomal locus (computed from eqs. 1-2 with exact coalescence probabilities, see Appendix A.2.6 for calculations of these probabilities) and **B.** X-linked locus (computed from eqs. A-51-A-53 with exact coalescence probabilities, see Appendix A.3.2 for calculations). Top: selection according to ratio of homozygotic effects in females and males (c_m/c_f) and the expected proportion of immigrants in each patch at each generation (($M_m + M_f$)/($n_m + n_f$)); other parameters: $n_m = n_f = 5$, $h_f = h_m = 0.2$, $M_m = M_f$. Bottom: selection according to ratio of homozygotic effects in females and males (c_m/c_f) and the expected proportion of males among immigrants in each patch at each generation ($M_m/(M_m + M_f)$); other parameters: $n_m = n_f = 5$, $h_f = h_m = 0.2$, $M_m + M_f = 0.1$.

We also derived the equilibrium allele frequency when polymorphism is favoured under limited dispersal (see eq. A-43 in Appendix A.2.7). In line with eq. (5), we find limited dispersal increases the equilibrium frequency of the allele with the weaker detrimental effect in homozygotes, irrespective of dominance (e.g. if $c_f > c_m$, the equilibrium frequency of the male-detrimental allele A increases with limited dispersal). In other words, where there is polymorphism, inbreeding leads selection to disproportionately direct equilibrium allele frequencies based on selective effects in homozygotes rather than heterozygotes.

3.1.4 Non-additive effects: sex-biased dispersal changes the nature of sexually antagonistic polymorphism

Analysing eq. (5) further, we find that the nature of sexually antagonistic alleles maintained by balancing selection depends on immigrant sex ratio (Fig. 3A bottom). Specifically, for polymorphism to be maintained under sex-specific dispersal, the cost of the allele antagonistic to the philopatric sex must be higher than the cost of the antagonistic allele in the dispersive sex (e.g. where dispersal is male-biased, $M_m/(M_m + M_f) > 0.5$,

polymorphism is maintained when $c_f > c_m$, see green region Fig. 3A bottom). In line with our analysis of the additive case (eq. 3), this is because sex-specific dispersal leads the philopatric sex to experience stronger kin competition, weakening the selective advantage of the allele favouring competitiveness in that sex relative to the other. As a result, sex-differences in dispersal proclivity lead to different types (i.e. competitiveness effects) of alleles being maintained under balancing selection, thereby influencing the nature of sexually antagonistic polymorphism.

3.1.5 Effects of limited dispersal are similar at X-linked and autosomal loci

We also derived the frequency change of allele A when linked to an X-chromosome (our results also apply to Z-linkage in ZW species; see Appendix A.3.1-A.3.2 for derivations). Our analysis of the resulting allelic dynamics, which can be found in Appendix A.3.3, shows that the effects of limited dispersal on X-linked sexual antagonism are broadly the same as for autosomes. Specifically, inbreeding also leads to more constrained polymorphism conditions on the X-chromosome (Figure 3B top), and sex-specific dispersal also favours adaptation in the more dispersive sex at the expense of the more philopatric one (so that polymorphism occurs only when fitness effects are greater in the philopatric sex, Fig. 3B bottom).

There are nevertheless some discrepancies between autosomal and X-linked positions due to the fact males only carry one copy of the X-chromosome. Firstly, the difference in copy number between the sexes means that male reproductive value for X-linked genes is half that of females (because on average, males transmit half as many X copies to their offspring as females). As a result, selection via female competition is twice as important as via male competition on the X-chromosome. The consequences of this can be most clearly seen by considering an X-linked additive locus ($h_f = 1/2$) with equal allelic effects in males and females ($c_f = c_m = c$). At such a locus, we show in Appendix A.3.3 that the change in p per generation is given by

$$\Delta p(p) = c F_X p(1-p) \left(2m_f(2-m_f) - m_m(2-m_m)\right), \tag{6}$$

where,

$$F_X = \frac{1+r}{1+r+6\left(M_m(1-r)+2M_f r\right)} \tag{7}$$

is the two-way coalescence probability for the X-chromosome (i.e. the probability any two genes sampled from the same sex in the same group before dispersal are IBD, in the limit of low dispersal and large group size, see Appendix A.3.2 for derivation of F_X). Eq. (6) shows that for the male beneficial, female detrimental allele to be favoured by selection, males must be twice as likely to disperse than females in order to compensate for their lower reproductive value for X-linked genes (i.e. $\Delta p(p) < 0$ for all $0 < p < 1$ when $m_m > 2m_f$, by contrast this occurs when $m_m > m_f$ on autosomes, see eq. 3).

Comparing the probabilities that autosomal and X-linked genes are IBD (eq. 4 vs. eq. 7) reveals a second effect of male hemizygosity: that X-linked genes in individuals from the same group tend to coalesce faster than autosomal genes, with the exception of where sex and immigrant ratios are strongly female biased (specifically, $F > F_X$ only when $r < 0.5$ and $M_m/(M_m+M_f) < 2r(2r-1)/(8r^2-5r-1)$, bottom left corner of Fig. 2A). This occurs because there are always fewer X copies than autosomes in a group. Furthermore, like for autosomes, X-linked coalescence probability increases when the adult and immigrant sex ratios are biased in a similar manner, however this occurs to a greater extent on the X when both ratios are biased towards males (Fig. 2B-C for exact probabilities). As a consequence, polymorphism conditions tend to be more restrictive on the X-chromosome than on autosomes with dispersal limitation (Fig. 3A-B top), especially when immigrants are more likely to be males (Fig. 3B bottom).

3.2 The interplay between sexually antagonistic selection and genetic drift under limited dispersal

By assuming an infinite number of groups, our analyses so far have ignored the influence of genetic drift, which is a pervasive evolutionary force in finite populations (Charlesworth and Charlesworth, 2010) and thought to

be especially relevant for intralocus sexual conflict (Connallon and Clark, 2012; Mullon et al., 2012). Predicting how sexually antagonistic variation is maintained under the joint effects of selection and genetic drift in subdivided population, however, is not straightforward. This is because dispersal limitation influences both balancing selection and genetic drift with conflicting implications for polymorphism. On one hand, because limited dispersal leads to kin competition, it weakens the strength of sexually antagonistic balancing selection relative to genetic drift. On the other, genetic isolation between patches increases the effective size (N_e) of a population and therefore reduces the strength of genetic drift relative to selection (in the absence of patch extinction, Caballero, 1994; Wang and Caballero, 1999).

3.2.1 A simulation approach

To investigate the conflicting effects of limited dispersal on the maintenance of polymorphism at sexually antagonistic loci through selection and genetic drift, we used individual-based simulations to track the dynamics of sexually antagonistic alleles at a di-allelic autosomal locus in a finite subdivided population (consisting of 50 groups; see Appendix B for simulation procedure with SLiM 3.3, Haller and Messer, 2019). In addition to allowing us to consider a finite population, using simulations also allows us to relax our assumption of weak selection and thus investigate the balance between sexual antagonism and genetic drift across multiple selection strengths. Starting with equal frequencies of male- and female-beneficial alleles in the population ($p = 0.5$ at Hardy-Weinberg equilibrium), we let simulations run until one allele is fixed and the other is lost (see green trajectories in Fig. 4A for examples of runs). To focus on the interaction between selection and genetic drift, we consider the case where allelic effects are additive ($h_m = h_f = 0.5$) and equal in both sexes ($c = c_m = c_f$), and where the adult sex ratio and dispersal are unbiased ($n_m = n_f$ and $m_m = m_f$). For each simulation replicate, we also tracked the F_{IT} and F_{ST} statistics (purple and orange dots in Fig. 4A for example).

3.2.2 Strong sexual antagonism and frequent dispersal maintains polymorphism in finite populations

In line with eq. (4), we observe that as dispersal probability decreases, inbreeding, F_{IT}, and genetic differentiation among groups, F_{ST}, both increase irrespective of the strength of balancing selection (Fig. 4B). In other words, as dispersal becomes limited we see a reduction in genetic variation within individuals and within groups, as well as an increase in genetic variation between groups. Such conditions are known to increase effective population size (Wang and Caballero, 1999). Accordingly, limited dispersal prolongs the maintenance of neutral genetic variation in our simulations ($c = 0$, black dashed line in Fig. 4C).

For sexually antagonistic loci ($c > 0$), the effect of dispersal on segregation time depends on the strength of selection. Where selection is weak ($c = 0.1$), the impact of limited dispersal on the time that sexually antagonistic polymorphism is maintained for is largely similar to that of a neutral locus, with variation maintained for longer when dispersal is limited (compare black-dashed and pale green in Fig. 4C). Thus, when selection is weak, the effect of limited dispersal on polymorphism is primarily due to its effect on N_e. Where selection is strong ($c = 0.2$), however, the relationship between segregation time and dispersal is quadratic, with polymorphism maintained for long periods when dispersal is either very strong or very weak (dark green curve in Fig. 4C). This suggests that the balance between the opposing effects of population subdivision on polymorphism (increasing N_e on one hand and decreasing the strength of balancing selection on the other) depends on how limited dispersal is. Specifically, when dispersal is mildly limited ($(M_m + M_f)/(n_m + n_f) \gtrsim 0.1$ in our simulations), the effect of reducing selection strength decreases segregation time more than the effect of inflating N_e increases it. Conversely, when dispersal is severely limited ($(M_m + M_f)/(n_m + n_f) \lesssim 0.01$), kin competition renders selection so weak that sexually antagonistic alleles segregate as though they were effectively neutral. In this case, the effect of limited dispersal in increasing N_e outweighs its effect on selection strength.

Figure 4: Allele segregation time and genetic variation in individual-based simulations. A. shows example simulation runs under neutrality ($c = c_m = c_f = 0$) and limited dispersal (when expected proportion of immigrants per group per generation, $(M_m + M_f)/(n_m + n_f) = 0.1$). Green trajectories show allele (A) frequency for different replicate runs, where the time taken for A to fix or be lost representing "segregation time" for that simulation. Purple and orange dots show F_{IT} and F_{ST} respectively at each generation during a simulation, while purple and orange lines represent time-averaged values for these quantities (eqs. B-1 and B-2 in Appendix B for details on how F_{IT} and F_{ST} are calculated). **B.** shows time-averaged F_{IT} and F_{ST} averaged across all (10,000) replicate runs under different levels of limited dispersal (number at top of each panel gives the expected proportion of immigrants in each group), and different strengths of sexually antagonistic selection (white circles represent neutrality $c = 0$, light and dark coloured represent $c = 0.1$ and $c = 0.2$ respectively). **C.** shows segregation time averaged across replicate runs as a function of total fraction of immigrants and selection strength (black dotted line for neutrality, solid light and dark green lines for $c = 0.1$ and $c = 0.2$ respectively). Other parameter values for simulations: $h_f = h_m = 0.5$, $n_m = n_f = 5$ (we also tested the case $n_m = n_f = 10$ and found similar results, Supplementary Figure 1; see Appendix B for more details on simulation procedure).

4 Discussion

Our analyses indicate that limited dispersal significantly impacts evolution at loci under sexual conflict, altering both the nature and scope of sexually antagonistic variation through three main pathways. Firstly, limited dispersal leads to kin competition as individuals of the same group are more likely to carry identical alleles than random individuals. Such competition between relatives weakens the strength of selection on competitive traits and hence of sexually antagonistic balancing selection. In addition, where dispersal is sex-specific, the strength of kin competition is asymmetric between the sexes and consequently biases the intersexual tug-of-war over allele frequencies in favour of the more dispersive sex (Fig. 3 Bottom). Secondly, limited dispersal causes inbreeding and thus reduced heterozygosity. This in turn limits the capacity of dominance effects to

promote polymorphism at sexually antagonistic loci (Fig. 3 Top). Thirdly, in finite populations, limited dispersal simultaneously reduces both the strength of genetic drift and the strength of sexually antagonistic selection. The implications of this for the maintenance of polymorphism in smaller populations depend on the strength of sexual antagonism, with limited dispersal promoting long-term polymorphism when sexual antagonism is weak and impeding it when sexual antagonism is strong (Fig. 4C).

Our results lead to several predictions. Firstly, in populations with sex-specific dispersal, we expect that sexually antagonistic polymorphism is characterised by greater antagonistic costs, and thus greater fitness variation attributable to sexual antagonism, in the non-dispersing sex compared to the dispersing one (e.g. when males disperse more, polymorphism is expected at loci with $c_f > c_m$). Although we are not aware of any empirical work explicitly investigating this, there is potentially relevant data from pedigree studies. In the yellow-pine chipmunk *(Tamias amoenus)*, a species which shows stronger kinship between neighbouring females than males (implying male-biased dispersal, Dobson, 1982; Schulte-Hostedde et al., 2001), females are under directional selection for increased body size, while males are not (Schulte-Hostedde et al., 2002). This suggests that in this species, males are closer to their fitness optimum (so that alleles deviating from this optimum incur a small c_m in males) than females (so that genetic variation for body size in females cause a large c_f). In the great reed warbler (*Acrocephalus arundinaceus*), which shows female-biased dispersal (Hansson et al., 2003), sexual antagonism has been described over wing length (Tarka et al., 2014). In line with our prediction, males are under stronger selection for wing length than females, and show larger additive genetic variance for fitness (Tables 1 and 2 in Tarka et al., 2014). However, the authors of this study also report a negative correlation between a male's wing length and the fitness of his female relatives (Table 3 in Tarka et al., 2014) but not the reverse (a correlation between female wing length and the fitness of male relatives), leading them to suggest that alleles beneficial for males are in fact costlier to females. These conflicting results highlight the difficulty in interpreting the strength of sexually antagonistic allele effects from quantitative genetics studies. More generally, further work is needed to discern the extent to which these observations reflect wider trends, and we suggest future studies consider the effect of dispersal regimes and mating systems on relatedness when formulating hypotheses on sexually antagonistic variation.

Secondly, we expect long-term sexually antagonistic polymorphism to accrue less in dispersal limited populations than in well-mixed populations. Furthermore, in dispersal limited species with small total population size (e.g. small fragmented populations), we expect more sexually antagonistic variation than neutral variation when selection and dispersal are both strong. Broadly speaking, we therefore expect different levels of sexually antagonistic variation between species according to population genetic structure. However, because of the close link between dispersal, mating system and the strength of sex-specific selection (Hamilton, 1967; Greenwood, 1980; Shields, 1987; Perrin and Mazalov, 2000; Li and Holman, 2018), taxa with contrasting spatial structures may also show differences in intensity of intralocus conflict, confounding interspecific patterns in sexually antagonistic variation. Instead, a more fine-scale test would be provided by a species with multiple populations of differing spatial structure but similar levels of sexual antagonism. One such potential study system is the Atlantic salmon (*Salmo salar*), a species lacking sex chromosomes in which QTL analyses have identified a polymorphic sexually antagonistic locus that is maintained through sex-specific dominance (Fig. 1D, Barson et al., 2015). Different salmon populations exhibit different levels of structure, with Scandinavian populations showing strong subdivision (Vähä et al., 2007, 2008) but North American populations appearing relatively panmictic (Wellband et al., 2019). Under the assumption that both salmon populations experience similar patterns of sexually antagonistic selection, this species would be well suited to test the effects of spatial subdivision on the amount of sexually antagonistic variation maintained in a population.

In addition to being useful to understand between-species and between-population patterns of sexually antagonistic variation, our results also have implications for variation within genomes. Much debate in studies of sexual antagonism has centred around the likelihood of different genomic regions accumulating sexually antagonistic polymorphisms, in particular whether such variation should be more common on the X-chromosome or autosomes (Rice, 1984; Chippindale et al., 2001; Gavrilets and Rice, 2006; Fry, 2010; Otto et al., 2011; Mullon et al., 2012; Jordan and Charlesworth, 2012). Previous results have pointed out potential disparities between the X-chromosome and autosomes in their proclivity to harbour sexually antagonistic variation due to male hemizygosity for the X, which leads to differences in selection (i.e. dominance effects act only through females for X-linked genes, Rice, 1984; Gavrilets and Rice, 2006; Fry, 2010; Jordan and Charlesworth, 2012), and differences in the strength of genetic drift (i.e. stronger genetic drift on the X owing to fewer copies

of X-chromosomes than autosomes, Charlesworth et al., 1987; Vicoso and Charlesworth, 2009; Mullon et al., 2012). Our findings indicate that population subdivision influences both of these differences. First, limited dispersal diminishes the importance of selection on heterozygotes and thus dominance effects in driving differences in variation between X-linked and autosomal locations (Fig. 3). Second, sex-specific dispersal and local demography alter F_{ST} for X- and autosomally-linked genes in different ways (Fig. 2) and hence influence the effective population size of the X-chromosome relative to an autosome. In particular, the X-linked and autosomal effective population sizes converge if the adult sex ratio within patches is sufficiently female-biased under limited dispersal (when $r \approx 0.15$, see p. 296 in Ramachandran et al., 2008). Taken together, these results suggest that conditions under which the X-chromosomes and autosomes differ in their propensity to retain sexually antagonistic alleles are less straightforward than predicted by panmictic models. Thus, limited dispersal may help explain the current patchy empirical support for either chromosome region harbouring disproportionate levels of sexually antagonistic variation (e.g. Chippindale et al., 2001; Lucotte et al., 2016; Grieshop and Arnqvist, 2018; Ruzicka et al., 2019).

Our analysis highlights how limited dispersal impacts sexual antagonism by introducing genetic correlations within individuals and groups (via F_{IT} and F_{ST}), skewing genotype frequencies away from Hardy-Weinberg proportions. Alongside limited dispersal, other factors can distort allele distributions away from Hardy Weinberg equilibrium and thus influence selection on sexual antagonistic alleles. In fact, previous theoretical studies have shown that by increasing and decreasing F_{IT} respectively, self-fertilisation (Jordan and Connallon, 2014; Tazzyman and Abbott, 2015) and assortative mating (Arnqvist, 2011; Kasimatis et al., 2019) affect the potential for sexually antagonistic polymorphism. Meanwhile, F_{ST} can be influenced by life-history traits other than dispersal, such as iteroparity (which increases genetic relatedness within groups, Taylor and Irwin, 2000). Interestingly, sex-asymmetries in relatedness also arise in systems where the sexes differ in the extent to which they are itero- or semelparous (e.g. one sex is long-lived while the other is short-lived, Van Cleve et al., 2010), suggesting that sex-differences in life-history strategies may have similar consequences for sexual antagonism as sex-specific dispersal.

Although our study was framed in the context of sexual antagonism, we also expect population subdivision to have implications for other forms of antagonistic evolution, such as antagonistic pleiotropy (where alleles have antagonistic effects on different fitness components, e.g. Williams, 1957; Rose, 1982; Kirkwood and Rose, 1991; Curtsinger et al., 1994; Williams and Day, 2003; Connallon and Clark, 2012). Here, models have also highlighted the importance of component-specific dominance effects, such that antagonistic effects are recessive for each fitness component, in facilitating balancing selection (Rose, 1982, 1985; Curtsinger et al., 1994). But these models assume populations are well-mixed and that genotype frequencies prior to antagonistic selection are at Hardy-Weinberg equilibrium (Rose, 1982, 1985; Curtsinger et al., 1994). From our results, we expect that limited dispersal will also reduce the scope for polymorphism from antagonistic pleiotropy and may therefore influence the evolution of a wide range of traits involved in fitness trade-offs (Roff, 2002), such as viability and expression of secondary sexual characters (Zahavi, 1975; Lande, 1981; Kirkpatrick, 1982; Grafen, 1990; Iwasa et al., 1991; Iwasa and Pomiankowski, 1994), resource acquisition (Van Noordwijk and de Jong, 1986), and reproductive investment and lifespan (Williams, 1957; Williams and Day, 2003; Nussey et al., 2013).

Our model, which has allowed us to produce tractable and intuitive results about intralocus sexual conflict under limited dispersal, relies on several assumptions that are of course unrealistic for many natural populations. In particular, while our model allows us to vary demographic parameters such as adult sex ratio and dispersal regime (immigrant sex ratio and total dispersal rate), these quantities are not free to vary among groups or evolve. In natural populations, however, parameters such as sex ratio and dispersal ratio are correlated by ecological dynamics, and their interdependence will lead them to co-evolve (reviewed in Li and Kokko, 2019). Investigating how such 'eco-evo feedbacks' between local demography and sexual antagonism shape mating system evolution would be an interesting avenue for future work (Svensson, 2019; Giery and Layman, 2019, for reviews; see de Vries and Caswell, 2019 for such an 'eco-evo' model of sexual antagonism under panmixia). In this context, it would also be relevant to relax our assumption that female fecundity is large in order to consider the effects of sex-specific reproductive variance (Mullon et al., 2014), which is common in nature (Clutton-Brock, 2007) and is especially important for selection in subdivided populations (Lehmann and Balloux, 2007).

In conclusion, through its effects on inbreeding, kin competition and genetic drift, limited dispersal qualitatively alters the segregation of sexually antagonistic alleles, narrowing the scope for sexually antagonistic polymorphism and changing the nature of intralocus conflict. Dispersal patterns, and in particular their sex-

specificity, are therefore a relevant consideration for the genetic and ecological conditions expected to lead to sexually antagonistic variation, as well as the nature of the alleles underlying such variation. More broadly, our results reinforce the general notion that spatial demography and population structure are important factors for population genetic dynamics and the maintenance of non-neutral variation (e.g. Wright, 1949; Hamilton, 1964; Hartl et al., 1997; Rousset, 2004; Charlesworth and Charlesworth, 2010).

Acknowledgements

We thank Max Reuter, Stu Wigby, and Carl Mackintosh for providing feedback on a previous version of the manuscript, Henry Barton for discussions on the empirical literature, and Ben Haller for generous advice on developing simulations in SLiM. CM is supported by a SNSF grant (No. PCEFP3181243).

References

Albert, A. Y. and Otto, S. P. (2005). Sexual selection can resolve sex-linked sexual antagonism. *Science*, 310(5745):119–121.

Arnqvist, G. (2011). Assortative mating by fitness and sexually antagonistic genetic variation. *Evolution: International Journal of Organic Evolution*, 65(7):2111–2116.

Barson, N. J., Aykanat, T., Hindar, K., Baranski, M., Bolstad, G. H., Fiske, P., Jacq, C., Jensen, A. J., Johnston, S. E., Karlsson, S., et al. (2015). Sex-dependent dominance at a single locus maintains variation in age at maturity in salmon. *Nature*, 528(7582):405.

Bonduriansky, R. and Chenoweth, S. F. (2009). Intralocus sexual conflict. *Trends in ecology & evolution*, 24(5):280–288.

Brommer, J. E., Kirkpatrick, M., Qvarnström, A., and Gustafsson, L. (2007). The intersexual genetic correlation for lifetime fitness in the wild and its implications for sexual selection. *PLoS one*, 2(8):e744.

Caballero, A. (1994). Developments in the prediction of effective population size. *Heredity*, 73(6):657.

Charlesworth, B. and Charlesworth, D. (2010). *Elements of Evolutionary Genetics*. Roberts and Company Publishers.

Charlesworth, B., Coyne, J. A., and Barton, N. H. (1987). The relative rates of evolution of sex chromosomes and autosomes. *The American Naturalist*, 130(1):113–146.

Chippindale, A. K., Gibson, J. R., and Rice, W. R. (2001). Negative genetic correlation for adult fitness between sexes reveals ontogenetic conflict in drosophila. *Proceedings of the National Academy of Sciences*, 98(4):1671–1675.

Clobert, J., Danchin, E., Dhondt, A. A., and Nichols, J. D., editors (2001). *Dispersal*. Oxford University Press.

Clutton-Brock, T. (2007). Sexual selection in males and females. *Science*, 318(5858):1882–1885.

Connallon, T. and Clark, A. G. (2012). A general population genetic framework for antagonistic selection that accounts for demography and recurrent mutation. *Genetics*, 190(4):1477–1489.

Connallon, T. and Clark, A. G. (2014). Evolutionary inevitability of sexual antagonism. *Proceedings of the Royal Society B: Biological Sciences*, 281(1776):20132123.

Curtsinger, J. W., Service, P. M., and Prout, T. (1994). Antagonistic pleiotropy, reversal of dominance, and genetic polymorphism. *The American Naturalist*, 144(2):210–228.

de Vries, C. and Caswell, H. (2019). Selection in two-sex stage-structured populations: Genetics, demography, and polymorphism. *Theoretical population biology*, 130:160–169.

Dobson, F. S. (1982). Competition for mates and predominant juvenile male dispersal in mammals. *Animal behaviour*, 30(4):1183–1192.

Eyer, P.-A., Blumenfeld, A. J., and Vargo, E. L. (2019). Sexually antagonistic selection promotes genetic divergence between males and females in an ant. *Proceedings of the National Academy of Sciences*, 116(48):24157–24163.

Foerster, K., Coulson, T., Sheldon, B. C., Pemberton, J. M., Clutton-Brock, T. H., and Kruuk, L. E. (2007). Sexually antagonistic genetic variation for fitness in red deer. *Nature*, 447(7148):1107.

Frank, S. A. (1998). *Foundations of Social Evolution*. Monogr. Behav. Ecol. Princeton University Press, Princeton.

Fry, J. D. (2010). The genomic location of sexually antagonistic variation: some cautionary comments. *Evolution: International Journal of Organic Evolution*, 64(5):1510–1516.

Gavrilets, S. and Rice, W. R. (2006). Genetic models of homosexuality: generating testable predictions. *Proceedings of the Royal Society B: Biological Sciences*, 273(1605):3031–3038.

Giery, S. T. and Layman, C. A. (2019). Ecological consequences of sexually selected traits: An eco-evolutionary perspective. *The Quarterly Review of Biology*, 94(1):29–74.

Grafen, A. (1990). Biological signals as handicaps. *Journal of theoretical biology*, 144(4):517–546.

Greenwood, P. J. (1980). Mating systems, philopatry and dispersal in birds and mammals. *Animal behaviour*, 28(4):1140–1162.

Grieshop, K. and Arnqvist, G. (2018). Sex-specific dominance reversal of genetic variation for fitness. *PLoS biology*, 16(12):e2006810.

Hager, R., Cheverud, J. M., Leamy, L. J., and Wolf, J. B. (2008). Sex dependent imprinting effects on complex traits in mice. *BMC evolutionary biology*, 8(1):303.

Haller, B. C. and Messer, P. W. (2019). Slim 3: forward genetic simulations beyond the wright–fisher model. *Molecular biology and evolution*, 36(3):632–637.

Hamilton, W. D. (1964). The genetical evolution of social behaviour. ii. *Journal of theoretical biology*, 7(1):17–52.

Hamilton, W. D. (1967). Extraordinary sex ratios. *Science*, 156(3774):477–488.

Hansson, B., Bensch, S., and Hasselquist, D. (2003). A new approach to study dispersal: immigration of novel alleles reveals female-biased dispersal in great reed warblers. *Molecular Ecology*, 12(3):631–637.

Hartl, D. L., Clark, A. G., and Clark, A. G. (1997). *Principles of population genetics*, volume 116. Sinauer associates Sunderland, MA.

Iwasa, Y. and Pomiankowski, A. (1994). The evolution of mate preferences for multiple sexual ornaments. *Evolution*, 48(3):853–867.

Iwasa, Y., Pomiankowski, A., and Nee, S. (1991). The evolution of costly mate preferences ii. the "handicap" principle. *Evolution*, 45(6):1431–1442.

Jaquiéry, J., Rispe, C., Roze, D., Legeai, F., Le Trionnaire, G., Stoeckel, S., Mieuzet, L., Da Silva, C., Poulain, J., Prunier-Leterme, N., Ségurens, B., Tagu, D., and Simon, J.-C. (2013). Masculinization of the X Chromosome in the Pea Aphid. *PLoS Genet*, 9(8):e1003690–15.

Jordan, C. Y. and Charlesworth, D. (2012). The potential for sexually antagonistic polymorphism in different genome regions. *Evolution: International Journal of Organic Evolution*, 66(2):505–516.

Jordan, C. Y. and Connallon, T. (2014). Sexually antagonistic polymorphism in simultaneous hermaphrodites. *Evolution*, 68(12):3555–3569.

Kasimatis, K. R., Ralph, P. L., and Phillips, P. C. (2019). Limits to genomic divergence under sexually antagonistic selection. *G3: Genes, Genomes, Genetics*, 9(11):3813–3824.

Kidwell, J., Clegg, M., Stewart, F., and Prout, T. (1977). Regions of stable equilibria for models of differential selection in the two sexes under random mating. *Genetics*, 85(1):171–183.

Kirkpatrick, M. (1982). Sexual selection and the evolution of female choice. *Evolution*, 36(1):1–12.

Kirkwood, T. B. and Rose, M. R. (1991). Evolution of senescence: late survival sacrificed for reproduction. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 332(1262):15–24.

Lande, R. (1981). Models of speciation by sexual selection on polygenic traits. *Proceedings of the National Academy of Sciences*, 78(6):3721–3725.

Lehmann, L. and Balloux, F. (2007). Natural selection on fecundity variance in subdivided populations: kin selection meets bet hedging. *Genetics*, 176(1):361–377.

Lewis, Z., Wedell, N., and Hunt, J. (2011). Evidence for strong intralocus sexual conflict in the indian meal moth, plodia interpunctella. *Evolution*, 65(7):2085–2097.

Li, X.-Y. and Holman, L. (2018). Evolution of female choice under intralocus sexual conflict and genotype-by-environment interactions. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 373(1757):20170425.

Li, X.-Y. and Kokko, H. (2019). Sex-biased dispersal: a review of the theory. *Biological Reviews*, 94(2):721–736.

Lucotte, E. A., Laurent, R., Heyer, E., Segurel, L., and Toupance, B. (2016). Detection of allelic frequency differences between the sexes in humans: a signature of sexually antagonistic selection. *Genome biology and evolution*, 8(5):1489–1500.

Mainguy, J., Coté, S. D., Festa-Bianchet, M., and Coltman, D. W. (2009). Father–offspring phenotypic correlations suggest intralocus sexual conflict for a fitness-linked trait in a wild sexually dimorphic mammal. *Proceedings of the Royal Society B: Biological Sciences*, 276(1675):4067–4075.

Mullon, C., Pomiankowski, A., and Reuter, M. (2012). The effects of selection and genetic drift on the genomic distribution of sexually antagonistic alleles. *Evolution: International Journal of Organic Evolution*, 66(12):3743–3753.

Mullon, C., Reuter, M., and Lehmann, L. (2014). The evolution and consequences of sex-specific reproductive variance. *Genetics*, 196(1):235–252.

Nussey, D. H., Froy, H., Lemaitre, J.-F., Gaillard, J.-M., and Austad, S. N. (2013). Senescence in natural populations of animals: widespread evidence and its implications for bio-gerontology. *Ageing research reviews*, 12(1):214–225.

Otto, S. P., Pannell, J. R., Peichel, C. L., Ashman, T.-L., Charlesworth, D., Chippindale, A. K., Delph, L. F., Guerrero, R. F., Scarpino, S. V., and McAllister, B. F. (2011). About par: the distinct evolutionary dynamics of the pseudoautosomal region. *Trends in Genetics*, 27(9):358–367.

Owen, A. (1953). A genetical system admitting of two distinct stable equilibria under natural selection. *Heredity*, 7(1):97.

Parker, G. et al. (1979). Sexual selection and sexual conflict. *Sexual selection and reproductive competition in insects*, 123:166.

Pearse, D. E., Barson, N. J., Nome, T., Gao, G., Campbell, M. A., Abadía-Cardoso, A., Anderson, E. C., Rundio, D. E., Williams, T. H., Naish, K. A., et al. (2019). Sex-dependent dominance maintains migration supergene in rainbow trout. *Nature ecology & evolution*, pages 1–12.

Perrin, N. and Mazalov, V. (2000). Local competition, inbreeding, and the evolution of sex-biased dispersal. *The American Naturalist*, 155(1):116–127.

Poissant, J., Wilson, A. J., and Coltman, D. W. (2010). Sex-specific genetic variance and the evolution of sexual dimorphism: a systematic review of cross-sex genetic correlations. *Evolution: International Journal of Organic Evolution*, 64(1):97–107.

Ramachandran, S., Rosenberg, N. A., Feldman, M. W., and Wakeley, J. (2008). Population differentiation and migration: coalescence times in a two-sex island model for autosomal and x-linked loci. *Theoretical population biology*, 74(4):291–301.

Rice, W. R. (1984). Sex chromosomes and the evolution of sexual dimorphism. *Evolution*, 38(4):735–742.

Roff, D. A. (2002). *Life history evolution.* Sinauer associates Sunderland, MA.

Rose, M. R. (1982). Antagonistic pleiotropy, dominance, and genetic variation. *Heredity*, 48(1):63–78.

Rose, M. R. (1985). Life history evolution with antagonistic pleiotropy and overlapping generations. *Theoretical Population Biology*, 28(3):342–358.

Rousset, F. (2004). *Genetic Structure and Selection in Subdivided Populations.* Princeton University Press, Princeton, NJ.

Roze, D. and Rousset, F. (2003). Selection and drift in subdivided populations: a straightforward method for deriving diffusion approximations and applications involving dominance, selfing and local extinctions. *Genetics*, 165(4):2153–2166.

Ruzicka, F., Hill, M. S., Pennell, T. M., Flis, I., Ingleby, F. C., Mott, R., Fowler, K., Morrow, E. H., and Reuter, M. (2019). Genome-wide sexually antagonistic variants reveal long-standing constraints on sexual dimorphism in fruit flies. *PLoS biology*, 17(4):e3000244.

Schulte-Hostedde, A. I., Gibbs, H. L., and Millar, J. S. (2001). Microgeographic genetic structure in the yellow-pine chipmunk (tamias amoenus). *Molecular Ecology*, 10(7):1625–1631.

Schulte-Hostedde, A. I., Millar, J. S., and Gibbs, H. L. (2002). Female-biased sexual size dimorphism in the yellow-pine chipmunk (tamias amoenus): sex-specific patterns of annual reproductive success and survival. *Evolution*, 56(12):2519–2529.

Shields, W. M. (1987). Dispersal and mating systems: investigating their causal connections. *Mammalian dispersal patterns: The effects of social structure on population genetics*, pages 3–24.

Spencer, H. G. and Priest, N. K. (2016). The evolution of sex-specific dominance in response to sexually antagonistic selection. *The American Naturalist*, 187(5):658–666.

Svensson, E. I. (2019). Eco-evolutionary dynamics of sexual selection and sexual conflict. *Functional Ecology*, 33(1):60–72.

Svensson, E. I., McAdam, A. G., and Sinervo, B. (2009). Intralocus sexual conflict over immune defense, gender load, and sex-specific signaling in a natural lizard population. *Evolution: International Journal of Organic Evolution*, 63(12):3124–3135.

Tarka, M., Åkesson, M., Hasselquist, D., and Hansson, B. (2014). Intralocus sexual conflict over wing length in a wild migratory bird. *The American Naturalist*, 183(1):62–73.

Taylor, P. D. and Irwin, A. J. (2000). Overlapping generations can promote altruistic behavior. *Evolution*, 54(4):1135–1141.

Tazzyman, S. J. and Abbott, J. K. (2015). Self-fertilization and inbreeding limit the scope for sexually antagonistic polymorphism. *J. Evol. Biol.*, 28(3):723–729.

Trochet, A., Courtois, E. A., Stevens, V. M., Baguette, M., Chaine, A., Schmeller, D. S., Clobert, J., and Wiens, J. J. (2016). Evolution of sex-biased dispersal. *The Quarterly Review of Biology*, 91(3):297–320.

Vähä, J.-P, Erkinaro, J., Niemelä, E., and Primmer, C. R. (2007). Life-history and habitat features influence the within-river genetic structure of atlantic salmon. *Molecular ecology*, 16(13):2638–2654.

Vähä, J.-P, Erkinaro, J., Niemelä, E., and Primmer, C. R. (2008). Temporally stable genetic structure and low migration in an atlantic salmon population complex: implications for conservation and management. *Evolutionary Applications*, 1(1):137–154.

Van Cleve, J., Feldman, M. W., and Lehmann, L. (2010). How demography, life history, and kinship shape the evolution of genomic imprinting. *The American Naturalist*, 176(4):440–455.

Van Noordwijk, A. J. and de Jong, G. (1986). Acquisition and allocation of resources: their influence on variation in life history tactics. *The American Naturalist*, 128(1):137–142.

Vicoso, B. and Charlesworth, B. (2009). Effective population size and the faster-x effect: an extended model. *Evolution: International Journal of Organic Evolution*, 63(9):2413–2426.

Wang, J. and Caballero, A. (1999). Developments in predicting the effective size of subdivided populations. *Heredity*, 82(2):212–226.

Wellband, K., Mérot, C., Linnansaari, T., Elliott, J., Curry, R. A., and Bernatchez, L. (2019). Chromosomal fusion and life history-associated genomic variation contribute to within-river local adaptation of atlantic salmon. *Molecular ecology*, 28(6):1439–1459.

Williams, G. C. (1957). Pleiotropy, natural selection, and the evolution of senescence. *Evolution*, 11(4):398–411.

Williams, P. D. and Day, T. (2003). Antagonistic pleiotropy, mortality source interactions, and the evolutionary theory of senescence. *Evolution*, 57(7):1478–1488.

Wright, S. (1922). Coefficients of inbreeding and relationship. *The American Naturalist*, 56(645):330–338.

Wright, S. (1931). Evolution in Mendelian Populations. *Genetics*, 16(2):290.

Wright, S. (1949). The genetical structure of populations. *Annals of eugenics*, 15(1):323–354.

Zahavi, A. (1975). Mate selection—a selection for a handicap. *Journal of theoretical Biology*, 53(1):205–214.

Appendix

A Allele frequency change in a subdivided dioecious population

In this appendix, we derive the change in frequency over one generation for an allele at an autosomal and sex-linked locus, assuming that selection is weak.

A.1 Model

We first provide further details on our modelling assumptions, which are necessary for our derivations. Recall we consider a dioecious population that is divided among a large number n_g of groups, each composed of n_m and n_f adult males and females, respectively. The life-cycle of this population is as follows. 1) Adults mate randomly within their group, females then produce a large number k of male and female juveniles in equal proportions (i.e. unbiased sex ratio at birth). 2) Independently of one another, each juvenile either remains in its natal group, or disperses to another randomly chosen group. 3) Adults die, and male and female juveniles compete to fill the n_m and n_f open breeding spots in each group (so competition occurs within the sexes and within groups).

Each individual carries a quantitative phenotype that is genetically-encoded and that determines the competitiveness of a juvenile to settle in a group (step 3 of the life cycle). We will separately consider evolution at two loci that influence this phenotype: at an autosomal and at a sex-linked locus (on the X-chromosome in an XY species or the Z chromosome in a ZW species). In either case, two alleles A and a with sexually antagonistic effects segregate in the population. We are interested in deriving the change in frequency in the whole population of the allele A. For mathematical convenience, the population is censused in juveniles before dispersal (between steps 1 and 2 of the life cycle).

A.2 Autosomal locus

A.2.1 Genotypes

To denote the genotype of males and females, we let $p_{ij1}^u \in \{0,1\}$ and $p_{ij2}^u \in \{0,1\}$ be the frequency of A at the maternally and paternally inherited locus, respectively, of juvenile $j \in \{1,\ldots,n_\mathrm{f}k/2\}$ of sex $u \in \{\mathrm{m},\mathrm{f}\}$ residing in group $i \in \{1,\ldots,n_\mathrm{g}\}$ (recall that k is the total fecundity of a female, so that within a group there are $n_\mathrm{f}k/2$ male and $n_\mathrm{f}k/2$ female juveniles after reproduction and before dispersal). The frequency of A in such an individual then is

$$p_{ij}^u = \frac{p_{ij1}^u + p_{ij2}^u}{2} \in \{0,1/2,1\}. \tag{A-1}$$

For example, if male 2 in group 1 has inherited A from its mother and a from its father, then $p_{121}^\mathrm{m} = 1$, $p_{122}^\mathrm{m} = 0$, and $p_{12}^\mathrm{m} = 1/2$. The average frequency of A in individuals of sex u can then be defined as

$$p^u = \overline{p_{ij}^u}, \tag{A-2}$$

where the upper bar means the average over all juveniles j and over all groups i throughout.

A.2.2 Phenotypes

The evolving phenotype determines the competitiveness of male and female juveniles. We assume that baseline competitiveness is 1 and that allele A is detrimental to males (causing a decrease in competitiveness δc_m in homozygotes and $\delta h_\mathrm{m} c_\mathrm{m}$ in heterozygotes, Table A-1). Conversely, allele a is detrimental to females causing a

	aa	Aa	AA
male phenotype	1	$1 - \delta h_{\mathrm{m}} c_{\mathrm{m}}$	$1 - \delta c_{\mathrm{m}}$
female phenotype	$1 - \delta c_{\mathrm{f}}$	$1 - \delta h_{\mathrm{f}} c_{\mathrm{f}}$	1

Table A-1: Male and female phenotypes according to genotype.

decrease in competitiveness δc_{f} in homozygotes and $\delta h_{\mathrm{f}} c_{\mathrm{f}}$ in heterozygotes, Table A-1). Parameters $c_{\mathrm{m}} > 0$ and $c_{\mathrm{f}} > 0$ respectively capture the cost of the detrimental allele in male and female homozygotes ($\delta > 0$ tunes the strength of selection), while parameters h_{m} and h_{f} determine the dominance or penetrance of the detrimental allele in males and females, respectively. We can then write the phenotype z_{ij}^{u} of juvenile j of sex u that resides in group i in terms of the frequency of allele A in this individual as

$$z_{ij}^{\mathrm{m}} = 1 - \delta c_{\mathrm{m}} \left(2 h_{\mathrm{m}} p_{ij}^{\mathrm{m}} + (1 - 2 h_{\mathrm{m}}) p_{ij1}^{\mathrm{m}} p_{ij2}^{\mathrm{m}} \right)$$
$$z_{ij}^{\mathrm{f}} = 1 + \delta c_{\mathrm{f}} \left(-1 + 2(1 - h_{\mathrm{f}}) p_{ij}^{\mathrm{f}} + (1 - 2(1 - h_{\mathrm{f}})) p_{ij1}^{\mathrm{f}} p_{ij2}^{\mathrm{f}} \right), \tag{A-3}$$

according to Table A-1.

A.2.3 Weighted allele frequency change

To track the dynamics of the frequencies of the A allele in the population, we consider the dynamics of a reproductive-value weighted average p of male and female frequencies,

$$p = \alpha p^{\mathrm{f}} + (1 - \alpha) p^{\mathrm{m}}, \tag{A-4}$$

where α and $1 - \alpha$ are the female and male reproductive values, respectively (e.g. Roze and Rousset, 2003). This weighting ensures that in the absence of selection (when $\delta = 0$), the change in p is zero. For an autosomal locus, males and females have equal reproductive values: $\alpha = 1/2$.

Following Roze and Rousset (2003) (adapting their eq. 35 for an autosomal locus) and Van Cleve et al. (2010) (their eq. 2), the change over one iteration of the life-cycle in weighted allele frequency change is

$$\Delta p = \frac{1}{n_{\mathrm{g}}} \sum_{i=1}^{n_{\mathrm{g}}} \frac{1}{n_{\mathrm{f}} k/2} \left[\alpha \left(\sum_{j=1}^{n_{\mathrm{f}} k/2} w_{ij}^{\mathrm{f} \to \mathrm{f}} \frac{p_{ij}^{\mathrm{f}}}{2} + \sum_{j=1}^{n_{\mathrm{f}} k/2} w_{ij}^{\mathrm{m} \to \mathrm{f}} \frac{p_{ij}^{\mathrm{m}}}{2} \right) + (1 - \alpha) \left(\sum_{j=1}^{n_{\mathrm{f}} k/2} w_{ij}^{\mathrm{f} \to \mathrm{m}} \frac{p_{ij}^{\mathrm{f}}}{2} + \sum_{j=1}^{n_{\mathrm{f}} k/2} w_{ij}^{\mathrm{m} \to \mathrm{m}} \frac{p_{ij}^{\mathrm{m}}}{2} \right) \right] - p, \tag{A-5}$$

where $w_{ij}^{u \to v}$ is the expected number of juveniles of sex v produced by juvenile j of sex u in group i (before dispersal) and the division of individual gene frequencies by 2 reflects that a gene has a probability of $1/2$ to be passed to an offspring at an autosomal locus (with no segregation bias). There are thus four relevant fitness functions: $w_{ij}^{\mathrm{f} \to \mathrm{f}}$, $w_{ij}^{\mathrm{m} \to \mathrm{f}}$, $w_{ij}^{\mathrm{f} \to \mathrm{m}}$, and $w_{ij}^{\mathrm{m} \to \mathrm{m}}$. We consider these fitness functions in more detail later (section A.2.5). But first note that when sex ratio at birth is equal, a juvenile of either sex is expected to produce the same numbers of male and female juvenile so that $w_{ij}^{\mathrm{f} \to \mathrm{f}} = w_{ij}^{\mathrm{f} \to \mathrm{m}} = w_{ij}^{\mathrm{f}}/2$ and $w_{ij}^{\mathrm{m} \to \mathrm{f}} = w_{ij}^{\mathrm{m} \to \mathrm{m}} = w_{ij}^{\mathrm{m}}/2$, where w_{ij}^{f} and w_{ij}^{m} are the expected total numbers of juveniles produces by a female and male juveniles, respectively. Substituting for these total fitness expressions and for reproductive value $\alpha = 1/2$ into eq. (A-5), we obtain

$$\Delta p = \frac{1}{4 n_{\mathrm{g}}} \sum_{i=1}^{n_{\mathrm{g}}} \frac{1}{n_{\mathrm{f}} k/2} \left[\sum_{j=1}^{n_{\mathrm{f}} k/2} w_{ij}^{\mathrm{f}} p_{ij}^{\mathrm{f}} + \sum_{j=1}^{n_{\mathrm{f}} k/2} w_{ij}^{\mathrm{m}} p_{ij}^{\mathrm{m}} \right] - p \tag{A-6}$$

for the change in weighted allele frequency.

A.2.4 Selection gradient

Next, we connect individual fitness with individual gene frequencies. To do so, first note that since competition is within groups and between individuals of the same sex, male and female fitness functions can in general be

written as functions of three relevant phenotypic values,

$$w_{ij}^u \left(z_{ij}^u, \overline{z}_{i0}^u, \overline{z}^u \right),$$ (A-7)

where z_{ij}^u is the phenotype of the focal individual whose fitness is being considered (indexed j of sex u in group i); $\overline{z}_{i0}^u$ is the average phenotype among the other individuals of the same sex born in group i (i.e. excluding the focal individual); and $\overline{z}^u$ is the average phenotype of individuals of sex u in the population (see eq. A-30 for an example of such fitness function). Second, we use the chain rule on these fitness functions to obtain,

$$
\begin{aligned}
w_{ij}^{\mathrm{f}} &= 2 + \delta \left(\frac{\partial w_{ij}^{\mathrm{f}}}{\partial z_{ij}^{\mathrm{f}}} \frac{\mathrm{d} z_{ij}^{\mathrm{f}}}{\mathrm{d}\delta} + \frac{\partial w_{ij}^{\mathrm{f}}}{\partial \overline{z}_{i0}^{\mathrm{f}}} \frac{\mathrm{d} \overline{z}_{i0}^{\mathrm{f}}}{\mathrm{d}\delta} + \frac{\partial w_{ij}^{\mathrm{f}}}{\partial \overline{z}^{\mathrm{f}}} \frac{\mathrm{d} \overline{z}^{\mathrm{f}}}{\mathrm{d}\delta} \right) + \mathscr{O}\left(\delta^2\right), \\
w_{ij}^{\mathrm{m}} &= 2 + \delta \left(\frac{\partial w_{ij}^{\mathrm{m}}}{\partial z_{ij}^{\mathrm{m}}} \frac{\mathrm{d} z_{ij}^{\mathrm{m}}}{\mathrm{d}\delta} + \frac{\partial w_{ij}^{\mathrm{m}}}{\partial \overline{z}_{i0}^{\mathrm{m}}} \frac{\mathrm{d} \overline{z}_{i0}^{\mathrm{m}}}{\mathrm{d}\delta} + \frac{\partial w_{ij}^{\mathrm{m}}}{\partial \overline{z}^{\mathrm{m}}} \frac{\mathrm{d} \overline{z}^{\mathrm{m}}}{\mathrm{d}\delta} \right) + \mathscr{O}\left(\delta^2\right)
\end{aligned}
$$ (A-8)

where 2 is the expected total number of juveniles produced by an individual under neutrality (i.e. one male and one female juvenile when $\delta = 0$) and all derivatives are evaluated at $\delta = 0$ (so at $z_{ij}^{\mathrm{f}} = \overline{z}_{i0}^{\mathrm{f}} = \overline{z}^{\mathrm{f}} = z_{ij}^{\mathrm{m}} = \overline{z}_{i0}^{\mathrm{m}} = \overline{z}^{\mathrm{m}} = 1$, from eq. A-3). The above equation may be further simplified by using the following relationships

$$
\begin{aligned}
\frac{\partial w_{ij}^{\mathrm{f}}}{\partial z_{ij}^{\mathrm{f}}} + \frac{\partial w_{ij}^{\mathrm{f}}}{\partial \overline{z}_{i0}^{\mathrm{f}}} + \frac{\partial w_{ij}^{\mathrm{f}}}{\partial \overline{z}^{\mathrm{f}}} &= 0 \\
\frac{\partial w_{ij}^{\mathrm{m}}}{\partial z_{ij}^{\mathrm{m}}} + \frac{\partial w_{ij}^{\mathrm{m}}}{\partial \overline{z}_{i0}^{\mathrm{m}}} + \frac{\partial w_{ij}^{\mathrm{m}}}{\partial \overline{z}^{\mathrm{m}}} &= 0,
\end{aligned}
$$ (A-9)

which hold because the population of each sex is constant, so that any fitness gain made by a focal individual due to a change of phenotype must be compensated by a decrease in fitness by the rest of the population (Rousset 2004, p. 96).

Substituting eq. (A-9) into eq. (A-8), we obtain

$$
\begin{aligned}
w_{ij}^{\mathrm{f}} &= 2 + \delta \frac{\partial w_{ij}^{\mathrm{f}}}{\partial z_{ij}^{\mathrm{f}}} \left(\frac{\mathrm{d} z_{ij}^{\mathrm{f}}}{\mathrm{d}\delta} - \frac{\mathrm{d} \overline{z}^{\mathrm{f}}}{\mathrm{d}\delta} \right) + \delta \frac{\partial w_{ij}^{\mathrm{f}}}{\partial \overline{z}_{i0}^{\mathrm{f}}} \left(\frac{\mathrm{d} \overline{z}_{i0}^{\mathrm{f}}}{\mathrm{d}\delta} - \frac{\mathrm{d} \overline{z}^{\mathrm{f}}}{\mathrm{d}\delta} \right) + \mathscr{O}\left(\delta^2\right), \\
w_{ij}^{\mathrm{m}} &= 2 + \delta \frac{\partial w_{ij}^{\mathrm{m}}}{\partial z_{ij}^{\mathrm{m}}} \left(\frac{\mathrm{d} z_{ij}^{\mathrm{m}}}{\mathrm{d}\delta} - \frac{\mathrm{d} \overline{z}^{\mathrm{m}}}{\mathrm{d}\delta} \right) + \delta \frac{\partial w_{ij}^{\mathrm{m}}}{\partial \overline{z}_{i0}^{\mathrm{m}}} \left(\frac{\mathrm{d} \overline{z}_{i0}^{\mathrm{m}}}{\mathrm{d}\delta} - \frac{\mathrm{d} \overline{z}^{\mathrm{m}}}{\mathrm{d}\delta} \right) + \mathscr{O}\left(\delta^2\right).
\end{aligned}
$$ (A-10)

In turn, plugging eq. (A-10) into eq. (A-6), we obtain that the allele frequency change is

$$
\begin{aligned}
\Delta p = \frac{1}{4}\delta \Bigg[&\frac{\partial w_{ij}^{\mathrm{f}}}{\partial z_{ij}^{\mathrm{f}}} \overline{\left(\frac{\mathrm{d} z_{ij}^{\mathrm{f}}}{\mathrm{d}\delta} - \frac{\mathrm{d} \overline{z}^{\mathrm{f}}}{\mathrm{d}\delta} \right) p_{ij}^{\mathrm{f}}} + \frac{\partial w_{ij}^{\mathrm{f}}}{\partial \overline{z}_{i0}^{\mathrm{f}}} \overline{\left(\frac{\mathrm{d} \overline{z}_{i0}^{\mathrm{f}}}{\mathrm{d}\delta} - \frac{\mathrm{d} \overline{z}^{\mathrm{f}}}{\mathrm{d}\delta} \right) p_{ij}^{\mathrm{f}}} \\
&+ \frac{\partial w_{ij}^{\mathrm{m}}}{\partial z_{ij}^{\mathrm{m}}} \overline{\left(\frac{\mathrm{d} z_{ij}^{\mathrm{m}}}{\mathrm{d}\delta} - \frac{\mathrm{d} \overline{z}^{\mathrm{m}}}{\mathrm{d}\delta} \right) p_{ij}^{\mathrm{m}}} + \frac{\partial w_{ij}^{\mathrm{m}}}{\partial \overline{z}_{i0}^{\mathrm{m}}} \overline{\left(\frac{\mathrm{d} \overline{z}_{i0}^{\mathrm{m}}}{\mathrm{d}\delta} - \frac{\mathrm{d} \overline{z}^{\mathrm{m}}}{\mathrm{d}\delta} \right) p_{ij}^{\mathrm{m}}} \Bigg] + \mathscr{O}\left(\delta^2\right),
\end{aligned}
$$ (A-11)

where the upper bar means the average over all juveniles and over all groups. From eq. (A-3), the averages in eq. (A-11) can be expressed in terms of products of allele frequencies within and between individuals:

$$\overline{\left(\frac{\mathrm{d} z_{ij}^{\mathrm{f}}}{\mathrm{d}\delta} - \frac{\mathrm{d} \overline{z}^{\mathrm{f}}}{\mathrm{d}\delta} \right) p_{ij}^{\mathrm{f}}} = c_{\mathrm{f}} \left[2(1 - h_{\mathrm{f}}) \left(\overline{(p_{ij}^{\mathrm{f}})^2} - (p^{\mathrm{f}})^2 \right) + (1 - 2(1 - h_{\mathrm{f}})) \left(\overline{p_{ij1}^{\mathrm{f}} p_{ij2}^{\mathrm{f}} p_{ij}^{\mathrm{f}}} - \overline{p_{ij1}^{\mathrm{f}} p_{ij2}^{\mathrm{f}}} p^{\mathrm{f}} \right) \right]$$ (A-12)

$$\overline{\left(\frac{\mathrm{d} \overline{z}_{i0}^{\mathrm{f}}}{\mathrm{d}\delta} - \frac{\mathrm{d} \overline{z}^{\mathrm{f}}}{\mathrm{d}\delta} \right) p_{ij}^{\mathrm{f}}} = c_{\mathrm{f}} \left[2(1 - h_{\mathrm{f}}) \left(\overline{p_{i0}^{\mathrm{f}} p_{ij}^{\mathrm{f}}} - (p^{\mathrm{f}})^2 \right) + (1 - 2(1 - h_{\mathrm{f}})) \left(\overline{p_{i01}^{\mathrm{f}} p_{i02}^{\mathrm{f}} p_{ij}^{\mathrm{f}}} - \overline{p_{ij1}^{\mathrm{f}} p_{ij2}^{\mathrm{f}}} p^{\mathrm{f}} \right) \right],$$ (A-13)

in females, where p_{i01}^{f} (and p_{i02}^{f}) is the average frequency of the A allele at the maternal (and paternal) locus of

the neighbouring females of a focal female indexed j in group i, and $p_{i0}^{\mathrm{f}} = (p_{i01}^{\mathrm{f}} + p_{i02}^{\mathrm{f}})/2$. Similarly,

$$\overline{\left(\frac{\mathrm{d}z_{ij}^{\mathrm{m}}}{\mathrm{d}\delta} - \frac{\mathrm{d}\overline{z}^{\mathrm{m}}}{\mathrm{d}\delta}\right)p_{ij}^{\mathrm{m}}} = -c_{\mathrm{m}}\left[2h_{\mathrm{m}}\left(\overline{(p_{ij}^{\mathrm{m}})^2} - (p^{\mathrm{m}})^2\right) + (1-2h_{\mathrm{m}})\left(\overline{p_{ij1}^{\mathrm{m}} p_{ij2}^{\mathrm{m}} p_{ij}^{\mathrm{m}}} - \overline{p_{ij1}^{\mathrm{m}} p_{ij2}^{\mathrm{m}}}p^{\mathrm{m}}\right)\right] \qquad \text{(A-14)}$$

$$\overline{\left(\frac{\mathrm{d}\overline{z}_{i0}^{\mathrm{m}}}{\mathrm{d}\delta} - \frac{\mathrm{d}\overline{z}^{\mathrm{m}}}{\mathrm{d}\delta}\right)p_{ij}^{\mathrm{m}}} = -c_{\mathrm{m}}\left[2h_{\mathrm{m}}\left(\overline{p_{i0}^{\mathrm{m}} p_{ij}^{\mathrm{m}}} - (p^{\mathrm{m}})^2\right) + (1-2h_{\mathrm{m}})\left(\overline{p_{i01}^{\mathrm{m}} p_{i02}^{\mathrm{m}} p_{ij}^{\mathrm{m}}} - \overline{p_{ij1}^{\mathrm{m}} p_{ij2}^{\mathrm{m}}}p^{\mathrm{m}}\right)\right], \qquad \text{(A-15)}$$

in males, where p_{i01}^{m} (and p_{i02}^{m}) is the average frequency of the A allele at the maternal (and paternal) locus of the neighbouring males of a focal male indexed j in group i, and $p_{i0}^{\mathrm{m}} = (p_{i01}^{\mathrm{m}} + p_{i02}^{\mathrm{m}})/2$. We first simplify eqs. (A-12) and (A-14). Using the properties of indicator variables (that $(p_{ij1}^u)^2 = p_{ij1}^u$ and $(p_{ij2}^u)^2 = p_{ij2}^u$), we have

$$(p_{ij}^u)^2 = \frac{1}{2}p_{ij}^u + \frac{1}{2}p_{ij1}^u p_{ij2}^u \qquad \text{(A-16)}$$

so that

$$\overline{(p_{ij}^u)^2} = \frac{1}{2}p^u + \frac{1}{2}\overline{p_{ij1}^u p_{ij2}^u}. \qquad \text{(A-17)}$$

Similarly, we have that

$$\overline{p_{ij1}^u p_{ij2}^u p_{ij}^u} = \overline{p_{ij1}^u p_{ij2}^u}. \qquad \text{(A-18)}$$

Substituting eqs. (A-17) and (A-18) into eqs. (A-12) and (A-14), these read as

$$\overline{\left(\frac{\mathrm{d}z_{ij}^{\mathrm{f}}}{\mathrm{d}\delta} - \frac{\mathrm{d}\overline{z}^{\mathrm{f}}}{\mathrm{d}\delta}\right)p_{ij}^{\mathrm{f}}} = c_{\mathrm{f}}\left[(1-p^{\mathrm{f}})\overline{p_{ij1}^{\mathrm{f}} p_{ij2}^{\mathrm{f}}} + (1-h_{\mathrm{f}})(1-2p^{\mathrm{f}})\left(p^{\mathrm{f}} - \overline{p_{ij1}^{\mathrm{f}} p_{ij2}^{\mathrm{f}}}\right)\right]$$

$$\overline{\left(\frac{\mathrm{d}z_{ij}^{\mathrm{m}}}{\mathrm{d}\delta} - \frac{\mathrm{d}\overline{z}^{\mathrm{m}}}{\mathrm{d}\delta}\right)p_{ij}^{\mathrm{m}}} = -c_{\mathrm{m}}\left[(1-p^{\mathrm{m}})\overline{p_{ij1}^{\mathrm{m}} p_{ij2}^{\mathrm{m}}} + h_{\mathrm{m}}(1-2p^{\mathrm{m}})\left(p^{\mathrm{m}} - \overline{p_{ij1}^{\mathrm{m}} p_{ij2}^{\mathrm{m}}}\right)\right]. \qquad \text{(A-19)}$$

Next, note that the different averages in eqs. (A-13), (A-15), and (A-19) correspond to different probabilities of genetic identity. For example, $\overline{p_{ij1}^u p_{ij2}^u}$ corresponds to the probability that two homologous gene copies randomly sampled in a juvenile of sex u before dispersal are both A. Further, these probabilities only need to be computed under neutrality to determine allele frequency change to the first order of δ (as they are already multiplied by δ in eqs. A-11). In such circumstances, we can relate probabilities of identity with probabilities of identity-by-descent of neutral genes, which are independent of the frequency of A in the infinite island model (see Roze and Rousset, 2003 for further considerations). Take $\overline{p_{ij1}^u p_{ij2}^u}$ again for example. Consider the two lineages of these genes backward in time: either the two lineages have stayed in the same group and coalesced (with a probability that we denote F_{IT} and which we will show later is equal in males and females, see eq. A-37) or have dispersed to different groups patches and have not coalesced (with probability $1 - F_{\mathrm{IT}}$). The probability that these two genes are A is then equal to $F_{\mathrm{IT}}p^u + (1 - F_{\mathrm{IT}})(p^u)^2$, where p^u is the frequency of A in the population of individuals of sex u. However, this frequency is equal in males and females under neutrality ($p^{\mathrm{m}} = p^{\mathrm{f}} = p$) so that we have

$$\overline{p_{ij1}^u p_{ij2}^u} = F_{\mathrm{IT}}p + (1 - F_{\mathrm{IT}})p^2 + \mathcal{O}(\delta). \qquad \text{(A-20)}$$

The probability F_{IT} is thus equivalent to Wright's coefficient of inbreeding in the infinite island model. We will later compute this coefficient explicitly in terms of demographic parameters in section A.2.6. First, we express the other two averages featuring in eqs. (A-13) and (A-15) ($\overline{p_{i0}^u p_{ij}^u}$ and $\overline{p_{i01}^u p_{i02}^u p_{ij}^u}$) in terms of coalescence probabilities.

The probability, $\overline{p_{i0}^u p_{ij}^u}$, that two genes randomly sampled in different juveniles of the same sex u within the same group are both A can be expressed as

$$\overline{p_{i0}^u p_{ij}^u} = F_{\mathrm{ST}}p + (1 - F_{\mathrm{ST}})p^2 + \mathcal{O}(\delta), \qquad \text{(A-21)}$$

where F_{ST} is the probability that the two lineages of our two genes have stayed in the same group and coalesced (which is equal when those two genes are sampled in males and in females, see eq. A-32 for more details). This probability is equal to Wright's F_{ST} within sexes in the infinite island model.

Finally, we need $\overline{p_{i01}^u p_{i02}^u p_{ij}^u}$, which corresponds to the probability that the two homologous genes of a juvenile of sex u and a third gene sampled in another juvenile of the same sex from the same group are all A. To compute this quantity, we let K be the probability that the three lineages of these genes have stayed in the same group and coalesced; and L be the probability that exactly two of those have coalesced (both probabilities are insensitive as to whether genes are sampled in males or females, see eq. A-39). With this notation, we have

$$\overline{p_{i01}^u p_{i02}^u p_{ij}^u} = Kp + Lp^2 + (1 - K - L)\, p^3 + \mathcal{O}(\delta)\,. \tag{A-22}$$

Further, from the relationships between the different coalescence events, we can write the probability L that exactly two of the genes of interest have coalesced as $L = F_{\mathrm{IT}} - K + 2(F_{\mathrm{ST}} - K)$, where $F_{\mathrm{IT}} - K$ is the probability that the two homologous gene copies coalesce together but not with the gene sampled in another juvenile, and $2(F_{\mathrm{ST}} - K)$ is the probability that the gene sampled in another juvenile and one gene from the homologous copies coalesce but not with the other copy. Substituting for L into eq. (A-22), we obtain

$$\overline{p_{i01}^u p_{i02}^u p_{ij}^u} = Kp + (F_{\mathrm{IT}} + 2F_{\mathrm{ST}} - 3K)\, p^2 + (1 - F_{\mathrm{IT}} - 2F_{\mathrm{ST}} + 2K)\, p^3 + \mathcal{O}(\delta) \tag{A-23}$$

for the probability that the two homologous genes of a juvenile of sex u and a third gene sampled in another juvenile of the same sex from the same group are all A.

Plugging eqs. (A-20), (A-21) and (A-23) into eqs. (A-13), (A-15) and (A-19), which are in turn substituted into eq. (A-11), we get that the change in allele frequency is given by

$$\Delta p(p) = \frac{\delta}{2} p(1 - p) \left(c_{\mathrm{f}} s_{\mathrm{f}}(p) - c_{\mathrm{m}} s_{\mathrm{m}}(p) \right) + \mathcal{O}\left(\delta^2\right), \tag{A-24}$$

where

$$s_{\mathrm{f}}(p) = \frac{1}{2} \frac{\partial w_{ij}^{\mathrm{f}}}{\partial z_{ij}^{\mathrm{f}}} \left[(1 - F_{\mathrm{IT}})p + F_{\mathrm{IT}} + (1 - h_{\mathrm{f}})(1 - 2p)(1 - F_{\mathrm{IT}}) \right] + \frac{1}{2} \frac{\partial w_{ij}^{\mathrm{f}}}{\partial \overline{z}_{i0}^{\mathrm{f}}} \left[K + 2(F_{\mathrm{ST}} - K)\left(p + (1 - h_{\mathrm{f}})(1 - 2p) \right) \right]$$

$$s_{\mathrm{m}}(p) = \frac{1}{2} \frac{\partial w_{ij}^{\mathrm{m}}}{\partial z_{ij}^{\mathrm{m}}} \left[(1 - F_{\mathrm{IT}})p + F_{\mathrm{IT}} + h_{\mathrm{m}}(1 - 2p)(1 - F_{\mathrm{IT}}) \right] + \frac{1}{2} \frac{\partial w_{ij}^{\mathrm{m}}}{\partial \overline{z}_{i0}^{\mathrm{m}}} \left[K + 2(F_{\mathrm{ST}} - K)\left(p + h_{\mathrm{m}}(1 - 2p) \right) \right],$$

$$\tag{A-25}$$

capture selection on the A allele due to its effect in females and males, respectively. We proceed to specify these selection effects in terms of demographic parameters and sex-specific dispersal, by specifying first the relevant fitness effects (i.e. fitness derivatives in section A.2.5), and second, relevant coalescence probabilities (in section A.2.6).

A.2.5 Fitness

According to the life-cycle described in section A.1, the expected number of female and male juveniles produced by a focal female juvenile can be decomposed as

$$w_{ij}^{\mathrm{f} \to \mathrm{f}} = w_{ij}^{\mathrm{f} \to \mathrm{m}} = s_{ij}^{\mathrm{f}} k/2 \tag{A-26}$$

where s_{ij}^{f} is the probability that this female juvenile survives competition to adulthood and $k/2$ is the number of male and female juveniles produced by an adult female. When the number of groups n_{g} is large and selection strength δ is small, the survival probability of a focal female is related to its phenotype z_{ij}^{f} (i.e. its competitiveness) according to

$$s_{ij}^{\mathrm{f}} = \frac{(1 - m_{\mathrm{f}})n_{\mathrm{f}} z_{ij}^{\mathrm{f}}}{z_{ij}^{\mathrm{f}} + (n_{\mathrm{f}} k/2 - 1)\left[(1 - m_{\mathrm{f}})\overline{z}_{i0}^{\mathrm{f}} + m_{\mathrm{f}}\overline{z}^{\mathrm{f}} \right]} + \frac{m_{\mathrm{f}} n_{\mathrm{f}} z_{ij}^{\mathrm{f}}}{z_{ij}^{\mathrm{f}} + (n_{\mathrm{f}} k/2 - 1)\overline{z}^{\mathrm{f}}}, \tag{A-27}$$

where m_{f} is the probability that a female juvenile disperses; $\overline{z}_{i0}^{\mathrm{f}}$ is the average phenotype among the other female juveniles born in the focal group (i.e. excluding the focal female); and $\overline{z}^{\mathrm{f}}$ is the average female phenotype

in the population. The first summand of eq. (A-27) is the probability that the focal female remains in its natal group (with probability $1 - m_f$) and survives, and the second, that she disperses (with probability m_f) and survives. In both cases, her survival depends on the ratio of her competitiveness (z_{ij}^f) and the expected average competitiveness in the group she is in.

Similarly, the expected number of male and female juveniles produced by a focal male juvenile is

$$w_{ij}^{m \to m} = w_{ij}^{m \to f} = s_{ij}^m (n_f k/2)/n_m, \tag{A-28}$$

where s_{ij}^m is the probability that this male juvenile survives to adulthood and $(n_f k/2)/n_m$ is the number of male and female juveniles produced by an adult male under random mating. Mirroring eq. (A-27), the survival probability of a focal male with phenotype z_{ij}^m can be written as

$$s_{ij}^m = \frac{(1 - m_m) n_m z_{ij}^m}{z_{ij}^m + (n_f k/2 - 1) \left[(1 - m_m) \overline{z}_{i0}^m + m_m \overline{z}^m \right]} + \frac{m_m n_m z_{ij}^m}{z_{ij}^m + (n_f k/2 - 1) \overline{z}^m}, \tag{A-29}$$

where m_m is the probability that a male juvenile disperses; $\overline{z}_{i0}^m$ is the average phenotype among the other male juveniles born in the focal group; and $\overline{z}^m$ is the average male phenotype in the population.

Substituting eq. (A-27) into eq. (A-26) and eq. (A-29) into eq. (A-28), we obtain that total female, w_{ij}^f, and male, w_{ij}^m, fitness (the expected total number of juveniles produced by a focal female and male juvenile) are given by

$$
\begin{aligned}
w_{ij}^f &= w_{ij}^{f \to f} + w_{ij}^{f \to m} = 2 \left(\frac{(1 - m_f) z_{ij}^f}{(1 - m_f) \overline{z}_{i0}^f + m_f \overline{z}^f} + \frac{m_f z_{ij}^f}{\overline{z}^f} \right) + \mathcal{O}(1/k), \\
w_{ij}^m &= w_{ij}^{m \to f} + w_{ij}^{m \to m} = 2 \left(\frac{(1 - m_m) z_{ij}^m}{(1 - m_m) \overline{z}_{i0}^m + m_m \overline{z}^m} + \frac{m_m z_{ij}^m}{\overline{z}^m} \right) + \mathcal{O}(1/k),
\end{aligned}
\tag{A-30}
$$

when fecundity k is large, which we will henceforth assume. From eq. (A-30), the direct and indirect fitness effects of female and male competitiveness are respectively given by

$$\frac{\partial w_{ij}^u}{\partial z_{ij}^u} = 2, \quad \frac{\partial w_{ij}^u}{\partial \overline{z}_{i0}^u} = -2(1 - m_u)^2, \tag{A-31}$$

for $u \in \{m, f\}$. Plugging eq. (A-31) into eq. (A-25), we finally obtain main text equations (2).

A.2.6 Coalescence probabilities at a neutral autosomal locus

In this section, we specify coalescence probabilities relevant for selection eq. (A-25) in terms of demographic parameters (numbers of males and females, n_m, n_f, and sex-specific dispersal, m_m, m_f). We do so using standard identity-by-descent arguments.

Pairwise coalescence probabilities. Consider first, $F_{ST}(t+1)$, the probability that two neutral genes randomly sampled at some arbitrary generation $t+1$ in different juveniles of the same sex u within the same group (before dispersal) coalesce. With probability 1/4 both sampled genes are maternally inherited; with probability 1/4, they are paternally inherited; and with probability 1/2, one is maternally inherited and the other paternally inherited. We can then write the probability that these two genes coalesce as

$$F_{ST}(t+1) = \frac{1}{4} \theta_f(t) + \frac{1}{4} \theta_m(t) + \frac{1}{2} \theta_{f,m}(t), \tag{A-32}$$

where $\theta_f(t)$ and $\theta_m(t)$ are respectively the probabilities that two maternally and paternally inherited genes coalesce; and $\theta_{f,m}(t)$ the probability that one paternal gene and one maternal gene (in different individuals) coalesce. Consider the first of these, $\theta_f(t)$, the probability that two maternally inherited genes coalesce. This probability will depend on whether or not the two juveniles that carry the sampled maternal genes have the

same mother: 1) If the focal juveniles have the same mother (which occurs with probability $1/n_{\mathrm{f}}$), then with probability $1/2$ the two genes are the same copy and they coalesce with probability 1, or, with probability $1/2$, the two genes are the homologous copies in the mother, in which case they coalesce with probability $F_{\mathrm{IT}}(t)$ (the probability that two homologous gene copies in a juvenile sampled at generation t coalesce); 2) If the focal juveniles have different mothers (which occurs with probability $1 - 1/n_{\mathrm{f}}$), then in order for the genes to coalesce both mothers must have remained philopatric (with probability $(1 - m_{\mathrm{f}})^2$) and their genes coalesce with probability $F_{\mathrm{ST}}(t)$. From these considerations, we thus have

$$\theta_{\mathrm{f}}(t) = \frac{1}{n_{\mathrm{f}}}\left(\frac{1}{2} + \frac{1}{2}F_{\mathrm{IT}}(t)\right) + \frac{n_{\mathrm{f}}-1}{n_{\mathrm{f}}}\left(1 - m_{\mathrm{f}}\right)^2 F_{\mathrm{ST}}(t). \tag{A-33}$$

Similarly, the probability that two paternally inherited genes coalesce is

$$\theta_{\mathrm{m}}(t) = \frac{1}{n_{\mathrm{m}}}\left(\frac{1}{2} + \frac{1}{2}F_{\mathrm{IT}}(t)\right) + \frac{n_{\mathrm{m}}-1}{n_{\mathrm{m}}}\left(1 - m_{\mathrm{m}}\right)^2 F_{\mathrm{ST}}(t), \tag{A-34}$$

where $1/n_{\mathrm{m}}$ is the probability that two juveniles have the same father. Finally, for one paternal gene and one maternal gene to coalesce, both the mother and father of the relevant juveniles must have remained philopatric, so that

$$\theta_{\mathrm{f,m}}(t) = (1 - m_{\mathrm{f}})(1 - m_{\mathrm{m}}) F_{\mathrm{ST}}(t). \tag{A-35}$$

Substituting eqs. (A-33)-(A-35) into eq. (A-32), we obtain a recurrence for $F_{\mathrm{ST}}(t)$,

$$
\begin{aligned}
F_{\mathrm{ST}}(t+1) = {}& \frac{1}{4}\left(\frac{1}{n_{\mathrm{f}}}\left(\frac{1}{2} + \frac{1}{2}F_{\mathrm{IT}}(t)\right) + \frac{n_{\mathrm{f}}-1}{n_{\mathrm{f}}}\left(1 - m_{\mathrm{f}}\right)^2 F_{\mathrm{ST}}(t)\right) \\
& + \frac{1}{4}\left(\frac{1}{n_{\mathrm{m}}}\left(\frac{1}{2} + \frac{1}{2}F_{\mathrm{IT}}(t)\right) + \frac{n_{\mathrm{m}}-1}{n_{\mathrm{m}}}\left(1 - m_{\mathrm{m}}\right)^2 F_{\mathrm{ST}}(t)\right) + \frac{1}{2}\left(1 - m_{\mathrm{f}}\right)\left(1 - m_{\mathrm{m}}\right) F_{\mathrm{ST}}(t)
\end{aligned}
\tag{A-36}
$$

that also depends on $F_{\mathrm{IT}}(t)$, the probability that two homologous gene copies in a juvenile sampled at generation t coalesce. This probability, meanwhile, satisfies the recurrence

$$F_{\mathrm{IT}}(t+1) = (1 - m_{\mathrm{f}})(1 - m_{\mathrm{m}}) F_{\mathrm{ST}}(t), \tag{A-37}$$

as for the two homologous gene copies of a juvenile to coalesce, both its mother and father must have remained philopatric, in which case the coalesce with probability $F_{\mathrm{ST}}(t)$.

Solving for the equilibrium, $F_{\mathrm{ST}}(t+1) = F_{\mathrm{ST}}(t) = F_{\mathrm{ST}}$ and $F_{\mathrm{IT}}(t) = F_{\mathrm{IT}}(t+1) = F_{\mathrm{IT}}$, using eqs. (A-36)-(A-37), we obtain

$$F_{\mathrm{ST}} = \frac{n_{\mathrm{f}} + n_{\mathrm{m}}}{n_{\mathrm{f}}(1 - m_{\mathrm{m}})(1 + m_{\mathrm{f}} - 2m_{\mathrm{m}}) + n_{\mathrm{m}}(1 - m_{\mathrm{f}})(1 + m_{\mathrm{m}} - 2m_{\mathrm{f}}) + 2n_{\mathrm{f}}n_{\mathrm{m}}(4 - m_{\mathrm{f}} - m_{\mathrm{m}})(m_{\mathrm{f}} + m_{\mathrm{m}})} \tag{A-38}$$
$$F_{\mathrm{IT}} = (1 - m_{\mathrm{f}})(1 - m_{\mathrm{m}}) F_{\mathrm{ST}}.$$

Substituting for $m_{\mathrm{f}} = M_{\mathrm{f}}/n_{\mathrm{f}}$, $m_{\mathrm{m}} = M_{\mathrm{m}}/n_{\mathrm{m}}$, $n_{\mathrm{f}} = (1 - r)n$ and $n_{\mathrm{m}} = rn$ in the above equation, we obtain F in eq. (4) of the main text in the limit $n \to \infty$.

Threeway coalescence probabilities. To compute the probability, K, that the two homologous genes of a juvenile (which we label as juvenile "1" for the sake of argument) and a third gene sampled in another juvenile (labelled juvenile "2") of the same sex sampled from the same group before dispersal all coalesce, consider that for this to happen, it is necessary that both parents of juvenile 1 are philopatric, which occurs with probability $(1 - m_{\mathrm{f}})(1 - m_{\mathrm{m}})$. Then, with probability $1/2$, the gene sampled in juvenile 2 is maternally inherited, in which case all three genes coalesce with a probability we call $\Theta_{\mathrm{f}}(t)$; with probability $1/2$, the gene sampled in juvenile 2 is paternally inherited, in which case all three genes coalesce with a probability we call $\Theta_{\mathrm{m}}(t)$, so that

$$K(t+1) = (1 - m_{\mathrm{f}})(1 - m_{\mathrm{m}})\left(\frac{1}{2}\Theta_{\mathrm{f}}(t) + \frac{1}{2}\Theta_{\mathrm{m}}(t)\right). \tag{A-39}$$

We proceed to specify $\Theta_{\mathrm{f}}(t)$ and $\Theta_{\mathrm{m}}(t)$.

Consider $\Theta_{\mathrm{f}}(t)$ first. If the gene sampled in juvenile 2 is maternally inherited, then whether it coalesces with the homologous genes of juvenile 1 depends on whether or not they have the same mother. If they have the same

mother (which occurs with probability $1/n_f$), then with probability $1/2$, the maternal genes of juveniles 1 and 2 are the same and all three genes coalesce with probability $F_{ST}(t)$, and with probability $1/2$, the maternal genes of juveniles 1 and 2 are different and all three genes coalesce with probability $K(t)$. If, however, juveniles have different mothers (which occurs with probability $1 - 1/n_f$), then for all three genes to coalesce, first the mother of juvenile 2 must be philopatric (which occurs with probability $(1 - m_f)$) and second, three genes sampled in three different juveniles must coalesce (we call $\kappa(t)$ the probability of this happening). The probability that two homologous genes of juvenile 1 and the maternally inherited gene of juvenile coalesce then reads as

$$\Theta_f(t) = \frac{1}{n_f}\left(\frac{1}{2}F_{ST}(t) + \frac{1}{2}K(t)\right) + \frac{n_f - 1}{n_f}(1 - m_f)\kappa(t). \tag{A-40}$$

Similarly, the probability that two homologous genes of juvenile 1 and the paternally inherited gene of juvenile coalesce then reads as

$$\Theta_m(t) = \frac{1}{n_m}\left(\frac{1}{2}F_{ST}(t) + \frac{1}{2}K(t)\right) + \frac{n_m - 1}{n_m}(1 - m_m)\kappa(t). \tag{A-41}$$

In order to compute $K(t+1)$, we therefore need an expression for $\kappa(t)$, the probability that three genes sampled in three different juveniles before dispersal coalesce. We can follow a similar argument as the one used to derive eq. (A-36). Specifically, conditioning on which genes are sampled in each juvenile (maternal or paternal), and whether these juveniles have parents in common, we obtain

$$\begin{aligned}
\kappa(t+1) = &\frac{1}{8}\left(\frac{1}{n_f^2}\left(\frac{1}{4} + \frac{3}{4}F_{IT}(t)\right) + 3\frac{n_f - 1}{n_f^2}(1 - m_f)^2\left(\frac{1}{2}F_{ST}(t) + \frac{1}{2}K(t)\right) + \frac{(n_f - 1)(n_f - 2)}{n_f^2}(1 - m_f)^3\kappa(t)\right) \\
&+ \frac{3}{8}(1 - m_f)(1 - m_m)(\Theta_f(t) + \Theta_m(t)) \\
&+ \frac{1}{8}\left(\frac{1}{n_m^2}\left(\frac{1}{4} + \frac{3}{4}F_{IT}(t)\right) + 3\frac{n_m - 1}{n_m^2}(1 - m_m)^2\left(\frac{1}{2}F_{ST}(t) + \frac{1}{2}K(t)\right) + \frac{(n_m - 1)(n_m - 2)}{n_m^2}(1 - m_m)^3\kappa(t)\right),
\end{aligned} \tag{A-42}$$

where the first line is the probability that all three genes are maternal and coalesce (decomposed according to whether the three sampled juveniles have the same mother, with probability $1/n_f^2$, two out of three have the same mother, with probability $3(n_f - 1)/n_f^2$, or when they have different mothers, with probability $(n_f - 1)(n_f - 2)/n_f^2$); the second line is the probability that two genes are maternal and one is paternal or that one is maternal and the other two are paternal and that they coalesce; and the third line is the probability that all three genes are paternal and coalesce (decomposed according to whether the three sampled juveniles have the same father, with probability $1/n_m^2$, two out of three have the same father, with probability $3(n_m - 1)/n_m^2$, or when they have different fathers, with probability $(n_m - 1)(n_m - 2)/n_m^2$).

Substituting eqs. (A-40) and (A-41) into (A-39) and (A-42) (with $F_{ST}(t)$ and $F_{IT}(t)$ at equilibrium eq. A-38) allows us to solve for the equilibrium coalescence probabilities $K = K(t+1) = K(t)$ and $\kappa = \kappa(t+1) = \kappa(t)$. Doing so we obtain a complicated expression for K, which when dispersal is weak and local population size is large is given by eq. (4) of the main text (i.e. substituting for $m_f = M_f/n_f$, $m_m = M_m/n_m$, $n_f = (1 - r)n$ and $n_m = rn$ and taking the limit $n \to \infty$).

A.2.7   Allele frequency at a polymorphic equilibrium

In this section we derive the equilibrium frequency p^* of the male-detrimental, female-beneficial A allele when polymorphism is favoured (i.e. eq. (5) of the main text holds). By plugging main text eq. (4) in to eqs. (1)-(2) and solving for $\Delta p(p^*) = 0$ for p^*, we find

$$p^* = p_\circ^*\left(1 + F(c_f - c_m)\right), \tag{A-43}$$

where F is given in eq. (4) and

$$p_\circ^* = \frac{(1 - h_f)c_f - h_m c_m}{(1 - 2h_f)c_f + (1 - 2h_m)c_m} \tag{A-44}$$

is the equilibrium frequency in a well-mixed population (see e.g. Box 3 in Mullon et al., 2012). Eq. (A-43) reveals that, compared with the well mixed case (when $F = 0$), limited dispersal ($F > 0$) leads to a greater equilibrium

frequency of the female-beneficial allele when $c_f > c_m$ (and vice-versa) and that this occurs irrespective of dominance effects.

A.3 Sex-linked locus

In this section, we derive and analyse the allele frequency change at a sex-linked locus and show that the effect of limited on allele segregation is largely the same as on autosomes (our results are summarised in section 3.1.5 of the main text). Without loss of generality, we assume that our population of interest belongs to an XY species (i.e. in which males are the heterogametic sex), so that the sexually antagonistic allele segregates on the X-chromosome. All our results can be applied to ZW species by simply exchanging male and females variables.

A.3.1 Weighted allele frequency change

Our argument follows the same argument used for an autosomal locus in section A.2. In terms of genotype and phenotype, females are specified in the same way as for an autosomal locus (eqs. A-1 and A-3). Males however are heterogametic and therefore have genotype and phenotype respectively specified by

$$
\begin{aligned}
p_{ij}^{\mathrm{m}} &= p_{ij1}^{\mathrm{m}} \in \{0,1\} \\
z_{ij}^{\mathrm{m}} &= 1 - \delta c_{\mathrm{m}} p_{ij}^{\mathrm{m}}.
\end{aligned}
\tag{A-45}
$$

Since males only pass their X linked genes to their daughters, the weighted allele frequency change is now given by

$$
\Delta p = \frac{1}{n_{\mathrm{g}}} \sum_{i=1}^{n_{\mathrm{g}}} \frac{1}{n_{\mathrm{f}} k/2} \left[\alpha \left(\sum_{j=1}^{n_{\mathrm{f}} k/2} w_{ij}^{\mathrm{f} \to \mathrm{f}} \frac{p_{ij}^{\mathrm{f}}}{2} + \sum_{j=1}^{n_{\mathrm{f}} k/2} w_{ij}^{\mathrm{m} \to \mathrm{f}} \frac{p_{ij}^{\mathrm{m}}}{2} \right) + (1-\alpha) \sum_{j=1}^{n_{\mathrm{f}} k/2} w_{ij}^{\mathrm{f} \to \mathrm{m}} p_{ij}^{\mathrm{f}} \right] - p
\tag{A-46}
$$

(compare with eq. A-5 for autosomes). Also due to male heterogamety, females and males now have reproductive values $\alpha = 2/3$ and $1-\alpha = 1/3$, respectively (see eq. A-4). Substituting for these reproductive values and using the fact that $w_{ij}^{\mathrm{f} \to \mathrm{f}} = w_{ij}^{\mathrm{f} \to \mathrm{m}} = w_{ij}^{\mathrm{f}}/2$ and $w_{ij}^{\mathrm{m} \to \mathrm{f}} = w_{ij}^{\mathrm{m}}/2$ for our model, eq. (A-46) becomes

$$
\Delta p = \frac{1}{6 n_{\mathrm{g}}} \sum_{i=1}^{n_{\mathrm{g}}} \frac{1}{n_{\mathrm{f}} k/2} \left[2 \sum_{j=1}^{n_{\mathrm{f}} k/2} w_{ij}^{\mathrm{f}} p_{ij}^{\mathrm{f}} + \sum_{j=1}^{n_{\mathrm{f}} k/2} w_{ij}^{\mathrm{m}} p_{ij}^{\mathrm{m}} \right] - p
\tag{A-47}
$$

(compare with eq. A-11 for autosomes).

Substituting fitness expansions eq. (A-10) into eq. (A-48), we obtain the following expression,

$$
\begin{aligned}
\Delta p = \frac{\delta}{6} \Bigg[&2 \frac{\partial w_{ij}^{\mathrm{f}}}{\partial z_{ij}^{\mathrm{f}}} \overline{\left(\frac{\mathrm{d} z_{ij}^{\mathrm{f}}}{d\delta} - \frac{\mathrm{d}\overline{z}^{\mathrm{f}}}{d\delta} \right) p_{ij}^{\mathrm{f}}} + 2 \frac{\partial w_{ij}^{\mathrm{f}}}{\partial \overline{z}_{i0}^{\mathrm{f}}} \overline{\left(\frac{\mathrm{d}\overline{z}_{i0}^{\mathrm{f}}}{d\delta} - \frac{\mathrm{d}\overline{z}^{\mathrm{f}}}{d\delta} \right) p_{ij}^{\mathrm{f}}} \\
&+ \frac{\partial w_{ij}^{\mathrm{m}}}{\partial z_{ij}^{\mathrm{m}}} \overline{\left(\frac{\mathrm{d} z_{ij}^{\mathrm{m}}}{d\delta} - \frac{\mathrm{d}\overline{z}^{\mathrm{m}}}{d\delta} \right) p_{ij}^{\mathrm{m}}} + \frac{\partial w_{ij}^{\mathrm{m}}}{\partial \overline{z}_{i0}^{\mathrm{m}}} \overline{\left(\frac{\mathrm{d}\overline{z}_{i0}^{\mathrm{m}}}{d\delta} - \frac{\mathrm{d}\overline{z}^{\mathrm{m}}}{d\delta} \right) p_{ij}^{\mathrm{m}}} \Bigg] + \mathscr{O}\left(\delta^2\right),
\end{aligned}
\tag{A-48}
$$

where the upper bar means the average over all juveniles and over all groups. The female related averages are computed in the same way as we did for autosomes (see section A.2.4), giving

$$
\begin{aligned}
\overline{\left(\frac{\mathrm{d} z_{ij}^{\mathrm{f}}}{d\delta} - \frac{\mathrm{d}\overline{z}^{\mathrm{f}}}{d\delta} \right) p_{ij}^{\mathrm{f}}} &= c_{\mathrm{f}} p(1-p) \left[(1-F_{\mathrm{IT}}) p + F_{\mathrm{IT}} + (1-h_{\mathrm{f}})(1-2p)(1-F_{\mathrm{IT}}) \right] + \mathscr{O}(\delta) \\
\overline{\left(\frac{\mathrm{d}\overline{z}_{i0}^{\mathrm{f}}}{d\delta} - \frac{\mathrm{d}\overline{z}^{\mathrm{f}}}{d\delta} \right) p_{ij}^{\mathrm{f}}} &= c_{\mathrm{f}} p(1-p) \left[K^{\mathrm{f}} + 2(F_{\mathrm{ST}}^{\mathrm{f}} - K^{\mathrm{f}}) \left(p + (1-h_{\mathrm{f}})(1-2p) \right) \right] + \mathscr{O}(\delta)
\end{aligned}
\tag{A-49}
$$

where F_{IT} is the probability that the two homologous gene copies at a neutral X-linked locus, sampled in a random juvenile female, are IBD (note that such a probability does not exist for males as they are heterogametic);

F_{ST}^f denotes the probability that two neutral X-linked are IBD when these are sampled in different female juveniles of the same group; and K^f is the probability that the two homologous gene copies in one juvenile female plus one copy in another randomly sampled juvenile female from the same group are all IBD.

For male related averages, we obtain from eq. (A-45)

$$\overline{\left(\frac{dz_{ij}^m}{d\delta} - \frac{d\overline{z}^m}{d\delta}\right)p_{ij}^m} = -\delta c_m p(1-p) + \mathcal{O}(\delta)$$

$$\overline{\left(\frac{d\overline{z}_{i0}^m}{d\delta} - \frac{d\overline{z}^m}{d\delta}\right)p_{ij}^m} = -\delta c_m F_{ST}^m p(1-p) + \mathcal{O}(\delta),$$
(A-50)

where F_{ST}^m is the probability that two neutral X-linked are IBD when these are sampled in different male juveniles of the same group defined as in the preceding paragraph (note that in contrast to the autosomal case, this probability is not necessarily equal to when genes are sampled in females, i.e. $F_{ST}^f \neq F_{ST}^m$, see section A.3.2 for more details).

Substituting eqs. (A-49) and (A-50) in eq. (A-48), we obtain that the change in weighted allele frequency at an X-linked locus is

$$\Delta p(p) = \frac{\delta}{3} p(1-p) \left(2c_f s_f(p) - c_m s_m(p)\right) + \mathcal{O}(\delta^2),$$
(A-51)

where

$$s_f(p) = \frac{1}{2}\frac{\partial w_{ij}^f}{\partial z_{ij}^f}\left[(1-F_{IT})p + F_{IT} + (1-h_f)(1-2p)(1-F_{IT})\right] + \frac{1}{2}\frac{\partial w_{ij}^f}{\partial \overline{z}_{i0}^f}\left[K + 2(F_{ST}^f - K)\left(p + (1-h_f)(1-2p)\right)\right]$$

$$s_m(p) = \frac{1}{2}\frac{\partial w_{ij}^m}{\partial z_{ij}^m} + \frac{1}{2}\frac{\partial w_{ij}^m}{\partial \overline{z}_{i0}^m}F_{ST}^m.$$
(A-52)

In turn, plugging fitness derivatives eq. (A-31) into eq. (A-52), we obtain

$$s_f(p) = (1-F_{IT})p + F_{IT} + (1-h_f)(1-2p)(1-F_{IT}) - (1-m_f)^2\left[K^f + 2\left(F_{ST}^f - K^f\right)\left(p + (1-h_f)(1-2p)\right)\right]$$

$$s_m(p) = 1 - (1-m_m)^2 F_{ST}^m.$$
(A-53)

While the general form $s_f(p)$ is unchanged from the autosomal case (see eq. 2 of main text), $s_m(p)$ no longer includes direct or indirect effects arising from male heterozygotes as male hemizygosity precludes the existence of male heterozygotes. Thus, the effect of selection in males on allele frequency change, $c_m \times s_m(p)$ in eq. (A-51), depends only on c_m and the strength of male kin competition (determined by $(1-m_m)^2 F_{ST}^m$ in eq. A-53). To analyse allele dynamics at an X-linked locus further, we compute the relevant coalescence probabilities in the next section.

A.3.2 Coalescence probabilities at a neutral X-linked locus

In this section, we calculate the coalescence probabilities that are relevant for selection (eq. A-52) at an X-linked locus, using similar arguments as those used for an autosomal locus (see section A.2.6).

Pairwise coalescence probabilities. Let us first consider the probabilities that two neutral X-linked genes sampled at generation $t+1$ in two neighbouring juvenile females, $F_{ST}^f(t+1)$, and males, $F_{ST}^m(t+1)$, coalesce. As for the autosomal case (see eq. A-32), we can decompose $F_{ST}^f(t+1)$ according to whether the two sampled genes in females are either both of maternal or paternal origin (in which case they coalesce with respective probabilities $\theta_f(t)$ and $\theta_m(t)$) or one is of maternal and the other of paternal origin (in which case they coalesce with respective probabilities $\theta_{f,m}(t)$ and $\theta_m(t)$). For X-linked genes sampled in two different males, $F_{ST}^m(t+1)$,

both genes are necessarily of maternal origin. We thus obtain a recurrence of the form

$$
\begin{aligned}
F_{\text{ST}}^{\text{f}}(t+1) &= \frac{1}{4}\theta_{\text{f}}(t) + \frac{1}{4}\theta_{\text{m}}(t) + \frac{1}{2}\theta_{\text{f,m}}(t) \\
F_{\text{ST}}^{\text{m}}(t+1) &= \theta_{\text{f}}(t).
\end{aligned}
\tag{A-54}
$$

Then, coalescence probabilities according to parental origin can be decomposed as

$$
\begin{aligned}
\theta_{\text{f}}(t) &= \frac{1}{n_{\text{f}}}\left(\frac{1}{2} + \frac{1}{2}F_{\text{IT}}(t)\right) + \frac{n_{\text{f}} - 1}{n_{\text{f}}}(1 - m_{\text{f}})^2 F_{\text{ST}}^{\text{f}}(t) \\
\theta_{\text{m}}(t) &= \frac{1}{n_{\text{m}}} + \frac{n_{\text{m}} - 1}{n_{\text{m}}}(1 - m_{\text{m}})^2 F_{\text{ST}}^{\text{m}}(t) \\
\theta_{\text{f,m}}(t) &= (1 - m_{\text{f}})(1 - m_{\text{m}}) F_{\text{ST}}^{\text{f,m}}(t),
\end{aligned}
\tag{A-55}
$$

where $F_{\text{IT}}(t)$ is the probability that the two homologous gene copies at a neutral X-linked locus in a randomly juvenile female at generation t coalesce, and $F_{\text{ST}}^{\text{f,m}}(t)$ is the coalescence probability for two X-linked genes, one sampled in a juvenile male and another in a juvenile female (from the same group) at generation t. In turn, these two probabilities satisfy,

$$
\begin{aligned}
F_{\text{IT}}(t+1) &= \theta_{\text{f,m}}(t) \\
F_{\text{ST}}^{\text{f,m}}(t+1) &= \frac{1}{2}\theta_{\text{f}}(t) + \frac{1}{2}\theta_{\text{f,m}}(t).
\end{aligned}
\tag{A-56}
$$

Plugging eq. (A-55) into eqs. (A-54) and (A-56) and solving simultaneously for the equilibrium $F_{\text{ST}}^{\text{f}} = F_{\text{ST}}^{\text{f}}(t+1) = F_{\text{ST}}^{\text{f}}(t)$, $F_{\text{ST}}^{\text{m}} = F_{\text{ST}}^{\text{m}}(t+1) = F_{\text{ST}}^{\text{m}}(t)$, $F_{\text{ST}}^{\text{f,m}} = F_{\text{ST}}^{\text{f,m}}(t+1) = F_{\text{ST}}^{\text{f,m}}(t)$, and $F_{\text{IT}} = F_{\text{IT}}(t+1) = F_{\text{IT}}(t)$ allows us to specify the relevant coalescence probabilities for selection on an additive allele ($h_{\text{f}} = 1/2$) in terms of demographic parameters and sex-specific dispersal. Because such a calculation leads to complicated (but analytical) expressions, we only present their approximation when patches are large and dispersal is weak in the main text (by substituting for $m_{\text{f}} = M_{\text{f}}/n_{\text{f}}$, $m_{\text{m}} = M_{\text{m}}/n_{\text{m}}$, $n_{\text{f}} = (1 - r)n$, and $n_{\text{m}} = rn$ and taking the limit $n \to \infty$, in which case we find $F_{\text{X}} = F_{\text{IT}} = F_{\text{ST}}^{\text{m}} = F_{\text{ST}}^{\text{f}}$, see eq. (7) in the main text for F_{X}).

Threeway coalescence probabilities. We now develop expressions for threeway coalescence probabilities. Recall that for selection (eq. A-52), we seek to compute K^{f}, which is the probability that the two homologous gene copies in one juvenile female plus one copy in another randomly sampled juvenile female from the same group are all IBD. We further introduce K^{m}, which is the probability that the two homologous gene copies in one juvenile female plus one copy in a randomly sampled juvenile male from the same group are IBD, and which will be necessary in our calculations. Using a similar argument as for the autosomal case (see eq. A-39), we can write,

$$
\begin{aligned}
K^{\text{f}}(t+1) &= (1 - m_{\text{f}})(1 - m_{\text{m}})\left(\frac{1}{2}\Theta_{\text{f}}(t) + \frac{1}{2}\Theta_{\text{m}}(t)\right) \\
K^{\text{m}}(t+1) &= (1 - m_{\text{f}})(1 - m_{\text{m}})\Theta_{\text{f}}(t),
\end{aligned}
\tag{A-57}
$$

where $\Theta_{\text{f}}(t)$ is the probability that all three relevant genes coalesce when the one gene sampled as a single copy is maternally inherited, and $\Theta_{\text{m}}(t)$ when it is paternally inherited. These probabilities can be expressed as

$$
\begin{aligned}
\Theta_{\text{f}}(t) &= \frac{1}{n_{\text{f}}}\left(\frac{1}{2}F_{\text{ST}}^{\text{f,m}} + \frac{1}{2}K^{\text{m}}(t)\right) + \frac{n_{\text{f}} - 1}{n_{\text{f}}}(1 - m_{\text{f}})\kappa^{\text{ffm}}(t) \\
\Theta_{\text{m}}(t) &= \frac{1}{n_{\text{m}}}F_{\text{ST}}^{\text{f,m}} + \frac{n_{\text{m}} - 1}{n_{\text{m}}}(1 - m_{\text{m}})\kappa^{\text{mmf}}(t),
\end{aligned}
\tag{A-58}
$$

where $\kappa^{\text{ffm}}(t)$ is the probability that three genes, randomly sampled in two different female and one male juvenile from the same group all coalesce, and $\kappa^{\text{mmf}}(t)$ is the coalescence probability for genes randomly sampled in two different male and one female juvenile from the same group. In turn, such probabilities can be written

as

$$\kappa^{\text{ffm}}(t+1) = \frac{1}{4}\left(\frac{1}{n_{\text{f}}^2}\left(\frac{1}{4}+\frac{3}{4}F_{\text{IT}}(t)\right)+3\frac{n_{\text{f}}-1}{n_{\text{f}}^2}(1-m_{\text{f}})^2\left(\frac{1}{2}F_{\text{ST}}^{\text{f}}(t)+\frac{1}{2}K^{\text{f}}(t)\right)+\frac{(n_{\text{f}}-1)(n_{\text{f}}-2)}{n_{\text{f}}^2}(1-m_{\text{f}})^3\kappa^{\text{fff}}(t)\right)$$

$$+\frac{1}{4}(1-m_{\text{f}})(1-m_{\text{m}})(2\Theta_{\text{f}}(t)+\Theta_{\text{m}}(t))$$

$$\kappa^{\text{mmf}}(t+1) = \frac{1}{2}\left(\frac{1}{n_{\text{f}}^2}\left(\frac{1}{4}+\frac{3}{4}F_{\text{IT}}(t)\right)+3\frac{n_{\text{f}}-1}{n_{\text{f}}^2}(1-m_{\text{f}})^2\left(\frac{1}{2}F_{\text{ST}}^{\text{f}}(t)+\frac{1}{2}K^{\text{f}}(t)\right)+\frac{(n_{\text{f}}-1)(n_{\text{f}}-2)}{n_{\text{f}}^2}(1-m_{\text{f}})^3\kappa^{\text{fff}}(t)\right)$$

$$+\frac{1}{2}(1-m_{\text{f}})(1-m_{\text{m}})\Theta_{\text{f}}(t),$$

$$(\text{A-59})$$

which depend on $\kappa^{\text{fff}}(t)$ and $\kappa^{\text{mmm}}(t)$, the coalescence probabilities for genes randomly sampled in three different female and male juveniles, respectively. In turn, these two latter probabilities themselves satisfy the recurrences

$$\kappa^{\text{fff}}(t+1) = \frac{1}{8}\left(\frac{1}{n_{\text{f}}^2}\left(\frac{1}{4}+\frac{3}{4}F_{\text{IT}}(t)\right)+3\frac{n_{\text{f}}-1}{n_{\text{f}}^2}(1-m_{\text{f}})^2\left(\frac{1}{2}F_{\text{ST}}^{\text{f}}(t)+\frac{1}{2}K^{\text{f}}(t)\right)+\frac{(n_{\text{f}}-1)(n_{\text{f}}-2)}{n_{\text{f}}^2}(1-m_{\text{f}})^3\kappa^{\text{fff}}(t)\right)$$

$$+\frac{3}{8}(1-m_{\text{f}})(1-m_{\text{m}})(\Theta_{\text{f}}(t)+\Theta_{\text{m}}(t))$$

$$+\frac{1}{8}\left(\frac{1}{n_{\text{m}}^2}+3\frac{n_{\text{m}}-1}{n_{\text{m}}^2}(1-m_{\text{m}})^2F_{\text{ST}}^{\text{f,m}}(t)+\frac{(n_{\text{m}}-1)(n_{\text{m}}-2)}{n_{\text{m}}^2}(1-m_{\text{m}})^3\kappa^{\text{mmm}}(t)\right)$$

$$\kappa^{\text{mmm}}(t+1) = \frac{1}{n_{\text{f}}^2}\left(\frac{1}{4}+\frac{3}{4}F_{\text{IT}}(t)\right)+3\frac{n_{\text{f}}-1}{n_{\text{f}}^2}(1-m_{\text{f}})^2\left(\frac{1}{2}F_{\text{ST}}^{\text{f}}(t)+\frac{1}{2}K^{\text{f}}(t)\right)+\frac{(n_{\text{f}}-1)(n_{\text{f}}-2)}{n_{\text{f}}^2}(1-m_{\text{f}})^3\kappa^{\text{fff}}(t).$$

$$(\text{A-60})$$

Substituting eqs. (A-58) into eqs. (A-57),(A-59) and (A-60) finally yields a dynamical system whose equilibrium we can solve to obtain the relevant quantity, $K^{\text{f}} = K^{\text{f}}(t+1) = K^{\text{f}}(t)$, for selection. In the high density-low migration limit (letting $m_{\text{f}} = M_{\text{f}}/n_{\text{f}}$, $m_{\text{m}} = M_{\text{m}}/n_{\text{m}}$, $n_{\text{f}} = (1-r)n$ and $n_{\text{m}} = rn$ and taking the limit $n \to \infty$), we obtain

$$K^{\text{f}} = \frac{F_{\text{X}}(1+r)}{1+r+3\big(M_{\text{m}}(1-r)+2M_{\text{f}}r\big)} \qquad (\text{A-61})$$

where F_{X} is given by eq. (7) in the main text.

A.3.3 Analysis of allele frequency change at a sex-linked locus

In this section, we analyse eqs. (A-51)-(A-53) using eqs. (7) and (A-61) assuming first additive and then non-additive allele effects, in order to understand the effects of limited dispersal on the segregation of sexually antagonistic alleles on the X-chromosome.

Additive effects. To obtain eq. (6) of the main text, we substitute eq. (7) into eqs. (A-51)-(A-53) with $h_{\text{f}} = 1/2$ and $c_{\text{f}} = c_{\text{m}} = c$.

Non-additive effects. Plugging eqs. (7) and (A-61) into eqs. (A-51)-(A-53) and deriving conditions such that $\Delta p'(0) > 0$ and $\Delta p'(1) < 0$, we find that the conditions favouring the maintenance of polymorphism under non-additive effects are

$$\frac{1}{2(1-h_{\text{f}})}\left(1+\frac{(1+r)(1-2h_{\text{f}})}{1+r+6(1-h_{\text{f}})\big(M_{\text{m}}(1-r)+2M_{\text{f}}r\big)}\right) < \frac{c_{\text{f}}}{c_{\text{m}}} < \frac{1}{2h_{\text{f}}}\left(1-\frac{(1+r)(1-2h_{\text{f}})}{1+r+6h_{\text{f}}\big(M_{\text{m}}(1-r)+2M_{\text{f}}r\big)}\right). \qquad (\text{A-62})$$

Note that if males and females are not dispersal limited (i.e. when M_{m} and M_{f} tend to infinity), eq. (A-62) reduces to the polymorphism condition of a well-mixed population (e.g. eq. 2 in Fry, 2010). However, as increasingly limited dispersal leads to inbreeding and a dearth of female heterozygotes, the polymorphism space

contracts (see left and right sides of the inequality A-62 as M_m and M_f decrease). Furthermore, because coalescence occurs faster on the X-chromosome than autosomes (main text eqs. 4 and 7), this narrowing of polymorphism conditions occurs more rapidly for X-linked sexually antagonistic loci (compare Fig. 3A and B Top).

As with autosomes, sex-biased dispersal at X-linked loci leads one sex to experience stronger kin competition and shifts the polymorphism space in favour of stronger antagonistic effects in the less dispersive sex (Fig. 3B Bottom). Notice, however, that the size of polymorphism conditions on the X-chromosome are affected by the immigrant sex ratio even when there are equal numbers of males and females ($r = 0.5$), becoming smaller when immigrants are male-biased. This is because coalescence probabilities (and thus frequency of inbreeding) on the X-chromosome are increased when males are the more dispersive sex (eq. 7).

Finally, solving for $\Delta p(p^*) = 0$ (from eqs. A-51-A-53 with eqs. 7 and A-61), we find that when eq. (A-62) holds and selection favours polymorphism, the equilibrium allele frequency p^* on an X-chromosome is given by

$$p^* = p^*_{\circ X} + \frac{1+r}{3\left[M_m(1-r) + 2M_f r\right]} \times \frac{c_f - c_m}{2(1 - 2h_f)c_f}, \tag{A-63}$$

where

$$p^*_{\circ X} = \frac{2(1 - h_f)c_f - c_m}{2(1 - 2h_f)c_f} \tag{A-64}$$

is the equilibrium in a well mixed population. Inspection of the last term of eq. (A-63) reveals that p^* will increase or decrease depending on whether the sign of $c_f - c_m$ is positive or negative respectively. Thus, eq. (A-63) shows that, relative to a well-mixed population, limited dispersal (low values of M_m and M_f) leads selection to favour the sex which feels the greatest cost of their detrimental allele independently of dominance (in this case h_f), like on autosomes (see eq. A-43).

B   Individual based simulations

We simulated a diploid population subdivided among $n_g = 50$ groups following the life-cycle elaborated in Appendix A.1 (our simulation code can be downloaded from the public repository XXX – determined upon acceptance for publication). Specifically, at the beginning of a generation, adult females in each patch produce a fixed number $k = 20$ of eggs. Each egg is fertilised by a randomly sampled male within the same patch to produce a zygote whose sex is determined at random (with probability 1/2 to be male and 1/2 to be a female). Each zygote disperses independently with sex-specific dispersal probability m_m and m_f for males and females, respectively. If a zygote disperses, it joins a patch other than its natal one at random (so each remaining patch has a probability $1/(n_g - 1)$ of being the destination of dispersal). After dispersal, zygotes within each patch compete intrasexually to form the new adult mating pool. To model this, n_m males and n_f females are sampled without replacement from the male and female zygotic pools in each patch, where the probability of an individual being sampled is weighted by the competitiveness of its genotype (according to Fig. 1A).

Simulations were initiated with both alleles at frequency 0.5 at Hardy-Weinberg equilibrium across the whole population and ran until one allele was fixed. 10,000 replicates were conducted for each parameter combination (proportion of immigrants: $(M_m + M_f)/(n_m + n_f) = 10^{-2.7}, 10^{-2.6}, \ldots, 10^{-0.1}, 1$; other parameters, see Fig. 4). At the end of each generation, we calculated F_{IT} as

$$F_{IT} = 1 - \frac{\sum_{i=1}^{n_g} \sum_{j=1}^{n} \frac{1}{n_g} \frac{1}{n} \left[p_{ij1}(1 - p_{ij2}) + (1 - p_{ij1})p_{ij2}\right]}{2p(1 - p)}, \tag{B-1}$$

where $n = n_m + n_f$ is the number of individuals of either sex in a given group, p_{ij1} and p_{ij2} is the frequency of A in individual indexed j of group i at the maternally and paternally inherited locus respectively, and $p = \sum_{i=1}^{n_g} \sum_{j=1}^{n} (p_{ij1} + p_{ij2})/(2n_g n)$ is the frequency of allele A in the whole population (so the fraction in eq. B-1 is the ratio of the observed number of heterozygotes to the expected number of heterozygotes under Hardy-Weinberg equilibrium). F_{ST} is calculated as the ratio of the genetic covariance between individuals within

groups to the total genetic variance in the population, i.e., as

$$F_{\text{ST}} = \frac{\sum_{i=1}^{n_{\text{g}}} \frac{1}{n_{\text{g}}} \left[\sum_{j=1}^{n} \sum_{k=j+1}^{n} \frac{2}{n(n-1)} (p_{ij} - p)(p_{ik} - p) \right]}{\sum_{i=1}^{n_{\text{g}}} \sum_{j=1}^{n} \frac{1}{n_{\text{g}}} \frac{1}{n} (p_{ij} - p)^2},$$

(B-2)

where $p_{ij} = (p_{ij1} + p_{ij2})/2$ gives the frequency of A in individual j in group i.

Supplementary Figure 1 : Allele segregation time in individual-based simulations with large groups.
Segregation time averaged across 10,000 replicate runs as a function of total fraction of immigrants and selection strength (black dotted line shows neutrality, green solid lines show, from lightest to darkest, $c = 0.05, c = 0.1$, and $c = 0.15$) where $n_{\mathrm{m}} = n_{\mathrm{f}} = 10$. We ran these simulations with weaker c values than Figure 4 for computational tractability given the increase in segregation time which comes with a larger population size. Other parameter values for simulations: $h_{\mathrm{f}} = h_{\mathrm{m}} = 0.5$; see Appendix B for more details on simulation procedure).