1

Establishment and maintenance of motor neuron identity via temporal modularity in terminal selector function

Yinan Li [1, 2], Anthony Osuma [1, 2], Edgar Correa [1, 3], Munachiso A. Okebalama[1], Pauline Dao[1], Olivia Gaylord [4], Jihad Aburas [1], Priota Islam [5, 6], André E.X. Brown [5, 6], Paschalis Kratsios [1, 2, 3, 4, 7]

[1] Department of Neurobiology, University of Chicago, Chicago, IL, USA

[2] Committee on Neurobiology, University of Chicago, Chicago, IL, USA

[3] Cell and Molecular Biology Program, University of Chicago, Chicago, IL, USA

[4] Committee on Development, Regeneration and Stem Cell Biology, University of Chicago, Chicago, IL, USA

[5] MRC London Institute of Medical Sciences, London, UK

[6] Institute of Clinical Sciences, Imperial College London, London, UK

[7] The Grossman Institute for Neuroscience, Quantitative Biology, and Human Behavior, University of Chicago, Chicago, IL, USA

Correspondence: pkratsios@uchicago.edu

ABSTRACT (150 words)

Terminal selectors are transcription factors (TFs) that establish during development and maintain throughout life post-mitotic neuronal identity. We previously showed that UNC-3/Ebf, the terminal selector of *C. elegans* cholinergic motor neurons (MNs), acts indirectly to prevent alternative neuronal identities (Feng et al., 2020). Here, we globally identify the direct targets of UNC-3. Unexpectedly, we find that the suite of UNC-3 targets in MNs is modified across different life stages, revealing "temporal modularity" in terminal selector function. In all larval and adult stages examined, UNC-3 is required for continuous expression of various protein classes (e.g., receptors, transporters) critical for MN function. However, only in late larvae and adults, UNC-3 is required to maintain expression of MN-specific TFs. Minimal disruption of UNC-3's temporal modularity via genome engineering affects locomotion. Another *C. elegans* terminal selector (UNC-30/Pitx) also exhibits temporal modularity, supporting the potential generality of this mechanism for the control of neuronal identity.

INTRODUCTION

Nervous system development is a multi-step process that culminates in the generation of distinct neuron types necessary for animal behavior. Seminal studies in many model systems have begun to elucidate the molecular mechanisms that control the early steps of neuronal development, such as specification of neural progenitors and generation of post-mitotic neurons (Catela and Kratsios, 2019, Doe, 2017, Greig et al., 2013, Jessell, 2000, Lodato and Arlotta, 2015, Perry et al., 2017). However, our understanding of the final steps of neuronal development is very rudimentary. Once neurons become post-mitotic, how do they acquire their unique functional features, such as neurotransmitter synthesis, electrical activity and signaling properties? And, perhaps more importantly, how do neurons maintain such features throughout post-embryonic life?

Terminal selectors represent one class of transcription factors (TFs) with continuous expression – from development through adulthood – in specific neuron types (Hobert, 2008, Hobert, 2011, Hobert, 2016b, Garcia-Bellido, 1975). A defining feature of terminal selectors is the ability to activate the expression of effector genes, whose protein products determine the terminally differentiated state, and thereby function, of a given neuron type. Such effector genes, herein referred to as "terminal identity genes", are expressed continuously in specific neuron types and endow them with distinct functional and phenotypic properties. Examples include neurotransmitter (NT) biosynthesis components, NT receptors, ion channels, neuropeptides, and adhesion molecules (Hobert, 2008, Hobert, 2011, Hobert, 2016b). Numerous studies support the idea that terminal selectors establish during development and maintain throughout life neuronal identity (and function) by activating expression of terminal identity genes (Deneris and Hobert, 2014, Hobert, 2008, Hobert, 2016b, Hobert and Kratsios, 2019). Multiple cases of terminal selectors have been described thus far in worms, flies, chordates and mice (Deneris and Hobert, 2014, Hobert and Kratsios, 2019, Konstantinides et al., 2018), suggesting high conservation of this type of regulators. Importantly, human mutations in terminal selectors and their effector target genes have been linked to either developmental or degenerative conditions of the nervous system (Deneris and Hobert, 2014).

However, the molecular mechanisms through which terminal selectors establish and maintain neuronal identity are poorly understood, in part due to two major challenges. First, the majority of studies follow a candidate approach focused on a

select subset of terminal identity genes (Flames and Hobert, 2009, Hobert, 2016a,

Lopes et al., 2012). Hence, the extent of terminal identity features and breadth of

biological processes controlled by terminal selectors remain largely unknown.

Addressing this knowledge gap requires unbiased methods for the identification of

terminal selector target genes, but such approaches have only been applied to a limited

number of terminal selectors to date (Corbo et al., 2010, Housset et al., 2013,

Kadkhodaei et al., 2013, Wyler et al., 2016, Yu et al., 2017). Second, the continuous

expression of terminal selectors represents an additional challenge because it is not

known whether their function - in a particular neuron type - remains the same, or

changes at different life stages. This is partly due to the difficulty of tracking individual

neurons in the complex vertebrate nervous system throughout embryonic and post-natal

life. Hence, longitudinal studies in simple model organisms are needed to determine

whether terminal selectors control an identical suite of target genes across different

stages (e.g., embryo, adult), or whether the suite of targets can change over time.

Addressing these two challenges may extend our knowledge of how terminal selectors

control neuronal identity, as well as advance our understanding of how cellular identity

is established and maintained.

This study focuses on UNC-3, the sole *C. elegans* ortholog of the Collier/Olf/Ebf

(COE) family of TFs (Dubois and Vincent, 2001). UNC-3 acts as a terminal selector in

cholinergic motor neurons (MNs) of the *C. elegans* ventral nerve cord (Kratsios et al.,

2012). Importantly, mutations in EBF3, a human ortholog of UNC-3, cause a

neurodevelopmental syndrome characterized by motor developmental delay (Blackburn

et al., 2017, Chao et al., 2017, Harms et al., 2017, Sleven et al., 2017). A previous study

proposed that UNC-3 controls cholinergic MN identity in *C. elegans* by activating directly

the expression of various terminal identity genes (e.g., acetylcholine biosynthesis

components, ion channels, NT receptors, neuropeptides), which were identified via a

candidate approach (Kratsios et al., 2012). More recently, it was demonstrated that

UNC-3 can also act indirectly to prevent the adoption of alternative neuronal identities

(Feng et al., 2020). Lastly, animals lacking *unc-3* gene activity display severe

locomotion defects (Brenner, 1974, Feng et al., 2020), suggesting UNC-3 may broadly

control gene expression in cholinergic MNs. However, an unbiased identification of

UNC-3 targets, as well as a longitudinal analysis of *unc-3* mutants are currently lacking.

Here, we employ chromatin immunoprecipitation followed by DNA sequencing

(ChIP-Seq) and report the identification of ~3,500 protein-coding genes as putative

direct targets of UNC-3. Protein class ontology analysis suggests that UNC-3, besides

terminal identity genes, also controls additional biological processes, such as neuronal

metabolism and downstream gene regulatory networks comprised of numerous TFs and

nucleic acid-binding proteins. These findings help obtain a comprehensive

understanding of terminal selector function.

Through a longitudinal analysis of *unc-3* mutants at embryonic, larval and adult

stages, we identified two groups of target genes with distinct temporal requirements for

UNC-3 in cholinergic MNs. One group encodes multiple classes of proteins (e.g.,

receptors, secreted molecules, TFs) that require UNC-3 for both embryonic initiation

and post-embryonic maintenance of their expression. Contrasting this stable mode of

regulation over time, a second group of targets consists exclusively of TFs (*cfi-1/Arid3a,*

bnc-1/BNC1-2, mab-9/Tbx20, ceh-44/CUX1-2, nhr-40/nuclear hormone receptor) that

do not require UNC-3 for initiation, but depend on UNC-3 activity for maintenance.

Hence, the suite of UNC-3 targets in cholinergic MNs is modified across different life

stages, a phenomenon we term "temporal modularity" in terminal selector function. To

provide mechanistic insights, we focused on the second group of targets and identified

a molecular mechanism for their *unc-3*-independent initiation that relies on Hox proteins.

Importantly, preventing UNC-3's ability to selectively maintain expression of a single TF

(*cfi-1/Arid3a*) from the second target group led to locomotion defects, indicating minimal

disruption of temporal modularity affects animal behavior. Lastly, we provide evidence

for temporal modularity in the function of UNC-30/PITX, the terminal selector of

GABAergic MN identity in *C. elegans* (Eastman et al., 1999, Jin et al., 1994). Because

terminal selectors have been identified in both invertebrate and vertebrate nervous

systems (Deneris and Hobert, 2014, Hobert and Kratsios, 2019), we hypothesize that

temporal modularity in their function may be a general mechanism for the establishment

and maintenance of neuronal identity.

RESULTS

Identifying the global targets of UNC-3 via ChIP-Seq

To identify in an unbiased manner putative UNC-3 target genes, we employed chromatin immunoprecipitation followed by DNA sequencing (ChIP-Seq). We used a reporter strain with in-frame GFP sequences inserted immediately upstream of the stop codon of the endogenous *unc-3* gene (**Figure 1A**). Expression of UNC-3::GFP fusion protein was observed in the nucleus of 53 cholinergic MNs (SAB subtype = 3 neurons, DA = 9, DB = 7, VA = 12, VB = 11, AS = 11) and 19 other neurons known to express *unc-3* (**Figure 1A**), indicating that this reporter faithfully recapitulates the endogenous *unc-3* expression pattern (Pereira et al., 2015, Prasad et al., 1998). Insertion of GFP does not detectably alter the function of UNC-3 since expression of known UNC-3 targets (*cho-1/ChT, unc-17/VAChT, ace-2/AChE*) is unaffected in *unc-3::gfp* animals (**Figure 1B**). Unlike *unc-3* null mutants, *unc-3::gfp* animals do not display locomotion defects. We therefore performed ChIP-seq on *unc-3::gfp* animals at larval stage 2 (L2) because all *unc-3* expressing neurons have been generated by this stage.

Our ChIP-seq dataset revealed strong enrichment of UNC-3 binding in the genome by identifying a total of 6,892 unique binding peaks (q-value cutoff: 0.05) (**Figure 1C** and **Figure 1 – figure supplement 1**). The majority of UNC-3 binding peaks (91.95%) are predominantly located between 0 and 3 kb upstream of a transcription start site (Kadkhodaei et al.), whereas only a small fraction is found in introns (2.05 %) or downstream of a gene (0.99%), suggesting UNC-3 chiefly acts at promoters and enhancers to regulate gene expression (**Figure 1D** and **1F**). Through *de novo* motif discovery analysis (see Materials and Methods), we identified a 12 bp pseudo-palindromic sequence overrepresented in the UNC-3 binding peaks (**Figure 1E**), which highly resembles the binding site of UNC-3 vertebrate orthologs (Treiber et al., 2010a, Treiber et al., 2010b, Wang and Reed, 1993, Wang et al., 1997). To test the quality of our ChIP-Seq results, we sought to determine whether UNC-3 binding peaks are present in the *cis*-regulatory region of all previously described UNC-3 targets in cholinergic MNs, because these neurons constitute the majority of *unc-3*-expressing cells (53 out of 72 cells). Previous studies identified 10 terminal identity genes as putative direct UNC-3 targets and 43 terminal identity genes whose expression is affected by genetic removal of *unc-3* (Kratsios et al., 2015, Kratsios et al., 2012). In the current study, we found UNC-3 binding peaks in 9 out of 10 (90%) direct UNC-3 targets

and 38 of the 43 (88.37%) downstream targets of UNC-3, indicating high quality in the

ChIP-Seq results (**Figure 2A** and **Figure 1 – figure supplement 1, Supplementary**

File 1). Moreover, ChIP-Seq for UNC-3 appears highly sensitive, as it identified peaks in

unc-3-dependent genes expressed in a limited number neurons (e.g., *glr-4/GluR* is

expressed in 4 out of 72 *unc-3+* neurons) (**Figure 2A**). In conclusion, our ChIP-Seq

experiment generated a comprehensive map of UNC-3 binding in the *C. elegans*

genome and provided biochemical evidence to the hypothesis that UNC-3 binds to the

cis-regulatory region of multiple terminal identity genes, consolidating UNC-3's function

as a terminal selector of cholinergic MN identity.

Our bioinformatic analysis of the UNC-3 binding peaks revealed 3,502 protein-

coding genes as putative UNC-3 targets (see Materials and Methods). To classify these

new targets, we performed gene ontology (GO) analysis focused on protein classes

using PANTHER (Mi et al., 2013). This analysis revealed three broad categories

(**Figure 2B, Supplementary File 2**). First, there is a preponderance (42.18 % of the

total number of UNC-3 targets classified by PANTHER) of terminal identity genes (e.g.,

114 transporter proteins, 111 receptors and trans-membrane proteins, 37 signaling

molecules, 11 cell adhesion molecules), suggesting that UNC-3 broadly affects multiple

features of neuronal terminal identity. The second overrepresented category (24.07% of

UNC-3 targets) contains a large number of proteins involved in the control of gene

expression, such as 239 nucleic acid binding proteins (16.77 %) and 104 TFs (7.3 %),

highlighting the possibility of an extensive network of gene regulatory factors

downstream of UNC-3. The third category (24.14 %) consists of genes coding for

various types of enzymes (e.g., hydrolases, ligases, oxidoreductases), suggesting a

new role for UNC-3 in neuronal metabolic pathways. Together, this analysis unravels

the breadth of biological processes potentially under the direct control of UNC-3.

The ChIP-Seq experiment was performed on endogenously tagged UNC-3, which

is expressed in 53 cholinergic MNs of the nerve cord and 19 other neurons located in

the *C. elegans* head and tail (Pereira et al., 2015). Since MNs are the majority (53 cells)

of *unc-3*-expressing cells (72 in total), a significant portion of the UNC-3-bound genes

may be expressed in MNs. To test this, we used available single-cell RNA-Seq data

(CeNGEN project: www.cengen.org) that identified 576 transcripts specifically enriched

in cholinergic MNs (Seth R Taylor, 2019). We found that 52.95% of these MN-

expressed genes are bound by UNC-3 and fall in the aforementioned gene categories

(**Supplementary File 3**), thereby constituting putative UNC-3 targets in cholinergic

MNs.

***Cis*-regulatory analysis reveals novel TFs as direct UNC-3 targets in motor**

neurons

Our ChIP-Seq results provide an opportunity to reveal new roles for UNC-3, beyond the

direct control of terminal identity genes. To this end, we focused on the 104 TFs

identified by ChIP-Seq as putative UNC-3 targets (**Figure 2B-C**). To functionally test

whether the UNC-3 bound DNA elements upstream of these TF-encoding genes carry

information critical for gene expression, we carried out a *cis*-regulatory analysis. We

isolated and fused to RFP 15 elements located upstream of 15 randomly selected TFs

of different families (e.g., homeobox, nuclear hormone receptors, Zn finger) (**Figure 2C,**

Table 1). We generated transgenic reporter animals and found that 10 of these *cis*-

regulatory elements were sufficient to drive RFP expression in ventral cord cholinergic

MNs (**Table 1**). Expression of five TFs (*nhr-1, nhr-19, nhr-49, zfh-2, ztf-26*) in ventral

cord MNs has not been previously described. The remaining 5 TFs (*cfi-1/Arid3a, bnc-*

1/Bnc1/2, mab-9/Tbx20, nhr-40, ceh-44/Cux1) are known to be expressed in subsets of

unc-3-positive MNs (Kerk et al., 2017, Pocock et al., 2008, Brozova et al., 2006), but our

analysis revealed the *cis*-regulatory elements that are sufficient for their MN expression.

Of note, some of these TF reporters are also expressed in *unc-3*-negative MNs of the

nerve cord, namely the GABAergic DD and VD neurons (**Table 1**). Next, we tested for

unc-3 dependency at the L4 stage, and found that 9 of these 10 TF reporters depend on

unc-3 for proper expression in cholinergic MNs (**Table 1**). Eight reporters (*nhr-1, nhr-40,*

zfh-2, ztf-26, ceh-44, cfi-1, bnc-1, mab-9) are positively controlled by *unc-3,* where*as*

one (*nhr-49*) is negatively regulated by *unc-3* (**Figure 3 – figure supplement 1, Table**

1). Together, this *cis*-regulatory analysis revealed novel TFs as direct UNC-3 targets in

cholinergic MNs. Besides its known role as an activator of terminal identity genes

(Kratsios et al., 2015, Kratsios et al., 2012), these findings suggest that UNC-3 can also

act directly to either activate or repress expression of multiple TF-encoding genes,

uncovering an extensive gene regulatory network downstream of UNC-3.

Temporal modularity of UNC-3 function in cholinergic motor neurons

Terminal selectors are continuously expressed, from development through adulthood, in

specific neuron types. However, it remains unclear whether – in the same neuron type –

a terminal selector controls an identical suite of targets across different life stages, or

the suite of targets can change over time. The case of UNC-3 offers an opportunity to

address this issue because its direct targets (terminal identity genes and newly

identified TFs [**Table 1**]) are continuously expressed in cholinergic MNs. However, *unc-*

3 dependency of terminal identity genes was mostly tested at a single developmental

stage, the last larval stage (L4) (Kratsios et al., 2012). We therefore performed a

longitudinal analysis to determine whether target gene dependency on *unc-3* remains

stable or changes at different life stages.

First, we tested 4 terminal identity genes (*acr-2/AChR, unc-129/TGFbeta, glr-*

4/GluR, unc-17/VAChT) at larval (L2, L4) and adult (day 1) stages and found that their

MN expression, at every stage, critically depends on UNC-3 (**Figure 3A**). However, a

different picture emerged after testing the 8 TF reporters that are positively regulated by

UNC-3 (**Table 1**). Similar to terminal identity genes, the expression of 3 TFs (*zfh-*

2/Zfhx3, ztf-26, nhr-1) critically depends on UNC-3 at every stage (L2, L4, adult) (**Figure**

3 – figure supplement 1), suggesting UNC-3 controls initiation and maintenance of

their expression. In striking contrast, the early expression (L2 stage) of five TFs (*cfi-1,*

bnc-1, mab-9, nhr-40, ceh-44) does not require UNC-3 (**Figure 3B**). However,

maintenance of their expression during late larval and/or adult stages does depend on

UNC-3 (**Figure 3B**). Hence, this longitudinal analysis revealed two groups of targets

with distinct requirements for UNC-3 at different life stages. One group consists of

terminal identity genes (*acr-2/AChR, unc-129/TGFbeta, glr-4/GluR, unc-17/VAChT*) and

TFs (*zfh-2/Zfhx3, ztf-26, nhr-1*) that require UNC-3 for both initiation and maintenance of

expression ("initiation and maintenance" module, **Figure 3A, C**). The second group

consists exclusively of TFs (*cfi-1/Arid3a, bnc-1/Bnc1, mab-9/Tbx20, ceh-44/Cux1, nhr-*

40) that depend on UNC-3 for maintenance, but not initiation ("maintenance-only"

module, **Figure 3B-C**).

Collectively, these findings suggest that, in cholinergic MNs, the suite of UNC-3

target genes can change at different life stages; distinct UNC-3 targets are controlled in

a modular manner over time ("initiation and maintenance" or "maintenance-only"

modules). To describe this phenomenon, we use the term "temporal modularity in UNC-

3 function" given that the function of a cell type-specific TF at a particular life stage is

determined by the suite of targets it controls at that stage (**Figure 3C**). In the following

sections, we hone in on a single target (*cfi-1/Arid3a*) from the "maintenance-only"

module, aiming to dissect the molecular mechanisms underlying the temporal

modularity of UNC-3 function in cholinergic MNs.

A distal enhancer is necessary for initiation and maintenance of *cfi-1/Arid3a*

expression in MNs

Our *cis*-regulatory analysis suggests that maintenance, but not initiation, of *cfi-1*

expression depends on UNC-3 (**Figure 3B**). We therefore hypothesized that the sole

UNC-3 binding peak on the *cfi-1* locus (located ~12 kb upstream) demarcates an

enhancer element selectively required for maintenance (**Figure 3B** and **4A**). If this were

to be the case, then it would be logical to assume that a separate *cis*-regulatory element

would control *cfi-1* initiation in MNs. To test this assumption, we conducted an unbiased

cis-regulatory analysis *in vivo* by generating a series of 12 transgenic reporter (GFP or

RFP) animals, with each reporter carrying small and contiguous DNA fragments

spanning a ~15 kb region (**Figure 4A**). Surprisingly, this analysis did not reveal a

separate initiation element. Instead, it identified a 2.5 kb distal element (reporter #7)

demarcated by the UNC-3 binding peak (**Figure 4A**) that drives RFP expression in

subsets of *unc-3*-expressing cholinergic MNs (DA, DB, VA, VB subtypes) at embryonic

(3-fold), larval and adult stages (**Figure 4A, C**). In addition, this 2.5 kb element also

showed expression at all these stages in *unc-3*-negative neurons of the nerve cord,

namely the GABAergic (DD, VD subtypes) MNs (**Figure 4A, C**), which will be discussed

later in Results. We conclude that this enhancer element is sufficient for initiation and

maintenance of *cfi-1* reporter expression in nerve cord MNs.

To test the necessity of this element, we first generated via CRISPR/Cas9 an

endogenous mNeonGreen (mNG) reporter allele for *cfi-1,* which also carries an auxin-

inducible degron (AID) tag (*mNG::AID::cfi-1*), enabling inducible depletion of CFI-1

(depletion experiments are described later in Results). Animals carrying the

mNG::AID::cfi-1 AID allele do not show any developmental phenotypes, suggesting that

the mNG::AID tag does not detectably alter *cfi-1* activity. This reporter showed

expression in subsets of *unc-3*-expressing MNs (DA, DB, VA, VB subtypes), GABAergic

nerve cord MNs (DD, VD subtypes), tail and head neurons, as well as head muscle

(**Figure 4A-C**), a pattern consistent with previous studies describing *cfi-1* expression

(Shaham and Bargmann, 2002, Kerk et al., 2017). We determined the onset of the

endogenous *cfi-1* reporter (*mNG::AID::cfi-1)* in MNs to be at the 3-fold embryonic stage,

coinciding with the onset of transgenic reporters (#7 and #8) containing the distal

enhancer (**Figure 4B**). Next, we employed CRISPR/Cas9 genome editing and deleted

769 bp that constitute the core of the UNC-3 binding peak (located ~12 kb upstream) in

the context of the endogenous *cfi-1* reporter (*mNG::AID::cfi-1$^{\Delta\ enhancer\ (769\ bp)}$*). We found

that *mNG::AID::cfi-1* expression is selectively eliminated in cholinergic (DA, DB, VA, VB)

and GABAergic (DD, VD) nerve cord MNs at all life stages examined (3-fold embryo,

L4, Day 1 adult) (**Figure 4A-B**, quantification of cholinergic MNs shown in **4D**).

We conclude that a distal enhancer (located ~12 kb upstream of *cfi-1*) is both

necessary and sufficient for *cfi-1* expression in nerve cord MNs. Genome editing

suggests that a 769 bp sequence within this enhancer is required for both initiation and

maintenance of *cfi-1* in nerve cord MNs (**Figure 4D**). In the ensuing sections, we test

the hypothesis that this enhancer integrates UNC-3 input for *cfi-1* maintenance in

cholinergic MNs, as well as UNC-3-independent input for *cfi-1* initiation in these

neurons.

UNC-3 maintains *cfi-1* expression in cholinergic MNs via direct activation of the

distal enhancer

The binding of UNC-3 to the distal enhancer strongly suggests UNC-3 acts directly to

maintain *cfi-1* expression in cholinergic MNs (**Figure 4A**). However, the UNC-3 peak is

spread across several hundred base pairs due to the inherently low ChIP-Seq

resolution; hence, the precise DNA sequences recognized by UNC-3 remained

unknown. Through bioinformatic analysis (see Materials and Methods), we identified 8

putative UNC-3 binding sites (COE motifs) within the 769 bp distal enhancer (**Figure**

4E). Using CRISPR/Cas9 technology, we simultaneously mutated all 8 motifs in the

context of the endogenous *cfi-1* reporter allele (*mNG::AID::cfi-1$^{8\ COE\ motifs\ mut}$*) by

substituting nucleotides known to be critical for DNA binding of UNC-3 orthologs

(Treiber et al., 2010a, Wang et al., 1993) (**Figure 4E**). During the L2 stage, expression

of *mNG* in MNs is not affected in *mNG::AID::cfi-1 $^{8\ COE\ motifs\ mut}$* animals, indicating early

cfi-1 expression occurs normally (**Figure 4F**). Intriguingly, *mNG* expression is

significantly down-regulated in cholinergic MNs at later larval (L4) and adult (day 1)

stages, resembling the phenotype of *unc-3* null mutants (**Figure 4F**). We conclude that

UNC-3 binds to the distal enhancer and directly acts through one or more of these 8

COE motifs to maintain *cfi-1* expression in cholinergic MNs.

Previous studies in the nervous system have shown that a TF can maintain its

own expression via transcriptional activation either by itself (positive auto-regulation), or

in partnership with other TFs (Leyva-Diaz and Hobert, 2019, Scott et al., 2005, Xue et

al., 1992). We found though that *cfi-1* does not auto-regulate and UNC-3 binding at the

distal enhancer occurs normally in *cfi-1* null mutants (**Figure 4 – figure supplement 1**),

excluding a potential involvement of CFI-1 in its own maintenance.

LIN-39 (Scr/Dfd/Hox4-5) and MAB-5 (Antp/Hox6-8) control *cfi-1* expression in

cholinergic MNs through the same distal enhancer

If UNC-3 exerts a maintenance role, what are the factors that initiate *cfi-1* expression in

MNs? Previous work showed that two Hox proteins, LIN-39 (Scr/Dfd/Hox4-5) and MAB-

5 (Antp/Hox6-8), control *cfi-1* expression in MNs (Kratsios et al., 2017). However,

analysis was performed at the last larval stage (L4) and transgenic *cfi-1* reporter

animals were used. Hence, it is unclear whether LIN-39 and MAB-5 are required for

initiation of endogenous *cfi-1*.

Because *lin-39; mab-5* double null mutants are viable (Liu and Fire, 2000), we

performed a longitudinal analysis and found that expression of the endogenous

mNG::AID::cfi-1 reporter in cholinergic MNs is severely affected at embryonic (3-fold),

larval (L2, L4) and adult (D1) stages (**Figure 5A-C**). Since onset of *cfi-1* expression

occurs at the 3-fold embryonic stage (**Figure 4B**), these results suggest LIN-39 and

MAB-5 are required for *cfi-1* initiation. Conversely, initiation of *cfi-1* expression is not

affected in a null mutant of *unc-3* (**Figure 5B-C**). Moreover, available ChIP-Seq data

from modENCODE (Boyle et al., 2014) indicate that LIN-39 and MAB-5 bind to the *cfi-1*

distal enhancer (**Figure 5A**). Expression of a 2.5 kb transgenic *cfi-1* reporter (reporter

#7) that carries the distal enhancer is significantly affected in *lin-39; mab-5* double

mutants at early larval (L2) stages (**Figure 5A, D**). These results strongly suggest that

LIN-39 and MAB-5 activate *cfi-1* expression directly. Although the DNA sequence of the

MAB-5 binding site is not known, mutation of a single, bioinformatically predicted LIN-39

binding site (wild-type: aaTTGAtg > mutated: aaGGGGtg) within the enhancer led to a

decrease in reporter gene expression at L2 (**Figure 5A, E**). This decrease was weaker

compared to *lin-39; mab-5* double mutants (**Figure 5C**), likely due to compensation by

MAB-5. Indeed, LIN-39 and MAB-5 appear to act synergistically because endogenous *cfi-1* expression (*mNG::AID::cfi-1*) is mildly affected in *lin-39* single mutants, but severely affected in *lin-39; mab-5* double mutants (**Figure 5F**). We conclude that the Hox proteins LIN-39 and MAB-5 are necessary for *cfi-1* initiation in cholinergic MNs (left panel, **Figure 5G**), and act through the same distal enhancer utilized by UNC-3 to maintain *cfi-1* (right panel, **Figure 5G**).

If Hox proteins were only required for initiation, then endogenous *cfi-1* expression should not be affected at later stages (L2, L4, D1 adults) in *lin-39; mab-5* double mutants. However, this is not the case (**Figure 5B-C**), suggesting that Hox proteins play an additional role at later stages. One possibility is that these proteins, like UNC-3, are also required for maintenance. To test this, we used the auxin-inducible degradation (AID) system and depleted the endogenous LIN-39 protein at the last larval stage (L4) by using a previously described *lin-39::mNG::AID* allele (Feng et al., 2020, Zhang et al., 2015). However, expression of *cfi-1* was unaffected in the adult (**Figure 5 – figure supplement 1**). This negative result could be attributed to low and undetectable levels of LIN-39 and/or functional compensation by MAB-5. We therefore used *lin-39; mab-5* double (null) mutants and crossed them to *unc-3* null animals. If Hox proteins, similar to UNC-3, are required for *cfi-1* maintenance in cholinergic MNs, stronger effects should be present in *unc-3*; *lin-39; mab-5* triple mutants compared to *unc-3* single mutants. Indeed, we found this to be the case in day 1 adult animals (graph on the right, **Figure 5C**). Supporting a maintenance role for Hox, mutation of the LIN-39 binding site within the *cfi-1* enhancer (reporter #8) led to a sustained decrease in reporter gene expression from L2 to adult stages (**Figure 5E**).

Together, our findings suggest the Hox proteins LIN-39 and MAB-5 control initiation and maintenance of *cfi-1* in cholinergic MNs via the same distal *cis*-regulatory region (enhancer) utilized by UNC-3 to maintain *cfi-1* (**Figure 5G**). However, this region bears distinct UNC-3 and LIN-39 binding sites.

cfi-1/Arid3a is required post-embryonically to maintain MN subtype identity

Expression of *cfi-1* in cholinergic (DA, DB, VA, VB) MNs is maintained throughout life by Hox and UNC-3 (**Figure 5G**). But why is it important to ensure continuous *cfi-1* expression? Although its function in VA and VB remains unknown, CFI-1 is required during early development to establish the identity of DA and DB subtypes by acting as a

transcriptional repressor (Kerk et al., 2017). In *cfi-1* null animals, glutamate receptor

subunit 4 (*glr-4/GluR),* a terminal identity gene normally activated by UNC-3 in another

MN subtype (SAB), becomes ectopically expressed in DA and DB neurons (**Figure 5 –**

figure supplement 2). Hence, early removal of *cfi-1* leads to DA and DB neurons

adopting a mixed identity. Whether CFI-1 is required post-embryonically to continuously

prevent DA and DB from obtaining a mixed identity is not known.

To enable CFI-1 protein depletion selectively at post-embryonic stages, we used

the *mNG::AID::cfi-1* reporter allele (**Figure 4A**), which also serves as a conditional allele

as it carries the auxin-inducible degron (AID) (Zhang et al., 2015). Auxin administration

at the first larval (L1) stage resulted in efficient depletion of *mNG::AID::cfi-1* expression,

which was undetectable 2 days later (**Figure 5 – figure supplement 2**). At the L3 stage

(2 days upon continuous auxin treatment), we observed ectopic expression of *glr-4* in

DA and DB neurons. These results suggest that CFI-1 is required post-embryonically to

prevent DA and DB neurons from adopting mixed identity, and underscore the critical

role of UNC-3 and Hox in maintaining *cfi-1* expression (**Figure 5G**).

Disruption of temporal modularity in UNC-3 function leads to locomotion defects

Animals carrying *unc-3* null alleles display severe locomotion defects (Feng et al.,

2020), likely due to combined defects in the expression of UNC-3 targets from both

"initiation and maintenance" and "maintenance-only" modules (**Figure 3C**). Genes from

the "initiation and maintenance" module include (among others) terminal identity genes

coding for ACh biosynthesis components (**Figure 3A**). Hence, it is conceivable that loss

of *unc-3* can lead to defects in ACh biosynthesis, likely contributing to locomotion

defects. However, it is unclear whether the expression of UNC-3 targets from the

"maintenance-only" module is critical for locomotion. To test this, we focused on c*fi-*

1/Arid3a and used the CRISPR-engineered allele (*mNG::AID::cfi-1* $^{8\ COE\ motifs\ mut}$) that

selectively affects maintenance, but not initiation, of *cfi-1* expression in cholinergic MNs

(**Figure 4E-F**). As controls, we used animals carrying: (**a**) the endogenous *cfi-1* reporter

allele (*mNG::AID::cfi-1),* (**b**) a putative null *cfi-1* allele (*ot786*) (Kerk et al., 2017), in

which *cfi-1* activity is affected in MNs and other neuron types of the motor circuit

(Pereira et al., 2015, Shaham and Bargmann, 2002), and (**c**) a deletion of the distal

enhancer (*mNG::AID::cfi-1$^{\Delta enhancer(769\ bp)}$*), in which both initiation and maintenance of *cfi-*

1 are abrogated in nerve cord MNs (**Figure 6A**). We performed a high-resolution

behavioral analysis of freely moving adult (day 2) animals of the above genotypes using automated multi-worm tracking technology (Javer et al., 2018b, Yemini et al., 2013). We found several features related to *C. elegans* locomotion (*e.g.,* body curvature, velocity) severely affected in *cfi-1 (ot786)* putative null animals (**Figure 6B-G**). Compared to *cfi-1* null mutants, animals carrying the *mNG::AID::cfi-1* [8 COE motifs mut] allele (selective disruption of *cfi-1* maintenance) display milder, but statistically significant locomotion defects in the adult (**Figure 6B-G**). As expected, these defects were also present in animals carrying the *mNG::AID::cfi-1* [Δenhancer (769 bp)] allele, in which both initiation and maintenance of *cfi-1* expression is affected. In summary, we specifically disrupted in cholinergic MNs the maintained expression of a single UNC-3 target (*cfi-1* from the "maintenance-only" module) and observed locomotion defects. This analysis suggests that minimal disruption of temporal modularity in UNC-3 function can affect animal behavior.

Hox proteins and UNC-3 control *bnc-1/BNC* expression in cholinergic motor neurons

Our *cis*-regulatory analysis suggested that five TFs (*cfi-1/Arid3a, bnc-1/BNC1-2, mab-9/Tbx20, ceh-44/CUX1-2, nhr-40/nuclear hormone receptor)* require UNC-3 selectively for maintenance (**Figure 3B**). An in-depth analysis of *cfi-1/Arid3a* revealed that Hox proteins (LIN-39, MAB-5) and UNC-3 ensure the continuous expression of *cfi-1* in subsets of *unc-3*-positive MNs (**Figure 3-5**). We next asked whether a similar mechanism applies to the regulation of *bnc-1*/BNC, which is also expressed in a subset of *unc-3*-positive MNs (VA, VB) and prevents them from adopting a mixed identity (**Figure 5 – figure supplement 3**) (Kerk et al., 2017). Using an endogenous reporter allele (*bnc-1::mNG::AID*), we found that LIN-39 and MAB-5 are required for *bnc-1* expression at all stages examined (L2, L4, day 1 adult; **Figure 5 – figure supplement 3**), suggesting a role for Hox in *bnc-1* initiation and maintenance. Next, we found that UNC-3 is absolutely required for *bnc-1* maintenance in the adult (day 1), albeit weaker effects were also observed at L2 and L4 (**Figure 5 – figure supplement 3**). Similar to *cfi-1*, these findings strongly suggest that endogenous *bnc-1* expression depends on Hox and UNC-3.

Temporal modularity of UNC-30/PITX function in GABAergic motor neurons

Is temporal modularity observed in the function of other terminal selectors? To address this, we focused on UNC-30/PITX, the terminal selector of GABAergic MN (DD, VD) identity in the *C. elegans* nerve cord (**Figure 7A**) (Jin et al., 1994). UNC-30/PITX is known to directly activate the expression of several terminal identity genes (e.g., *unc-25*/GAD [glutamic acid decarboxylase], *unc-47*/VAGT [vesicular GABA transporter])(Eastman et al., 1999), but a longitudinal analysis of target gene expression in *unc-30* null animals is lacking. Using reporter strains and methodologies similar to those used for UNC-3, we found that terminal identity gene (*unc-25*/GAD, *unc-47*/VAGT) expression is affected in GABAergic MNs of *unc-30* mutants at all stages examined (3-fold embryo, L2, L4, adult [day 1]) (**Figure 7B, D**), suggesting a requirement for initiation and maintenance. As mentioned earlier (**Figure 4C**), *cfi-1* is also expressed in GABAergic MNs (**Figure 7A**). Since UNC-3 is required for *cfi-1* maintenance in cholinergic MNs, we asked whether UNC-30/PITX plays a similar role to maintain *cfi-1* expression in GABAergic MNs. We found no effect at the embryonic stage (3-fold), but progressively stronger effects at larval (L2, L4) and adult (day 1) stages (**Figure 7C**), indicating that, in GABAergic MNs, UNC-30/PITX is selectively required for the maintenance, but not initiation of *cfi-1* expression. Taken together, our findings indicate that, like UNC-3 in cholinergic MNs (**Figure 3C**), the function of UNC-30 in GABAergic MNs is organized into two modules (module #1: initiation and maintenance; module #2: maintenance-only) (**Figure 7D-E**), suggesting temporal modularity may be a shared feature among terminal selector type-TFs.

To gain mechanistic insights, we analyzed available UNC-30 ChIP-Seq data at the L2 stage (Yu et al., 2017). UNC-30 binds to the *cis*-regulatory region of terminal identity genes (*unc-25*/GAD, *unc-47*/VAGT) (**Figure 7B**), confirming previous observations (Eastman et al., 1999). UNC-30 also binds to the same distal enhancer of *cfi-1* in GABAergic MNs, as UNC-3 does in cholinergic MNs (**Figure 7C**). However, the UNC-30 binding sites are distinct from the UNC-3 sites in this enhancer (**Figure 7C**). CRISPR-mediated deletion of this enhancer abolished *cfi-1* expression in both cholinergic and GABAergic MNs (**Figure 4A-B**). This finding suggests that maintenance of *cfi-1* expression in two different neuron types relies on the same enhancer receiving UNC-30/PITX input in GABAergic MNs and UNC-3/EBF input in cholinergic MNs. Interestingly, these results provide an example of "enhancer pleiotropy" (Sabaris et al.,

499   2019), in which the same *cis*-regulatory element is used to control gene expression in
500   different neuron types.
501
502
503

DISCUSSION

Terminal selectors are continuously expressed TFs that determine the identity and function of individual neuron types (Hobert, 2008, Hobert, 2016b). However, the breadth of biological processes controlled by terminal selectors remains unclear. Moreover, whether terminal selectors control an identical suite of target genes across different life stages, or the suite of targets can change over time is largely unexplored. Filling such knowledge gaps can help us understand how cellular identity is established during development and maintained throughout life, a fundamental goal in the field of developmental biology. In this study, we focus on UNC-3/Ebf, the terminal selector of cholinergic MNs in the *C. elegans* nerve cord. Through ChIP-Seq, we identify in an unbiased manner the direct targets of UNC-3, uncovering the breadth of biological processes potentially controlled by this terminal selector. Unexpectedly, we find two groups of target genes with distinct temporal requirements for UNC-3 in cholinergic MNs. One group encodes different classes of proteins (e.g., receptors, transporters) that require UNC-3 activation at all life stages examined. However, a second group exclusively encodes MN-specific TFs that selectively require UNC-3 for maintenance of their expression at late larval and adult stages. Hence, the suite of UNC-3 targets in cholinergic MNs is modified across different life stages, a phenomenon we term "temporal modularity" in terminal selector function. Minimal disruption of this modularity by selectively removing the ability of UNC-3 to maintain expression of a single target gene (the TF *cfi-1/Arid3a* from the second group) led to locomotion defects, highlighting the necessity of temporal modularity for animal behavior.

Temporal modularity in terminal selector function may represent a general principle for neuronal subtype diversity

Why is there a need for temporal modularity in the function of continuously expressed TFs, such as terminal selectors? The case of UNC-3 suggests that temporal modularity is necessary for generating and maintaining neuronal subtype diversity. UNC-3 is continuously expressed in six MN subtypes (SAB, DA, DB, VA, VB, AS) of the *C. elegans* ventral nerve cord (**Figure 4C**) (Pereira et al., 2015). However, most of UNC-3 targets from either module (modules # 1 and #2, **Figure 3C**) are expressed in some, but not all, of these subtypes. By ensuring the continuous expression of genes from module #1 ("initiation and maintenance" module, **Figure 3C**), such as terminal identity genes

[(Kratsios et al., 2012) and this study], UNC-3 consolidates the unique identity of each

MN subtype. However, target genes from module #2 ("maintenance-only" module,

Figure 3C) escape initiation by UNC-3, but do require UNC-3 for maintenance.

Interestingly, all module #2 targets code for TFs (*cfi-1/Arid3a, bnc-1/BNC1-2, mab-*

9/Tbx20, ceh-44/Cux1, nhr-40), specifically expressed in subsets of the six subtypes

(SAB, DA, DB, VA, VB, AS) ((Kerk et al., 2017) and this study). Three of these factors

(*cfi-1/Arid3a, bnc-1/Bnc1, mab-9/Tbx20)* are thought to act as transcriptional repressors

to prevent the adoption of mixed MN identity (Kerk et al., 2017). For example, loss of *cfi-*

1 during early development results in "mixed" identity of DA and DB subtypes, as these

cells, in addition to their normal molecular signature, also acquire expression of genes

(e.g., *glr-4*/GluR) normally expressed in other MNs (**Figure 5 – figure supplement 2**).

Through protein depletion experiments at post-embryonic stages, we show here that

CFI-1 is continuously required to prevent DA and DB neurons from acquiring a mixed

identity. Hence, UNC-3 indirectly maintains the unique identity of individual MN

subtypes (DA, DB) by selectively maintaining expression of the TF *cfi-1* (from module

#2) during late larval and adult stages. In addition, UNC-3 acts directly to control DA

and DB identity by ensuring the continuous expression of *unc-129/TGFb*, a DA- and

DB-specific terminal identity gene (from module #1) (**Figure 3A**, **Figure 5 – figure**

supplement 2). Hence, temporal modularity in UNC-3 function can be envisioned as a

"double safety" mechanism for the generation and maintenance of MN diversity.

The mechanism of temporal modularity may represent a general paradigm of

how terminal selectors establish and maintain subtype identity within a class of

functionally related neurons. Indeed, the same terminal selector is often continuously

expressed in multiple subtypes of a given neuronal class (Hobert and Kratsios, 2019). In

mice, for example, the terminal selectors Nurr1 and Pet1 are expressed in several

subtypes of dopaminergic and serotonergic neurons, respectively (Kadkhodaei et al.,

2009, Okaty et al., 2015). However, future studies are needed to determine whether

temporal modularity in the function of these terminal selectors is necessary for

establishing and maintaining neuronal subtype identity.

Temporal modularity offers key insights into how terminal selectors control

neuronal identity over time

The prevailing hypothesis of how terminal selectors establish and maintain neuronal

identity is that they bind constantly, from development through adulthood, on the *cis-*

regulatory region of terminal identity genes, and thereby continuously activate their

expression (Deneris and Hobert, 2014, Hobert, 2008, Hobert and Kratsios, 2019). For

most terminal selectors, however, biochemical evidence for binding and longitudinal

analysis of terminal selector mutant animals are currently lacking. Our analysis on

module #1 genes (e.g., terminal identity genes) supports the aforementioned

hypothesis; ChIP-Seq for UNC-3 demonstrated binding on these genes (**Figure 3A, C**).

Moreover, our longitudinal genetic analysis together with a previous report that

selectively removed UNC-3 activity in adult MNs (Kerk et al., 2017) strongly suggest

that UNC-3 is continuously required to maintain terminal identity gene expression.

Interestingly, the analysis on module #2 genes provides new mechanistic insights

into how terminal selectors control neuronal identity over time. Instead of constantly

activating the same set of genes, as predicted by the above hypothesis, terminal

selectors can also modify the suite of their target genes at different life stages (temporal

modularity). This is based on the finding that UNC-3 is selectively required for

maintenance, not initiation, of five conserved TF-encoding genes from module #2 (*cfi-*

1/Arid3a, bnc-1/BNC1-2, mab-9/Tbx20, ceh-44/Cux1, nhr-40) (**Figure 3B-C**). Moreover,

by honing in on one TF (*cfi-1/Arid3a),* we identify a mechanism that enables initiation

and maintenance of module #2 genes (**Figure 5G**). That is, the Hox proteins LIN-39 and

MAB-5 control initiation of *cfi-1/Arid3a* in cholinergic MNs independently of UNC-3, but

cfi-1 maintenance in these same neurons depends on both Hox and UNC-3 (**Figure**

5G). Mechanistically, we propose that Hox-dependent initiation and Hox/UNC-3-

dependent maintenance of *cfi-1* are "funneled" through the same distal enhancer, which

bears both Hox and UNC-3 binding sites. This suggests embryonic initiation and post-

embryonic maintenance of expression of a particular gene, in a specific cell type, can be

achieved by distinct TF combinations acting upon the same *cis*-regulatory region

(enhancer) (**Figure 5G**). This somewhat surprising mechanism differs from previous fly

and mouse studies reporting distinct and physically separated *cis*-regulatory regions

necessary for either initiation or maintenance of cell type-specific gene expression

(Ellmeier et al., 2002, Johnson et al., 2011, Manzanares et al., 2001, Pfeffer et al.,

2002, Rhee et al., 2016).

In summary, our findings critically extend the mechanisms underlying UNC-3

function. Previous work demonstrated that, in cholinergic MNs, UNC-3 not only

activates expression of terminal identity genes (Kratsios et al., 2012), but also prevents

alternative neuronal identities (Feng et al., 2020). Here, we report that the suite of UNC-

3 targets in these neurons can be modified at different life stages, offering key insights

into how terminal selectors control neuronal identity over time.

Hox proteins collaborate with stage-specific TFs to establish and maintain MN

terminal identity

During early neuronal development, Hox proteins are essential for cell survival,

neuronal diversity, and circuit assembly (Baek et al., 2013, Catela et al., 2016, Estacio-

Gomez and Diaz-Benjumea, 2014, Karlsson et al., 2010, Miguel-Aliaga and Thor, 2004,

Moris-Sanz et al., 2015, Philippidou and Dasen, 2013). However, their post-embryonic

neuronal functions remain elusive. Moreover, Hox proteins are often continuously

expressed in multiple cell types of a given body region (Baek et al., 2013, Hutlet et al.,

2016, Takahashi et al., 2004), raising the question of how do they achieve cell type-

specificity in their function. Our findings on mid-body Hox proteins LIN-39 and MAB-5

begin to address this question.

619   LIN-39 and MAB-5 are continuously expressed in multiple cell types located at the

C. elegans mid-body region (Clark et al., 1993, Cowing and Kenyon, 1992, Feng et al.,

2020, Maloof and Kenyon, 1998). We find that LIN-39 and MAB-5 exert a cell type-

specific function; they are required to initiate (in embryo) and maintain (post-

embryonically) *cfi-1* expression in specific subsets of cholinergic MNs (DA, DB, VA, VB;

Figure 4C). Such specificity likely arises through collaboration with distinct TFs

responsible for either initiation or maintenance of *cfi-1* (**Figure 5G**). Supporting this

scenario, Hox (LIN-39, MAB-5) and UNC-3 are co-expressed in DA, DB, VA, and VB

neurons (Feng et al., 2020). Moreover, UNC-3 is selectively required for *cfi-1*

maintenance (not initiation) in these neurons (**Figure 5G**). We surmise that other, yet-

to-be-identified factors collaborate with Hox proteins at early stages to initiate *cfi-1*

expression specifically in DA, DB, VA, and VB neurons. Such initiation-specific factors

likely act through the same distal enhancer because its deletion completely abolished

cfi-1 expression in these neurons at early (and late) stages (**Figure 4D, 5G**). In

summary, we propose that Hox proteins collaborate with distinct TFs over time, that is,

initiation-specific factors and the terminal selector UNC-3, to ensure continuous

expression of *cfi-1/Arid3a* in specific subtypes of cholinergic MNs. This mechanism may

extend to the regulation of terminal identity genes, as we previously showed that LIN-39

and UNC-3 are required to maintain expression of *acr-2/* human CHRNA1

(acetylcholine receptor subunit) and *unc-77/* human NALCN (sodium channel) in

cholinergic MNs (Feng et al., 2020). Altogether, these findings offer mechanistic insights

into the recently proposed hypothesis that Hox proteins in *C. elegans* collaborate with

terminal selectors to establish and maintain neuronal terminal identity (Kratsios et al.,

2017, Zheng et al., 2015).

Limitations of this study

By conducting a longitudinal analysis for 14 UNC-3 target genes, we identified two

groups with distinct temporal requirements (modules #1 and #2 in **Figure 3C**). It is likely

though that UNC-3 temporally controls other targets though additional modules. For

example, animals lacking *unc-3* display severe axon guidance defects in cholinergic

MNs (Prasad et al., 1998), but the underlying mechanisms remain unknown. Since axon

guidance molecules are often expressed in a transient fashion during early neuronal

development, we speculate that UNC-3 may transiently activate expression of such

molecules, a possibility that would add another temporal module in UNC-3 function.

In addition, the breadth of biological processes controlled by terminal selectors

remains largely unknown. Our ChIP-Seq analysis potentially implicates UNC-3 in a

range of biological processes. First, this dataset significantly extends previous reports

on the role of UNC-3 in neuronal terminal identity (Kim et al., 2005, Kratsios et al., 2012)

by identifying hundreds of terminal identity genes (42.18% of total ChIP-Seq hits) as

putative UNC-3 targets. Second, the ChIP-Seq dataset suggests new roles for UNC-3 in

neuronal metabolic pathways (24.14% of UNC-3 target genes code for enzymes) and

gene regulatory networks (24.07% of UNC-3 targets are TFs and nucleic acid-binding

proteins). However, future RNA-Sequencing studies in *unc-3*-depleted MNs are

necessary to correlate gene expression changes with UNC-3 ChIP-Seq data, and

thereby identify *bona fide* targets and biological processes under UNC-3 control.

Temporal modularity may be a shared feature among continuously expressed TFs

This study suggests that the suite of targets of two *C. elegans* terminal selectors (UNC-

3/Ebf and UNC-30/Pitx) can be modified over time, providing evidence for temporal

modularity in their function. Given that terminal selectors, as well as other neuron type-

specific TFs with continuous expression, have been described in both invertebrate and
vertebrate species (Deneris and Hobert, 2014, Hobert and Kratsios, 2019, Mayer et al.,
2018, Mi et al., 2018), it will be interesting to determine the potential generality of the
temporal mechanism described here. Supporting this possibility, the terminal selector of
serotonergic neurons in mice (Pet-1) activates expression of serotonin biosynthesis
proteins during development, but appears to switch transcriptional targets at later life
stages (Wyler et al., 2016). Outside the nervous system, cell type-specific TFs with
continuous expression have been described in worms, flies and mice (Pikkarainen et
al., 2004, Soler et al., 2012, Wiesenfahrt et al., 2016, Zhou et al., 2017). Future studies
will determine whether the principle of temporal modularity is widely employed for the
control of cell type identity.
680
681
682

MATERIALS AND METHODS

C. elegans **strain culture**

Worms were grown at 20°C or 25°C on nematode growth media (NGM) plates supplied

with *E. coli* OP50 as food source (Brenner, 1974). A list of all *C. elegans* strains used is

provided in **Supplementary file 4**.

Generation of transgenic animals carrying transcriptional fusion reporters

Reporter gene fusions for *cis*-regulatory analyses and validation of newly identified

UNC-3 target genes were made with PCR fusion (Hobert, 2002). Genomic regions were

amplified and fused to the coding sequence of *tagrfp* followed by the *unc-54* 3' UTR. To

mutate the LIN-39 binding motif, the reporter fusion was first introduced into the pCR-

XL-TOPO vector using the TOPO XL PCR cloning kit (Invitrogen). Then, mutagenesis

PCR was performed, and single clones containing plasmids that carry the desired

mutation were isolated. PCR fusion DNA fragments were injected into young adult *pha-*

1(e2123) hermaphrodites at 50 ng/μl together with *pha-1* (pBX plasmid) as co-injection

marker (50 ng/μl).

Chromatin Immunoprecipitation (ChIP)

ChIP assay was performed as previously described (Yu et al. 2017; Zhong et al. 2010)

with the following modifications. Synchronized L1 *unc-3 (ot839[unc-3::gfp])* worms and

N2 worms were cultured on 10 cm plates seeded with OP50 at 20°C overnight. Late L2

worms were cross-linked and resuspended in FA buffer supplemented with protease

inhibitors (150 mM NaCl, 10 μl 0.1 M PMSF, 100 μl 10% SDS, 500 μl 20% N-Lauroyl

sarsosine sodium, 2 tablets of cOmplete ULTRA Protease Inhibitor Cocktail [Roche

Cat.# 05892970001] in 10ml FA buffer). For each IP experiment, 200 μl worm pellet was

collected. The sample was then sonicated using a Covaris S220 at the following

settings: 200 W Peak Incident Power, 20% Duty Factor, 200 Cycles per Burst for 1 min.

Samples were transferred to centrifuge tubes and spun at the highest speed for 15 min.

The supernatant was transferred to a new tube, and 5% of the material was saved as

input and stored at -20°C. The remainder was incubated with 2 μl GFP antibody (Abcam

Cat.# ab290) at 4°C overnight. Wild-type (N2) worms do not carry the GFP tag and

serve as negative control. The *unc-3 (ot839[unc-3::gfp])* CRIPSR generated allele was

used in order to immunoprecipitate the endogenous UNC-3 protein. On the next day, 20

ul Dynabeads Protein G (1004D) was added to the immunocomplex, which was then

incubated for 2 hr at 4°C. The beads then were washed at 4°C twice with 150 mM NaCl

FA buffer (5 min each), and once with 1M NaCl FA buffer (5 min). The beads were

transferred to a new centrifuge tube and washed twice with 500 mM NaCl FA buffer (10

min each), once with TEL buffer (0.25 M LiCl, 1% NP-40, 1% sodium deoxycholate,

1mM EDTA, 10 mM Tris-HCl, pH 8.0) for 10 min, and twice with TE buffer (5 min each).

The immunocomplex was then eluted in 200 µl elution buffer (1% SDS in TE with 250

mM NaCl) by incubating at 65°C for 20 min. The saved input samples were thawed and

treated with the ChIP samples as follows. One (1) µl of 20 mg/ml proteinase K was

added to each sample and the samples were incubated at 55°C for 2 hours then 65°C

overnight (12-20 hours) to reverse cross-link. The immonuprecipitated DNA was purified

with Ampure XP beads (A63881) according to manufacturer's instructions.

ChIP-sequencing data analysis

Unique reads were mapped to the *C. elegans* genome (ce10) with bowtie2 (Langmead

and Salzberg, 2012). Peak calling was then performed with MACS2 (minimum q-value

cutoff for peak detection: 0.005). For visualization purposes, the sequencing depth was

normalized to 1x genome coverage using bamCoverage provided by deepTools

(Ramirez et al., 2016), and peak signals were shown in IGV. Heatmap of peak coverage

in regard to UNC-3 enrichment center was generated with NGSplot (Shen et al., 2014).

The average profile of peaks binding to TSS region was generated with ChIPseeker (Yu

et al., 2015). To study the distribution of peaks genome-wide, the peaks were annotated

using annotatePeaks.pl provided by Homer (Heinz et al., 2010), and each peak was

assigned to a gene with the nearest TSS. For *de novo* motif discovery, sequences

containing 100bp around the centers of each peak (from -50bp to +50bp) were

extracted and supplied to findMotifsGenome.pl provided by Homer.

Protein Class Ontology analysis using PANTHER

Protein Class Ontology analysis was performed on 1,478 UNC-3-bound genes out of

the 3,502 protein-coding genes. The number of genes is significantly lower than the

number of peaks because Panther analysis only considers genes with known protein

class terms.

Targeted genome editing

The *cfi-1* endogenous reporter strain *kas16 [mNG::AID::cfi-1]* was generated by

employing CRISPR/Cas9 genome editing and inserting

the *mNG::3xFLAG::AID* cassette immediately after the ATG of *cfi-1*. The *cfi-1* enhancer

knock-out allele *mNG::AID::cfi-1*$^{\Delta enhancer\ (769\ bp)}$ was generated by using two guide RNAs

flanking the *cfi-1* enhancer to guide excision of the genomic region, which was then

followed by homology dependent repair (HDR) to create a 769 bp deletion (-11,329 bp

to -12,097 bp). The UNC-3 binding motif mutation allele *mNG::AID::cfi-1*$^{8\ COE\ motifs}$

^{mut} was generated by creating nucleotide substitutions in the repair template, which

carries homology arms complementary to the *cfi-1* enhancer region and is then

introduced into the genome through HDR.

Microscopy

Imaging slides were prepared by anesthetizing worms with sodium azide (NaN$_3$, 100

mM) and mounting them on a 4% agarose pad on glass slides. Images were taken with

an automated fluorescence microscope (Zeiss, Axio Imager Z2). Images containing

several z stacks (0.50 μm intervals between stacks) were taken with Zeiss Axiocam 503

mono using the ZEN software (Version 2.3.69.1000, Blue edition). Representative

images are shown following max-projection of 2-5 μm Z-stacks using the maximum

intensity projection type. Image reconstruction was performed with Image J (Schindelin

et al., 2012).

Motor neuron subtype identification

The identification of specific MN subtypes expressing a given UNC-3 target gene was

assessed based on the following: (a) co-localization with fluorescent reporters that label

specific MN subtypes; (b) Invariant position of neuronal cell bodies along the ventral

nerve cord; (c) Birth order of specific motor neuron subtypes (e.g. during

embryogenesis or post-embryogenesis); (d) Total cell numbers in each motor neuron

subtype.

Bioinformatic prediction of binding motifs

Information of the LIN-39 binding motif is curated in the Catalog of Inferred Sequence

Binding Preferences database (http://cisbp.ccbr.utoronto.ca). To predict and identify

LIN-39 binding motifs and UNC-3 binding motifs (identified in this paper) in the *cfi-1*

enhancer (-11,391 bp to -12,146 bp), we utilized tools provided by MEME (Multiple

Expectation maximization for Motif Elicitation) bioinformatics suite (http://meme-

suite.org/), and performed FIMO (Find Individual Motif Occurrences) motif scanning

analysis.

Temporally controlled protein degradation

Temporally controlled protein degradation was achieved with the auxin-inducible

degradation system (Zhang et al., 2015). TIR1 expression was driven by the

ubiquitously active *eft-3* promoter in the transgene *ieSi57 [eft-3prom::tir1]* or a

transgene that drives TIR1 selectively in neurons (*otTi28*). To induce degradation of

LIN-39 and CFI-1 proteins, the following alleles were used: *lin-39 (kas9 [lin-*

39::mNG::AID]), and cfi-1 (kas16 [mNG::AID::cfi-1]). L1 or L4 worms were grown at 20

˚C on NGM plates coated with 4 nM auxin (indole-3-acetic acid [IAA] dissolved in

ethanol) or ethanol (negative control) for 1 day or 4 days before tested (see figure

legends for more details). All plates were shielded from light.

Worm tracking

Worms were maintained as mixed stage populations by chunking on NGM plates with

E. coli OP50 as the food source. Worms were bleached and the eggs were allowed to

hatch in M9 buffer to arrest as L1 larvae. L1 larvae were refed on OP50 and allowed to

grow to day 2 or adulthood. On the day of tracking, five worms were picked from the

incubated plates to each of the imaging plates (see below) and allowed to habituate for

30 minutes before recording for 15 minutes. Imaging plates are 35 mm plates with 3.5

mL of low-peptone (0.013% Difco Bacto) NGM agar (2% Bio/Agar, BioGene) to limit

bacteria growth. Plates are stored at 4°C for at least two days before use. Imaging

plates are seeded with 50µl of a 1:10 dilution of OP50 in M9 the day before tracking and

left to dry overnight with the lid on at room temperature.

Behavioral feature extraction and analysis

All videos were analyzed using Tierpsy Tracker to extract each worm's position and

posture over time (Javer et al., 2018a). These postural data were then converted into a

set of behavioral features selected from a large set of features as previously described

(Javer et al., 2018b). For each strain comparison, we performed unpaired two-sample t-

tests independently for each feature. The false discovery rate was controlled at 5%

across all strain and feature comparisons using the Benjamini Yekutieli procedure.

Statistical analysis

For data quantification, graphs show values expressed as mean ± standard error mean

(SEM) of animals. The statistical analyses were performed using the unpaired t-test

(two-tailed). Calculations were performed using the GraphPad QuickCalcs online

software (http://www.graph- pad.com/quickcalcs/). Differences with $p<0.05$ were

considered significant.

Data accessibility

The accession number for the UNC-3 ChIP-seq data is GEO: GSE143165

ACKNOWLEDGEMENTS

We thank the *Caenorhabditis* Genetics Center (CGC), which is funded by NIH Office of

Research Infrastructure Programs (P40 OD010440), for providing strains. We thank the

lab of Oliver Hobert for providing the OH14930 strain and Kaiyuan Tang for generating

reporter gene constructs. We are grateful to Edwin Ferguson, Oliver Hobert, Daniele

Canzio, Catarina Catela, and Weidong Feng for comments on this manuscript. This

work was supported by a training grant [University of Chicago Initiative for Maximizing

Student Development (IMSD), 2R25GM109439-06] to A.O, a training grant (T32

GM007183) to E. C., a Whitehall Foundation grant to P.K., grants from National Institute

of Neurological Disorders and Stroke (NINDS) of the NIH (Award Numbers

R00NS084988 and R21NS108505) to P.K., and a Medical Research Council grant MC-

A658-5TY30 to A.E.X.B.

AUTHOR CONTRIBUTIONS

Y. L., Data curation, Investigation, Visualization, Methodology, Writing—original draft,

review and editing; A.O., E. C., Formal analysis, Validation, Investigation; M.O., P. D.,

O. G., J. A., P.I., Investigation; A.E.X.B., Project administration, Writing—editing;

P. K., Conceptualization, Supervision, Investigation, Funding acquisition, Project

administration, Writing— original draft, review and editing.

DECLARATION OF INTERESTS

The authors declare no competing interests.

LEGENDS OF FIGURES AND TABLES

Figure 1. Mapping UNC-3 binding genome-wide with ChIP-Seq.

A: Diagram illustrating the endogenous reporter allele of UNC-3. GFP is inserted

immediately upstream of *unc-3*'s stop codon. Below, a representative image at L2 stage

showing expression of UNC-3::GFP fusion protein in cholinergic MN nuclei. Region

highlighted in dashed box is enlarged. Scale bar, 20 µm.

B: Quantification of terminal identity gene markers that report expression of known

UNC-3 targets (*cho-1/ChT, unc-17/VAChT, ace-2/AChE*) in WT and *ot839 [unc-3::gfp]*

animals at the L4 stage (N = 15). N. S.: not significant.

C: Heatmap of UNC-3 ChIP-Seq signal around 1.0 kb of the center of the binding peak.

D: Summary plot of UNC-3 ChIP-Seq signal with a 95% confidence interval (grey area)

around 3.0 kb of the TSS. The average signal peak is detected at ~200 bp upstream of

the TSS.

E: *de novo* motif discovery analysis of 6,892 UNC-3 binding peaks identifies a 12 bp-

long UNC-3 binding motif.

F: Pie chart summarizes genomic distribution of UNC-3 ChIP-Seq signal.

Figure 2. Global analyses of UNC-3 ChIP-Seq data.

A: Snapshots of UNC-3 ChIP-Seq and input (negative control) signals at the *cis*-

regulatory regions of known UNC-3 targets (*cho-1/ChT, unc-17/VAChT, acr-2/AChR,*

glr-4/GluR).

B: Graph summarizing protein class ontology analysis of putative target genes of UNC-

3 identified by ChIP-Seq. Out of the 3,502 protein-coding UNC-3 targets, 1,425 encode

for proteins with known protein class terms and these were the ones considered by

PANTHER. This analysis classifies UNC-3 targets into 3 broad categories: terminal

identity genes, gene expression regulators, and enzymes.

C: Pie chart breaking down TF families that show UNC-3 binding.

Figure 3. Terminal identity genes and transcription factors display distinct

temporal requirements for UNC-3.

A: Top: snapshots of UNC-3 ChIP-Seq and input (negative control) signals at the *cis*-

regulatory regions of 4 cholinergic terminal identity genes (*acr-2/AChR, unc-129/TFGb,*

glr-4/GluR, and *unc-17/VAChT*). The length of DNA elements included in each reporter

is shown. Bottom: quantification of terminal identity gene reporters in WT and *unc-3*

(n3435) animals at 3 different developmental stages – L2, L4, and day 1 adults (N ⩾

12). UNC-3 is required for both initiation and maintenance of all 4 terminal identity

genes. *** $p < 0.001$.

B: Top: snapshots of UNC-3 ChIP-Seq and input (negative control) signals at the *cis-*

regulatory regions of 5 transcription factors (*cfi-1/Arid3a, bnc-1/Bnc1, nhr-40, mab-*

9/Tbx20, and ceh-44/Cux1). The length of DNA elements included in each reporter is

shown. Middle: representative images of WT L4 animals showing expression of the

transgenic reporters in MNs. Scale bar, 20 µm. Bottom: quantification of transcription

factor reporters in WT and *unc-3 (n3435)* animals at 3 different developmental stages –

L2, L4, and young adults (day 1 or day 3) (N ⩾ 12). UNC-3 is required for maintenance,

but not initiation of the 5 TFs. N.S.: not significant, * $p < 0.05$, *** $p < 0.001$.

C: Schematic summarizing the phenomenon of temporal modularity in UNC-3 function.

The first module consists of terminal identity genes and TFs (**Figure S2**), which require

UNC-3 for both initiation and maintenance of gene expression. The second module

consists exclusively of TFs that require UNC-3 only for maintenance.

Figure 4. UNC-3 acts through a distal enhancer to maintain *cfi-1* expression in

cholinergic motor neurons.

A: Top: Snapshots of UNC-3 ChIP-Seq and input (negative control) signals at the *cis-*

regulatory region of *cfi-1*. The grey bar highlights an UNC-3 binding peak located ~12 kb

upstream of the TSS of *cfi-1* (-11,391 bp to -12,146 bp). Bottom: schematic showing the

strategy of constructing *cfi-1* reporters. Twelve transcriptional fusion reporters ([-1 bp to

-8,170 bp], #2 [993 bp to 3,764 bp], #3 [547 bp to -1,173 bp], #4 [-1,164 bp to -2,875

bp], #5 [-2,865 bp to -6,141 bp], #6 [-8,162 bp to -11,346 bp], #7 [-11,329 bp to -13,824

bp], #8 [-11,329 bp to -11,881 bp], #9 [-11,851 bp to -12,234 bp], #10 [-12,223 bp to -

12,722 bp], #11 [-12,705 bp to -13,284 bp], and #12 [-13,263 bp to -13,824 bp]) carry

cis-regulatory regions fused to fluorescent reporters (GFP or RFP). The endogenous

reporter alleles (*mNG::AID::cfi-1* and *mNG::AID::cfi-1* $^{\Delta enhancer\ (769\ bp)}$) have an in-frame

fluorescent protein mNeonGreen (mNG) insertion immediately after the ATG of *cfi-1*.

The enhancer KO allele *mNG::AID::cfi-1* $^{\Delta enhancer\ (769\ bp)}$ carries a 769 bp deletion (-

11,329 bp to -12,097 bp). Table on the right summarizes the expression pattern of each

reporter allele at L4 stage. N ⩾ 12.   +: reporter expressed, –: reporter not expressed,

+/–: reporter partially expressed in the respective neurons. Number of independent transgenic lines tested for each reporter is shown on the right.

B: Representative images showing the expression of reporter #7, reporter #8, *mNG::AID::cfi-1*, and *mNG::AID::cfi-1* $^{\Delta enhancer\ (769\ bp)}$ at specific life stages. Areas highlighted in dashed boxes are enlarged and presented on the right side of each picture. The onset of *cfi-1* expression occurs at the 3-fold embryonic stage. mNG+ MNs are annotated with arrowheads. Scale bars, 5 µm (3-fold embryos); 20 µm (larvae and adults).

C: Schematic summarizing the expression pattern of *cfi-1* and *unc-3* in nerve cord MNs.

D: Quantification of the number of cholinergic MNs expressing endogenous *cfi-1* (*mNG::AID::cfi-1)* in WT and animals carrying the enhancer deletion (*mNG::AID::cfi-1* $^{\Delta enhancer\ (769\ bp)}$). Deletion of the enhancer element located ~12 kb upstream of the TSS of *cfi-1* completely abolishes *cfi-1* expression in MNs at all tested stages. A red fluorescent marker (*ttr-39::mCherry*) for GABAergic MNs was used to exclude these neurons from the quantification. Cholinergic MNs expressing *cfi-1* were positive for mNG and negative for mCherry.

E: Bioinformatic analysis predicted 8 UNC-3 binding sites (COE motifs, shown as pink boxes) in the *cfi-1* enhancer region, which displays UNC-3 binding (-11,391 bp to -12,146 bp). Using CRISPR/Cas9, these 8 motifs were mutated by substituting duplets or triplets of nucleotides as shown below, thereby generating the strain *cfi-1 (syb1856 [mNG::AID::cfi-1* $^{8\ COE\ motifs\ mut}$]).

F: Quantification of the number of cholinergic MNs expressing the endogenous *cfi-1* reporter (*mNG::AID::cfi-1)* in WT and *unc-3 (n3435)* animals, as well as in animals with mutated COE motifs (*mNG::AID::cfi-1* $^{8\ COE\ motifs\ mut}$) at L2, L4, and day 1 adult stages (N ⩾ 12). N.S.: not significant, *** p < 0.001.

Figure 5. UNC-3 and Hox control *cfi-1* expression in cholinergic MNs.

A: A snapshot of UNC-3 (L2 stage), LIN-39 (L3 stage), and MAB-5 (L2 stage) ChIP-Seq signals at the *cfi-1* locus. UNC-3, LIN-39, and MAB-5 bind to the same *cfi-1* enhancer (highlighted in gray). Below: Schematics illustrating the reporters used in the rest of the figure.

B: Representative images showing the expression of the *mNG::AID::cfi-1* in WT, *unc-3 (n3435)*, and *lin-39 (n1760); mab-5 (e1239)* animals during 3-fold embryonic, L2, L4,

and day 1 adult stages. *cfi-1* is expressed in 4 cholinergic MN subtypes (DA, DB, VA,

and VB). DA and DB are born embryonically, while VA and VB are born post-

embryonically. Areas highlighted in dashed boxes are enlarged and presented on the

right side of each picture. mNG+ MNs are annotated with arrowheads. Scale bars, 5 μm

(3-fold embryos); 20 μm (larvae and adults).

C: Quantification of the number of cholinergic MNs expressing the endogenous *cfi-1*

reporter (*mNG::AID::cfi-1)* in WT animals, *unc-3 (n3435)* mutants, *lin-39 (n1760); mab-5*

(e1239) double mutants, and *unc-3 (n3435); lin-39 (n1760); mab-5 (e1239)* triple

mutants during 3-fold embryonic, L2, L4, and day 1 adult stages (N $\geqslant$ 12). N.S.: not

significant, *** $p < 0.001$. A red fluorescent marker (*ttr-39::mCherry*) for GABAergic MNs

was used to exclude these neurons from the quantification. Cholinergic MNs expressing

cfi-1 were positive for mNG and negative for mCherry.

D: Quantification of the expression of transgenic *cfi-1* reporter #7 in WT and *lin-39*

(n1760); mab-5 (e1239) animals at L2 stage (N = 13). Reporter expression is strongly

affected in *lin-39 (-); mab-5 (-)* double mutants. *** $p < 0.001$.

E: Quantification of the WT transgenic reporter #8 and the same reporter with the LIN-

39 binding site mutated (point mutations) at larval (L2) and adult (D3) stages (N $\geqslant$ 13). *

$p < 0.05$.

F: Quantification of the expression of the *mNG::AID::cfi-1* allele in WT animals, *lin-39*

(n1760) single mutants, and *lin-39 (n1760); mab-5 (e1239)* double mutants at the L2

stage (N $\geqslant$ 12). While the number of cholinergic MNs with *cfi-1* expression is mildly

decreased in *lin-39* single mutants, more severe effects are observed in double

mutants. *** $p < 0.001$.

G: Schematic summarizing the mechanisms underlying initiation and maintenance of

cfi-1 expression in cholinergic MNs.

Figure 6. Disruption of temporal modularity in UNC-3 function leads to

locomotion defects

A: Schematics illustrating four *cfi-1* alleles tested for behavioral analysis.

B-G: Examples of six locomotion features significantly disrupted in animals carrying a

putative null *(ot786)* allele for *cfi-1*. Locomotion analysis was performed on day 2 adult

worms. Animals lacking *cfi-1* expression (initiation and maintenance) specifically in MNs

(*mNG::AID::cfi-1* $^{\Delta enhancer\ (769\ bp)}$) and animals unable to maintain *cfi-1* expression in

cholinergic MNs (*mNG::AID::cfi-1*$^{8\ COE\ motifs\ mut}$) display locomotion defects when

compared to control *mNG::AID::cfi-1* animals. As expected, these defects are milder

when compared to animals carrying the *cfi-1* (*ot786*) allele. Panels B-F show locomotion

features related to body curvature, whereas panel G shows radial velocity of the neck. A

detailed description of each locomotion feature is provided below:

(**B**) curvature_std_midbody_w_backward_abs_10th: 10th percentile of the absolute

value of the standard deviation of the curvature of the midbody, while worm is moving

backwards.

(**C**) curvature_tail_w_backward_abs_90th: 90th percentile of the absolute value of the

curvature of the tail, while worm is moving backwards.

(**D**) curvature_std_midbody_w_backward_50th: 50th percentile of the standard deviation

of the curvature of the midbody, while worm is moving backwards.

(**E**) curvature_std_hips_abs_10th: 10th percentile of the absolute value of the standard

deviation of the curvature of the hips.

(**F**) curvature_std_hips_abs_50th: 50th percentile of the absolute value of the standard

deviation of the curvature of the hips

(**G**) d_rel_to_body_radial_vel_neck_w_forward_90th: 90th percentile of the derivative of

radial velocity of the neck relative to the centroid of the midbody points, while worm is

moving forwards.

1006

1007

Figure 7. Temporal modularity in UNC-30/Pitx function in GABAergic MNs.

A: Schematic summarizing the expression of *cfi-1/Arid3a, unc-3/Ebf, unc-30/Pitx, unc-*

25/GAD, and *unc-47/VGAT* in MN subtypes of the *C. elegans* ventral nerve cord.

B: Schematic showing time of birth and cell body position of GABAergic nerve cord

MNs. DD neurons are born embryonically. VD neurons are born post-embryonically.

C: Bioinformatic analyses predict 4 UNC-30 binding sites (yellow boxes) in the *cfi-1*

enhancer. The location of UNC-3 and LIN-39 binding sites are also shown.

D: Top: snapshots of UNC-30 ChIP-Seq and input (negative control) signals at the *cis-*

regulatory regions of 2 GABAergic terminal identity genes (*unc-25/GAD, unc-47/VGAT*).

Bottom: quantification of the expression of transgenic reporters in WT and *unc-30*

(e191) animals at 4 different developmental stages – 3-fold embryo, L2, L4, and day 1

adults (N = 15). UNC-30 is required for both initiation and maintenance of *unc-25/GAD*
and *unc-47/VGAT*. *** p < 0.001.

E: Top: a snapshot of UNC-3 ChIP-Seq and UNC-30 ChIP-Seq signals at the *cfi-1*
locus. Bottom: quantification of the number of MNs expressing the endogenous reporter
mNG::AID::cfi-1 in WT and *unc-30 (e191)* animals. All *cfi-1*-expressing MNs in the
ventral cord (cholinergic and GABAergic MNs) were counted in 3-fold embryos due to a
lack of a specific marker that labels GABAergic MNs in embryos. Expression of *cfi-1*
specifically in GABA neurons was quantified at L2, L4, and day 1 adult stages (N ⩾ 12).
At those stages, cholinergic MNs were identified based on a fluorescent marker (*cho-*
1::mChOpti), which are ruled out during scoring. GABAergic MNs were scored positive
for *cfi-1* expression when the *mNG::AID::cfi-1* (green) expression co-localized with *ttr-*
39::mCherry (red). N.S.: not significant, *** p < 0.001.

1031
1032

Table 1: A summary of the *cis*-regulatory analysis to identify novel transcription
factors controlled by UNC-3. Reporter alleles of each TF were built and examined for
expression pattern. TFs that are expressed in MNs were further tested for UNC-3
dependency. Ten of 16 tested TFs (62.5%) show expression in MNs, of which 9 require
UNC-3 activity for normal expression. Two reporters were generated for *nhr-1* and *nhr-*
40 because two distinct UNC-3 ChIP-Seq peaks were found in the *cis*-regulatory region
of these genes. N/A: Not applicable.

1040

 LEGENDS OF SUPPLEMENTARY FIGURES

Figure 1 – figure supplement 1. UNC-3 ChIP-Seq results yield genome-wide

enrichment of UNC-3.

A: Fingerprint plot indicating localized, strong enrichment of UNC-3 in the genome.

Specifically, when counting the reads contained in 93% of all genomic bins (data point

0.93, 0.5 on UNC-3 ChIP curve), only 50% of the maximum number of reads are

reached, which indicates 7% of the genome contains half of total sequencing reads from

the ChIP sample.

B: Snapshots of UNC-3 ChIP-Seq and input (negative control) signals at the *cis*-

regulatory regions of known UNC-3 targets (*del-1, acc-4, twk-13, dbl-1, unc-77,* and

twk-40).

1052

Figure 3 – figure supplement 1. UNC-3 directly controls the expression of several

TF reporters in MNs.

A-C: Top: snapshots of UNC-3 ChIP-Seq and input (negative control) signal at the *cis*-

regulatory regions of 5 TF-encoding gens (*nhr-1, ztf-26, zfh-2, nhr-49, nhr-19*).

Transgenic RFP reporters contain the *cis*-regulatory regions bound by UNC-3, as well

as flanking sequences. Middle: representative images of WT L4 animals showing

expression of the transgenic reporters in nerve cord MNs. Scale bar, 20 µm. Bottom:

quantification of TF reporters in WT and *unc-3 (n3435)* animals at 3 different

developmental stages – L2, L4, and day 3 adult. *nhr-1, ztf-26,* and *zfh-2* are activated

by UNC-3 **(A)**, while *nhr-49* is repressed by UNC-3 **(B)** and *nhr-19* does not appear to

be controlled by UNC-3 **(C)**. N ⩾ 15. N.S.: not significant, * $p < 0.05$, ** $p < 0.01$, *** $p <$

0.001.

D: Three TF reporters (*ccch-3, ztf-17, nhr-1*) are not expressed in nerve cord MNs, but

show expression in head neurons. Top: snapshots of UNC-3 ChIP-seq and input

(negative control) signals at the *cis*-regulatory regions of *ccch-3, ztf-17,* and *nhr-1*.

Transgenic RFP reporters contain the *cis*-regulatory regions bound by UNC-3. Bottom:

representative images of WT L4 animals showing expression of the transgenic reporters

in some unidentified neurons of the head. It is known that UNC-3 is expressed in some

head neurons.

1072

1073

Figure 4 – figure supplement 1: CFI-1 does not auto-regulate.

A: Snapshots of UNC-3 ChIP-seq signal at the *cfi-1* locus in WT and *cfi-1 (ot786)*

animals. UNC-3 binds to the *cfi-1* enhancer normally in *cfi-1 (ot786)* mutants.

B: Quantification of the number of MNs expressing the transcriptional fusion reporter #7

(*cfi-1$^{2.5kb}$::RFP*) in WT worms and *cfi-1 (ot786)* mutants (N = 14). No significant

difference was detected between WT and *cfi-1 (ot786)*, suggesting that the expression

of reporter #7 (*cfi-1$^{2.5kb}$::RFP*) is not altered upon genetic removal of *cfi-1*. N.S.: not

significant.

1082

1083

Figure 5 – figure supplement 1. Auxin-inducible depletion of LIN-39 at larval stage

4 (L4) does not affect *cfi-1* expression in nerve cord MNs.

A: Representative images showing expression of LIN-39::mNG::AID after treatment with

ethanol and auxin (negative control). LIN-39::mNG::AID is degraded and mNG

fluorescent signal becomes undetectable in MNs, when worms are imaged after 4 days

of auxin treatment. Arrowheads indicate MN nuclei in the nerve cord.

B: Knock-down of *lin-39::mNG::AID* starting at the L4 stage (onset of auxin treatment at

L4) did not affect *cfi-1$^{2.5kb}$::RFP* expression in adult animals 4 days later (N ⩾ 13). N.S.:

not significant. The number of MNs expressing *cfi-1$^{2.5kb}$::RFP* was quantified.

1093

Figure 5 – figure supplement 2. CFI-1 is required post-embryonically to maintain

DA and DB neuronal identities.

A: CFI-1 regulates DA and DB MN identity by repressing the glutamate receptor subunit

glr-4/GluR and possibly other genes. In *cfi-1 (-)* mutants, *glr-4/GluR* becomes

ectopically activated by UNC-3 in DA and DB MNs, and these neurons adopt a mixed

molecular identity.

B: Representative images showing expression of mNG::AID::CFI-1 after treatment with

ethanol and auxin. Upon two days of continuous auxin treatment (onset of treatment at

L1), mNG::AID::CFI-1 is degraded and mNG expression in MNs (arrowheads) becomes

undetectable

C: Quantification of MNs expressing *glr-4::RFP* was performed on L4 worms (2 days

after the onset of auxin treatment). Some mild hypomorphic effects in the expression of

glr-4::RFP was observed in the EtOH group (negative control), potentially due to mild

reduction in CFI-1 levels triggered by TIR1 even in the absence of auxin. A significant

increase in the number of MNs expressing *glr-4::RFP* was evident in auxin-treated

worms in comparison to control animals. WT and *cfi-1 (-)* data are also provided for

comparison. N $\geqslant$ 20. N.S.: not significant, ** p < 0.01.

1111

Figure 5 – figure supplement 3: Hox proteins and UNC-3 control *bnc-1* expression

in VA and VB neurons.

A: Schematic summarizing the expression of *bnc-1/Bnc1* and *unc-3/Ebf* in MN subtypes

of the *C. elegans* ventral nerve cord.

B: A snapshot of UNC-3 ChIP-seq and input (negative control) signals at the *cis-*

regulatory region of *bnc-1*.

C: Quantification of expression of the endogenous reporter *bnc-1::mNG::AID* in WT,

unc-3 (n3435), and *lin-39 (n1760); mab-5 (e1239)* animals during L2, L4, and day 1

adult stages (N $\geqslant$ 12). *** p < 0.001.

D: Representative images of the endogenous reporter *bnc-1::mNG::AID* in WT, *unc-3*

(e151), and *lin-39 (n1760); mab-5 (e1239)* animals during L2, L4, and day 1 adult

stages. *bnc-1* is expressed in two cholinergic MN subtypes (VA and VB). Areas

highlighted in dashed boxes are enlarged and presented on the right side of each

picture. Arrowheads point to nuclei of VA and VB neurons that express the reporter.

Above the white dashed line lies the intestine, which is autofluorescent in the green

channel. Scale bar, 20 µm.

1128

1129

LEGENDS OF SUPPLEMENTARY FILES

Supplementary File 1: Table summarizing UNC-3 ChIP-Seq signal distribution at the *cis*-regulatory regions of previously identified UNC-3 targets.

Supplementary File 2: Table summarizing the results of protein class ontology analysis over novel target genes of UNC-3. In total, 1,425 genes are classified into 25 protein classes.

Supplementary File 3: UNC-3 binds to the *cis*-regulatory region of numerous genes expressed in cholinergic MNs.

Supplementary File 4: A list of all *C. elegans* strains described in this study.

REFERENCES

BAEK, M., ENRIQUEZ, J. & MANN, R. S. 2013. Dual role for Hox genes and Hox co-factors in conferring leg motoneuron survival and identity in Drosophila. *Development,* 140**,** 2027-38.

BLACKBURN, P. R., BARNETT, S. S., ZIMMERMANN, M. T., COUSIN, M. A., KAIWAR, C., PINTO, E. V. F., NIU, Z., FERBER, M. J., URRUTIA, R. A., SELCEN, D., KLEE, E. W. & PICHURIN, P. N. 2017. Novel de novo variant in EBF3 is likely to impact DNA binding in a patient with a neurodevelopmental disorder and expanded phenotypes: patient report, in silico functional assessment, and review of published cases. *Cold Spring Harb Mol Case Stud,* 3**,** a001743.

BOYLE, A. P., ARAYA, C. L., BRDLIK, C., CAYTING, P., CHENG, C., CHENG, Y., GARDNER, K., HILLIER, L. W., JANETTE, J., JIANG, L., KASPER, D., KAWLI, T., KHERADPOUR, P., KUNDAJE, A., LI, J. J., MA, L., NIU, W., REHM, E. J., ROZOWSKY, J., SLATTERY, M., SPOKONY, R., TERRELL, R., VAFEADOS, D., WANG, D., WEISDEPP, P., WU, Y. C., XIE, D., YAN, K. K., FEINGOLD, E. A., GOOD, P. J., PAZIN, M. J., HUANG, H., BICKEL, P. J., BRENNER, S. E., REINKE, V., WATERSTON, R. H., GERSTEIN, M., WHITE, K. P., KELLIS, M. & SNYDER, M. 2014. Comparative analysis of regulatory information and circuits across distant species. *Nature,* 512**,** 453-6.

BRENNER, S. 1974. The genetics of Caenorhabditis elegans. *Genetics,* 77**,** 71-94.

BROZOVA, E., SIMECKOVA, K., KOSTROUCH, Z., RALL, J. E. & KOSTROUCHOVA, M. 2006. NHR-40, a Caenorhabditis elegans supplementary nuclear receptor, regulates embryonic and early larval development. *Mech Dev,* 123**,** 689-701.

CATELA, C. & KRATSIOS, P. 2019. Transcriptional mechanisms of motor neuron development in vertebrates and invertebrates. *Dev Biol*.

CATELA, C., SHIN, M. M., LEE, D. H., LIU, J. P. & DASEN, J. S. 2016. Hox Proteins Coordinate Motor Neuron Differentiation and Connectivity Programs through Ret/Gfralpha Genes. *Cell Rep,* 14**,** 1901-15.

CHAO, H. T., DAVIDS, M., BURKE, E., PAPPAS, J. G., ROSENFELD, J. A., MCCARTY, A. J., DAVIS, T., WOLFE, L., TORO, C., TIFFT, C., XIA, F., STONG, N., JOHNSON, T. K., WARR, C. G., UNDIAGNOSED DISEASES, N., YAMAMOTO, S., ADAMS, D. R., MARKELLO, T. C., GAHL, W. A., BELLEN, H. J., WANGLER, M. F. & MALICDAN, M. C. 2017. A Syndromic Neurodevelopmental Disorder Caused by De Novo Variants in EBF3. *Am J Hum Genet,* 100**,** 128-137.

CLARK, S. G., CHISHOLM, A. D. & HORVITZ, H. R. 1993. Control of cell fates in the central body region of C. elegans by the homeobox gene lin-39. *Cell,* 74**,** 43-55.

CORBO, J. C., LAWRENCE, K. A., KARLSTETTER, M., MYERS, C. A., ABDELAZIZ, M., DIRKES, W., WEIGELT, K., SEIFERT, M., BENES, V., FRITSCHE, L. G., WEBER, B. H. & LANGMANN, T. 2010. CRX ChIP-seq reveals the cis-regulatory architecture of mouse photoreceptors. *Genome Res,* 20**,** 1512-25.

COWING, D. W. & KENYON, C. 1992. Expression of the homeotic gene mab-5 during Caenorhabditis elegans embryogenesis. *Development,* 116**,** 481-90.

DENERIS, E. S. & HOBERT, O. 2014. Maintenance of postmitotic neuronal cell identity. *Nat Neurosci,* 17**,** 899-907.

DOE, C. Q. 2017. Temporal Patterning in the Drosophila CNS. *Annu Rev Cell Dev Biol,* 33**,** 219-240.

DUBOIS, L. & VINCENT, A. 2001. The COE--Collier/Olf1/EBF--transcription factors:
structural conservation and diversity of developmental functions. *Mech Dev,* 108**,** 3-
12.
EASTMAN, C., HORVITZ, H. R. & JIN, Y. 1999. Coordinated transcriptional regulation of the
unc-25 glutamic acid decarboxylase and the unc-47 GABA vesicular transporter by
the Caenorhabditis elegans UNC-30 homeodomain protein. *J Neurosci,* 19**,** 6225-34.
ELLMEIER, W., SUNSHINE, M. J., MASCHEK, R. & LITTMAN, D. R. 2002. Combined deletion of
CD8 locus cis-regulatory elements affects initiation but not maintenance of CD8
expression. *Immunity,* 16**,** 623-34.
ESTACIO-GOMEZ, A. & DIAZ-BENJUMEA, F. J. 2014. Roles of Hox genes in the patterning of
the central nervous system of Drosophila. *Fly (Austin),* 8**,** 26-32.
FEDERICI, T. & BOULIS, N. M. 2006. Gene-based treatment of motor neuron diseases.
Muscle Nerve, 33**,** 302-23.
FENG, W., LI, Y., DAO, P., ABURAS, J., ISLAM, P., ELBAZ, B., KOLARZYK, A., BROWN, A. E. &
KRATSIOS, P. 2020. A terminal selector prevents a Hox transcriptional switch to
safeguard motor neuron identity throughout life. *Elife,* 9.
FLAMES, N. & HOBERT, O. 2009. Gene regulatory logic of dopamine neuron differentiation.
Nature, 458**,** 885-9.
GARCIA-BELLIDO, A. 1975. Genetic control of wing disc development in Drosophila. *Ciba
Found Symp,* 0**,** 161-82.
GREIG, L. C., WOODWORTH, M. B., GALAZO, M. J., PADMANABHAN, H. & MACKLIS, J. D.
2013. Molecular logic of neocortical projection neuron specification, development
and diversity. *Nat Rev Neurosci,* 14**,** 755-69.
HARMS, F. L., GIRISHA, K. M., HARDIGAN, A. A., KORTUM, F., SHUKLA, A., ALAWI, M., DALAL,
A., BRADY, L., TARNOPOLSKY, M., BIRD, L. M., CEULEMANS, S., BEBIN, M., BOWLING,
K. M., HIATT, S. M., LOSE, E. J., PRIMIANO, M., CHUNG, W. K., JUUSOLA, J., AKDEMIR,
Z. C., BAINBRIDGE, M., CHARNG, W. L., DRUMMOND-BORG, M., ELDOMERY, M. K.,
EL-HATTAB, A. W., SALEH, M. A., BEZIEAU, S., COGNE, B., ISIDOR, B., KURY, S.,
LUPSKI, J. R., MYERS, R. M., COOPER, G. M. & KUTSCHE, K. 2017. Mutations in EBF3
Disturb Transcriptional Profiles and Cause Intellectual Disability, Ataxia, and Facial
Dysmorphism. *Am J Hum Genet,* 100**,** 117-127.
HEINZ, S., BENNER, C., SPANN, N., BERTOLINO, E., LIN, Y. C., LASLO, P., CHENG, J. X., MURRE,
C., SINGH, H. & GLASS, C. K. 2010. Simple combinations of lineage-determining
transcription factors prime cis-regulatory elements required for macrophage and B
cell identities. *Mol Cell,* 38**,** 576-89.
HOBERT, O. 2002. PCR fusion-based approach to create reporter gene constructs for
expression analysis in transgenic C. elegans. *Biotechniques,* 32**,** 728-30.
HOBERT, O. 2008. Regulatory logic of neuronal diversity: terminal selector genes and
selector motifs. *Proc Natl Acad Sci U S A,* 105**,** 20067-71.
HOBERT, O. 2011. Regulation of terminal differentiation programs in the nervous system.
Annu Rev Cell Dev Biol, 27**,** 681-96.
HOBERT, O. 2016a. A map of terminal regulators of neuronal identity in Caenorhabditis
elegans. *Wiley Interdiscip Rev Dev Biol,* 5**,** 474-98.
HOBERT, O. 2016b. Terminal Selectors of Neuronal Identity. *Curr Top Dev Biol,* 116**,** 455-75.
HOBERT, O. & KRATSIOS, P. 2019. Neuronal identity control by terminal selectors in
worms, flies, and chordates. *Curr Opin Neurobiol,* 56**,** 97-105.
HOUSSET, M., SAMUEL, A., ETTAICHE, M., BEMELMANS, A., BEBY, F., BILLON, N. &
LAMONERIE, T. 2013. Loss of Otx2 in the adult retina disrupts retinal pigment
epithelium function, causing photoreceptor degeneration. *J Neurosci,* 33**,** 9890-904.

HUTLET, B., THEYS, N., COSTE, C., AHN, M. T., DOSHISHTI-AGOLLI, K., LIZEN, B. & GOFFLOT,
F. 2016. Systematic expression analysis of Hox genes at adulthood reveals novel
patterns in the central nervous system. *Brain Struct Funct,* 221**,** 1223-43.
JAVER, A., CURRIE, M., LEE, C. W., HOKANSON, J., LI, K., MARTINEAU, C. N., YEMINI, E.,
GRUNDY, L. J., LI, C., CH'NG, Q., SCHAFER, W. R., NOLLEN, E. A. A., KERR, R. &
BROWN, A. E. X. 2018a. An open-source platform for analyzing and sharing worm-
behavior data. *Nat Methods,* 15**,** 645-646.
JAVER, A., RIPOLL-SANCHEZ, L. & BROWN, A. E. X. 2018b. Powerful and interpretable
behavioural features for quantitative phenotyping of Caenorhabditis elegans. *Philos*
Trans R Soc Lond B Biol Sci, 373.
JESSELL, T. M. 2000. Neuronal specification in the spinal cord: inductive signals and
transcriptional codes. *Nat Rev Genet,* 1**,** 20-9.
JIN, Y., HOSKINS, R. & HORVITZ, H. R. 1994. Control of type-D GABAergic neuron
differentiation by C. elegans UNC-30 homeodomain protein. *Nature,* 372**,** 780-3.
JOHNSON, S. A., HARMON, K. J., SMILEY, S. G., STILL, F. M. & KAVALER, J. 2011. Discrete
regulatory regions control early and late expression of D-Pax2 during external
sensory organ development. *Dev Dyn,* 240**,** 1769-78.
KADKHODAEI, B., ALVARSSON, A., SCHINTU, N., RAMSKOLD, D., VOLAKAKIS, N.,
JOODMARDI, E., YOSHITAKE, T., KEHR, J., DECRESSAC, M., BJORKLUND, A.,
SANDBERG, R., SVENNINGSSON, P. & PERLMANN, T. 2013. Transcription factor
Nurr1 maintains fiber integrity and nuclear-encoded mitochondrial gene expression
in dopamine neurons. *Proc Natl Acad Sci U S A,* 110**,** 2360-5.
KADKHODAEI, B., ITO, T., JOODMARDI, E., MATTSSON, B., ROUILLARD, C., CARTA, M.,
MURAMATSU, S., SUMI-ICHINOSE, C., NOMURA, T., METZGER, D., CHAMBON, P.,
LINDQVIST, E., LARSSON, N. G., OLSON, L., BJORKLUND, A., ICHINOSE, H. &
PERLMANN, T. 2009. Nurr1 is required for maintenance of maturing and adult
midbrain dopamine neurons. *J Neurosci,* 29**,** 15923-32.
KARLSSON, D., BAUMGARDT, M. & THOR, S. 2010. Segment-specific neuronal subtype
specification by the integration of anteroposterior and temporal cues. *PLoS Biol,* 8**,**
e1000368.
KERK, S. Y., KRATSIOS, P., HART, M., MOURAO, R. & HOBERT, O. 2017. Diversification of C.
elegans Motor Neuron Identity via Selective Effector Gene Repression. *Neuron,* 93**,**
80-98.
KIM, K., COLOSIMO, M. E., YEUNG, H. & SENGUPTA, P. 2005. The UNC-3 Olf/EBF protein
represses alternate neuronal programs to specify chemosensory neuron identity.
Dev Biol, 286**,** 136-48.
KONSTANTINIDES, N., KAPURALIN, K., FADIL, C., BARBOZA, L., SATIJA, R. & DESPLAN, C.
2018. Phenotypic Convergence: Distinct Transcription Factors Regulate Common
Terminal Features. *Cell,* 174**,** 622-635 e13.
KRATSIOS, P., KERK, S. Y., CATELA, C., LIANG, J., VIDAL, B., BAYER, E. A., FENG, W., DE LA
CRUZ, E. D., CROCI, L., CONSALEZ, G. G., MIZUMOTO, K. & HOBERT, O. 2017. An
intersectional gene regulatory strategy defines subclass diversity of C. elegans
motor neurons. *Elife,* 6.
KRATSIOS, P., PINAN-LUCARRE, B., KERK, S. Y., WEINREB, A., BESSEREAU, J. L. & HOBERT,
O. 2015. Transcriptional coordination of synaptogenesis and neurotransmitter
signaling. *Curr Biol,* 25**,** 1282-95.
KRATSIOS, P., STOLFI, A., LEVINE, M. & HOBERT, O. 2012. Coordinated regulation of
cholinergic motor neuron traits through a conserved terminal selector gene. *Nat*
Neurosci, 15**,** 205-14.

LANGMEAD, B. & SALZBERG, S. L. 2012. Fast gapped-read alignment with Bowtie 2. *Nat*
Methods, 9**,** 357-9.
LEYVA-DIAZ, E. & HOBERT, O. 2019. Transcription factor autoregulation is required for
acquisition and maintenance of neuronal identity. *Development,* 146.
LIU, J. & FIRE, A. 2000. Overlapping roles of two Hox genes and the exd ortholog ceh-20 in
diversification of the C. elegans postembryonic mesoderm. *Development,* 127**,** 5179-
90.
LODATO, S. & ARLOTTA, P. 2015. Generating neuronal diversity in the mammalian cerebral
cortex. *Annu Rev Cell Dev Biol,* 31**,** 699-720.
LOPES, R., VERHEY VAN WIJK, N., NEVES, G. & PACHNIS, V. 2012. Transcription factor LIM
homeobox 7 (Lhx7) maintains subtype identity of cholinergic interneurons in the
mammalian striatum. *Proc Natl Acad Sci U S A,* 109**,** 3119-24.
MALOOF, J. N. & KENYON, C. 1998. The Hox gene lin-39 is required during C. elegans vulval
induction to select the outcome of Ras signaling. *Development,* 125**,** 181-90.
MANZANARES, M., BEL-VIALAR, S., ARIZA-MCNAUGHTON, L., FERRETTI, E., MARSHALL, H.,
MACONOCHIE, M. M., BLASI, F. & KRUMLAUF, R. 2001. Independent regulation of
initiation and maintenance phases of Hoxa3 expression in the vertebrate hindbrain
involve auto- and cross-regulatory mechanisms. *Development,* 128**,** 3595-607.
MAYER, C., HAFEMEISTER, C., BANDLER, R. C., MACHOLD, R., BATISTA BRITO, R., JAGLIN,
X., ALLAWAY, K., BUTLER, A., FISHELL, G. & SATIJA, R. 2018. Developmental
diversification of cortical inhibitory interneurons. *Nature,* 555**,** 457-462.
MI, D., LI, Z., LIM, L., LI, M., MOISSIDIS, M., YANG, Y., GAO, T., HU, T. X., PRATT, T., PRICE, D. J.,
SESTAN, N. & MARIN, O. 2018. Early emergence of cortical interneuron diversity in
the mouse embryo. *Science,* 360**,** 81-85.
MI, H., MURUGANUJAN, A., CASAGRANDE, J. T. & THOMAS, P. D. 2013. Large-scale gene
function analysis with the PANTHER classification system. *Nat Protoc,* 8**,** 1551-66.
MIGUEL-ALIAGA, I. & THOR, S. 2004. Segment-specific prevention of pioneer neuron
apoptosis by cell-autonomous, postmitotic Hox gene activity. *Development,* 131**,**
6093-105.
MORIS-SANZ, M., ESTACIO-GOMEZ, A., SANCHEZ-HERRERO, E. & DIAZ-BENJUMEA, F. J.
2015. The study of the Bithorax-complex genes in patterning CCAP neurons reveals
a temporal control of neuronal differentiation by Abd-B. *Biol Open,* 4**,** 1132-42.
OKATY, B. W., FRERET, M. E., ROOD, B. D., BRUST, R. D., HENNESSY, M. L., DEBAIROS, D.,
KIM, J. C., COOK, M. N. & DYMECKI, S. M. 2015. Multi-Scale Molecular Deconstruction
of the Serotonin Neuron System. *Neuron,* 88**,** 774-91.
PEREIRA, L., KRATSIOS, P., SERRANO-SAIZ, E., SHEFTEL, H., MAYO, A. E., HALL, D. H.,
WHITE, J. G., LEBOEUF, B., GARCIA, L. R., ALON, U. & HOBERT, O. 2015. A cellular and
regulatory map of the cholinergic nervous system of C. elegans. *Elife,* 4.
PERRY, M., KONSTANTINIDES, N., PINTO-TEIXEIRA, F. & DESPLAN, C. 2017. Generation and
Evolution of Neural Cell Types and Circuits: Insights from the Drosophila Visual
System. *Annu Rev Genet,* 51**,** 501-527.
PFEFFER, P. L., PAYER, B., REIM, G., DI MAGLIANO, M. P. & BUSSLINGER, M. 2002. The
activation and maintenance of Pax2 expression at the mid-hindbrain boundary is
controlled by separate enhancers. *Development,* 129**,** 307-18.
PHILIPPIDOU, P. & DASEN, J. S. 2013. Hox genes: choreographers in neural development,
architects of circuit organization. *Neuron,* 80**,** 12-34.
PIKKARAINEN, S., TOKOLA, H., KERKELA, R. & RUSKOAHO, H. 2004. GATA transcription
factors in the developing and adult heart. *Cardiovasc Res,* 63**,** 196-207.

POCOCK, R., MIONE, M., HUSSAIN, S., MAXWELL, S., PONTECORVI, M., ASLAM, S., GERRELLI, D., SOWDEN, J. C. & WOOLLARD, A. 2008. Neuronal function of Tbx20 conserved from nematodes to vertebrates. *Dev Biol,* 317**,** 671-85.

PRASAD, B. C., YE, B., ZACKHARY, R., SCHRADER, K., SEYDOUX, G. & REED, R. R. 1998. unc-3, a gene required for axonal guidance in Caenorhabditis elegans, encodes a member of the O/E family of transcription factors. *Development,* 125**,** 1561-8.

RAMIREZ, F., RYAN, D. P., GRUNING, B., BHARDWAJ, V., KILPERT, F., RICHTER, A. S., HEYNE, S., DUNDAR, F. & MANKE, T. 2016. deepTools2: a next generation web server for deep-sequencing data analysis. *Nucleic Acids Res,* 44**,** W160-5.

RHEE, H. S., CLOSSER, M., GUO, Y., BASHKIROVA, E. V., TAN, G. C., GIFFORD, D. K. & WICHTERLE, H. 2016. Expression of Terminal Effector Genes in Mammalian Neurons Is Maintained by a Dynamic Relay of Transient Enhancers. *Neuron,* 92**,** 1252-1265.

SABARIS, G., LAIKER, I., PREGER-BEN NOON, E. & FRANKEL, N. 2019. Actors with Multiple Roles: Pleiotropic Enhancers and the Paradigm of Enhancer Modularity. *Trends Genet,* 35**,** 423-433.

SCHINDELIN, J., ARGANDA-CARRERAS, I., FRISE, E., KAYNIG, V., LONGAIR, M., PIETZSCH, T., PREIBISCH, S., RUEDEN, C., SAALFELD, S., SCHMID, B., TINEVEZ, J. Y., WHITE, D. J., HARTENSTEIN, V., ELICEIRI, K., TOMANCAK, P. & CARDONA, A. 2012. Fiji: an open-source platform for biological-image analysis. *Nat Methods,* 9**,** 676-82.

SCOTT, M. M., KRUEGER, K. C. & DENERIS, E. S. 2005. A differentially autoregulated Pet-1 enhancer region is a critical target of the transcriptional cascade that governs serotonin neuron development. *J Neurosci,* 25**,** 2628-36.

SETH R TAYLOR, G. S., MOLLY REILLY, LORI GLENWINKEL, ABIGAIL POFF, REBECCA MCWHIRTER, CHUAN XU, ALEXIS WEINREB, MANASA BASAVARAJU, STEVEN J COOK, ALEC BARRETT, ALEXANDER ABRAMS, BERTA VIDAL, CYRIL CROS, IBNUL RAFI, NENAD SESTAN, MARC HAMMARLUND, OLIVER HOBERT, DAVID M. MILLER III 2019. Expression profiling of the mature C. elegans nervous system by single-cell RNA-Sequencing. *bioRxiv*.

SHAHAM, S. & BARGMANN, C. I. 2002. Control of neuronal subtype identity by the C. elegans ARID protein CFI-1. *Genes Dev,* 16**,** 972-83.

SHEN, L., SHAO, N., LIU, X. & NESTLER, E. 2014. ngs.plot: Quick mining and visualization of next-generation sequencing data by integrating genomic databases. *BMC Genomics,* 15**,** 284.

SLEVEN, H., WELSH, S. J., YU, J., CHURCHILL, M. E., WRIGHT, C. F., HENDERSON, A., HORVATH, R., RANKIN, J., VOGT, J., MAGEE, A., MCCONNELL, V., GREEN, A., KING, M. D., COX, H., ARMSTRONG, L., LEHMAN, A., NELSON, T. N., DECIPHERING DEVELOPMENTAL DISORDERS, S., STUDY, C., WILLIAMS, J., CLOUSTON, P., HAGMAN, J. & NEMETH, A. H. 2017. De Novo Mutations in EBF3 Cause a Neurodevelopmental Syndrome. *Am J Hum Genet,* 100**,** 138-150.

SOLER, C., HAN, J. & TAYLOR, M. V. 2012. The conserved transcription factor Mef2 has multiple roles in adult Drosophila musculature formation. *Development,* 139**,** 1270-5.

TAKAHASHI, Y., HAMADA, J., MURAKAWA, K., TAKADA, M., TADA, M., NOGAMI, I., HAYASHI, N., NAKAMORI, S., MONDEN, M., MIYAMOTO, M., KATOH, H. & MORIUCHI, T. 2004. Expression profiles of 39 HOX genes in normal human adult organs and anaplastic thyroid cancer cell lines by quantitative real-time RT-PCR system. *Exp Cell Res,* 293**,** 144-53.

TREIBER, N., TREIBER, T., ZOCHER, G. & GROSSCHEDL, R. 2010a. Structure of an Ebf1:DNA
complex reveals unusual DNA recognition and structural homology with Rel
proteins. *Genes Dev,* 24**,** 2270-5.
TREIBER, T., MANDEL, E. M., POTT, S., GYORY, I., FIRNER, S., LIU, E. T. & GROSSCHEDL, R.
2010b. Early B cell factor 1 regulates B cell gene networks by activation, repression,
and transcription- independent poising of chromatin. *Immunity,* 32**,** 714-25.
WANG, M. M. & REED, R. R. 1993. Molecular cloning of the olfactory neuronal transcription
factor Olf-1 by genetic selection in yeast. *Nature,* 364**,** 121-6.
WANG, M. M., TSAI, R. Y., SCHRADER, K. A. & REED, R. R. 1993. Genes encoding components
of the olfactory signal transduction cascade contain a DNA binding site that may
direct neuronal expression. *Mol Cell Biol,* 13**,** 5805-13.
WANG, S. S., TSAI, R. Y. & REED, R. R. 1997. The characterization of the Olf-1/EBF-like HLH
transcription factor family: implications in olfactory gene regulation and neuronal
development. *J Neurosci,* 17**,** 4149-58.
WIESENFAHRT, T., BERG, J. Y., OSBORNE NISHIMURA, E., ROBINSON, A. G., GOSZCZYNSKI,
B., LIEB, J. D. & MCGHEE, J. D. 2016. The function and regulation of the GATA factor
ELT-2 in the C. elegans endoderm. *Development,* 143**,** 483-91.
WYLER, S. C., SPENCER, W. C., GREEN, N. H., ROOD, B. D., CRAWFORD, L., CRAIGE, C.,
GRESCH, P., MCMAHON, D. G., BECK, S. G. & DENERIS, E. 2016. Pet-1 Switches
Transcriptional Targets Postnatally to Regulate Maturation of Serotonin Neuron
Excitability. *J Neurosci,* 36**,** 1758-74.
XUE, D., FINNEY, M., RUVKUN, G. & CHALFIE, M. 1992. Regulation of the mec-3 gene by the
C.elegans homeoproteins UNC-86 and MEC-3. *EMBO J,* 11**,** 4969-79.
YEMINI, E., JUCIKAS, T., GRUNDY, L. J., BROWN, A. E. & SCHAFER, W. R. 2013. A database of
Caenorhabditis elegans behavioral phenotypes. *Nat Methods,* 10**,** 877-9.
YU, B., WANG, X., WEI, S., FU, T., DZAKAH, E. E., WAQAS, A., WALTHALL, W. W. & SHAN, G.
2017. Convergent Transcriptional Programs Regulate cAMP Levels in C. elegans
GABAergic Motor Neurons. *Dev Cell,* 43**,** 212-226 e7.
YU, G., WANG, L. G. & HE, Q. Y. 2015. ChIPseeker: an R/Bioconductor package for ChIP peak
annotation, comparison and visualization. *Bioinformatics,* 31**,** 2382-3.
ZHANG, L., WARD, J. D., CHENG, Z. & DERNBURG, A. F. 2015. The auxin-inducible
degradation (AID) system enables versatile conditional protein depletion in C.
elegans. *Development,* 142**,** 4374-84.
ZHENG, C., JIN, F. Q. & CHALFIE, M. 2015. Hox Proteins Act as Transcriptional Guarantors to
Ensure Terminal Differentiation. *Cell Rep,* 13**,** 1343-1352.
ZHOU, Q., LIU, M., XIA, X., GONG, T., FENG, J., LIU, W., LIU, Y., ZHEN, B., WANG, Y., DING, C. &
QIN, J. 2017. A mouse tissue transcription factor atlas. *Nat Commun,* 8**,** 15089.
1424

Figure 4

A

UNC-3 ChIP

B

Figure 5

A

B

cfi-1::RFP

N.S.

Number of RFP+ MNs

EtOH   Auxin

lin-39::mNG::AID;
eft-3::tir1

Table 1: A summary of the *cis*-regulatory analysis to identify novel transcription factors controlled by UNC-3. Reporter alleles of each TF were built and examined for expression pattern. TFs that are expressed in MNs were further tested for UNC-3 dependency. Ten of 16 tested TFs (62.5%) show expression in MNs, of which 9 require UNC-3 activity for normal expression. Two reporters were generated for *nhr-1* and *nhr-40* because two distinct UNC-3 ChIP-Seq peaks were found in the *cis*-regulatory region of these genes. N/A: Not applicable.

	Reporter	TF family	*cis*-regulatory region included	RFP expression	*unc-3* dependence	Expression	Expression in Cholinergic MNs	Expression in GABAergic MNs	# of lines
1	*nhr-1* (second peak)	Nuclear hormone receptor	-838 bp ~ 1,535 bp	YES	Positively regulated	VNC, Head and Tail Neurons	YES	YES	2
2	*nhr-40* (second peak)	Nuclear hormone receptor	4,356 bp ~ 5,525 bp	YES	Positively regulated	VNC, Head and Tail Neurons	YES	YES	2
3	*mab-9*	T-box	-5,561 bp ~ -3,773 bp	YES	Positively regulated	VNC, Head and Tail Neurons	YES	YES	2
4	*ztf-26*	Zinc finger	910 bp ~ 3,394 bp	YES	Positively regulated	VNC and Head Neurons	YES	NO	2
5	*ceh-44*	C. elegans homeobox	1,605 bp ~ 3,111 bp	YES	Positively regulated	VNC, Head and Tail Neurons	YES	Not determined	2
6	*zfh-2*	Zinc finger	12,048 bp ~ 13,549 bp	YES	Positively regulated	VNC, Head and Tail Neurons	YES	NO	2
7	*cfi-1*	AT-rich interaction domain	-13,824 bp ~ -11,329 bp	YES	Positively regulated	VNC, Head and Tail Neurons	YES	YES	2
8	*bnc-1*	Zinc finger	-1,800 bp ~ -1 bp	YES	Positively regulated	VNC and Tail Neurons	YES	NO	2
9	*nhr-49*	Nuclear hormone receptor	-750 bp ~ 75 bp	YES	Negatively regulated	VNC, Head and Tail	NO	YES	2

						Neurons			
10	*nhr-1* (first peak)	Nuclear hormone receptor	-7,452 bp ~ -5,871 bp	YES	Not tested	Head and Tail Neurons	N/A	N/A	2
11	*nhr-19*	Nuclear hormone receptor	1,261 bp ~ 2,294 bp	YES	NO	VNC, Head and Tail Neurons	Not determined	Not determined	2
12	*nhr-40* (first peak)	Nuclear hormone receptor	919 bp ~ 1,839 bp	YES	NO	VNC, Head and Tail Neurons	Not determined	Not determined	2
13	*ccch-3*	Zinc finger	-385 bp ~ -1 bp	YES	Not tested	Head Neurons	N/A	N/A	2
14	*ztf-17*	Zinc finger	-1,031 bp ~ 3 bp	YES	Not tested	Head Neurons, Muscles, and Intestine	N/A	N/A	2
15	*nhr-47*	Nuclear hormone receptor	-675 bp ~ 171 bp	NO	N/A	N/A	N/A	N/A	2
16	*unc-55*	Nuclear receptor	-2,105 bp ~ -1,327 bp	NO	N/A	N/A	N/A	N/A	2
17	*ztf-13*	Zinc finger	-1,392 bp ~ -1 bp	NO	N/A	N/A	N/A	N/A	2
18	*ztf-14*	Zinc finger	-2,087 bp ~ -531 bp	NO	N/A	N/A	N/A	N/A	2