Whole genome sequencing and phylogenetic analysis of SARS-CoV-2 strains from the index and early patients with COVID-19 in Dubai, United Arab Emirates, 29 January to 18 March 2020

Ahmad Abou Tayoun[1,2 ‡], Tom Loney[2 ‡], Hamda Khansaheb[3], Sathishkumar Ramaswamy[1], Divinlal Harilal[1], Zulfa Omar Deesi[4], Rupa Murthy Varghese[4], Hanan Al Suwaidi[2], Abdulmajeed Alkhajeh[3], Laila Mohamed AlDabal[5], Mohammed Uddin[2,6], Rifat Hamoudi[7], Rabih Halwani[7], Abiola Senok[2], Qutayba Hamid[8], Norbert Nowotny[2,9*], Alawi Alsheikh-Ali[2*]

[1]Al Jalila Genomics Center, Al Jalila Children's Hospital, Dubai, United Arab Emirates.

[2]College of Medicine, Mohammed Bin Rashid University of Medicine and Health Sciences, Dubai, United Arab Emirates.

[3]Medical Education & Research Department, Dubai Health Authority, Dubai, United Arab Emirates.

[4]Microbiology and Infection Control Unit, Pathology and Genetics Department, Latifa Women and Children Hospital, Dubai Health Authority, Dubai, United Arab Emirates.

[5]Medical Affairs Department, Rashid Hospital, Dubai Health Authority, Dubai, United Arab Emirates.

[6]The Centre for Applied Genomics, The Hospital for Sick Children, Toronto, ON, Canada

[7]Clinical Sciences Department, College of Medicine, University of Sharjah, Sharjah, United Arab Emirates.

[8]College of Medicine, University of Sharjah, Sharjah, United Arab Emirates.

[9]Institute of Virology, University of Veterinary Medicine Vienna, Vienna, Austria.

*Corresponding authors: Norbert Nowotny, Norbert.Nowotny@vetmeduni.ac.at; Alawi Alsheikh-Ali, Alawi.Alsheikhali@mbru.ac.ae.

‡ Joint first authorship.

Author Contributions

AAA, AT, NN, and TL conceived the study and drafted the protocol. All authors provided critical input into the protocol. AAA, AAK, and HK coordinated the ethical approval and sample retrieval. RV and ZD conducted the RT-qPCR analysis. AT, SR, and DH performed the whole genome sequencing and phylogenetic analysis. HK performed data extraction from the medical records and TL completed data analysis for the manuscript. AT and TL drafted the manuscript, AAA and NN refined it before AAK, AS, DH HA, HK, MU, QH, RH, RHA, RV, SR, and ZD, provided comments and feedback on the first draft. All authors read and approved the final version of the manuscript.

Competing Interest Statement: Authors have no conflicts of interest to disclose.

This file includes:

> Main Text
> Figures 1 to 2
> Table 1

Word Count: 2,638

Reference Count: 11

Significance

Dubai in the United Arab Emirates has become the major international tourism and travel hub within the Middle East connecting the far East and the West. Understanding the early spread of SARS-CoV-2 in Dubai will, therefore, inform the global dynamics of the COVID19 pandemic spread. Compared to the rest of the world, a very limited number of SARS-CoV-2 genome sequences have been published from the Middle East. This study establishes SARS-CoV-2 full genome sequences from the index and early COVID19 patients in the UAE, which are analyzed in the context of virus introductions, chain of transmissions, and possible links to earlier strains in the rest of the world. This study complements the world map of the pandemic evolution.

Abstract

Whole genome sequencing and phylogenetic analysis of SARS-CoV-2 strains from the index and early patients with COVID-19 in Dubai (United Arab Emirates; UAE) showed multiple spatiotemporal introductions from Asia, Europe, and the Middle East. We combine genetic, demographic, and clinical information from early patients to show that the majority of introductions were from Europe and the Middle East/Iran, and to provide evidence for early community-based transmission. We further catalogue new mutations in SARS-CoV-2 isolates in the UAE. Our findings can be used to further understand the global transmission network of SARS-CoV-2.

Introduction

In December 2019, several cases of a new respiratory illness (now called COVID-19) were reported in the city of Wuhan (Hubei province, China) and in January 2020 it was confirmed these infections were caused by a novel coronavirus subsequently named SARS-CoV-2 [1-2]. On 12 March 2020, the ongoing SARS-CoV-2 outbreak was declared a pandemic by the World Health Organization (WHO) [3]. To date (13 May 2020), there have been over 4.2 million laboratory-confirmed cases of COVID-19 and more than ~290,000 deaths in 187 countries [4].

Dubai in the United Arab Emirates (UAE) is a cosmopolitan metropolis that has become a popular tourist destination and home to one of the busiest airport hubs in the world connecting the east with the west [4]. Currently, the UAE has reported 19,661 confirmed cases and 203 deaths associated with COVID-19 (1.0% case fatality; 13 May 2020) [5]. In view of Dubai's important tourism and travel connections, we attempted to characterize the full-genome sequence of SARS-CoV-2 strains from the index and early patients with COVID-19 in Dubai to gain a deeper understanding of the molecular epidemiology of the outbreak in Asia, Europe, and the Middle East.

Results

Patient Cohort and SARS-CoV-2 Whole Genome Sequencing

The 49 patients included in this study were the earliest confirmed cases in the UAE. The time period of 29 January to 18 March 2020 was specifically selected to focus on early SARS-CoV-2 viral introductions into the UAE. The first index patient in the UAE was reported on the 29 January 2020. Subsequently, Emirates airlines suspended flights to and from 30 destinations across the world from 18 March and Dubai airport was closed to passenger flights on 25 March 2020; hence, patients after 18 March 2020 were expected to be more likely a result of community transmission as

opposed to imported infections. In brief, the index patient with COVID-19 in the UAE was a female Chinese tourist (aged 63 years) travelling from Wuhan with her daughter-in-law and two grandchildren to visit her son working in Dubai. The Chinese family arrived in Dubai on 16 January 2020 and the grandmother, son, daughter-in-law, and one grandchild all tested positive on the 29 January 2020 (**Table 1**). Over the next seven weeks, there were multiple new travel-related possible introductions of the virus into the UAE from European and Iranian tourists and residents, and a cluster of cases amongst residents with no travel history (**Table 1**). Of the 49 COVID-19 patients in Dubai, 22 had a travel history from Europe including Austria, France, Germany, Italy, Ireland, Norway, and the United Kingdom (**Table 1**). Nearly two-thirds (63.3%) of patients were male and 61.2% were aged between 20 and 44 years reflecting the young age structure of the UAE population [4]. Majority of patients (88%) were asymptomatic or had mild symptoms and only four required intensive care with invasive ventilation. There were strong occupational links in several patients including a woman who flew from Italy to Dubai for business (L5752), two airline cabin crew with a positive travel history from Austria and Italy (L1014, L1076, respectively), and two members of a professional cycling team (L9333, L9756) who tested positive during the UAE 2020 Cycle Tour. The seven-stage cycle tour was cancelled on 27 February 2020 following the fifth stage when numerous riders and support staff tested positive and all riders and teams had a positive travel history from Europe following previous cycle races within the last 7-14 days.

SARS-CoV-2 whole genome sequencing was performed on all 49 COVID-19 patient samples. Only genomes with almost complete coverage (n=25, **Methods**) were used for phylogenetic analysis. The 25 genomes were obtained from cases with disease onsets in late January (n=1), early February (n=1), late February (n=6), early March (n=8), and late March (n=9). Of those, 17 (68%) were males, and age ranges were 10-40 years (n=16), 40-59 years (n=5), >60 years (n=2), and unreported for two individuals.

Phylogenetic analysis

To understand early viral transmission in Dubai in the global context, we performed phylogenetic analysis on the 25 novel viral genomes we sequenced from early patients in the UAE (**Table 1**) in this study (**Methods**) along with 157 largely complete SARS-CoV-2 genomes deposited in GISAID from different countries between December 2019 and early March 2020 [6] (**Table S1)**.

Consistent with multiple independent introductions, the UAE SARS-CoV-2 isolates were distributed across the phylogenetic tree (**Figure 1**). The majority (76%) clustered with clades A2a (48%) and A3 (28%) which are largely composed of isolates from COVID19 patients in Europe and Iran, respectively. This clearly suggests that the major introductions into the UAE during the early phase of the pandemic originated from Europe and the Middle East/Iran.

Supporting its European origin, all individuals with the A2a clade isolates were mostly European and/or with recent travel history to a European country, mainly to Italy (n=4), Germany (n=3), United Kingdom (n=2), Spain (n=1), and Norway (n=1) (**Table 1** and **Figure 2**). Onset of symptoms reported in this group was within or after the second week of March (**Table 1**) suggesting that the viral infections in this group could have occurred during late February to early March. Of note, a SARS-CoV-2 isolate submitted from Mexico (GISAID ID: EPI_ISL_412972) was 100% identical to that of an Italian expatriate working in the UAE (L0881), while another submitted in Germany (GISAID ID: EPI_ISL_412912) differed by a single mutation (**Figure 1**). All three individuals had a recent travel history to Italy and overlapping infection time frames (late February – early March). Within this group, isolates from patients L1758, L0484, and L2185 were identical (**Figure 2**) suggesting a possible common direct source of transmission.

Isolates in the A3 clade were obtained from five individuals with travel history to Iran (L2409, L6627, L0904, L0184, and L4682) and one Indian resident (L0231) and one Indian tourist (L0068) (**Figure 2)**. Onset of symptoms for the five individuals with travel history in this group was reported to be

around 21-24 February (**Table 1**). Patient L0231 had no travel history and reported symptoms onset on 7 March highly suggesting a possible community-based transmission event. Interestingly, all but one isolate obtained from patient L4682 – the only patient in this group with severe clinical presentation – shared a common ancestral strain identical to that obtained from patient L2409. The SARS-CoV-2 isolate from L4682 had two unique missense variants in the *ORF1ab* gene (**Table S2**) which might be worth investigating for any possible biological effect(s). Consistent with its Iranian origin, a SARS-CoV-2 sequence submitted by the University of Sydney (GISAID ID: EPI_ISL_412975) on 28 February 2020 differed by only two mutations from that of L2409, and both this Iranian male tourist and the Australian male had a recent travel history to Iran. We speculate that individuals with travel history to Iran around this time frame (L8386, L6867, and L3280), for whom a full viral genome sequence could not be obtained, were also very likely to cluster within the A3 clade.

Only one viral strain obtained from L5630, a family member of the early Chinese index patient, belonged to the B2 clade. Although we did not obtain full viral genome sequences from the other members of that Chinese family, we expect that all had a similar strain to L5630. Interestingly, our data do not suggest any transmission of this clade at least among the earliest patients (**Figure 2**) included in this study which is consistent with the reported early detection and isolation of this family. This finding also supports the notion of secondary source(s) for the ongoing local transmission.

The remaining five isolates did not belong to A2a, A3, B2, or any of the clades on nextrain.org as of 12 May 2020, suggesting earlier introduction(s). Those isolates were obtained from four Asians, two residents (L4280, L6599) and two tourists (L4184, L9766), and one Czech resident (L1014) working as an airline cabin crew with travel history to Austria (**Table 1**). Consistent with the Asian predominance among this patient group and the fewer (1 or 2) mutations for most of their isolates (4 out of 5) relative to the Wuhan reference genome (**Figure 2**), several early viral strains submitted in Asia clustered very closely to this group (**Figure 1**). L4280 was the first sequenced patient without travel history and became infected after transporting a work colleague with severe/critical symptoms to hospital (L0826; not sequenced; no travel history). Patient L0826 reported symptom onset on 22 January suggesting that community-based transmission started in the UAE in early-to-mid January. L6599 was an Indian expatriate living with three other Filipino and Sri Lankan expatriates (L3715, L2771, L8480) (**Table 1**). All four individuals had no documented recent travel history suggesting local transmission, and although full viral genome sequences could not be obtained from all but patient L6599, it is very likely that all have related isolates.

In aggregate, we identified 70 variants relative to the reference GenBank SARS-CoV-2 sequence NC_045512.2. The majority of these variants were missense (n=41) with the most frequent nucleotide change being C>T (n=33), and more than half (38/70) were localized in the *ORF1ab* gene (**Table S2**). Notably, 17 out of the 70 variants were novel as they were not identified in the Chinese National Center for Bioinformation Database (https://bigd.big.ac.cn/ncov/variation/annotation; last accessed April 29, 2020). Of those, 12 were coding missense variants distributed across the *ORF1ab* (n=8), *S* (n=1), *ORF7a* (n=1), and the *N* (n=2) genes. Assessing the biological significance of these variants is critical for determining the efficacy of any future therapies and/or vaccines.

Discussion

Our findings suggest multiple independent spatiotemporal introductions of SARS-CoV-2 into the UAE from Asia, Europe, and the Middle East. We show that the majority of introductions (78%) were from Iran and Europe during two different time frames (mid-late February and early March, respectively). Although we show evidence for possible local transmission within the Middle Eastern/Iranian isolates, it will be important to sequence further isolates at subsequent dates to determine whether these introductions succeeded in seeding more clustering and whether such

clustering was affected by proactive and vigilant public health measures. Nation-wide public health measures included immediate isolation of positive cases, rapid and comprehensive contact tracing and testing, transitioning to online learning for schools and universities, implementing work-from-home protocols across all sectors, nationwide disinfection campaigns, and shielding priority populations including older adults, people of determination, and at-risk individuals with underlying health conditions (e.g. cancer patients).

Six isolates (22%) did not cluster with the European or Iranian groups and represented earlier introductions which did not appear to seed larger clusters in our sampled cohort. However, additional sequencing is needed to determine the extent of community transmission, especially given that our data strongly suggest that the earliest patient (early to mid-January) in the UAE could have been a secondary infection from one of those introductions.

The new mutations and novel amino acid changes found in the viruses identified in the UAE warrant further investigation to explore whether they influence viral characteristics, especially pathogenicity, or provide important information for vaccine development. One of the major strengths of the study was the non-biased representative sample of early cases, including the index family cluster, in Dubai from the only central testing lab, along with detailed demographic and clinical information (i.e. age, gender, travel history, disease severity, symptoms onset, etc.). Limitations included the inability to conduct full whole genome sequencing on more samples most likely due to low viral load issues, although we were able to deduce the origin of transmission in most of those individuals based on travel history. Regardless, this study contributes important molecular epidemiological data that can be used to further understand the global transmission network of SARS-CoV-2.

Materials and Methods

Human Subjects and Ethics Approval

The electronic medical records of the earliest 49 patients with laboratory confirmed SARS-CoV-2 from 29 January to 18 March 2020 were reviewed and sociodemographic and clinical data extracted using the WHO case report form. This study was approved by the Dubai Scientific Research Ethics Committee - Dubai Health Authority (approval number #DSREC-04/2020_02).

SARS-CoV-2 Whole Genome Sequencing

All 49 COVID-19 patients tested positive for SARS-CoV-2 by RT-qPCR in the central Dubai Health Authority virology laboratory where viral RNA was extracted from nasopharyngeal swabs following the QIAamp Viral RNA Mini or the EZ1 DSP Virus Kits (Qiagen, Hilden, Germany). Extracted genomic RNA from all samples were then subjected to shotgun metagenomic sequencing using the Illumina (San Diego, CA, USA) next-generation sequencing (NGS) platform at Al Jalila Children's Specialty Hospital, Dubai, UAE. Approximately 1000ng of input RNA from each patient sample was used to prepare sequencing-ready libraries following the TruSeq Stranded Total RNA Library Prep Gold kit from Illumina (San Diego, CA, USA). Human ribosomal RNA was first depleted, and then the remaining RNA underwent fragmentation, reverse transcription (using the SuperScript II Reverse Transcriptase Kit from Invitrogen, Carisbad, USA), adaptor ligation, and amplification. Libraries were sequenced using the Illumina SP Reagent kit (2 X 150 cycles) and the NovaSeq platform (San Diego, CA, USA). Sample L5630 underwent a target enrichment approach where double stranded DNA (synthesized using the QuantiTect Reverse Transcription Kit from Qiagen, Hilden, Germany) was amplified using 26 overlapping primer sets covering most of SARS-CoV-2 genome [7]. PCR products were then sheared by ultra-sonication (Covaris LE220-plus series, MA, USA) and prepared for sequencing using the SureSelectXT Library Preparation kit (Agilent, CA, USA). This library was sequenced using the MiSeq Micro Reagent Kit, V2 (2 X 150 cycles) on the MiSeq platform (Illumina, San Diego, CA, USA).

SARS-CoV-2 Genome Assembly

High quality (>Q30) sequencing reads, obtained through shotgun or targeted enrichment, were trimmed and then aligned to the reference SARS-CoV-2 genome from Wuhan, China (GenBank accession number: NC_045512.2) using a custom-made bioinformatics pipeline (**Fig. S1**). Assembled genomes with at least 20X average coverage across most nucleotide positions (56-29,797) were used for subsequent phylogenetic analysis (**Table S3**). A total of 25 viral genomes (24 by shotgun and 1 by target enrichment) met this inclusion criterion and were submitted to the Global Initiative on Sharing All Influenza Data (GISAID) database under accession IDs: EPI_ISL_435119-435142 (**Table S1**).

Phylogenetic analysis

We downloaded 157 global non-UAE sequences (**Table S1**) with largely complete genomes (nucleotide positions 56-29,797) submitted to GISAID EpiCoV (https://www.epicov.org/) between December 2019 and 04 March 2020 [6]. All 182 sequences, including the 25 UAE sequences generated in this study, were analyzed using Nexstrain [8], which consists of Augur v6.4.3 pipeline for multiple sequence alignment (*MAFFT* v7.455 [9]) and phylogenetic tree construction (*IQtree* v1.6.12 [10]). Tree topology was assessed using the fast bootstrapping function with 1000 replicates. Tree visualization and annotations were performed in *FigTree* v1.4.4 [11] for **Figure 1** and in auspice v2.13.0 tool [8] for **Figure 2**. SARS-CoV-2 clades annotations were performed in auspice v2.13.0 and cross-checked with nextstrain.org as of 12 May 2020.

References

1. Ashour HM, Elkhatib WF, Rahman MM, Elshabrawy HA. Insights into the Recent 2019 Novel Coronavirus (SARS-CoV-2) in Light of Past Human Coronavirus Outbreaks. *Pathogens.* 9(3):E186 (2020).
2. Uddin M, Mustafa F, Rizvi TA, Loney T, Al Suwaidi H, Al Marzouqi A, et al. SARS-CoV-2/COVID-19: Viral Genomics, Epidemiology, Vaccines, and Therapeutic Interventions. *Viruses.* 12(5):526 (2020).
3. World Health Organization (WHO). Coronavirus disease 2019 (COVID-19) Situation Report – 52. Geneva: WHO; 12 Mar 2020. Available from: https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200312-sitrep-52-covid-19.pdf?sfvrsn=e2bfc9c0_4
4. Johns Hopkins Center for Systems Sciences and Engineering. COVID19 Dashboard. Available from: https://www.arcgis.com/apps/opsdashboard/index.html#/bda7594740fd40299423467b48e9ecf6
5. Loney T, Aw TC, Handysides DG, Ali R, Blair I, Grivna M, et al. An analysis of the health status of the United Arab Emirates: the 'Big 4' public health issues. *Glob Health Action.*;6:20100 (2013).
6. Forster P, Forster L, Renfrew C, Forster M. Phylogenetic network analysis of SARS-CoV-2 genomes. *Proc Natl Acad Sci U S A.* 117(17):9241-9243 (2020).
7. Wu F, Zhao S, Yu B, Chen Y-M, Wang W, Song Z-G, et al. A new coronavirus associated with human respiratory disease in China. *Nature.* 579(7798):265-269 (2020).
8. Hadfield J, Megill C, Bell SM, Huddleston J, Potter B, Callender C, et al. Nextstrain: real-time tracking of pathogen evolution. *Bioinformatics.* 34(23):4121-4123 (2018).
9. Katoh K, Misawa K, Kuma K, Miyata T. MAFFT: A Novel Method for Rapid Multiple Sequence Alignment Based on Fast Fourier Transform. *Nucleic Acids Res.* 15;30(14):3059-66.
10. Chernomor O, von Haeseler A, Quang Minh B. Terrace Aware Data Structure for Phylogenomic Inference from Supermatrices. *Systematic Biology.* 65(6):997-1008 (2016).

11. Rambaut. A. FigTree 1.4.2 Software. *Institute of Evolutionary Biology, Univ. Edinburg.*

Figures

Figure 1. Phylogenetic relationships of SARS-CoV-2 isolates from early patients in Dubai and early global strains. A maximum likelihood phylogeny of 182 SARS-CoV-2 genomes (157 obtained from GISAID as of early Match and 25 genomes in this study). Bootstrap support values >70% supporting major branches are shown. The 3 non-UAE isolates in clades A2a and A3 namely, Mexico/CDMX-InDRE_01, Germany/Baden/Wuerttemberg-1, and Australia/NSW05 are the GISAID ID: EPI_ISL_412972, GISAID ID: EPI_ISL_412912, and GISAID ID: EPI_ISL_412975, respectively, referred to in the main text. Scale bar represents number of nucleotide substitutions per site. UAE = United Arab Emirates.

Figure 2. Phylogenetic relationship of early SARS-CoV-2 isolates in the UAE. A maximum likelihood phylogenetic tree of all 25 UAE SARS-CoV-2 sequences, generated in this study, is shown. The two Wuhan genomes (Wuhan-Hu-1/2019, GISAID ID: EPI_ISL_402125 and Wuhan/WH01/2019, GISAID ID: EPI_ISL_406798) were used as reference genomes (blue filled circles). UAE viral strains (black filled circles) were labeled with sample ID and Nationality (in brackets). Bootstrap support values >70% supporting major branches are shown. Branch lengths mark divergence from the reference Wuhan SARS-CoV-2 genome (GenBank accession number: NC_045512.2) in mutations numbers, while branch color represents travel history. Travel history to Europe for patients UAE/L1758, UAE/L0484, UAE/L2185, and UAE/L0881 is obscured by vertical lines. UAE = United Arab Emirates.

Table 1. Sociodemographic and clinical characteristics of the index and early patients (n=49) with laboratory-confirmed SARS-CoV2 in Dubai, United Arab Emirates, 29 January-18 March 2020

Study ID	Age (years)	Sex	Nationality	Resident /Tourist	Travel History	Symptom Onset Date 2020‡	Severity	Self-Reported History Fever/Cough	ICU Admission/ Ventilator	Full Genome Sequence	GISAID Accession ID
L8156	63	Female	China	Tourist	Wuhan, China	09 Jan	Asymp/Mild	No/Yes	No/No	No	-
L8497	38	Female	China	Tourist	Wuhan, China	24 Jan	Asymp/Mild	Yes/No	No/No	No	-
L5630*	9	Male	China	Tourist	Wuhan, China	28 Jan	Asymp/Mild	No/No	No/No	Yes	EPI_ISL_435137
L8205	36	Male	China	Resident	Wuhan, China	28 Jan	Asymp/Mild	No/No	No/No	No	-
L0826	42	Male	Philippines	Resident	None/Contact with positive case	22 Jan	Severe/Critical	Yes/Yes	Yes/Yes	No	-
L4280	36	Male	India	Resident	None/Contact with positive case	08 Feb	Asymp/Mild	No/No	No/No	Yes	EPI_ISL_435134
L3715	34	Male	Philippines	Resident	None/Contact with positive case	16 Feb	Asymp/Mild	No/No	No/No	No	-
L2771	25	Male	Sri Lanka	Resident	None/Contact with positive case	16 Feb	Asymp/Mild	No/No	No/No	No	-
L8480	21	Male	Sri Lanka	Resident	None/Contact with positive case	16 Feb	Asymp/Mild	No/No	No/No	No	-
L8386	70	Male	Iran	Tourist	Iran	17 Feb	Severe/Critical	No/Yes	Yes/Yes	No	-
L6599	41	Male	Pakistan	Resident	None/Contact with positive case	21 Feb	Asymp/Mild	No/No	No/No	Yes	EPI_ISL_435138
L0904	35	Male	Iran	Tourist	Iran	21 Feb	Asymp/Mild	Yes/Yes	No/No	Yes	EPI_ISL_435126
L6867	60	Female	Iran	Tourist	Iran	22 Feb	Asymp/Mild	No/Yes	No/No	No	-
L2409	59	Male	Iran	Tourist	Iran	23 Feb	Asymp/Mild	Yes/No	No/No	Yes	EPI_ISL_435131
L0184	64	Female	Iran	Tourist	Iran	23 Feb	Asymp/Mild	Yes/Yes	No/No	Yes	EPI_ISL_435121

L4682	64	Male	Bahrain	Tourist	Iran	24 Feb	Severe/Critical	Yes/No	Yes/Yes	Yes	EPI_ISL_435135
L6627	29	Male	China	Resident	Iran	24 Feb	Asymp/Mild	Yes/No	No/No	Yes	EPI_ISL_435139
L1802	38	Male	Dominica	Resident	Iran and Pakistan	28 Feb	Moderate	Yes/Yes	No/No	No	-
L7100	44	Female	Iran	Resident	Iran	29 Feb	Asymp/Mild	Yes/No	No/No	No	-
L5752	38	Female	Italy	Tourist	Italy	04 Mar	Asymp/Mild	No/No	No/No	No	-
L8477	36	Female	India	Resident	US/contact with positive case	05 Mar	Asymp/Mild	Yes/No	No/No	No	-
L5970	70	Female	Italy	Tourist	Milan, Italy	05 Mar	Severe/Critical	Yes/No	Yes/Yes	No	-
L3280	3	Male	Saudi Arabia	Tourist	Iran	07 Mar	Asymp/Mild	No/No	No/No	No	-
L7356	23	Male	Slovakia	Resident	China and Asia	07 Mar	Asymp/Mild	No/No	No/No	Yes	EPI_ISL_435140
L0509	36	Male	Kuwait	Resident	Austria and Germany	07 Mar	Asymp/Mild	Yes/No	No/No	No	-
L0231	54	Male	India	Resident	None	07 Mar	Moderate	Yes/Yes	No/No	Yes	EPI_ISL_435122
L9766	28	Female	Indonesia	Tourist	Indonesia	07 Mar	Asymp/Mild	Yes/Yes	No/No	Yes	EPI_ISL_435142
L4184	28	Female	India	Tourist	Ireland and France	08 Mar	Asymp/Mild	Yes/No	No/No	Yes	EPI_ISL_435133
L0484	28	Male	Canada	Resident	Germany	09 Mar	Asymp/Mild	Yes/Yes	No/No	Yes	EPI_ISL_435123
L1758	30	Male	Lebanon	Resident	Germany	10 Mar	Asymp/Mild	No/No	No/No	Yes	EPI_ISL_435129
L5621	58	Male	Australia	Tourist	Italy	10 Mar	Asymp/Mild	Yes/No	No/No	Yes	EPI_ISL_435136
L9440	35	Male	UK	Resident	Italy	10 Mar	Asymp/Mild	No/No	No/No	Yes	EPI_ISL_435141
L0860	46	Male	France	Resident	US through France	10 Mar	Asymp/Mild	No/Yes	No/No	No	-
L2185	46	Male	Norway	Tourist	Norway	12 Mar	Asymp/Mild	No/No	No/No	Yes	EPI_ISL_435130
L9928	42	Female	Spain	Resident	Spain	13 Mar	Asymp/Mild	Yes/No	No/No	No	-

L0068	33	Male	India	Tourist	India	13 Mar	Asymp/Mild	Yes/No	No/No	Yes	EPI_ISL_435120
L1014	27	Female	Czech Rep.	Resident	Austria	13 Mar	Asymp/Mild	Yes/Yes	No/No	Yes	EPI_ISL_435127
L1076	30	Female	Italy	Resident	Italy	14 Mar	Asymp/Mild	No/No	No/No	Yes	EPI_ISL_435128
L0000	27	Female	Serbia	Resident	UK/contact with positive case	14 Mar	Asymp/Mild	No/No	No/No	Yes	EPI_ISL_435119
L3779	32	Female	Spain	Resident	Spain	14 Mar	Asymp/Mild	Yes/Yes	No/No	Yes	EPI_ISL_435132
L7355	20	Male	UAE	Resident	UK	14 Mar	Asymp/Mild	No/Yes	No/No	No	-
L0549	24	Female	Lebanon	Resident	None/contact with positive case	15 Mar	Asymp/Mild	No/No	No/No	No	-
L9768	19	Male	UAE	Resident	UK	18 Mar	Asymp/Mild	No/Yes	No/No	Yes	EPI_ISL_435143
L4519	54	Male	Iran	Resident	Iran	NR	NR	NR	NR	No	-
L9333	26	Male	Italy	Resident	Italy	NR	NR	NR	NR	No	-
L9756	53	Male	Italy	Resident	Italy	NR	NR	NR	NR	No	-
L0879	NR	Female	Jordan	Resident	Germany	NR	NR	NR	NR	Yes	EPI_ISL_435124
L0880	NR	Female	Germany	Resident	Germany	NR	NR	NR	NR	No	-
L0881	NR	Male	Italy	Resident	Italy	NR	NR	NR	NR	Yes	EPI_ISL_435125

*There was a second grandchild in the Chinese family cluster from Wuhan who remained negative throughout all testing. ‡Self-reported onset of symptoms extracted from medical records; NR, Not Reported; UK, United Kingdom; UAE, United Arab Emirates; US, United States.