Power and limits of selection genome scans on temporal data from a selfing population

Miguel Navascués[1,2,3,✉], **Arnaud Becheler**[1,*], **Laurène Gay**[4], **Joëlle Ronfort**[4], **Karine Loridon**[4], and **Renaud Vitalis**[1,2]

[1]CBGP, INRAE, CIRAD, IRD, Montpellier SupAgro, Univ Montpellier, Montpellier, France
[2]Institut de Biologie Computationnelle, Montpellier, France
[3]Human Evolution, Department of Organismal Biology, Uppsala University, Uppsala, Sweden
[4]AGAP, INRAE, CIRAD, IRD, Montpellier SupAgro, Univ Montpellier, Montpellier, France
[*]current affiliation: Department of Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, USA

Tracking genetic changes of populations through time allows a more direct study of the evolutionary processes acting on the population than a single contemporary sample. Several statistical methods have been developed to characterize the demography and selection from temporal population genetic data. However, these methods are usually developed under the assumption of outcrossing reproduction and might not be applicable when there is substantial selfing in the population. Here, we focus on a method to detect loci under selection based on a genome scan of temporal differentiation, adapting it to the particularities of selfing populations. Selfing reduces the effective recombination rate and can extend hitch-hiking effects to the whole genome, erasing any local signal of selection on a genome scan. Therefore, selfing is expected to reduce the power of the method. By means of simulations, we evaluate the performance of the method under scenarios of adaptation from new mutations or standing variation at different rates of selfing. We find that the detection of loci under selection in predominantly selfing populations remains challenging even with the adapted method. Still, selective sweeps from standing variation on predominantly selfing populations can leave some signal of selection around the selected site thanks to historical recombination before the sweep. Under this scenario, ancestral advantageous alleles at low frequency leave the strongest local signal, while new advantageous mutations leave no local footprint of the sweep.

selective sweep | neutrality test | temporal differentiation | *Medicago truncatula*

Introduction

Several evolutionary processes (such as migration, selection or drift) can change the genetic make-up of populations through time. Thus, patterns of genetic diversity can inform us about the evolutionary history of the populations (see Pool *et al.* 2010, for a review). However, observing the genetic diversity changes through time (instead of at a single time point) can provide more precise information about the evolutionary processes in action.

Since the beginning of the 20th century, researchers have used repeated observations of hereditary characters in the same populations (e.g. color patters in *Diabrotica soror*, Kellogg and Bell 1904) or subfossil records (e.g. banding patterns in *Cepaea* snails, Diver 1929) to study evolution. An emblematic example is the time-series data on the frequency of the *medionigra* phenotype in a population of the moth *Callimorpha dominula*, which inspired the discussion on the prevalence of selection over drift between Fisher and Wright (Fisher and Ford 1947; Wright 1948) and has continued to offer insight into the evolutionary process through recent re-analyses (e.g. Foll *et al.* 2014, and references therein).

The technological advances in molecular genetics have allowed these temporal studies to switch from Mendelian characters to polytene chromosomes (e.g. Dobzhansky 1943), isozymes (e.g. Yamazaki 1971) and, eventually, to high throughput DNA sequencing (e.g. Frachon *et al.* 2017). Indeed, molecular genetics has opened the door to the study of short-generation-time microorganisms (Biek *et al.* 2015), ancient samples (e.g. subfossil samples, museum and herbaria specimens; Leonardi *et al.* 2017) and experimental populations (Schlötterer *et al.* 2015), allowing for an increasing availability of temporal population genetic data.

Temporal population genetic data allows to study the change of allele frequencies through time. In the absence of migration, mutation and selection, these changes are the product of genetic drift. As such, they can be used to estimate the effective population size, N_e, either with moment based (e.g. Krimbas and Tsakas 1971; Nei and Tajima 1981; Waples 1989) or likelihood-based approaches (e.g. Anderson *et al.* 2000; Williamson and Slatkin 1999). If a sample from the source of migration is available, it is also possible to co-estimate migration and drift with an extension of the likelihood method (Wang and Whitlock 2003). These methods all assume short timescales and low mutation rates, so that no new mutations arrive during the studied period. However, for temporal data over larger scales in which mutations can no longer be neglected, it is also possible to co-estimate substitution rate with effective population size (Drummond *et al.* 2002; Rambaut 2000). Finally, like for migration or mutation rates, it is also

possible to make inferences about selection. Two strategies can be followed. First, three or more temporal samples may be used to separate the random component (drift) from the systematic component (selection) in the changes of allele frequencies (e.g. Bollback *et al.* 2008; Buffalo and Coop 2019; Feder *et al.* 2014). Alternatively, if only two temporal samples are available, loci under selection may be detected by an outlier approach (e.g. Goldringer and Bataillon 2004, described in more detail below).

Most of the statistical methods in population genetics, including those mentioned above, have been developed for outcrossing populations. Methods specifically adapted to selfing populations are scarce (Hartfield *et al.* 2017). Nevertheless, many plants reproduce through selfing or partial selfing (see Whitehead *et al.* 2018, for a recent overview), including an important proportion of the species considered in temporal monitoring programs (e.g. supplementary text S1). Selfing, by increasing homozygosity and linkage disequilibrium, shapes the genetic diversity of population in a particular way (Golding and Strobeck 1980; Vitalis and Couvet 2001). Notably, it generates repeated multilocus genotypes that can persists over several generations due to the lack of effective recombination (Jullien *et al.* 2019). The dynamics of adaptation is also impacted by selfing, with a general reduction in the efficacy of selection (e.g. Burgarella *et al.* 2015). In the light of these drastic effects, there is a need for development or adaptation of methods to take into account different mating systems.

Even if we can adapt methods to release the assumption of random mating, distinguishing selection from demography in highly selfing species may be problematic because of the reduced effective recombination due to self-fertilization. Selective sweeps in these populations could involve a genome-wide hitch-hiking effect, which prevents any difference in genetic diversity between the neutral and adaptive regions. If that were the case, even methods adapted to selfing would fail to detect regions under selection, which questions the relevance of temporal genome scans in predominantly selfing populations. On the other hand, a temporal genome scan on a highly selfing *Arabidopsis thaliana* population ($\sigma \approx 0.94$) revealed several outliers regions with compelling evidence for the action of selection on them (Frachon *et al.* 2017). Therefore, in the planning of future research, there is a need to understand under which circumstances selfing imposes a limit for the detection of loci under selection.

In this work, we introduce several modifications to the temporal genome scan approach proposed by Goldringer and Bataillon (2004) to take into account partial self-fertilization. Then, by means of simulated data, we evaluate the performance of that method for the estimation of the effective population size (the first step of this genome scan) and the detection of regions

under selection under different scenarios of adaptation (from new mutations or from standing variation) and selfing. Our results highlight the importance of taking into account the mating system in the analysis of population genetic data. They also highlight a threshold beyond which loci under selection can be detected for highly selfing populations. We applied the approach to a population of the predominantly selfing species *Medicago truncatula* and re-discuss some of the results from Frachon *et al.* (2017) temporal genome scan on *Arabidopsis thaliana*.

Materials and Methods

Overview of the Genome Scan Method. In a single isolated population, allele frequencies change through time under the action of selection, that acts upon specific loci, and genetic drift, that acts upon the whole genome. In order to identify loci subjected to selection, we use a procedure inspired from the test for homogeneity of differentiation across loci by Goldringer and Bataillon (2004). The principle is that, in the case of complete neutrality across loci, all sampled markers should provide estimates of genetic differentiation drawn from the same distribution. Assuming a single isolated population, this distribution depends on the strength of genetic drift, that is, on the length of the period, τ in number of generations, and on the effective population size, N_{e} (see Table S1 for a summary of notation). On the other hand, if some of the studied polymorphisms are under selection or linked to selected variants, we expect some heterogeneity in the distribution of differentiation values, because directional selection induces larger values of genetic differentiation than expected under the neutral case. The approach we describe uses the expected distribution of temporal differentiation to identify those polymorphisms showing outlier values compared to a neutral expectation. In Frachon *et al.* (2017), we already made some modifications (to account for the uncertainty of the initial allele frequency) to the approach proposed by Goldringer and Bataillon (2004) for the standard (random-mating or haploid) case. Here, we present further modifications for a more general case including partial self-fertilization.

Estimation of effective population size. The estimated magnitude of drift between two time samples is used as a null model in this temporal genome scan method. Temporal differentiation can be measured by estimating the F_{ST} with the analysis of variance approach proposed by Weir and Cockerham (1984). Weir and Cockerham's (1984) analysis of variance partitions variance within individuals, among individuals within population and between populations, allowing to account for the correlation of allele identity within individuals due to selfing in the F_{ST} estimate. Temporal F_{ST} can be used

to estimate N_e as $\hat{N}_e = \frac{\tau(1-\hat{F}_{ST})}{4\hat{F}_{ST}}$ (Frachon *et al.* 2017; Skoglund *et al.* 2014).

Building null distribution of drift. In order to test the homogeneity between the focal-locus l and genome-wide differentiation, the null distribution for single locus F_{ST}^l is build through simulations of drift. For these simulations, Goldringer and Bataillon (2004) considered the observed allele frequency in the sample as the initial allele frequency in the population, π_0 (from this point subscript 0 indicates values at time $t = 0$). This approach ignores the uncertainty due to sampling. Instead, in Frachon *et al.* (2017), we improved this step by assuming that allele counts observed in a sample of n_0 diploid individuals come from a binomial distribution $B(2n_0, \pi_0)$, where π_0 is the (unknown) allele frequency in the population. Using Bayes inversion formula and assuming a uniform prior for the allele frequency, this allows to sample from the posterior probability distribution with $\text{Beta}(k_0 + 1, 2n_0 - k_0 + 1)$, where k_0 is the observed count of the reference allele in the sample. However, this assumes that allele copies within an individual are independent samples from the population. In (partially) selfing populations, gene copies are not independent samples, but individuals are. Genotype counts observed in the simulated sample of n_0 individuals can be modelled as coming from a multinomial distribution $\text{Mult}(n_0, \boldsymbol{\gamma_0})$, where $\boldsymbol{\gamma_0}$ are the genotype frequencies in the population. Similarly to Frachon *et al.* (2017), assuming the same prior probability for the three genotype frequencies, we sample genotype frequencies in the population from the posterior probability distribution with $\text{Dir}(\boldsymbol{K_0}, \boldsymbol{1})$, where $\boldsymbol{K_0}$ is the observed genotype counts in the sample of the focal locus at time $t = 0$.

Allele frequencies π_t at subsequent generations ($t \in [1, \tau]$) were simulated following a binomial distribution as $\pi_t \sim B(2\hat{N}_e, \pi_{t-1})/2\hat{N}_e$, where $\hat{N}_e$ is the genome wide estimate of the effective population size and π_0 is determined by $\boldsymbol{\gamma_0}$. Simulated genotype counts in sample at time $t = \tau$, $\boldsymbol{K_\tau^*}$, were taken from a multinomial distribution, $\boldsymbol{K_\tau^*} \sim \text{Mult}(n_\tau, \boldsymbol{\gamma_\tau})$, where n_τ is the sample size (in number of diploid individuals) at time $t = \tau$ and $\boldsymbol{\gamma_\tau}$:

$$\gamma_{\text{AA},\tau} = \pi_\tau^2 + \hat{F}_{IS}(1 - \pi_\tau)\pi_\tau$$

$$\gamma_{\text{Aa},\tau} = 2(1 - \pi_\tau)\pi_\tau(1 - \hat{F}_{IS})$$

$$\gamma_{\text{aa},\tau} = (1 - \pi_\tau)^2 + \hat{F}_{IS}(1 - \pi_\tau)\pi_\tau$$

are the genotype frequencies in the populations in function of the allele frequency π_τ and inbreeding coefficient F_{IS} (Haldane 1924), using the Weir and Cockerham's (1984) multilocus inbreeding coefficient estimate. For each simulation, we estimated the genetic differentiation F_{ST}^*. The proportion of F_{ST}^* equal or larger than the observed F_{ST}^l provides an estimate of the p-value for the test.

Preliminary results revealed that filtering loci according to minor allele frequency (MAF) was required to assure a uniform distribution of p-values from neutral sites (Fig. S1). Distribution of p-values is important for studies at the genomic scale where thousands of loci are tested (see François *et al.* 2016, for a review). In these studies false discovery rate (FDR) is estimated to control for multiple testing and this FDR estimation assumes the uniformity of p-values under the null model (Storey 2002). The criterion to filter loci was to have a minimum global MAF: loci with $\min\left(\frac{p_0 + p_\tau}{2}, 1 - \frac{p_0 + p_\tau}{2}\right) < 0.05$ were removed from the dataset. Thus, loci to be tested were chosen based on their genetic diversity, a bias that has to be taken into account in the test. This was done by discarding drift simulations that produced $\min\left(\frac{p_0^* + p_\tau^*}{2}, 1 - \frac{p_0^* + p_\tau^*}{2}\right) < 0.05$, where p_t^* is the frequency of the reference allele in the sample at time t in the simulation.

Simulations. We produced simulated data using the individual-based forward population genetics simulator SLiM 1.8 (Messer 2013). We considered a single isolated population sampled twice at the beginning and end of a time interval of τ generations. The population size $N = 500$ (number of diploid individuals) and the selfing rate σ were constant through time, and the effective population size was $N_e = \frac{(2-\sigma)N}{2}$ under neutrality. The genome was composed by two linkage groups of size 2.5×10^8 base pairs. The neutral mutation rate per base pair was $\mu = 10^{-8}$ and the recombination rate between base pairs was $r = 10^{-8}$. The recombination rate between the two linkage groups was 0.5.

In order to start our simulations with a population at mutation-drift equilibrium, SLiM was run for $20N_e$ generations. After that period of neutral evolution, two different selective scenarios were simulated: adaptation from new mutations or from standing variation. In the first case a new advantageous dominant mutation was introduced at $t = 0$ with a selection coefficient s at a random position in the genome. In the case of selection on standing variation, a random neutral polymorphic site was chosen at $t = 0$ and a selection coefficient s was assigned randomly to one of the two alleles, which also becomes dominant. The initial conditions of the simulations of selection on standing variation are, therefore, variable. The initial frequency of the allele under selection is known to affect the signal of the sweep: an advantageous allele starting at high frequency is expected to leave a lower signal than one starting at a lower frequency (Berg and Coop 2015; Innan and Kim 2004). This will likely affect the power of our test. We therefore studied the effect of the initial frequency of the advantageous allele in scenarios of adaptation from standing variation with additional simulations, where the site under selection was randomly chosen among those sites with the required allele frequency. The fre-

quency of an allele is expected to correlate with its age (Kimura and Ohta 1973), and older alleles can accumulate more mutations in their neighborhood and recombine with the background variation. That could favour the presence of a local signal of selection (Fig. S2). Because ancestral alleles are older than derived alleles, we also studied the effect of the nature of the allele in the set of simulations focusing on adaptation from standing variation.

In order to ease comparisons across scenarios, the advantageous allele was also set as dominant. Indeed, in outcrossing populations adapting from new mutation, advantageous dominant alleles are expected to spread faster than recessive ones, an effect known as Haldane's sieve. However, this effect is reduced or absent in predominantly selfing populations (Charlesworth 1992; Ronfort and Glémin 2013) and populations adapting from standing variation (Orr and Betancourt 2001).

Simulated populations were sampled at generations $t = 0$ and $t = \tau$, with samples sizes $n_0 = n_\tau = 50$ diploid individuals. Data for 10,000 polymorphic loci were taken randomly from all polymorphic sites in the sample, except for the locus under selection that was always included in the data.

Different scenarios were considered by exploring values of selfing rate $\sigma \in [0, 0.5, 0.75, 0.8, 0.85, 0.9, 0.95, 0.99, 1]$, selection coefficient $s \in [0, 0.1, 0.2, 0.3, 0.4, 0.5]$, duration of period of selection $\tau \in [5, 10, 25, 50, 100, 200]$ (in generations), type of selection (neutral, new mutation or standing variation) and, in the case of selection from standing variation, whether the locus was chosen randomly among all loci or among the loci with the required initial frequency ($\pi_0 \in [0.1, 0.5, 0.9]$) and nature (ancestral or derived) of the allele becoming advantageous. For each scenario, 100 simulation replicates were performed. Replicates in which the advantageous allele was lost were discarded and replaced by additional replicates.

Analysis of simulated data. For each simulation replicate, data were analysed as described in the two previous sections. In addition, effective population size was also estimated from F_C (following Waples 1989) for comparison. For a given scenario, the true positive rate was estimated from test results at loci under selection. The false positive rate was estimated from the test results at the neutral loci on the linkage group that does not include the locus under selection. Neutral loci on the linkage group with the locus under selection were used to characterize the footprint of selection due to hitch-hiking by estimating the positive rate as a function of the distance to the locus under selection. In order to quantify the variance of the selection signal, loci were bootstrapped and the 95% quantile interval for the proportion of positive tests was calculated. Positive tests were defined with an arbitrary threshold of

Fig. 1. Effective population size estimates from temporal differentiation. Estimates of N_e from F_{ST} were obtained from simulated data of populations of $N = 500$ diploid individuals, sampled twice with $\tau = 25$ generations between samples and selfing rate σ. **(a)** Neutral selfing population. **(b)** Outcrossing population under selection: a new advantageous mutation appears at generation $t = 0$ with coefficient of selection s. Samples are made of 50 diploid individuals genotyped at 10,000 biallelic markers (including the locus under selection), but only loci with a global MAF over 0.05 are used in the estimate (see details in Materials and Methods). Boxplot for estimates from 100 simulation replicates. Dashed line marks true effective population size, N_e, in panel **(a)** and census size, N, in panel **(b)**.

p-value<0.001. In order to control for multiple testing, FDR estimates were quantified with q-values for each locus, following Storey (2002) with the *qvalue* R package (Storey *et al.* 2019). Manhattan and QQ plots were generated with *qqman* R package (Turner 2017).

As explained in the previous section, the genetic diversity around the selected locus could influence the outlier test in a scenario of adaptation from standing variation in a predominantly sefing species (Fig. S2). In such scenarios, we therefore examined the structure of genetic diversity at the beginning of the sweep. This allowed us to characterize the effects of historical recombination previous to the outset of the selective sweep. For each simulated population, the haplotypes, defined as the whole linkage group under selection, were classified in two groups: one group for haplotypes carrying the advantageous allele and one group for haplotypes with the neutral allele. Genetic diversity (measured as the average expected heterozygosity per bp) within the haplotypes carrying the advantageous allele and differ-

Fig. 2. Power and false positive rate of the genome scan for increasing decision thresholds. Receiver operating characteristic curve is estimated from 100 replicates of simulated data of one population of $N = 500$ diploid individuals, sampled twice with $\tau = 25$ generations between samples, selection coefficient $s = 0.5$ and selfing rate σ. **(a)** Selection on new mutation. **(b)** Selection on standing variation. Circles mark the values for positive threshold of p-value>0.001, which is the threshold used in Fig. 3.

entiation between the two groups of haplotypes (measured as the F_{ST}) were calculated on 3,000 bp windows at increasing distance from the locus under selection.

Real data application. *Medicago truncatula* is an annual, predominantly selfing species (Siol *et al.* 2008) of the legume family (Fabaceae), found around the Mediterranean Basin. We conducted a temporal survey in a population located in Cape Corsica (42°58.406' North, 9°22,015' East, 362 m.a.s.l.). Samples of around 100 pods were collected in 1987 and 2009 along three transects running across the population, with at least one meter distance between each pod collected in order

to avoid over-sampling the progeny of a single individual. Using these samples, we collected leaf material from 64 plants from 1987 and 96 plants from 2009 grown in greenhouse for DNA extraction. Between 100 mg and 200 mg of young leaves were ground in liquid nitrogen in a Qiagen Retsch Tissue Lyser (Qiagen N. V., Hilden, Germany) during 2×1 min at 30 Hz. The fine powder was mixed with 600 µl of pre-heated lysis buffer consisting of 100 mM Tris-HCl pH 8.0, 20 mM EDTA pH 8.0, 1.4 M NaCl, 2% (w/v) CTAB, 1% (w/v) PVP40 and 1% (w/v) sodium bisulphite plus 8 µl of 10 mg/ml RNAse per sample added extemporaneously. After incubation for 20 min at 65°C under medium shaking, 600 µl of chloroform were added and mixed with a vortex mixer. Each sample was centrifuged for 15 min at 10,000 g and 10°C and the upper phase was transferred into a new tube and then mixed with 60 µl of 3M sodium acetate and 600 µl of cold isopropanol. DNA was precipitated by another centrifugation (30 min, 15,000 g, 4°C), rinsed with 300 µl of 70% cold ethanol, centrifuged again (10 min, 15,000 g, 4°C), dried for 10 min at room temperature and resuspended in 100 µl of sterile deionized water. The genotyping was performed using two SNP chips specifically developed for *M. truncatula* (Loridon *et al.* 2013). Out of the 1920 SNPs, 137 were located in genes encoding flowering time, 721 in other candidate genes, in particular genes involved in symbiosis and 1062 at random positions of the genome. The 1920 SNPs were widespread across all eight linkage groups (Table S2). The genotyping assays were achieved at GenoToul (Genomic Platform in Toulouse, France) and at the BioMedical Genomics Center (Minneapolis, University of Minnesota, USA) using GoldenGate Assay (Fan *et al.* 2006, 2003) and respectively Illumina's VeraCode technology (Lin *et al.* 2009) or Bead Array technology. Data generated from the BeadXpressTM reader (384 SNP×480 DNA) or BeadArray Reader (1536 SNP×192 DNA) were analyzed as detailed in Loridon *et al.* (2013).

These data were analysed using the temporal genome scan described in this paper, providing estimates of the effective population size (from F_{ST}) and the selfing rate (from F_{IS}) of the population. For loci with a global MAF higher than 0.05, we obtained a p-value for the test of homogeneity. In order to control for multiple testing, FDR estimates were quantified with q-values for each locus, following Storey (2002).

Results

Accuracy of N_e estimates. In neutral scenarios, estimates of effective population size derived from F_{ST} performed reasonably well, decreasing in the presence of partial selfing following the N_e theoretical expectation (Fig. 1a). Selfing also caused an increase of error in the estimation of N_e. In the presence of selection, estimates of effective population size decreased with the strength of the selective coefficient of the causal muta-

Fig. 3. Selection footprint on the selected chromosome under different scenarios. Proportion of positive tests in function of distance to the locus under selection. Estimates obtained from 100 replicates of simulated data of one population of $N = 500$ diploid individuals, sampled twice with τ generations between samples and selection coefficient s and selfing rate σ. **(a)** Adaptation from new mutation, $\tau = 25$, $s = 0.5$. **(b)** Adaptation from standing variation, $\tau = 25$, $s = 0.5$. **(c)** Adaptation from new mutation, $\tau = 25$, $\sigma = 0$. **(d)** Adaptation from new mutation, $s = 0.5$, $\sigma = 0$. Distance measured in centimorgans (cM). Coloured areas mark the 95% bootstrap interval.

tion (Fig. 1b). As expected, effective population size estimates from F_C showed a clear bias in the presence of partial self-fertilization while the bias was negligible in estimates from F_{ST} (Fig. S3a).

Footprint of selection. The temporal genome scan discriminated well between selected and neutral (on a separate linkage group) sites in most scenarios (Fig. 2). Yet, in the case of selection on new mutations, extreme selfing rate decreased the performance of the test (Fig. 2a). In scenarios of selection on standing variation the performance of the test was lower (Fig. 2b). In addition, the effect of selfing was more complex in the case of selection on standing variation. Moderate levels of selfing seemed to improve the discrimination capacity while the performance for high levels of selfing was similar to the one for outcrossing simulations. Only complete selfing reduced dramatically the discrimination capacity of the method.

These results, however, only consider the causal polymorphisms and completely independent neutral variants, ignoring linked sites that could have been subject to hitch-hiking. In practice, the signal for the de-

tection of regions under selection comes mainly from those linked sites, that can create a local excess of outlier loci testing positive around the site under selection (even if the later is not in the data). We therefore examined the footprint of selection at increasing distance from the advantageous allele, within a linkage group. As expected, the highest probability of positive test was at the locus under selection; then probability decreased with distance and reached very low values, similar to those for neutral loci on a separate linkage group (Fig. 3). The distance at which the hitch-hiking effect disappears depends on the scenario, being smaller for selection on scenarios of standing variation and scenarios with high selfing rate, and larger for scenarios on new mutation and scenarios of intermediate selfing rate. Under a scenario of adaptation from new mutations, the overall strength of the signal decreased with increasing selfing rate (Fig. 3a) and decreasing the selection coefficient (Fig. 3c). The time sample interval also influenced the strength of the signal, with intermediate values being more favourable for the detection of outlier loci (Fig. 3d).

The results for the signal of hitch-hiking mirror those

of the power to detect the causal site. In the case of selection on standing variation, there was an overall reduction of the signal compared to scenarios of selection on new mutation (Fig. 3a,b) except for highly selfing populations ($\sigma \geq 0.95$). Nevertheless, as in the scenario of adaptation from new mutations, predominantly selfing populations had a weaker signal of selection than outcrossing populations. However, the strength of the signal for population with intermediate levels of selfing ($\sigma = 0.5$, $\sigma = 0.8$) was similar or even higher than outcrossing populations (Fig. 3b).

The initial frequency of the advantageous allele and whether it is ancestral or derived also influenced the strength of the signal in predominantly selfing populations ($\sigma = 0.95$, Fig. 4). As expected, the lower the initial frequency, the stronger was the signal of selection, but this was modulated by the nature of the allele. With low initial frequencies, selection on the ancestral allele left a stronger signal. Symmetrically, the signal was stronger for derived alleles if the initial frequency was high. To understand these results, we examined the genetic diversity and the differentiation between the group of haplotypes carrying the advantageous mutation and the group not carrying it at the outset of the sweep. We found that this genetic differentiation decreases with the distance from the selected site, with higher differentiation for scenarios with stronger signal of selection (i.e. when the advantageous allele is ancestral with low initial frequency). Besides, ancestral alleles were associated to more diverse local haplotypes than derived alleles at the same frequency. These results show that not further recombination during the sweep is required to generate a local signal of selection.

Medicago truncatula. From the 1920 SNPs genotyped, 1224 were polymorphic in the sample and 987 had a global MAF higher than 0.05. Effective population size was estimated at $\hat{N}_e = 42$ (from $\hat{F}_{ST} = 0.207$) and selfing rate at $\hat{\sigma} = 0.986$ (from $\hat{F}_{IS} = 0.972$). The test of homogeneity did not identify any SNP as a strong candidate for being under selection: the lowest p-value was 0.04 with a corresponding q-value of 0.48 (Fig. 5 and Table S2).

Discussion

Estimating effective population size under selfing and selection. It is well known that self-fertilization reduces the effective size of populations. Our results show that Weir and Cockerham's (1984) F_{ST} allows to measure the amount of drift between two temporal samples, even in the presence of selfing. However, the precision of the estimates diminishes with the rate of selfing. We hypothesize that it is the consequence of the effect of reduced effective recombination in selfing, which reduces the number of loci with independent evolutionary histories. This effect is most extreme for completely selfing populations, where the whole genome behaves as a single locus, reducing dramatically the information available in the data. In order to test this hypothesis, we estimated N_e for a similar set of simulations under an unrealistic model where each locus was simulated independently (supplementary text S2). On those simulations, precision of N_e estimates was no longer reduced by selfing (Fig. S3b).

An alternative and common way to estimate N_e from temporal data uses the standardized variance in allele frequencies (F_C, Nei and Tajima 1981). This was the approach originally proposed for the temporal genome scan by Goldringer and Bataillon (2004). However, estimates from F_C suffer from a bias (Fig. S3) because this approach assumes that the $2n$ gene copies (in a sample of n diploid individuals) are independent draws from the population gene pool, and uses a binomial distribution to model it (Waples 1989). However, in the case of partially or predominantly selfing populations, the two gene copies within an individual are not independent samples from the population, which explains the bias of the estimate. Using Weir and Cockerham's (1984) F_{ST} estimate corrects for this bias.

Strong selection also reduces the effective population size by reducing the number of breeding individuals or increasing the variance of reproductive success among individuals (Robertson 1961; Santiago and Caballero 1995). This is reflected in the estimation of N_e, which decreases with the strength of the selective coefficient of the mutation under selection in our simulations (Fig. 1b). Note that, in our case, this effect is not only driven by the local increase of temporal genetic differentiation around the site under selection. The strong selection considered in our simulations increases F_{ST} genome-wide, even in regions unlinked to the site under selection (e.g. distance of about 50 centimorgan or larger; Fig. S5). The combined effect of selfing (i.e. increased linkage disequilibrium) and selection produced very strong drift compared to a neutrally evolving population with the same census size (Fig. S5).

The effective population size estimated in Cape Corsica *M. truncatula* was extremely low, in agreement with the results from microsatellites data on the same population ($\hat{N}_e = 20$, Jullien *et al.* 2019). However, such small N_e estimates might reflect other processes in addition to drift (e.g. immigration, Jullien *et al.* 2019). Indeed, the distribution of p-values from the genome scan shows a departure from the expected distribution which could indicate an inappropriate null model (Fig. S7a). This deficit of low p-values means that the outlier test is too conservative because the null distribution of temporal F_{ST} is wider than the distribution observed genomewide. The QQ plots for simulated populations show that, as the selfing rate increases, the test progressively shifts from delivering and excess of low p-values to delivering a deficit of low p-values (Fig. S7b), as observed for *M. truncatula*. This implies that the test is not appropriate for such high

Fig. 4. Influence of derived/ancestral state and initial frequency on selection footprint. Proportion of positive test in function of distance to the locus under selection. Estimates obtained from 100 replicates of simulated data of one population of $N = 500$ diploid individuals, sampled twice with $\tau = 25$ generations between samples and selection coefficient $s = 0.5$. Adaptation on standing variation, selfing rate $\sigma = 0.95$.

Fig. 5. Genome-scan for selection based on temporal differentiation in cape Corsica *Medicago truncatula* population. $-\log_{10}(p\text{-value})$ of the simulation-based test of the null hypothesis that the locus-specific differentiation measured at each SNP is only due to genetic drift. Only SNP markers with MAF>0.05 and known position at the reference genome are shown. Red line marks p-value=0.001, the significance threshold used for figures 3 and 4.

levels of selfing ($\sigma > 0.99$), where the whole genome evolves as a single locus, but the method can perform well with slightly lower selfing rates ($\sigma = 0.95$). In addition, the assumption that temporal data collected from the same geographical location belong to a single continuous population is likely to be wrong in many cases, even more so for predominantly selfing populations, where the spatial structure is usually very strong **?**. Jullien (2019) proposes to address this issue by performing demographic inference using approximate Bayesian computation (Csilléry *et al.* 2010) that can potentially discriminate isolated from admixed populations, and jointly estimate drift, migration and selfing parameters.

In our simulations we have considered that individual genotypes are available. However, in many temporal experiments, sequencing is performed for pools of individuals (Pool-seq; e.g. Franks *et al.* 2016) to reduce the costs. Estimating effective population sizes remains possible, using Pool-seq adapted F_{ST} and F_C estimates (Hivert *et al.* 2018; Jónás *et al.* 2016). However, since individual information is lost, these estimates cannot take into account the within individual allele identity correlation due to selfing. When studying a selfing species, we therefore recommend working with individuals genotypes. If Pool-seq is necessary due to budget limitations, we advise to follow a similar procedure as in Frachon *et al.* (2017) and reproduce seeds by selfing for one or more generations so that sequenced individuals are completely homozygous and can be treated as effectively haploid samples. However, this approach is likely to only give good results with predominantly selfing species (with a small proportion of heterozygous loci to be removed) and divides the amount of data by two.

Limits imposed by selfing to temporal genome scans of selection.

Our results show that the search of regions under selection with temporal F_{ST} genome scans in predominantley selfing populations can be challenging. In the case of complete selfing the whole genome behaves as single locus and selection, if present, affects the whole genome. In partially selfing populations, the strength of the signal left by selection depends on whether the advantageous allele had the time to recombine with different genetic backgrounds. In a population with a selfing rate as high as $\sigma = 0.95$, outcrossing events are unlikely to occur during the sweep of a new beneficial mutation and produce no effective recombination if they happen between close relatives sharing the same homozygous genotype. As a results, like completely selfing populations, the selective sweep will reduce genetic diversity on the whole genome and will leave no local signal of selection.

On the contrary, when adaptation proceeds through standing variation, historical recombination can put an allele on different genetic backgrounds before it becomes advantageous and sweeps. In that case, the sweeping haplotypes are only similar among them and different from the haplotypes in the rest of the population around the site under selection. As the distance to the selected site increases, the diversity of the sweeping haplotypes raises and the differentiation with the other haplotypes segregating in the population decreases due to the action of historical recombination (Fig. S4). These differences along the genome create the local signal of selection, even if no further recombination occurs during the sweep.

Selection on standing variation is usually associated to a weak signal of selection (i.e. a soft sweep, Hermisson and Pennings 2005). Our results indeed confirm this expectation for outcrossing populations (compare Fig. 3a,b for $\sigma = 0$). Interestingly, for predominantly selfing populations, the situation is reversed. Yet, the

strength of the local signal of selection, and therefore the power of the outlier test depends not only on the initial frequency of the advantageous allele, but also on its genetic background. Consider an advantageous derived allele at low frequency. Because its low frequency, it is likely a young allele (Fig. S6) and few mutations will have accumulated around it. The selected haplotypes will therefore display low diversity and little differentiation from the deleterious haplotypes (Fig. S4). Thus, we expect that only few mutations will show a significant change in allele frequencies around the site under selection (e.g. Fig. S2a). On the other hand, if the ancestral allele (at low frequency) becomes advantageous, it will be likely to be on a very old lineage (since a high frequency derived allele is expected to be old). With more time for mutations to accumulate on both lineages, creating diversity that is unique to the haplotypes under selection. This will lead to more sites with a significant allele frequency change around the causal mutation (e.g. Fig. S2b). In the case of selection on an allele starting at high frequency, the situation is reversed. The lowest signal is for selection on high frequency ancestral alleles that not only will have a small change on allele frequency but also carry diversity common to the whole population. Finally, low frequency ancestral alleles might have the potential to create the strongest signal, but they are scarce in an equilibrium population (Fig. S6) so they might not be the most frequent case of sweep from standing variation.

Although temporal genome scans in predominantly selfing species are able to reveal the footprint of selection on standing variation, how prevalent is adaptation from pre-existing variation in selfing populations remains an open question. In the short-term, selection is most likely acting on standing variation (Barrett and Schluter 2008), but there are not specific predictions for selfing populations. Theoretical models predict an overall limitation of adaptation in selfers compared to outcrossers (Hartfield and Glémin 2016). However, selfers can hold high levels of cryptic genetic variation for polygenic traits, due to negative linkage disequilibrium. This diversity may allow a selfing population adapt to changing conditions as quickly as outcrossing populations (Clo *et al.* 2019). Indeed, there is some evidence of selective sweeps in selfing species (Bonhomme *et al.* 2015; Huber *et al.* 2014). In agreement with this, the temporal genome scan in a predominantly selfing population of *A. thaliana* showed footprints of selection (Frachon *et al.* 2017). Such positive results could have been favoured by the relatively large genetic variation of this population (Baron *et al.* 2015; Frachon *et al.* 2017). Based on the simulation results we present here, we can assume that this population has been adapting from standing variation over a short period of time (eight generations) rather than from many new mutations occurring during (or shortly before) the studied period.

The absence of evidence for selection in the *M. truncat-*

ula dataset is manifest. It is reasonable to attribute this results to the extreme selfing rate estimate ($\hat{\sigma} = 0.986$) in this population, for which our simulation approach gives little hope to detect any signal of selection. Indeed, the same selfing lineages are observed throughout the study period of 22 generations (Jullien *et al.* 2019), which further suggest that almost no recombination occurs in the population. If selection, either on new or pre-existing mutations, has occurred, it has changed the frequencies of multilocus genotypes without leaving different any local signal along the genome. This would have reduced dramatically the effective population size, which is consistent with our extremely low estimate $\hat{N}_e = 42$. However, the present analysis cannot be conclusive on any of these hypotheses and the observed diversity changes might be better explained by a demographic model including genetic exchanges with neighbouring populations, as discussed by Jullien *et al.* (2019).

Conclusions. Identifying regions under selection with a temporal genome scan can fail for several reasons. First, timing is paramount. A sample at the beginning and at the end of the selective sweep would be ideal for detection of selection. However, the start and duration of the sweep and the sampling times cannot be synchronized except for some experimental evolution studies. Frequency of the advantageous allele at the beginning of the sweep can also reduce the chances of capturing any signal. On top of that, selfing presents itself as a strong additional difficulty for this task, reducing the efficacy of selection (reduced N_e) and extending hitchhiking effects that blur the distinction between neutral and selected regions. Nevertheless, scenarios of adaptation from standing variation can leave some signal of selection in highly selfing populations. Curiously, adaptation from new mutation under the same highly selfing scenarios leave no signal, reversing the general expectation for the footprint of selection from "hard" and "soft" sweeps. Chance plays an important role in the success of a genome scan. Therefore, researchers should focus on the factors that can be controlled, such as the use of adapted methods to selfing species, to increase the probability of a positive outcome.

ADDITIONAL INFORMATION

Author Contributions. Miguel Navascués, Laurène Gay, Joëlle Ronfort and Renaud Vitalis conceived and designed the research. MN and RV developed the temporal F_{ST} scan for partially selfing populations and the code implementing it. Arnaud Becheler and MN evaluated the performance of the method. Karine Loridon, LG and JR produced the *Medicago truncatula* data set. MN wrote the article with the help of all authors that critically reviewed and approved the text.

Competing Interests. The authors declare there are no competing interests.

Data Availability. The method described in this work is implemented in DriftTest, a C program available at Zenodo (Navascués and Vitalis 2018). An R script (R Core Team 2018) to reproduce the simulations and analyses in this work is available at Zenodo (Becheler *et al.* 2018).

Funding. This research was developed under the SelfAdapt project, funded by INRA metaprogramme "Adaptation of Agriculture and Forests to Climate Change" (ACCAF).

ACKNOWLEDGEMENTS
A large part of the analyses presented in this work was performed on the CBGP HPC computational platform; we thank the platform manager, Alexandre Dehne Garcia, for being ever-helpful.

References

Anderson, E. C., E. G. Williamson, and E. A. Thompson, 2000 Monte Carlo evaluation of the likelihood for N_e from temporally spaced samples. Genetics **156**: 2109–2118.

Baron, E., J. Richirt, R. Villoutreix, L. Amsellem, and F. Roux, 2015 The genetics of intra- and interspecific competitive response and effect in a local population of an annual plant species. Functional Ecology **29**: 1361–1370. doi: 10.1111/1365-2435.12436.

Barrett, R. D. H. and D. Schluter, 2008 Adaptation from standing genetic variation. Trends in Ecology & Evolution **23**: 38–44. doi: 10.1016/j.tree.2007.09.008.

Becheler, A., R. Vitalis, and M. Navascués, 2018 DriftTest-Evaluation. A suite of R scripts to evaluate the performance of DriftTest for detection of selection with temporal data. Zenodo doi: 10.5281/zenodo.1194667.

Berg, J. J. and G. Coop, 2015 A coalescent model for a sweep of a unique standing variant. Genetics **201**: 707–725. doi: 10.1534/genetics.115.178962.

Biek, R., O. G. Pybus, J. O. Lloyd-Smith, and X. Didelot, 2015 Measurably evolving pathogens in the genomic era. Trends in Ecology & Evolution **30**: 306–313. doi: 10.1016/j.tree.2015.03.009.

Bollback, J. P., T. L. York, and R. Nielsen, 2008 Estimation of $2N_es$ from temporal allele frequency data. Genetics **179**: 497–502. doi: 10.1534/genetics.107.085019.

Bonhomme, M., S. Boitard, H. San Clemente, B. Dumas, N. Young, *et al.*, 2015 Genomic signature of selective sweeps illuminates adaptation of *Medicago truncatula* to root-associated microorganisms. Molecular Biology and Evolution **32**: 2097–2110. doi: 10.1093/molbev/msv092.

Buffalo, V. and G. Coop, 2019 The linked selection signature of rapid adaptation in temporal genomic data. Genetics **213**: 1007–1045. doi: 10.1534/genetics.119.302581.

Burgarella, C., P. Gayral, M. Ballenghien, A. Bernard, P. David, *et al.*, 2015 Molecular evolution of freshwater snails with contrasting mating systems. Molecular Biology and Evolution **32**: 2403–2416. doi: 10.1093/molbev/msv121.

Charlesworth, B., 1992 Evolutionary rates in partially self-fertilizing species. American Naturalist pp. 126–148. doi: 10.1086/285406.

Clo, J., J. Ronfort, and D. A. Awad, 2019 Hide and seek: hidden genetic variance contributing to the adaptive potential of selfing populations. bioRxiv p. 810515. doi: 10.1101/810515.

Csilléry, K., M. G. Blum, O. E. Gaggiotti, and O. François, 2010 Approximate Bayesian computation (ABC) in practice. Trends in Ecology & Evolution **25**: 410–418. doi: 10.1016/j.tree.2010.04.001.

Diver, C., 1929 Fossil records of Mendelian mutants. Nature **124**: 183. doi: 10.1038/124183a0.

Dobzhansky, T., 1943 Genetics of natural populations IX. Temporal changes in the composition of populations of *Drosophila pseudoobscura*. Genetics **28**: 162–186.

Drummond, A. J., G. K. Nicholls, A. G. Rodrigo, and W. Solomon, 2002 Estimating mutation parameters, population history and genealogy simultaneously from temporally spaced sequence data. Genetics **161**: 1307–1320.

Etterson, J. R., S. J. Franks, S. J. Mazer, R. G. Shaw, N. L. S. Gorden, *et al.*, 2016 Project Baseline: An unprecedented resource to study plant evolution across space and time. American Journal of Botany **103**: 164–173. doi: 10.3732/ajb.1500313.

Fan, J., K. L. Gunderson, M. Bibikova, J. M. Yeakley, J. Chen, *et al.*, 2006 Illumina universal bead arrays. In *Methods in Enzymology*, volume 410 of *DNA Microarrays, Part A: Array Platforms and Wet-Bench Protocols*, pp. 57–73, Academic Press. doi: 10.1016/S0076-6879(06)10003-8.

Fan, J.-B., A. Oliphant, R. Shen, B. G. Kermani, F. Garcia, *et al.*, 2003 Highly parallel SNP genotyping. Cold Spring Harbor Symposia on Quantitative Biology **68**: 69–78. doi: 10.1101/sqb.2003.68.69.

Feder, A. F., S. Kryazhimskiy, and J. B. Plotkin, 2014 Identifying signatures of selection in genetic time series. Genetics **196**: 509–522. doi: 10.1534/genetics.113.158220.

Fisher, R. A. and E. B. Ford, 1947 The spread of a gene in natural conditions in a colony of the moth *Panaxia dominula* L. Heredity **1**: 143–174. doi: 10.1038/hdy.1947.11.

Foll, M., H. Shim, and J. D. Jensen, 2014 WFABC: a Wright-Fisher ABC-based approach for inferring effective population sizes and selection coefficients from time-sampled data. Molecular Ecology Resources **15**: 87–98. doi: 10.1111/1755-0998.12280.

Frachon, L., C. Libourel, R. Villoutreix, S. Carrere, C. Glorieux, *et al.*, 2017 Intermediate degrees of synergistic pleiotropy drive adaptive evolution in ecological time. Nature Ecology & Evolution **1**: 1551–1561. doi: 10.1038/s41559-017-0297-1.

Franks, S. J., N. C. Kane, N. B. O'Hara, S. Tittes, and J. S. Rest, 2016 Rapid genome-wide evolution in *Brassica rapa* populations following drought revealed by sequencing of ancestral and descendant gene pools. Molecular Ecology **25**: 3622–3631. doi: 10.1111/mec.13615.

François, O., H. Martins, K. Caye, and S. D. Schoville, 2016 Controlling false discoveries in genome scans for selection. Molecular Ecology **25**: 454–469. doi: 10.1111/mec.13513.

Golding, G. B. and C. Strobeck, 1980 Linkage disequilibrium in a finite population that is partially selfing. Genetics **94**: 777–789.

Goldringer, I. and T. Bataillon, 2004 On the distribution of temporal variations in allele frequency consequences for the estimation of effective population size and the detection of loci undergoing selection. Genetics **168**: 563–568. doi: 10.1534/genetics.103.025908.

Goodwillie, C. and E. Stewart, 2013 Cleistogamy and hybridization in two subspecies of *Triodanis perfoliata* (Campanulaceae). Rhodora **115**: 42–60. doi: 10.3119/12-01.

Haldane, J. B. S., 1924 A mathematical theory of natural and artificial selection. Part II. The influence of partial self-fertilisation, inbreeding, assortative mating, and selective fertilisation on the composition of mendelian populations, and on natural selection. Biological Reviews **1**: 158–163. doi: 10.1111/j.1469-185X.1924.tb00546.x.

Hartfield, M., T. Bataillon, and S. Glémin, 2017 The evolutionary interplay between adaptation and self-fertilization. Trends in Genetics **33**: 420–431. doi: 10.1016/j.tig.2017.04.002.

Hartfield, M. and S. Glémin, 2016 Limits to adaptation in partially selfing species. Genetics **203**: 959–974. doi: 10.1534/genetics.116.188821.

Hermisson, J. and P. S. Pennings, 2005 Soft sweeps: Molecular population genetics of adaptation from standing genetic variation. Genetics **169**: 2335–2352. doi: 10.1534/genetics.104.036947.

Hivert, V., R. Leblois, E. J. Petit, M. Gautier, and R. Vitalis, 2018 Measuring genetic differentiation from pool-seq data. Genetics **210**: 315–330. doi: 10.1534/genetics.118.300900.

Huber, C. D., M. Nordborg, J. Hermisson, and I. Hellmann, 2014 Keeping it local: Evidence for positive selection in Swedish *Arabidopsis thaliana*. Molecular Biology and Evolution **31**: 3026–3039. doi: 10.1093/molbev/msu247.

Innan, H. and Y. Kim, 2004 Pattern of polymorphism after strong artificial selection in a domestication event. Proceedings of the National Academy of Sciences **101**: 10667–10672. doi: 10.1073/pnas.0401720101.

Jullien, M., 2019 *Analyse temporelle de la diversité en régime autogame : Approches théorique et empirique*. PhD, Montpellier SupAgro, Montpellier.

Jullien, M., M. Navascués, J. Ronfort, K. Loridon, and L. Gay, 2019 Structure of multilocus genetic diversity in predominantly selfing populations. Heredity. doi: 10.1038/s41437-019-0182-6.

Jónás, Á., T. Taus, C. Kosiol, C. Schlötterer, and A. Futschik, 2016 Estimating the effective population size from temporal allele frequency changes in experimental evolution. Genetics **204**: 723–735. doi: 10.1534/genetics.116.191197.

Kellogg, V. L. and R. G. Bell, 1904 Studies of variation in insects. Proceedings of the Washington Academy of Sciences **6**: 203–332.

Kimura, M. and T. Ohta, 1973 The age of a neutral mutant persisting in a finite population. Genetics **75**: 199–212.

Kon, K. F. and W. M. Blacklow, 1990 Polymorphism, outcrossing and polyploidy in *Bromus diandrus* and *B. rigidus*. Australian Journal of Botany **38**: 609–618. doi: 10.1071/bt9900609.

Krimbas, C. B. and S. Tsakas, 1971 The genetics of *Dacus oleae*. V. Changes of esterase polymorphism in a natural population following insecticide control—selection or drift? Evolution **25**: 454–460. doi: 10.1111/j.1558-5646.1971.tb01904.x.

Larson, S. R., E. Cartier, C. L. McCracken, and D. Dyer, 2001 Mode of reproduction and amplified fragment length polymorphism variation in purple needlegrass (*Nassella pulchra*): utilization of natural germplasm sources. Molecular Ecology **10**: 1165–1177. doi: 10.1046/j.1365-294X.2001.01267.x.

Leonardi, M., P. Librado, C. Der Sarkissian, M. Schubert, A. H. Alfarhan, *et al.*, 2017 Evolutionary patterns and processes: Lessons from ancient DNA. Systematic Biology **66**: e1–e29. doi: 10.1093/sysbio/syw059.

Lin, C. H., J. M. Yeakley, T. K. McDaniel, and R. Shen, 2009 Medium- to high-throughput SNP genotyping using VeraCode microbeads. In *DNA and RNA Profiling in Human Blood: Methods and Protocols*, edited by P. Bugert, Methods in Molecular Biology, pp. 129–142, Humana Press, Totowa, NJ. doi: 10.1007/978-1-59745-553-4_10.

Loridon, K., C. Burgarella, N. Chantret, F. Martins, J. Gouzy, *et al.*, 2013 Single-nucleotide polymorphism discovery and diversity in the model legume *Medicago truncatula*. Molecular Ecology Resources **13**: 84–95. doi: 10.1111/1755-0998.12021.

Messer, P. W., 2013 SLiM: Simulating evolution with selection and linkage. Genetics **194**: 1037–1039. doi: 10.1534/genetics.113.152181.

Navascués, M. and R. Vitalis, 2018 DriftTest v1.0.4. a computer program to detect selection from temporal genetic differentiation. Zenodo doi: 10.5281/zenodo.1194663.

Nei, M. and F. Tajima, 1981 Genetic drift and estimation of effective population size. Genetics **98**: 625–640.

Novak, S. J., R. N. Mack, and D. E. Soltis, 1991 Genetic variation in *Bromus tectorum* (Poaceae): Population differentiation in its North American range. American Journal of Botany **78**: 1150–1161. doi: 10.2307/2444902.

Orr, H. A. and A. J. Betancourt, 2001 Haldane's sieve and adaptation from the standing genetic variation. Genetics **157**: 875–884.

Pool, J. E., I. Hellmann, J. D. Jensen, and R. Nielsen, 2010 Population genetic inference from genomic sequence variation. Genome Research **20**: 291–300. doi: 10.1101/gr.079509.108.

R Core Team, 2018 *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.

Rambaut, A., 2000 Estimating the rate of molecular evolution: incorporating non-contemporaneous sequences into maximum likelihood phylogenies. Bioinformatics **16**: 395–399. doi: 10.1093/bioinformatics/16.4.395.

Robertson, A., 1961 Inbreeding in artificial selection programmes. Genetics Research **2**: 189–194. doi: 10.1017/S0016672300000690.

Ronfort, J. and S. Glémin, 2013 Mating system, Haldane's sieve, and the domestication process. Evolution **67**: 1518–1526. doi: 10.1111/evo.12025.

Sanders, T. B. and J. L. Hamrick, 1980 Variation in the breeding system of *Elymus canadensis*. Evolution **34**: 117–122. doi: 10.1111/j.1558-5646.1980.tb04794.x.

Santiago, E. and A. Caballero, 1995 Effective size of populations under selection. Genetics **139**: 1013–1030.

Schemske, D. W., 1978 Evolution of reproductive characteristics in *Impatiens* (Balsaminaceae): The significance of cleistogamy and chasmogamy. Ecology **59**: 596–613. doi: 10.2307/1936588.

Schlötterer, C., R. Kofler, E. Versace, R. Tobler, and S. U. Franssen, 2015 Combining experimental evolution with next-generation sequencing: a powerful tool to study adaptation from standing genetic variation. Heredity **114**: 431–440. doi: 10.1038/hdy.2014.86.

Siol, M., J. M. Prosperi, I. Bonnin, and J. Ronfort, 2008 How multilocus genotypic pattern helps to understand the history of selfing populations: a case study in *Medicago truncatula*. Heredity **100**: 517–525. doi: 10.1038/hdy.2008.5.

Skoglund, P., P. Sjödin, T. Skoglund, M. Lascoux, and M. Jakobsson, 2014 Investigating population history using temporal genetic differentiation. Molecular Biology and Evolution **31**: 2516–2527. doi: 10.1093/molbev/msu192.

Storey, J. D., 2002 A direct approach to false discovery rates. Journal of the Royal Statistical Society: Series B (Statistical Methodology) **64**: 479–498. doi: 10.1111/1467-9868.00346.

Storey, J. D., A. J. Bass, A. Dabney, and D. Robinson, 2019 *qvalue: Q-value estimation for false discovery rate control*. doi: 10.18129/B9.bioc.qvalue.

Turner, S., 2017 *qqman: Q-Q and Manhattan plots for GWAS data*. R package version 0.1.4.

Vitalis, R. and D. Couvet, 2001 Two-locus identity probabilities and identity disequilibrium in a partially selfing subdivided population. Genetical research **77**: 67–81.

Wang, J. and M. C. Whitlock, 2003 Estimating effective population size and migration rates from genetic samples over space and time. Genetics **163**: 429–446.

Waples, R. S., 1989 A generalized approach for estimating effective population size from temporal changes in allele frequency. Genetics **121**: 379–391.

Weir, B. S. and C. C. Cockerham, 1984 Estimating F-statistics for the analysis of population structure. Evolution **38**: 1358–1370. doi: 10.2307/2408641.

Whitehead, M. R., R. Lanfear, R. J. Mitchell, and J. D. Karron, 2018 Plant mating systems often vary widely among populations. Frontiers in Ecology and Evolution **6**: 38. doi: 10.3389/fevo.2018.00038.

Williamson, E. G. and M. Slatkin, 1999 Using maximum likelihood to estimate population size from temporal changes in allele frequencies. Genetics **152**: 755–761.

Wright, S., 1948 On the roles of directed and random changes in gene frequency in the genetics of populations. Evolution **2**: 279–294. doi: 10.1111/j.1558-5646.1948.tb02746.x.

Yamazaki, T., 1971 Measurement of fitness at the esterase-5 locus in *Drosophila pseudoobscura*. Genetics **67**: 579–603.

Supplementary Information

S1: Predominantly selfing species in project Baseline

Project Baseline will track the evolution of natural populations from more than 60 plant species in the next 50 years through seed collection (www.baselineseedbank.org, Etterson *et al.* 2016). Many of the species included are capable of self-fertilization and at least six of them reproduce predominantly by selfing: *Bromus diandrus* (Kon and Blacklow 1990), *Bromus tectorum* (Novak *et al.* 1991), *Elymus canadensis* (Sanders and Hamrick 1980), *Impatiens pallida* (Schemske 1978), *Stipa pulchra* (Larson *et al.* 2001) and *Triodanis biflora* (Goodwillie and Stewart 2013).

S2: Simulation of independent loci

Additional simulations were performed to show the effect in the estimation of N_e of non-independent sample of gene copies within individuals (i.e. F_{IS}) without the effect of linkage disequilibrium due to selfing. The simulation approach is very similar to the simple drift model used to build null distribution of the test described in the main text. The main difference is that initial allele frequency is set as a fixed parameter. Simulations are used to generate temporal data at 10,000 biallelic loci per pseudo-observed dataset. Initial allele frequencies (π_0) are fixed to 0.5 for all loci. Allele frequency π in generation t were simulated with a binomial distribution as $\pi_t \sim B(2N_e, \pi_{t-1})/2N_e$, where N_e is the effective population size in number diploid individuals, for generations $t \in [1, \tau]$. Thus, each locus is simulated independently from each other, ignoring the linkage among them that should have occurred due to selfing. Genotype counts in samples, K_t^*, at time $t = 0$ and $t = \tau$ are taken from a multinomial distribution, $K_t^* \sim \text{Mult}(n_t, \gamma_t)$, where n_t is the sample size (in number of diploid individuals) at time t and γ_t:

$$\gamma_{AA,t} = \pi_t^2 + F_{IS}(1 - \pi_t)\pi_t$$

$$\gamma_{Aa,t} = 2(1 - \pi_t)\pi_t(1 - F_{IS})$$

$$\gamma_{aa,t} = (1 - \pi_t)^2 + F_{IS}(1 - \pi_t)\pi_t$$

are the genotype frequencies in the populations in function of the inbreeding coefficient F_{IS}, which is determined by the selfing rate $F_{IS} = \sigma/(2 - \sigma)$. Simulations were performed with parameters values for $N_e = 500$ diploid individuals, $\tau = 25$ generations, $n_0 = n_{25} = 50$ diploid individuals and selfing rate, σ, had values of 0, 0.5, 0.75, 0.8, 0.85, 0.9, 0.925, 0.95, 0.975 or 1. Results are presented in Fig. S3.

Table S1. Notation.

symbol	meaning
γ_t	genotype frequencies in the population at time t
π_t	allele frequency (in the population) at generation t
σ	rate of reproduction by selfing in the population
τ	time in generations between the two samples
AA, Aa, aa	the three possible genotypes of a bi-allelic locus
F_{ST}^l	observed differentiation statistic F_{ST} at focal locus l
K_t	observed genotype counts in the sample at time t
K_t^*	simulated genotype counts in the sample at time t
k_t	observed count of the reference allele in sample at time t
N	census population size in number of diploid individuals
N_e	effective population size in number of diploid individuals
n_t	sample size (number of diploid individuals) at time t
p_t	frequency of the reference allele in the sample at generation t; $p_t = k_t/2n_t$
p_t^*	simulated frequency of the reference allele in the sample at generation t; $p_t = k_t/2n_t$
s	selection coefficient of the adaptive locus
t	time, measured in generations, with $t = 0$ for first sample and $t = \tau$ for last sample

Table S2. Genome scan results for *M. truncatula*.

This table is provided in a supplementary file.

Fig. S1. Effect of a filter on minor allele frequency (MAF) on the distribution of p-values. p-values calculated from 100 simulation replicates of a neutral outcrossing population of $N = 500$ diploid individuals, sampled twice with $\tau = 25$ generations between samples. The blue histogram shows the distribution of p-value using all loci, with corresponding blue line in the QQ-plot. The orange histogram shows the distribution of p-value using loci with a minimum global allele frequency of 0.05, with corresponding orange line in the QQ-plot.

Fig. S2. Schematic representation of the consequences of selection on standing variation in the temporal pattern of genetic diversity. Gene genealogies with samples carrying the derived allele represented with a circle filled in black and samples carrying the ancestral allele represented with a circle filled in white. Mutations are represented as coloured squares over the genealogy at their time of mutation; the site under selection is represented with a circle filled in black. Lineages of the genealogy subject to positive selection are represented in red. Next to each genealogy a schematic representation of the sequence alignment of the samples is presented with each sequence represented as a line at the same height that the corresponding sample of the genealogy. Polymorphisms are represented with the corresponding coloured squares in the genealogy, indicating the sequences that carry the derived alleles. Sequences carrying the allele under positive selection are represented with a red line. Asterisks signal polymorphic sites showing large (*) and extreme (***) changes in allele frequency between the two temporal samples. This figure represents only a small region around the locus under selection that did not recombine through the period considered (hence can be represented as a single genealogy). **(a)** Gene genealogy of two temporal samples taken just before a low frequency derived allele becomes advantageous and after the sweep. The mutation is young, as expected for a low frequency derived allele (Fig. S6), so the lineages carrying the allele that will become advantageous had little time to accumulate mutations and become distinctive from other haplotypes in the population. Most of the alleles on the haplotypes that swept and are found in the second sample on high frequency were already at high frequency in the first sample. **(b)** Gene genealogy of two temporal samples taken just before a low frequency ancestral allele becomes advantageous and after the sweep. The mutation is old, as expected for a high frequency derived allele (Fig. S6); thus, the split between the lineages carrying the ancestral and derived allele is very old. Several mutations have accumulated in both lineages, making them very distinctive. Several alleles on the haplotypes that swept and are found in the second sample on high frequency were at low frequency in the first sample, creating a strong local signal of selection around the selected site.

Fig. S3. Effective population size estimates from temporal differentiation from F_C and F_{ST}. Estimates obtained from simulated data of populations of $N = 500$ diploid individuals, sampled twice with $\tau = 25$ generations between samples and 100 estimates at independent simulation replicates. Dashed line marks true $N_e = \frac{N(\sigma-2)}{2}$. **(a)** Simulation performed with SLiM (Messer 2013) as described in the main text and presented partially in Fig. 1. **(b)** Simulations performed with a simple model of independent loci with initial allele frequency of 0.5, as described in supplementary text S2.

Fig. S4. Effect of recombination occurring before the selective sweep on differentiation and diversity of the haplotypes carrying the advantageous allele. Genetic patters found in a simulated population of $N = 500$ diploid individuals reproducing predominantly by selfing ($\sigma = 0.95$) after $20N_e$ generations. **(a)** Differentiation is measured as the F_{ST} between haplotypes carrying the advantageous allele and those carrying the alternative allele. **(b)** Diversity is measured as the average heterozygosity per bp. Both differentiation and diversity were calculated on 3,000 bp windows at different distances to the loci under selection. The mean from 100 simulation replicates is shown.

Fig. S5. Genetic differentiation as a function of the distance to locus under slection. Estimates of F_{ST} (mean across 100 simulation replicates) from polymorphisms in a window of size 1000Kb ($\approx$1cM) at different distances to locus under selection. Simulation of populations of $N = 500$ diploid individuals, sampled twice with $\tau = 25$ generations between samples and different rates of selfing, σ. A new advantageous mutation appears at generation $t = 0$ with coefficient of selection $s = 0.5$. Samples are made of 50 diploid individuals genotyped at 10,000 biallelic markers (including the locus under selection). Boxplot for 100 estimates from simulation replicates. For reference, expected F_{ST} values for $\sigma = 0$ (black) and $\sigma = 1$ (red) are shown with dashed lines.

Fig. S6. Relationship between frequency (π) and age of mutation. Bidimensional histogram for the frequency and age (in generations) of all mutations in a simulated population of $N = 500$ diploid individuals reproducing predominantly by selfing ($\sigma = 0.95$) after $20N_e$ generations. Theoretical expectation of age $\left(\frac{-2\pi}{1-\pi} \ln(\pi) \frac{1}{2N_e} \right)$ is shown with an orange line (Kimura and Ohta 1973).

Fig. S7. Departure from expectations of the p-value distribution for *Medicago truncatula* genome scan. (a) QQ plots for *Medicago truncatula* (this study) and *Arabidopsis thaliana* (Fig. S21 Frachon *et al.* 2017). For *M. truncatula*, p-values estimated for 987 SNP loci that were polymorphic in the sample with a global minor allele frequency higher than 0.05 (Table S2). Null model considered $N_e = 42$ and $\sigma = 0.99$. **(b)** QQ plots for simulations under a scenario of adaptation from new mutations in a population of $N = 500$ diploid individuals sampled twice with $\tau = 25$ generations between samples and different rates of selfing, $sigma$. A new advantageous mutation appears at generation $t = 0$ with coefficient of selection $s = 0.5$. Samples are made of 50 diploid individuals genotyped at 10,000 biallelic markers (including the locus under selection). The plot consider the distribution of p-values from 100 simulation replicates.