# Cast away on Mindoro island: population dynamics of the critically endangered tamaraw (*Bubalus mindorensis*) at Mounts Iglit–Baco Natural Park

C. Bonenfant[1], J. Burton[2], R. Boyles[3], and E. Schütz[4]

1 Université de Lyon, F-69 000, Lyon ; Université Lyon 1 ; CNRS, UMR 5558, Laboratoire "Biométrie et Biologie Évolutive", F-69 622, Villeurbanne, France.
2 IUCN-SSC / Asian Wild Cattle Specialist Group, Global Wildlife Conservation, Texas, USA
3 Department of Environment and Natural Resources of the Philippines, Occidental Mindoro
4 D'Aboville Foundation and Demo Farm Inc, Manila, Philippines

**Keywords**

counts, density-dependence, exponential model, Mindoro dwarf buffalo, protection measures, poaching, population growth.

**Correpondence**

Christophe Bonenfant, `christophe.bonenfant@univ-lyon1.fr` – ORCID 0000-0002-9924-419X

# Abstract

Endangered species, despite living at low population density, may undergo density-dependent feed-backs in case of successful recovery or marked reduction in range. When at work, density-dependence dynamics increases extinction risks and hamper conservation efforts. The last population of the critically endangered tamaraw (*Bubalus mindorensis*) lives on a $< 3,000$ha in Mounts Iglit-Baco Natural Park (MIBNP) with very limited expansion possibilities. Tamaraw abundance has been monitored on a yearly basis from animal counts by Philippine authorities since the year 2000. Consistent with its protected status by law, the MIBNP tamaraw population has been increasing in size at an average rate of $+6\%$ per year, which we found to be relatively low compared to other similar-sized *Bovinea* species. Population growth was strikingly spatially structured within MIBNP with a population growth close to $+16\%$ in the core area of protection, while a reduction of abundance of $-27\%$ was measured at the periphery of the species range inside MIBNP. Highly concerning is the fact that the annual population growth rate progressively decreased significantly over the years since 2008, which we interpreted as an evidence of density-dependence. This isolated tamaraw population is currently experiencing a contraction of its range at MIBNP, likely caused by anthropogenic pressures forcing large herbivores to live at relatively high density in the core zone of the monitoring where protection is most effective. Our study highlights that beyond the encouraging results of a continuous growth over the last two decades, the MIBNP tamaraw population remains subject to uncertainty of its long term viability.

## INTRODUCTION

2 Low population abundance is a characteristic shared by most endangered species

living in the wild (), and one of the foremost criteria used for the evaluation of

4 conservation statuses (Mace *et al.* 2008; Neel *et al.* 2012). The particular

emphasize given on population abundance in conservation finds its roots in the

6 fact that population viability (Lande *et al.* 2006), and many ecological and

demographic processes depend on the number of animals in a population.

8 Small-sized populations are of concerns because its dynamics differ in several

ways from what is generally reported at higher abundance (Caughley 1994;

10 Mugabo *et al.* 2013). When population size is small, the random death of a few

individuals may precipitate the extinction of a population, a process referred to

12 as demographic stochasticity (Shaffer 1981; Lande 1993). Similarly, when

mating partners are too few to meet and reproduction fails, demographic

14 stochasticity generates the so-called Allee effects (Courchamp *et al.* 1999), that

is a decrease of the population growth at low population abundance (*i.e.*

16 demographic component, see Stephens *et al.* (1999)). On the other hand,

classical density-dependent effects, the decrease of the population growth rate

18 with density (Nicholson 1933), on population dynamics are often overlooked in

conservation because they are *a priori* expected to occur at high population

20 abundance. Increasing density in populations of endangered species may,

however, occur incidentally following a range reduction caused by habitat loss or

22 successful protection measures, and in turn triggers undesirable ecological

consequences for its conservation.

24    Density-dependence is a pervasive demographic responses among large

**3**

mammal populations (Fowler 1987; Bonenfant *et al.* 2009). In addition to

depressing recruitment rates rapidly, density-dependence can increase mortality

of adults as population density approaches the carrying capacity (Eberhardt

1977; 2002). Density-dependence can generate a diversity of dynamics including

chaotic, cyclic and stable dynamics (May 1976) and, in practice, the strength and

shape of the density-dependence response strongly determines the fate and

viability of populations at risk of extinction (Beissinger & Westphal 1998; Henle

*et al.* 2004). With rising density, individuals may die from starvation because of

limiting food resources, but agonistic interactions and exposure to high stress

levels magnify the effect of other mortality sources. For instance, at high

population density, epizootics are more likely to emerge and to spread quickly in

the populations (). Because high population density magnifies the effect of

environmental variation on demographic rates (Hones & Clutton-Brock 2007),

the risks of massive mortality events are much higher if the population is

maintained in this state on the long term. Finally, at high population density, the

number of animals that can be removed, for reinforcement or captive breeding

purposes, without decreasing population size becomes smaller as the population

growth rate converges to 0 (Boyce 1989). Obviously an early detection of any

signs of density-dependence is of the upmost importance for the management

and conservation of animal species to limit the risk of catastrophic events.

The tamaraw (*Bubalus mindorensis*) is a large mammalian herbivore endemic

to the Island of Mindoro and the only wild cattle of the Philippine archipelago

(Heude 1888; Custodio *et al.* 1996). Despite the tamaraw being officially

protected by law in 1954 and the creation of the Mts Iglit-Baco Natural Park

(MIBNP) protected area in 1970 (Maala 2001), its distribution range has been

**4**

dramatically shrinking over the last decades. Originally found across the whole

island of Mindoro with a gross estimate of 10,000 individuals in 1900 (Long

*et al.* 2018), the tamaraw has suffered from intensive land conversion for

agriculture and logging industry, which negative effects on wildlife were further

exacerbated by trophy hunting and disease outbreaks spread out by domestic

cattle (Maala 2001). In recent years, the main putative causes of tamaraw

reduction in distribution range are believed to be a combination of poaching

from lowlander Filipinos, together with land encroachment, and traditional

hunting activities conducted by upland indigenous communities who share their

living space with the large herbivore (). As a consequence, the species is now

found in few isolated populations scattered across Mindoro, with MIBNP

hosting the largest number (Matsubayashi *et al.* 2010; Wilson & Mittermeier

2011; Long *et al.* 2018). The MIBNP tamaraw population has been restricted to

one location of less than <3,000 ha in 2018, that we referred to as the Core Zone

of the Monitoring (CMR, see Fig. 1).

    In an attempt to reinforce tamaraw protection, wildlife managers came to a

much needed agreement with residing indigenous people in 2016, declaring a

1,600 ha no-hunting zone area (NHZ) within the CZM (Fig. 1). An unforeseen

consequence of the NHZ establishment for tamaraw seems to be an increase of

pressure by traditional hunting at its direct boundaries. MIBNP rangers regularly

report setting of snare and spear traps, and violation of the NHZ agreement in

the last year is not uncommon with several deaths of tamaraws confirmed ().

Another response of large herbivores to hunting is space use modification to

reduce the risk of being killed (Chassagneux *et al.* 2019; Marantz *et al.* 2016).

For tamaraw, the ecological consequences of poaching and traditional hunting

are threefold. (*i*) Mortality of animals is likely higher at the periphery than inside

the NHZ, especially during the rainy season at the time of hunting season by

residing indigenous communities. (*i*) Consequently, dispersion of individuals

outside of the CZM becomes risky and leads to almost no emigration, so that the

MIBNP tamaraw population is considered demographically closed. (*iii*) Because

harvesting prevents the geographic expansion of the population, available space

for tamaraws is constrained within the 3,000 ha CZM at best, which must set an

upper limit to the maximum number of individuals these CZM can support.

Given its critical importance for the conservation of these iconic species, the

MIBNP tamaraw population has been the cornerstone of all past, present and

future conservation plans (Maala 2001; Long *et al.* 2018). Unfortunately, the

population dynamics of tamaraws in MIBNP has not been documented so far,

nor do we have any estimation of its average population growth rate (noted *r*).

Yet a better knowledge about its ecology and population dynamics is

fundamental to assess its current status, strengthen protection measures and to

guide future conservation strategy with evidence-based information (Caughley

1994; Norris 2004) (Stewart *et al.* 2005; Sutherland *et al.* 2004). Every year

since 2000, the Tamaraw Conservation Program (TCP), a banner program of the

Philippine Department of Environment and Natural Resources (DENR), has

been conducting tamaraw counts on 18 vantage points distributed across the

CZM in MIBNP, yielding an annual index of relative population abundance.

These annual counts make the only available source of information about

tamaraw population trends and dynamics for officials and conservation agencies.

Here we present a detailed analysis of the yearly counts at MIBNP to provide the

first estimate of the average population growth (*r*) of tamaraw. We also used this

**6**

19 years-long time series of abundance to test the following predictions:

1. This dwarf buffalo of Mindoro is among the rarest large herbivore of
   south-east Asia, justifying its 'critically endangered' status (CR) on the
   IUCN Red List (IUCN2019 2019). The tamaraw is therefore under the
   highest status of protection with substantial conservation effort in the last
   decade. Because protection measures aimed specifically at increasing the
   abundance of tamaraw, we predict the MIBNP population should increase
   in size since 2000 and hence, $r$ to be statistically $> 0$. Similarly, we also
   expect the agreement of no hunting zone with IPs to reduce related
   mortality, and hence predict a higher average growth rate ($r$) after than
   before 2016;

2. Because the no-hunting agreement area might create a spatial demarcation
   between risky and safe habitats, tamaraws may adjust their space use to
   limit mortality risks by concentrating in the center of the CZM. In
   addition, animals living on the boundaries of the CZM or moving outside
   of the NHZ are subject to a higher risk of poaching or to be injured by
   traps. We hence expect local growth rate to differ in space and predict a
   decrease of $r$ while getting away from the main rangers base camps and
   patrolling routes (namely VP Loibfo and Magawang in the center of the
   CZM (Fig. 1), where the monitoring is more prominent and disturbance
   lower, towards the limits of the CZM;

3. Given the absence of large mammalian predators on Mindoro and
   successful mitigation of direct threats thanks to protection measures in the
   context of a restricted range the MIPBN tamaraw population could be

approaching the carrying capacity of the CZM. Under this hypothesis, we expect the occurrence of density-dependence at MIBNP, *i.e.* a decrease of the annual population growth rate ($r$) with population abundance;

## MATERIAL AND METHODS

### Study site

The study area is located within the Mounts Iglit-Baco Natural Park (N12°54′ E121°13′) in the south-central part of Mindoro which, with 106,665 ha, is the island largest protected area. MIBNP shelters the largest number of tamaraw in a single site located on the south-west edge of the Protected Area. (Fig. 1). Tamaraw distribution range is now restricted to less than 3,000 ha, which represents approx. 3% the MIBNP surface. We will refer to this area as the Core Zone of the Monitoring (CZM) because this is where most of the patrolling and observations are carried out by the Park's officers. There is currently 15 park rangers divided into two or three teams to monitor the area on a regular basis, following regular distinct patrolling routes centered around the three base camps. (Fig. 1).

The CZM area is a rolling grassland plateau with an average elevation of 800m a.s.l., dominated by cogon (*Imperata cylindrica*) and wild sugarcane (*Saccharum spontaneum*), and interspersed with numerous wooded creeks, secondary forest fragments and steep hills. In recent years, the invasive Siam weed (*Chromolaena odorata*, locally called hagonoy) has been quickly spreading in the grassland and now covers an unknown fraction of the tamaraw habitat. The tropical climate is strongly seasonal with a rainy season spanning

from June to October (mean temperature 26.5°C; mean rainfall: 150 mm), and a

hot dry season from November to May (mean temperature 29.8°C; mean rainfall:

5 mm). Apart from humans, the only potential predator to the tamaraw is the

reticulated python (*Malayopython reticulatus*) that is capable of preying upon

calves or yearlings but no such observation have been made so far.

## Tamaraw counts

Since 2000 and for every single year since then, a standardized protocol of

tamaraw counts has been carried out within MIBNP. The tamaraw counting area

encompasses 18 vantage points, covering nearly 2,200 ha within the CZM

(Fig. 1). The method is rather invasive for the MIBNP ecosystem because it

involves the burning of grasslands a few weeks ahead of each count, in order to

increase visibility and to attract tamaraws to areas with highly palatable and

attractive regrowing grasses. Each year, between 1,200 and 1,500 ha of

grasslands were intentionally burned for the purpose of monitoring tamaraw

abundance. Individual tamaraws were counted simultaneously at the 18 vantage

points for two hours at dusk and dawn, and for four consecutive days, in late

March or early April. For the 8 sessions and at each vantage points, at least two

observers spotted and recorded all animal seen on field paper sheets, writing

down observation time, approximate location, and assessed sex and age-category

of individuals (split into calves, yearlings, sub-adults and adults). Right after the

completion of the 8 sessions, count data were cross-checked and cleaned for

obvious double observations, and the total number of different animals seen was

then derived for each vantage point $s$ and year $t$.

## Data analyses

For populations of long-lived and endangered species, robust and accurate estimations of $r$ remains difficult to obtain. Knowledge about $r$ requires either the estimation of demographic rates combined into a Leslie matrix (Leslie 1945) or the availability of long time series of population abundance, preferably on an annual basis. In the absence of natural marking, capturing individuals of an endangered species to make them identifiable with tags often face strong and justified resistance for ethical reasons, mostly because of unavoidable mortality risks. Consequently, in the best case the monitoring of the population abundance of endangered species relies on an estimator accounting for detection probability of animals (*e.g.* Kéry & Schmidt 2008), but most of time a simple index of abundance is carried out ().

Working with abundance index such as annual counts to derive an empirical estimates of $r$ however comes with some serious methodological issues (Îto 1972)(Lindley 2003). Over the last decades, population counts have repeatedly been associated with a large sampling variance (*e.g.* Caughley 1977), and coefficient of variation for population abundance as high as 30% were reported for species 1, species 2 for instance. From a conservation biologist viewpoint, a large sampling variance in the input data for estimating $r$ is a major pitfall because of the over-optimistic results it leads to. Both the point estimate and its precision are overestimated in the presence of sampling variance (Lindley 2003). Despite the availability of appropriate statistical methods to account for the sampling variance of population counts (Kalman filter,  De Valpine & Hastings 2002), such tools remain seldom used in practice for conservation.

Here we analyzed the number of seen tamaraws and judged as different

2   individuals by observers for each of the 18 vantage points, split by years ($C_{s,t}$,

where indice $s \in \{1, \ldots, 18\}$ stands for the vantage point, and

4   $t \in \{2000, \ldots, 2019\}$ stands for the time in years. We implemented a state-space

model to tease apart process from sampling variances, making the assumption

6   that double and under-counts compensate and are randomly distributed from

year to years. We worked in a Bayesian framework (*e.g.* Kéry & Schaub 2011)

8   but note this is fully equivalent to a Kalman filter with a frequentist approach. To

do so we defined $N_{s,t}$ as the unobserved 'true' abundance linked to $C_{s,t}$ by a

10   random effect corresponding to the process variance ($\sigma_c^2$), on the log scale to

ensure positive values for abundance. Our baseline model was:

$$\log(C_{s,t}) \sim \mathcal{N}(\log(N_{s,t}), \sigma_c^2)$$

12   Accordingly, we computed the population growth rate $r_{s,t}$ of the tamaraw

population at each vantage points *s*, as:

$$r_{s,t} = \log\left(\frac{N_{s,t+1}}{N_{s,t}}\right),$$

14   from counts corrected for the sampling variance, hence returning an unbiased

estimation of the population growth rate. The overall population growth of the

16   tamaraw population of MIBNP is then given by:

$$\bar{r} = \frac{1}{19} \sum_{t=1}^{19} \log\left(\frac{N_{t+1}}{N_t}\right),$$

whereby $N_t = \sum_{s=1}^{18} N_{s,t}$. We informed $N_t$ with the total number of animals seen

each count session ($C_{tot}$) and linked counts at the VP and population levels with

the following observation equation: $C_{tot} \sim \mathcal{N}(N_t/p, \sigma_{tot}^2)$. The consolidated

number of tamaraws $C_t$ being much smaller than total number seen $C_t$ from

which double-counts were removed, we included a constant proportion

parameter $p$, similar to a "detection rate" (which is not). By doing so, we could

use the available information about tamaraw counts fully, including both the

knowledge of experienced rangers and the raw number of detected tamaraws

with no *a priori* interpretation. In addition, we could input missing values at the

VP level for years 2000 to 2003 and 2005 thanks to the $C_{tot}$ constrain on $N_t$.

To test our third hypothesis we investigated potential density-dependence in

the annual growth rate by fitting a logistic growth model to the total number of

tamaraws seen per year $N_t$:

$$\log(N_{t+1}) = \log(N_t) + r_m \times \left(1 - \frac{N_t}{K_l}\right),$$

where $r_m$ is the maximum population growth for the tamaraw, and $K$ is the

carrying capacity of the MIBNP. Likewise, we tested if the annual population

growth rate would decrease in time with population density by fitting a

Gompertz model:

$$\log(N_{t+1}) = \log(N_1) + \log\left(\frac{K_G}{N_1}\right) \times \left(1 - e^{-\beta \times t}\right),$$

Because the pair of parameters $r_m$ and $K_l$ (logistic), and $\beta$ and $K_G$

(Gompertz), are not fully separately identifiable (Lebreton & Gimenez 2013), we

**12**

used informative priors for $r_m$ and $\beta$ based on the reported maximum population

growth for other large bovids (see Traill *et al.* 2007,  for a similar approach).

Accordingly, we set a moderately informative prior value of mean 0.3 and a

precision of 0.1 in our models (Table 2). We assessed the different models by

computing explicitly, during the model fit, the probability for our abundance

time series to be generated by each of the three demographic models

(exponential, Gompertz and logistic).

We fitted our statistical model to the tamaraw count data with JAGS 4.0

(Plummer *et al.* 2003), with 3 MCMC chains and a burn-in of 40,000 iterations.

We obtained parameter estimates with an additional 30,000 iterations and a

thinning factor of 5 (hence a total of 6,000 MCMC samples). We checked model

convergence graphically by investigating the mixing of MCMC chains, along

with the $\hat{R}$ statistics (Brooks & Gelman 1998). With the exception of $r_m$ (see

above), we set uninformative priors with large variance for all other parameters

to be estimated, and checked the sensitivity of our results to the average value of

the priors by replicating the analyses with variable mean and initial values for

prior distributions. We report all parameters estimates as the mean of the

posterior distributions along with the 95% percentile for the credible intervals

following Louis & Zeger (2008):  $_{95\%low}\, estimate\, _{95\%up}$.

Note that despite the fact that we analysed the number of seen tamaraws as

described above, we chose not to display such numbers on figures and to report a

standardized abundance ($C_{z,t}/max(C_{z,t})$) instead. This transformation applies to

the estimated carrying capacities of the Gompertz and logistic models,

$K/max(C_{z,t})$, hence reported values for $K$ are $> 1$. We did not provide raw

counts to avoid a likely confusion between the index of relative tamaraw

**13**

abundance we worked with, and a real population size estimator that was not

2  implemented at MIBNP. Such a variable transformation has no incidence on the

biological interpretation of the results though.

4  **RESULTS**

From our baseline model of tamaraw counts, we estimated an average annual

6  growth rate of $\bar{r} = {}_{0.02} 0.06 {}_{0.12}$ for the MIBNP population from 2000 to 2019,

which associated variance process equals $\sigma_{proc} = {}_{0.25} 0.37 {}_{0.49}$ (Fig. 2). As

8  expected for count data, the sampling variance was large $\sigma_c = {}_{7.32} 9.31 {}_{9.98}$

suggesting frequent double counts of the same individuals. That the MIBNP

10 tamaraw population increased in size by c.a. 6% per year and was statistically

different from 0 is a clear support of our first hypothesis predicting increase of

12 abundance thanks to successful protection efforts at MIBNP.

Considering the vantage point specific growth rates ($r_s$), we observed a

14 marked variability in space (Table 2; Fig. 3) considering that growth rates ranged

between $r_s = {}_{-2.52} -0.32 {}_{0.20}$ at Tarzan VP and $r_s = {}_{0.03} 0.19 {}_{0.35}$ at Inubon VP.

16 Vantage point specific growth rates were spatially structured at MIBNP (Fig 3).

With no exception, $r_s < 0$ were located at the periphery of the NHZ (Iyan,

18 Tarzan, Saligue East, Mibluan, Fangandatan) although $r_s$ of a few vantage points

was $> 0$ despite being located on the NHZ limit too (Nagbobong, Malibayong,

20 Talafu West and East). Together these results are in accordance with our

hypothesis of an on-going contraction of the tamaraw range at MIBNP after

22 2000, and the disappearance of individuals at the periphery of the core zone of

the monitoring.

**14**

Analytical evidence for density-dependence in the tamaraw dynamics

between year 2008 and 2019 was moderate to high. We explicitly estimated the

probability $\pi$ that the exponential, Gompertz and logistic models would be the

underlying process generating the observed variation in tamaraw abundance

(proxy of Baye's factor). Our different models did have different probability to

have generated to observed data with, respectively $\pi = 0.80$, $\pi = 0.00$ and

$\pi = 0.20$ for the exponential, Gompertz and logistic model respectively. The

estimated coefficients of the Gompertz model lent statistical support for the

hypothesis of a gradual decrease in the annual population growth with time

($\beta = {}_{0.07} 0.09 {}_{0.11}$), and estimated carrying capacity to $K_G = {}_{1.12} 1.33 {}_{1.64}$.

Similarly, point estimates for the two parameters of the logistic growth function

were $r_m = {}_{0.09} 0.11 {}_{0.13}$ for the maximum population growth rate and

$K_l = {}_{1.14} 1.35 {}_{1.61}$ for the carrying capacity, very close to $K_G$ estimates.

## DISCUSSION

The isolated tamaraw population in MIBNP has been increasing in size since

2000 (Fig. 2). It is not doubt for such an highly endangered species that a

positive growth rate of 0.06 over 19 years of monitoring is a great success, both

locally and for the global conservation effort. The devil lies in the details though,

and two additional results darken this otherwise positive picture of tamaraw

population dynamics and conservation. First, a progressive decrease of the

average growth rate suggests density-dependent effects at the population level.

Furthermore, we found support for the hypothesis of a contraction of the

tamaraw distribution, evidenced by a clear spatially structured pattern of the

population growth rates. The population dynamics we document is clearly

calling for new and urgent conservation actions if the future of tamaraw is to be

ensured.

## Population growth

The average population growth rate of $\bar{r} = 0.06$ we found gives strong evidence

in favor of our hypothesis of successful conservation policies of tamaraw over

the past two decades. Yet, an annual growth of 6% is relatively low compared to

other similar-sized or even larger large herbivores (Table 2). For instance, a

population of African buffalo (*Syncerus caffer*) in Southern Africa grew at a rate

of 12% per year for 28 years (Jolles 2007), and values close to 30% per year

have regularly been recorded for bison (*Bison bison*) or banteng (*Bos javanicus*)

despite these species are between 2 and 2.5 times larger in size than the tamaraw.

From this brief comparative approach, we could expect a substantially higher

average growth rate for tamaraws, suggesting that specific environmental factors

are limiting its population dynamic in MIBNP.

A first limiting factor could stand in the fact that the most available and

accessible biotope for the tamaraw in MIBNP is grassland dominated (83% of

the CMZ) by strong pioneer competitive species such as cogon and wild

sugarcane; both grasses becoming de facto prominent in the tamaraw diet. While

providing tamaraws with abundant and high quality forage at the early stages of

plant growth, the grassland could rapidly become of much lower nutritive value

during the dry season. Once grass growth is complete, stems exceeds tamaraw

height (300cm vs. $<$ 120 cm respectively) and dry out to become unpalatable,

raising the question of food resource accessibility and availability for tamaraws

**16**

and the ambivalent role of fire regime. On the one hand, regular burnings

rejuvenate vegetation and promote regeneration of attractive new grass shoots

few months ahead of the rainy season, when animals might look for alternative

marginal habitats such as forest or creeks for complementary feeding purposes.

On the other hand, previous observation suggest that regular fire could promote

Siam weed expansion (Nath *et al.* 2019), which is indeed rapidly spreading on

the study site. Like other grazing herbivores (Rozen-Rechels *et al.* 2017),

tamaraw avoid this alien plant that could in turn decrease the CMZ carrying

capacity (**?**). More generally tamaraw feeding preference is not fully understood

and might be a mixed-feeding or a browsing herbivore (*sensu* Hofmann 1989).

While being *de facto* restricted on grassland habitats, tamaraw could be

ecologically flexible and adapted to forested habitats, as recent assessment of a

new tamaraw population in mountain forest habitats in the upland inner Mindoro

Island would suggest (Schütz 2019). In other words, the CZM might actually

provide ecological conditions whose intrinsic limits are artificially reduced, so

far, thanks to long lasting and highly disturbing habitat management

policies;detrimental impact starting progressively to exceed beneficial effects.

An additional factor related to human activities could account for the low

tamaraw population growth rate we found. Tamaraw has been traditionally

hunted by IPs and both intentional and unintentional casualties still happen.

Besides tamaraw is sought after for bushmeat by lowlander poachers. Both

threats are hard to control in the field, are likely pervasive across MIBNP, and

lead to an increased mortality of all age-classes of tamaraws to an unknown

extent. Positive growth over the past two decades is a clear sign of successful

protection measures thanks to the constant presence of rangers and the recurrent

**17**

patrols that have likely contributed to decrease harvesting pressure within the

CZM. However, the large heterogeneity in the observed growth rates with

negative values at the periphery of the counting area might reveal a lower

protection efficiency when moving away from the main patrolling routes and

base camps. Actually, the CZM area functions like a source-sink system with

central areas nearby base camps concentrating most of the vantage points with

positive growth rates, with local value as high as 16% (Table 1), more in line

with the average growth we review for large bovids (Table 2).

## Density-dependence

At first glance, the gradual decrease of the tamaraw population growth rate with

overall abundance at MIBNP would suggest some sort of density-dependence,

that is a change in demographic rates lowering the number of individuals added

to the population from year to year (Fowler 1987; Sinclair 1989). Predicted

growth rates from the lowest to the highest abundance varied between $r_t = 0.11$

and $r_t = 0.01$ for tamaraw at MIBNP. Close to zero population growth rates are

expected when the *per capita* food resource limits the energy budget of

individuals (Begon 1984). Detection of density-dependence in MIBNP is

worrying because it suggests that tamaraws could start suffering from food

resource shortage. In absence of safe corridors for individuals to leave the CZM

and the adverse effects of Siam weed on forage quantity, we cannot expect this

population to keep on increasing in size for the next decade as we report between

2000 and 2019. From current data, we could estimate a carrying capacity of 1.4

meaning the current MIBNP tamaraw abundance has already reached almost 2/3

of the carrying capacity. This first explanation is the *biological* hypothesis of

**18**

density-dependence, which is well known and repeatedly shown in many

2 populations of large mammals (Bonenfant *et al.* 2009). In absence of

demographic rates estimations of tamaraws at MIBNP we however hit the limits

4 of the so-called pattern-oriented approach (Krebs 2002). In other words we are

left with the interpretation of a pattern, *i.e.* a temporal variation of the annual

6 population growth rate with no way to evidence real changes in demographic

rates with abundance, as the process-oriented approach would allow.

8      Consequently, an alternative interpretation of the density-dependence pattern

that we report for tamaraw is possible, which we describe as the *count bias*

10 hypothesis. Visual counts of animals, as it is conducted in MIBNP, are

notoriously known to be inaccurate and prone to underestimation of the real

12 abundance (Andersen 1953; Laake et al. 1999). Tamaraw counts were not

designed to be used with a population abundance estimator such as returned by

14 the distance sampling (Buckland *et al.* 2004) or capture-recapture methods

(Schwarz & Seber 1999), and therefore failed to account for the imperfect

16 detection of individuals. The proportion of undetected tamaraws could increase

with their overall abundance, a previously reported effect on red deer (*Cervus*

18 *elaphus* Garel *et al.* 2010) and roe deer (*Capreolus capreolus*, Pellerin et al.

2018). Alternatively, different cognitive biases (Baddeley et al. 2004) could

20 generate similar patterns if, for instance, observers unconsciously pull the total

number of animals downward when consolidating count data. At this stage,

22 counts at year $t$ could likely influence the results of counts at year $t + 1$,

assuming observers are conservative in their expectation of the tamaraw annual

24 population growth. The monitoring of additional biological variables in the field

such as the reproductive success of females could help at teasing apart the two

**19**

hypotheses.

## 2  Conclusion

Current conservation measures assume tamaraw population growth will continue
in the next decade, without a need to mitigate increased pressures thought to be
impacting the periphery of the CZM or to address the density dependence
pressure on the population. Our results are informing new strategic planning for
protection and management within MIBNP, as part of the MIBNP Management
Plan and the island-wide Tamaraw Conservation Management and Action Plan.
Firstly, there is two main aspects that must be urgently addressed: (a) increase
the available space for the species so to mitigate density-dependence effect and
maintain positive growth rate, that must be combined with (b) increasing the
effectiveness of ranger patrols in the periphery area of the CZM to reduce the
recurrent poaching pressure from Filipino lowlanders and losses due to
traditional hunting. These targets request to collaborate closely with residing IP
communities on a more sustainable land-use and traditional hunting system.
Indeed, a more effective protection and suitable habitat management might in
turn increase the average growth rate at MINBP, and exacerbate the observed
density-dependence demographic responses of tamaraws. Secondly, when
investigating the feasibility of translocating animals from this source population
to other sites, estimated demographic rates should be incorporated into
population modelling for extraction number that maintain a sustainable yield and
the tamaraw population viability. Indeed, the maximum sustainable yield is
found at $K/2$ in density-dependent models, not when population abundance
reaches carrying capacity $K$.

**20**

Our findings exemplifies how detailed analyses of relative population

abundance data, even without absolute population size estimators, can contribute

to determining future conservation measures and are essential for large mammal

conservation. As a greater proportion of wild cattle species are inhabiting in

human impacted or intensively managed landscapes, there is an increasing need

to understand the interaction between anthropogenic activities and the population

dynamics of protected species. Similar to the tamaraw case is the banteng (*Bos*

*javanicus*) population of Alas Purwo National Park, Indonesia. In spite of an

increasing abundance, this population has become consolidated close to an

artificially maintained grazing area, which may be causing increased predation

pressure of juveniles, and the population dynamics and carrying capacity are not

yet known (JB, pers. obs). Clearly, the protection of large herbivores may hence

come at the costs of undesirable ecological effects, rendering their conservation

even more difficult.

## ACKNOWLEDGMENTS

A lot of people are to thank here. We are grateful to Fernando Garcìa for producing

maps used in this paper. Special thanks are extended to Alvaro Gonzalez.

## REFERENCES

Begon, M. (1984). Density and individual fitness: asymmetric competition. In: *Evolutionary Ecology* (ed. Shorrocks, B.). Blackwell, vol. 23, pp. 175–194. TY - CONF Publication Type: Conference Proceedings Language: English.

Beissinger, S.R. & Westphal, M.I. (1998). On the use of demographic models of population viability in endangered species management. *The Journal of wildlife management*, pp. 821–841.

Bonenfant, c., Gaillard, J.M., Coulson, T., Festa-Bianchet, M., Loison, A., Garel, M.,

Loe, L.E., Blanchard, P., Pettorelli, N., Owen-Smith, N., Du Toit, J. & Duncan, P. (2009). Empirical evidence of density-dependence in popualtions of large herbivores. *Adv. Ecol. Res.*, 41, 313–357.

Boyce, M.S. (1989). *The Jackson elk herd : intensive wildlife management in North America*. Cambridge University Press, New York. Mark S. Boyce. Includes index. Bibliography: p. [274]-292.

Brooks, S.P. & Gelman, A. (1998). General methods for monitoring convergence of iterative simulations. *Journal of Computational and Graphical Statistics*, 7, 434–455.

Buckland, S.T., Anderson, D.R., Burnham, K.P., Laake, J.L., Borchers, D.L. & Thomas, L. (2004). *Advanced distance sampling*. Oxford University Press.

Buys, D. & Dott, H. (1991). Population fluctuations and breeding of eland *Taurotragus oryx* in a western Transvaal nature reserve. *Koedoe*, 34, 31–36.

Caughley, G. (1977). *Analysis of vertebrate populations*. Wiley, London. Graeme Caughley. 'A Wiley-Interscience Publication' Includes index. Bibliography: p.217-228.

Caughley, G. (1994). Directions in conservation biology. *Journal of animal ecology*, pp. 215–244.

Chassagneux, A., Calenge, C., Siat, V., Mortz, P., Baubet, E. & Saïd, S. (2019). Proximity to the risk and landscape features modulate female red deer movement patterns over several days after drive hunts. *Wildlife Biology*, 2019.

Choquenot, D. (1993). Growth, body condition and demography of wild banteng (*Bos javanicus*) on Cobourg Peninsula, northern Australia. *Journal of Zoology*, 231, 533–542.

Cornélis, D., Melletti, M., Korte, L., Ryan, S.J., Mirabile, M., Prin, T. & Prins, H.H. (2014). *African buffalo Syncerus caffer (Sparrman, 1779)*, Cambridge University Press, pp. 326–372.

Courchamp, F., Clutton-Brock, T. & Grenfell, B. (1999). Inverse density dependence and the allee effect. *Trends in ecology & evolution*, 14, 405–410.

Custodio, C.C., Lepiten, M.V. & Heaney, L.R. (1996). *Bubalus mindorensis*. *Mammalian Species*, pp. 1–5.

De Valpine, P. & Hastings, A. (2002). Fitting population models incorporating process noise and observation error. *Ecological Monographs*, 72, 57–76.

Eberhardt, L.L. (1977). Optimal policies for conservation of large mammals, with special references to marine ecosystems. *Environmental Conservation*, 4, 205–212.

Eberhardt, L.L. (2002). A paradigm for population analysis of long-lived vertebrates. *Ecology*, 83, 2841–2854.

Fowler, C.W. (1987). A review of density dependence in populations of large mammals. *Curr. Mammal.*, 1, 401–441.

Fuller, J.A., Garrott, R.A., White, P., Aune, K.E., Roffe, T.J. & Rhyan, J.C. (2007). Reproduction and survival of Yellowstone bison. *Journal of Wildlife Management*, 71, 2365–2372.

Garel, M., Bonenfant, C., Hamann, J.L., Klein, F. & Gaillard, J.M. (2010). Are abundance indices derived from spotlight counts reliable to monitor red deer *Cervus elaphus* populations? *Widlife Biology*, ??, ??

Gates, C. & Larter, N. (1990). Growth and dispersal of an erupting large herbivore population in northern Canada: the Mackenzie wood bison (*Bison bison athabascae*). *Arctic*, pp. 231–238.

Haque, N. (1990). *Study on the ecology of wild ungulates of Keoladeo national Park Bharatpur, Rajasthan*. Ph.D. thesis, Aligarh Muslim University.

Heinen, J.T. & Paudel, P.K. (2015). On the translocation of wild asian buffalo (**?**) in Nepal: are feral backcrosses worth conserving? *Conservation Science*, 3.

Henle, K., Sarre, S. & Wiegand, K. (2004). The role of density regulation in extinction processes and population viability analysis. *Biodiversity and Conservation*, 13, 9–52.

Heude, P. (1888). Études sur les ruminants de l'Asie orientale. cerfs des Philippines et de l'Indo-Chine. *Mémoires Concernant l'Histoire Naturelle de l'Empire Chinois par des Péres de la Compagnie de Jésus*, 2, 1–10.

Hofmann, R.R. (1989). Evolutionnary steps of ecophysiological adaptation and diversification of ruminants: a comparative view of their digestive system. *Oecologia*, 78, 443–457.

Hone, J., Duncan, R.P. & Forsyth, D.M. (2010). Estimates of maximum annual population growth rates ($r_m$) of mammals and their application in wildlife management. *Journal of Applied Ecology*, 47, 507–514.

Hones, J. & Clutton-Brock, T. (2007). Climate, food, density and wildlife population growth rate. *J. Anim. Ecol.*, ?, ?–?

Îto (1972). On the methods for determining density-dependence by means of regression. *Oecologia*, 10, 776–778.

IUCN2019 (2019). The IUCN Red List of threatened species. Version 2019-2.

**23**

Jolles, A.E. (2007). Population biology of african buffalo (*Syncerus caffer*) at Hluhluwe-iMfolozi Park, South Africa. *African Journal of Ecology*, 45, 398.

Kemp, N., Dilger, M., Burgess, N. & Van Dung, C. (1997). The saola *Pseudoryx nghetinhensis* in Vietnam – new information on distribution and habitat preferences, and conservation needs. *Oryx*, 31, 37–44.

Kéry, M. & Schaub, M. (2011). *Bayesian population analysis using WinBUGS: a hierarchical perspective*. Academic Press.

Kéry, M. & Schmidt, B. (2008). Imperfect detection and its consequences for monitoring for conservation. *Community Ecology*, 9, 207–216.

Koláčková, K., Hejcmanova, P., Antonínová, M. & Brandl, P. (2011). Population management as a tool in the recovery of the critically endangered Western Derby eland *Taurotragus derbianus* in Senegal, Africa. *Wildlife Biology*, 17, 299–311.

Krasiński, Z.A. (1978). Dynamics and structure of the european bison population in the białowieża Primeval Forest. *Acta Theriologica*, 23, 3–48.

Krebs, C.J. (2002). Two complementary paradigms for analysing populations dynamics. *Philosophical Transactions of the Royal Society of London, Series B*, 357, 1211–1219.

Lande, R. (1993). Risks of population extinction from demographic and environmental stochasticity and random catastrophes. *The American Naturalist*, 142, 911–927.

Lande, R., Engen, S., Sæther, B. & Coulson, T.N. (2006). Estimating density dependence from time series of population age structure. *American Naturalist*, 168, 76–87.

Larter, N., Sinclair, A., Ellsworth, T., Nishi, J. & Gates, C. (2000). Dynamics of reintroduction in an indigenous large ungulate: the wood bison of northern Canada. 3, 299–309.

Lebreton, J.D. & Gimenez, O. (2013). Detecting and estimating density dependence in wildlife populations. *Journal of Wildlife Management*, 77, 12–23.

Leslie, P.H. (1945). On the use of matrices in population mathematics. *Biometrika*, 33, 182–212.

Long, B., Schütz, E., Appleton, M., Gonzales, A., Holland, J., Lees, C., Marandola, E., Natural, V.C.J., Pineda-David, M.T.J., Salao, C., Slade, J., Tan, E.H.P., Tiongson, L. & Young, S. (2018). Review of tamaraw (*Bubalus mindorensis*) status and conservation actions. *BULLtin*, 1, 18–34.

**24**

Louis, T.A. & Zeger, S.L. (2008). Effective communication of standard errors and confidence intervals. *Biostatistics*, 10, 1–2.

Maala, C.P. (2001). Endangered Philippine wildlife species with special reference to the Philippine eagle (*Pithecophaga jefferyi*) and tamaraw (*Bubalus mindorensis*). *Journal of International Development and Cooperation*, 8, 1–17.

Mace, G.M., Collar, N.J., Gaston, K.J., Hilton-Taylor, C., Akçakaya, H.R., Leader-Williams, N., Milner-Gulland, E.J. & Stuart, S.N. (2008). Quantification of extinction risk: Iucn's system for classifying threatened species. *Conservation biology*, 22, 1424–1442.

Marantz, S.A., Long, J.A., Webb, S.L., Gee, K.L., Little, A.R. & Demarais, S. (2016). Impacts of human hunting on spatial behavior of white-tailed deer (*Odocoileus virginianus*). *Canadian Journal of Zoology*, 94, 853–861.

Mathur, V.B. (1991). *The ecological interaction between habitat composition, habitat quality and abundance of some wild ungulates in India*. Ph.D. thesis, University of Oxford.

Matsubayashi, H., Boyles, R.M., Salac, R.L., Del Barrio, A., Cruz, L., Garcia, R.A., Ishihara, S. & Kanai, Y. (2010). Present status of tamaraw (*Bubalus mindorensis*) in Mt. Aruyan, Mindoro, Philippines. *Tropics*, 18, 167–170.

May, R. (1976). Simple mathematical models with very complicated dynamics. *Nature*, 261, 459–469.

Mugabo, M., Perret, S., Legendre, S. & Le Galliard, J.F. (2013). Density-dependent life history and the dynamics of small populations. *Journal of Animal Ecology*, 82, 1227–1239.

Mysterud, A., Bartoń, K., Jedrzejewska, B., Krasiński, Z., Niedziałkowska, M., Kamler, J., Yoccoz, N. & Stenseth, N. (2007). Population ecology and conservation of endangered megafauna: the case of european bison in białowieża Primeval Forest, Poland. *Animal Conservation*, 10, 77–87.

Nath, A., Sinha, A., Lahkar, B.P. & Brahma, N. (2019). In search of aliens: Factors influencing the distribution of *Chromolaena odorata* L. and Mikania micrantha Kunth in the Terai grasslands of Manas National Park, India. *Ecological engineering*, 131, 16–26.

Neel, M.C., Leidner, A.K., Haines, A., Goble, D.D. & Scott, J.M. (2012). By the Numbers: How is Recovery Defined by the US Endangered Species Act? *BioScience*, 62, 646–657.

Nicholls, A., Viljoen, P., Knight, M. & Van Jaarsveld, A. (1996). Evaluating population persistence of censused and unmanaged herbivore populations from the Kruger National Park, South Africa. *Biological Conservation*, 76, 57–67.

Nicholson, A.J. (1933). The balance of animal populations. *J. Anim. Ecol.*, S2, 132–178.

Norris, K. (2004). Managing threatened species: the ecological toolbox, evolutionary theory and declining-population paradigm. *Journal of Applied Ecology*, 41, 413–426.

Ottichilo, W.K., De Leeuw, J., Skidmore, A.K., Prins, H.H. & Said, M.Y. (2000). Population trends of large non-migratory wild herbivores and livestock in the Masai Mara ecosystem, Kenya, between 1977 and 1997. *African journal of Ecology*, 38, 202–216.

Plummer, M. *et al.* (2003). Jags: A program for analysis of bayesian graphical models using gibbs sampling. In: *Proceedings of the 3rd international workshop on distributed statistical computing*. Vienna, Austria., vol. 124, p. 10.

Rozen-Rechels, D., te Beest, M., Dew, L.A., le Roux, E., Druce, D.J. & Cromsigt, J.P. (2017). Contrasting impacts of an alien invasive shrub on mammalian savanna herbivores revealed on a landscape scale. *Diversity and Distributions*, 23, 656–666.

Schwarz, C. & Seber, G.A.F. (1999). A review of estimating animal abundance III. *Statistical Science*, 14, 427–456.

Schütz, E. (2019). Tamaraw verification surveys in the upper Amnay watershed region, Mindoro. *BULLtin*, 2, 6–15.

Shaffer, M.L. (1981). Minimum population sizes for species conservation. *BioScience*, 31, 131–134.

Sinclair, A.R.E. (1989). Population regulation in animals. In: *Ecological concepts : the contribution of ecology to an understanding of the natural world* (eds. Cherrett, J.M. & Bradshaw, A.D.). Blackwell Scientific Publications, Oxford, England., pp. 197–241.

Singer, F.J. & Norland, J.E. (1994). Niche relationships within a guild of ungulate species in Yellowstone National Park, Wyoming, following release from artificial controls. *Canadian Journal of Zoology*, 72, 1383–1394.

Smith, E. (1954). Quantitative aspects of population growth in the dynamic of growth processes (ej boel, ed.).

Steinmetz, R., Chutipong, W., Seuaturien, N., Chirngsaard, E. & Khaengkhetkarn, M. (2010). Population recovery patterns of southeast asian ungulates after poaching. *Biological Conservation*, 143, 42–51.

Stephens, P.A., Sutherland, W.J. & Freckleton, R.P. (1999). What is the allee effect? *Oikos*, pp. 185–190.

Stewart, G.B., Coles, C.F. & Pullin, A.S. (2005). Applying evidence-based practice in conservation management: lessons from the first systematic review and dissemination projects. *Biological Conservation*, 126, 270–278.

Sutherland, W.J., Pullin, A.S., Dolman, P.M. & Knight, T.M. (2004). The need for evidence-based conservation. *Trends in Ecology & Evolution*, 19, 305–308.

Traill, L.W., Bradshaw, C.J. & Brook, B.W. (2007). Minimum viable population size: a meta-analysis of 30 years of published estimates. *Biological conservation*, 139, 159–166.

Van Houtan, K.S., Halley, J.M., Van Aarde, R. & Pimm, S.L. (2009). Achieving success with small, translocated mammal populations. *Conservation Letters*, 2, 254–262.

Wilson, D. & Mittermeier, R. (2011). *Handbook of the mammals of the world. Vol. 2. Hoofed Mammals*. Lynx Edicions, Barcelona.

## TABLE CAPTIONS

**Table 1** Annual growth rate (*r*) of tamaraw abundance as estimated at each vantage counts in Mounts Iglit-Baco Natural Park located on
Mindoro island, Philippines, between 2006 and 2019. We used a Bayesian state-space model to account for sampling variance in animal
counts and reported the posterior mean estimation and its associated 95% credible interval (CI). Note the marked spatial variation in the
temporal trend of tamaraw abundance, ranging from a rapid decline at Tarzan to a marked increase at Bato Fidel or Inubon (see also Fig 3).

**Table 2** Estimated annual growth rates (*r*) of some large *bovinae* populations with body size comparable to the endangered tamaraw
(*Bubalus mindorensis*). The observed population growth rate of tamaraws at Mounts Iglit-Baco Natural Park, Mindoro, Philippines,
between 2000 and 2019 of $r = 0.06$ lies at the lower range of reported values.

**Table 1**

| Vantage point | r | 2.5% | 97.5% |
|---|---|---|---|
| Loibfo | 0.12 | −0.01 | 0.25 |
| Magawang | 0.12 | −0.02 | 0.24 |
| Bayokbok | 0.12 | −0.01 | 0.25 |
| Bato Fidel | 0.15 | 0.01 | 0.27 |
| Inubon | 0.15 | 0.02 | 0.28 |
| Mibluan | 0.13 | −0.01 | 0.26 |
| Nagbobong | 0.02 | −0.14 | 0.19 |
| Fangandatan | −0.03 | −0.24 | 0.14 |
| Anyayos | 0.08 | −0.06 | 0.21 |
| Lanas I | 0.10 | −0.05 | 0.24 |
| Lyan | −0.02 | −0.24 | 0.16 |
| Tarzan | −0.18 | −0.42 | 0.02 |
| Talafu East | 0.06 | −0.10 | 0.20 |
| Talafu West | 0.07 | −0.08 | 0.21 |
| Malitwang | 0.08 | −0.08 | 0.22 |
| Lanas II | 0.02 | −0.13 | 0.17 |
| Saligi East | −0.18 | −0.45 | 0.04 |
| Malibayong | 0.02 | −0.16 | 0.19 |
| Saligi East | −0.46 | −2.98 | 0.17 |

**Table 2**

| Species | Location | *r* | Reference |
| --- | --- | --- | --- |
| *Bison bison* | Mackenzie Bison Sanctuary, Canada | 0.21 | Gates & Larter (1990) |
| *Bison bison* | Mackenzie Bison Sanctuary, Canada | 0.20 | Larter *et al.* (2000) |
| *Bison bison* | Yellowstone, Wyoming, USA | 0.07 | Fuller *et al.* (2007) |
| *Bison bison* | Yellowstone, Wyoming, USA | 0.30 | Singer & Norland (1994) |
| *Bison bonasus* | Białowieża, Poland | 0.07 | Mysterud *et al.* (2007)[1] |
| *Bison bonasus* | Białowieża, Poland | 0.18 | Krasiński (1978)[1] |
| *Bos gaurus* | Thung Yai Naresuan Sanctuary, Thailand | 0.31 | Steinmetz *et al.* (2010) |
| *Bos javanicus* | Not applicable, model prediction | 0.32 | Hone *et al.* (2010) |
| *Bos javanicus* | Cobourg peninsula, Australia | 0.00 | (Choquenot 1993) |
| *Bos taurus* | commander la ref !! | 0.36 | Smith (1954) |
| *Boselaphus tragocamelus* | Kanha National Park, India | 0.18 | Mathur (1991) |
| *Boselaphus tragocamelus* | Keoladoe National Park, India | 0.02 - 0.06 | Haque (1990) |
| *Bubalus arnee* | | 0.02 - 0.06 | Khatri et al. (2012) |
| *Bubalus arnee* | Koshi Tappu Wildlife Reserve, Nepal | 0.03 | Heinen & Paudel (2015) |
| *Bubalus bubalis* | | 0.03 | Heinen (1993) |
| *Bubalus bubalis* | | ? | Freeland and Boulton (1990) |
| *Bubalus mindorensis* | Mount Iglit-Baco National Park, Philipines | 0.06 | This study |
| *Pseudoryx nghetinhensis* | Vietnam & Laos | 0.05 | Kemp *et al.* (1997)[b] |
| *Syncerus caffer* | Hluhluwe-iMfolozi Park, South Africa | 0.12 | Jolles (2007) |
| *Syncerus caffer* | Zakouma National Park, Chad | 0.08 | Cornélis *et al.* (2014) |
| *Taurotragus oryx* | Lombard Nature Reserve, South Africa | 0.12 | Buys & Dott (1991)[c] |
| *Taurotragus oryx* | South Africa | 0.16 | Van Houtan *et al.* (2009) |
| *Taurotragus oryx* | Kruger National Park, South Africa | −0.02 | Nicholls *et al.* (1996) |
| *Taurotragus oryx* | Masai Mara Ecosystem, Kenya | −0.11 - 0.07 | Ottichilo *et al.* (2000) |
| *Taurotragus derbianus* | Senegal | 0.31 | Koláčková *et al.* (2011) |

**FIGURE CAPTIONS**

2  **Figure 1**  General location of the Mounts Iglit-Baco Natural Park on Mindoro

island, Philippines. The current geographical range of the endangered tamaraw

4  (*Bubalus mindorensis*) population is limited to a 2,500ha area (salmon color

area) of which 1,250ha consists in a no-hunting zone (red dotted line) since

6  2016, hence offering the highest level of protection for the species. Every year

since 2000, taramaw counts have been carried out in March–April on 18

8  different vantage points (green circles) for the purpose of monitoring population

abundance trends.

10  **Figure 2**  Time series of tamaraw (*Bubalus mindorensis*) abundance in Mounts

Iglit-Baco Natural Park on Mindoro island, Philippines, from years 2000 to 2019.

12  Raw counts (black solid line) and counts corrected for sampling variance (blue

solid line) are presented with the associated 95% credible interval. Over the 13

14  years of monitoring, tamaraw abundance increased at a rate of 6% per year on

average.

16  **Figure 3**  Spatial variation in the local growth rate of tamaraw (*Bubalus

mindorensis*) abundance in Mounts Iglit-Baco Natural Park on Mindoro island,

18  Philippines, from years 2000 to 2019. With a general positive growth since 2000,

the situation in the core area of the monitoring zone is in sharp contrast with the

20  important decrease in abundance observed at the periphery. Movement of

animal toward less disturbed areas or illegal hunting outside of the no-hunting

22  agreement zone could account for this marked spatial structure of the tamaraw

population dynamics.

**Figure 1**

**Figure 2**

# Figure 3