Cast away on Mindoro island: population dynamics of the critically endangered tamaraw (*Bubalus mindorensis*) at Mounts Iglit–Baco Natural Park

C. Bonenfant[1], A. Rutschmann[2], J. Burton[3], R. Boyles[4], and E. Schütz[5]

1 Université de Lyon, F-69 000, Lyon ; Université Lyon 1 ; CNRS, UMR 5558, Laboratoire "Biométrie et Biologie Évolutive", F-69 622, Villeurbanne, France.
2 School of Biological Sciences, University of Auckland, Auckland, New Zealand
3 IUCN SSC Asian Wild Cattle Specialist Group, Global Wildlife Conservation, Texas, USA
4 Department of Environment and Natural Resources of the Philippines, Occidental Mindoro
5 D'Aboville Foundation and Demo Farm Inc, Manila, Philippines

Keywords

counts, density-dependence, exponential model, Mindoro dwarf buffalo, protection measures, poaching, population growth.

Correpondence

Christophe Bonenfant, `christophe.bonenfant@univ-lyon1.fr` – ORCID 0000-0002-9924-419X

Abstract

Endangered species, despite living at low population density, may undergo density-dependent feed-backs in case of successful recovery or marked reduction in range. When at work, density-dependence dynamics increases extinction risks and hamper conservation efforts. The largest population of the critically endangered tamaraw (*Bubalus mindorensis*) lives on a $< 3,000$ha in Mounts Iglit-Baco Natural Park (MIBNP) with very limited expansion possibilities. Tamaraw abundance has been monitored on a yearly basis from animal counts by Philippine authorities since the year 2000. Consistent with its protected status by law, the MIBNP tamaraw population has been increasing in size at an average rate of $+6\%$ per year, which we found to be relatively low compared to other similar-sized *Bovinea* species. Population growth was strikingly spatially structured within MIBNP with a population growth close to $+16\%$ in the core area of protection, while a reduction of abundance of -27% was measured at the periphery of the species range inside MIBNP. Highly concerning is the fact that the annual population growth rate progressively decreased significantly over the years since 2008, which we interpreted as an evidence of density-dependence. This isolated tamaraw population is currently experiencing a contraction of its range in MIBNP, likely caused by anthropogenic pressures forcing large herbivores to live at relatively high density in the core zone of the monitoring where protection is most effective. Our study highlights that beyond the encouraging results of a continuous growth over the last two decades, the MIBNP tamaraw population remains subject to uncertainty of its long term viability.

INTRODUCTION

In conservation planning, population abundance is critical and focuses the attention of field biologists and wildlife management institutions. Low population abundance is considered as one of the foremost criteria used for the evaluation of conservation statuses (Mace *et al.* 2008; Neel *et al.* 2012) . The International Union for Conservation of Nature (IUCN) would for instance classify any species with very small population sizes as threatened, and so without evidence for past, present or future decline (Mace *et al.* 2008). Also most populations of endangered wild species are indeed isolated and fragmented, and hence are of small size. The repeated focus on abundance in conservation biology has its roots in the fact that population viability strongly depends on the number of animals in a population (Lande *et al.* 2006). Overall, the ecology of small populations is particularly relevant to the genetics, evolution and demography of threatened species (Korn 1994).

In terms of populations dynamics, small populations differ in several ways from what is generally reported at higher abundance (Caughley 1994; Mugabo *et al.* 2013). When population size is small, the random death of a few individuals may precipitate the extinction of a population, a process referred to as demographic stochasticity (Shaffer 1981; Lande 1993). Similarly, when mating partners are too few to meet and reproduction fails, demographic stochasticity generates the so-called Allee effects (Courchamp *et al.* 1999), that is a decrease of the population growth at low population abundance (*i.e.* demographic component, see Stephens *et al.* (1999)). Yet vulnerable populations can also suffer from classical density-dependent effects whereby population

growth rate decreases with animal density (Nicholson 1933), driving the fate and

viability of populations at risk of extinction (Beissinger & Westphal 1998; Henle

et al. 2004) . The classical density-dependent effects on population dynamics

and demographic rates are often overlooked in conservation because they are *a*

priori expected to occur mainly at high population abundance. Although

counter-intuitive for threatened populations, increasing abundance could,

however, occur incidentally following a range contraction caused by habitat loss,

the continuous and long-term reintroduction of individuals for reinforcement, or

following successful protection measures. High abundance can potentially

trigger undesirable ecological and demographic consequences for its

conservation.

Density-dependence is a pervasive demographic responses among large

mammals (Fowler 1987; Bonenfant *et al.* 2009). In addition to depressing

recruitment rates rapidly, mortality of adults may increase as population density

approaches the carrying capacity (Eberhardt 1977; 2002). Many changes in

behaviour and life history traits of individuals account for those observed

changes in demographic rates (Bonenfant *et al.* 2009; Choquenot 1991; Van der

Wal *et al.* 2014). With rising density, individuals die from starvation because of

reduction in the availability and accessibility of food resources by conspecifics.

Agonistic interactions between individuals generate high stress levels which, in

turns, magnify the effect of other mortality sources (Peterson & Black 1988).

For instance, at high population density, epizootics are more likely to emerge and

to spread quickly in populations (). Higher densities may also force

non-dominant animals to occupy low-quality territories, ultimately decreasing

the average productivity and survival of the population (Krüger & Lindström

4

2001). Finally, because high population density exacerbates the effect of

2 environmental variation on demographic rates (Hones & Clutton-Brock 2007),

the risks of massive mortality events are much higher if the population is

4 maintained in this state on the long term. Given its numerous biological

consequences, an early detection of density-dependence is of the upmost

6 importance for the management and conservation of animal species to limit the

risk of catastrophic events.

8   The tamaraw (*Bubalus mindorensis*) is a large mammalian herbivore endemic

to the Island of Mindoro (Heude 1888; Custodio *et al.* 1996). Despite the

10 tamaraw being officially protected by law in 1954 and the creation of the Mts

Iglit-Baco Natural Park (MIBNP) protected area in 1970 (Maala 2001), its

12 distribution range has been dramatically shrinking over the last decades.

Originally found across the whole island of Mindoro with a gross estimate of

14 10,000 individuals in 1900 (Long *et al.* 2018), the tamaraw has suffered from

intensive land conversion for agriculture and logging industry, trophy hunting,

16 disease outbreaks spread out by domestic cattle (Maala 2001), and traditional

hunting conducted by upland indigenous communities who share their living

18 space with the large herbivore (Long *et al.* 2018). As a consequence, the species

is now found in a few isolated populations scattered across Mindoro, with

20 MIBNP hosting the largest number (Matsubayashi *et al.* 2010; Wilson &

Mittermeier 2011; Long *et al.* 2018). The MIBNP tamaraw population has been

22 restricted to one location of less than <3,000 ha in 2018, that we referred to as

the Core Zone of the Monitoring (CZM, see Fig. 1).

24   Given its critical importance for the conservation of this iconic species, the

MIBNP tamaraw population has been the cornerstone of all past, present and

5

future conservation plans (Maala 2001; Long *et al.* 2018). A better knowledge

about its ecology and population dynamics is fundamental to assess its current

conservation status, to strengthen protection measures and to guide future

conservation strategy with evidence-based information (Caughley 1994; Norris

2004; Stewart *et al.* 2005; Sutherland *et al.* 2004). Unfortunately, the population

dynamics of tamaraws in MIBNP has not been documented so far, nor do we

have any estimation of its average population growth rate. Yet it is likely that

density-dependence of demographic parameters will affect the population in the

near future because hunting and human disturbance prevent this population from

expanding outside of the CZM.

Here we took advantage of a 19 year-long time series of abundance data to

present the first detailed analysis of the tamaraw population dynamics in the

CZM of the MIBNP. We test the hypothesis that this population of large

herbivores could be facing negative effects of density. We predict (*i*) that because

of the substantial conservation effort in the last decade, overall population

growth rate (noted *r*) should be statistically > 0. Yet (*ii*), because the no-hunting

agreement area creates a spatial demarcation between risky and safe habitats, we

expect local growth rates to differ in space and predict a decrease of r while

getting away from the main rangers' base camps and patrolling routes. Finally

(*iii*), given the absence of direct threats, we predict that the MIPBN tamaraw

population could be approaching the carrying capacity of the CZM and expect a

negative association between *r* and tamaraw abundance within the NHZ.

MATERIAL AND METHODS

Study site

The study area is located within the Mounts Iglit-Baco Natural Park (N12°54′ E121°13′) in the south-central part of Mindoro which, with 106,665 ha, is the island largest protected area. MIBNP shelters the largest number of tamaraw in a single site located on the south-west edge of the protected area (Fig. 1). Tamaraw distribution range is now restricted to less than 3,000 ha, which represents approx. 3% the MIBNP surface. This area is the Core Zone of the Monitoring because this is where most of the patrolling and observations are carried out by the Park's officers. There are currently 15 park rangers divided into two or three teams to monitor the area on a regular basis, following regular distinct patrolling routes centered around the three base camps.

In an attempt to reinforce tamaraw protection, wildlife managers came to a much needed agreement with residing indigenous people in 2016, declaring a 1,600 ha no-hunting zone area (NHZ) within the CZM (Fig. 1). An unforeseen consequence of the NHZ establishment for tamaraw seems to be an increase of pressure by traditional hunting at its direct boundaries. MIBNP rangers regularly report setting of snare and spear traps, and violation of the NHZ agreement in the last year is not uncommon with several deaths of tamaraws confirmed (Boyles comm. pers.). Another possible response of large herbivores to hunting is space use modification to reduce the risk of being killed (Chassagneux *et al.* 2019; Marantz *et al.* 2016).

The CZM area is a rolling grassland plateau with an average elevation of 800m a.s.l., dominated by cogon (*Imperata cylindrica*) and wild sugarcane

(*Saccharum spontaneum*), and interspersed with numerous wooded creeks,

secondary forest fragments and steep hills. In recent years, the invasive Siam

weed (*Chromolaena odorata*, locally called hagonoy) has been quickly

spreading in the grassland and now covers an unknown fraction of the tamaraw

habitat. The tropical climate is strongly seasonal with a rainy season spanning

from June to October (mean temperature 26.5°C; mean rainfall: 150 mm), and a

hot dry season from November to May (mean temperature 29.8°C; mean rainfall:

5 mm). Apart from humans, the only potential predator to the tamaraw is the

reticulated python (*Malayopython reticulatus*) that is capable of preying upon

calves or yearlings but no such observation have been made so far.

Tamaraw counts

Every year since 2000, the Tamaraw Conservation Program (TCP), a high profile

program of the Philippine Department of Environment and Natural Resources

(DENR), has been conducting tamaraw counts across the CZM in MIBNP,

yielding an annual index of relative population abundance. The tamaraw

counting area consists in 18 vantage points, covering nearly 2,200 ha within the

CZM (Fig. 1). The method is rather invasive for the MIBNP ecosystem because

it involves the burning of grasslands a few weeks ahead of each count, in order to

increase visibility and to attract tamaraws to areas with highly palatable and

attractive regrowing grasses. Each year, between 1,200 and 1,500 ha of

grasslands were intentionally burned for the purpose of monitoring tamaraw

abundance.

Individual tamaraws were counted simultaneously at the 18 vantage points

for two hours at dusk and dawn, and for four consecutive days, in late March or

early April. For the 8 sessions and at each vantage points, at least two observers

spotted and recorded all animal seen on field paper sheets, writing down

observation time, approximate location, and assessed sex and age-category of

individuals (split into calves, yearlings, sub-adults and adults). Right after the

completion of the 8 sessions, count data were cross-checked and cleaned for

obvious double observations, and the total number of different animals seen was

then derived for each vantage point s and year t. These annual counts make the

only available source of information about tamaraw population trends and

dynamics for officials and conservation agencies.

Data analyses

For populations of long-lived and endangered species, robust and accurate

estimations of r remains difficult to obtain. Knowledge about r requires either

the estimation of demographic rates combined into a Leslie matrix (Leslie 1945)

or the availability of long time series of population abundance, preferably on an

annual basis. In the absence of natural marking, capturing individuals of an

endangered species to make them identifiable with tags often face strong and

justified resistance for ethical reasons, mostly because of unavoidable mortality

risks. Consequently, in the best case the monitoring of the population abundance

of endangered species relies on an estimator accounting for detection probability

of animals (*e.g.* Kéry & Schmidt 2008), but most of time a simple index of

abundance is carried out.

Working with abundance index such as annual counts to derive an empirical

estimates of r however comes with some serious methodological issues (Îto

1972; Lindley 2003). Over the last decades, population counts have repeatedly

9

been associated with a large sampling variance (*e.g.* Caughley 1977), and

2 coefficient of variation for population abundance as high as 30% were reported

for species 1, species 2 for instance. From a conservation biologist viewpoint, a

4 large sampling variance in the input data for estimating *r* is a major pitfall

because of the over-optimistic results it leads to. Both the point estimate and its

6 precision are overestimated in the presence of sampling variance (Lindley 2003).

Despite the availability of appropriate statistical methods to account for the

8 sampling variance of population counts (Kalman filter, De Valpine & Hastings

2002), such tools remain seldom used in practice for conservation.

10   Here we analyzed the number of observed tamaraws and judged as different

individuals by observers for each of the 18 vantage points, split by years ($C_{s,t}$,

12 where indice $s \in \{1, \dots, 18\}$ stands for the vantage point, and

$t \in \{2000, \dots, 2019\}$ stands for the time in years. We implemented a state-space

14 model to tease apart process from sampling variances, making the assumption

that double and under-counts compensate and are randomly distributed from

16 year to years. We worked in a Bayesian framework (*e.g.* Kéry & Schaub 2011)

but note this is fully equivalent to a Kalman filter with a frequentist approach. To

18 do so we defined $N_{s,t}$ as the unobserved 'true' abundance linked to $C_{s,t}$ by a

random effect corresponding to the process variance (σ_c^2), on the log scale to

20 ensure positive values for abundance. Our baseline model was:

$$\log(C_{s,t}) \sim \mathcal{N}(\log(N_{s,t}), \sigma_c^2)$$

Accordingly, we computed the population growth rate $r_{s,t}$ of the tamaraw

22 population at each vantage points *s*, as:

$$r_{s,t} = \log\left(\frac{N_{s,t+1}}{N_{s,t}}\right),$$

from counts corrected for the sampling variance, hence returning an unbiased

estimation of the population growth rate. The overall population growth of the

tamaraw population of MIBNP is then given by:

$$\bar{r} = \frac{1}{19}\sum_{t=1}^{19}\log\left(\frac{N_{t+1}}{N_t}\right),$$

whereby $N_t = \sum_{s=1}^{18} N_{s,t}$. We informed N_t with the total number of animals seen

each count session (C_{tot}) and linked counts at the VP and population levels with

the following observation equation: $C_{tot} \sim \mathcal{N}(N_t/p, \sigma_{tot}^2)$. The consolidated

number of tamaraws C_t being much smaller than total number seen $C_t ot$ from

which double-counts were removed, we included a constant proportion

parameter p, similar to a "detection rate" (which is not). By doing so, we could

use the available information about tamaraw counts fully, including both the

knowledge of experienced rangers and the raw number of detected tamaraws

with no *a priori* interpretation. In addition, we could input missing values at the

VP level for years 2000 to 2003 and 2005 thanks to the C_{tot} constrain on N_t.

To test our third hypothesis we investigated potential density-dependence in

the annual growth rate by fitting a logistic growth model to the total number of

tamaraws seen per year N_t:

$$\log(N_{t+1}) = \log(N_t) + r_m \times \left(1 - \frac{N_t}{K_l}\right),$$

11

where r_m is the maximum population growth for the tamaraw, and K is the

carrying capacity of the MIBNP. Likewise, we tested if the annual population

growth rate would decrease in time with population density by fitting a

Gompertz model:

$$\log(N_{t+1}) = \log(N_1) + \log\left(\frac{K_G}{N_1}\right) \times \left(1 - e^{-\beta \times t}\right),$$

Because the pair of parameters r_m and K_l (logistic), and β and K_G

(Gompertz), are not fully separately identifiable (Lebreton & Gimenez 2013), we

used informative priors for r_m and β based on the reported maximum population

growth for other large bovids (see Traill *et al.* 2007, for a similar approach).

Accordingly, we set a moderately informative prior value of mean 0.3 and a

precision of 0.1 in our models (Table 2). We assessed the different models by

computing explicitly, during the model fit, the probability for our abundance

time series to be generated by each of the three demographic models

(exponential, Gompertz and logistic).

We fitted our statistical model to the tamaraw count data with JAGS 4.0

(Plummer *et al.* 2003), with 3 MCMC chains and a burn-in of 40,000 iterations.

We obtained parameter estimates with an additional 30,000 iterations and a

thinning factor of 5 (hence a total of 6,000 MCMC samples). We checked model

convergence graphically by investigating the mixing of MCMC chains, along

with the $\hat{R}$ statistics (Brooks & Gelman 1998). With the exception of r_m (see

above), we set uninformative priors with large variance for all other parameters

to be estimated, and checked the sensitivity of our results to the average value of

the priors by replicating the analyses with variable mean and initial values for

12

prior distributions. We report all parameters estimates as the mean of the
posterior distributions along with the 95% percentile for the credible intervals
following Louis & Zeger (2008): $_{95\%low}\,estimate\,_{95\%up}$.

   Note that despite the fact that we analysed the number of seen tamaraws as
described above, we chose not to display such numbers on figures and to report a
standardized abundance ($C_{z,t}/max(C_{z,t})$) instead. This transformation applies to
the estimated carrying capacities of the Gompertz and logistic models,
$K/max(C_{z,t})$, hence reported values for K are > 1. We did not provide raw
counts to avoid a likely confusion between the index of relative tamaraw
abundance we worked with, and a real population size estimator that was not
implemented at MIBNP. Such a variable transformation has no incidence on the
biological interpretation of the results though.

RESULTS

From our baseline model of tamaraw counts, we estimated an average annual
growth rate of $\bar{r} = {}_{0.02}\,0.06\,_{0.12}$ for the MIBNP population from 2000 to 2019,
which associated variance process equals $\sigma_{proc} = {}_{0.25}\,0.37\,_{0.49}$ (Fig. 2). As
expected for count data, the sampling variance was large $\sigma_c = {}_{7.32}\,9.31\,_{9.98}$
suggesting frequent double counts of the same individuals. That the MIBNP
tamaraw population increased in size by c.a. 6% per year and was statistically
different from 0 is a clear support of our first hypothesis predicting increase of
abundance thanks to successful protection efforts at MIBNP.

   Considering the vantage point specific growth rates (r_s), we observed a
marked variability in space (Table 2; Fig. 3) considering that growth rates ranged

13

between $r_s = {}_{-2.52} -0.32 {}_{0.20}$ at Tarzan VP and $r_s = {}_{0.03} 0.19 {}_{0.35}$ at Inubon VP.

Vantage point specific growth rates were spatially structured at MIBNP (Fig 3).
With no exception, $r_s < 0$ were located at the periphery of the NHZ (Iyan,
Tarzan, Saligue East, Mibluan, Fangandatan) although r_s of a few vantage points
was > 0 despite being located on the NHZ limit too (Nagbobong, Malibayong,
Talafu West and East). Together these results are in accordance with our
hypothesis of an on-going contraction of the tamaraw range at MIBNP after
2000, and the disappearance of individuals at the periphery of the CZM.

Analytical evidence for density-dependence in the tamaraw dynamics
between year 2008 and 2019 was moderate to high. We explicitly estimated the
probability π that the exponential, Gompertz and logistic models would be the
underlying process generating the observed variation in tamaraw abundance
(proxy of Baye's factor). Our different models did have different probability to
have generated to observed data with, respectively $\pi = 0.80$, $\pi = 0.00$ and
$\pi = 0.20$ for the exponential, Gompertz and logistic model respectively. The
estimated coefficients of the Gompertz model lent statistical support for the
hypothesis of a gradual decrease in the annual population growth with time
($\beta = {}_{0.07} 0.09 {}_{0.11}$), and estimated carrying capacity to $K_G = {}_{1.12} 1.33 {}_{1.64}$.
Similarly, point estimates for the two parameters of the logistic growth function
were $r_m = {}_{0.09} 0.11 {}_{0.13}$ for the maximum population growth rate and
$K_l = {}_{1.14} 1.35 {}_{1.61}$ for the carrying capacity, very close to K_G estimates.

DISCUSSION

The isolated tamaraw population in MIBNP has been increasing in size since 2000 (Fig. 2). Undoubtedly, for such an highly endangered species a positive growth rate of 0.06 over 19 years of monitoring is a great success, both locally and for the global conservation effort. Yet, the devil lies in the details though, and two additional results darken this otherwise positive picture of tamaraw population dynamics and conservation. First, our results highlight a progressive decrease of the average growth rate, suggesting density-dependent effects at the population level. Furthermore, we found support for the hypothesis of a contraction of the tamaraw distribution, evidenced by a clear spatially structured pattern of the population growth rates. The population dynamics we document is clearly calling for new and urgent conservation actions if the future of tamaraw is to be ensured.

Long-run Positive Growth of Tamaraws

The average population growth rate of $\bar{r} = 0.06$ we found gives strong evidence in favor of our hypothesis of successful conservation policies of tamaraw over the past two decades. Yet, an annual growth of 6% is relatively low compared to other similar-sized or even larger cattle (Table 2). For instance, a population of African buffalo (*Syncerus caffer*) in Southern Africa grew at a rate of 12% per year for 28 years (Jolles 2007), and values close to 30% per year have regularly been recorded for bison (*Bison bison*) or banteng (*Bos javanicus*) despite these species are between 2 and 2.5 times larger in size than the tamaraw. From this brief comparative approach, we could expect a substantially higher average

15

growth rate for tamaraws if the population was in a colonizing phase. Given the

tamaraw is threatened, our results suggest that specific environmental factors are

limiting its dynamic in MIBNP.

A first limiting factor could stand in the fact that the most available and

accessible biotope for the tamaraw in MIBNP is grassland (83% of the CMZ)

dominated by strong pioneer competitive species such as cogon and wild

sugarcane; both grasses becoming *de facto* prominent in the tamaraw diet. While

providing tamaraws with abundant and high quality forage at the early stages of

plant growth, the grassland could rapidly become of much lower nutritive value

during the dry season. Once grass growth is complete, stems exceeds tamaraw

height (300cm vs. < 120 cm respectively) and dry out to become unpalatable,

raising the question of food resource accessibility and availability for a large

population of tamaraws.

Dry vegetation also questions the ambivalent role of fire regime. On the one

hand, regular burning rejuvenate vegetation and promote regeneration of

attractive new grass shoots few months ahead of the rainy season, when animals

might look for alternative marginal habitats such as forest or creeks for

complementary feeding purposes. On the other hand, previous observation

suggest that regular fire could promote Siam weed expansion (Nath *et al.* 2019),

which is indeed rapidly spreading on the study site. Like other grazing

herbivores (Rozen-Rechels *et al.* 2017), tamaraw avoid this invasive plant, an

issue that could ultimately decrease the CMZ carrying capacity (Dumalisile &

Somers 2017). More generally tamaraw feeding preference is not fully

understood and the species might be a mixed-feeding or a browsing herbivore

(*sensu* Hofmann 1989). While being *de facto* restricted on grassland habitats,

16

tamaraw are suspected to be ecologically flexible, as recent assessment of a new

tamaraw population in mountain forest habitats in the upland inner Mindoro

Island would suggest (Schütz 2019). In other words, the CZM might actually

provide ecological conditions whose intrinsic limits are artificially reduced, so

far, thanks to long lasting and highly disturbing habitat management

policies;detrimental impact starting progressively to exceed beneficial effects.

Possible Impacts of Human Disturbance and Hunting

An additional factor, related to human activities, could also account for the low

tamaraw population growth rate we report. Tamaraw has always been

traditionally hunted by IPs and despite agreement of mutual effort of

conservation, both intentional and unintentional casualties still happen. Besides

tamaraw is sought after for bushmeat by lowlander poachers. Both threats are

hard to control in the field, are likely pervasive across MIBNP, and lead to an

increased mortality of all age-classes of tamaraws to an unknown extent.

Positive growth over the past two decades is a clear sign of successful protection

measures thanks to the constant presence of rangers and the recurrent patrols that

have likely contributed to decrease harvesting pressure within the CZM.

However, the marked spatial heterogeneity we document on the observed growth

rates with negative values at the periphery of the counting area might reveal a

lower protection efficiency when moving away from the main patrolling routes

and base camps. Actually, the CZM area functions like a source-sink system

with central areas nearby base camps concentrating most of the vantage points

with positive growth rates, with local value as high as 16% (Table 1), more in

line with the average growth we review for large bovids (Table 2).

17

Space Needed for Sustaining Population Growth

2   At first glance, the gradual decrease of the tamaraw population growth rate over
the survey coupled with the overall abundance observed at MIBNP reflects the

4 signature of some sort of density-dependence: a change in demographic rates
lowering the number of individuals added to the population from year to year

6 (Fowler 1987; Sinclair 1989). Predicted growth rates from the lowest to the
highest abundance varied between $r_t = 0.11$ and $r_t = 0.01$ for tamaraw at

8 MIBNP. Detection of density-dependence in MIBNP is worrying because it is
usually the sign of increased competition between individuals for food or

10 territorial space. In other word, our results suggest that tamaraws could already
be suffering from food resource shortage. Similar to the tamaraw case is the

12 banteng (*Bos javanicus*) population of Alas Purwo National Park, Indonesia. In
spite of an increasing abundance, this population has become consolidated close

14 to an artificially maintained grazing area, which may be causing increased
predation pressure of juveniles, and the population dynamics and carrying

16 capacity are not yet known (JB, pers. obs).

In absence of safe corridors for individuals to safely emigrate from the CZM

18 and in the presence of adverse effects of Siam weed on forage quantity, we
cannot expect this population to keep on increasing in size for the next decade as

20 we report between 2000 and 2019. From current data, we could estimate a
carrying capacity of 1.4 meaning the current MIBNP tamaraw abundance has

22 already reached almost 2/3 of the carrying capacity. This explanation is the
biological hypothesis of density-dependence, which is well known and

24 repeatedly detected in many populations of large mammals (Bonenfant *et al.*

2009). Yet, in absence of trustful demographic rates estimations of tamaraws at

MIBNP we hit the limits of the so-called pattern-oriented approach (Krebs

2002), with no way to assert real changes in demographic rates with abundance.

Consequently, an alternative interpretation of the density-dependence pattern

that we report for tamaraw is possible, which we describe as the *count bias*

hypothesis. Visual counts of animals, as it is conducted in MIBNP, are

notoriously known to be inaccurate and prone to underestimation of the real

abundance (Andersen 1953; **?**) (Andersen 1953; Laake et al. 1999). Tamaraw

counts were not designed to be used with a population abundance estimator such

as returned by the distance sampling (Buckland *et al.* 2004) or capture-recapture

methods (Schwarz & Seber 1999), and therefore failed to account for the

imperfect detection of individuals. The proportion of undetected tamaraws could

increase with their overall abundance, a previously reported effect on red deer

(*Cervus elaphus* Garel *et al.* 2010) and roe deer (*Capreolus capreolus*, Pellerin et

al. 2018). Alternatively, different cognitive biases (Baddeley et al. 2004) could

generate similar patterns if, for instance, observers unconsciously pull the total

number of animals downward when consolidating count data. At this stage,

counts at year t could likely influence the results of counts at year $t + 1$,

assuming observers are conservative in their expectation of the tamaraw annual

population growth. The monitoring of additional biological variables in the field

such as the reproductive success of females could help at teasing apart the two

hypotheses (Morellet *et al.* 2007).

Implication for tamaraw conservation

Current conservation measures assume tamaraw population growth will continue in the next decade, without a need to mitigate increased pressures thought to be impacting the periphery of the CZM or to address the density dependence pressure on the population. Our results call for a adjustment to strategic planning for protection and management within MIBNP, as part of the MIBNP Management Plan and the island-wide Tamaraw Conservation Management and Action Plan (DENR 2019). Firstly, two main aspects must be urgently addressed: (*i*) increase the available space for the species so to mitigate density-dependence effect and maintain positive growth rate, that must be combined with (*ii*) increasing the effectiveness of ranger patrols in the periphery area of the CZM to reduce the recurrent poaching pressure from Filipino lowlanders and losses due to traditional hunting. Both targets request to collaborate closely with residing IP communities on a more sustainable land-use and traditional hunting system. Indeed, a more effective protection and suitable habitat management might in turn increase the average growth rate at MINBP, and exacerbate the observed density-dependence demographic responses of tamaraws. Secondly, when investigating the feasibility of translocating animals from this source population to other sites, estimated demographic rates should be incorporated into population modelling for extraction number that maintain a sustainable yield and the tamaraw population viability. Indeed, the maximum sustainable yield is found at $K/2$ in density-dependent models, not when population abundance reaches carrying capacity K.

Conclusion

Despite increased efforts of land conservation, range contractions and habitat fragmentation are still one of the most important threat to large animals population (Ripple *et al.* 2015). As a consequence, a greater proportion of wild herbivores (and their natural predators) are expected to be impacted by intensively managed landscapes. So far, conservation responses essentially consist in protecting large areas of land, a more than praiseworthy approach, that nevertheless have limits, as illustrated by our results. In any case, positive growth rates detected for a species in a conservation area could potentially hide a darker picture. First, the apparent growth of a population, trapped in a too small conservation area, can be the pervasive effect of range contraction rather than the effect of increased reproduction and survival. Second, because of this, the population could actually face important demographic challenges that, in the long term, could lead to new threats. In this context, there is an increasing need to understand the interaction between anthropogenic activities and the population dynamics of protected species. Clearly, the protection of large herbivores may hence come at the costs of undesirable ecological effects, rendering their conservation even more difficult. On the bright side, our findings also exemplifies how detailed analyses of relative population abundance data, even without absolute population size estimators, can contribute to determining future conservation measures and are essential for large mammal conservation.

ACKNOWLEDGMENTS

A lot of people are to thank here. We are grateful to Fernando Garcìa for producing maps used in this paper. Special thanks are extended to Alvaro Gonzalez. We must acknowledge the invaluable help of all rangers, students and volunteers who contributed to the annual tamaraw counts.

REFERENCES

Andersen, N.J. (1953). Analysis of a Danish roe deer population. *Danish Review of Game Biology*, 2, 121–155.

Beissinger, S.R. & Westphal, M.I. (1998). On the use of demographic models of population viability in endangered species management. *The Journal of wildlife management*, pp. 821–841.

Bonenfant, c., Gaillard, J.M., Coulson, T., Festa-Bianchet, M., Loison, A., Garel, M., Loe, L.E., Blanchard, P., Pettorelli, N., Owen-Smith, N., Du Toit, J. & Duncan, P. (2009). Empirical evidence of density-dependence in popualtions of large herbivores. *Adv. Ecol. Res.*, 41, 313–357.

Brooks, S.P. & Gelman, A. (1998). General methods for monitoring convergence of iterative simulations. *Journal of Computational and Graphical Statistics*, 7, 434–455.

Buckland, S.T., Anderson, D.R., Burnham, K.P., Laake, J.L., Borchers, D.L. & Thomas, L. (2004). *Advanced distance sampling*. Oxford University Press.

Buys, D. & Dott, H. (1991). Population fluctuations and breeding of eland *Taurotragus oryx* in a western Transvaal nature reserve. *Koedoe*, 34, 31–36.

Caughley, G. (1977). *Analysis of vertebrate populations*. Wiley, London. Graeme Caughley. 'A Wiley-Interscience Publication' Includes index. Bibliography: p.217-228.

Caughley, G. (1994). Directions in conservation biology. *Journal of animal ecology*, pp. 215–244.

Chassagneux, A., Calenge, C., Siat, V., Mortz, P., Baubet, E. & Saïd, S. (2019). Proximity to the risk and landscape features modulate female red deer movement patterns over several days after drive hunts. *Wildlife Biology*, 2019.

Choquenot, D. (1991). Density-dependent growth, body condition, and demography in feral donkeys: testing the food hypothesis. *Ecology*, 72, 805–813.

Choquenot, D. (1993). Growth, body condition and demography of wild banteng (*Bos javanicus*) on Cobourg Peninsula, northern Australia. *Journal of Zoology*, 231, 533–542.

Cornélis, D., Melletti, M., Korte, L., Ryan, S.J., Mirabile, M., Prin, T. & Prins, H.H. (2014). *African buffalo Syncerus caffer (Sparrman, 1779)*, Cambridge University Press, pp. 326–372.

Courchamp, F., Clutton-Brock, T. & Grenfell, B. (1999). Inverse density dependence and the allee effect. *Trends in ecology & evolution*, 14, 405–410.

Custodio, C.C., Lepiten, M.V. & Heaney, L.R. (1996). *Bubalus mindorensis*. *Mammalian Species*, pp. 1–5.

De Valpine, P. & Hastings, A. (2002). Fitting population models incorporating process noise and observation error. *Ecological Monographs*, 72, 57–76.

Dumalisile, L. & Somers, M.J. (2017). The effects of an invasive alien plant (*Chromolaena odorata*) on large African mammals. *Nature Conservation Research.* , 2, 102–108.

Eberhardt, L.L. (1977). Optimal policies for conservation of large mammals, with special references to marine ecosystems. *Environmental Conservation*, 4, 205–212.

Eberhardt, L.L. (2002). A paradigm for population analysis of long-lived vertebrates. *Ecology*, 83, 2841–2854.

Fowler, C.W. (1987). A review of density dependence in populations of large mammals. *Curr. Mammal.*, 1, 401–441.

Fuller, J.A., Garrott, R.A., White, P., Aune, K.E., Roffe, T.J. & Rhyan, J.C. (2007). Reproduction and survival of Yellowstone bison. *Journal of Wildlife Management*, 71, 2365–2372.

Garel, M., Bonenfant, C., Hamann, J.L., Klein, F. & Gaillard, J.M. (2010). Are abundance indices derived from spotlight counts reliable to monitor red deer *Cervus elaphus* populations? *Widlife Biology*, ??, ??

Gates, C. & Larter, N. (1990). Growth and dispersal of an erupting large herbivore population in northern Canada: the Mackenzie wood bison (*Bison bison athabascae*). *Arctic*, pp. 231–238.

Haque, N. (1990). *Study on the ecology of wild ungulates of Keoladeo national Park Bharatpur, Rajasthan*. Ph.D. thesis, Aligarh Muslim University.

Heinen, J.T. & Paudel, P.K. (2015). On the translocation of wild asian buffalo (**?**) in Nepal: are feral backcrosses worth conserving? *Conservation Science*, 3.

23

Henle, K., Sarre, S. & Wiegand, K. (2004). The role of density regulation in extinction processes and population viability analysis. *Biodiversity and Conservation*, 13, 9–52.

Heude, P. (1888). Études sur les ruminants de l'Asie orientale. cerfs des Philippines et de l'Indo-Chine. *Mémoires Concernant l'Histoire Naturelle de l'Empire Chinois par des Péres de la Compagnie de Jésus*, 2, 1–10.

Hofmann, R.R. (1989). Evolutionnary steps of ecophysiological adaptation and diversification of ruminants: a comparative view of their digestive system. *Oecologia*, 78, 443–457.

Hone, J., Duncan, R.P. & Forsyth, D.M. (2010). Estimates of maximum annual population growth rates (r_m) of mammals and their application in wildlife management. *Journal of Applied Ecology*, 47, 507–514.

Hones, J. & Clutton-Brock, T. (2007). Climate, food, density and wildlife population growth rate. *J. Anim. Ecol.*, ?, ?–?

Îto (1972). On the methods for determining density-dependence by means of regression. *Oecologia*, 10, 776–778.

Jolles, A.E. (2007). Population biology of african buffalo (*Syncerus caffer*) at Hluhluwe-iMfolozi Park, South Africa. *African Journal of Ecology*, 45, 398.

Kemp, N., Dilger, M., Burgess, N. & Van Dung, C. (1997). The saola *Pseudoryx nghetinhensis* in Vietnam – new information on distribution and habitat preferences, and conservation needs. *Oryx*, 31, 37–44.

Kéry, M. & Schaub, M. (2011). *Bayesian population analysis using WinBUGS: a hierarchical perspective*. Academic Press.

Kéry, M. & Schmidt, B. (2008). Imperfect detection and its consequences for monitoring for conservation. *Community Ecology*, 9, 207–216.

Koláčková, K., Hejcmanova, P., Antonínová, M. & Brandl, P. (2011). Population management as a tool in the recovery of the critically endangered Western Derby eland *Taurotragus derbianus* in Senegal, Africa. *Wildlife Biology*, 17, 299–311.

Korn, H. (1994). Genetic, demographic, spatial, environmental and catastrophic effects on the survival probability of small populations of mammals. In: *Minimum Animal Populations* (ed. Remmert, H.). Springer, pp. 33–49.

Krasiński, Z.A. (1978). Dynamics and structure of the european bison population in the białowieża Primeval Forest. *Acta Theriologica*, 23, 3–48.

Krebs, C.J. (2002). Two complementary paradigms for analysing populations dynamics. *Philosophical Transactions of the Royal Society of London, Series B*, 357, 1211–1219.

24

Krüger, O. & Lindström, J. (2001). Habitat heterogeneity affects population growth in goshawk *Accipiter gentilis*. *Journal of Animal Ecology*, 70, 173–181.

Lande, R. (1993). Risks of population extinction from demographic and environmental stochasticity and random catastrophes. *The American Naturalist*, 142, 911–927.

Lande, R., Engen, S., Sæther, B. & Coulson, T.N. (2006). Estimating density dependence from time series of population age structure. *American Naturalist*, 168, 76–87.

Larter, N., Sinclair, A., Ellsworth, T., Nishi, J. & Gates, C. (2000). Dynamics of reintroduction in an indigenous large ungulate: the wood bison of northern Canada. 3, 299–309.

Lebreton, J.D. & Gimenez, O. (2013). Detecting and estimating density dependence in wildlife populations. *Journal of Wildlife Management*, 77, 12–23.

Leslie, P.H. (1945). On the use of matrices in population mathematics. *Biometrika*, 33, 182–212.

Lindley, S.T. (2003). Estimation of population growth and extinction parameters from noisy data. *Ecological Applications*, 13, 806–813.

Long, B., Schütz, E., Appleton, M., Gonzales, A., Holland, J., Lees, C., Marandola, E., Natural, V.C.J., Pineda-David, M.T.J., Salao, C., Slade, J., Tan, E.H.P., Tiongson, L. & Young, S. (2018). Review of tamaraw (*Bubalus mindorensis*) status and conservation actions. *BULLtin*, 1, 18–34.

Louis, T.A. & Zeger, S.L. (2008). Effective communication of standard errors and confidence intervals. *Biostatistics*, 10, 1–2.

Maala, C.P. (2001). Endangered Philippine wildlife species with special reference to the Philippine eagle (*Pithecophaga jefferyi*) and tamaraw (*Bubalus mindorensis*). *Journal of International Development and Cooperation*, 8, 1–17.

Mace, G.M., Collar, N.J., Gaston, K.J., Hilton-Taylor, C., Akçakaya, H.R., Leader-Williams, N., Milner-Gulland, E.J. & Stuart, S.N. (2008). Quantification of extinction risk: Iucn's system for classifying threatened species. *Conservation biology*, 22, 1424–1442.

Marantz, S.A., Long, J.A., Webb, S.L., Gee, K.L., Little, A.R. & Demarais, S. (2016). Impacts of human hunting on spatial behavior of white-tailed deer (*Odocoileus virginianus*). *Canadian Journal of Zoology*, 94, 853–861.

Mathur, V.B. (1991). *The ecological interaction between habitat composition, habitat quality and abundance of some wild ungulates in India*. Ph.D. thesis, University of Oxford.

25

Matsubayashi, H., Boyles, R.M., Salac, R.L., Del Barrio, A., Cruz, L., Garcia, R.A., Ishihara, S. & Kanai, Y. (2010). Present status of tamaraw (*Bubalus mindorensis*) in Mt. Aruyan, Mindoro, Philippines. *Tropics*, 18, 167–170.

Morellet, N., Gaillard, J.M., Hewison, A.J.M., Ballon, P., Boscardin, Y., Duncan, P., Klein, F. & Maillard, D. (2007). Indicators of ecological change: new tools for managing populations of large herbivores. *J. Appl. Ecol.*, 44, 634—643.

Mugabo, M., Perret, S., Legendre, S. & Le Galliard, J.F. (2013). Density-dependent life history and the dynamics of small populations. *Journal of Animal Ecology*, 82, 1227–1239.

Mysterud, A., Bartoń, K., Jedrzejewska, B., Krasiński, Z., Niedziałkowska, M., Kamler, J., Yoccoz, N. & Stenseth, N. (2007). Population ecology and conservation of endangered megafauna: the case of european bison in białowieża Primeval Forest, Poland. *Animal Conservation*, 10, 77–87.

Nath, A., Sinha, A., Lahkar, B.P. & Brahma, N. (2019). In search of aliens: Factors influencing the distribution of *Chromolaena odorata* L. and Mikania micrantha Kunth in the Terai grasslands of Manas National Park, India. *Ecological engineering*, 131, 16–26.

Neel, M.C., Leidner, A.K., Haines, A., Goble, D.D. & Scott, J.M. (2012). By the Numbers: How is Recovery Defined by the US Endangered Species Act? *BioScience*, 62, 646–657.

Nicholls, A., Viljoen, P., Knight, M. & Van Jaarsveld, A. (1996). Evaluating population persistence of censused and unmanaged herbivore populations from the Kruger National Park, South Africa. *Biological Conservation*, 76, 57–67.

Nicholson, A.J. (1933). The balance of animal populations. *J. Anim. Ecol.*, S2, 132–178.

Norris, K. (2004). Managing threatened species: the ecological toolbox, evolutionary theory and declining-population paradigm. *Journal of Applied Ecology*, 41, 413–426.

Ottichilo, W.K., De Leeuw, J., Skidmore, A.K., Prins, H.H. & Said, M.Y. (2000). Population trends of large non-migratory wild herbivores and livestock in the Masai Mara ecosystem, Kenya, between 1977 and 1997. *African journal of Ecology*, 38, 202–216.

Peterson, C.H. & Black, R. (1988). Density-dependent mortality caused by physical stress interacting with biotic history. *American Naturalist*, 131, 257–270.

Plummer, M. *et al.* (2003). Jags: A program for analysis of bayesian graphical models using gibbs sampling. In: *Proceedings of the 3rd international workshop on distributed statistical computing*. Vienna, Austria., vol. 124, p. 10.

26

Ripple, W.J., Newsome, T.M., Wolf, C., Dirzo, R., Everatt, K.T., Galetti, M., Hayward, M.W., Kerley, G.I., Levi, T., Lindsey, P.A. *et al.* (2015). Collapse of the world's largest herbivores. *Science advances*, 1, e1400103.

Rozen-Rechels, D., te Beest, M., Dew, L.A., le Roux, E., Druce, D.J. & Cromsigt, J.P. (2017). Contrasting impacts of an alien invasive shrub on mammalian savanna herbivores revealed on a landscape scale. *Diversity and Distributions*, 23, 656–666.

Schwarz, C. & Seber, G.A.F. (1999). A review of estimating animal abundance III. *Statistical Science*, 14, 427–456.

Schütz, E. (2019). Tamaraw verification surveys in the upper Amnay watershed region, Mindoro. *BULLtin*, 2, 6–15.

Shaffer, M.L. (1981). Minimum population sizes for species conservation. *BioScience*, 31, 131–134.

Sinclair, A.R.E. (1989). Population regulation in animals. In: *Ecological concepts : the contribution of ecology to an understanding of the natural world* (eds. Cherrett, J.M. & Bradshaw, A.D.). Blackwell Scientific Publications, Oxford, England., pp. 197–241.

Singer, F.J. & Norland, J.E. (1994). Niche relationships within a guild of ungulate species in Yellowstone National Park, Wyoming, following release from artificial controls. *Canadian Journal of Zoology*, 72, 1383–1394.

Steinmetz, R., Chutipong, W., Seuaturien, N., Chirngsaard, E. & Khaengkhetkarn, M. (2010). Population recovery patterns of southeast asian ungulates after poaching. *Biological Conservation*, 143, 42–51.

Stephens, P.A., Sutherland, W.J. & Freckleton, R.P. (1999). What is the allee effect? *Oikos*, pp. 185–190.

Stewart, G.B., Coles, C.F. & Pullin, A.S. (2005). Applying evidence-based practice in conservation management: lessons from the first systematic review and dissemination projects. *Biological Conservation*, 126, 270–278.

Sutherland, W.J., Pullin, A.S., Dolman, P.M. & Knight, T.M. (2004). The need for evidence-based conservation. *Trends in Ecology & Evolution*, 19, 305–308.

Traill, L.W., Bradshaw, C.J. & Brook, B.W. (2007). Minimum viable population size: a meta-analysis of 30 years of published estimates. *Biological conservation*, 139, 159–166.

Van Houtan, K.S., Halley, J.M., Van Aarde, R. & Pimm, S.L. (2009). Achieving success with small, translocated mammal populations. *Conservation Letters*, 2, 254–262.

27

Van der Wal, E., Laforge, M.P. & McLoughlin, P.D. (2014). Density dependence in
social behaviour: home range overlap and density interacts to affect conspecific
encounter rates in a gregarious ungulate. *Behavioral Ecology and Sociobiology*, 68,
383–390.

Wilson, D. & Mittermeier, R. (2011). *Handbook of the mammals of the world. Vol. 2.
Hoofed Mammals*. Lynx Edicions, Barcelona.

TABLE CAPTIONS

Table 1 Annual growth rate (*r*) of tamaraw abundance as estimated at each vantage counts in Mounts Iglit-Baco Natural Park located on Mindoro island, Philippines, between 2006 and 2019. We used a Bayesian state-space model to account for sampling variance in animal counts and reported the posterior mean estimation and its associated 95% credible interval (CI). Note the marked spatial variation in the temporal trend of tamaraw abundance, ranging from a rapid decline at Tarzan to a marked increase at Bato Fidel or Inubon (see also Fig 3).

Table 2 Estimated annual growth rates (*r*) of some large *bovinae* populations with body size comparable to the endangered tamaraw (*Bubalus mindorensis*). The observed population growth rate of tamaraws at Mounts Iglit-Baco Natural Park, Mindoro, Philippines, between 2000 and 2019 of $r = 0.06$ lies at the lower range of reported values.

Table 1

Vantage point	r	2.5%	97.5%
Loibfo	0.12	−0.01	0.25
Magawang	0.12	−0.02	0.24
Bayokbok	0.12	−0.01	0.25
Bato Fidel	0.15	0.01	0.27
Inubon	0.15	0.02	0.28
Mibluan	0.13	−0.01	0.26
Nagbobong	0.02	−0.14	0.19
Fangandatan	−0.03	−0.24	0.14
Anyayos	0.08	−0.06	0.21
Lanas I	0.10	−0.05	0.24
Lyan	−0.02	−0.24	0.16
Tarzan	−0.18	−0.42	0.02
Talafu East	0.06	−0.10	0.20
Talafu West	0.07	−0.08	0.21
Malitwang	0.08	−0.08	0.22
Lanas II	0.02	−0.13	0.17
Saligi East	−0.18	−0.45	0.04
Malibayong	0.02	−0.16	0.19
Saligi East	−0.46	−2.98	0.17

Table 2

Species	Location	r	Reference
Bison bison	Mackenzie Bison Sanctuary, Canada	0.21	Gates & Larter (1990)
Bison bison	Mackenzie Bison Sanctuary, Canada	0.20	Larter *et al.* (2000)
Bison bison	Yellowstone, Wyoming, USA	0.07	Fuller *et al.* (2007)
Bison bison	Yellowstone, Wyoming, USA	0.30	Singer & Norland (1994)
Bison bonasus	Białowieża, Poland	0.07	Mysterud *et al.* (2007)[1]
Bison bonasus	Białowieża, Poland	0.18	Krasiński (1978)[1]
Bos gaurus	Thung Yai Naresuan Sanctuary, Thailand	0.31	Steinmetz *et al.* (2010)
Bos javanicus	Not applicable, model prediction	0.32	Hone *et al.* (2010)
Bos javanicus	Cobourg peninsula, Australia	0.00	Choquenot (1993)
Boselaphus tragocamelus	Kanha National Park, India	0.18	Mathur (1991)
Boselaphus tragocamelus	Keoladoe National Park, India	0.02 - 0.06	Haque (1990)
Bubalus arnee		0.02 - 0.06	Khatri et al. (2012)
Bubalus arnee	Koshi Tappu Wildlife Reserve, Nepal	0.03	Heinen & Paudel (2015)
Bubalus bubalis		0.03	Heinen (1993)
Bubalus bubalis		?	Freeland and Boulton (1990)
Bubalus mindorensis	Mount Iglit-Baco National Park, Philipines	0.06	This study
Pseudoryx nghetinhensis	Vietnam & Laos	0.05	Kemp *et al.* (1997)[b]
Syncerus caffer	Hluhluwe-iMfolozi Park, South Africa	0.12	Jolles (2007)
Syncerus caffer	Zakouma National Park, Chad	0.08	Cornélis *et al.* (2014)
Taurotragus oryx	Lombard Nature Reserve, South Africa	0.12	Buys & Dott (1991)[c]
Taurotragus oryx	South Africa	0.16	Van Houtan *et al.* (2009)
Taurotragus oryx	Kruger National Park, South Africa	−0.02	Nicholls *et al.* (1996)
Taurotragus oryx	Masai Mara Ecosystem, Kenya	−0.11 - 0.07	Ottichilo *et al.* (2000)
Taurotragus derbianus	Senegal	0.31	Koláčková *et al.* (2011)

FIGURE CAPTIONS

2   **Figure 1** General location of the Mounts Iglit-Baco Natural Park on Mindoro

island, Philippines. The current geographical range of the endangered tamaraw

4   (*Bubalus mindorensis*) population is limited to a 2,500ha area (salmon color

area) of which 1,250ha consists in a no-hunting zone (red dotted line) since

6   2016, hence offering the highest level of protection for the species. Every year

since 2000, taramaw counts have been carried out in March–April on 18

8   different vantage points (green circles) for the purpose of monitoring population

abundance trends.

10 **Figure 2** Time series of tamaraw (*Bubalus mindorensis*) abundance in Mounts

Iglit-Baco Natural Park on Mindoro island, Philippines, from years 2000 to 2019.

12 Raw counts (black solid line) and counts corrected for sampling variance (blue

solid line) are presented with the associated 95% credible interval. Over the 13

14 years of monitoring, tamaraw abundance increased at a rate of 6% per year on

average.

16 **Figure 3** Spatial variation in the local growth rate of tamaraw (*Bubalus

mindorensis*) abundance in Mounts Iglit-Baco Natural Park on Mindoro island,

18 Philippines, from years 2000 to 2019. With a general positive growth since 2000,

the situation in the core area of the monitoring zone is in sharp contrast with the

20 important decrease in abundance observed at the periphery. Movement of

animal toward less disturbed areas or illegal hunting outside of the no-hunting

22 agreement zone could account for this marked spatial structure of the tamaraw

population dynamics.

Figure 1

Figure 2

Figure 3