Cast away on Mindoro island: could density dependence threaten tamaraw (*Bubalus mindorensis*) on the long term?

C. Bonenfant[1], A. Rutschmann[2], J. Burton[3], R. Boyles[4], F. García[5], A. Tilker[6,7], and E. Schütz[5]

Université de Lyon, F-69 000, Lyon ; Université Lyon 1 ; CNRS, UMR 5558, Laboratoire "Biométrie et Biologie Évolutive", F-69 622, Villeurbanne, France.
School of Biological Sciences, University of Auckland, Auckland, New Zealand.
IUCN SSC Asian Wild Cattle Specialist Group, Global Conservation, Texas, USA.
Department of Environment and Natural Resources of the Philippines, Occidental Mindoro.
D'Aboville Foundation and Demo Farm Inc, Manila, Philippines.
Re:wild, Austin, Texas USD
Leibniz Institute for Zoo and Wildlife Research, Berlin, Germany.
5 D'ABOVILLE Foundation and Demo Farm Inc, Manila, Philippines.

Keywords

counts, exponential model, Mindoro dwarf buffalo, protection measures, poaching, population dynamics.

Correpondence

Christophe Bonenfant, `christophe.bonenfant@cnrs.fr` – ORCID 0000-0002-9924-419X

Abstract

Endangered species, despite often living at low population densities, may undergo unexpected density-dependent feedbacks in the case of successful recovery or marked reduction in range. Because density-dependence dynamics can increase risk of extinction, these effects can hamper conservation efforts. In this study, we analyse the dynamics of the largest population of the tamaraw (*Bubalus mindorensis*), a critically endangered species endemic to the Mindoro island, Philippines. The population is located within a $< 3,000$ ha area in Mounts Iglit-Baco Natural Park, with limited expansion possibilities. We took advantage of a 21 year time series of tamaraw counts to estimate annual population growth rate and possible density-dependence, accounting for sampling errors in the counts. The tamaraw population has been increasing in size at an average rate of $+5\%$ per year, as would be expected given its protected status by law. Population growth showed strong spatial structuring, with a population growth close to $+10\%$ in the core area of protection, and a reduction of abundance of -5% at the periphery of its range, inside the protected area. This range constriction is concerning because our best population dynamic model suggests significant negative density-dependence (Bayes factor = 0.9). The contraction of tamaraw range is likely caused by anthropogenic pressures forcing herbivores to live at relatively high densities in the core zone where protection is most effective. Our study highlights the fact that, despite the continuous population growth over the last two decades, the long-term viability of the Mounts Iglit-Baco Natural Park tamaraw population remains uncertain.

INTRODUCTION

Assessing population abundance is an important component of conservation planning because of the strong link between population size and likelihood of extinction (Boyce 1992). Accordingly, low population abundance is one of the primary criteria used for the evaluation of conservation status (Mace *et al.* 2008; Neel *et al.* 2012). Indeed, the effects of small population size on species persistence has been identified as one of the two central paradigms of conservation biology (Caughley 1994). Research has shown that the population dynamics of small populations differ in several ways from what is reported at higher abundance (Caughley 1994; Mugabo *et al.* 2013). When population size is small, demographic stochasticity – the random death of a few individuals – is more influential on the population dynamics and its viability than in populations that are larger in size (Shaffer 1981; Lande 1993). Similarly, when mating partners are too few to meet and reproduction fails, demographic stochasticity generates the so-called Allee effect (Courchamp *et al.* 1999), characterized by a decrease of population growth at low population abundance (*i.e.* demographic component, see Stephens *et al.* (1999)). Yet demographic stochasticity and the Allee effect are not the only threats to small populations; other density-dependent effects may also put these vulnerable populations at risk.

Classically, negative density-dependence describes situations where population growth rate is altered by increasing animal density (Nicholson 1933), with crowding driving the fate and viability of populations at risk of extinction (Beissinger & Westphal 1998; Henle *et al.* 2004). Negative density-dependent effects on population dynamics and demographic rates are often overlooked in

conservation for a simple reason: they are *a priori* expected to occur at high

population abundance, a counter-intuitive situation for threatened populations.

Yet, even in small populations, increasing abundance can occur rapidly as the

result of (*i*) a range contraction caused by habitat loss, (*ii*) the reintroduction of

individuals for population reinforcement, (*iii*) the disruption of natural

emigration due to habitat fragmentation, or (*iv*) following successful protection

measures without the possibility for the population to expand geographically.

Regardless of the underlying reason, negative density dependence can put the

population at risk of extinction by limiting its annual growth, or by leading to

undesirable ecological and demographic effects that hinder conservation efforts.

There are multiple well-documented negative density-dependence effects

(Eberhardt 1977; 2002; Fowler 1987; Bonenfant *et al.* 2009), most of which are

the result of behaviour and life history trait changes as population density

approaches carrying capacity (Bonenfant *et al.* 2009; Choquenot 1991; Van der

Wal *et al.* 2014). First, with increasing population density, individuals may die

from starvation because of a reduction in the availability and accessibility of

food resources by conspecifics. Such antagonistic interactions may generate high

stress levels which, in turns, magnify the effect of other mortality sources

(Peterson & Black 1988; Hone & Clutton-Brock 2007). For instance, at high

population density, epizootics are more likely to emerge and to spread quickly in

populations (Langwig *et al.* 2012). Second, higher densities may also force

non-dominant animals to occupy low-quality territories, ultimately decreasing

the average productivity and survival of the population (Krüger & Lindström

2001). Given its numerous biological consequences, early detection of

density-dependence is of the utmost importance for the management and

conservation of threatened species.

The tamaraw (*Bubalus mindorensis*) is a large herbivore species endemic to

the island of Mindoro, Philippines (Heude 1888; Custodio *et al.* 1996). The

species was originally found across the entire island of Mindoro, with a total

estimate of 10 000 individuals in 1 900 (Long *et al.* 2018). Since then, tamaraw

populations have been negatively impacted by land conversion for agriculture

and logging, trophy hunting, disease outbreaks spread out by domestic cattle

(Maala 2001), and traditional hunting conducted by upland indigenous

communities who share their living space with the species (Long *et al.* 2018).

Despite being officially protected by law since 1954, as well as the creation of

the sizeable Mounts Iglit-Baco Natural Park (MIBNP) protected area in 1970 to

protect its key habitat (Maala 2001), tamaraw range has decreased substantially

over the last decades. As a consequence, the tamaraw is now found in a few

isolated populations scattered across Mindoro, with MIBNP holding the largest

population (Matsubayashi *et al.* 2010; Wilson & Mittermeier 2011; Long *et al.*

2018), restricted to a single location <3 000 ha hereafter referred to as the core

zone of monitoring (CZM).

Given its importance for the conservation of this iconic species, the MIBNP

tamaraw population has been the cornerstone of past, present, and future

conservation plans (Maala 2001; Long *et al.* 2018). In an attempt to reinforce

tamaraw protection, wildlife managers came to an agreement with residing

indigenous people communities in 2016, declaring a 1 600 ha no-hunting zone

area within the CZM (Fig. 2). Unfortunately, one unforeseen consequence of the

no-hunting area establishment is an increase in hunting pressure at its

boundaries. Park rangers regularly report the setting of snare and spear traps in

5

the border areas, and violation of the no-hunting zone agreement is not

uncommon, with several deaths of tamaraws confirmed in the last years (Boyles

comm. pers). Furthermore, observations by rangers suggest that tamaraw

population density is growing within the no-hunting zone, which could increase

the risks of crowding. In this context, it is possible that negative

density-dependence of demographic parameters will affect the population in the

near future, undermining long-term conservation strategies.

Here, we took advantage of a 21 year time series of abundance data to

provide a detailed analysis of the tamaraw population dynamics in the CZM of

MIBNP. We predict (*i*) that because of the substantial conservation effort in the

last decade, overall population growth rate (noted *r*) should be statistically > 0.

Yet (*ii*), because the no-hunting agreement area creates a spatial demarcation

between risky and safe habitats, we expect local growth rates to differ in space,

and predict a decrease of *r* in areas further from the main rangers' base camps

and patrolling routes. Finally (*iii*), given the intensive conservation efforts in the

CZM, and relative safety of tamaraw in the CZM, we predict a negative

association between *r* and tamaraw abundance within the no-hunting zone as a

result of negative density dependence effects. To test the first two hypotheses, we

computed the population growth rate of the tamaraw population at two spatial

scales: over the entire CZM, and at specific locations used in annual tamaraw

counts (hereafter referred to as vantage points). To test our third hypothesis of

density-dependent demographic responses, we fitted and compared three

population dynamics models: the baseline exponential model (no

density-dependence), a Ricker model to test for a linear relationship between *r*

and annual abundance, and a particular formulation of Gompertz model to test

6

133   for a decrease of *r* through time.

MATERIAL AND METHODS

Study site

The Mounts Iglit-Baco Natural Park (N12°54′, E121°13′) is a 106 655 ha

protected area of the Philippines in the south-central part of the island of

Mindoro. The Mounts Iglit-Baco Natural Park harbors the largest population of

tamaraw, found in a single 3 000 ha area located on the south-west edge of the

protected area (Fig. 2). This 3 000 ha area demarcates our study site, and is

where most patrolling and observation activities are carried out (the so-called

CZM). In 2016, a 1 600 ha area inside the CZM was declared as a no-hunting

zone with the consensus of local indigenous people. There are currently around

15 park rangers, which are divided into two or three teams to monitor the area on

a regular basis. The teams patrol routes within the CZM, centered around the

three permanent base camps. The CZM and no-hunting zone represent approx. 3

and 1.5% of MIBNP, respectively.

The CZM area is a rolling grassland plateau with an average elevation of 800

m a.s.l., dominated by cogon (*Imperata cylindrica*) and wild sugarcane

(*Saccharum spontaneum*), and interspersed with numerous wooded creeks,

secondary forest fragments, and steep hills. In recent years, the invasive Siam

weed (*Chromolaena odorata*) has been rapidly spreading in the grassland area

and now covers an unknown fraction of the tamaraw habitat. The tropical

climate is strongly seasonal with a rainy season from June to October (mean

temperature 26.5°C; mean rainfall: 150 mm), and a hot dry season from

7

November to May (mean temperature 29.8°C; mean rainfall: 5 mm). Apart from
humans, the only potential predator to tamaraw is the reticulated python
(*Malayopython reticulatus*) that is capable of preying upon calves or yearlings,
though no such predation observations had been made so far.

Tamaraw counts

Every year since 2000, the Tamaraw Conservation Program (TCP), a program of
the Philippine Department of Environment and Natural Resources (DENR), has
conducted tamaraw counts across the CZM, yielding an annual index of relative
population abundance. The tamaraw counting area consists of 18 vantage points,
covering nearly 2 200 ha within the CZM (Fig. 2). The tamaraw counting
process is rather invasive for the MIBNP ecosystem because it involves the
burning of grasslands a few weeks ahead of each count to increase visibility and
to attract tamaraw to areas with nutrient-rich young grasses. For the past 21
years, between 1 200 and 1 500 ha of grasslands were intentionally burned each
year for the purpose of monitoring tamaraw abundance.

During the annual counts, individual tamaraws were counted simultaneously
at the 18 vantage points for one hour and a half at dusk and dawn over a period
of four consecutive days in late March or early April. For the 8 sessions and at
each vantage point, at least two observers spotted and recorded all animals seen,
writing down observation time, approximate location, and sex and age-category
of individuals (split into calves, yearlings, sub-adults, and adults). This sampling
design remained unchanged for the 21 years, though the number of observers
could vary from year to year. Immediately after the completion of the 8 sessions,
raw data were cross-checked and cleaned for potential double observations. The

8

total number of different animals corrected for double counts was then derived

for each vantage point and year, and constitute the consolidated counts C. These

annual counts comprise the only available source of information about tamaraw

population trends and dynamics for officials to use for conservation management

decisions. Note that, because of the COVID-19 pandemic, the tamaraw counts in

2020 did not take place and we missed detailed counts (vantage point level) for

years 2000 to 2003 and 2005 following the loss of field data sheets.

Data analyses

For populations of long-lived and endangered species, robust and accurate

estimations of the population growth rate (r) remain difficult to obtain.

Knowledge about r requires either the estimation of demographic rates combined

into a Leslie matrix (Leslie 1945) or the availability of long-term series of

population abundance, preferably on an annual basis. Because we were unable to

mark tamaraw, both for legal and ethical reasons (Putman 1995), we could not

individually identify animals, and thus it was not possible to apply methods of

abundance estimations based on capture-recapture (see Schwarz & Seber 1999,

for a review). We therefore used auto-regressive statistical models to analyze the

annuals counts. Working with abundance index data to derive an empirical

estimates of r, however, comes with serious methodological issues (Îto 1972;

Lindley 2003). Over the last decades, population counts have repeatedly been

associated with a large sampling variance (*e.g.* Caughley 1977; Morellet *et al.*

2007), and coefficients of variation for population abundance close to 30% have

been reported for various large mammal species (Lubow & Ransom 2016). From

a conservation perspective, a large sampling variance in the input data for

estimating *r* is a major pitfall because it leads to over-optimistic results, usually

from an over-estimation of both the point estimate and its precision (Lindley

2003). Despite the availability of appropriate statistical methods to account for

the sampling variance of population counts (Kalman filter, De Valpine &

Hastings 2002), such tools remain seldom used in practice in conservation.

Here, we analyzed the consolidated number of observed tamaraws (*C*), as

judged by observers to be different individuals, for each of the 18 vantage points,

split by years ($C_{s,t}$, where index $s \in \{1, \dots, 18\}$ stands for the vantage point, and

$t \in \{2000, \dots, 2021\}$ stands for the time in years. We implemented a state-space

model to tease apart process from sampling variance, making the assumption

that double and under-counts compensate and are randomly distributed from

year to year. We worked in a Bayesian framework (*e.g.* Kéry & Schaub 2011)

but note this is fully equivalent to a Kalman filter with a frequentist approach. To

do so, we defined $N_{s,t}$ as the unobserved 'true' abundance linked to $C_{s,t}$ by a

random effect corresponding to the process variance (σ_c^2) on the log scale to

ensure positive values for abundance. Our baseline observation model was:

$$\log(C_{s,t}) \sim \mathcal{N}(\log(N_{s,t}), \sigma_c^2)$$

Accordingly, we computed the population growth rate $r_{s,t}$ of the tamaraw

population at each vantage points *s*, as:

$$r_{s,t} = \log\left(\frac{N_{s,t+1}}{N_{s,t}}\right),$$

from counts corrected for the sampling variance, thus returning an unbiased

10

²²³ estimation of the population growth rate. The overall population growth of the

²²⁴ tamaraw population of MIBNP is then given by:

$$\bar{r} = \frac{1}{21} \sum_{t=1}^{21} \log\left(\frac{N_{t+1}}{N_t}\right),$$

²²⁵ where $N_t = \sum_{s=1}^{18} N_{s,t}$. We input missing values at the vantage point level (years

²²⁶ 2000–2003 and 2005) thanks to the sum constraint on $N_{s,t}$, and the recursive

²²⁷ equation linking $N_{s,t}$ and $N_{s,t+1}$. By back-casting missing values of tamaraw

²²⁸ abundance, we could estimate population growth at each vantage point (r_s) with

²²⁹ the same number of years and in turn, test for the effect of human disturbance at

²³⁰ the periphery of the CZM on the population dynamics.

²³¹   To test our third hypothesis we investigated potential density-dependence in

²³² the annual growth rate by fitting two models classically used to detect negative

²³³ density dependence (Dennis & Taper 1994): the Ricker and a Gompertz growth

²³⁴ model, to be compared with an exponential growth model. The exponential

²³⁵ growth model was fitted implicitly for the estimation of the average population

²³⁶ growth rate $\bar{r}$ and served as our baseline model:

$$\log(N_{t+1}) = \log(N_t) + \bar{r},$$

²³⁷   Then we fitted a Ricker growth model to the total number of tamaraws seen

²³⁸ per year N_t as follows:

$$\log(N_{t+1}) = \log(N_t) + r_m \times \left(1 - \frac{N_t}{K_l}\right),$$

239 where r_m is the maximum population growth for the tamaraw, and K_l is the

240 carrying capacity of the MIBNP. Likewise, we tested if the annual population

241 growth rate would decrease in time with population density, or any other

242 unmeasured variable varying in time, by fitting a particular formulation of the

243 Gompertz model (Norton's formulation, see Tjørve & Tjørve 2017):

$$\log(N_{t+1}) = \log(N_1) + \log\left(\frac{K_G}{N_1}\right) \times \left(1 - e^{-\beta \times t}\right),$$

Because the pair of parameters r_m and K_l (Ricker) and β and K_G (Gompertz)

are not separately identifiable (Lebreton & Gimenez 2013), we used informative

priors for r_m and β based on the reported maximum population growth for other

large bovids (see Traill *et al.* 2007, for a similar approach). Accordingly, we set

a moderately informative prior value of mean 0.3 and a precision of 0.1 in our

models (Table 1). We tested the occurrence of density-dependence processes in

the MIBNP tamaraw population by explicitly computing the probability (π_k) for

our abundance time series to be generated by each of the three candidate

demographic models during the model fit. At each MCMC step, a multinomial

variable stated which of the exponential, Gompertz and Ricker model was the

most likely underlying model given the data by taking the value one and zero

otherwise. We estimated the three probabilities π_k as the number of times each

models received a one divided by the number of MCMC iterations.

We fitted our statistical model to the tamaraw count data with JAGS 4.0

(Plummer *et al.* 2003), with 3 MCMC chains and a burn-in of 40,000 iterations.

We obtained parameter estimates with an additional 30 000 iterations and a

thinning factor of 5 (therefore giving a total of 6 000 MCMC samples). We

12

checked model convergence visually by investigating the mixing of MCMC

chains, along with the $\hat{R}$ statistics (Brooks & Gelman 1998). With the exception

of r_m (see above), we set uninformative priors with large variance for all other

parameters to be estimated, and checked the sensitivity of our results to the

average value of the priors by replicating the analyses with variable mean and

initial values for prior distributions. We report all parameters estimates as the

mean of the posterior distributions along with the 95% percentile for the credible

intervals following Louis & Zeger (2008): $_{95\%low}\, estimate\, _{95\%up}$.

Note that despite the fact that we analysed the number of tamaraws as

described above, we chose not to display such numbers on figures and to report a

standardized abundance ($C_t' = C_t/max(C_t)$) instead. This transformation applies

to the estimated carrying capacities of the Gompertz and Ricker models,

$K' = K/max(C_t)$. We did not provide raw counts to avoid confusion between the

index of relative tamaraw abundance we worked with, and a real population size

estimator that was not implemented at MIBNP. With this variable transformation

K_l and K_G are unitless and can no longer be interpreted as the maximum number

of tamaraws but the maximum relative abundance (C_t') we can expect to observe

at MIBNP. Reported values of the carrying capacities K_l' and K_g' can hence take

values > 1 because the carrying capacity has not yet been reached. Biologically

speaking, we put more emphasis on how far from the carrying capacity the

current abundance of tamaraws is (C_t/K) than of the number on animals *per se*.

RESULTS

From our baseline model of tamaraw counts, we estimated an average annual growth rate of $\bar{r} = {}_{0.01} 0.05 {}_{0.09}$ for the MIBNP population from 2000 to 2021, with an associated variance process equal to $\sigma_{proc} = {}_{0.25} 0.31 {}_{0.38}$ (Fig. 1). As expected for count data, the sampling variance was large $\sigma_c = {}_{7.18} 9.21 {}_{9.98}$, suggesting frequent double counts of the same individuals. For the vantage point specific growth rates (r_s), we observed a marked variability in space (Fig. 2; see Supplementary Table S1) as shown by the fact that growth rates ranged between $r_s = {}_{-0.42} -0.18 {}_{0.20}$ at Tarzan and $r_s = {}_{0.00} 0.14 {}_{0.26}$ at Bato Fidel and Bayokbok vantage points. Without exception, $r_s < 0$ were located at the periphery of the no-hunting area (Iyan, Tarzan, Saligue East, Mibluan, Fangandatan), although r_s of a few vantage points were > 0 despite being located on the no-hunting zone border (Nagbobong, Malibayong, Talafu West and East). The mean growth rate outside of the no-hunting area was $r = {}_{0.34} -0.05 {}_{0.08}$ (9 vantage points) while tamaraw abundance increased at rate of $r = {}_{0.04} 0.09 {}_{0.15}$ inside (9 vantage points).

We explicitly estimated the probabilities π_k that the exponential, Ricker, or Gompertz models would be the underlying process generating the observed variation in tamaraw abundance (proxy of Bayes factor). The estimated probabilities for the three population dynamics models given our count data were $\pi_1 = 0.10$, $\pi_2 = 0.90$ and $\pi_3 = 0.00$, respectively, supporting our hypothesis of density-dependence feedbacks in the tamaraw population. Point estimates for the two parameters of the Ricker model were $r_m = {}_{0.07} 0.08 {}_{0.09}$ for the maximum population growth rate and $K'_l = {}_{1.16} 1.37 {}_{1.62}$ for the carrying capacity.

14

Although Gompertz model received no statistical support, the estimated

coefficients were statistically significant and read $\beta = {}_{0.04}\, 0.05\, {}_{0.07}$, and

$K'_G = {}_{1.20}\, 1.41\, {}_{1.66}$ for the carrying capacity. Note that the statistical support for

the Ricker model held for the dynamics of the core area only, that is, when

considering only the 9 vantage points inside the no-hunting area ($\pi_1 = 0.02$,

$\pi_2 = 0.83$ and $\pi_3 = 0.15$).

DISCUSSION

The isolated tamaraw population in MIBNP has been increasing in size since

2000 (Fig. 1). Undoubtedly, a positive growth rate of 0.05 over 21 years of

monitoring is a noteworthy success for such a highly endangered species, both

for local populations and for global conservation efforts. However, the devil lies

in the details, and two additional results darken this otherwise positive picture.

First, we found support for the hypothesis of a contraction of the tamaraw

distribution, evidenced by a clear spatially structured pattern in the population

growth rates. Second, our results highlight a progressive decrease of the average

growth rate, suggesting density-dependent effects at the population level. The

population dynamics we document clearly indicate that new and urgent

conservation actions are needed to secure the future of the tamaraw population in

Mindoro.

Long-term Positive Growth of Tamaraws

326 The average population growth rate of $\bar{r} = 0.05$ gives strong evidence that

327 successful conservation policies over the past two decades have led to an

15

increase in tamaraw abundance. Nonetheless, an annual growth of 5% is

relatively low compared to other similar-sized or even larger cattle species

(Table 1). For instance, a population of African buffalo (*Syncerus caffer*) in

South Africa grew at a rate of 12% per year for 28 years (Jolles 2007), and

values close to 30% per year have regularly been recorded for bison (*Bison*

bison) and banteng (*Bos javanicus*), despite the fact that these species are

between 2 and 2.5 times larger in size than the tamaraw (Wilson & Mittermeier

2011). From this brief comparative approach, we would expect a substantially

higher average growth rate for tamaraws if the population was in a colonizing

phase. Given that the tamaraw is secure within the no-hunting area, our results

suggest that specific environmental factors are limiting its growth in the CZM.

A first limiting factor could be related to the fact that the most available and

accessible habitat for the tamaraw in MIBNP is grassland (83% of the CMZ),

which is dominated by competitive pioneer plant species such as cogon and wild

sugarcane. Both grasses appear to be *de facto* prominent in the tamaraw diet.

Although these grasses provide tamaraws with abundant and high quality forage

at the early stages of plant growth, in later stages of development these species

are characterized by much lower nutritive value, especially during the dry

season. Observations in the field indicate that once grass growth is complete, the

stems exceeds tamaraw height (300cm vs. < 120 cm, respectively) and dry out,

thus becoming both inaccessible and unpalatable. It is therefore possible that

lack of food resource accessibility and availability could be limiting the growth

rate for the population.

The regular burning of the grassland habitat for population count may also be

impacting plant composition in the CZM, with implications for tamaraw

16

population dynamics. On the one hand, regular burning promotes the

regeneration of new grass shoots, providing a nutrient-rich food source for

tamaraw. On the other hand, previous observations suggest that regular burning

could promote Siam weed expansion (Nath *et al.* 2019), which appears to be

rapidly spreading throughout the study site. Like other grazing herbivores

(Rozen-Rechels *et al.* 2017), tamaraw avoid this invasive plant, and so the

increase of Siam weed could ultimately decrease the CMZ carrying capacity

(Dumalisile & Somers 2017). Although burning creates high-quality forage in

the short-term, such an intensive habitat management practice is likely to cause a

progressive decrease in habitat quality over the long term.

Possible Impacts of Human Disturbance and Hunting

While the positive growth of the tamaraw population over the past two decades

is evidence of a successful decrease in hunting pressure in the CZM, human

activities could also account for the low tamaraw population growth rate we

report. Tamaraw have always been traditionally hunted by indigenous people in

Mindoro, and despite the "no-hunting agreement" established in 2016, both

intentional and unintentional tamaraw deaths still occur. Moreover, the tamaraw

is sought after for bushmeat by lowlander poachers. Because illegal activities are

difficult to control in MIBNP, it has likely led to an increased mortality of all

age-classes of tamaraws to an unknown, but perhaps considerable extent.

In the early years of the tamaraw monitoring, the apparent population growth

(Fig. 1) likely resulted from immigrating individuals from outside of the CMZ,

and a high survival and reproduction rate inside. Today a different demographic

process could be at work, with adult females gradually concentrating at the heart

17

of the no-hunting area to avoid human disturbance. Conversely, a limited number

of young individuals and adult males may venture outside of the no-hunting area,

putting themselves at greater risk of dying. The net movements of animals in and

out of the no-hunting area could lead to an apparent continuous growth of

tamaraw abundance since the beginning of monitoring efforts, despite a shift in

the underlying population dynamics.

The marked spatial heterogeneity we document, with negative growth rates at

the periphery of the counting area, could indeed indicate a lower protection

efficiency in areas further away from the main patrolling routes and ranger base

camps. Such a result supports our second hypothesis, suggesting an unexpected

contraction of the tamaraw range at MIBNP, and the disappearance of

individuals at the periphery of the CZM. In other words, the CZM area may

function like a source-sink system, with the more central and better patrolled

areas exhibiting positive growth rates, and more peripheral areas with higher

hunting pressure serving as population sinks. Again, our findings provide some

support for this idea, as local growth rates in the core area of the no-hunting zone

have values as high as 14% (Fig. 2), which is more in line with the average

growth rates that we reviewed for large bovids (Table 1).

Space Needed for Sustaining Population Growth

396 The gradual decrease of the tamaraw population growth rate in MIBNP, coupled

397 with the overall abundance in the CZM, suggests a type of density-dependence is

398 occurring, where a change in demographic rates lowers the number of

399 individuals added to the population from year to year (Fowler 1987; Sinclair

400 1989). Predicted growth rates from the lowest to the highest abundance varied

18

between $r = 0.08$ and $r = 0.01$ for tamaraw. From the current data, our best model suggests a carrying capacity of 1.4, meaning that only 40% of growth is left possible before reaching the carrying capacity. If competition for food or mating becomes strong then the main tamaraw population may experience an increased net loss of animals in the near future. Tamaraws may already be experiencing a food resource shortage, which could make the population more prone to parasitism (Stien *et al.* 2002) or epizootics (Davidson & Doster 1997).

Another behaviour associated with high population abundance is dispersal (Matthysen 2005). In mammals, sub-adult individuals are more likely to disperse at high than low population abundance (e.g. Loe *et al.* 2009, for an example on red deer *Cervus elaphus*). At the high population abundance we documented in MIBNP, we would expect a greater number of dispersing individuals to venture outside of the no-hunting zone, where the risk of mortality is high because of hunting or poaching. In addition to jeopardizing the fate of the population by decreasing survival probabilities of sub-adults that are key to maintaining the future of the population, the empirical evidence for density-dependence that we report would limit the ability of this population to act as a source. With a combination of low recruitment and low survival of sub-adults, old individuals would quickly become over-represented in the population, which could depress growth even further.

Unfortunately, in the absence of individual monitoring of tamaraws to estimate demographic rates independently of abundance, the real cause of the density-dependence we report will be uncertain. We are currently unable to determine if the gradual decrease in growth rate is the consequence of an increased mortality or dispersal rate, a reduced reproduction, or some

19

combination of these factors. The difficulty of teasing apart different

demographic processes is a well-known limit of the so-called pattern-oriented

approach (*sensu* Krebs 2002) when working with time series of abundance.

Further investigation of these possibilities is needed, as each situation would

require different management approaches.

Implications for Conservation

The main conservation goals for the tamaraw are (1) to ensure its long-term

viability, and (2) support the expansion of the population within MIBNP and

potentially in other sites. Current conservation measures assume that the

tamaraw population growth will continue in the next decade. The detection of

density-dependence, however, suggests continued population growth will not be

the case. In this situation, the dynamics of the tamaraw will change in the years

to come if resources become the main limiting factor inside the no-hunting zone,

forcing animals to disperse to, and potentially get killed in the outer zone.

Several actions must be implemented by managers and authorities to address this

issue in the short term. The priorities should be to expand the no-hunting zone or

to create safety corridors through consensual agreement with indigenous people,

both implemented in the context of creating a more sustainable land-use and

traditional hunting system. Arriving at such an agreement with indigenous

people will be a long process, but these conversations are already ongoing.

Extending the no-hunting area will ultimately increase the sustainable population

size, thus making it more viable and maintaining more genetic diversity.

Likewise, allowing for dispersing individuals to leave the CZM and possibly

settle in distant areas would relieve density dependence pressure on the core

population.

However, one could argue that such conservation actions only represent a

short-term solution, as even if the size of the no-hunting area was increased, the

tamaraw population would eventually grow towards the carrying capacity. We

acknowledge this, but believe that these short-term conservation actions are still

important, as they would help buy valuable time that is necessary to devise more

long-term strategies. Any long-term strategy will likely involve active population

management through translocation. Two of the four known populations of

tamaraw on Mindoro (Aruyan Malate and Mount Calavite) have extremely small

population sizes, and are thus believed to be non-viable. Transporting a small

number of animals to these populations would improve their viability, while

simultaneously reducing the effects of density-dependence for the main

population at MIBNP. The number of tamaraw that are translocated should be

determined by the number of recruited individuals each year, so that ultimately

there is no decrease in population size in the MIBNP. We acknowledge that

translocation of tamaraw may be a perceived by stakeholders as a risky

proposition, but a preliminary feasibility study for translocation and captive

breeding is a realistic option/ an appropriate next step. Ultimately, however,

preventing the extinction of tamaraw will require decision makers to balance

short-term actions with long-term strategies that involve active population

management.

Conclusion

472 Securing large areas of habitat is often an effective conservation approach, but

473 may not always be a viable strategy to protect highly threatened species, as

21

illustrated by our study on the tamaraw population in MIBNP. More generally,

we show that positive growth rates could potentially hide a darker picture for

some threatened species. First, the growth of a population that is confined to a

small conservation area may reflect a range contraction as a result of

anthropogenic pressures, rather than the effect of increased abundance *per se*.

Second, because of this, the population could face important demographic

challenges that, in the long-term, lead to new threats. In this context, there is an

increasing need to understand the interaction between anthropogenic activities

and the population dynamics of threatened species. Our results suggest that, in

some situations, conservation strategies designed to protect small, isolated

populations may lead to long-term demographic issues that undermine species

recovery. On the positive side, our findings also show how detailed analyses of

relative population abundance data, even without absolute population size

estimators, can help provide information that assists in the development of

conservation measures that are essential for large mammal conservation.

ACKNOWLEDGMENTS

We are grateful to Alvaro Gonzalez. We also acknowledge the invaluable help of all

rangers, students and volunteers who contributed to the annual tamaraw counts. Thanks

are extended to Jean-Michel Gaillard for his critical and constructive reading of the

manuscript.

REFERENCES

Beissinger, S.R. & Westphal, M.I. (1998). On the use of demographic models of
population viability in endangered species management. *The Journal of Wildlife Management*, 62, 821–841.

22

Bonenfant, C., Gaillard, J.M., Coulson, T., Festa-Bianchet, M., Loison, A., Garel, M.,
Loe, L.E., Blanchard, P., Pettorelli, N., Owen-Smith, N., Du Toit, J. & Duncan, P.
(2009). Empirical evidence of density-dependence in populations of large herbivores.
Advances in Ecological Research, 41, 313–357.

Boyce, M.S. (1992). Population viability analysis. *Annual Review of Ecology and*
Systematics, 23, 481–506.

Brooks, S.P. & Gelman, A. (1998). General methods for monitoring convergence of
iterative simulations. *Journal of Computational and Graphical Statistics*, 7, 434–455.

Buys, D. & Dott, H. (1991). Population fluctuations and breeding of eland *Taurotragus*
oryx in a western Transvaal nature reserve. *Koedoe*, 34, 31–36.

Caughley, G. (1977). *Analysis of vertebrate populations*. Wiley, London. Graeme
Caughley. 'A Wiley-Interscience Publication' Includes index. Bibliography:
p.217-228.

Caughley, G. (1994). Directions in conservation biology. *Journal of Animal Ecology*,
63, 215–244.

Choquenot, D. (1991). Density-dependent growth, body condition, and demography in
feral donkeys: testing the food hypothesis. *Ecology*, 72, 805–813.

Choquenot, D. (1993). Growth, body condition and demography of wild banteng (*Bos*
javanicus) on Cobourg Peninsula, northern Australia. *Journal of Zoology*, 231,
533–542.

Cornélis, D., Melletti, M., Korte, L., Ryan, S.J., Mirabile, M., Prin, T. & Prins, H.H.
(2014). *African buffalo Syncerus caffer (Sparrman, 1779)*, Cambridge University
Press, pp. 326–372.

Courchamp, F., Clutton-Brock, T.H. & Grenfell, B. (1999). Inverse density dependence
and the Allee effect. *Trends in Ecology & Evolution*, 14, 405–410.

Custodio, C.C., Lepiten, M.V. & Heaney, L.R. (1996). *Bubalus mindorensis*.
Mammalian Species, pp. 1–5.

Davidson, W. & Doster, G. (1997). Health characteristics and white-tailed deer
population density in the southeastern United States. In: *The Science of*
overabundance: deer ecology and population management (eds. McShea, W.J.,
Underwood, H.B. & Rappole, J.H.). Smithsonian Institution Press, Washington, DC
(USA), pp. 164–184.

De Valpine, P. & Hastings, A. (2002). Fitting population models incorporating process
noise and observation error. *Ecological Monographs*, 72, 57–76.

Dennis, B. & Taper, M. (1994). Density dependence in time series observations of
natural populations: estimation and testing. *Ecological Monographs*, 64, 205–224.

Dumalisile, L. & Somers, M.J. (2017). The effects of an invasive alien plant
(*Chromolaena odorata*) on large African mammals. *Nature Conservation Research*,
2, 102–108.

Eberhardt, L.L. (1977). Optimal policies for conservation of large mammals, with
special references to marine ecosystems. *Environmental Conservation*, 4, 205–212.

Eberhardt, L.L. (2002). A paradigm for population analysis of long-lived vertebrates.
Ecology, 83, 2841–2854.

Fowler, C.W. (1987). A review of density dependence in populations of large mammals.
Current Mammalogy, 1, 401–441.

Fuller, J.A., Garrott, R.A., White, P., Aune, K.E., Roffe, T.J. & Rhyan, J.C. (2007).
Reproduction and survival of Yellowstone bison. *Journal of Wildlife Management*,
71, 2365–2372.

Gates, C. & Larter, N. (1990). Growth and dispersal of an erupting large herbivore
population in northern Canada: the Mackenzie wood bison (*Bison bison athabascae*).
Arctic, pp. 231–238.

Haque, N. (1990). *Study on the ecology of wild ungulates of Keoladeo national Park
Bharatpur, Rajasthan*. Ph.D. thesis, Aligarh Muslim University.

Heinen, J.T. & Paudel, P.K. (2015). On the translocation of wild asian buffalo *Bubalis
arnee* in Nepal: are feral backcrosses worth conserving? *Conservation Science*, 3,
11–19.

Henle, K., Sarre, S. & Wiegand, K. (2004). The role of density regulation in extinction
processes and population viability analysis. *Biodiversity and Conservation*, 13, 9–52.

Heude, P. (1888). Études sur les ruminants de l'Asie orientale. cerfs des Philippines et
de l'Indo-Chine. *Mémoires Concernant l'Histoire Naturelle de l'Empire Chinois par
des Pères de la Compagnie de Jésus*, 2, 1–10.

Hone, J. & Clutton-Brock, T.H. (2007). Climate, food, density and wildlife population
growth rate. *Journal of Animal Ecology*, 76, 367–367.

Hone, J., Duncan, R.P. & Forsyth, D.M. (2010). Estimates of maximum annual
population growth rates (r_m) of mammals and their application in wildlife
management. *Journal of Applied Ecology*, 47, 507–514.

Îto (1972). On the methods for determining density-dependence by means of regression.
Oecologia, 10, 776–778.

Jolles, A.E. (2007). Population biology of african buffalo (*Syncerus caffer*) at Hluhluwe-iMfolozi Park, South Africa. *African Journal of Ecology*, 45, 398.

Kemp, N., Dilger, M., Burgess, N. & Van Dung, C. (1997). The saola *Pseudoryx nghetinhensis* in Vietnam – new information on distribution and habitat preferences, and conservation needs. *Oryx*, 31, 37–44.

Kéry, M. & Schaub, M. (2011). *Bayesian population analysis using WinBUGS: a hierarchical perspective*. Academic Press, Amsterdam, the Netherlands.

Koláčková, K., Hejcmanova, P., Antonínová, M. & Brandl, P. (2011). Population management as a tool in the recovery of the critically endangered Western Derby eland *Taurotragus derbianus* in Senegal, Africa. *Wildlife Biology*, 17, 299–311.

Krasiński, Z.A. (1978). Dynamics and structure of the european bison population in the białowieża Primeval Forest. *Acta Theriologica*, 23, 3–48.

Krebs, C.J. (2002). Two complementary paradigms for analysing populations dynamics. *Philosophical Transactions of the Royal Society of London, Series B*, 357, 1211–1219.

Krüger, O. & Lindström, J. (2001). Habitat heterogeneity affects population growth in goshawk *Accipiter gentilis*. *Journal of Animal Ecology*, 70, 173–181.

Lande, R. (1993). Risks of population extinction from demographic and environmental stochasticity and random catastrophes. *American Naturalist*, 142, 911–927.

Langwig, K.E., Frick, W.F., Bried, J.T., Hicks, A.C., Kunz, T.H. & Marm Kilpatrick, A. (2012). Sociality, density-dependence and microclimates determine the persistence of populations suffering from a novel fungal disease, white-nose syndrome. *Ecology Letters*, 15, 1050–1057.

Larter, N., Sinclair, A., Ellsworth, T., Nishi, J. & Gates, C. (2000). Dynamics of reintroduction in an indigenous large ungulate: the wood bison of northern Canada. *Animal Conservation*, 3, 299–309.

Lebreton, J.D. & Gimenez, O. (2013). Detecting and estimating density dependence in wildlife populations. *Journal of Wildlife Management*, 77, 12–23.

Leslie, P.H. (1945). On the use of matrices in population mathematics. *Biometrika*, 33, 182–212.

Lindley, S.T. (2003). Estimation of population growth and extinction parameters from noisy data. *Ecological Applications*, 13, 806–813.

Loe, L.E., Mysterud, A., Veiberg, V. & Langvatn, R. (2009). Negative density-dependent emigration of males in an increasing red deer population. *Proceedings of the Royal Society B: Biological Sciences*, 276, 2581–2587.

25

Long, B., Schütz, E., Appleton, M., Gonzales, A., Holland, J., Lees, C., Marandola, E.,
Natural, V.C.J., Pineda-David, M.T.J., Salao, C., Slade, J., Tan, E.H.P., Tiongson, L.
& Young, S. (2018). Review of tamaraw (*Bubalus mindorensis*) status and
conservation actions. *BULLetin*, 1, 18–34.

Louis, T.A. & Zeger, S.L. (2008). Effective communication of standard errors and
confidence intervals. *Biostatistics*, 10, 1–2.

Lubow, B.C. & Ransom, J.I. (2016). Practical bias correction in aerial surveys of large
mammals: Validation of hybrid double-observer with sightability method against
known abundance of feral horse (*Equus caballus*) populations. *PLoS One*, 11,
e0154902.

Maala, C.P. (2001). Endangered Philippine wildlife species with special reference to the
Philippine eagle (*Pithecophaga jefferyi*) and tamaraw (*Bubalus mindorensis*). *Journal
of International Development and Cooperation*, 8, 1–17.

Mace, G.M., Collar, N.J., Gaston, K.J., Hilton-Taylor, C., Akçakaya, H.R.,
Leader-Williams, N., Milner-Gulland, E.J. & Stuart, S.N. (2008). Quantification of
extinction risk: IUCN's system for classifying threatened species. *Conservation
Biology*, 22, 1424–1442.

Mathur, V.B. (1991). *The ecological interaction between habitat composition, habitat
quality and abundance of some wild ungulates in India*. Ph.D. thesis, University of
Oxford.

Matsubayashi, H., Boyles, R.M., Salac, R.L., Del Barrio, A., Cruz, L., Garcia, R.A.,
Ishihara, S. & Kanai, Y. (2010). Present status of tamaraw (*Bubalus mindorensis*) in
Mt. Aruyan, Mindoro, Philippines. *Tropics*, 18, 167–170.

Matthysen, E. (2005). Density-dependent dispersal in birds and mammals. *Ecography*,
28, 403–416.

Morellet, N., Gaillard, J.M., Hewison, A.J.M., Ballon, P., Boscardin, Y., Duncan, P.,
Klein, F. & Maillard, D. (2007). Indicators of ecological change: new tools for
managing populations of large herbivores. *Journal of Applied Ecology*, 44, 634—643.

Mugabo, M., Perret, S., Legendre, S. & Le Galliard, J.F. (2013). Density-dependent life
history and the dynamics of small populations. *Journal of Animal Ecology*, 82,
1227–1239.

Mysterud, A., Bartoń, K.A., Jedrzejewska, B., Krasiński, Z.A., Niedziałkowska, M.,
Kamler, J.F., Yoccoz, N.G. & Stenseth, N.C. (2007). Population ecology and
conservation of endangered megafauna: the case of european bison in Białowieża
Primeval Forest, Poland. *Animal Conservation*, 10, 77–87.

Nath, A., Sinha, A., Lahkar, B.P. & Brahma, N. (2019). In search of aliens: Factors
influencing the distribution of *Chromolaena odorata* L. and Mikania micrantha Kunth
in the Terai grasslands of Manas National Park, India. *Ecological Engineering*, 131,
16–26.

Neel, M.C., Leidner, A.K., Haines, A., Goble, D.D. & Scott, J.M. (2012). By the
numbers: How is recovery defined by the US endangered species act? *BioScience*, 62,
646–657.

Nicholls, A., Viljoen, P., Knight, M. & Van Jaarsveld, A. (1996). Evaluating population
persistence of censused and unmanaged herbivore populations from the Kruger
National Park, South Africa. *Biological Conservation*, 76, 57–67.

Nicholson, A.J. (1933). The balance of animal populations. *Journal of Animal Ecology*,
S2, 132–178.

Ottichilo, W.K., De Leeuw, J., Skidmore, A.K., Prins, H.H. & Said, M.Y. (2000).
Population trends of large non-migratory wild herbivores and livestock in the Masai
Mara ecosystem, Kenya, between 1977 and 1997. *African Journal of Ecology*, 38,
202–216.

Peterson, C.H. & Black, R. (1988). Density-dependent mortality caused by physical
stress interacting with biotic history. *American Naturalist*, 131, 257–270.

Plummer, M. *et al.* (2003). JAGS: A program for analysis of bayesian graphical models
using gibbs sampling. In: *Proceedings of the 3rd international workshop on
distributed statistical computing*. Vienna, Austria, vol. 124, p. 10.

Putman, R. (1995). Ethical considerations and animal welfare in ecological field studies.
Biodiversity & Conservation, 4, 903–915.

Rozen-Rechels, D., te Beest, M., Dew, L.A., le Roux, E., Druce, D.J. & Cromsigt, J.P.
(2017). Contrasting impacts of an alien invasive shrub on mammalian savanna
herbivores revealed on a landscape scale. *Diversity and Distributions*, 23, 656–666.

Schwarz, C. & Seber, G.A.F. (1999). A review of estimating animal abundance III.
Statistical Science, 14, 427–456.

Shaffer, M.L. (1981). Minimum population sizes for species conservation. *BioScience*,
31, 131–134.

Sinclair, A.R.E. (1989). Population regulation in animals. In: *Ecological concepts : the
contribution of ecology to an understanding of the natural world* (eds. Cherrett, J.M.
& Bradshaw, A.D.). Blackwell Scientific Publications, Oxford, England., pp.
197–241.

Singer, F.J. & Norland, J.E. (1994). Niche relationships within a guild of ungulate species in Yellowstone National Park, Wyoming, following release from artificial controls. *Canadian Journal of Zoology*, 72, 1383–1394.

Steinmetz, R., Chutipong, W., Seuaturien, N., Chirngsaard, E. & Khaengkhetkarn, M. (2010). Population recovery patterns of southeast asian ungulates after poaching. *Biological Conservation*, 143, 42–51.

Stephens, P.A., Sutherland, W.J. & Freckleton, R.P. (1999). What is the Allee effect? *Oikos*, pp. 185–190.

Stien, A., Irvine, R.J., Ropstad, E., Halvorsen, O., Langvatn, R. & Albon, S.D. (2002). The impact of gastrointestinal nematodes on wild reindeer: experimental and cross-sectional studies. *Journal of Animal Ecology*, 71, 937–945.

Tjørve, K.M. & Tjørve, E. (2017). The use of Gompertz models in growth analyses, and new Gompertz-model approach: An addition to the unified-Richards family. *PloS one*, 12, e0178691.

Traill, L.W., Bradshaw, C.J.A. & Brook, B.W. (2007). Minimum viable population size: a meta-analysis of 30 years of published estimates. *Biological Conservation*, 139, 159–166.

Van Houtan, K.S., Halley, J.M., Van Aarde, R. & Pimm, S.L. (2009). Achieving success with small, translocated mammal populations. *Conservation Letters*, 2, 254–262.

Van der Wal, E., Laforge, M.P. & McLoughlin, P.D. (2014). Density dependence in social behaviour: home range overlap and density interacts to affect conspecific encounter rates in a gregarious ungulate. *Behavioral Ecology and Sociobiology*, 68, 383–390.

Wilson, D. & Mittermeier, R. (2011). *Handbook of the mammals of the world. Vol. 2. Hoofed Mammals*. Lynx Edicions, Barcelona.

TABLE CAPTIONS

Table 1 Estimated annual growth rates (r) of some large *bovinae*

populations with body size comparable to the endangered tamaraw

(*Bubalus mindorensis*). The observed population growth rate of

tamaraws at Mounts Iglit-Baco Natural Park, Mindoro, Philippines,

between 2000 and 2019 of $r = 0.05$ lies at the lower range of reported

values. Note some populations may be in a very different ecological

context (colonization, saturation) making comparisons .

Table 1

Species	Location	*r*	Reference
Bison bison	Mackenzie Bison Sanctuary, Canada	0.21	Gates & Larter (1990)
Bison bison	Mackenzie Bison Sanctuary, Canada	0.20	Larter *et al.* (2000)
Bison bison	Yellowstone, Wyoming, USA	0.07	Fuller *et al.* (2007)
Bison bison	Yellowstone, Wyoming, USA	0.30	Singer & Norland (1994)
Bison bonasus	Białowieża, Poland	0.07	Mysterud *et al.* (2007)[1]
Bison bonasus	Białowieża, Poland	0.18	Krasiński (1978)[1]
Bos gaurus	Thung Yai Naresuan Sanctuary, Thailand	0.31	Steinmetz *et al.* (2010)
Bos javanicus	Not applicable, model prediction	0.32	Hone *et al.* (2010)
Bos javanicus	Cobourg peninsula, Australia	0.00	Choquenot (1993)
Boselaphus tragocamelus	Kanha National Park, India	0.18	Mathur (1991)
Boselaphus tragocamelus	Keoladoe National Park, India	0.02 - 0.06	Haque (1990)
Bubalus arnee		0.02 - 0.06	Khatri et al. (2012)
Bubalus arnee	Koshi Tappu Wildlife Reserve, Nepal	0.03	Heinen & Paudel (2015)
Bubalus bubalis		0.03	Heinen (1993)
Bubalus bubalis		?	Freeland and Boulton (1990)
Bubalus mindorensis	Mount Iglit-Baco National Park, Philipines	0.05	This study
Pseudoryx nghetinhensis	Vietnam & Laos	0.05	Kemp *et al.* (1997)[b]
Syncerus caffer	Hluhluwe-iMfolozi Park, South Africa	0.12	Jolles (2007)
Syncerus caffer	Zakouma National Park, Chad	0.08	Cornélis *et al.* (2014)
Taurotragus oryx	Lombard Nature Reserve, South Africa	0.12	Buys & Dott (1991)[c]
Taurotragus oryx	South Africa	0.16	Van Houtan *et al.* (2009)
Taurotragus oryx	Kruger National Park, South Africa	−0.02	Nicholls *et al.* (1996)
Taurotragus oryx	Masai Mara Ecosystem, Kenya	−0.11 - 0.07	Ottichilo *et al.* (2000)
Taurotragus derbianus	Senegal	0.31	Koláčková *et al.* (2011)

FIGURE CAPTIONS

Figure 1 Time series of tamaraw (*Bubalus mindorensis*) abundance in Mounts Iglit-Baco Natural Park on Mindoro island, Philippines, from years 2000 to 2021 (with the exception of 2020 because of the COVID pandemic). Raw counts (black solid line) and counts corrected for sampling variance (gray solid line) are presented with the associated 95% credible intervals. Over the 21 years of monitoring, tamaraw abundance increased at a rate of 5% per year on average.

Figure 2 Spatial variation in the local growth rate of tamaraw (*Bubalus mindorensis*) abundance in Mounts Iglit-Baco Natural Park (MINBNP) on Mindoro island, Philippines, from years 2000 to 2021.

Figure 1

Figure 2

713 **SUPPLEMENTARY MATERIAL**

714 **Table S1** Estimates and 95% credible intervals of tamaraw local growth rate as

715 observed on the 18 vantage points used to count the tamaraw population at Mounts

Iglit-Baco National Park (Mindoro Island, Philippines) between 2000 and 2021.

Vantage point	r	2.5%	97.5%
Loibfo	0.12	−0.02	0.24
Magawang	0.11	−0.02	0.24
Bayokbok	0.14	0.00	0.26
Bato Fidel	0.14	0.00	0.26
Inubon	0.11	−0.02	0.24
Mibluan	0.08	−0.06	0.22
Nagbobong	0.02	−0.13	0.17
Fangandatan	0.00	−0.19	0.16
Anyayos	0.07	−0.07	0.20
Lanas I	0.07	−0.08	0.21
Iyan	−0.04	−0.25	0.14
Tarzan	−0.18	−0.42	0.02
Talafu East	0.05	−0.10	0.19
Talafu West	0.05	−0.10	0.19
Malitwang	0.07	−0.09	0.21
Lanas II	0.01	−0.14	0.15
Saligue East[1]	−0.08	−0.40	0.15
Malibayong	−0.01	−0.20	0.15
Tangle	−0.48	−2.94	0.16

[1]Counts at this vantage point stopped in 2019, replaced by Tangle