Cast away on Mindoro island: could density dependence threaten tamaraw (*Bubalus mindorensis*) on the long term?

C. Bonenfant[1], A. Rutschmann[2], J. Burton[3], R. Boyles[4], F. García[5], A. Tilker[6,7], and E. Schütz[5]

Université de Lyon, F-69 000, Lyon ; Université Lyon 1 ; CNRS, UMR 5558, Laboratoire "Biométrie et Biologie Évolutive", F-69 622, Villeurbanne, France.
School of Biological Sciences, University of Auckland, Auckland, New Zealand
IUCN SSC Asian Wild Cattle Specialist Group, Global Conservation, Texas, USA
Department of Environment and Natural Resources of the Philippines, Occidental Mindoro
D'Aboville Foundation and Demo Farm Inc, Manila, Philippines
Re:wild, Austin, Texas USD
Leibniz Institute for Zoo and Wildlife Research, Berlin, Germany
5 D'Aboville Foundation and Demo Farm Inc, Manila, Philippines

Keywords

counts, exponential model, Mindoro dwarf buffalo, protection measures, poaching, population dynamics.

Correpondence

Christophe Bonenfant, `christophe.bonenfant@univ-lyon1.fr` – ORCID 0000-0002-9924-419X

Abstract

Endangered species, despite often living at low population densities, may undergo unexpected density-dependent feedbacks in the case of successful recovery or marked reduction in range. Because density-dependence dynamics can increase risk of extinction, these effects can hamper conservation efforts. In this study, we analyse the dynamics of the largest population of the tamaraw (*Bubalus mindorensis*), a Critically Endangered species endemic to the Philippines. The population is located within a $< 3,000$ha area in Mounts Iglit-Baco Natural Park (MIBNP), with limited expansion possibilities. We took advantage of a 21 year time series of tamaraw counts to estimate annual population growth rate and possible density-dependence using state-space models in a Bayesian framework, hence accounting for sampling errors in the counts. We found that the MIBNP tamaraw population has been increasing in size at an average rate of $+5\%$ per year, as would be expected given its protected status by law. However, this growth rate is relatively low compared to other similar-sized *Bovinea* species. Population growth showed strong spatial structuring, with a population growth close to $+14\%$ in the core area of protection, and a reduction of abundance of -27% at the periphery of its range inside MIBNP. That the annual population growth rate has progressively decreased since 2008 is concerning, and we interpret this result to be a sign of density-dependence. The isolated tamaraw population in MIBNP is currently experiencing a contraction of its range, likely caused by anthropogenic pressures forcing tamaraw to live at relatively high densities in the core zone where protection is most effective. Our study highlights the fact that, despite the

35   encouraging results as indicated by continuous population growth over the last

36   two decades, the long-term viability of the MIBNP tamaraw population remains

37   uncertain.

INTRODUCTION

Assessing population abundance is an important component of conservation planning because of the strong link between population size and likelihood of extinction (Boyce 1992). Accordingly, low population abundance is one of the primary criteria used for the evaluation of conservation status (Mace *et al.* 2008; Neel *et al.* 2012). The International Union for Conservation of Nature (IUCN), for example, classifies any species with a small population size as threatened, regardless of past or projected future population declines (Mace *et al.* 2008). Indeed, the effects of small population size on species persistence has been identified as one of the two central paradigms of conservation biology (Caughley 1994). Research has shown that the population dynamics of small populations differ in several ways from what is reported at higher abundance (Caughley 1994; Mugabo *et al.* 2013). When population size is small, the random death of a few individuals may precipitate the entire population toward extinction, a process referred to as demographic stochasticity (Shaffer 1981; Lande 1993). Similarly, when mating partners are too few to meet and reproduction fails, demographic stochasticity generates the so-called Allee effect (Courchamp *et al.* 1999), characterized by a decrease of population growth at low population abundance (*i.e.* demographic component, see Stephens *et al.* (1999)). Yet demographic stochasticity and the Allee effect are not the only threats to small populations; other density-dependent effects may also put these vulnerable populations at risk.

Classically, negative density-dependence describes situations where population growth rate is altered by increasing animal density (Nicholson 1933),

with crowding driving the fate and viability of populations at risk of extinction

(Beissinger & Westphal 1998; Henle *et al.* 2004). Negative density-dependent

effects on population dynamics and demographic rates are often overlooked in

conservation for a simple reason: they are *a priori* expected to occur at high

population abundance, a counter-intuitive situation for threatened populations.

Yet, even in small populations, increasing abundance can occur rapidly as the

result of (*i*) a range contraction caused by habitat loss, (*ii*) the reintroduction of

individuals for population reinforcement (*iii*) the disruption of natural emigration

due to habitat fragmentation, or (*iv*) following successful protection measures

without the possibility for the population to expand geographically. Regardless

of the underlying reason, negative density dependence can put the population at

risk by preventing its ability to reach full carrying capacity, and may ultimately

cause undesirable ecological and demographic effects that hinder conservation

efforts.

There are multiple well-documented negative density-dependence effects

(Eberhardt 1977; 2002; Fowler 1987; Bonenfant *et al.* 2009), most of which are

the result of behaviour and life history trait changes as population density

approaches carrying capacity (Bonenfant *et al.* 2009; Choquenot 1991; Van der

Wal *et al.* 2014). First, with increasing population density, individuals may die

from starvation because of a reduction in the availability and accessibility of

food resources by con-specifics. Such antagonistic interactions may generate

high stress levels which, in turns, magnify the effect of other mortality sources

(Peterson & Black 1988). For instance, at high population density, epizootics are

more likely to emerge and to spread quickly in populations (Langwig *et al.*

2012). Second, higher densities may also force non-dominant animals to occupy

low-quality territories, ultimately decreasing the average productivity and

survival of the population (Krüger & Lindström 2001). Finally, because high

population density exacerbates the effect of environmental variation on

demographic rates (Hones & Clutton-Brock 2007), the risks of massive mortality

events are much higher if the population is maintained close to carrying capacity

on the long term. Given its numerous biological consequences, early detection of

density-dependence is of the utmost importance for the management and

conservation of threatened species.

The tamaraw (*Bubalus mindorensis*) is a type of wild cattle species endemic

to the island of Mindoro, Philippines (Heude 1888; Custodio *et al.* 1996).

Despite being officially protected by law since 1954, as well as the creation of

the sizeable Mts Iglit-Baco Natural Park (MIBNP) protected area in 1970 to

protect its key habitat, (Maala 2001), tamaraw range has decreased substantially

over the last decades. The species was originally found across the entire island

of Mindoro, with a total estimate of 10,000 individuals in 1900 (Long *et al.*

2018). Since then, however, tamaraw populations have been negatively impacted

by land conversion for agriculture and logging, trophy hunting, disease outbreaks

spread out by domestic cattle (Maala 2001), and traditional hunting conducted

by upland indigenous people (IP) communities who share their living space with

the species (Long *et al.* 2018). As a consequence, the tamaraw is now found in a

few isolated populations scattered across Mindoro, with MIBNP holding the

largest number (Matsubayashi *et al.* 2010; Wilson & Mittermeier 2011; Long

et al. 2018). The MIBNP tamaraw population is restricted to a single location of

less than <3,000 ha, hereafter referred to as the Core Zone of Monitoring (CZM,

see Fig. 2).

6

Given its critical importance for the conservation of this iconic species, the MIBNP tamaraw population has been the cornerstone of past, present, and future conservation plans (Maala 2001; Long *et al.* 2018). In an attempt to reinforce tamaraw protection, wildlife managers came to an agreement with IP living in the reserve in 2016, declaring a 1,600 ha no-hunting zone area (NHZ) within the CZM (Fig. 2). Unfortunately, one unforeseen consequence of the NHZ establishment for tamaraw is an increased pressure by traditional hunting and poaching at its immediate boundaries. MIBNP rangers regularly report the setting of snare and spear traps in the border areas, and violation of the NHZ agreement is not uncommon, with several deaths of tamaraws confirmed in the last year (Boyles comm. pers). Furthermore, observations by rangers suggest that tamaraw population density is growing within the NHZ, which could increase the risks of crowding. In this context, it is possible that negative density-dependence of demographic parameters will affect the population in the near future, undermining long-term conservation strategies. The population dynamics of tamaraws in MIBNP has been poorly documented so far, and a detailed examination of the population growth rate is therefore needed.

Here, we took advantage of a 21 year time series of abundance data to provide a detailed analysis of the tamaraw population dynamics in the CZM of MIBNP. We test the hypothesis that this large herbivore population could be facing negative density-dependence effects. We predict (*i*) that because of the substantial conservation effort in the last decade, overall population growth rate (noted *r*) should be statistically > 0. Yet (*ii*), because the no-hunting agreement area creates a spatial demarcation between risky and safe habitats, we expect local growth rates to differ in space and predict a decrease of *r* in areas further

7

from the main rangers' base camps and patrolling routes. Finally (*iii*), given the

intensive conservation efforts in the CZM, and relative safety of tamaraw in the

CZM, we predict a negative association between r and tamaraw abundance

within the NHZ as a result of negative density dependence effects. To test the

first two hypotheses, we computed the population growth rate of the tamaraw

population at two spatial scales: over the entire CZM, and at specific locations

used in annual tamaraw counts (hereafter referred to as vantage points). To test

our third hypothesis of density-dependent demographic responses, we fitted and

compared three population dynamics models: the baseline exponential model

(no density-dependence), a logistic model to test for a linear relationship

between r and annual abundance, and a particular formulation of Gompertz's

model to test for a decrease of r in time.

MATERIAL AND METHODS

Study site

The study area is located within the Mounts Iglit-Baco Natural Park (N12°54′

E121°13′) in the south-central part of the island of Mindoro. MIBNP harbors the

largest population of tamaraw, found in a single 3,000 ha area located on the

south-west edge of the protected area (Fig. 2) where most patrolling and

observation activities are carried out. In 2016, a 1,600 ha area inside the CZM

was declared as a no-hunting zone with the consensus of local IPs. There are

currently 15 park rangers, which are divided into two or three teams to monitor

the area on a regular basis. The teams patrol regular routes within the CZM,

centered around the three permanent base camps. The CZM and NHZ areas

represent approx. 3 and 1.5% of MIBNP, respectively.

The CZM area is a rolling grassland plateau with an average elevation of

800m a.s.l., dominated by cogon (*Imperata cylindrica*) and wild sugarcane

(*Saccharum spontaneum*), and interspersed with numerous wooded creeks,

secondary forest fragments, and steep hills. In recent years, the invasive Siam

weed (*Chromolaena odorata*, locally called hagonoy) has been rapidly spreading

in the grassland area and now covers an unknown fraction of the tamaraw

habitat. The tropical climate is strongly seasonal with a rainy season from June

to October (mean temperature 26.5°C; mean rainfall: 150 mm), and a hot dry

season from November to May (mean temperature 29.8°C; mean rainfall: 5

mm). Apart from humans, the only potential predator to tamaraw is the

reticulated python (*Malayopython reticulatus*) that is capable of preying upon

calves or yearlings, though no such predation observations had been made so far.

Tamaraw counts

Every year since 2000, the Tamaraw Conservation Program (TCP), a program of

the Philippine Department of Environment and Natural Resources (DENR), has

conducted tamaraw counts across the CZM, yielding an annual index of relative

population abundance. The tamaraw counting area consists of 18 vantage points,

covering nearly 2,200 ha within the CZM (Fig. 2). The tamaraw counting

process is rather invasive for the MIBNP ecosystem because it involves the

burning of grasslands a few weeks ahead of each count to increase visibility and

to attract tamaraw to areas with nutrient-rich young grasses. For the past 21

years, between 1,200 and 1,500 ha of grasslands were intentionally burned for

the purpose of monitoring tamaraw abundance.

9

During the annual counts, individual tamaraws were counted simultaneously

at the 18 vantage points for two hours at dusk and dawn over a period of four

consecutive days in late March or early April. For the 8 sessions and at each

vantage points, at least two observers spotted and recorded all animals seen,

writing down observation time, approximate location, and sex and age-category

of individuals (split into calves, yearlings, sub-adults, and adults). This sampling

designed remained unchanged for the 21 years, albeit the number of observers

could vary from year to year. Immediately after the completion of the 8 sessions,

raw data (i.e. the number of seen animals T) were cross-checked and cleaned for

potential double observations. The total number of different animals corrected

for double counts was then derived for each vantage point and year, and

constitute the consolidated counts C. These annual counts comprise the only

available source of information about tamaraw population trends and dynamics

for officials to use for conservation management decisions. Note that, because of

the COVID-19 pandemic, the tamaraw counts in 2020 did not take place.

Data analyses

For populations of long-lived and endangered species, robust and accurate

estimations of r remain difficult to obtain. Knowledge about r requires either the

estimation of demographic rates combined into a Leslie matrix (Leslie 1945) or

the availability of long-term series of population abundance, preferably on an

annual basis. Because we could not individually identify tamaraws, neither mark

them for ethical reasons (Mccarthy & Parris 2008), applying methods of

abundance estimations based on capture-recapture (see Schwarz & Seber 1999,

for a review) was not possible, an only annuals counts could be analyzed with

10

auto-regressive statistical models. Working with abundance index data to derive

an empirical estimates of *r*, however, comes with serious methodological issues

(Îto 1972; Lindley 2003). Over the last decades, population counts have

repeatedly been associated with a large sampling variance (*e.g.* Caughley 1977;

Morellet *et al.* 2007), and coefficients of variation for population abundance

close to 30% have been reported (Lubow & Ransom 2016). From a conservation

perspective, a large sampling variance in the input data for estimating *r* is a

major pitfall because it leads to over-optimistic results, usually from an

over-estimation of both the point estimate and its precision (Lindley 2003).

Despite the availability of appropriate statistical methods to account for the

sampling variance of population counts (Kalman filter, De Valpine & Hastings

2002), such tools remain seldom used in practice in conservation.

Here, we analyzed the consolidated number of observed tamaraws (*C*), as

judged by observers to be different individuals, for each of the 18 vantage points,

split by years ($C_{s,t}$, where indice $s \in \{1, \ldots, 18\}$ stands for the vantage point, and

$t \in \{2000, \ldots, 2019\}$ stands for the time in years. We implemented a state-space

model to tease apart process from sampling variances, making the assumption

that double and under-counts compensate and are randomly distributed from

year to year. We worked in a Bayesian framework (*e.g.* Kéry & Schaub 2011)

but note this is fully equivalent to a Kalman filter with a frequentist approach. To

do so, we defined $N_{s,t}$ as the unobserved 'true' abundance linked to $C_{s,t}$ by a

random effect corresponding to the process variance (σ_c^2) on the log scale to

ensure positive values for abundance. Our baseline observation model was:

$$\log(C_{s,t}) \sim \mathcal{N}(\log(N_{s,t}), \sigma_c^2)$$

11

231   Accordingly, we computed the population growth rate $r_{s,t}$ of the tamaraw

232 population at each vantage points s, as:

$$r_{s,t} = \log\left(\frac{N_{s,t+1}}{N_{s,t}}\right),$$

233 from counts corrected for the sampling variance, thus returning an unbiased

234 estimation of the population growth rate. The overall population growth of the

235 tamaraw population of MIBNP is then given by:

$$\bar{r} = \frac{1}{19}\sum_{t=1}^{19}\log\left(\frac{N_{t+1}}{N_t}\right),$$

where $N_t = \sum_{s=1}^{18} N_{s,t}$. We informed N_t with the total number of animals seen

each count session (T_t) and linked counts at the vantage point and population

levels with the following observation equation: $T_t \sim \mathcal{N}(N_t/p, \sigma_{tot}^2)$. Because the

consolidated number of tamaraws C_t was much smaller than the total number

seen T_t before double-counts were removed, we included a constant proportion

parameter p, similar to a "detection rate". We acknowledge C_t and T_t are

strongly correlated and certainly share the same information about tamaraw

abundance. However, this approach allowed us to fully use all available

information about tamaraw counts fully, including both the knowledge of

experienced rangers and the raw number of detected tamaraws with no *a priori*

interpretation. In addition, we could input missing values at the vantage point

level for years 2000 to 2003 and 2005 thanks to the T_t constraint on N_t. The data

imputation of abundance back in early years of monitoring allowed us to

compare the spatio-temporal variation in growth rate r_i at the sub-population

scale, and to quantity the effect of human disturbance at the periphery of the

CZM.

To test our third hypothesis we investigated potential density-dependence in

the annual growth rate by fitting two models classically used to detect negative

density dependence (Dennis & Taper 1994) the logistic and a Gompertz growth

model, to be compared with an exponential growth model. The exponential

growth model was fitted implicitly for the estimation of the average population

growth rate $\bar{r}$:

$$\log(N_{t+1}) = \log(N_t) + \bar{r},$$

258   Then we fitted a logistic growth model to the total number of tamaraws seen

259 per year N_t as follows:

$$\log(N_{t+1}) = \log(N_t) + r_m \times \left(1 - \frac{N_t}{K_l}\right),$$

260 where r_m is the maximum population growth for the tamaraw, and K_l is the

261 carrying capacity of the MIBNP. Likewise, we tested if the annual population

262 growth rate would decrease in time with population density by fitting a particular

263 formulation of the Gompertz model (Norton's formulation, see Tjørve & Tjørve

264 2017):

$$\log(N_{t+1}) = \log(N_1) + \log\left(\frac{K_G}{N_1}\right) \times \left(1 - e^{-\beta \times t}\right),$$

265   Because the pair of parameters r_m and K_l (logistic) and β and K_G (Gompertz)

13

are not separately identifiable (Lebreton & Gimenez 2013), we used informative

priors for r_m and β based on the reported maximum population growth for other

large bovids (see Traill *et al.* 2007, for a similar approach). Accordingly, we set

a moderately informative prior value of mean 0.3 and a precision of 0.1 in our

models (Table 1). We tested the occurrence of density-dependence processes in

the MIBNP tamaraw population by explicitly computing the probability (π_k) for

our abundance time series to be generated by each of the three candidate

demographic models during the model fit. At each MCMC step, a multinomial

variable stated which of the exponential, Gompertz and logistic model was the

most likely underlying model given the data by taking the value one and zero

otherwise. The three probabilities π_k were estimated as the number of times each

models received a one divided by the number of MCMC iterations.

We fitted our statistical model to the tamaraw count data with JAGS 4.0

(Plummer *et al.* 2003), with 3 MCMC chains and a burn-in of 40,000 iterations.

We obtained parameter estimates with an additional 30,000 iterations and a

thinning factor of 5 (therefore giving a total of 6,000 MCMC samples). We

checked model convergence visually by investigating the mixing of MCMC

chains, along with the $\hat{R}$ statistics (Brooks & Gelman 1998). With the exception

of r_m (see above), we set uninformative priors with large variance for all other

parameters to be estimated, and checked the sensitivity of our results to the

average value of the priors by replicating the analyses with variable mean and

initial values for prior distributions. We report all parameters estimates as the

mean of the posterior distributions along with the 95% percentile for the credible

intervals following Louis & Zeger (2008): $_{95\%low}\,estimate\,_{95\%up}$.

Note that despite the fact that we analysed the number of tamaraws as

14

described above, we chose not to display such numbers on figures and to report a

standardized abundance ($C_t' = C_t/max(C_t))$) instead. This transformation applies

to the estimated carrying capacities of the Gompertz and logistic models,

$K' = K/max(C_t)$. We did not provide raw counts to avoid confusion between the

index of relative tamaraw abundance we worked with, and a real population size

estimator that was not implemented at MIBNP. With this variable transformation

K_l and K_G can no longer be interpreted as the maximum number of tamaraws but

the maximum relative abundance (C_t') we can expect to observe at MIBNP.

Reported values of the carrying capacities K_l' and K_g' can hence take values > 1

because the carrying capacity has not yet been reached. Biologically speaking,

we put more emphasize on how far from the carrying capacity the current

abundance of tamaraws is (C_t/K) than of the number of animals *per se*.

RESULTS

From our baseline model of tamaraw counts, we estimated an average annual

growth rate of $\bar{r} = {}_{0.01}\,0.05\,{}_{0.09}$ for the MIBNP population from 2000 to 2021,

with an associated variance process equal to $\sigma_{proc} = {}_{0.25}\,0.31\,{}_{0.38}$ (Fig. 2). As

expected for count data, the sampling variance was large $\sigma_c = {}_{7.18}\,9.21\,{}_{9.98}$,

suggesting frequent double counts of the same individuals. That the MIBNP

tamaraw population increased in size by c.a. 5% per year and was statistically

different from 0 gives clear support of our first hypothesis, which predicted that

tamaraw abundance has increased as a result of successful protection efforts in

MIBNP.

For the vantage point specific growth rates (r_s), we observed a marked

variability in space (Fig. 3; see Supplementary Table S1) as shown by the fact

that growth rates ranged between $r_s = {}_{-0.18} -0.42 {}_{0.20}$ at Tarzan vantage point

and $r_s = {}_{0.00} 0.14 {}_{0.26}$ at Bato Fidel and Bayokbok vantage points. Vantage point

specific growth rates showed strong spatial structuring (Fig 3). Without

exception, $r_s < 0$ were located at the periphery of the NHZ (Iyan, Tarzan,

Saligue East, Mibluan, Fangandatan), although r_s of a few vantage points was

> 0 despite being located on the NHZ border (Nagbobong, Malibayong, Talafu

West and East). Together, these results support our hypothesis suggesting an

ongoing contraction of the tamaraw range at MIBNP after 2000, and the

disappearance of individuals at the periphery of the CZM.

We explicitly estimated the probabilities π_k that the exponential, logistic, or

Gompertz models would be the underlying process generating the observed

variation in tamaraw abundance (proxy of Baye's factor). The estimated

probabilities for the three population dynamics model were $\pi_1 = 0.10$, $\pi_2 = 0.90$

and $\pi_3 = 0.00$, respectively, supporting our hypothesis of density-dependence

feed-backs in the tamaraw population of MIBNP. Point estimates for the two

parameters of the logistic growth function were $r_m = {}_{0.07} 0.08 {}_{0.09}$ for the

maximum population growth rate and $K'_l = {}_{1.16} 1.37 {}_{1.62}$ for the carrying

capacity. Although Gompertz model received no statistical support, the

estimated coefficients were ($\beta = {}_{0.04} 0.05 {}_{0.07}$), and $K'_G = {}_{1.20} 1.41 {}_{1.66}$ for the

carrying capacity.

DISCUSSION

The isolated tamaraw population in MIBNP has been increasing in size since 2000 (Fig. 1). Undoubtedly, a positive growth rate of 0.05 over 21 years of monitoring is a great success for such a highly endangered species, both locally and for global conservation efforts. However, the devil lies in the details, and two additional results darken this otherwise positive picture. First, we found support for the hypothesis of a contraction of the tamaraw distribution, evidenced by a clear spatially structured pattern in the population growth rates. Second, our results highlight a progressive decrease of the average growth rate, suggesting density-dependent effects at the population level. The population dynamics we document clearly indicate that new and urgent conservation actions are needed to secure the future of the tamaraw population in MIBNP.

Long-run Positive Growth of Tamaraws

The average population growth rate of $\bar{r} = 0.05$ gives strong evidence that successful conservation policies over the past two decades have led to an increase in tamaraw abundance. Nonetheless, an annual growth of 5% is relatively low compared to other similar-sized or even larger cattle species (Table 1). For instance, a population of African buffalo (*Syncerus caffer*) in South Africa grew at a rate of 12% per year for 28 years (Jolles 2007), and values close to 30% per year have regularly been recorded for bison (*Bison bison*) and banteng (*Bos javanicus*), despite the fact that these species are between 2 and 2.5 times larger in size than the tamaraw. From this brief comparative approach, we would expect a substantially higher average growth

17

rate for tamaraws if the population was in a colonizing phase. Given that the

tamaraw is secure within the NHZ, our results suggest that specific

environmental factors are limiting its growth rates in MIBNP.

A first limiting factor could be related to the fact that the most available and

accessible habitat for the tamaraw in MIBNP is grassland (83% of the CMZ),

which is dominated by competitive pioneer species such as cogon and wild

sugarcane. Both grasses appear to be *de facto* prominent in the tamaraw diet.

Although these grasses provide tamaraws with abundant and high quality forage

at the early stages of plant growth, in later stages of development these species

are characterized by much lower nutritive value, especially during the dry

season. Observations in the field indicate that once grass growth is complete, the

stems exceeds tamaraw height (300cm vs. < 120 cm, respectively) and dry out,

thus becoming both inaccessible and unpalatable. It is therefore possible that

lack of food resource accessibility and availability could be limiting the growth

rate for the population.

The regular burning of the grassland habitat for population count may also be

impacting plant composition in the CZM, with implications for tamaraw

population dynamics. On the one hand, regular burning promotes the

regeneration of new grass shoots, providing a nutrient-rich food source for

tamaraw. On the other hand, previous observations suggest that regular burning

could promote Siam weed expansion (Nath *et al.* 2019), which appears to be

rapidly spreading throughout the study site. Like other grazing herbivores

(Rozen-Rechels *et al.* 2017), tamaraw avoid this invasive plant, and so the

increase of Siam weed could ultimately decrease the CMZ carrying capacity

(Dumalisile & Somers 2017). In addition to promoting the spread of a

18

non-palatable invasive species, it is possible that, more generally, the regular burning creates ecological conditions that are unfavorable for tamaraw. Although burning creates high-quality forage in the short-term, such an intensive habitat management practice is likely to cause a progressive decrease in habitat quality over the long term.

Here it should be noted that tamaraw feeding preference is not well understood, and the species might be a mixed-feeding or a browsing herbivore (*sensu* Hofmann 1989). Indeed, tamaraw are suspected to be ecologically flexible, as indicated by the identification of a new tamaraw population in montane forest habitat in the upland inner Mindoro (Schütz 2019). In this context, the CZM might actually provide ecological conditions whose intrinsic limits are artificially reduced due to the long-lasting and highly invasive habitat management policies, and that the negative impacts of these practices are progressively starting to exceed their short-term beneficial effects.

Possible Impacts of Human Disturbance and Hunting

Human activities could also account for the low tamaraw population growth rate we report. Tamaraw have always been traditionally hunted by IP in Mindoro, and despite the "no-hunting agreement" established in 2016, both intentional and unintentional tamaraw deaths still occur. Moreover, the tamaraw is sought after for bushmeat by lowlander poachers. Because hunting from IP and poaching by lowlanders is difficult to control in MIBNP, it has likely lead to an increased mortality of all age-classes of tamaraws to an unknown, but perhaps considerable, extent. While the positive growth of the tamaraw population over the past two decades is evidence of a successful decrease in hunting pressure in

19

the CZM, the marked spatial heterogeneity we document, with negative growth

rates at the periphery of the counting area, could indicate a lower protection

efficiency in areas further away from the main patrolling routes and ranger base

camps. Furthermore, this result support our hypothesis suggesting an unexpected

contraction of the tamaraw range at MIBNP and the disappearance of individuals

at the periphery of the CZM. In other words, the CZM area may function like a

source-sink system, with the more central and better patrolled areas exhibiting

positive growth rates, and more peripheral areas with higher hunting pressure

serving as population sinks. Again, our findings provide some support for this

idea, as local growth rates in the core area of the NHZ have values as high as

14% (Fig. 2), which is more in line with the average growth rates that we

reviewed for large bovids (Table 1).

Space Needed for Sustaining Population Growth

The gradual decrease of the tamaraw population growth rate in MIBNP, coupled

with the overall abundance in the CZM, suggests a type of density-dependence is

occurring, where a change in demographic rates lowers the number of

individuals added to the population from year to year (Fowler 1987; Sinclair

1989). Predicted growth rates from the lowest to the highest abundance varied

between $r = 0.08$ and $r = 0.01$ for tamaraw at MIBNP. From the current data,

our best model suggested a carrying capacity of 1.4, meaning that only 40% of

growth is left possible before reaching the carrying capacity in MIBNP. This

explanation is the *biological* hypothesis of density-dependence, which is well

known and repeatedly detected in populations of large mammals (Bonenfant

et al. 2009). Yet, in the absence of accurate demographic rate estimations for the

20

tamaraw population in MIBNP, we encounter the limits of the so-called

pattern-oriented approach (Krebs 2002), and are unable to link real changes in

demographic rates with population abundance. Consequently, we are unable to

determine if tamaraws experience increased mortality, reduced reproduction, or

are more willing to disperse. Further investigation of these possibilities is

needed, as each situation would require different management approaches.

Another behaviour associated with high population abundance is dispersal

(Matthysen 2005). In mammals, sub-adult individuals are more likely to disperse

at high than low population abundance (e.g. Loe *et al.* 2009, for an example on

red deer). At the high population abundance we documented in MIBNP, we

would expect a greater number of dispersing individuals to venture outside of the

NHZ, where the risk of mortality is high because of traditional hunting. We

expect that, if competition for food or mating becomes strong, as is suggested by

our results, then the main tamaraw population may experience an increased net

loss of animals in the near future. In addition to jeopardizing the fate of the

population by decreasing survival probabilities of sub-adults that are key to

maintaining the future of the population, the empirical evidence for

density-dependence that we report would limit the ability of this population to

act as a source. With a combination of low recruitment and low survival of

sub-adults, old individuals would quickly become over-represented in the

population, which could depress growth even further.

Implication for tamaraw conservation

The main conservation goals for the MIBNP tamaraw populations are (1) to

ensure its long-term viability, and (2) the expansion of the population, either

21

through natural emigration or through transcolation. In regards to population

expansion, evidence of density-dependence in MIBNP is worrisome because it is

usually associated with increased competition between individuals for food or

territorial space. Our results therefore suggest that tamaraws may already be

experiencing a food resource shortage, which could make the population more

prone to parasitism (Stien *et al.* 2002) or epizootics (Davidson & Doster 1997).

Such results indicate that the tamaraw population is becoming potentially less

viable on the long-run. A similar situation has been reported for a banteng (*Bos*

javanicus) population in Alas Purwo National Park, Indonesia. Despite an

increasing abundance, the banteng population has become confined to an

artificially maintained grazing area, which has caused increased predation

pressure of juveniles (JB, pers. obs).

Current conservation measures assume that the tamaraw population growth

will continue in the next decade, without a need to mitigate increased pressures

thought to be impacting the periphery of the CZM or to address the pressure of

density dependence on the population. Our results indicate that strategic

adjustments need to be made as part of the MIBNP Management Plan

specifically, and the island-wide Tamaraw Conservation Management and Action

Plan (DENR 2019) more generally. First, two main issues must be urgently

addressed: (*i*) increase the available space for the species to mitigate the effects

of density-dependence and maintain a positive growth rate, which must be

combined with (*ii*) increasing the effectiveness of ranger patrols in the periphery

area of the CZM to reduce the recurrent poaching pressure from Filipino

lowlanders and losses due to traditional hunting. Both targets need to involve

close collaboration with local IP communities to create conditions for a more

sustainable land-use and traditional hunting system. With more effective

protection and suitable habitat management measures in place, it is possible that

the average tamaraw growth rate will increase in MINBP, and that the

density-dependent demographic effects that we report can be reversed. Second,

when investigating the feasibility of translocating animals from this source

population to other sites, our estimation of the annual growth rates and the

documented density-dependence responses should be taken into account. For

example, the number of tamaraws removed should be equal to the number of

recruited individuals each year that is necessary for the population not to

decrease.

Conclusion

Despite increased efforts to protect key habitats for threatened species, ongoing

habitat loss, degradation, and fragmentation are still significant threats to large

mammal populations globally (Ripple *et al.* 2015). Consequently, a greater

number of wild herbivores (and their natural predators) are expected to be

impacted by intensively managed landscapes in the future. So far, conservation

responses have primarily focused on protecting large areas of land. Securing

large areas of habitat is often an effective conservation approach, but may not

always be a viable strategy to protect highly threatened species, as illustrated our

study on the tamaraw population in MIBNP. More generally, we show that

positive growth rates for a threatened species could potentially hide a darker

picture. First, the growth of a population that is confined to a small conservation

area may reflect a range contraction as a result of anthropogenic pressures, rather

than the effect of increased reproduction and survival. Second, because of this,

23

the population could face important demographic challenges that, in the long

term, lead to new threats. In this context, there is an increasing need to

understand the interaction between anthropogenic activities and the population

dynamics of protected species. Our results suggest that, in some situations,

conservation strategies designed to protect small, isolated populations may lead

to long-term demographic issues that undermine species recovery. On the

positive side, our findings also show how detailed analyses of relative population

abundance data, even without absolute population size estimators, can help

provide information that assists in the development of conservation measures

that are essential for large mammal conservation.

ACKNOWLEDGMENTS

A lot of people are to thank here. We are grateful to Alvaro Gonzalez. We must

acknowledge the invaluable help of all rangers, students and volunteers who contributed

to the annual tamaraw counts.

REFERENCES

Beissinger, S.R. & Westphal, M.I. (1998). On the use of demographic models of
population viability in endangered species management. *The Journal of wildlife
management*, pp. 821–841.

Bonenfant, c., Gaillard, J.M., Coulson, T., Festa-Bianchet, M., Loison, A., Garel, M.,
Loe, L.E., Blanchard, P., Pettorelli, N., Owen-Smith, N., Du Toit, J. & Duncan, P.
(2009). Empirical evidence of density-dependence in popualtions of large herbivores.
Adv. Ecol. Res., 41, 313–357.

Boyce, M.S. (1992). Population viability analysis. *Annual Review of Ecology and
Systematics*, 23, 481–506.

Brooks, S.P. & Gelman, A. (1998). General methods for monitoring convergence of
iterative simulations. *Journal of Computational and Graphical Statistics*, 7, 434–455.

Buys, D. & Dott, H. (1991). Population fluctuations and breeding of eland *Taurotragus oryx* in a western Transvaal nature reserve. *Koedoe*, 34, 31–36.

Caughley, G. (1977). *Analysis of vertebrate populations*. Wiley, London. Graeme Caughley. 'A Wiley-Interscience Publication' Includes index. Bibliography: p.217-228.

Caughley, G. (1994). Directions in conservation biology. *Journal of animal ecology*, pp. 215–244.

Choquenot, D. (1991). Density-dependent growth, body condition, and demography in feral donkeys: testing the food hypothesis. *Ecology*, 72, 805–813.

Choquenot, D. (1993). Growth, body condition and demography of wild banteng (*Bos javanicus*) on Cobourg Peninsula, northern Australia. *Journal of Zoology*, 231, 533–542.

Cornélis, D., Melletti, M., Korte, L., Ryan, S.J., Mirabile, M., Prin, T. & Prins, H.H. (2014). *African buffalo Syncerus caffer (Sparrman, 1779)*, Cambridge University Press, pp. 326–372.

Courchamp, F., Clutton-Brock, T. & Grenfell, B. (1999). Inverse density dependence and the allee effect. *Trends in ecology & evolution*, 14, 405–410.

Custodio, C.C., Lepiten, M.V. & Heaney, L.R. (1996). *Bubalus mindorensis*. *Mammalian Species*, pp. 1–5.

Davidson, W. & Doster, G. (1997). Health characteristics and white-tailed deer population density in the southeastern united states.

De Valpine, P. & Hastings, A. (2002). Fitting population models incorporating process noise and observation error. *Ecological Monographs*, 72, 57–76.

Dennis, B. & Taper, M. (1994). Density dependence in time series observations of natural populations: estimation and testing. *Ecological Monographs*, 64, 205–224.

Dumalisile, L. & Somers, M.J. (2017). The effects of an invasive alien plant (*Chromolaena odorata*) on large African mammals. *Nature Conservation Research.* , 2, 102–108.

Eberhardt, L.L. (1977). Optimal policies for conservation of large mammals, with special references to marine ecosystems. *Environmental Conservation*, 4, 205–212.

Eberhardt, L.L. (2002). A paradigm for population analysis of long-lived vertebrates. *Ecology*, 83, 2841–2854.

Fowler, C.W. (1987). A review of density dependence in populations of large mammals. *Curr. Mammal.*, 1, 401–441.

Fuller, J.A., Garrott, R.A., White, P., Aune, K.E., Roffe, T.J. & Rhyan, J.C. (2007).
Reproduction and survival of Yellowstone bison. *Journal of Wildlife Management*,
71, 2365–2372.

Gates, C. & Larter, N. (1990). Growth and dispersal of an erupting large herbivore
population in northern Canada: the Mackenzie wood bison (*Bison bison athabascae*).
Arctic, pp. 231–238.

Haque, N. (1990). *Study on the ecology of wild ungulates of Keoladeo national Park*
Bharatpur, Rajasthan. Ph.D. thesis, Aligarh Muslim University.

Heinen, J.T. & Paudel, P.K. (2015). On the translocation of wild asian buffalo (**?**) in
Nepal: are feral backcrosses worth conserving? *Conservation Science*, 3.

Henle, K., Sarre, S. & Wiegand, K. (2004). The role of density regulation in extinction
processes and population viability analysis. *Biodiversity and Conservation*, 13, 9–52.

Heude, P. (1888). Études sur les ruminants de l'Asie orientale. cerfs des Philippines et
de l'Indo-Chine. *Mémoires Concernant l'Histoire Naturelle de l'Empire Chinois par*
des Péres de la Compagnie de Jésus, 2, 1–10.

Hofmann, R.R. (1989). Evolutionnary steps of ecophysiological adaptation and
diversification of ruminants: a comparative view of their digestive system. *Oecologia*,
78, 443–457.

Hone, J., Duncan, R.P. & Forsyth, D.M. (2010). Estimates of maximum annual
population growth rates (r_m) of mammals and their application in wildlife
management. *Journal of Applied Ecology*, 47, 507–514.

Hones, J. & Clutton-Brock, T. (2007). Climate, food, density and wildlife population
growth rate. *J. Anim. Ecol.*, ?, ?–?

Îto (1972). On the methods for determining density-dependence by means of regression.
Oecologia, 10, 776–778.

Jolles, A.E. (2007). Population biology of african buffalo (*Syncerus caffer*) at
Hluhluwe-iMfolozi Park, South Africa. *African Journal of Ecology*, 45, 398.

Kemp, N., Dilger, M., Burgess, N. & Van Dung, C. (1997). The saola *Pseudoryx*
nghetinhensis in Vietnam – new information on distribution and habitat preferences,
and conservation needs. *Oryx*, 31, 37–44.

Kéry, M. & Schaub, M. (2011). *Bayesian population analysis using WinBUGS: a*
hierarchical perspective. Academic Press.

Koláčková, K., Hejcmanova, P., Antonínová, M. & Brandl, P. (2011). Population
management as a tool in the recovery of the critically endangered Western Derby
eland *Taurotragus derbianus* in Senegal, Africa. *Wildlife Biology*, 17, 299–311.

26

Krasiński, Z.A. (1978). Dynamics and structure of the european bison population in the
białowieża Primeval Forest. *Acta Theriologica*, 23, 3–48.

Krebs, C.J. (2002). Two complementary paradigms for analysing populations dynamics.
Philosophical Transactions of the Royal Society of London, Series B, 357,
1211–1219.

Krüger, O. & Lindström, J. (2001). Habitat heterogeneity affects population growth in
goshawk *Accipiter gentilis*. *Journal of Animal Ecology*, 70, 173–181.

Lande, R. (1993). Risks of population extinction from demographic and environmental
stochasticity and random catastrophes. *The American Naturalist*, 142, 911–927.

Langwig, K.E., Frick, W.F., Bried, J.T., Hicks, A.C., Kunz, T.H. & Marm Kilpatrick, A.
(2012). Sociality, density-dependence and microclimates determine the persistence of
populations suffering from a novel fungal disease, white-nose syndrome. *Ecology*
letters, 15, 1050–1057.

Larter, N., Sinclair, A., Ellsworth, T., Nishi, J. & Gates, C. (2000). Dynamics of
reintroduction in an indigenous large ungulate: the wood bison of northern Canada. 3,
299–309.

Lebreton, J.D. & Gimenez, O. (2013). Detecting and estimating density dependence in
wildlife populations. *Journal of Wildlife Management*, 77, 12–23.

Leslie, P.H. (1945). On the use of matrices in population mathematics. *Biometrika*, 33,
182–212.

Lindley, S.T. (2003). Estimation of population growth and extinction parameters from
noisy data. *Ecological Applications*, 13, 806–813.

Loe, L.E., Mysterud, A., Veiberg, V. & Langvatn, R. (2009). Negative density-dependent
emigration of males in an increasing red deer population. *Proceedings of the Royal*
Society B: Biological Sciences, 276, 2581–2587.

Long, B., Schütz, E., Appleton, M., Gonzales, A., Holland, J., Lees, C., Marandola, E.,
Natural, V.C.J., Pineda-David, M.T.J., Salao, C., Slade, J., Tan, E.H.P., Tiongson, L.
& Young, S. (2018). Review of tamaraw (*Bubalus mindorensis*) status and
conservation actions. *BULLtin*, 1, 18–34.

Louis, T.A. & Zeger, S.L. (2008). Effective communication of standard errors and
confidence intervals. *Biostatistics*, 10, 1–2.

Lubow, B.C. & Ransom, J.I. (2016). Practical bias correction in aerial surveys of large
mammals: Validation of hybrid double-observer with sightability method against
known abundance of feral horse (equus caballus) populations. *PLoS One*, 11,
e0154902.

Maala, C.P. (2001). Endangered Philippine wildlife species with special reference to the
Philippine eagle (*Pithecophaga jefferyi*) and tamaraw (*Bubalus mindorensis*). *Journal*
of International Development and Cooperation, 8, 1–17.

Mace, G.M., Collar, N.J., Gaston, K.J., Hilton-Taylor, C., Akçakaya, H.R.,
Leader-Williams, N., Milner-Gulland, E.J. & Stuart, S.N. (2008). Quantification of
extinction risk: Iucn's system for classifying threatened species. *Conservation*
biology, 22, 1424–1442.

Mathur, V.B. (1991). *The ecological interaction between habitat composition, habitat*
quality and abundance of some wild ungulates in India. Ph.D. thesis, University of
Oxford.

Matsubayashi, H., Boyles, R.M., Salac, R.L., Del Barrio, A., Cruz, L., Garcia, R.A.,
Ishihara, S. & Kanai, Y. (2010). Present status of tamaraw (*Bubalus mindorensis*) in
Mt. Aruyan, Mindoro, Philippines. *Tropics*, 18, 167–170.

Matthysen, E. (2005). Density-dependent dispersal in birds and mammals. *Ecography*,
28, 403–416.

Mccarthy, M.A. & Parris, K.M. (2008). Optimal marking of threatened species to
balance benefits of information with impacts of marking. *Conservation Biology*, 22,
1506–1512.

Morellet, N., Gaillard, J.M., Hewison, A.J.M., Ballon, P., Boscardin, Y., Duncan, P.,
Klein, F. & Maillard, D. (2007). Indicators of ecological change: new tools for
managing populations of large herbivores. *J. Appl. Ecol.*, 44, 634—643.

Mugabo, M., Perret, S., Legendre, S. & Le Galliard, J.F. (2013). Density-dependent life
history and the dynamics of small populations. *Journal of Animal Ecology*, 82,
1227–1239.

Mysterud, A., Bartoń, K., Jedrzejewska, B., Krasiński, Z., Niedziałkowska, M., Kamler,
J., Yoccoz, N. & Stenseth, N. (2007). Population ecology and conservation of
endangered megafauna: the case of european bison in białowieża Primeval Forest,
Poland. *Animal Conservation*, 10, 77–87.

Nath, A., Sinha, A., Lahkar, B.P. & Brahma, N. (2019). In search of aliens: Factors
influencing the distribution of *Chromolaena odorata* L. and Mikania micrantha Kunth
in the Terai grasslands of Manas National Park, India. *Ecological engineering*, 131,
16–26.

Neel, M.C., Leidner, A.K., Haines, A., Goble, D.D. & Scott, J.M. (2012). By the
Numbers: How is Recovery Defined by the US Endangered Species Act? *BioScience*,
62, 646–657.

Nicholls, A., Viljoen, P., Knight, M. & Van Jaarsveld, A. (1996). Evaluating population
persistence of censused and unmanaged herbivore populations from the Kruger
National Park, South Africa. *Biological Conservation*, 76, 57–67.

Nicholson, A.J. (1933). The balance of animal populations. *J. Anim. Ecol.*, S2, 132–178.

Ottichilo, W.K., De Leeuw, J., Skidmore, A.K., Prins, H.H. & Said, M.Y. (2000).
Population trends of large non-migratory wild herbivores and livestock in the Masai
Mara ecosystem, Kenya, between 1977 and 1997. *African journal of Ecology*, 38,
202–216.

Peterson, C.H. & Black, R. (1988). Density-dependent mortality caused by physical
stress interacting with biotic history. *American Naturalist*, 131, 257–270.

Plummer, M. *et al.* (2003). Jags: A program for analysis of bayesian graphical models
using gibbs sampling. In: *Proceedings of the 3rd international workshop on
distributed statistical computing*. Vienna, Austria., vol. 124, p. 10.

Ripple, W.J., Newsome, T.M., Wolf, C., Dirzo, R., Everatt, K.T., Galetti, M., Hayward,
M.W., Kerley, G.I., Levi, T., Lindsey, P.A. *et al.* (2015). Collapse of the world's
largest herbivores. *Science advances*, 1, e1400103.

Rozen-Rechels, D., te Beest, M., Dew, L.A., le Roux, E., Druce, D.J. & Cromsigt, J.P.
(2017). Contrasting impacts of an alien invasive shrub on mammalian savanna
herbivores revealed on a landscape scale. *Diversity and Distributions*, 23, 656–666.

Schwarz, C. & Seber, G.A.F. (1999). A review of estimating animal abundance III.
Statistical Science, 14, 427–456.

Schütz, E. (2019). Tamaraw verification surveys in the upper Amnay watershed region,
Mindoro. *BULLtin*, 2, 6–15.

Shaffer, M.L. (1981). Minimum population sizes for species conservation. *BioScience*,
31, 131–134.

Sinclair, A.R.E. (1989). Population regulation in animals. In: *Ecological concepts : the
contribution of ecology to an understanding of the natural world* (eds. Cherrett, J.M.
& Bradshaw, A.D.). Blackwell Scientific Publications, Oxford, England., pp.
197–241.

Singer, F.J. & Norland, J.E. (1994). Niche relationships within a guild of ungulate
species in Yellowstone National Park, Wyoming, following release from artificial
controls. *Canadian Journal of Zoology*, 72, 1383–1394.

Steinmetz, R., Chutipong, W., Seuaturien, N., Chirngsaard, E. & Khaengkhetkarn, M.
(2010). Population recovery patterns of southeast asian ungulates after poaching.
Biological Conservation, 143, 42–51.

Stephens, P.A., Sutherland, W.J. & Freckleton, R.P. (1999). What is the allee effect?
Oikos, pp. 185–190.

Stien, A., Irvine, R.J., Ropstad, E., Halvorsen, O., Langvatn, R. & Albon, S.D. (2002).
The impact of gastrointestinal nematodes on wild reindeer: experimental and
cross-sectional studies. *J. Anim. Ecol.*, 71, 937–945.

Tjørve, K.M. & Tjørve, E. (2017). The use of gompertz models in growth analyses, and
new gompertz-model approach: An addition to the unified-richards family. *PloS one*,
12, e0178691.

Traill, L.W., Bradshaw, C.J. & Brook, B.W. (2007). Minimum viable population size: a
meta-analysis of 30 years of published estimates. *Biological conservation*, 139,
159–166.

Van Houtan, K.S., Halley, J.M., Van Aarde, R. & Pimm, S.L. (2009). Achieving success
with small, translocated mammal populations. *Conservation Letters*, 2, 254–262.

Van der Wal, E., Laforge, M.P. & McLoughlin, P.D. (2014). Density dependence in
social behaviour: home range overlap and density interacts to affect conspecific
encounter rates in a gregarious ungulate. *Behavioral Ecology and Sociobiology*, 68,
383–390.

Wilson, D. & Mittermeier, R. (2011). *Handbook of the mammals of the world. Vol. 2.*
Hoofed Mammals. Lynx Edicions, Barcelona.

TABLE CAPTIONS

Table 1 Estimated annual growth rates (r) of some large *bovinae* populations with body size comparable to the endangered tamaraw (*Bubalus mindorensis*). The observed population growth rate of tamaraws at Mounts Iglit-Baco Natural Park, Mindoro, Philippines, between 2000 and 2019 of $r = 0.05$ lies at the lower range of reported values.

Table 1

Species	Location	*r*	Reference
Bison bison	Mackenzie Bison Sanctuary, Canada	0.21	Gates & Larter (1990)
Bison bison	Mackenzie Bison Sanctuary, Canada	0.20	Larter *et al.* (2000)
Bison bison	Yellowstone, Wyoming, USA	0.07	Fuller *et al.* (2007)
Bison bison	Yellowstone, Wyoming, USA	0.30	Singer & Norland (1994)
Bison bonasus	Białowieża, Poland	0.07	Mysterud *et al.* (2007)[1]
Bison bonasus	Białowieża, Poland	0.18	Krasiński (1978)[1]
Bos gaurus	Thung Yai Naresuan Sanctuary, Thailand	0.31	Steinmetz *et al.* (2010)
Bos javanicus	Not applicable, model prediction	0.32	Hone *et al.* (2010)
Bos javanicus	Cobourg peninsula, Australia	0.00	Choquenot (1993)
Boselaphus tragocamelus	Kanha National Park, India	0.18	Mathur (1991)
Boselaphus tragocamelus	Keoladoe National Park, India	0.02 - 0.06	Haque (1990)
Bubalus arnee		0.02 - 0.06	Khatri et al. (2012)
Bubalus arnee	Koshi Tappu Wildlife Reserve, Nepal	0.03	Heinen & Paudel (2015)
Bubalus bubalis		0.03	Heinen (1993)
Bubalus bubalis		?	Freeland and Boulton (1990)
Bubalus mindorensis	Mount Iglit-Baco National Park, Philipines	0.05	This study
Pseudoryx nghetinhensis	Vietnam & Laos	0.05	Kemp *et al.* (1997)[b]
Syncerus caffer	Hluhluwe-iMfolozi Park, South Africa	0.12	Jolles (2007)
Syncerus caffer	Zakouma National Park, Chad	0.08	Cornélis *et al.* (2014)
Taurotragus oryx	Lombard Nature Reserve, South Africa	0.12	Buys & Dott (1991)[c]
Taurotragus oryx	South Africa	0.16	Van Houtan *et al.* (2009)
Taurotragus oryx	Kruger National Park, South Africa	−0.02	Nicholls *et al.* (1996)
Taurotragus oryx	Masai Mara Ecosystem, Kenya	−0.11 - 0.07	Ottichilo *et al.* (2000)
Taurotragus derbianus	Senegal	0.31	Koláčková *et al.* (2011)

FIGURE CAPTIONS

Figure 1 Time series of tamaraw (*Bubalus mindorensis*) abundance in Mounts

Iglit-Baco Natural Park on Mindoro island, Philippines, from years 2000 to 2021

(with the exception of 2020 because of the COVID pandemic). Raw counts

(black solid line) and counts corrected for sampling variance (gray solid line) are

presented with the associated 95% credible intervals. Over the 21 years of

monitoring, tamaraw abundance increased at a rate of 5% per year on average.

Figure 2 Spatial variation in the local growth rate of tamaraw (*Bubalus*

mindorensis) abundance in Mounts Iglit-Baco Natural Park on Mindoro island,

Philippines, from years 2000 to 2019.

Figure 1

Figure 2

740 **SUPPLEMENTARY MATERIAL**

741 **Table S1** Estimates and 95% credible intervals of tamaraw local growth rate as

742 observed on the 18 vantage points used to count the tamaraw population at Mounts

Iglit-Baco National Park (Mindoro Island, Philippines) between 2000 and 2021.

Vantage point	r	2.5%	97.5%
Loibfo	0.12	−0.02	0.24
Magawang	0.11	−0.02	0.24
Bayokbok	0.14	0.00	0.26
Bato Fidel	0.14	0.00	0.26
Inubon	0.11	−0.02	0.24
Mibluan	0.08	−0.06	0.22
Nagbobong	0.02	−0.13	0.17
Fangandatan	0.00	−0.19	0.16
Anyayos	0.07	−0.07	0.20
Lanas I	0.07	−0.08	0.21
Iyan	−0.04	−0.25	0.14
Tarzan	−0.18	−0.42	0.02
Talafu East	0.05	−0.10	0.19
Talafu West	0.05	−0.10	0.19
Malitwang	0.07	−0.09	0.21
Lanas II	0.01	−0.14	0.15
Saligue East[1]	−0.08	−0.40	0.15
Malibayong	−0.01	−0.20	0.15
Tangle	−0.48	−2.94	0.16

[1]Counts at this vantage point stopped in 2019, replaced by Tangle