The Multilocus Multispecies Coalescent: A Flexible New Model of Gene Family Evolution

Qiuyi Li[1,†], Celine Scornavacca[2,†], Nicolas Galtier[2], and Yao-Ban Chan[1,*]

[1] *School of Mathematics and Statistics / Melbourne Integrative Genomics, The University of Melbourne, Melbourne, 3010, Australia*
[2] *Institut des Sciences de l'Evolution, Université Montpellier, CNRS, IRD, EPHE, Montpellier, 34095, France*
[*]yaoban@unimelb.edu.au

[†] *These authors contributed equally to this work.*

Abstract

Incomplete lineage sorting (ILS), the interaction between coalescence and speciation, is a process which generates incongruences between gene trees and species trees, as do gene duplication (D), transfer (T) and loss (L). These processes are usually modelled independently, but in reality, ILS can affect gene copy number polymorphism, i.e., interfere with DTL. This has been previously recognised, but not treated in a satisfactory way, mainly because DTL events are naturally modelled forward-in-time, while ILS requires backward-in-time coalescent modelling. Here we consider the joint action of ILS and DTL on the gene tree/species tree problem in all its complexity. In particular, we show that the interaction between ILS and duplications/transfers only can result in patterns usually interpreted as resulting from gene loss, and that the realised rate of D, T and L becomes non-homogeneous in time when ILS is taken into account. We introduce algorithmic solutions to these problems. Our new model, the *multilocus multispecies coalescent* (MLMSC), which also accounts for any level of linkage between loci, generalises the multispecies coalescent model and offers a versatile, powerful framework for proper simulation and inference of gene family evolution.

Key words: Gene duplication, gene loss, horizontal gene transfer, incomplete lineage sorting, multispecies coalescent, hemiplasy, recombination

Species trees and gene trees are two important kinds of phylogenetic trees which are key to the study of gene and genome evolution. A species tree depicts the evolutionary history of a set of organisms, whereas a gene tree depicts the evolutionary history of a gene family within a set of organisms.

When considering speciations as the only possible events shaping species histories (e.g., no hybridisation or reassortment possible), internal nodes of species trees represent only speciation events, and branch lengths represent divergence times or substitutions per site. Nodes of gene trees, in contrast, can be the result of a combination of diverse processes, such as variation in gene copy number or gene transfers between species, in addition to speciations. For this reason, gene trees often differ from species trees, which can be seen as a problem — if the goal is to infer the species tree — or a source of information — if the goal is to study molecular evolution.

These evolutionary processes can include several 'gene-range' events such as gene duplications, gene losses, and horizontal gene transfers. A gene duplication (D) is an event in which a single gene copy gives rise to two copies at distinct loci: the parent locus and a new (child) locus. In contrast, a gene loss (L) removes a gene from the genome. Horizontal gene transfer (T) occurs when a gene from one species enters the genome of another contemporary species, which can occur (frequently in bacteria, for example) through a number of biological mechanisms such as transformation, transduction, and conjugation. Collectively, we refer to these processes as 'DTL'. These events can occur multiple times, allowing the gene tree to possibly differ greatly from the species tree. Examples of these events are given in Figure 1.

In addition to these events, a gene tree can also be incongruent to the species tree due to a phenomenon known as incomplete lineage sorting (ILS, Maddison, 1997). When a population of individuals undergoes several speciations in a relatively short time,

(a) Gene duplication. The duplication is represented by a white square, and the new gene lineage created by the duplication is coloured red.

(b) Horizontal gene transfer. The transfer origin is represented by a white triangle, and the new gene lineage created by the transfer is coloured blue.

(c) Gene loss. The loss event is represented by a cross, and the lost lineage is dashed.

Fig. 1: Tree representations of a gene duplication, a horizontal gene transfer, and a gene loss, respectively. The gene lineages (finer lines) evolve within a species tree (outer 'tubes').

polymorphism (different alleles) maintained throughout this time may eventually fix in different descendant lineages. This can produce discrepancies between the gene tree and species tree. ILS is more likely to occur in branches of the species tree (i.e., ancestral species) that represent small time spans and/or large population sizes (Pamilo and Nei, 1988). An example of ILS is given in Figure 2. Hemiplasy (Avise and Robinson, 2008) is a term used to refer to the species tree/gene tree conflicts that result from incomplete lineage sorting.

(a) Population-genetic view. Dots represent individuals and rows of dots correspond to generations.

(b) Gene evolution representation

Fig. 2: Example of incomplete lineage sorting (ILS). The original population contains a single white allele for the gene of interest. First, a mutation leads to a new black allele at the locus, then the first speciation takes place, rapidly followed by a second one. As the white and black alleles still coexist when the second speciation takes place, both alleles may be fixed in separate descendant species, resulting in a gene history which differs from the species history.

Coalescent theory (Kingman, 1982) provides a genealogical interpretation of ILS

that helps to connect this phenomenon to gene tree-species tree discordance. A key point is that the age of the common ancestor of two gene copies sampled in two different species is older (in the absence of DTL) than the time of speciation between the two species. This is due to the existence of polymorphism in the ancestral species. When speciations occur far apart in time from each other, ancestral polymorphism only creates differences in branch lengths, not in topology, between gene trees and species trees. If, however, two or more speciations occur in a time interval of the order of coalescence times, then coalescence and speciation may interact. This can cause not only branch lengths, but also topology, to differ between gene trees and species trees, as in Figure 2.

The multispecies coalescent model (MSC, Rannala and Yang, 2003) predicts the effect of ILS on gene tree branch lengths and topology as a function of the effective population size and timing of speciations. ILS and the MSC have received much attention after the discovery that classical phylogenetic methods are inconsistent in a subset of the parameter space (Kubatko and Degnan, 2007; Roch and Steel, 2015), and in the context of the study of convergent evolution (Guerrero and Hahn, 2018).

So far, ILS has mainly been considered separately from other sources of conflict between gene and species trees. However, ILS can interact in complex and often unintuitive ways with the processes of gene duplication, loss, and transfer, as first noted by Rasmussen and Kellis (2012). This is because DTL events spend some time in a polymorphic stage in a population, when individuals differ from each other in terms of gene copy number, before they become fixed. If speciations occur during this transient period of polymorphism, then the issue of lineage sorting becomes even more complex than in the one-locus case. For example, an allele which is not yet fixed can be lost, as shown in Figure 3. It is also possible that a newly created locus does not fix in all descendant species, as shown in Figure 4. Thus modelling ILS together with DTL requires greater flexibility than simply modelling each process individually. We refer to the discrepancies in gene copy number that result from the interaction between ILS and DTL as 'copy number

₇₈ hemiplasy' (or CNH for short). We note here that Rasmussen and Kellis' model considers

₇₉ DTL and ILS independently, and does not model hemiplasy. Therefore, it cannot handle

₈₀ the scenarios shown in Figures 3 and 4.

(a) Population-genetic view   (b) Gene evolution representation

Fig. 3: Example of ILS interacting with loss. With only one loss event, two descendant species may end up with an empty locus due to the presence of ILS.

original locus   new locus

Fig. 4: Example of copy number hemiplasy. A duplication arises in an ancestral species, and then two successive speciation events occur. This results in some species retaining two copies of the gene, while others have only one, without any gene loss event. Here, the original gene descends to all extant species (A, B, and C) in the original locus, but the duplicated gene only fixes in species A and B. Species B ends up with the same number of gene copies as A, while being more closely related to C.

₈₁ Discrepancies between a gene tree and a species tree due to DTL and/or ILS are

₈₂ frequently analysed via mappings from the gene tree into the species tree, called

₈₃ *reconciliations* (Goodman et al., 1979). More formally, a reconciliation between a gene tree

₈₄ G and species tree S is a mapping of the nodes of G into the nodes (or a sequence of

[85] nodes) of S (Doyon et al., 2011), respecting some constraints that depend on the

[86] evolutionary model chosen. This gives rise to the problem of reconciliation inference, where

[87] we seek to reconstruct the 'true' reconciliation from the gene and species trees.

[88]   There are two main paradigms for reconciliation inference: the parsimonious and

[89] the probabilistic (Doyon et al., 2011). In the parsimonious approach, we assign a cost to

[90] each evolutionary event, and search for a most parsimonious reconciliation, i.e., one which

[91] induces the lowest total cost. In the probabilistic approach, a stochastic model of evolution

[92] is assumed, and either the reconciliation with the maximum likelihood under this model is

[93] found, or a Bayesian approach is used to sample the posterior reconciliation space. In

[94] general, the probabilistic approach is more accurate but less time efficient, since it

[95] typically requires the estimation of many parameters, e.g., effective population sizes and

[96] branch lengths.

[97]   In our discussion, it is important to distinguish between three concepts:

[98] • The *model* of evolution — a specification of what can (and cannot) happen in the
[99]   evolution of a gene family;

[100] • A statistical specification of the model, which we call here the *implementation* of the
[101]   model, which is necessary to support a probabilistic method; and

[102] • A *method* to choose an optimal reconciliation under some criterion (parsimonious or
[103]   probabilistic).

[104] Every reconciliation method assumes a particular model of gene family evolution, limiting

[105] the potential sources of gene tree-species tree discrepancy. However, the underlying model

[106] is not always explicitly specified, which is necessary for a proper understanding and

[107] comparison of reconciliation methods.

[108]   The first reconciliation models considered only gene duplications and losses as the

[109] sources of discrepancy (Goodman et al., 1979; Zhang, 1997; Arvestad et al., 2004; Durand

[110] et al., 2006; Rasmussen and Kellis, 2011). More recently, horizontal gene transfer has been

included in these models (Arvestad et al., 2009; Doyon et al., 2010; David and Alm, 2011;

Tofigh et al., 2011; Sjöstrand et al., 2013), increasing both their complexity and realism.

Statistically, these models are usually implemented by birth-death processes running inside

the species tree. On the other hand, the multispecies coalescent (Rannala and Yang, 2003)

implements a model where discrepancies between gene and species trees are only due to

ILS.

Only a few models (and corresponding methods to infer reconciliations) exist which

attempt to unify these processes. The DLCoalRecon and DLCpar methods (Rasmussen

and Kellis, 2012; Wu et al., 2014; Du et al., 2019; Mawhorter et al., 2019) consider ILS

together with duplications and losses, overlaid on a model called DLCoal, which we refer

to in this paper as the *locus tree model*. SimPhy (Mallo et al., 2015) is a simulator based

on the locus tree model that additionally considers transfers. Schrempf and Szöllősi (2018)

also consider these possible events, but use a different model that we call here the

haplotype tree model. Lastly, the IDTL (Chan et al., 2017) and Notung (Stolzer et al.,

2012) methods incorporate both ILS and transfers with duplications and losses, but again

using different underlying models. Each of these models are slightly different from each

other, and some of the papers (Chan et al., 2017; Du et al., 2019) compare the various

models. However, they all have their limitations, which are discussed in Section "Existing

Models of Evolution". In particular, none of them can appropriately model copy number

hemiplasy, although CNH is at the heart of the ILS/DTL interaction.

When modelling ILS together with DTL, one must also consider the issue of genetic

linkage between gene copies. A new gene copy arising from a duplication often appears

close to the parent gene on the chromosome, in which case the two loci are expected to

follow correlated coalescent processes. The strength of the correlation is controlled by the

amount of recombination between the two loci; in the absence of recombination, the two

genealogies will be the same. The joint coalescent process of partly linked loci in a single

population is well characterised (Hudson, 1983), but the connection with speciation and

¹³⁸ ILS has only been rarely considered so far, the model of Slatkin and Pollack (2006) being
¹³⁹ one notable exception. (See Section "Generating a Haplotype Forest" for details about
¹⁴⁰ their model.) A realistic model of gene family evolution should account for the possible
¹⁴¹ existence of linkage between duplicated gene copies.

¹⁴² In this paper, we propose a new gene family evolution model, called the *multilocus*
¹⁴³ *multispecies coalescent*, or MLMSC for short. (This should not be confused with the
¹⁴⁴ multilocus coalescent, or MLC for short, introduced by Rasmussen and Kellis (2012).) This
¹⁴⁵ model generalises the multispecies coalescent to gene families, and is designed to capture
¹⁴⁶ all possible scenarios that can arise through ILS, DTL, and any interaction between these
¹⁴⁷ processes. The MLMSC combines forward- and backward-in-time modelling in order to
¹⁴⁸ properly account for copy number hemiplasy and linkage between loci. Importantly, we
¹⁴⁹ show that the realised rates of D, T and L become non-homogeneous as these processes
¹⁵⁰ interact with ILS, and we introduce a solution to this problem. The MLMSC model is
¹⁵¹ more flexible and predicts a more diverse range of biological patterns than existing models
¹⁵² of gene family evolution.

Existing Models of Evolution

¹⁵⁴ We consider the problem of modelling gene family evolution in a phylogenetic
¹⁵⁵ context. Given a set of species, we assume that we have sampled exactly one haploid
¹⁵⁶ genome per species, and these genomes contain all the gene copies descending from one
¹⁵⁷ particular gene present at the root of the species tree, i.e., a complete gene family. We aim
¹⁵⁸ to model the genealogical relationships between these gene copies, assuming that it has
¹⁵⁹ been shaped by speciations, duplications, transfers, losses and ILS. In this section, we first
¹⁶⁰ review the existing models which have previously addressed this problem.

Locus Tree Model

The reconciliation methods DLCoal and DLCpar are both based on the locus tree model, DLCoal (Rasmussen and Kellis, 2012). In this model, when a gene duplication occurs, the child and parent gene copies evolve independently of each other, i.e., the two loci are unlinked. Biologically, this arises when there is a sufficient level of recombination between the two loci that they can be considered to evolve completely independently.

To implement this model statistically, given a species tree, we first generate a tree under the multispecies coalescent model on the species tree. At each duplication, the species tree is copied from that point onwards and attached to the original species tree (where it may be subject to further duplications), whereas the species tree is truncated at each loss. This produces the so-called *locus tree*, which depicts the bifurcating evolution of all loci containing a copy of the gene.

The gene tree is then constructed by applying a multispecies coalescent process within the locus tree, with the caveat that only one gene copy can 'travel' along the branch connecting a duplicated locus back to its parent. In other words, all the lineages in one particular locus must coalesce into one lineage before coalescing with a copy in its parent locus — the so-called *bounded coalescent process*. This does not permit copy number hemiplasy — a duplicated gene is transmitted to all descendant species at the new locus, barring gene loss. Under this model, ILS does not interfere with gene duplication and loss in generating variation in gene copy number among species.

An example of this model is given in Figure 5, in which a new locus (with descendants in B and C) is created by duplication. The duplicated gene is then fixed in all species at the new locus.

It is easy to extend this formulation to include gene transfers, and indeed SimPhy (Mallo et al., 2015) is a simulator that implements the locus model, extended to transfers (note that SimPhy does not provide a method to find an optimal reconciliation).

Fig. 5: An example of the locus tree model.

Haplotype Tree Model

The haplotype tree model was introduced by Schrempf in a talk at the SMBE 2018 conference (Schrempf and Szöllősi, 2018). In contrast to the locus tree model, the haplotype tree model assumes that there is no recombination. This implies that a duplicated gene must undergo the exact same genealogical history as its parent gene — a strong assumption.

To implement this model statistically, we first generate a tree under the multispecies coalescent model on the species tree, obtaining a so-called *haplotype tree*. The gene tree is then obtained by performing a duplication-loss birth-death process on the haplotype tree. An example is given in Figure 6. Note that, while some sort of copy number hemiplasy is allowed in this model, it is restricted: a gene which duplicates must be sorted into exactly the same descendants as its parent gene. For instance, it is not allowed for a duplicated gene to be sorted into all descendant species, if the parent gene is not.

Fig. 6: An example of the haplotype tree model.

IDTL Model

The IDTL reconciliation method (Chan et al., 2017) takes a parsimony approach to the reconciliation problem. It defines events forward in time and assumes an underlying gene family evolution model which shares some elements with both the locus and haplotype tree models. Note that, while the model is specified, no statistical implementation has been specified by Chan et al. (2017), since this is not necessary for a parsimony method.

One possible implementation of the underlying model is to generate a tree under the multispecies coalescent model on the species tree, followed by a DTL birth-death process, as is done in the haplotype tree model. However, for each new locus, a new multispecies coalescent is performed as in the locus tree model. As in the haplotype tree model, a limited form of copy number hemiplasy is allowed where a duplicate gene must be sorted into exactly the same descendant species as the parent gene. An example of this is given in Figure 7.

This model deals with recombination in an inconsistent way; this will be discussed further in Section "MLMSC vs IDTL Model".

Notung

The Notung method (Vernot et al., 2008; Stolzer et al., 2012) was one of the earliest methods to unify the processes of DTL and ILS. It is primarily intended to be used with non-binary species trees, i.e., situations where the branching order is unclear. Because of this, ILS in Notung is only allowed at a polytomy (node with more than two descending

Fig. 7: An example of the IDTL model. An allele which is sorted into species *A*, *B*, and *C* is duplicated. The duplicated gene must then be sorted into the same species, but the coalescent process may be different from the parent gene, resulting in a subtree (coloured red) with differing topology.

²²⁰ lineages) in the species tree, and each possible sorting of genes at the polytomy is

²²¹ considered to be equally likely. Thus there is no explicit modelling of alleles co-existing

²²² inside a species branch. As with the IDTL method, Notung is a parsimony method and

²²³ does not have a formal statistical implementation.

The MLMSC Model

²²⁴

²²⁵   The models of gene family evolution in the literature do not appropriately model

²²⁶ copy number hemiplasy or linked loci. Here we introduce a new gene family evolution

²²⁷ model, the *multilocus multispecies coalescent* (MLMSC), for this purpose.

Assumptions

²²⁸

²²⁹   We first clearly set out the assumptions in the model, which helps to distinguish the

²³⁰ possible scenarios that can arise. One of the primary conceptual difficulties in unifying ILS

²³¹ with a DTL model has been that ILS is traditionally implemented by a backwards-in-time

²³² coalescent model, whereas the birth-death process used to implement DTL events runs

²³³ forwards in time. The MLMSC model is based on a Wright-Fisher model for the

²³⁴ population-level genealogical process, which also runs forwards in time and is therefore

easier to merge with a birth-death process. To calculate the genealogy of alleles, the coalescent model is still used in practice, as it is far more efficient than the Wright-Fisher; however, the Wright-Fisher is used to specify the possible scenarios.

In the MLMSC model:

1. The evolution of the gene inside the species is considered forward in time from one common ancestral gene towards the contemporary genes.

2. Duplication (D), horizontal gene transfer (T), and loss (L) are the possible events which can shape gene histories. (ILS is not considered to be an event, but rather a name for the discrepancies arising from population-level processes.)

3. Each species evolves as a panmictic population. Within a population, in each generation each member is descended (and thus inherits all genes) from any member of the previous generation. Several alleles may be maintained in a population in any generation.

4. When a speciation occurs, two independent new populations are created with the same loci as the parent species. Each member of these populations is descended from a randomly selected member of the parent species. If several alleles are present in the parental population, a sequence of rapid speciations may thus result in topological discrepancies between the gene and species trees, or ILS. (Items 3 and 4 describe the ordinary multispecies coalescent.)

5. When a duplication occurs:

    (a) An individual in a population is chosen to have the duplication to occur within its genome. If this individual carries a copy of the gene, a new locus is created in the genome at the given species. For this locus, the population contains one individual that exhibits the duplicated gene, while the remainder have no gene copy at this locus.

(b) The duplication can be of two types:

    i. *unlinked duplication*, where the new locus is unlinked to the parent locus, and thus is independently transmitted across generations;

    ii. *linked duplication*, where the new locus is linked to the parent locus. In this case, the new locus is inherited together with the parent locus, except for when there is a recombination event: when a recombination event occurs, a single individual in the population has different parents in the two loci.

(c) The duplicated gene may consequently fix in any subset (including none or all) of the descendants of the species containing the duplication (CNH).

6. When a transfer occurs, the same process is followed as for a duplication, except that the transferred gene appears in a new locus of a contemporary species of the origin species, and this locus is always unlinked to any existing locus.

7. When a loss occurs, an individual in a population is chosen to have the loss to occur within its genome, if a copy of the gene is present. This gene may or may not be an allele. The loss may consequently fix in any subset (including none or all) of the descendants of the species containing the loss (CNH). If a population loses its last copy of a gene in a given locus in a species, that locus is lost in that species.

The Model

The MLMSC model, which generalises the multispecies coalescent model to gene families, is an exact coalescent-based simulation of the genealogies that arise from a Wright-Fisher model in each species branch with constant rates of duplications, transfers, and losses. The major innovations in this model are:

1. Modelling copy number hemiplasy with the *incomplete coalescent*;

2. Modelling linked loci with recombination with the *linked coalescent*;

3. Simulating surviving DTL events under the correct rates with the *coalescent-rate*
process.

We now outline each of these ideas in turn. A full specification of the model is left
to Section "Full Model Specification". Firstly, we introduce the concepts of *unilocus trees*
and *haplotype forests and trees*.

Unilocus trees model the history of duplications, transfers, and speciations. We assume
that each duplication or transfer event creates a new locus. This new locus may be
inherited by the descendants of the ancestral species, S, in which it was created, and only
by these. For each newly created locus, we define its *unilocus* tree as the subtree of the
species tree rooted at S, i.e., including S and its descendants. The leaves of a unilocus tree
thus correspond to the same newly-created locus in different species (i.e., the leaves below
S). The history of gene duplications, gene transfers, and speciations in the MLMSC is
therefore stored as a collection of disconnected unilocus trees.

Note that this collection only slightly differs from the locus tree of the DLCoal
model (see above): a collection of unilocus trees can equivalently be seen as a decorated
locus tree, where the decorations are duplications/locus changes. For ease of
understanding, in the following we consider them as disconnected trees.

Haplotype forests and trees model the genealogies of lineages within a unilocus tree. We
consider that we have sampled a single haploid genome per species, meaning that, for a
given locus, the number of gene copies carried by a given species is zero or one. We model
the genealogical relationships of gene copies across species, for a given locus, via a so-called
haplotype forest, which we define as a collection of *haplotype trees*, where a haplotype tree
depicts the genealogy of gene copies for a subset of species. A haplotype forest contains
either one haplotype tree or a set of disconnected haplotype trees, i.e., haplotype trees
whose sets of leaves are disjoint. Generally, only one of the trees in the haplotype forest
will actually carry a copy of the gene, in which case we refer to it as *the* haplotype tree for

the locus.

Haplotype forests and trees will be used to correctly model the presence of different alleles in the populations (via ILS, CNH and losses). Note that in our model, DTL events are modelled forwards in time and then haplotypes are modelled backwards in time.

Modelling Copy Number Hemiplasy

To model the interaction between D, T, and ILS, and allow for copy number hemiplasy, we introduce a new process called the *incomplete multispecies coalescent.* In this model, when generating the haplotype tree for a given locus, we do not require that all extant genes have to coalesce to their most recent common ancestor (MRCA) by the time of origination, i.e., at the root of the unilocus tree. Rather, we simply stop the coalescent process at the time of origination of the locus, thus producing a haplotype forest. One of these trees is then randomly chosen to be the haplotype tree, which is thus joined with the haplotype forest of the originating locus (for more details on how the joining is performed, see below). The root of this tree represents the first individual carrying the new copy of the gene. In this way, a duplicated/transferred gene copy does not have to be transmitted to all descendant species, and CNH can occur. An example of this process is shown in Figure 8.

Not requiring an MRCA to be reached has a further implication, which is that a locus can be entirely lost via sorting effects only. Consider the example in Figure 9: in a locus 1, originated at time t_1, a duplication occurs at time t_2, resulting in the origination of locus 2. Under the incomplete coalescent, it might be that the duplicating lineage fails to coalesce with any lineages in locus 1 by time t_1. This can be interpreted as a duplication arising in an individual who does not actually contain any gene copy at that locus — i.e., an impossibility. In this case, the duplication is discarded and no new locus is created. A similar situation can occur with transfers, or a combination of the two types of event.

Additionally, to model copy number hemiplasy, we must be able to lose an allele which is not completely fixed in a population. To achieve this, losses are simulated on gene

Fig. 8: An example of the incomplete multispecies coalescent. Two trees (the red one and dashed one) are generated in the new locus using the multispecies coalescent. The red tree is randomly chosen to be the haplotype tree for this locus, while the dashed tree is part of the haplotype forest but is not the haplotype tree. The root of the red tree represents the individual carrying the new copy.

Fig. 9: The new haplotype tree does not coalesce with the haplotype forest in the parent locus by the time of its creation, so the duplication is discarded. (Here we use the convention "dashed implies not chosen as haplotype tree".)

lineages (branches of the haplotype trees), rather than on loci (as is done in the locus tree model).

Modelling Linked Loci

Some duplications, and all transfers, appear in *unlinked* loci; that is, the genealogy of the population (and thus the evolution of the gene) in the new locus is independent from that at the parent locus. This is a reasonable assumption which occurs in many cases, for example with large-scale duplications. However, some duplications may produce a new locus which is physically close to the parent locus on the chromosome, in which case their evolution will be dependent. We call these *linked* loci.

To model linked loci, we introduce a new process called the *linked coalescent* to generate the haplotype forest and tree in a linked locus created by duplication. In this process, the haplotype forest in the parent locus is mirrored into the new unilocus tree and used as a pre-existing genealogy which can be followed by the lineages in the new locus. The incomplete coalescent is then run, during which gene lineages can coalesce with each other, as usual, or with the pre-existing genealogy. In this way, the genealogy of the new locus depends on the genealogy of the existing locus; in the absence of recombination, the two will be identical.

To model the effects of recombination, lineages in the new locus which are coalesced with the pre-existing genealogy can 'uncoalesce' from it, representing a recombination event between the two loci. In contrast, lineages in the new locus which coalesce with each other cannot uncoalesce, because they represent actual lineages. By varying the rate at which recombination events occur, we can model any level of dependence between loci. An example of this process is given in Figure 10.

A duplication may either be *ancestral* (the duplicating individual is a direct ancestor of a sampled individual in the genealogy of the parent locus), or *non-ancestral* (the lineage of the duplicating individual does not survive, or is not sampled, in the parent locus). See Figure 11 for examples. For unlinked loci, it can be assumed that only non-ancestral duplications occur, as the probability of survival in the parent locus, which is independent of the existence of the duplicated locus, is extremely small and can be safely

Fig. 10: The linked coalescent. On the left, we have the original unilocus tree and its haplo-type forest (the haplotype tree is solid and other trees in the haplotype forest are dashed); on the right, the new unilocus tree with a linked coalescent process run on it. The coalescent process starts linked with the genealogy copied from the parent locus in all species, then 'un-coalesces' due to recombination (black circles) in all species at different times, coalesces back to the mirrored genealogy (black stars) in A and the ancestor of B and C, and, finally, the lineages that descend to A and B coalesce with each other (black diamond). In the middle, we show why the coalescent process starts linked and then 'uncoalesces': suppose we have individuals with two linked loci (the left half of the circle corresponds to the first loci, the right half to the second). A recombination happens at generation $i + 1$. From the drawing, it should be clear why the lineages of the two genes are linked below the recombination and unlinked above.

ignored. However, when modelling linked loci, it is important to distinguish between these two cases. By associating lineages in the new locus with lineages in the parent locus, the linked coalescent provides a natural way to determine if a linked duplication is ancestral or non-ancestral.

Simulating Surviving Events

It is important to realise that, here, we do not want to simulate all DTL events that occur, because the vast majority of these events will simply fail to fix in the population, and thus be unobserved. Instead, we only wish to simulate events which survive to the present day and are observed in at least one sampled individual. (We use the terms 'surviving' and 'observed' interchangeably, as a lineage which survives in an unsampled individual is undetectable.) In order to do this, we must consider the probability of

(a) Ancestral duplication

(b) Non-ancestral duplication

Fig. 11: Ancestral vs non-ancestral duplications.

survival of each event.

In simpler models (e.g., the locus tree model), the survival probability is constant in time and across lineages, but unrealistic assumptions are needed for this to occur. In the MLMSC model, the survival probability is not constant. This can be seen by considering the following simple example: a duplication occurs 'just before' a speciation into two species leaves. In this case, the duplication can survive because it is observed in either of the descendant species (Figure 12a,b) and possibly (though unlikely) in both (Figure 12c). Since the probabilities of being observed in either species are close to independent, the total survival probability is roughly twice that of a duplication which occurs in a terminal branch of the species tree (Figure 12d).

In our model, we develop the *coalescent-rate process*, which simulates the exact rate of surviving events, under the Wright-Fisher model, anywhere in the species tree. The simplest version of this process — applied to unlinked duplications — runs an incomplete coalescent in a unilocus tree, and then simulates events at constant rate on the branches of the trees obtained in this way. The events are then considered to occur in the corresponding branches of the unilocus tree (see Figure 13). This allows us to simulate

Fig. 12: Although duplications occur at a constant rate, they do not survive with constant probability. (a-b) Duplication 'just before' a speciation. A duplication in either of the filled individuals will survive (we use the convention that the dashed line is not chosen as the haplotype tree). (c) Only a duplication in one of the individuals will survive, but this scenario requires a near-immediate coalescence and is unlikely. (d) Duplication in a species leaf. Only a duplication in one of the individuals can survive.

surviving duplications at the correct rate. Some modifications must be made for transfers and linked duplications, and we give further details below in Section "Full Model Specification". We have implemented a simulator based on this model, available at

https://github.com/QiuyiLi/MLMSC.

Model Comparison

In this section, we discuss the limitations of the current models in the literature, and show how the MLMSC model is subject to none of these limitations. Our discussion is summarised by, but not limited to, Table 1.

MLMSC vs Locus Tree Model

The locus tree model simulates losses on the locus tree, instead of the haplotype tree. Assigning a loss to a locus, instead of a branch of the haplotype tree, means that the

Fig. 13: An example of the coalescent-rate process. Firstly, 'temporary' trees are sampled from the multispecies coalescent, and then events are sampled at constant rate on the branches of these trees. Finally, the 'temporary' trees are removed and the sampled events are considered to occur in the corresponding branches of the unilocus tree.

	Transfers	CNH	Recombination	Statistical implementation	Reconciliation method
MLMSC	✓	✓	All	✓	✗
Locus tree model	✗	✗	Infinite	✓	✓
Haplotype tree model	✗	✗	None	✓	✗
IDTL	✓	✗	Limited	✗	✓
Notung	✓	✗	Infinite	✗	✓

Table 1: Comparison of the models in the literature.

loss of an allele due to lineage sorting cannot be modelled. Consider Figure 14a: here, two alleles were present in the ancestral population of B, C, and D, one of which was subsequently lost, i.e., replaced by the null allele. The null allele is then sorted into species B and C. The locus tree model can only produce this gene tree by invoking at least two losses, as in Figure 14b. In order to properly capture this scenario, losses must be placed on the haplotype tree rather than the locus (or unilocus) tree, which is exactly what is done in the MLMSC model.

In a similar vein, the locus tree model also assigns duplications to the locus tree. These duplications must then rejoin the haplotype tree via the multispecies coalescent; in other words, they are assumed to be non-ancestral. As discussed above, this is a reasonable consequence of the model assumption that all loci evolve independently (i.e., are unlinked),

(a) An allele is lost, with the null allele being sorted into species B and C.

(b) The locus tree model must invoke at least two losses to reproduce this gene tree.

Fig. 14: The locus tree model cannot model lost alleles.

since the probability of an ancestral unlinked duplication is vanishingly small. However, in the more realistic case that some duplicated loci may be linked to their parent loci, the survival of a duplicated gene is correlated to the survival of its parent gene, and the probability of an ancestral linked duplication, knowing that the new duplicate exists, is non-negligible. Thus it is important to model both ancestral and non-ancestral duplications, as the MLMSC model does.

Finally, in the locus tree model, a duplicated gene is either lost, or fixed in all possible descendant species. This means there can be no copy number hemiplasy. For example, the locus tree model cannot model the scenario in Figure 15a: an additional loss is needed, as in Figure 15b. By use of the incomplete coalescent, the MLMSC model can model both scenarios.

(a) The duplication fixes only in species B without a loss.

(b) The locus tree model must infer a loss in species C.

Fig. 15: The locus tree model cannot model copy number hemiplasy.

MLMSC vs Haplotype Tree Model

In the haplotype tree model, a duplicated gene is assumed to have exactly the same genealogy as its parent gene; the model cannot model duplicated genes with different coalescent histories. For this to occur, we must assume that there is no recombination between loci, which is too restrictive. It is more realistic and flexible to allow the loci to be linked, where evolution in the loci are dependent but not necessarily identical, or unlinked, where evolution is completely independent. This is done in the MLMSC model.

Because the haplotype tree model applies the coalescent first, duplications and losses are assigned to gene lineages, rather than loci as in the locus tree model. It is reasonable to apply duplications to gene lineages when there is no recombination (and thus only ancestral duplications can be observed). But, as discussed above, a more realistic model is to allow recombination, and model both ancestral and non-ancestral duplications, as done by the MLMSC model. On the other hand, the disadvantages of the locus tree model do not apply to the haplotype tree model; for example, it can model both scenarios in Figure 14.

The haplotype tree model also does not fully allow for copy number hemiplasy; instead, keeping with the assumption of fully dependent loci, it enforces a restricted version wherein a duplicate must undergo the same coalescent process, and therefore be sorted into the same species as the parent gene. For example, the haplotype tree model cannot model the scenario in Figure 16a, where the duplicated gene is sorted into different species than the parent gene. It also cannot model the scenario in Figure 16b, where the duplicated gene undergoes a different genealogy from the parent gene. The MLMSC model can accommodate both of these scenarios, assuming either unlinked loci or linked loci with recombination.

(a) The haplotype tree model cannot have a duplication which is sorted into different species from the parent lineage.

(b) The haplotype tree model cannot have a duplication which has a different genealogy than the parent lineage.

Fig. 16: Limits of the haplotype tree model. Duplicated lineages are in red, while the parent lineages are in black.

MLMSC vs IDTL Model

The IDTL model offers a kind of 'halfway house' between the locus tree and haplotype tree models, where duplications, transfers, and losses are applied to gene lineages rather than loci, and duplicates can be sorted in different ways from their parent genes. However, some of the assumptions made in the model are computationally convenient but biologically questionable — for example, a duplicated gene must be sorted into the same species as its parent gene (i.e., no recombination allowed), but it may be sorted in a different way (i.e., recombination is allowed). For example, Figure 15a is also not allowed by the IDTL model. The MLMSC model is based explicitly on a model of the coalescent with recombination, and therefore incorporates recombination events in each species explicitly, allowing a greater range of scenarios.

MLMSC vs Notung

As discussed above, Notung only allows ILS at a polytomy, and each possible sorting of genes at the polytomy is equivalent and equally likely; there is no 'correct' sorting which agrees with a specified species tree. In particular, this means that ILS is not penalised, and so any gene tree-species tree discrepancy which can be attributed to ILS is attributed to it, with the remaining discrepancies then explained by DTL. In contrast,

MLMSC allows ILS everywhere in the tree, with probabilities based on the branch lengths of the species tree, and balances that with the DTL processes. It also specifies an underlying 'true' binary species tree which is always the most likely outcome for the gene sorting. Internal branch lengths in the MLMSC can be made arbitrarily short, which effectively covers what Notung represents by polytomies.

For computational convenience, the Notung model assumes that whenever a transfer occurs, the parent lineage must survive to the present day. This means that transfers must be ancestral. (In reconciliation terminology, this means that there are no 'transfer-loss' events.) As discussed above, this is not a realistic assumption, as only the transferred lineage needs to be observed; in fact, assuming selective neutrality, the chance of the parent lineage also surviving in the parent locus is $O(\frac{1}{2N})$, where $2N$ is the effective population size. The MLMSC model has no such restriction and can model transfer events accurately.

Discussion

The MLMSC model generalises the multispecies coalescent to include duplications, transfers, and losses. By using the incomplete coalescent, haplotype forests, and disjoint unilocus trees, the MLMSC accounts for scenarios in which ancestral gene copy number polymorphisms are incompletely sorted among the descendant species, in contrast with existing models. By allowing for both linked and unlinked loci, the MLMSC model can also model more complex evolutionary scenarios than the existing models, while taking recombination into account in a natural way. Finally, the MLMSC recognises the fact, so far ignored, that the realised rate of duplication, transfer and loss becomes non-homogeneous if ILS is at work.

MLMSC is a particularly versatile model, which captures many evolutionary processes: speciation, gene duplication, gene loss, gene transfer, and genetic linkage. In this respect, MLMSC is more powerful than any of the existing models of gene family evolution (see Table 1). Note that what one can and cannot model has a strong impact on the

accuracy of the estimation of species and gene trees, as pointed out for instance by Boussau and Scornavacca (2020). We suggest that, in the presence of CNH, existing reconciliation methods might lead to biased estimates of gene trees, species trees, and/or evolutionary parameters, since every instance of a non-fixed duplication would incorrectly cost one loss if CNH is not explicitly accounted for.

Not all these processes, however, are necessarily relevant to every biological system. The prevalence of ILS, and consequently of CNH, depends on the ratio of effective population size to branch lengths (Pamilo and Nei, 1988), both of which vary by orders of magnitude among taxa and datasets. Scornavacca and Galtier (2017), for instance, suggested that ILS is only a minor determinant of conflicts between gene trees in phylogenetic analyses of the Mammalia clade, in which effective population sizes are relatively small (Romiguier et al., 2014) and branches typically represent millions of generations. DTL alone might be a sufficient model of gene family evolution at this scale. However, even in mammals, studies of closely related species have demonstrated the potential importance of ILS when short time frames are considered (Hobolth et al., 2011; Kutschera et al., 2014), suggesting that, as larger and larger data sets are generated and analysed, the pertinence of MLMSC should increase. Similarly, the modelling of genetic linkage between duplicates might be superfluous when relatively small and ancient gene families are considered, but should be crucial for the analyses of large and dynamic gene families, such as venom toxins (Fry et al., 2009) or olfactive receptors (Niimura et al., 2014), in which paralogous gene copies are often organized in clusters across the genome (Olender et al., 2020).

To assess the impact of CNH on gene evolution, we simulated gene trees on the fungal species tree used by Rasmussen and Kellis (2012), using three different duplication and loss rates (10^{-10}, 5×10^{-10}, and 10×10^{-10}, duplication and loss rates are assumed to be equal), three effective population sizes (10^7, 5×10^7, and 10×10^7), and 0.9 years per generation. For comparison, Rasmussen and Kellis (2012) used a duplication rate of

7.32×10^{-10}, a loss rate of 8.59×10^{-10}, an effective population size of 10^7, and 0.9 years

per generation. For each set of parameters, we ran 500 simulations and calculated the

proportion of unilocus trees for which the haplotype forest contained more than one

haplotype tree (i.e., CNH occurred). From Figure 17, we see that, for this data set, CNH is

far from negligible, and as expected, its impact is higher when the effective population size

is large — ranging from a bit less than 15% for $N_e = 10^7$, to more than 50% for $N_e = 10^8$.

On the other hand, the proportion of CNH stays constant when varying the duplication

and loss rate. This indicates that CNH is an important phenomenon which must be taken

into account when modelling biological systems.

Fig. 17: The proportion of CNH in our simulated data, varying the duplication and loss rate
and the effective population size.

An implicit assumption in the MLMSC is that we can determine the linked or

unlinked status of each locus solely with respect to its parent locus. In this respect, it is not

fully cognisant of the linear structure of the chromosome. For example, if a locus (say locus

1) has two linked duplications in rapid succession (say loci 2 and 3), then one of the three

loci must lie between the other two. If the order is (say) 1–2–3, then a recombination event

between loci 1 and 2 must imply a recombination event between loci 1 and 3; this kind of

dependency is not modelled by the MLMSC, which effectively represents the relations

between linked loci as a tree rather than a linear structure. Moreover, in a scenario such as this, it may not be reasonable to assume that the recombination rate between loci 1 and 2 is equal to the recombination rate between 1 and 3. A full model which incorporates the linear structure of the chromosome would have to explicitly model the position of the gene copies, which we have elected not to do here for reasons of computational convenience.

We note that our goal here is not to correctly model the recombination to its full extent, but to handle the case where the original locus and the duplicated one are next to each other and may thus have linked histories. In addition to the above, recombination is only allowed between (not within) duplicates. Traditional population genetic models, such as the coalescent with recombination and its approximations such as the sequentially Markov coalescent, permit more flexibility on this matter: recombination is allowed anywhere within a linear genome, and no pre-knowledge about the genealogy in neighbour loci is required. These models also have their drawbacks: all genomes are assumed to be aligned, with insertions and creations of new loci not allowed, and they assume that recombination occurs at a constant rate throughout the genome, ignoring the existence of hotspots or variation in rates due to gene function. One could say that our model does include this somewhat by considering a locus as a non-recombinant segment and only allowing recombination outside it.

Another limitation of our model is that it only accounts for a portion of the events shaping species and gene histories. For example, reticulated events such as hybridisation or reassortment, whole genome duplication events, and gene conversions are not modelled. Neither is "transfer from the dead" (Szöllősi et al., 2013); although this would be a relatively easy inclusion, we have elected not to do this in order to maintain the elegance of our model. Indeed, it is unclear whether the conclusions from Szöllősi et al. (2013) apply when population-level processes are explicitly modelled, as we do here; in this case, an alternative interpretation of their result that "almost all transfers involve speciation" could be that "almost all transfer involve divergence", i.e., are non-ancestral.

⁵⁶² The development of the MLMSC model opens up several avenues for further

⁵⁶³ research. Firstly, because the model produces a more realistic and powerful framework for

⁵⁶⁴ simulating gene family evolution, it can be used to compare the accuracy of current

⁵⁶⁵ reconciliation algorithms in a 'neutral' setting. Secondly, the simulations themselves can be

⁵⁶⁶ explored under biologically realistic parameter values to ascertain the relative importance

⁵⁶⁷ of population-level processes to macro-evolutionary ones. Finally, we hope that the

⁵⁶⁸ MLMSC model and its statistical implementation will be the starting block for the

⁵⁶⁹ development of new reconciliation methods.

Full Model Specification

⁵⁷¹ In this section, we detail a full statistical implementation of the MLMSC model.

⁵⁷² The model starts with the species tree as the unilocus tree for the original locus. Then we

⁵⁷³ alternately generate a haplotype tree and forest for that locus, or new loci and unilocus

⁵⁷⁴ trees, until all simulated duplication and transfer events are accounted for. The haplotype

⁵⁷⁵ trees are then concatenated together to form the full gene tree.

⁵⁷⁶ To ease the reading of this very dense section, we provide a full example of these

⁵⁷⁷ operations in Figures 18, 19, and 20. The formal statistical details are provided after the

⁵⁷⁸ example. The pseudocode, to which we refer in the main text, is given in Algorithms 1 and

⁵⁷⁹ 2 at the end of the section. Additional details/comments/justifications are given within

⁵⁸⁰ boxes, which can be glossed over depending on the reader's objectives.

Fig. 18: Modelling gene evolution in the original unilocus tree. (a) The species tree (top) and original unilocus tree ('tubes', on the bottom), within which a haplotype tree (finer lines) is generated according to the standard multispecies coalescent. (b) Simulating unlinked duplications, linked duplications and transfers in the unilocus tree with coalescent-rate processes, and losses on the haplotype tree. Here, only one unlinked duplication and one loss are simulated, with no other events. (c) The unilocus tree is decorated with the unlinked duplication, and the haplotype tree is truncated at the loss.

Fig. 19: Modelling gene evolution in the first child locus. Here, we follow the duplicated gene in Figure 18. (a) We generate the new unilocus tree ('tubes', on the right) by copying the original unilocus tree ('tubes', on the left) from the time of the duplication. The new haplotype forest (finer lines, on the right) is generated according to the incomplete multispecies coalescent; the red tree is chosen to be the new haplotype tree. (b) Simulating events in the new unilocus tree, as in Figure 18b. Here, only one transfer and one linked duplication are sampled, with no other events. (c) The new haplotype tree is joined to the haplotype tree in the original locus, and the new unilocus tree is decorated with the simulated events.

Fig. 20: Modelling the evolution of the loci derived from the first child locus. Here, we follow the duplicated gene (a) and transferred gene (b) in Figure 19. (a) We generate the new unilocus tree ('tubes', on the right) by copying the parent unilocus tree ('tubes', on the centre) from the time of the duplication. The new haplotype tree (finer lines, on the right) is generated according to the linked multispecies coalescent. This tree is linked, in the parent locus, to a tree in the haplotype forest which is not the haplotype tree, and is thus discarded. (b) We apply a similar procedure for the transferred gene, which joins the parent haplotype tree and is thus kept. The new event sampling procedure (far right) does not simulate any new event. This concludes our example. (c) The final gene tree within the species tree. (d) The final gene tree.

Generating a Haplotype Tree for the Original Locus

We start with a species tree S (line 1, Algorithm 1). In the original locus (which we denote as locus 0), the unilocus tree S_0 is the original species tree S. For this locus only, the haplotype tree G_0 is generated according to the standard multispecies coalescent, starting from a single copy of the gene in each leaf of the tree (line 2, Algorithm 1). We also set the haplotype forest $\mathcal{G}_0$ to be the set $\{G_0\}$ (line 3, Algorithm 1).

> **Note.** For simplicity, we assume (here and in all further coalescents) that the effective population size $2N$ is constant over time and across species. The process can easily be generalised to more variable cases, with the caveat that the population size is a property of the species and must therefore remain the same across loci.

For loci created by duplication or transfer, a more complex process is required (line 4, Algorithm 1). We first describe how to generate new loci and unilocus trees, then return to generating haplotype trees within those unilocus trees.

Simulating Events

Suppose that we are simulating events from a locus l, with unilocus tree S_l, haplotype tree G_l, and haplotype forest $\mathcal{G}_l$. We are given a set of parameters $\boldsymbol{r} = (r_d, r_t, r_l, r_r, p_u)$, where r_d, r_t, and r_l are the rates of duplications, transfers, and losses respectively, r_r is the rate of recombination for linked duplications (see Section "Generating a Haplotype Forest"), and p_u is the (fixed) probability that a duplication will be unlinked.

> **Details.** Unilocus and haplotype trees have branch lengths given in coalescent units, and by convention the present day is time $t = 0$, and t increases as we go backwards in time. The rates r_d, r_t, r_l, and r_r are given in units of events per individual per locus per coalescent unit.

Losses are sampled from the haplotype tree at constant rate; duplications and transfers are sampled from the unilocus tree, and their rates need to be scaled by the probability of survival of the event.

Details. Recall that, as discussed in Section "Simulating Surviving Events", we only wish to model surviving DTL events, not all of them. A duplication may either be ancestral (the duplicating individual is a direct ancestor of a sampled lineage in the parent locus), or non-ancestral. Either of these kinds of duplication can survive, and so we can consider a duplicating individual to be any one of the population, and thus sample duplications from the unilocus tree. Likewise, (surviving) transfers can also be sampled from the unilocus tree.

On the other hand, a loss in a lineage which does not survive to the present day will not be observed; in other words, losses are always ancestral. Hence losses should be sampled from the haplotype tree at constant rate r_l (line 4, Algorithm 2). The constant rate is justified because a population which contains multiple alleles will lose each of them at the same rate as a completely fixed gene, and thus the overall loss rate will be proportional to the number of existing lineages.

In order to sample surviving duplications and transfers, we must consider both the rate at which the events occur, and their probability of being observed. We assume that the events (all events, not surviving events) occur at a constant rate per individual per locus per coalescent unit, as given by the parameters. Thus the overall rate of each event across a population is $2N$ multiplied by the per-individual rate. We then need to scale this rate by the probability of survival of the event. This probability differs for unlinked duplications, transfers, and linked duplications. We consider each in turn.

Simulating unlinked duplications (line 1, Algorithm 2)

Note. The locus tree model only considers unlinked duplications. Because no CNH is allowed, the survival probability of a duplication is constant, and since unlinked duplications occur at a constant rate, surviving unlinked duplications can also be simulated at a constant rate. In the MLMSC model, we use the incomplete coalescent to model CNH, so this property no longer holds.

In the MLMSC model, unlinked duplications occur at a constant rate of $2Np_ur_d$. To calculate the survival probability of an unlinked duplication at a locus l, consider a duplication which occurs at a certain time and species in S_l, and consider the sampled individuals in the extant descendants of this species. Each of these individuals will have a single direct ancestor at the time of the duplication, which may or may not be distinct from each other. The duplication will be observed if and only if one of these ancestors is the duplicating individual in a population of size $2N$. The probability that such a duplication is observed is therefore $\frac{1}{2N}$ multiplied by the expected number of lineages at the time of the duplication, under the multispecies coalescent.

613   It is difficult to obtain a closed formula for this number, but there is a simple way

614 to simulate at the correct rate. We call this a *coalescent-rate process*, which we now detail.

> **Details.** To simulate unlinked duplications, we run a (incomplete) multispecies coalescent within S_l, and then sample events with a constant-rate Poisson process at rate $p_u r_d$ from the branches of the resulting coalescent trees. These events will be considered as duplications at the corresponding branch and time in S_l. An example of this process is given in Figure 13.
>
> At any time, the coalescent trees will, on average, trivially have the expected number of lineages under the coalescent, and so the resulting rate is correct. We must also specify a rate for the coalescent-rate process, which is the rate at which events are sampled from the branches of the coalescent trees (in this case $p_u r_d$).

616   *Simulating transfers (line 2, Algorithm 2)* —   Transfers (which are always

617 unlinked) can be simulated using a similar coalescent-rate process to that of duplications,

618 but with some modifications.

> **Details.** It is important to note that the coalescent-rate process reflects the probability of a gene lineage surviving if it *appears* (by duplication or transfer) in a particular species. A duplication always appears in the same species that it originates from, but for a transfer, this is not the case. Thus the coalescent-rate process must be used to select the target species of the transfer, not its origin species. Additionally, while a transfer from locus l must originate from within S_l, it need not appear in S_l, but could appear in any contemporary species different from the origin species.
>
> In order to simulate surviving transfers from locus l, we thus perform a coalescent-rate process with rate r_t over the *entire* species tree S. Each resulting event marks the target species of a transfer, and we subsequently choose an origin species uniformly at random from *all* other contemporaneous species. If this origin species does not lie in S_l, the entire transfer is discarded. An example of this process is given in Figure 21.
>
> In practice, it is less efficient to simulate many transfers and discard some, so we take advantage of the fact that transfers originate from each species at an equal rate, since we assume that all population sizes are equal. Hence, we scale the overall transfer rate to each species by the proportion of possible origin species which are also contained in S_l, and then select an origin species uniformly at random from the possible origin species in S_l. It is easy to see that this is equivalent to the process described above, but without having to discard some simulated transfers.

620   *Simulating linked duplications (line 3, Algorithm 2)* —   In a linked locus created

621 by duplication, the probability of the duplication surviving is dependent on the genealogy

622 of the parent locus. However, the principle is the same: a duplication will survive if and

(a) Transfer targets (represented by black triangles) are generated using a coalescent-rate process over the entire species tree: an incomplete coalescent process is run on the species tree, and transfer targets are sampled on the coalescent trees generated in this way (left). The coalescent trees are then discarded and the transfer targets kept (right).

(b) For each transfer target, we simulate a transfer origin (represented by a white triangle, left). Transfers originating from outside the unilocus tree (depicted as shaded), e.g., the lower transfer in our example, are discarded (right).

Fig. 21: Simulating transfers from a given unilocus tree (with leaves B and C, shaded) by first simulating targets (a) and then origins (b). The transfer must originate from inside the unilocus tree, since we are simulating transfers *from* it. In (b), the new unilocus tree derived from the only retained transfer (with leaf A) is shown in a lighter shading on the right.

623   only if it is the direct ancestor of a sampled lineage in the new locus. The linked coalescent

624   (described in Sections "Modelling Linked Loci" and "Generating a Haplotype Forest")

625   directly models the genealogy in such a locus, and thus it can be substituted in place of

626   the ordinary coalescent when generating linked duplications. For brevity, we refer to this as

627   a *coalescent-rate process in the presence of* $\mathcal{G}_l$.

Details. To simulate surviving linked duplications originating from locus l, we run a linked coalescent in the unilocus tree S_l, allowing the lineages to coalesce and uncoalesce (through recombination) with the existing haplotype forest $\mathcal{G}_l$. We then sample linked duplications as events at constant rate $(1 - p_u)r_d$ on the branches of the resulting coalescent forest. These events are then interpreted as duplications at the corresponding branch and time inside S_l.

Note. Before continuing, we comment briefly on duplication and transfer rates. For a duplication/transfer to 'survive', there are three things which need to happen: 1. the event occurs; 2. the duplicated/transferred individual has a copy of the gene at the locus; and 3. we observe a descendant of the duplicated/transferred individual.
We have a fixed rate for point 1 (i.e., the rates r_d and r_t); the coalescent-rate process handles point 3 by incorporating the probability to observe the duplicated/transferred individual; and point 2 is accounted for by rejection sampling, i.e., potentially discarding loci when joining the haplotype tree of the new locus with that of the original locus (we will discuss this later in the text, see Figures 26b, 27b and 27c). This implies that the 'actual' rates (i.e., the rates of surviving events which duplicate or transfer a copy of the gene) will be lower than the rates r_d, r_t. The actual relation between these pairs of rates is highly non-trivial.

630   *Generating new loci and unilocus trees (lines 11–12, Algorithm 2)*

631   When a new locus is created, we create a unilocus tree for it, which shows the

632   evolution of all species which could possibly contain the locus (i.e., all descendants of the

633   species where the locus is created). This is the subtree of the species tree starting from the

634   time (and branch) of the creation of the locus. We then generate duplication, loss, and

635   transfer events, as detailed below.

Details. In order to formally describe this process, we first introduce some notation. Given a tree T, $V(T)$ and $E(T)$ are the sets of their nodes and edges (branches) respectively. For a node $v \in V(T)$, e_v represents the unique edge in $E(T)$ which has v as its target node (the node at the bottom of the edge).

Now, given a unilocus tree S_l and the corresponding haplotype tree G_l and haplotype forest $\mathcal{G}_l$, we simulate losses as described above, producing a sequence of events $e^l = \{(b_i, t_i)\}$, where $b_i \in E(G_l)$, and $t_i \geqslant 0$ is the time of occurrence of the loss (how the haplotype tree and forest are generated is described in the next section). Note that we must have $t_2 < t_i < t_1$, where t_1 and t_2 are the times of the top and bottom nodes of branch b_i respectively. At each event, we decorate G_l at that branch and time. An example of this process is shown in Figure 22.

We then simulate (as described above):

1. unlinked duplications in $e^{ud} = \{(b_i, t_i)\}$, where $b_i \in E(S_l)$;

2. transfers in $e^t = \{(b_i, t_i)\}$, where $b_i \in E(S)$;

3. linked duplications in $e^{ld} = \{(b_i, t_i)\}$, where $b_i \in E(S_l)$.

In all cases, the same constraints on t_i given for losses apply. At each duplication, we decorate S_l at that branch and time, and at each transfer, we choose and decorate the origin branch of S_l at that time. An example of the generation of unlinked duplications is given in Figure 23.

Fig. 22: Losses are sampled from the haplotype tree with constant rate. A, B, and C are species labels, while a, b, and c are gene labels. Here, we have $e^l = \{(e_a, t_1)\}$.

637   Now, the effect of each event is applied in a forwards-in-time order. Losses truncate

638 the haplotype tree, while duplications and transfers create new loci and unilocus trees.

Fig. 23: Duplications are sampled with a coalescent-rate process on the unilocus tree. Here, we have $\boldsymbol{e^{ud}} = \{(e_Y, t_2), (e_A, t_3)\}$.

Details. At each loss event, the haplotype tree is truncated on branch b_i at time t_i, with further events on the same or descendant branches of G_l (but not S_l) having no effect. (Note that the haplotype forest is *not* truncated.)

At each duplication event $(b_i, t_i) \in \boldsymbol{e^{ud}} \cup \boldsymbol{e^{ld}}$, a new locus m is created, with a unilocus tree S_m which is a copy of the subtree of S_l which starts at time t_i on the species branch b_i (see Figure 24).

Likewise, at each transfer event $(b_i, t_i) \in \boldsymbol{e^t}$, b_i represents the target of the transfer, not the source (as discussed above). The new locus will then have a unilocus tree created, which is a copy of the subtree of S starting from branch b_i at the transfer time t_i.

(a) For the duplication at time t_2, the new unilocus tree is the parent unilocus tree with the branch e_Y slightly shorter, i.e., starting at time t_2 instead of T.

(b) For the duplication at time t_3, the new unilocus tree is the leaf branch e_A of the parent unilocus tree, starting at time t_3.

Fig. 24: The new unilocus trees generated by the events in Figure 22: (a) a duplication at (e_Y, t_2) and (b) a duplication at (e_A, t_3). The unilocus trees are depicted as shadings on the species tree, with height equal to the duplication time.

Generating a Haplotype Forest (lines 13–19, Algorithm 2)

Once we have generated the unilocus tree for a new locus, we must then simulate the haplotype tree and forest. If the new locus is unlinked to the parent locus (i.e., it is created by transfer or unlinked duplication), then the two loci evolve completely independently. We can then use the incomplete coalescent process as specified in Section "Modelling Copy Number Hemiplasy". For the new locus m, the haplotype forest $\mathcal{G}_m$ is the collection of trees produced by the incomplete coalescent, while the haplotype tree G_m is the single tree selected by the incomplete coalescent (lines 14–15, Algorithm 2).

Suppose a locus l has a linked duplication, creating a new locus m. We need to consider that the genealogies of the two loci are dependent, and take this into account when constructing the haplotype tree and forest (lines 17–19, Algorithm 2). We have some genealogical information from the parent locus l, represented by the haplotype forest $\mathcal{G}_l$.

> **Details.** Since S_m is a copy of a subtree of S_l, we can identify all subtrees of the haplotype forest $\mathcal{G}_l$ which lie in the subtree of S_l corresponding to S_m. Formally, if locus m is created from an event (b_i, t_i), then we take all subtrees of $\mathcal{G}_l$ which lie in the species branch b_i and its descendants, truncated so that they start at time t_i. These subtrees are then copied to the new locus m (see Figure 25b for an example).

We now run the 'linked multispecies coalescent' in the new unilocus tree S_m, which we now detail.

Details. We start from a single lineage in each extant species, as with the ordinary multispecies coalescent. Because there is only one sampled individual per population, the lineage in each species begins coalesced to the copied subtree which has that species as a leaf. By construction, each extant species has one leaf in $\mathcal{G}_l$, so there is always exactly one possibility for this.

The coalescent then proceeds backwards in time, where the lineages which are coalesced with a copied subtree follow the same genealogy as that subtree. In particular, lineages which are coalesced with copied subtrees will coalesce with each other when the corresponding subtrees coalesce in the parent locus. However, with a constant rate r_r, each lineage which is coalesced with a copied subtree may have a recombination event (i.e., a recombination occurs between the two loci). If there are no recombination events, the haplotype forest in the new locus will be identical to that of the parent locus (Figure 25a).

When a recombination event occurs, the lineage in the new locus becomes 'uncoalesced' from the copied subtree, representing the fact that the individual in the population had different ancestors for the two loci. This lineage now becomes 'free' (not coalesced with a copied subtree) and is also followed backwards in time, where it may coalesce with a copied subtree of $\mathcal{G}_l$, or with another lineage in the new locus (which may itself be coalesced with a copied subtree, or not). Note that lineages in the new locus m which coalesce with each other cannot uncoalesce, since they exist in the same locus; a lineage can only uncoalesce with a copied subtree from the parent locus l, because a copied subtree represents a 'backbone' for the process rather than a lineage of the new locus. An example of this process is given in Figure 25b.

This process continues backwards in time, until we reach the time of creation of locus m.

This produces the haplotype forest $\mathcal{G}_m$ for the new locus. At this time, we will have a certain number $|\mathcal{G}_m|$ of lineages, some of which may be coalesced with copied subtrees. We choose one uniformly at random, and the descendant tree becomes the new haplotype tree G_m.

This process only produces the haplotype tree and forest in the new locus m. The haplotype tree must then be joined back to the haplotype tree in the parent locus, which we detail in Section "Assembling the Full Gene Tree" below.

(a) Each lineage begins coalesced with a copied subtree. Here no recombination events occur, so the new haplotype forest is identical to the parent one.

(b) When a recombination event occurs, the lineage 'uncoalesces' from the copied subtree. It may then subsequently coalesce with another lineage or copied subtree.

Fig. 25: The linked multispecies coalescent. The subtrees of the haplotype forest in the parent locus are copied to the new locus (dashed black lines). Then, a new multispecies coalescent (in red) is run, with possible recombination events. Finally, one of the red lineages is chosen to be the root of the new haplotype tree. (On the left side of the figure, we use the convention "dashed implies not chosen as haplotype tree".)

Note. Observe how this process produces the appropriate results for either extreme of the recombination rate: if $r_r = 0$ (no recombination), then no recombination events occur, and the haplotype tree G_m in the new locus will be a subtree of $\mathcal{G}_l$ (it is possible however that it will not be a subtree of G_l, in which case the duplication does not carry a copy of the gene and must be discarded; see the next section). On the other hand, as $r_r \to \infty$ ('infinite' recombination), then any coalescence with $\mathcal{G}_l$ will immediately become uncoalesced, so the haplotype tree G_m will be completely independent from $\mathcal{G}_l$ as desired.

The linked coalescent is very similar to the model of Slatkin and Pollack (2006), with a few subtle differences. In our model, we have fixed haplotype information for one locus, and simulate the second locus conditioned on that information, whereas the model of Slatkin and Pollack considers both loci simultaneously. They are also used in different contexts, as Slatkin and Pollack (2006) are concerned only with the probability that the two haplotype trees will be identical. Finally, their model is limited to only 3 species; certainly there is no theoretical barrier to extending it to more species, but it quickly becomes too complicated for theoretical analysis, rather than simulation.

⁶⁶⁵   If a locus generates no duplications or transfers, it will not produce any new loci.

⁶⁶⁶ Thus almost surely, the process of generating new unilocus and haplotype trees/forests will

⁶⁶⁷ stop. When this happens, we assemble the full gene tree by concatenating each haplotype

⁶⁶⁸ tree in a created locus to the haplotype tree in its parent locus.

(a) Here the red subtree is chosen to be the new haplotype tree. As it has coalesced with the haplotype tree copied from the parent locus, it is joined directly to that tree at the creation time of the new locus.

(b) Here the blue subtree is chosen to be the new haplotype tree. It has coalesced with a copied subtree which is not the haplotype tree in the parent locus, so it is discarded.

Fig. 26: If a duplication is ancestral, the lineage is joined directly to subtree with which it has coalesced. (On the left side of the figure, we use the convention "dashed implies not chosen as haplotype tree".)

Details. Recall that linked duplications may be ancestral (the duplicating lineage survives in the parent locus) or non-ancestral (otherwise). Suppose that we wish to re-attach the haplotype tree from a locus m created by a linked duplication, to its parent locus l. Once the haplotype tree and forest is generated in locus m, we can recognise three cases:

1. The lineage which is chosen to be the haplotype tree G_m in locus m is coalesced (at the time of creation of locus m) with (a copy of) the haplotype tree G_l in locus l. Then the duplication is an ancestral duplication, and we attach G_m to G_l at the lineage and time where the duplication occurred (Figure 26a, lines 20–21, Algorithm 2).

2. The lineage which is chosen to be the haplotype tree G_m is coalesced with a copied subtree of $\mathcal{G}_l$ which is not the haplotype tree G_l. This corresponds to an ancestral duplication of a lineage in locus l which does not carry a copy of the gene. Thus locus m will not carry a copy of the gene, and the locus (along with the duplication, and all loci derived from m) is discarded (Figure 26b, lines 22–23, Algorithm 2).

3. The lineage which is chosen to be the haplotype tree G_m is not coalesced with a copied subtree of $\mathcal{G}_l$. Then the duplication is a non-ancestral duplication, and is joined as specified below.

Events which create an unlinked locus (transfers and unlinked duplications) can always be considered to be non-ancestral; since the new locus is independent of the parent locus, the fact that the gene survives in the new locus gives no information about whether it survives in the parent locus (i.e., is ancestral). Therefore the probability of survival in the parent locus is extremely small ($O(\frac{1}{2N})$), and can be safely ignored.

Now, if locus m is created by a non-ancestral event (without loss of generality we will say it is a duplication), the duplicated gene does not survive in the parent locus l. Therefore we can treat the duplicating individual as simply another member of the population in the originating branch of S_l at the duplication time t_m. We follow this lineage backwards in time (in locus l) via the multispecies coalescent to where it coalesces with $\mathcal{G}_l$. Again, there are three cases (this corresponds to lines 24–25, Algorithm 2):

1. The lineage coalesces with the haplotype tree G_l. We then attach G_m to G_l at the point where the coalescence occurs (Figure 27a).

2. The lineage coalesces with an element of $\mathcal{G}_l$ which is not the haplotype tree G_l. This corresponds to a lineage with no copy of the gene, so locus m is discarded (Figure 27b).

3. The lineage does not coalesce with $\mathcal{G}_l$ by the time of creation of locus l. Again, this lineage does not carry a copy of the gene, so locus m is discarded (Figure 27c).

If we do attach G_m to G_l, we must finally check that the lineage from the root of G_m to the point where it coalesces with G_l has no losses on it, which we do by running a constant-rate loss process (with rate r_l) on that branch. If a loss exists on that branch, the new locus m (and all derived loci) must be discarded (lines 26–28, Algorithm 2).

(a) Here the green subtree is chosen to be the new haplotype tree. It coalesces with the haplotype tree in the parent locus, and is joined at the coalescent point.

(b) Here the green subtree is chosen to be the new haplotype tree. It coalesces with a lineage which is not in the haplotype tree in the parent locus, so it is discarded.

(c) Here the green subtree is chosen to be the new haplotype tree. It does not coalesce by the time of creation of the parent locus, so it is discarded.

Fig. 27: If a duplication is non-ancestral, we follow the lineage backwards in time until it coalesces with the haplotype forest in the parent locus. (On the left side of the figure, we use the convention "dashed implies not chosen as haplotype tree".)

⁶⁷⁰   This process is done backwards in time order by first attaching (or discarding) each

⁶⁷¹ haplotype tree with no child loci to the haplotype trees of their parent loci, and then those

to the haplotype trees of their parent loci, and so on until the full gene tree is assembled.

The full pseudocode for this algorithm is given in Algorithms 1 and 2. For clarity,

we have chosen to be brief in the specification of the multispecies coalescent processes in

lines 14, 18, and 25; for full details, refer to the above discussion. In particular, in line 25,

the incomplete coalescent "joining" may fail to actually join G_m to G_l^+, as shown in

Figures 27b and 27c, in which case locus m is discarded.

Algorithm 1 CONSTRUCT_GENE_TREE

Input: Species tree S with branch lengths in coalescent units, event parameters $\mathbf{r} = (r_d, r_t, r_l, r_r, p_u)$

Output: Gene tree G

1: $S_0 \leftarrow S$
2: $G_0 \leftarrow$ multispecies coalescent within S_0
3: $\mathcal{G}_0 \leftarrow \{G_0\}$
4: $G_0^+ \leftarrow$ ADD_NEW_LOCI$(S_0, G_0, \mathcal{G}_0, G_0, \mathbf{r})$
5: **return** G_0^+

Algorithm 2 ADD_NEW_LOCI
___

Input: Unilocus tree S_l, haplotype tree G_l, haplotype forest $\mathcal{G}_l$, partial gene tree G_l^+, event parameters $\mathbf{r}$

Output: Gene tree G_l^+

1:   $\boldsymbol{e^{ud}} \leftarrow$ D coalescent-rate process on S_l with rate parameter $p_u r_d$
2:   $\boldsymbol{e^t} \leftarrow$ T coalescent-rate process on S with rate parameter r_t, scaled by the proportion of origin species in S_l
3:   $\boldsymbol{e^{ld}} \leftarrow$ D coalescent-rate process in the presence of $\mathcal{G}_l$ on S_l with rate parameter $(1-p_u)r_d$ and recombination rate r_r
4:   $\boldsymbol{e^l} \leftarrow$ L constant-rate process on G_l with rate parameter r_l
5:   **for** $(b,t) \in \boldsymbol{e^{ud}} \cup \boldsymbol{e^{ld}} \cup \boldsymbol{e^t} \cup \boldsymbol{e^l}$ in time order **do**
6:      **if** $(b,t) \in \boldsymbol{e^l}$ **then**   ▷ here $b \in E(G_l)$
7:          $g \leftarrow$ a unary node on G_l^+ at time t along the branch b
8:          $G_l^+ \leftarrow$ cut G_l^+ from g
9:          remove all events from $\boldsymbol{e^l}$ below b which occur at a later time
10:     **else if** $(b,t) \in \boldsymbol{e^{ud}} \cup \boldsymbol{e^{ld}} \cup \boldsymbol{e^t}$ **then**   ▷ here $b \in E(S)$
11:          create a new locus m
12:          $S_m \leftarrow$ subtree of S_l rooted at b at time t
13:          **if** $(b,t) \in \boldsymbol{e^{ud}} \cup \boldsymbol{e^t}$ **then**
14:             $\mathcal{G}_m \leftarrow$ incomplete coalescent within S_m to time t
15:             $G_m \leftarrow$ an element of $\mathcal{G}_m$
16:          **else if** $(b,t) \in \boldsymbol{e^{ld}}$ **then**
17:             $\mathcal{G}'_m \leftarrow$ all subtrees of $\mathcal{G}_l$ within S_m
18:             $\mathcal{G}_m \leftarrow$ linked coalescent in the presence of $\mathcal{G}'_m$ within S_m to time t, with recombination rate r_r
19:             $G_m \leftarrow$ an element of $\mathcal{G}_m$
20:          **if** $(b,t) \in \boldsymbol{e^{ld}}$ and G_m has coalesced with a subtree of G_l at time t **then**
21:             attach G_m to G_l^+ at time t
22:          **else if** $(b,t) \in \boldsymbol{e^{ld}}$ and G_m has coalesced with a subtree of $\mathcal{G}_l$ at time t **then**
23:             discard locus m
24:          **else**
25:             $G_l^+ \leftarrow$ incomplete coalescent "joining" of G_m to G_l^+ in the presence of $\mathcal{G}_l$ within S_l (keep G_l^+)
26:               $\boldsymbol{e^l} \leftarrow$ L constant-rate process from root of G_m to G_l^+ with rate parameter r_l
27:               **if** $|\boldsymbol{e^l}| > 0$ **then**
28:                  discard locus m
29:          $G_l^+ \leftarrow$ ADD_NEW_LOCI$(S_m, G_m, \mathcal{G}_m, G_l^+, \mathbf{r})$
30: **return** G_l^+

Acknowledgements

YBC and CS acknowledge the Partenariat Hubert Curien (Hubert Curien Partnership) for providing grants for collaborative travel. QL would like to acknowledge Geoffrey Law for providing assistance with programming.

REFERENCES

Arvestad, L., Berglund, A.-C., Lagergren, J., and Sennblad, B. (2004). Gene tree reconstruction and orthology analysis based on an integrated model for duplications and sequence evolution. In *RECOMB 2004*, pages 326–335. ACM.

Arvestad, L., Lagergren, J., and Sennblad, B. (2009). The gene evolution model and computing its associated probabilities. *J. ACM*, 56(2):1–44.

Avise, J. C. and Robinson, T. J. (2008). Hemiplasy: a new term in the lexicon of phylogenetics. *Syst. Biol.*, 57(3):503–507.

Boussau, B. and Scornavacca, C. (2020). Reconciling Gene trees with Species Trees. In Scornavacca, C., Delsuc, F., and Galtier, N., editors, *Phylogenetics in the Genomic Era*, pages 3.2:1–3.2:23. No commercial publisher | Authors open access book.

Chan, Y., Ranwez, V., and Scornavacca, C. (2017). Inferring incomplete lineage sorting, duplications, transfers and losses with reconciliations. *J. Theor. Biol.*, 432:1–13.

David, L. A. and Alm, E. J. (2011). Rapid evolutionary innovation during an archaean genetic expansion. *Nature*, 469(7328):93.

Doyon, J.-P., Ranwez, V., Daubin, V., and Berry, V. (2011). Models, algorithms and programs for phylogeny reconciliation. *Brief. Bioinform.*, 12(5):392–400.

Doyon, J.-P., Scornavacca, C., Gorbunov, K. Y., Szöllősi, G. J., Ranwez, V., and Berry, V. (2010). An efficient algorithm for gene/species trees parsimonious reconciliation with losses, duplications and transfers. In *RECOMB International Workshop on Comparative Genomics*, pages 93–108. Springer.

Du, H., Ong, Y. S., Knittel, M., Mawhorter, R., Liu, N., Gross, G., Tojo, R., Libeskind-Hadas, R., and Wu, Y.-C. (2019). Multiple optimal reconciliations under the duplication-loss-coalescence model. *IEEE/ACM Trans. Comput. Biol. Bioinform.*

Durand, D., Halldórsson, B. V., and Vernot, B. (2006). A hybrid micro–macroevolutionary approach to gene tree reconstruction. *J. Comput. Biol.*, 13(2):320–335.

Fry, B. G., Roelants, K., Champagne, D. E., Scheib, H., Tyndall, J. D., King, G. F., Nevalainen, T. J., Norman, J. A., Lewis, R. J., Norton, R. S., et al. (2009). The toxicogenomic multiverse: convergent recruitment of proteins into animal venoms. *Annu Rev Genomics Hum Genet*, 10:483–511.

Goodman, M., Czelusniak, J., Moore, G. W., Romero-Herrera, A. E., and Matsuda, G. (1979). Fitting the gene lineage into its species lineage, a parsimony strategy illustrated by cladograms constructed from globin sequences. *Syst. Biol.*, 28(2):132–163.

Guerrero, R. F. and Hahn, M. W. (2018). Quantifying the risk of hemiplasy in phylogenetic inference. *Proc. Natl. Acad. Sci. U.S.A*, 115(50):12787–12792.

Hobolth, A., Dutheil, J. Y., Hawks, J., Schierup, M. H., and Mailund, T. (2011). Incomplete lineage sorting patterns among human, chimpanzee, and orangutan suggest recent orangutan speciation and widespread selection. *Genome Res.*, 21(3):349–356.

Hudson, R. R. (1983). Properties of a neutral allele model with intragenic recombination. *Theor. Popul. Biol.*, 23(2):183–201.

Kingman, J. F. (1982). On the genealogy of large populations. *J. Appl. Probab.*, 19(A):27–43.

Kubatko, L. S. and Degnan, J. H. (2007). Inconsistency of phylogenetic estimates from concatenated data under coalescence. *Syst. Biol.*, 56(1):17–24.

Kutschera, V. E., Bidon, T., Hailer, F., Rodi, J. L., Fain, S. R., and Janke, A. (2014). Bears in a forest of gene trees: phylogenetic inference is complicated by incomplete lineage sorting and gene flow. *Mol. Biol. Evol.*, 31(8):2004–2017.

Maddison, W. P. (1997). Gene trees in species trees. *Syst. Biol.*, 46(3):523–536.

Mallo, D., de Oliveira Martins, L., and Posada, D. (2015). Simphy: phylogenomic simulation of gene, locus, and species trees. *Syst. Biol.*, 65(2):334–344.

Mawhorter, R., Liu, N., Libeskind-Hadas, R., and Wu, Y.-C. (2019). Inferring pareto-optimal reconciliations across multiple event costs under the duplication-loss-coalescence model. *BMC Bioinform.*, 20(20):639.

Niimura, Y., Matsui, A., and Touhara, K. (2014). Extreme expansion of the olfactory receptor gene repertoire in african elephants and evolutionary dynamics of orthologous gene groups in 13 placental mammals. *Genome Res.*, 24(9):1485–1496.

Olender, T., Jones, T. E., Bruford, E., and Lancet, D. (2020). A unified nomenclature for vertebrate olfactory receptors. *BMC Evol. Biol.*, 20:1–12.

Pamilo, P. and Nei, M. (1988). Relationships between gene trees and species trees. *Mol. Biol. Evol*, 5(5):568–583.

Rannala, B. and Yang, Z. (2003). Bayes estimation of species divergence times and ancestral population sizes using DNA sequences from multiple loci. *Genetics*, 164(4):1645–1656.

Rasmussen, M. D. and Kellis, M. (2011). A bayesian approach for fast and accurate gene tree reconstruction. *Mol. Biol. Evol.*, 28(1):273–290.

Rasmussen, M. D. and Kellis, M. (2012). Unified modeling of gene duplication, loss, and coalescence using a locus tree. *Genome Res.*, 22(4):755–765.

Roch, S. and Steel, M. (2015). Likelihood-based tree reconstruction on a concatenation of aligned sequence data sets can be statistically inconsistent. *Theor. Popul. Biol.*, 100:56–62.

Romiguier, J., Gayral, P., Ballenghien, M., Bernard, A., Cahais, V., Chenuil, A., Chiari, Y., Dernat, R., Duret, L., Faivre, N., et al. (2014). Comparative population genomics in animals uncovers the determinants of genetic diversity. *Nature*, 515(7526):261–263.

Schrempf, D. and Szöllősi, G. J. (2018). Phylogenetic incongruences - opportunities to improve the reconstruction of a dated tree of life.

Scornavacca, C. and Galtier, N. (2017). Incomplete lineage sorting in mammalian phylogenomics. *Syst. Biol.*, 66(1):112–120.

Sjöstrand, J., Arvestad, L., Lagergren, J., and Sennblad, B. (2013). Genphylodata: realistic simulation of gene family evolution. *BMC Bioinform.*, 14(1):209.

Slatkin, M. and Pollack, J. L. (2006). The concordance of gene trees and species trees at two linked loci. *Genetics*, 172(3):1979–1984.

Stolzer, M., Lai, H., Xu, M., Sathaye, D., Vernot, B., and Durand, D. (2012). Inferring duplications, losses, transfers and incomplete lineage sorting with nonbinary species trees. *Bioinformatics*, 28(18):i409–i415.

Szöllősi, G. J., Tannier, E., Lartillot, N., and Daubin, V. (2013). Lateral gene transfer from the dead. *Syst. Biol.*, 62(3):386–397.

Tofigh, A., Hallett, M., and Lagergren, J. (2011). Simultaneous identification of duplications and lateral gene transfers. *IEEE/ACM Trans. Comput. Biol. Bioinform. (TCBB)*, 8(2):517–535.

Vernot, B., Stolzer, M., Goldman, A., and Durand, D. (2008). Reconciliation with non-binary species trees. *J. Comput. Biol.*, 15(8):981–1006.

Wu, Y.-C., Rasmussen, M. D., Bansal, M. S., and Kellis, M. (2014). Most parsimonious reconciliation in the presence of gene duplication, loss, and deep coalescence using labeled coalescent trees. *Genome Res.*, 24(3):475–486.

Zhang, L. (1997). On a Mirkin-Muchnik-Smith conjecture for comparing molecular phylogenies. *J. Comput. Biol.*, 4(2):177–187.