Tradeoffs between mobility and infectiousness in the spatial spread of an emerging

pathogen

Kate E. Langwig[1]*, J. Paul White[2], Katy L. Parise[3], Heather M. Kaarakka[2], Jennifer A.

Redell[2], John E. DePue[4], William H. Scullon[5], Jeffrey T. Foster[3], A. Marm Kilpatrick[6,Ψ],

Joseph R. Hoyt[1,Ψ]

Affiliations:

[1] Department of Biological Sciences, Virginia Polytechnic Institute, Blacksburg, VA,

24060 USA

[2] Wisconsin Department of Natural Resources, Madison, WI, 53703 USA

[3] Pathogen and Microbiome Institute, Northern Arizona University, Flagstaff, AZ 86011,

USA

[4] Michigan Department of Natural Resources, Baraga, MI 49870, USA

[5] Michigan Department of Natural Resources, Norway, MI 49908, USA

[6] Department of Ecology and Evolutionary Biology, University of California, Santa Cruz,

CA 95064 USA

*Corresponding author: klangwig@gmail.com; [Ψ]these authors contributed equally

24   **Abstract**

25   1. Emerging infectious diseases can have devastating effects on host communities,

26   causing population collapse and species extinctions. The timing of novel pathogen

27   arrival into naïve species communities can have consequential effects that shape

28   the trajectory of epidemics through populations. Pathogen introductions are often

29   presumed to occur when hosts are highly mobile. However, spread patterns can be

30   influenced by a multitude of other factors including host body condition and

31   infectiousness.

32   2. White-nose syndrome (WNS) is a seasonal emerging infectious disease of bats,

33   which is caused by the fungal pathogen *Pseudogymnoascus destructans*. Within-

34   site transmission of *P. destructans* primarily occurs over winter, however the

35   influence of bat mobility and infectiousness on the seasonal timing of pathogen

36   spread to new populations is unknown. We combined data on host population

37   dynamics and pathogen transmission from 22 bat communities to investigate the

38   timing of pathogen arrival, in relation to host mobility and infectiousness, and the

39   consequences of varying pathogen arrival times on disease impacts.

40   3. We found that midwinter arrival of the fungus predominated spread patterns,

41   suggesting that bats were most likely to spread *P. destructans* when they are

42   highly infectious, but have reduced mobility. In communities where *P.*

43   *destructans* was detected in early winter, one species suffered higher fungal

44   burdens and experienced more severe declines than at sites where the pathogen

45   was detected later in the winter, suggesting that the timing of pathogen

46   introduction had consequential effects for some bat communities. We also found

evidence of source-sink population dynamics over winter, suggesting some

movement among sites occurs during hibernation, even though bats at northern

latitudes were thought to be fairly immobile during this period. Winter emergence

behavior symptomatic of white-nose syndrome may further exacerbate these

winter bat movements to uninfected areas.

4. Our results suggest that low infectiousness during host migration may have

reduced the rate of expansion of this deadly pathogen, and that elevated

infectiousness outweighs differences in seasonal mobility. Furthermore, our

results highlight the importance of both accurate estimation of the timing of

pathogen spread and the consequences of varying arrival times to prevent and

mitigate the effects of infectious diseases.

1. Introduction:

The seasonality of pathogen spread is important for understanding, predicting, and

controlling disease outbreaks. Pathogens infecting highly mobile hosts often have rapid

rates of spread (Conner & Miller 2004; Altizer *et al.* 2006; Altizer, Bartel & Han 2011;

Dalziel, Pourbohloul & Ellner 2013). However, key tradeoffs exist between mobility and

disease, which can affect the likelihood that hosts spread pathogens during periods of

high mobility (Kiesecker *et al.* 1999; Norris & Evans 2000; Wendland *et al.* 2010;

Shakhar & Shakhar 2015). For example, migration can affect host body condition and

immune status which could increase host susceptibility to infection (Altizer, Bartel &

Han 2011). However, once a highly mobile host becomes infected, behavioral responses,

such as sickness behavior, are likely to decrease mobility (Van Gils *et al.* 2007;

Bouwman & Hawley 2010). Pathogen spread should therefore be most likely to occur

when hosts are both highly infectious and mobile, but given tradeoffs caused by sickness

behavior, peak mobility and infectiousness may not temporally align (Shakhar & Shakhar

2015). Understanding the relative importance of these factors in the seasonal timing of

pathogen spread is critical because varying pathogen arrival times can have consequential

effects on hosts that shape the trajectory of epidemics (Dalziel *et al.* 2018).

White-nose syndrome (WNS) is an annual seasonal epidemic that occurs in bat

communities during winter (Langwig *et al.* 2015a; Hoyt *et al.* 2020). The fungus that

causes WNS, *Pseudogymnoascus destructans*, is psychrophilic and grows only at cool

temperatures (1–20°C) (Verant *et al.* 2012) which restricts fungal replication and

infection into bats' epidermal tissue to periods when bats are in torpor (e.g. cool their

body temperatures <20°C) (Meteyer *et al.* 2009; Langwig *et al.* 2015a; Langwig *et al.*

2016). White-nose syndrome disrupts bat physiology during hibernation, causing

increased arousals that result in bats prematurely burning through fat stores before the

end of winter (Lorch *et al.* 2011; Reeder *et al.* 2012; Warnecke *et al.* 2012; Warnecke *et*

al. 2013; Verant *et al.* 2014). The spread of *P. destructans* has resulted in dramatic

declines in bat communities across North America, and threatens several species with

extinction (Frick *et al.* 2010; Langwig *et al.* 2012; Frick *et al.* 2015; Langwig *et al.*

2016). Seasonal changes in host behavior and physiology paired with seasonal

differences in pathogen growth may influence the timing of pathogen spread to new

communities.

The seasonal timing of *P. destructans* spread among sites is unknown but may

occur during autumn when bats are highly mobile during mating and move among caves

and mines (Davis & Hitchcock 1965; Cope & Humphrey 1977; Thomas, Fenton &

Barclay 1979; Glover & Altringham 2008), which are known reservoirs of the fungus

(Hoyt *et al.* 2020). Autumn swarm behavior is an important source of gene flow when

individuals move large distances between otherwise disconnected populations and

contact among mating individuals could lead to pathogen transmission (Veith *et al.* 2004;

Arnold 2007; Glover & Altringham 2008; Rossiter *et al.* 2012; Wilder, Kunz & Sorenson

2015). However, the spread of WNS has occurred relatively slowly compared with other

disease systems with similarly mobile hosts (Kilpatrick *et al.* 2006; Kilpatrick, LaDeau &

Marra 2007). In addition, *P. destructans* is less prevalent on bats during autumn, and

infected individuals typically have reduced infection levels (i.e. 100–1000-fold lower

than in midwinter; (Langwig *et al.* 2015a)), suggesting that spread may occur outside of

the highly mobile autumn period. If infectiousness is more important than mobility in *P.*

destructans spread among sites, then pathogen introduction to new communities could

occur during winter when fungal loads on bats are highest (Langwig *et al.* 2015a).

However, most species are thought to be relatively sedentary during winter, particularly

at northern latitudes where temperatures fall well below freezing (Davis & Hitchcock

1965; Davis 1970; Fenton & Barclay 1980; Fujita & Kunz 1984; Caceres & Barclay

2000). Therefore, further investigation is needed to disentangle the importance of host

mobility and infectiousness in the spread of *P. destructans*.

We investigated the timing of arrival of *P. destructans* into 22 hibernating bat

communities in the Midwestern U.S. We hypothesized that pathogen invasion would be

more likely to occur during winter when bats are less mobile but highly infectious,

resulting in the first detections of *P. destructans* at each site occurring in late, rather than

early winter. Given the potential importance of seasonally varying arrival times on

disease dynamics, we also assessed the influence of timing of *P. destructans*

introductions on disease severity, hypothesizing that earlier introductions of *P.*

destructans would result in higher infection prevalence, fungal loads, and population

impacts.

2. Methods:

2.1 Study sites and data collection

We studied patterns of *P. destructans* arrival at 22 sites in the Midwestern U.S.

Over a 4-year period, we collected data on population dynamics and infection status of

four hibernating bat species (*Myotis lucifugus*, *Myotis septentrionalis*, *Eptesicus fuscus*,

and *Perimyotis subflavus*). We used epidermal swab sampling to determine the presence

and abundance of *P. destructans* on bats at two time points during hibernation

(November - early hibernation, and March - late hibernation). During each visit, we

counted all bats present and identified bats to the species level. In addition, we installed

HOBO temperature and humidity loggers at 1–4 locations within a site to determine roost

temperature and humidity.

2.2 Sample testing

We sampled bats using a standardized protocol (Langwig *et al.* 2015a) and stored

swabs in RNAlater® for sample preservation until extraction. We tested samples for *P.*

destructans DNA using real-time PCR (Muller *et al.* 2013) and quantified fungal loads

based on the cycle threshold (C_t) value to estimate a fungal load on each bat, with a cut-

off of 40 cycles. Quantification of serial dilutions of the DNA from 10 ng to 1000 fg

resulted in C_t scores ranging from 17.33 to 30.74 and a quantification relationship of $C_t =$

$-3.348*\log_{10}(P.\ destructans[\text{ng}]) + 22.049$, $r^2 = 0.986$.

2.3 Statistical analysis

We assessed our power to detect *P. destructans* arrival at a site by calculating an

expected early prevalence at each site where *P. destructans* was not detected in early

hibernation based on a weighted mean prevalence that incorporated the average

prevalence of each species at sites where *P. destructans* was detected in early hibernation

multiplied by the proportional abundance of a given species at each site. We then

calculated the probability of missing *P. destructans* at a site as the probability of getting

all negatives in bats given the expected prevalence at a site multiplied by the probability

of missing *P. destructans* in the unsampled bats (Appendix).

We investigated the effect of timing of *P. destructans* arrival on late winter

prevalence by fitting a generalized linear mixed model with a binomial distribution and a

logit link. We included fixed effects for the timing of *P. destructans* detection (early or

late hibernation) interacting with the effect of species, and included site as a random

effect. We also examined the effect of timing of *P. destructans* detection on fungal loads

using a linear mixed model with species and timing of detection as interacting fixed

effects, and site as a random effect. Lastly, we assessed the effect of the timing of *P.*

destructans introduction on $\log_{10}$ bat population growth rates, calculated as the annual

change in late winter counts at each site, using linear mixed models with species

interacting with timing of detection as fixed effects and site as a random effect. We

explored the inclusion of autumn prevalence as a continuous variable (0 for all sites

where *P. destructans* was not detected, prevalence ranged between 0.01 and 0.25 for sites

where *P. destructans* was detected during autumn) and the results were qualitatively

similar to those shown, so we present the discretized (early or late winter *P. destructans*

detection) for simplicity of visualization and interpretation.

Lastly, we investigated evidence of overwinter movements of bats among sites.

At each site, we calculated our response variable, overwinter λ, as the proportional

change in the number of bats of each species at a site by dividing the late hibernation

count by the early hibernation count. We then used a linear mixed model to assess the

relationship between $\log_{10}$(overwinter λ) and $\log_{10}$(early hibernation colony size) with an

additive or interactive effect of species and site as a random effect.

3. Results:

During early hibernation, we sampled 706 total bats from 22 sites in the year that

P. destructans was first detected (site mean +/- SE = 67.4 +/- 1.79). We first detected *P.*

destructans in early hibernation at 32% of sites (7/22), whereas in 68% of sites (15/22),

the detection of *P. destructans* first occurred during our late hibernation visit (Figure 1).

A total of 468 bats (site mean +/- SE = 31.1 +/- 1.25) were sampled from sites that were

negative for *P. destructans* during early hibernation and became positive later in the same

winter. After accounting for the relatively high proportion of bats sampled in each site

(an average of 41% of the total bats at each site) and variable estimated "true" prevalence

among sites (weighted mean prevalence of the species sampled from sites where *P.*

destructans was detected in early hibernation: mean = 0.07, range = 0.05, 0.10), the mean

probability of missing *P. destructans* at each negative site was 0.063 (median = 0.039,

range: 0.005–0.3266), suggesting fairly good power to detect *P. destructans* presence.

Across sites where *P. destructans* was first detected during early hibernation, we

were more likely to detect *P. destructans* on *M. lucifugus*, *M. septentrionalis*, and *E.*

fuscus than on *P. subflavus* (*P. subflavus* intercept: -1.82 +/- 0.66, *M. lucifugus* coeff:

1.79 +/- 0.38 P < 0.0001, *E. fuscus* coeff: 1.40 +/- 0.51, P = 0.006, *M. septentrionalis*

coef: 1.44 +/- 0.39, P = 0.0002, Appendix). In sites where *P. destructans* was first

detected in late hibernation, we were also less likely to detect *P. destructans* on *P.*

subflavus than on any other species (*P. subflavus* intercept: -1.69 +/- 0.31, *M. lucifugus*

coeff: 0.94 +/- 0.22 P < 0.0001, *E. fuscus* coeff: 0.66 +/- 0.30, P = 0.0264, *M.*

septentrionalis coeff: 0.84 +/- 0.26, P = 0.0014, Appendix).

The timing of *P. destructans* introduction influenced disease dynamics and

population impacts for some, but not all, bat species (Figure 2). For *M. septentrionalis*,

prevalence of *P. destructans*, fungal loads, and population impacts during late winter

were significantly higher at sites where *P. destructans* was first detected in early

hibernation (one-tailed P-values = 0.032, 0.005, 0.0004, respectively, Appendix). For *M.*

lucifugus, prevalence was also significantly higher in sites where *P. destructans* was first

detected in early hibernation (one-tailed P = 0.04, Appendix), although the effect on

fungal loads and population impacts was less clear (Appendix). There was no clear effect

of the timing of *P. destructans* introduction on prevalence, fungal loads, or population

impacts for either *E. fuscus* or *P. subflavus*, possibly because these species had lower

prevalence and fungal loads than *M. septentrionalis* and *M. lucifugus* in the first year of

detection (Figure 2, Appendix).

We found that prior to the arrival of WNS, there were significant changes in

population counts between early and late hibernation (Figure 3). Across species, as

colony size decreased in early hibernation, immigration increased (early hibernation

colony size coeff: -0.15 +/- 0.06, t = -2.66) such that smaller colonies had proportionally

more immigrants than larger colonies, indicative of a general spreading out of bats across

sites during the hibernation season. We found no support for an interactive model over an

additive model (ΔAIC = -9.63), suggesting no clear indication that the slope of this effect

differed among species. The changes in overwinter counts occurred despite frigid

minimum temperatures that were consistently below 5°C throughout this period (Figure

S1), suggesting that some movement among hibernacula continues to occur during winter

when bats are most infected with *P. destructans*.

Lastly, we explored the effect of multiple covariates on the probability that *P.*

destructans arrived at a site outside of or during the hibernation season. We found that

higher mean relative humidity increased the probability that *P. destructans* was first

detected at a site during early hibernation (coeff: 0.22 +/- 0.09, P = 0.026), and found no

clear effects of any of the other covariates we explored. However, there were non-

significant trends such that sites with higher overwinter mobility (larger proportional

increases in the bat populations) were more likely to be infected during the hibernation

season than sites with lower overwinter mobility (coeff: -1.36 +/- 1.53, P = 0.38), and

sites with warmer mean temperatures were more likely to be infected in early hibernation

(coeff: 0.21 +/- 0.13, P = 0.12; Figure S2, Appendix).

Figure 1

(a) Change in overwinter prevalence of *Pseudogymnoascus destructans* in the year that *P. destructans* was first detected at a site. Lines shows fits of a generalized linear mixed model with a binomial distribution with timing of first detection interacting with species and an additive effect of date with site as a random effect. (b) Twice as many sites had the first detection of *P. destructans* in late winter as in early winter.

Figure 2

The timing of *Pseudogymnoascus destructans* introduction (early or late hibernation)

influences late winter infection prevalence +/- SE (a), fungal loads (b), and population

impacts +/- SE (c), in some species. In sites where *P. destructans* was first detected

during late hibernation *M. septentrionalis* had significantly higher infection prevalence,

fungal loads, and population impacts by the end of winter in the first year of WNS.

Figure 3

Pre-WNS immigration rates show dynamic populations during winter. Panels show bat

species and colors indicate different bat populations. Lines show the fitted relationship

between $\log_{10}$ early winter counts of bats and the rate of change in the same population

over winter. Despite cold (<5°C) temperatures, there was considerable immigration and

emigration among hibernation sites, consistent with bats moving from larger to smaller

colonies over winter.

4. Discussion:

Our results indicate that host infectiousness was more important in determining

timing of pathogen spread than host mobility. We found that *P. destructans* was more

likely to be introduced to sites during winter hibernation than during autumn, suggesting

that elevated infectiousness overwinter outweighed the increase in activity and long-

distance movements that occur during autumn. Bats were previously thought to be

restricted in their movement among sites during hibernation due to extremely cold

temperatures (< -5C; Fig S1) and a lack of prey availability. However, we found evidence

that limited movement may be occurring among hibernacula during the winter, and given

their increased levels of infection, provide a higher probability of successful invasion to

new sites. The lower infectiousness of bats during autumn when they are highly mobile

has likely decreased the rate of geographic expansion of the fungus as the majority of

spread occurs during the hibernation period. As *P. destructans* moves into less connected

western bat populations (Wilder, Kunz & Sorenson 2015; Lorch *et al.* 2016), spread

could be further dampened by larger distances among colonies (Wilder, Kunz &

Sorenson 2015; Weller *et al.* 2018).

Although the absolute absence of a pathogen is difficult to confirm, large sample

sizes paired with the differential consequences of the timing of *P. destructans*

introduction on disease dynamics, suggest that we did not simply fail to detect *P.*

destructans during early winter. Our results also demonstrate that the timing of *P.*

destructans introduction was more important for some species than others. Early

introduction of *P. destructans* resulted in higher mortality in *M. septentrionalis*

populations, but not *M. lucifugus.* Previous laboratory studies have shown that infection

duration prior to death was 90-120 days for *M. lucifugus*. The absence of clear mortality

in *M. lucifugus* populations while *M. septentrionalis* suffered from higher impacts

suggests that *M. septentrionalis* may have a shorter time from infection until death.

Infection in *P. subflavus* and *E. fuscus* were lower in the first year of WNS than *M. spp.*,

likely due to low environmental contamination (Langwig *et al.* 2015b; Hoyt *et al.* 2020)

and fewer contacts over the hibernation period (Hoyt *et al.* 2018).

We found evidence of substantial overwinter movement among hibernacula prior

to the arrival of WNS, with a general trend of bats spreading out from larger to smaller

colony size sites. Other studies have found that some species (especially the larger *E.*

fuscus) can move in very cold temperatures (Klüg-Baerwald *et al.* 2016), although winter

flight of *Myotis* spp. is thought to be a relatively rare occurrence (Davis & Hitchcock

1965). In addition, symptoms of WNS include mid-hibernation emergence onto the

winter landscape (Carr, Bernard & Stiver 2014; Bernard & McCracken 2017). This

sickness behavior may increase movements between hibernacula and the spread of *P.*

destructans.

Our results also support that environmental conditions of hibernacula may be

important determinants of *P. destructans* establishment into new communities outside of

the hibernation season. While bats with low fungal loads are likely moving among

uninfected hibernacula during autumn, humidity and potentially temperature appear to be

important in determining whether *P. destructans* can successfully establish in a

population. Given the limited number of sites where *P. destructans* arrived outside of the

hibernation season, it was difficult to ascertain the importance of a number of factors that

have been implicated by other studies (Wilder *et al.* 2011; Maher *et al.* 2012; Wilder,

Kunz & Sorenson 2015), and therefore additional research is needed to determine which

site-specific differences may influence the timing of pathogen spread.

Importantly, many studies rely on disease stage (e.g. invading, epidemic or

endemic) to draw inferences about geographic disease risk, population declines, and

community persistence. Given the high probability of *P. destructans* introduction to new

sites during winter, early and midwinter surveys could miss the introduction of *P.*

destructans, and thus falsely conclude pathogen absence during the first year of arrival.

Our results suggest that the arrival time of *P. destructans* can substantially influence

dynamics, and therefore could be responsible for unexplained variance in transmission,

impacts, and structure of remnant populations when sites are grouped by disease stage.

Our ability to predict the timing and patterns of pathogen spread are fundamental

to the prevention and control of infectious disease outbreaks. The differential timing of

initial pathogen arrival can have important effects on disease dynamics and lasting

impacts to populations. This study suggests that host infectiousness is an important factor

in determining successful pathogen spread and that incorrectly attributing pathogen

spread to periods of high mobility would have masked the true underlying causes of

among site variation in transmission and population impacts. Future studies examining

the spatial and seasonal patterns of pathogen movement should consider the multitude of

factors that might influence spread patterns.

Acknowledgements: We thank Steffany Yamada, MI DNR, Eric McMaster and the many

landowners for site access. The research was funded by NSF grants DEB-1115895, DEB-

1336290, DEB-1911853 and the USFWS (F17AP00591).

Author Contributions: KEL analyzed the data and wrote the original draft; KEL, AMK

and JRH revised the manuscript with input from all authors; KLP and JTF tested the

samples and KEL, JPW, HMK, JAR, JED, WHS, AMK, and JRH collected the data.

Data Availability Statement:

Data will archived at VTData Repository upon manuscript acceptance or reviewer

request.

Altizer, S., Bartel, R. & Han, B.A. (2011) Animal Migration and Infectious Disease
Risk. *Science,* **331**, 296-302.
Altizer, S., Dobson, A., Hosseini, P., Hudson, P., Pascual, M. & Rohani, P. (2006)
Seasonality and the dynamics of infectious diseases. *Ecology Letters,* **9**, 467-
484.
Arnold, B.D. (2007) Population structure and sex-biased dispersal in the forest
dwelling vespertilionid bat, Myotis septentrionalis. *The American Midland*
Naturalist, **157**, 374-384.
Bernard, R.F. & McCracken, G.F. (2017) Winter behavior of bats and the progression
of white-nose syndrome in the southeastern United States. *Ecology and*
Evolution, **7**, 1487-1496.
Bouwman, K.M. & Hawley, D.M. (2010) Sickness behaviour acting as an evolutionary
trap? Male house finches preferentially feed near diseased conspecifics.
Biology Letters, **6**, 462-465.
Caceres, M.C. & Barclay, R.M.R. (2000) Myotis septentrionalis. *Mammalian Species*, 1-
4.
Carr, J.A., Bernard, R.F. & Stiver, W.H. (2014) Unusual bat behavior during winter in
Great Smoky Mountains National Park. *Southeastern naturalist,* **13**, N18-N21.
Conner, M.M. & Miller, M.W. (2004) Movement patterns and spatial epidemiology of
a prion disease in mule deer population units. *Ecological Applications,* **14**,
1870-1881.
Cope, J.B. & Humphrey, S.R. (1977) SPRING AND AUTUMN SWARMING BEHAVIOR
IN INDIANA BAT, MYOTIS-SODALIS. *Journal of Mammalogy,* **58**, 93-95.
Dalziel, B.D., Kissler, S., Gog, J.R., Viboud, C., Bjørnstad, O.N., Metcalf, C.J.E. & Grenfell,
B.T. (2018) Urbanization and humidity shape the intensity of influenza
epidemics in US cities. *Science,* **362**, 75-79.
Dalziel, B.D., Pourbohloul, B. & Ellner, S.P. (2013) Human mobility patterns predict
divergent epidemic dynamics among cities. *Proceedings of the Royal Society*
B: Biological Sciences, **280**, 20130763.

Davis, W. (1970) Hibernation: ecology and physiological ecology. *Biology of Bats,* **1,**
265 - 300.
Davis, W.H. & Hitchcock, H.B. (1965) Biology and migration of the bat, Myotis
lucifugus, in New England. *Journal of Mammalogy,* **46,** 296-313.
Fenton, M.B. & Barclay, R.M.R. (1980) Myotis lucifugus. *Mammalian Species,* 1-8.
Frick, W.F., Pollock, J.F., Hicks, A.C., Langwig, K.E., Reynolds, D.S., Turner, G.G.,
Butchkoski, C.M. & Kunz, T.H. (2010) An emerging disease causes regional
population collapse of a common north american bat species. *Science,* **329,**
679-682.
Frick, W.F., Puechmaille, S.J., Hoyt, J.R., Nickel, B.A., Langwig, K.E., Foster, J.T., Barlow,
K.E., Bartonička, T., Feller, D., Haarsma, A.J., Herzog, C., Horacek, I., van der
Kooij, J., Mulkens, B., Petrov, B., Reynolds, R., Rodrigues, L., Stihler, C.W.,
Turner, G.G. & Kilpatrick, A.M. (2015) Disease alters macroecological
patterns of North American bats. *Global Ecology and Biogeography,* **24,** 741-
749.
Fujita, M.S. & Kunz, T.H. (1984) Pipistrellus subflavus. *Mammalian Species,* 1-6.
Glover, A.M. & Altringham, J.D. (2008) Cave selection and use by swarming bat
species. *Biological Conservation,* **141,** 1493-1504.
Hoyt, J.R., Langwig, K.E., Sun, K., Parise, K.L., Li, A., Wang, Y., Huang, X., Worledge, L.,
Miller, H. & White, J.P. (2020) Environmental reservoir dynamics predict
global infection patterns and population impacts for the fungal disease
white-nose syndrome. *Proceedings of the National Academy of Sciences,* **117,**
7255-7262.
Hoyt, J.R., Langwig, K.E., White, J.P., Kaarakka, H., Redell, J., Kurta, A., DePue, J.,
Scullon, W.H., Parise, K.L., Foster, J.T., Frick, W.F. & Kilpatrick, A.M. (2018)
Cryptic connections illuminate pathogen transmission within community
networks. *Nature.*
Kiesecker, J.M., Skelly, D.K., Beard, K.H. & Preisser, E. (1999) Behavioral reduction of
infection risk. *Proceedings of the National Academy of Sciences of the United*
States of America, **96,** 9165-9168.
Kilpatrick, A.M., Chmura, A.A., Gibbons, D.W., Fleischer, R.C., Marra, P.P. & Daszak, P.
(2006) Predicting the global spread of H5N1 avian influenza. *Proceedings of*
the National Academy of Sciences of the United States of America, **103,** 19368-
19373.
Kilpatrick, A.M., LaDeau, S.L. & Marra, P.P. (2007) Ecology of west nile virus
transmission and its impact on birds in the western hemisphere. *Auk,* **124,**
1121-1136.
Klüg-Baerwald, B.J., Gower, L.E., Lausen, C. & Brigham, R. (2016) Environmental
correlates and energetics of winter flight by bats in Southern Alberta,
Canada. *Canadian Journal of Zoology,* **94,** 829-836.
Langwig, K.E., Frick, W.F., Bried, J.T., Hicks, A.C., Kunz, T.H. & Marm Kilpatrick, A.
(2012) Sociality, density-dependence and microclimates determine the
persistence of populations suffering from a novel fungal disease, white-nose
syndrome. *Ecology Letters,* **15,** 1050-1057.

Langwig, K.E., Frick, W.F., Hoyt, J.R., Parise, K.L., Drees, K.P., Kunz, T.H., Foster, J.T. &
Kilpatrick, A.M. (2016) Drivers of variation in species impacts for a multi-
host fungal disease of bats. *Phil. Trans. R. Soc. B,* **371,** 20150456.
Langwig, K.E., Frick, W.F., Reynolds, R., Parise, K.L., Drees, K.P., Hoyt, J.R., Cheng, T.L.,
Kunz, T.H., Foster, J.T. & Kilpatrick, A.M. (2015a) Host and pathogen ecology
drive the seasonal dynamics of a fungal disease, white-nose syndrome.
Proceedings of the Royal Society B: Biological Sciences, **282.**
Langwig, K.E., Hoyt, J.R., Parise, K.L., Kath, J., Kirk, D., Frick, W.F., Foster, J.T. &
Kilpatrick, A.M. (2015b) Invasion dynamics of white-nose syndrome white-
nose syndrome fungus, midwestern United States, 2012-2014. *Emerg Infect*
Dis, **21.**
Lorch, J.M., Meteyer, C.U., Behr, M.J., Boyles, J.G., Cryan, P.M., Hicks, A.C., Ballmann,
A.E., Coleman, J.T.H., Redell, D.N., Reeder, D.M. & Blehert, D.S. (2011)
Experimental infection of bats with Geomyces destructans causes white-nose
syndrome. *Nature,* **480,** 376-378.
Lorch, J.M., Palmer, J.M., Lindner, D.L., Ballmann, A.E., George, K.G., Griffin, K.,
Knowles, S., Huckabee, J.R., Haman, K.H., Anderson, C.D., Becker, P.A.,
Buchanan, J.B., Foster, J.T. & Blehert, D.S. (2016) First Detection of Bat White-
Nose Syndrome in Western North America. *mSphere,* **1,** e00148-00116.
Maher, S.P., Kramer, A.M., Pulliam, J.T., Zokan, M.A., Bowden, S.E., Barton, H.D.,
Magori, K. & Drake, J.M. (2012) Spread of white-nose syndrome on a network
regulated by geography and climate. *Nature Communications,* **3.**
Meteyer, C.U., Buckles, E.L., Blehert, D.S., Hicks, A.C., Green, D.E., Shearn-Bochsler, V.,
Thomas, N.J., Gargas, A. & Behr, M.J. (2009) Histopathologic criteria to
confirm white-nose syndrome in bats. *Journal of Veterinary Diagnostic*
Investigation, **21,** 411-414.
Muller, L.K., Lorch, J.M., Lindner, D.L., O'Connor, M., Gargas, A. & Blehert, D.S. (2013)
Bat white-nose syndrome: a real-time TaqMan polymerase chain reaction
test targeting the intergenic spacer region of Geomyces destructans.
Mycologia, **105,** 253-259.
Norris, K. & Evans, M.R. (2000) Ecological immunology: life history trade-offs and
immune defense in birds. *Behavioral Ecology,* **11,** 19-26.
Reeder, D.M., Frank, C.L., Turner, G.G., Meteyer, C.U., Kurta, A., Britzke, E.R., Vodzak,
M.E., Darling, S.R., Stihler, C.W., Hicks, A.C., Jacob, R., Grieneisen, L.E.,
Brownlee, S.A., Muller, L.K. & Blehert, D.S. (2012) Frequent Arousal from
Hibernation Linked to Severity of Infection and Mortality in Bats with White-
Nose Syndrome. *Plos One,* **7.**
Rossiter, S.J., Zubaid, A., MOHD - ADNAN, A., Struebig, M.J., Kunz, T.H., Gopal, S., Petit,
E.J. & Kingston, T. (2012) Social organization and genetic structure: insights
from codistributed bat populations. *Molecular Ecology,* **21,** 647-661.
Shakhar, K. & Shakhar, G. (2015) Why Do We Feel Sick When Infected—Can
Altruism Play a Role? *Plos Biology,* **13,** e1002276.
Thomas, D.W., Fenton, M.B. & Barclay, R.M.R. (1979) Social behavior of the Little
brown bat, Myotis lucifugus. I. mating behavior. *Behavioral Ecology and*
Sociobiology, **6,** 129-136.

Van Gils, J.A., Munster, V.J., Radersma, R., Liefhebber, D., Fouchier, R.A. & Klaassen, M.
451   (2007) Hampered foraging and migratory performance in swans infected
452   with low-pathogenic avian influenza A virus. *Plos One,* **2,** e184.
453   Veith, M., Beer, N., Kiefer, A., Johannesen, J. & Seitz, A. (2004) The role of swarming
454   sites for maintaining gene flow in the brown long-eared bat (Plecotus
455   auritus). *Heredity,* **93,** 342.
456   Verant, M.L., Boyles, J.G., Waldrep, W., Wibbelt, G. & Blehert, D.S. (2012)
457   Temperature-dependent growth of Geomyces destructans, the fungus that
458   causes bat white-nose syndrome. *Plos One,* **7**.
459   Verant, M.L., Carol, M.U., Speakman, J.R., Cryan, P.M., Lorch, J.M. & Blehert, D.S.
460   (2014) White-nose syndrome initiates a cascade of physiologic disturbances
461   in the hibernating bat host. *BMC physiology,* **14,** 10.
462   Warnecke, L., Turner, J.M., Bollinger, T.K., Lorch, J.M., Misra, V., Cryan, P.M., Wibbelt,
463   G., Blehert, D.S. & Willis, C.K.R. (2012) Inoculation of bats with European
464   Geomyces destructans supports the novel pathogen hypothesis for the origin
465   of white-nose syndrome. *Proceedings of the National Academy of Sciences of*
466   *the United States of America,* **109,** 6999-7003.
467   Warnecke, L., Turner, J.M., Bollinger, T.K., Misra, V., Cryan, P.M., Blehert, D.S.,
468   Wibbelt, G. & Willis, C.K.R. (2013) Pathophysiology of white-nose syndrome
469   in bats: a mechanistic model linking wing damage to mortality. *Biology*
470   *Letters,* **9**.
471   Weller, T.J., Rodhouse, T.J., Neubaum, D.J., Ormsbee, P.C., Dixon, R.D., Popp, D.L.,
472   Williams, J.A., Osborn, S.D., Rogers, B.W. & Beard, L.O. (2018) A review of bat
473   hibernacula across the western United States: Implications for white-nose
474   syndrome surveillance and management. *Plos One,* **13**.
475   Wendland, L.D., Wooding, J., White, C.L., Demcovitz, D., Littell, R., Berish, J.D., Ozgul,
476   A., Oli, M.K., Klein, P.A., Christman, M.C. & Brown, M.B. (2010) Social behavior
477   drives the dynamics of respiratory disease in threatened tortoises. *Ecology,*
478   **91,** 1257-1262.
479   Wilder, A.P., Frick, W.F., Langwig, K.E. & Kunz, T.H. (2011) Risk factors associated
480   with mortality from white-nose syndrome among hibernating bat colonies.
481   *Biology Letters,* **7,** 950-953.
482   Wilder, A.P., Kunz, T.H. & Sorenson, M.D. (2015) Population genetic structure of a
483   common host predicts the spread of white‐nose syndrome, an emerging
484   infectious disease in bats. *Molecular Ecology,* **24,** 5495-5506.
485