White pupae genes in the Tephritids *Ceratitis capitata, Bactrocera dorsalis* and
Zeugodacus cucurbitae: a story of parallel mutations

Short title: **Genetic mutations causing white pupae phenotypes**

Ward CM[a,1], Aumann RA[b,1], Whitehead MA[c], Nikolouli K[d], Leveque G [e,f], Gouvi G[d,g], Fung E[h],
Reiling SJ[e], Djambazian H[e], Hughes MA[c], Whiteford S[c], Caceres-Barrios C[d], Nguyen TNM[a,k],
Choo A[a], Crisp P[a,h], Sim S[i], Geib S[i], Marec F[j], Häcker I[b], Ragoussis J[e], Darby AC[c], Bourtzis
K[d,*], Baxter SW[k,*], Schetelig MF[b,*]

[a] School of Biological Sciences, University of Adelaide, Australia, 5005

[b] Justus-Liebig-University Gießen, Institute for Insect Biotechnology, Department of Insect
Biotechnology in Plant Protection, Winchesterstr. 2, 35394 Gießen, Germany

[c] Centre for Genomic Research, Institute of Integrative Biology, The Biosciences Building, Crown Street,
Liverpool, L69 7ZB, United Kingdom

[d] Insect Pest Control Laboratory, Joint FAO/IAEA Division of Nuclear Techniques in Food and
Agriculture, Seibersdorf, A-1400 Vienna, Austria

[e] McGill University Genome Centre, McGill University, Montreal, Quebec, Canada

[f] Canadian Centre for Computational Genomics (C3G), McGill University, Montreal, Quebec, Canada

[g] Department of Environmental Engineering, University of Patras, 2 Seferi str., 30100 Agrinio, Greece

[h] South Australian Research and Development Institute, Waite Road, Urrbrae, South Australia 5064

[i] USDA-ARS Daniel K. Inouye US Pacific Basin Agricultural Research Center, 64 Nowelo Street, Hilo,
Hawaii, 96720, USA

[j] Biology Centre, Czech Academy of Sciences, Institute of Entomology, Branišovská 31, 370 05 České
Budějovice, Czech Republic

[k] Bio21 Molecular Science and Biotechnology Institute, School of BioSciences, University of Melbourne,
Australia, 3010

1 Authors contributed equally to the study

* Corresponding authors:

k.bourtzis@iaea.org
simon.baxter@unimelb.edu.au
marc.schetelig@agrar.uni-giessen.de

Contributions

RA, CMW, CC, PC, SS, SG, IH, JR, ACD, KB, SWB, MFS designed research; CMW, RA,
MAW, KN, GG, EF, SR, MAH, CC, TNMN, AC, SS, SG, ACD, KB, SWB, MFS performed
research; RA, CMW, HD, GL, FM, JR, KB, SWB, MFS contributed new reagents/analytic tools;
CMW, RA, MAW, KN, GL, GG, HD, SW, TNMN, AC, SS, SG, IH, JR, ACD, KB, SWB, MFS
analyzed data; RA, CMW, KN, GL, GG, SR, SW, AC, SS, SG, IH, JR, ACD, KB, SWB, MFS
wrote the paper.

Abstract (<250 words)

Flooding insect pest populations with huge numbers of sterilized males is an effective mean of biological control as they mate with, but cannot fertilize, wild females. The greatest challenge of the sterile insect technique (SIT) is the removal of unrequired factory reared females prior to sterilization and release. Spontaneous white-pupae (wp^-) color mutations have been integrated as a dimorphic selectable marker into historical SIT strains for three major tephritid fruit fly pests, *Bactrocera dorsalis*, *Ceratitis capitata* and *Zeugodacus cucurbitae*. Here we identify parallel genetic mutations causing the phenotype in all three species using diverse experimental approaches. The *B. dorsalis wp^-* locus was introgressed into a related species, *Bactrocera tryoni,* and whole-genome sequencing identified a 37 bp truncating mutation within a gene containing a Major Facilitator Superfamily domain (MFS). In *C. capitata* and *Z. cucurbitae*, cytogenetics, comparative genomics and transcriptomic analysis of strains carrying brown and white pupae phenotypes identified an 8,150 bp insertion of a putative transposon into the *C. capitata* MFS ortholog and a 13 bp deletion in the *Z. cucurbitae* ortholog. In *B. tryoni* CRISPR/Cas9-mediated knock-out of the putative *Bt_wp* developed mosaic white and brown puparium colors in G_0, and G_1 progeny with recessive white pupae phenotypes. In *C. capitata,* complementation crosses of CRISPR-induced wp^- mutants to flies carrying the naturally occurring recessive wp^- mutation confirmed the role of the gene. Gene editing technology carries the potential for engineering white pupae phenotypes and generating dimorphic SIT strains in other tephritids or insect pest species.

Keywords (> 3 words)

introgression, chromosomal inversions, Sterile Insect Technique, genetic sexing strains, pest control, dimorphic markers

Significance statement (<120 words)

Mass releases of sterilized male insects within sterile insect technique programs have helped

suppress insect pest populations since the 1970s. In the major horticultural pests *Bactrocera*

dorsalis, Ceratitis capitata, and *Zeugodacus cucurbitae,* a key phenotype white pupae (wp)

has been used for decades to selectively remove females before releases, yet the gene

responsible remained unknown. Here we use classical and modern genetic approaches to

identify and functionally characterize causal wp^- mutations in distantly related fruit fly species

and show it can be used to rapidly generate novel wp^- strains. The conserved phenotype and

independent nature of the wp^- mutations suggest that this technique can provide a generic

approach to produce sexing strains in other significant medical and agricultural insect pests.

Introduction

Tephritid species, including the Mediterranean fruit fly (medfly) *Ceratitis capitata*, the oriental fruit fly *Bactrocera dorsalis*, the melon fly *Zeugodacus cucurbitae* and the Queensland fruit fly *Bactrocera tryoni*, are major agricultural pests worldwide (1). The sterile insect technique (SIT) is a species-specific and environment-friendly approach to control their populations, which has been successfully applied as a component of area-wide integrated pest management programs (2-4). The efficacy and cost-effectiveness of these large-scale operational SIT applications has been significantly enhanced by the development and use of genetic sexing strains (GSS) for Medfly, *B. dorsalis* and *Z. cucurbitae* (5, 6).

A GSS requires two principal components: a selectable marker, which could be phenotypic or conditionally lethal, and the linkage of the wild type allele of this marker to the male sex, ideally as close as possible to the male determining region. In a GSS, males are heterozygous and phenotypically wild type, whilst females are homozygous for the mutant allele thus facilitating sex separation (6-8). Pupal color was one of the first phenotypic traits exploited as a selectable marker for the construction of GSS. In all three species, brown is the typical pupae color. However, naturally occurring color mutants such as white pupae (*wp*) (9) and dark pupae (*dp*) (10) have occurred in the field or laboratory stocks. The *wp* locus was successfully used as a selectable marker to develop GSS for *C. capitata*, *B. dorsalis* and *Z. cucurbitae* (6, 11, 12), however, its genetic basis has never been resolved.

Biochemical studies provided evidence that the white pupae phenotype in medfly is due to a defect in the mechanism responsible for the transfer of catecholamines from the hemolymph to the puparial cuticle (13). In addition, classical genetic studies showed that the *wp* phenotype is due to a recessive mutation in an autosomal gene located on chromosome 5 of the medfly genome (9, 14). The development of translocation lines combined with deletion and transposition mapping and advanced cytogenetic studies allowed the localization of the gene responsible for the wp phenotype on the right arm of chromosome 5, at position 59B of the trichogen polytene chromosome map (15). In the same series of experiments, the *wp* locus was shown to be tightly linked to a *temperature-sensitive lethal (tsl)* gene (position 59B-61C), which is the second selectable marker of the VIENNA 7 and VIENNA 8 GSS currently used in all medfly SIT operational programs worldwide (7, 15).

The genetic stability of a GSS is a major challenge, mainly due to recombination phenomena taking place between the selectable marker and the translocation breakpoint. To address this risk, a chromosomal inversion called D53 was induced and integrated into the medfly VIENNA 8 GSS (VIENNA 8^{D53+}) (6, 8). Cytogenetic analysis indicated that the D53 inversion spans a large region of chromosome 5 (50B-59C on trichogen polytene chromosome map) with the *wp* locus being inside the inversion, close to its right breakpoint (6).

Extensive genetic and cytogenetic studies facilitated the development of a physical map of
the medfly genome (8, 16). The annotated gene set provided opportunities for the identification
of genes or loci-associated mutant phenotypes, such as the *wp* and *tsl*, used for the
construction of GSS (16, 17). Salivary gland polytene chromosome maps developed for *C.*
capitata, *B. dorsalis*, *Z. cucurbitae,* and *B. tryoni* show that their homologous chromosomes
exhibit similar banding patterns. In addition, *in situ* hybridization analysis of several genes
confirmed that there is extensive shared synteny, including the right arm of chromosome 5
where the *C. capitata wp* gene is localized (8). Interestingly, two recent studies identified SNPs
associated with the *wp* phenotype in *C. capitata* and *Z. cucurbitae* that were also on
chromosome 5 (18, 19).

In the present study, we employed different strategies involving genetics, cytogenetics,
genomics, transcriptomics, gene editing and bioinformatics to identify independent natural
mutations in a novel gene responsible for pupal coloration in three tephritid species of major
agricultural importance, *C. capitata*, *B. dorsalis*, and *Z. cucurbitae*. We then functionally
characterized causal mutations within this gene in *C. capitata* and *B. tryoni* resulting in
development of new white pupae strains. Due to its conserved nature (20) and widespread
occurrence in many insect species of agricultural and medical importance, we also discuss
the potential use of this gene as a generic selectable marker for the construction of GSS for
SIT applications.

Results

Resolving the *B. dorsalis wp* locus with interspecific introgression. The *B. dorsalis white pupae* phenotype was introgressed into *B. tryoni* to generate a strain referred to as the *Bactrocera Introgressed Line* (*BIL,* Fig. S1). To determine the proportion of *B. dorsalis* genome introgressed into *BIL*, whole genome sequence data from male and female *B. dorsalis*, *B. tryoni,* and *BIL* individuals were analyzed. Paired end Illumina short read data from single *B. oleae* males (SRR826808) and females (SRR826807) were used as an outgroup. Single copy orthologs across the genome (n = 1,846) were used to reconstruct the species topology revealing species-specific monophyly (Fig. 1A) consistent with previously published phylogenies (21, 22). Reconstruction also showed monophyly between *B. tryoni* and *BIL* across 99.2% of gene trees suggesting the majority of loci originally introgressed from *B. dorsalis* have been removed during backcrosses.

Genomes were partitioned into 100 kb windows and pairwise absolute genetic distance (d_{XY}) calculated between each species and *BIL* to estimate admixture. *B. dorsalis* was found to be highly similar to a small proportion of the *BIL* genome (Fig. 1B; purple), as indicated by d_{XY} values approaching the median value of *B. dorsalis* vs *B. tryoni* (Fig. 1B; orange).

Two formal tests for introgression were also carried out, the f estimator $\widehat{f_d}$ (Fig. 1C) and topology weighting (Fig. 1D). Three distinct local evolutionary histories (Fig. 1D) were tested using d_{XY} and topology weighting across the *B. dorsalis wp* Quantitative Trait Locus (QTL) i) *BIL* is closest to *B. tryoni* (Fig. 1D; purple, expected across most of the genome), ii) *BIL* is closest to *B. dorsalis* (Fig. 1D; orange, expected at the *wp⁻* locus), and iii) *BIL* is closest to *B. oleae* (Fig. 1D; green, a negative control). Across the nuclear genome the species topology was supported in 98.82% of windows. Both $\widehat{f_d}$ and topology weighting confirmed a lack of widespread introgression from *B. dorsalis* into *BIL* with few (n = 42) discordant outlier windows. Genomic windows discordant across all three tests were considered candidate regions for the *wp* mutation. Four scaffolds accounting for 1.18% of the *B. dorsalis* genome met these criteria and only two, NW_011876372.1 and NW_011876398.1, showed homozygous introgression consistent with a recessive white pupae phenotype (Fig. S2).

To resolve breakpoints within the *B. dorsalis wp* QTL, a 10 kb windowed analysis across NW_011876398.1 and NW_011876372.1 was performed using d_{XY} (Fig. 1E), topology weighting (Fig. 1F) and $\widehat{f_d}$ (Fig. 1G). The maximum range of the introgressed locus was 4.49 Mbp (NW_011876398.1 was 2.9-5.94 Mb and NW_011876372.1 was 0-1.55 Mb) (Fig. 1E-G). The *wp⁻* QTL was further reduced to a 2.71 Mb region containing 113 annotated protein coding genes through analyzing nucleotide diversity (π) among eight pooled BIL genomes (3.8 Mb on NW_011876398.1 to 0.73 Mb on scaffold NW_011876372.1, Fig. S2).

Fig. 1. Characterization of total introgression from *B. dorsalis* into the *Bactrocera Introgressed Line* and identification of the *white pupae* locus. (A) Species tree constructed from 1,846 single copy ortholog gene trees for four haplotypes of *B. oleae, B. dorsalis, B. tryoni* and *BIL.* Branches corresponding to *BIL* individuals are shown in blue*.* All nodes were well supported with posterior probabilities >0.97. (B) Nei's absolute genetic distance (d_{XY}) calculated for tiled 100 kb windows across the genome between *B. tryoni* vs *BIL (*Bt vs BIL*); B. tryoni* vs *B. dorsalis* (*Bt* vs *Bd*)*; B. tryoni* vs *B. oleae* (*Bt* vs *Bo*) and *BIL* vs *B. oleae* (*BIL* vs *Bo*). (C) The introgression estimator ($\widehat{f_d}$) calculated across tiled 100 kb windows to identify regions of disproportionately shared alleles between *BIL* and *B. dorsalis*, $\widehat{f_d}$ (*Bt*, *BIL*, *Bd*; *Bo*). (D) The three evolutionary hypothesis/topologies of interest to identify introgressed regions and their representation across the genome: species (purple, 98.82%), introgression (orange, 1.04%) and a negative control tree (green, 0.14%). (E) Nei's absolute genetic distance (d_{XY}) calculated for tiled 10 kb windows across the candidate *wp* locus colors follow the legend in (D). (F) Topology weighting for each topology shown in (D) calculated for 1 kb tiled local trees across the candidate *wp* locus. (G) The introgression estimator ($\widehat{f_d}$) calculated across tiled 10 kb windows, $\widehat{f_d}$ (*Bt*, *BIL*, *Bd*; *Bo*).

Resolving the *C. capitata* D53 inversion breakpoints and *wp* locus with genome sequencing and *in situ* hybridization. Previous cytogenetic studies determined the localization of the gene responsible for the white pupae phenotype on the right arm of chromosome 5, at position 59B of the trichogen polytene chromosome map (15). The equivalent of position 59B is position 76B of the salivary gland polytene chromosome map, inside but close to the right breakpoint of the D53 inversion (69C-76B on the salivary gland polytene chromosome map). Long read sequencing data were generated of the wild type strain Egypt II (EgII), the inversion line D53 and the genetic sexing strain VIENNA 8 (without

the inversion; VIENNA 8$^{D53-|-}$) (Table S1) to enable a comparison of the genomes and locate

the breakpoints of the D53 inversion, to subsequently narrow down the target region, and to

identify *wp* candidate genes.

Chromosome 5-specific markers (16) were used to identify the EgII_Ccap3.2.1 scaffold_5

as complete chromosome 5. Candidate D53 breakpoints in EgII scaffold_5 were identified

using the alignment of three genome datasets EgII, VIENNA 8$^{D53-|-}$, and D53 (see methods).

The position of the D53 inversion breakpoints was located between 25,455,334 – 25,455,433

within a scaffold gap (left breakpoint), and at 61,880,224 bp in a scaffolded contig (right

breakpoint) on EgII chromosome 5 (Ccap3.2.1; accession GCA_902851445). The region

containing the causal *wp* gene was known to be just next to the right breakpoint. Cytogenetic

analysis and *in situ* hybridization using the wild type EgII strain and the D53 inversion line

confirmed the overall structure of the inversion, covering the area of 69C-76B on the salivary

gland polytene chromosomes (Fig. 2), as well as the relative position of markers residing

inside and outside the breakpoints (Fig. 2 and Fig. S3). PCRs using two primer pairs flanking

the predicted breakpoints (Fig. S4C) and subsequent sequencing confirmed the exact

sequence of the breakpoints. Using a primer combination specific for the chromosome 5 wild

type status confirmed in the WT in EgII flies and VIENNA 7$^{D53+|-}$ GSS males, which are

heterozygous for the inversion. Correspondingly, these amplicons were not present in D53

males and females or in VIENNA 7$^{D53+|+}$ GSS females (all homozygous for the inversion) (Fig.

S4A-B). Positive signals for the inversion were detected in D53 and VIENNA 7^{D53+} GSS males

and females, but not in WT flies using an inversion-specific primer pair (Fig. S4D-E).

Fig. 2. Genomic positioning of the D53 inversion on chromosome 5 of *C. capitata*. (A)
Chromosome scale assembly of *C. capitata* EgII chromosome 5. Shown are the positions of *in situ*
mapped genes *white* (*w*), *6-phosphogluconate dehydrogenase* (*Pgd*), *glucose-6-phosphate 1-*
dehydrogenase (*Zw*) and *sex lethal* (*Sxl*), the position of the D53 inversion breakpoints (blue), and the
relative position of *white pupae* (*wp*) on the polytene chromosome map of chromosome 5 (23) and the
PacBio-Hi-C EgII scaffold_5, representing the complete chromosome 5 (genome Ccap3.2.1, accession
GCA_902851445). The position of the *yellow* gene (*y*, LOC101455502) was confirmed on chromosome
5 70A by *in situ* hybridization, despite its sequence not been included in the scaffold assembly. (D)
Schematic illustration of chromosome 5 without (EgII, WT) and with (D53) D53 inversion. The inverted
part of chromosome 5 is shown in light blue. Two probes, one inside (*y*, 70A) and one outside (*Pgd*,
68B) of the left inversion breakpoint were used to verify the D53 inversion breakpoints by *in situ*
hybridization. WT EgII is shown in (B) and (C), D53 in (E) and (F).

Genome and transcriptome sequencing lead to a single candidate *wp⁻* gene in *B. dorsalis, C. capitata, and Z. cucurbitae*.

Orthologs within the QTL of *B. dorsalis, C. capitata* and scaffolds previously identified to segregate with the *wp⁻* phenotype in *Z. cucurbitae* (NW_011863770.1 and NW_011863674.1 (18)) were investigated for high effect mutations under the assumption that a null mutation in a conserved gene results in the *wp⁻* phenotype. A single ortholog contained fixed indels absent from wild type strains was identified in each species. *White pupae B. dorsalis* and *BIL* strains showed a 37 bp frame-shift deletion in the first coding exon of LOC105232189 introducing a premature stop codon 210 bp from the transcription start site (Fig. 3). Presence of the deletion was confirmed *in silico* using whole genome resequencing from the *wp* and wildtype mapped to the reference, and by *de novo* assembly of Illumina RNAseq data transcripts (Fig. 3).

In *C. capitata*, *wp* individuals D53 Nanopore read alignment on EgII showed an independent approximate 8,150 bp insertion into the third exon of LOC101451947 disrupting proper gene transcription 822 bp from the transcription start site (Fig. 3). The insertion sequence is flanked by identical repeats, suggesting that it may originate from a transposable element insertion. The *C. capitata* mutation was confirmed *in silico,* as in *B. dorsalis,* using whole genome sequencing and RNAseq data (Fig. 3).

Transcriptome data from the white pupae-based genetic sexing strain of *Z. cucurbitae* revealed a 13 bp deletion in the third exon of LOC105216239 on scaffold NW_011863770.1 introducing a premature stop codon (Fig. 3).

The candidate *white pupae* gene in all three species had a reciprocal best BLAST hit to the putative metabolite transport protein CG14439 in *Drosophila melanogaster* and contains a Major Facilitator-like superfamily domain (MFS_1, pfam07690), suggesting a general function as a metabolite transport protein. *In situ* hybridization on polytene chromosomes of *B. dorsalis*, *C. capitata* and *Z. cucurbitae* was used to confirm the presence of the *wp* locus in the same syntenic position on the right arm of chromosome 5 (Fig. 3). Therefore, all three species show a mutation in the same positional orthologous gene likely to be responsible for the phenotype in all three genera.

Fig. 3. Identification of the *wp* mutation in the transcriptomes of *B. dorsalis*, *C. capitata* and *Z. cucurbitae*. The grey graphs show expression profiles from the candidate *wp* loci in WT (*wp*⁺) and mutant (*wp*⁻) flies at the immobile pupae stages of (A) *B. dorsalis*, (B) *C. capitata* and (C) *Z. cucurbitae*. The gene structure (not drawn to scale) is indicated below as exons (arrows labelled E1-E4) and introns (dashed lines). The positions of independent *wp* mutations (*Bd*: 37 bp deletion, *Cc*: approximate 8,150 bp insertion, *Zc*: 13 bp deletion) are marked with black boxes in the expression profiles and are shown in detail below the gene models based on *de novo* assembly of RNAseq data from WT and white pupae phenotype individuals. *In situ* hybridization on polytene chromosomes (right) confirmed the presence of the *wp* locus on the right arm of chromosome 5 in all three species (arrows in micrographs).

CRISPR/Cas9 knockout of a Major Facilitator Superfamily gene in *B. tryoni* and *C. capitata* causes the white pupae phenotype. An analogous *B. dorsalis wp*⁻ mutation was developed in *B. tryoni* by functional knockouts of the putative *Bt_wp* using the CRISPR/Cas9 system. A total of 591 embryos from the Ourimbah laboratory strain were injected using two guides with recognition sites in the first coding exon of this gene (Fig. 4A). Injected embryos surviving to adulthood (n = 19, 3.2%) developed with either wild type brown (n=12) or somatically mosaic white-brown puparia (n = 7, Fig. S5). Surviving G_0 adults were individually backcrossed into the Ourimbah strain, resulting in potentially $wp^{+/-(CRISPR)}$ heterozygous brown pupae (Fig. 4C). Five independent G_0 crosses were fertile (three mosaic white-brown and two brown pupae phenotypes), G_1 offspring were sibling mated and visual inspection of G_2

progeny revealed that three families contained *white pupae* individuals. Four distinct frameshift mutations were observed in screened G_2 progeny (Fig. 4A) suggesting functional KO of putative *Bt_wp* is sufficient to produce the *white pupae* phenotype in *B. tryoni*. Capillary sequencing of cloned *Bt_MFS* amplicons revealed deletions ranging from a total of 4 bp to 155 bp, summed across the two guide recognition sites, introducing premature stop codons.

In *C. capitata,* CRISPR/Cas9 gene editing was used to knock-out the orthologous gene and putative *Cc_wp*, LOC101451947, to confirm that it causes a wp^- phenotype. A mix of recombinant Cas9 protein and the gRNA_MFS, targeting the third exon and thereby the MFS domain of the presumed *Cc_wp* CDS (Fig. 4B), was injected into 588 EgII WT embryos of which 96 developed to larvae and 67 pupated. All injected G_0 pupae showed brown pupal color. In total, 29 G_0 males and 34 females survived to adulthood (9.3%) and were backcrossed individually or in groups (see material and methods) to a strain carrying the naturally occurring white pupae mutation ($wp^{-(nat)}$; strain #1402_22m1B) (24) (Fig. 4D). As w*hite pupae* is known to be monogenic and recessive in *C. capitata*, this complementation assay was used to reveal whether the targeted gene is responsible for the naturally occurring white pupae phenotype or if the mutation is located in a different gene. G_1 offspring would only show white pupae phenotypes if *Cc_wp* was indeed the *white pupae* gene, knocked-out by the CRISPR approach, and complemented by the natural mutation through the backcross ($wp^{-(nat)|-(CRISPR)}$). In the case that the *Cc_wp* is not the gene carrying the natural wp^- mutation, a brown phenotype would be observed for all offspring. Here, five out of 13 crosses, namely M1, M3, F2, F3, and F4, produced white pupae phenotype offspring. The crosses generated 221, 159, 70, 40, and 52 G_1 pupae, of which 10, 30, 16, 1, and 1 pupa respectively, were white. Fifty-seven flies emerged from white pupae were analyzed via non-lethal genotyping, and all of them showed mutation events within the target region. Overall, eight different mutation events were seen, including deletions ranging from 1-9 bp and a 46 bp deletion combined with a 2 bp insertion (Fig. 4B). Five mutation events (B, D, E, G, H) caused frameshifts and premature stop codons. The remaining three (A, C, F), however, produced deletions of only one to three amino acids. To make $wp^{-(CRISPR)}$ mutations homozygous, mutants were either inbred (mutation C) (Fig. 4D) or outcrossed to WT EgII (mutation A-H), both in groups according to their genotype. This demonstrated that *Cc_wp* is the gene carrying the $wp^{-(nat)}$, and that even the loss of a single amino acid without a frameshift at this position can cause the white pupae phenotype. Offspring from outcrosses of mutation A, D, and H, as well as offspring of the inbreeding (mutation C), were genotyped via PCR, and $wp^{+|-(CRISPR)}$ and $wp^{-(CRISPR)|-(CRISPR)}$ positive flies were inbred to establish pure new *white pupae* lines.

Fig. 4. CRISPR/Cas9-based generation of homozygous *wp*-(CRISPR) lines in *B. tryoni* and *C. capitata*. A schematic structure of the *wp* CDS exons (E1, E2, E3, E4) including the MFS domain in *B. tryoni* (A) and *C. capitata* (B) are shown. Positions of gRNAs targeting the first and third exon in *B. tryoni* and *C. capitata,* respectively, are indicated by green arrows. Nucleotide and amino acid sequences of mutant *wp* alleles identified in G_1 individuals are compared to the WT reference sequence in *B. tryoni* (A) and *C. capitata* (B). Deletions are shown as dashes, alterations on protein level leading to premature stop codons are depicted as asterisks highlighted in black. Numbers on the right side represent InDel sizes (bp = base pairs). Crossing schemes to generate homozygous *wp*-(CRISPR) lines in *B. tryoni* (C) and *C. capitata* (D) show different strategies to generate *wp* strains. Bright field images of empty puparia are depicted for both species. Genotype schematics and corresponding PCR analysis (for *C. capitata*) validating the presence CRISPR-induced (orange) and natural (blue, for *C. capitata*) *wp* mutations are shown next to the images of the puparia. (C) Injected G_0 *B. tryoni* were backcrossed to the Ourimbah laboratory strain resulting in uniformly brown G_1 offspring (depicted as illustration because no images were acquired during G_1). G_1 inbreeding led to G_2 individuals homozygous for the white pupae phenotype. (D) Injected WT G_0 flies were crossed to flies homozygous for the naturally occurring *wp*- allele (*wp*-(nat)). *wp*-(nat) (457 bp amplicon) and *wp*-(CRISPR) or WT (724 bp amplicon) alleles were identified by multiplex PCR (left lane; L = NEB 2log ladder). White pupae phenotypes in G_1 indicated positive CRISPR events. G_2 flies with a white pupae phenotype that were homozygous for *wp*-(CRISPR) allele were used to establish lines.

Discussion

White pupae (*wp*) was first identified in *C. capitata* as a spontaneous mutation and was subsequently adopted as a phenotypic marker of fundamental importance for the construction of GSS for SIT (6, 9). Full penetrance expressivity and recessive inheritance rendered *wp* the marker of choice for GSS construction in two additional tephritid species, *B. dorsalis* and *Z. cucurbitae* (11, 12), allowing automated sex sorting based on pupal color. This was only possible because spontaneous *wp* mutations occur at relatively high rates either in the field or in mass rearing facilities and can easily be detected (6, 9). Despite the easy detection and establishment of *wp* mutants in these three species, similar mutations have not been detected in other closely or distantly related species such as *B. tryoni*, *B. oleae,* or *Anastrepha ludens* despite large screens being conducted. In addition to being a visible GSS marker used to separate males and females, the *wp* phenotype is also important for detecting and removing recombinants in cases where sex separation is based on a conditional lethal gene such as the *tsl* gene in the medfly VIENNA 7 or VIENNA 8 GSS (6, 7). However, it took more than 20 years from the discovery and establishment of the *wp* mutants to the large-scale operational use of the medfly VIENNA 8 GSS for SIT applications (6, 9) and the genetic nature of the *wp* mutation remained unknown. The discovery of the underlying *wp* mutations and the availability of CRISPR/Cas genome editing would allow the fast recreation of such phenotypes and sexing strains in other insect pests. Isolation of the *wp* gene would also facilitate future efforts towards the identification of the closely linked *tsl* gene.

Using an integrated approach consisting of genetics, cytogenetics, genomics, transcriptomics and bioinformatics, we identified the white pupae (*wp*) genetic locus in three major tephritid agricultural pest species, *B. dorsalis*, *C. capitata,* and *Z. cucurbitae*.

Our study clearly shows the power of employing different strategies for gene discovery, one of which was species hybridization. In *Drosophila*, hybridization of different species has played a catalytic role in the deep understanding of species boundaries and the speciation processes, including the evolution of mating behavior and gene regulation (25-29). In our study, we took advantage of two closely related species, *B. dorsalis* and *B. tryoni*, which can produce fertile hybrids and be backcrossed for consecutive generations. This allowed the introgression of the *wp* mutant locus of *B. dorsalis* into *B. tryoni,* resulting in the identification of the introgressed region, including the causal *wp* mutation via whole-genome resequencing and advanced bioinformatic analysis.

In *C. capitata*, we exploited two essential pieces of evidence originating from previous genetic and cytogenetic studies: the localization of wp to region 59B and 76B on chromosome 5 in the trichogen cells and salivary gland polytene chromosome map, respectively (15, 30), and its position close to the right breakpoint of the large inversion D53 (6). This data prompted

us to undertake a comparative genomic approach to identify the exact position of the right
breakpoint of the D53 inversion, which would bring us in the vicinity of the *wp* gene. Coupled
with comparative transcriptomic analysis, this strategy ensured that the analysis indeed
tracked the specific *wp* locus on the right arm of chromosome 5, instead of any mutation in
another, random locus which may participate in the pigmentation pathway and therefore result
in the same phenotype. Functional characterization via CRISPR/Cas9-mediated knock-out
resulted in the establishment of new white pupae strains in *C. capitata* and *B. tryoni* and
confirmed that this gene is responsible for the pupal coloration in these tephritid species.
Interestingly, the wp phenotype is based on three independent and very different natural
mutations of this gene, a rather large and transposon-like insertion in *C. capitata*, but only
small deletions in the two other tephritids, *B. dorsalis* and *Z. cucurbitae*. Also in medlfy,
however, CRSIRP-induced in-frame deletions of one or three amino acids in the MFS domain
were sufficient to induce the wp phenotype, underlining the importance of this domain for
correct pupal coloration.

It is worth noting, that in the first stages of this study, we employed two additional
approaches, which did not allow us to successfully narrow down the *wp* genomic region to the
desired level. The first was based on Illumina sequencing of libraries produced from laser
micro-dissected (Y;5) mitotic chromosomes that carry the wild type allele of the *wp* gene
through a translocation from the fifth chromosome to the Y. This dataset from the medfly
VIENNA 7 GSS was comparatively analyzed to wild type (Egypt II) Y and X chromosomes,
and the complete genomes of Egypt II, VIENNA 7 $^{D53-}$ GSS, and a D53 inversion line in an
attempt to identify the chromosomal breakpoints of the translocation and/or inversion, which
are close to the *wp* locus (Table S2). However, this effort was not successful due to the short
Illumina reads and the lack of a high-quality reference genome. The second approach was
based on individual scale whole-genome resequencing/genotyping, and identifying fixed loci
associated with pupal color phenotypes, which complemented the QTL analysis (19). Seven
loci associated with SNPs and larger deletions linked to the white pupae phenotype were
analyzed based on their respective mutations and literature searches for their potential
involvement in pigmentation pathways. However, we could not identify a clear link to the pupal
coloration as shown by *in silico*, molecular, and *in situ* hybridization analysis (Supplementary
text S1, Fig. S6 and S7, Table S3).

The *wp* gene is a member of a major facilitator superfamily (MFS). Orthologs of *white pupae*
are present in 146 of 148 insect species aggregated in OrthoDB v9 (20) present in all orders
and single copy in 133 species. Furthermore, wp is included in the benchmarking universal
single copy ortholog (BUSCO) gene set for insecta and according to OrthoDB v10 (31) has a
below average evolutionary rate (0.87, OrthoDB group 42284at50557) suggesting an
important and evolutionarily conserved function (Fig. S8). Its ortholog in *Bombyx mori*, *mucK*,

was shown to participate in the pigmentation at the larval stage (32) whereas in *D.*
melanogaster peak expression is during the prepupal stage after the larva has committed to
pupation (33), which is the stage where pupal cuticle sclerotization and melanization occurs.
It is known that the insect cuticle consists of chitin, proteins, lipids and catecholamines, which
act as cross-linking agents thus contributing to polymerization and the formation of the
integument (34). Interestingly, the sclerotization and melanization pathways are connected
and this explains the different mechanical properties observed in different medfly pupal color
strains with the "dark" color cuticles to be harder than the "brown" ones and the latter harder
than the "white" color ones (35). The fact that the white pupae mutants are unable to transfer
catecholamines from the haemolymph to the cuticle is perhaps an explanation for the lack of
the brown pigmentation (13).

The discovery of the long-sought *wp* gene in this study and the recent discovery of the
Maleness-on-the-Y (*MoY*) gene, which determines the male sex in several tephritids (36),
opens the way for the development of a 'generic approach' for the construction of GSS for
other species. Using CRISPR/Cas-based genome editing approaches, we can: (a) induce
mutations in the *wp* orthologues of SIT target species and establish lines with *wp* phenotype
and (b) link the rescue alleles as closely as possible to the *MoY* region. Given that the *wp*
gene is present in diverse insect species including agricultural insect pests and mosquito
disease vectors, this approach would allow more rapid development of GSS in SIT target
species. In principle, these GSS will have higher fertility compared to the semi-sterile
translocation lines (6). In addition, these new generation GSS will be more stable since the
rescue allele will be tightly linked to the male determining region thus eliminating
recombination which can jeopardize the genetic integrity of any GSS. The concept of the
'generic approach' can also be applied in species which lack a typical Y chromosome such as
Aedes aegypti and *Aedes albopictus*. In these species, the rescue allele should be transferred
close to the male determining gene (*Nix*) and the M locus (37, 38). It is hence important for
this 'generic approach' to identify regions close enough to the male determining loci to ensure
the genetic stability of the GSS and to allow the proper expression of the rescue alleles. In the
present study, we have already shown that CRISPR/Cas9-induced mutations resulting in the
white pupae phenotype can be developed in SIT target species and the resulting strains
provide already new opportunities for GSS based on visible markers.

Material and methods

All sequence libraries prepared during this study are publicly available on NCBI within the ENA BioProject PRJEB36344/ERP119522 (accession numbers ERS4426857 - ERS4426873, ERS4426994 - ERS4427029, ERS4519515; see table S1) and the BioProject PRJNA629430 (SRA accessions SRR11649127 - SRR11649132; see Supplementary text 1).

Insect rearing. *C. capitata*, *B. dorsalis* and *Z. cucurbitae* fly strains were maintained at $25\pm1°C$, 48% RH and 14/10 h light/dark cycle. They were fed with a mixture of sugar and yeast extract (3v:1v) and water. Larvae were reared on a gel diet, containing carrot powder (120 g/l), agar (3 g/l), yeast extract (42 g/l), benzoic acid (4 g/l), HCl (25%, 5.75 ml/l) and ethyl-4-hydroxybenzoate (2.86 g/l). Flies were anesthetized with N_2 or CO_2 for screening, sexing, and the setup of crosses. To slow down the development during the non-lethal genotyping process (*C. capitata*), adult flies were kept at 19°C, 60% RH, and 24 h light for this period (1-4 d).

    B. tryoni flies were obtained from New South Wales Department of Primary Industries (NSW DPI), Ourimbah, Australia and reared at $25 \pm 2°C$, $65 \pm 10\%$ RH and 14/10 h light/dark cycle. Flies were fed with sugar, Brewer's yeast and water and larvae were reared on a gel diet, containing Brewer's yeast (204 g/l), sugar (121 g/l), methyl p-hydroxy benzoate (2 g/l), citric acid (23 g/l), wheat germ oil (2 g/l), sodium benzoate (2 g/l) and agar (10 g/l).

Introgression and bioinformatic identification of a natural *wp* mutation from *B. dorsalis* in *B. tryoni*. Interspecific crosses between *B. tryoni* and *white pupae B. dorsalis* were carried out between male *B. tryoni* ($wp^{+|+}$) and female *B. dorsalis* ($wp^{-|-}$). The G_1 $wp^{+|-}$ hybrids developed with brown puparia and were mass crossed. G_2 $wp^{-|-}$ females were backcrossed into *B. tryoni* $wp^{+|+}$ males and backcrossing was then repeated for five additional times to produce the *white pupae Bactrocera Introgressed Line* (BIL, Fig. S1).

    Genome sequencing using Illumina NovaSeq (2 x 150 bp, Deakin University) was performed on a single male and female from the *B. dorsalis wp* strain, *B. tryoni*, and the *BIL* (~ 26X) and two pools of five *BIL* individuals (~32X). Quality control of each sequenced library was carried out using FastQC v0.11.6 (https://www.bioinformatics.babraham.ac.uk/projects/fastqc/) and aggregated using ngsReports v1.3 (39). Adapter trimming was carried out using Trimmomatic v0.38 (https://academic.oup.com/bioinformatics/article/30/15/2114/2390096) and paired reads were mapped to the *B. dorsalis* reference genome (GCF_000789215.1) using NextGenMap v0.5.5 (40) under default settings. Mapped data were sorted and indexed using SAMtools, and deduplication was carried out using Picard MarkDuplicates v2.2.4 (https://github.com/broadinstitute/picard). Genotypes were called on single and pooled

libraries separately with ploidy set to two and ten respectively using Freebayes v1.0.2 (41).
Each strain was set as a different population in Freebayes. Genotypes with less than five
genotype depth were set to missing and sites with greater than 20% missing genotypes or
indels filtered out using BCFtools v1.9 (42). Conversion to the genomic data structure (GDS)
format was carried out using SeqArray v1.26.2 (43) and imported into the R package geaR
v0.1 (https://github.com/CMWbio/geaR) for population genetic analysis.

Single copy orthologs were identified in the *B. dorsalis* reference annotated proteins (NCBI
Bactrocera dorsalis Annotation Release 100) with BUSCO v3 (44, 45) using the dipteran gene
set (44). Nucleotide alignments of each complete single copy ortholog were extracted from
the called genotype set using geaR v0.1 and gene trees built using RAxML v8.2.10 (46) with
a GTR+G model. Gene trees were then imported into Astral III v5.1.1 (47) for species tree
estimation. Genome scans of absolute genetic divergence (d_{XY}), nucleotide diversity (π), and
the f estimator $\widehat{f_d}$ were carried out using geaR v0.1. Two levels of analysis were carried out:
i) genome wide scans of non-overlapping 100 kb windows and ii) locus scans of 10 kb tiled
windows. Local phylogenies were built for nucleotide alignments of non-overlapping 1 kb
windows using RAxML v8.2.10 with a GTR+G model and topology weighting was calculated
using TWISST (48).

Introgressed regions (i.e. candidate *wp* loci) were identified by extracting windows in the
genome wide scan with topology weighting and fd greater than 0.75 and visually inspecting
the 'locus scan' data set for d_{XY}, fd and topology weighting patterns indicative of introgression.
Nucleotide alignments of all genes within candidate *B. dorsalis* introgressed regions were
extracted from the GDS using geaR and visually inspected for fixed mutations in *B. dorsalis*
wp, BIL individuals and the two BIL pools. Candidate genes were then searched by tBLASTn
against the *D. melanogaster* annotated protein set to identify putative functions and functional
domains were annotated using HMMer (49). Mapped read depth was calculated around
candidate regions using SAMtools depth v1.9 (50) and each sample's read depth was
normalized to the sample maximum to inspect putative deletions. Called genotypes were
confirmed by *de novo* genome assembly of the *B. dorsalis wp* genome using MaSuRCA v3.3
(51) under default settings. The *de novo* scaffold containing LOC105232189 was identified
using the BLASTn algorithm. *In silico* exon-intron boundaries were then manually annotated
in Geneious v11 (52).

Identification, characterization and molecular analysis of the D53 inversion and *wp*
locus in *C. capitata*.
Characteristics of C. capitata *strains used for this study*
Egypt II (EgII) is a wild type laboratory strain. D53 is a homozygous strain with an irradiation-
induced inversion covering the area 69C-76B on the salivary gland polytene chromosome map

(50B-59C on the trichogen cells polytene chromosome map). VIENNA 7 and VIENNA 8 are

two GSS in which two (Y;5) translocations, on the region 58B and 52B of the trichogen cells

polytene chromosome map respectively, have resulted to the linkage of the wild type allele of

the *wp* and *tsl* genes to the male determining region of the Y chromosome. Thus, VIENNA 7

and VIENNA 8 males are heterozygous in the *wp* and *tsl* loci but phenotypically wild type while

VIENNA 7 and VIENNA 8 females are homozygous for the mutant alleles and phenotypically

white pupae and they die when exposed at elevated temperatures. The VIENNA 7 and

VIENNA 8 GSS can be constructed with and without the D53 inversion (VIENNA 7/8 $^{D53+}$ or $^{D53-}$

). When the GSS have the inversion, females are homozygous ($^{D53+/+}$) for D53 while males are

heterozygous ($^{D53+/-}$) (6, 8, 16).

Whole genome sequencing of C. capitata *strains*

High-molecular-weight (HMW) DNA was extracted from *C. capitata* lines (males and females

of the wild type EgII strain, the VIENNA 7 $^{D53-/-}$ and VIENNA 8 $^{D53-/-}$ GSS and the inversion line

D53) and sequenced. Freshly emerged, virgin and unfed males and females were collected

from all strains. For 10X Genomics linked read and Nanopore sequencing, the HMW was

prepared as follows: twenty individuals of each sex and strain were pooled, ground in liquid

nitrogen, and HMW DNA was extracted using the QIAGEN Genomic tip 100/G kit (Qiagen,

Germany). For PacBio Sequel an EgII line was created with single pair crossing and

subsequent sibling-mating for six generations. In all generations adult and larval diet contained

100 µg/ml tetracycline. HMW DNA from G_6 individuals was prepared as follows: five males

from this EgII line were pooled and ground in liquid nitrogen, and HMW DNA was extracted

using the phenol/chloroform Phase Lock Gel™ tubes (QuantaBio) (53). DNA for Illumina

applications was extracted from individual flies (Table S1).

PacBio *de novo* sequencing: samples were purified with AMPure beads (Beckman Coulter,

UK) (0.6 volumes) and QC checked for concentration, size, integrity and purity using Qubit

(Qiagen, UK), Fragment Analyser (Agilent Technologies) and Nanodrop (Thermo Fisher)

machines. The samples were then processed without shearing using the PacBio Express kit

1 for library construction and an input of 4 µg DNA following the manufacturer's protocol. The

final library was size-selected using the Sage Blue Pippin (Sage Sciences) 0.75% cassette U1

marker in the range of 25-80 kb. The final library size and concentrations were obtained on

the Fragment Analyser before being sequenced using the Sequel 1 2.1 chemistry with V4

primers at a loading on plate concentration of 6 pM and 10 h movie times.

For Nanopore sequencing, the ligation sequencing kits SQK-LSK109 or SQK-RAD004

were used as recommended by the manufacturer (Oxford Nanopore Technologies, Oxford,

United Kingdom). Starting material for the ligation library preparation were 1 - 1.5 µg HMW

gDNA for the ligation libraries and 400 ng for the rapid libraries. The prepared libraries were

loaded onto FLO-PRO002 (R9.4) flow cells. Data collection was carried out using a
PromethION Beta with live high accuracy base calling for up to 72 h and with mux scan
intervals of 1.5 h. Each sample was sequenced at least twice. Data generated were 7.7 Gb
for EgII male, 31.09 Gb for D53 male, 26.72 Gb for VIENNA 7[D53-|-] male, and 24.83 Gb for
VIENNA 8 [D53-|-] male. Run metrics are shown in Supplementary Table S5.

The PacBio data were assembled using CANU (54) with two parameter settings: the first
to avoid haplotype collapsing (genomeSize=500m corOutCoverage=200 "batOptions=-dg 3 -
db 3 -dr 1 -ca 500 -cp 50") and the second to merge haplotypes together (genomeSize=500m
corOutCoverage=200 correctedErrorRate=0.15). The genome completeness was assessed
with BUSCO v3 (44, 45) using the dipteran gene set (44). The two assemblies were found to
be "duplicated" due to alternative haplotypes. To improve the contiguity and reduce duplication
haploMerger2 was used (55). The new assembly was then retested with BUSCO v3 and used
scaffolding. Phase Genomics[TM] Hi-C libraries were made by Phase genomics from males
(n=2) of the same family used for PacBio sequencing. Initial scaffolding was completed by
Phase Genomics, but edited using the Salsa (56) and 3D-DNA
(https://github.com/theaidenlab/3d-dna) software. The resulting scaffolds were allocated a
chromosome number using chromosome specific marker described previously (16). Specific
attention was made to the assembly and scaffolding of chromosome 5. Two contig
misassemblies were detected by the Hi-C data and fitted manually. The new assembly
(EgII_Ccap3.2.1) was then validated using the Hi-C data. Genes were called using the
Funannotate software making use of the Illumina RNAseq data generated by this project;
mRNA mapping to the genome is described below.

D53 breakpoint analysis in C. capitata
To identify possible breakpoint positions, the Nanopore D53 fly assembly contig_531 was
mapped onto the EgII_scaffold_5 (from the EgII_CCAP3.2_CANU_Hi-C_scaffolds.fasta
assembly) using MashMap v2.0 (https://github.com/marbl/MashMap). This helped to visualize
the local alignment boundaries (Fig. S9A). MashMap parameters were set to kmer size = 16;
window size = 100; segment length = 500; alphabet = DNA; percentage identity threshold =
95%; filter mode = one-to-one.

Subsequent to this, and to help confirm the exact location of the identified breakpoints,
minimap2 (v2.17, https://github.com/lh3/minimap2) was used to align D53 as well as VIENNA
8[D53-|-] and VIENNA 7[D53-|-] Nanopore reads onto the EgII scaffold_5 reference (Fig. S9B-C).
Minimap2 parameters for Nanopore reads were: minimap2 -x map-ont -A 1 -a --MD -L -t 40.
Samtools (v1.9, https://github.com/samtools/samtools) was used to convert the
alignment.sam to .bam and prepare the alignment file to be viewed in the Integrative Genomics
Viewer (IGV, http://software.broadinstitute.org/software/igv/). The expectation was to see a

leftmost breakpoint in D53 read set alignments but not in VIENNA 8[D53-|-] and VIENNA 7[D53-|-]

when compared to the EgII reference (Fig.S9B). Due to an assembly gap in the EgII scaffold_5

sequence, the exact location of the leftmost inversion breakpoint was not conclusive using this

approach.

A complementary approach was then used to facilitate detection of the leftmost inversion

breakpoint in the D53 inversion line. Minimap2 was again used, but here D53 contig_531 was

used as reference for the mapping of EgII male PacBio reads as well as VIENNA [8D53-|-] male

and VIENNA 7 D53-|- male Nanopore reads (Fig. S10). Minimap2 parameters for PacBio

reads were: minimap2 -x map-pb -A 1 -a --MD -L -t 40. Minimap2 parameters for Nanopore

reads were: minimap2 -x map-ont -A 1 -a --MD -L -t 40. Samtools (v1.9,

https://github.com/samtools/samtools) was used to convert the alignment .sam to .bam and

prepare the alignment file to be viewed in the Integrative Genomics Viewer (IGV,

http://software.broadinstitute.org/software/igv/). The expectation was to see a common

breakpoint for all three of the above read set alignments when compared to the D53 genome

in the area of the inversion. Position ~3,055,294 was identified in the D53 contig_531 as the

most likely leftmost breakpoint.

To determine the rightmost breakpoint, D53 male and VIENNA 8 [D53-|-] and VIENNA 7[D53-|-]

male nanopore reads were aligned on the EgII_scaffold_5 sequence. The expectation was to

see a breakpoint in D53 read set alignments but not in VIENNA 7[D53-|-] and VIENNA 8 [D53-|-]. This

is the case here, since read alignments coming from both sides of the inversion are truncated

at one position (Fig. S9C). Findings from genome version EgII_Ccap3.2 were extrapolated to

the manually revised genome version EgII_Ccap3.2.1.

Predicted D53 inversion breakpoints were verified via PCRs in EgII, D53, and VIENNA 7[D53+]

GSS male and female flies, using PhusionFlash Polymerase in a 10 µl reaction volume [98°C,

10 s; 30 cycles of (98°C, 1 s; 56°C, 5 s; 72°C, 35 s); 72°C, 1 min] (Fig. S4). The primer pair

for the right breakpoint was designed based on EgII sequence information, primers for the left

breakpoint were designed based on D53 sequence information. The wild type status of

chromosome 5 (EgII male and female, VIENNA 7 [D53+|-] male) was amplified using primer pairs

P_1794 and P_1798 (1,950 bp) and P_1795 and P_1777 (690 bp) (Table S4). Chromosome

5 with the inversion (D53 male and female, VIENNA 7 [D53+|-] male and VIENNA 7 [D53+|+] female)

was verified using primer pairs P_1777 and P_1798 (1,188 bp) and P_1794 and P_1795

(1,152 bp) and amplicon sequencing.

RNA extraction for transcriptomic analysis of C. capitata, B. dorsalis *and* Z. cucurbitae

Samples of *C. capitata*, *B. dorsalis* and *Z. cucurbitae* species were collected for RNA

extraction (Table S1) at 3rd instar larval and pre-pupal stages. Total RNA was extracted by

homogenizing three larvae of *C. capitata* and *B. dorsalis* and a single larvae of *Z. cucurbirtae*

in liquid nitrogen*,* and then using the RNeasy Mini kit (Qiagen). Three replicates per strain and

time point were performed. mRNA was isolated using the NEBNext polyA selection and the

Ultra II directional RNA library preparation protocols from NEB and sequenced on the Illumina

NovaSeq 6000 using duel indexes as 150 bp paired end reads (library insert 500 bp).

Individual libraries were sequenced to provide >1 million paired end reads per sample. Each

replicate was then assembled separately using Trinity (57). The assembled transcripts from

Trinity were mapped to the Ccap3.2 genome using minimap (58) (parameters -ax splice:hq -

uf). The Illumina reads were mapped with STAR (59). IGV v 2.6 (60) was used to view all data

at a genomic and gene level. Given that the white pupae GSS (12, 61) was used to collect

samples for RNA extraction from single larvae of *Z. cucurbitae*, larval sex was confirmed by a

maleness-specific PCR on the *MoY* gene of *Z. cucurbitae* (36) using cDNA synthesized with

the OneStep RT-PCR Kit (Qiagen) and the primer pair ZcMoY1F and ZcMoY1R (Table S4)

amplifying a 214 bp fragment. Conditions for a 25 µl PCR reaction using the 1× *Taq* PCR

Master Mix kit (Qiagen) were: [95°C, 5 min; 30 cycles of (95°C, 1 min; 51°C, 1 min; 72°C, 1

min); 72°C, 10 min]. Presence of a PCR product indicated a male sample. Each, male and

female sample was a pool of three individuals. Three replicates per strain and time point were

collected.

Functional and cytogenetic verification of medfly D53 inversion and tephritid *wp* genes.

Polytene chromosomes for *in situ* hybridization were prepared from third-instar larvae salivary

glands as described previously (62). In brief, the glands were dissected in 45% acetic acid

and placed on a coverslip in a drop of 3:2:1 solution (3 parts glacial acetic acid: 2 parts water:

1-part lactic acid) until been transparent (approximately 5 min). The coverslip was picked up

with a clean slide. After squashing, the quality of the preparation was checked by phase

contrast microscope. Satisfactory preparations were left to flatten overnight at -20°C and

dipped into liquid nitrogen until the babbling stopped. The coverslip was immediately removed

with razor blade and the slides were dehydrated in absolute ethanol, air dried and kept at room

temperature.

Probes were prepared by PCR. Single adult flies were used to extract DNA with the Extract

me kit (Blirt SA), following the manufacturer's protocol. NanoDrop spectrometer was used to

assess the quantity and quality of the extracted DNA which was then stored at -20°C until

used. Primers (P1790/1791, P1821/1822, Pgd_probe_F/R, vg1_probe_F/R, Sxl_probe_F/R,

y_probe_F/R, zw_probe_F/R, P1633/1634, Zc_F/R, Bd_F/R, P1395/1396, P1415/1416;

Table S4) were designed for each targeted gene using Primer 3 and/or Geneious Prime

programs. PCR was performed in a 25 µl reaction volume using 12.5 µl PCR Master mix 2x

Kit (Thermo Fisher Scientific), 60-80 ng DNA, and the following PCR settings [94°C, 5 min; 35

cycles of (94°C, 45 s; 56°C, 30 s; 72°C, 90 s); 72°C, 1 min].

The labelling of the probes was carried out according to the instruction manual of the Dig DNA labelling kit (Roche). *In situ* hybridization was performed as described previously (63). In brief, before hybridization, stored chromosome preparations were hydrated by placing them for 2 min at each of the following ethanol solutions: 70%, 50% and 30%. Then they were placed in 2× SSC at room temperature for 2 min. The stabilization of the chromosomes was done by placing them in 2× SSC at 65°C for 30 min, denaturing in 0.07 M NaOH 2 min, washing in 2× SSC for 30 sec, dehydrating (2 min in 30%, 50%, 70% and 95% ethanol), and air drying. Hybridization was performed on the same day by adding 15 µl of denatured probe (boiled for 10 min and ice-chilled). Slides were covered with a siliconized coverslip, sealed with rubber cement, and incubated at 45°C overnight in a humid box. At the end of incubation, the coverslip was floated off in 2x SSC and the slide washed in 2x SSC for 3 x 20 min at 53°C.

After 5 min wash in Buffer 1 (100 mM tris-HCl pH 7.5/ 1.5 M NaCl), the preparations were in Blocking solution (Blocking reagent 0.5% in Buffer 1) for 30 min, and then washed for 1 min in Buffer 1. The antibody mix was added to each slide and a coverslip was added. Then the slides were incubated in a humid box for 45 min at room temperature, following 2× 15 min washes in Buffer 1, and a 2 min wash in detection buffer (100 mM Tris-HCl pH 9.5/ 100 mM NaCl). The color was developed with 1 ml of NBT/BCIP solution during a 40 min incubation in the dark at room temperature. The removal of the NBT/BCIP solution was done by rinsing in water twice. Hybridization sites were identified and photographed using a 40× objective or 100× oil objectives (phase or bright field), with reference to the salivary gland chromosome maps (64).

Gene editing and generation of homozygous *wp⁻* strains in *B. tryoni* and *C. capitata*. For CRISPR/Cas9 gene editing in *B. tryoni,* purified Cas9 protein (Alt-R® S.p. Cas9 Nuclease V3, #1081058, 10 µg/µl) and guide RNAs (customized Alt-R® CRISPR/Cas9 crRNA, 2 nmol and Alt-R® CRISPR/Cas9 tracrRNA, #1072532, 5nmol) were obtained from Integrated DNA Technologies (IDT) and stored at -20°C. The guide RNAs were individually resuspended to a 100 µM stock solution with nuclease-free duplex buffer and stored at -20°C before use. The two customized 20 bp crRNA sequences (Bt_MFS-1 and Bt_MFS-2) (Table S4) were designed using CRISPOR (65). Injection mixes for microinjection of *B. tryoni* embryos comprise of 300 ng/µl Cas9 protein, 59 ng/µl of each individual crRNA, 222 ng/µl tracrRNA and 1x injection buffer (0.1 mM sodium phosphate buffer pH 6.8, 5 mM KCl) in a final volume of 10 µl. The guide RNA complex containing the two crRNAs and tracrRNA was initially prepared by heating the mix at 95°C for 5 min before cooling to room temperature. The Cas9 enzyme along with the rest of the injection mix components were then added to the guide RNA complex and incubated at room temperature for 5 minutes to assemble the RNP complexes. Microinjections were performed in *B. tryoni* Ourimbah laboratory strain embryos that were

collected over 1 hour time period and prepared for injection as previously described (22).
Injections were performed under paraffin oil using borosilicate capillary needles (#30-0038,
Harvard Apparatus) drawn out on a Sutter P-87 flaming/brown micropipette puller and
connected to an air-filled 20 ml syringe, a manual MM-3 micromanipulator (Narishige) and a
CKX31-inverted microscope (Olympus). Microscope slides with the injected embryos were
placed on agar in a Petri dish that was then placed in a vented container containing moist
paper towels at 25°C (± 2°C). Hatched first instar larvae were removed from the oil and
transferred to larval food. Individual G_0 flies were crossed to six virgin flies from the Ourimbah
laboratory strain and eggs were collected overnight for two consecutive weeks. G_1 flies were
then allowed to mate *inter se* and eggs were collected in the same manner. G_2 pupae were
then analyzed phenotypically and separated according to color of pupae (brown, mosaic or
white).

For *C. capitata* CRISPR/Cas9 gene editing, lyophilized Cas9 protein (PNA Bio Inc, CP01)
was reconstituted to a stock concentration of 1 µg/µl in 20 mM Hepes, 150 mM KCl, 2%
sucrose and 1 mM DTT (pH 7.5) and stored at -80°C until use. A guide RNA (gRNA_MFS),
targeting the third CDS exon of *CcMFS* was designed and tested for potential off target effects
using Geneious Prime (52) and the *C. capitata* genome annotation Ccap2.1. *In silico* target
site analysis predicted an on-target activity score of 0.615 (scores are between 0 and 1; the
higher the score the higher the expected activity (66) and zero off-targets sites in the medfly
genome. Guide RNA was synthesized by *in vitro* transcription of linear double-stranded DNA
template as previously described (67) using primers P_1753 and P_369 (Table S4), Q5 HF
polymerase, and a Bio-Rad C1000 Touch thermal cycler [98°C, 30 s; 35 cycles of (98°C, 10
s; 58°C, 20 s; 72°C, 20 s); 72°C, 2 min] and HiScribe™ T7 High Yield RNA Synthesis. Injection
mixes for microinjection of embryos contained 360 ng/µl Cas9 protein (1 µg/µl, dissolved in its
formulation buffer), 200 ng/µl gRNA_MFS and an end-concentration of 300 mM KCl according
to previous studies (67, 68). The mix was freshly prepared on ice followed by an incubation
step for 10 min at 37°C to allow pre-assembly of gRNA-Cas9 ribonucleoprotein complexes
and stored on ice prior to injections. Microinjections were conducted in wild type EgII *C.*
capitata embryos. Eggs were collected over a 30-40 min time period and prepared for injection
as previously described (67). Injections were performed using siliconized quartz glass needles
(Q100-70-7.5; LOT171381; Science Products, Germany), drawn out on a Sutter P-2000 laser-
based micropipette puller. The injection station consisted of a manual micromanipulator (MN-
151, Narishige), an Eppendorf FemtoJet 4i microinjector, and an Olympus SZX2-TTR
microscope (SDF PLAPO 1xPF objective). The microscope slide with the injected embryos
was placed in a Petri dish containing moist tissue paper in an oxygen chamber (max. 2 psi).
Hatched first instar larvae were transferred from the oil to larval food.

As complementation assay, reciprocal crosses between surviving G_0 adults and virgin *white*

pupae strain #1402_22m1B (pBac_*fa_attP-TREhs43-Cctra-I-hid^{Ala5}-SV40_a_PUb-nls-EGFP-*

SV40) ($wp^{-(nat)}$) (24) were set up either single paired (six cages) or in groups of seven to ten

flies (seven cages). Eggs were collected three times every 1-2 days. Progeny (G_1) exhibiting

the *white pupae* phenotype ($wp^{-(nat)|-(CRISPR)}$) were assayed via non-lethal genotyping and

sorted according to mutation genotype (see Fig. 4). Genotypes 'A-H' were group-backcrossed

into WT EgII ($wp^{+|+}$), genotype 'C' siblings massed crossed. Eggs were collected 4 times every

1-2 days. Generation G_2 flies were analyzed via PCR using three primers, specific for wp^+ and

$wp^{-(CRISPR)}$ or $wp^{-(nat)}$ allele size respectively. Offspring of outcross cages showed brown pupae

phenotype and either $wp^{+|-(nat)}$ or $wp^{+|-(CRISPR)}$ genotype. In order to make mutations A, D, and

H homozygous, 40 flies (25 females, 15 males) were genotyped each, and $wp^{+|-(CRISPR)}$ positive

flies were inbred (mutation A: 15 females, 7 males, mutation D: 12 females, 7 males, mutation

H: 11 females, 8 males). G_3 showing white pupae phenotype was homozygous for $wp^{-(CRISPR)}$

mutations A, D, or H, respectively and was used to establish lines. Inbreeding of mutation C

$wp^{-(nat)|-(CRISPR)}$ flies produced only white pupae offspring, based on either $wp^{-(nat)|-(nat)}$, $wp^{-(nat)|-}$

$^{(CRISPR)}$ or $wp^{-(CRISPR)|-(CRISPR)}$. 94 flies (46 females, 48 males) were genotyped, homozygous wp^-

$^{(CRISPR)}$ were inbred to establish a line (13 females, 8 males).

Molecular analyses of *wp* mutants and mosaics. In *B. tryoni,* genomic DNA was isolated

for genotyping from G_2 pupae using the DNeasy Blood and Tissue Kit (Qiagen). PCR

amplicons spanning both BtMFS guide recognition sites were generated using Q5 polymerase

(NEB) with primers BtMFS_5primeF and BtMFS_exon2R (Table S4). Products were purified

using MinElute PCR Purification Kit (Qiagen), ligated into pGEM-t-easy vector (Promega) and

transformed into DH5α cells. Plasmids were purified with Wizard *Plus* SV Minipreps

(Promega) and sequenced.

In *C. capitata,* non-lethal genotyping was performed to identify parental genotypes before

setting up crosses. Therefore, genomic DNA was extracted from single legs of G_1 and G_2 flies

following a protocol established by Carvalho et al. (69). Single legs of anesthetized flies were

cut at the proximal femur using scissors. Legs were then homogenized by ceramic beads and

50 µl buffer (10 mM Tris-Cl, pH 8.2, 1 mM EDTA, 25 mM NaCl) for 15 s (6 m/s) using a

FastPrep-24^{TM} 5G homogenizer. Then, 30 µl buffer and 1.7 µl proteinase-K (2.5 U/mg) were

added, incubated for 1 h at 37°C and the reaction stopped by 4 min at 98°C. The reaction mix

was cooled down on ice and used for PCR. For G_1 flies, PCR on *wp* was performed in a 25 µl

reaction volume using the Dream*Taq* polymerase, primer pair P_1643 and P_1644 (Table

S4), and 3.75 µl reaction mix. Different amplicon sizes are expected for brown (wp^+ and

$wp^{-(CRISPR)}$) and white pupae ($wp^{-(nat)}$) alleles, 724 bp and 8,872 bp in size, respectively. The

$wp^{-(nat)}$ amplicon was excluded via PCR settings [95°C, 3 min; 35 cycles of (95°C, 30 s; 56°C,

30 s; 72°C, 1 min); 72°C, 5 min]. The 724 bp PCR product was verified by agarose

electrophoresis and purified from the PCR reaction using the DNA Clean & Concentrator™-5

kit. PCR products were sequenced using primer P_1644 and sequences analyzed using the

Geneious Prime Software Package (6). In generation G_2, flies were analyzed using multiplex

PCR with primers P_1657, P_1643, and P_1644 (Table S4), to distinguish between the wp^{-}

$^{(nat)}$ (457 bp; P_1643 and P_1657), and $wp^{-(CRISPR)}$ alleles (724 bp; P_1643 and P1644) using

the PCR protocol [95°C, 3 min; 35 cycles of (95°C, 30 s; 56°C, 30 s; 72°C, 1 min); 72°C, 5

min]).

Image acquisition. Images of *B. tryoni* pupae were taken with an Olympus SZXI6

microscope, Olympus DP74 camera and Olympus LF-PS2 light source using the Olympus

stream basic 2.3.3 software. Images of *C. capitata* pupae were taken with a Keyence digital

microscope VHX-5000. Image processing was conducted with Adobe Photoshop CS5.1

software to apply moderate changes to image brightness and contrast. Changes were applied

equally across the entire image and throughout all images.

Ethics approval and consent to participate

Not applicable

Consent for publication

Not applicable

Availability of data and material

All data generated or analyzed are included in this article.

Declaration of interest

none

Funding

This study was financially supported by the Joint FAO/IAEA Insect Pest Control Subprogramme of the Joint FAO/IAEA Division of Nuclear Techniques in Food and Agriculture. The project has also been funded by the Horticulture Innovation Australia (FF17000, FF18002), using funds from the Australian Government, and co-investment from Macquarie University and South Australian Research and Development Institute (SARDI). SWB was supported by the Australian Research Council (FT140101303) and Hermon Slade Foundation grant HSF 18/6. Furthermore, this work was supported by the Emmy Noether program of the German Research Foundation (SCHE 1833/1-1; to MFS) and the LOEWE Center for Insect Biotechnology and Bioresources of the Hessen State Ministry for Higher Education, Research and the Arts (HMWK; to MFS). The work was also supported by Canadian Foundation for Innovation and Genome Canada Genome Technology Platform awards and the International Atomic Energy Agency research contact no. 23358 as part of the Coordinated Research Project "Generic approach for the development of genetic sexing strains for SIT applications" (JR).

Acknowledgements

This study was benefitted from discussions at International Atomic Energy Agency funded meetings for the Coordinated Research Projects (CRPs) and particularly the CRP on "Generic approach for the development of genetic sexing strains for SIT applications". The authors also wish to thank Tanja Rehling, Jakob Martin and Johanna Rühl for technical assistance and Germano Sollazzo for helping with injections and primers design (Justus-Liebig University Gießen and Insect Pest Control Laboratory); Elena Isabel Cancio Martinez, Thilakasiri Dammalage, Sohel Ahmad and Gülizar Pillwax for insect rearing (Insect Pest Control Laboratory), Shu-Huang Chen (McGill University) for technical assistance with Nanopore library preparations; and Arjen van´t Hof (University of Liverpool) for constructing libraries from micro-dissected chromosomes for Illumina sequencing.

References

1. A. S. Robinson, G. Hooper, *Fruit flies: their biology, natural enemies, and control*
(Elsevier, Amsterdam; New York, 1989), vol. 1, pp. 372.
2. D. M. Suckling *et al.*, Eradication of tephritid fruit fly pest populations: outcomes and
prospects. *Pest Manag Sci* **72**, 456-465 (2016).
3. V. A. Dyck *et al.*, "Management of area-wide integrated pest management
programmes that integrate the sterile insect technique" in *Sterile insect technique -*
principles and practice in area-wide integrated pest management, V. A. Dyck, J.
Hendrichs, A. S. Robinson, Eds. (Springer, Dordrecht, NL, 2005), pp. 525-545.
4. M. Vreysen, A. S. Robinson, J. Hendrichs, *Area-wide control of insect pests: from*
research to field implementation. M. Vreysen, A. S. Robinson, J. Hendrichs, Eds.
(Springer, Dordrecht, NL, 2007).
5. P. Rendon, D. McInnis, D. Lance, J. Stewart, Medfly (Diptera: Tephritidae) genetic
sexing: large-scale field comparison of males-only and bisexual sterile fly releases in
Guatemala. *Journal of Economic Entomology* **97**, 1547-1553 (2004).
6. G. Franz, "Genetic sexing strains in Mediterranean fruit fly, an example for other
species amenable to large-scale rearing for the sterile insect technique" in *Sterile*
insect technique - principles and practice in area-wide integrated pest management,
V. A. Dyck, J. Hendrichs, A. S. Robinson, Eds. (Springer, Dordrecht, NL, 2005), pp.
427-451.
7. A. A. Augustinos *et al.*, *Ceratitis capitata* genetic sexing strains: laboratory evaluation
of strains from mass-rearing facilities worldwide. *Entomologia Experimentalis et*
Applicata **164**, 305-317 (2017).
8. A. Zacharopoulou *et al.*, A review of more than 30 years of cytogenetic studies of
Tephritidae in support of sterile insect technique and global trade. *Entomologia*
Experimentalis et Applicata **164**, 204-225 (2017).
9. Y. Rössler, The genetics of the Mediterranean fruit fly: a "white pupae" Mutant.
Annals of the Entomological Society of America **72**, 583-585 (1979).
10. Y. Rössler, Y. Koltin, The genetics of the Mediterranean fruit fly, *Ceratitis capitata*:
three morphological mutations. *Annals of the Entomological Society of America* **69**,
604-608 (1976).
11. S. D. McCombs, S. H. Saul, Linkage analysis of five new genetic markers of the
oriental fruit fly, *Bactrocera dorsalis* (Diptera: Tephritidae). *The Journal of heredity*
83, 199-203 (1992).
12. D. O. McInnis *et al.*, Development of a pupal color-based genetic sexing strain of the
melon fly, *Bactrocera cucurbitae* (Coquillett) (Diptera: Tephritidae). *Annals of the*
Entomological Society of America **97**, 1026-1033 (2004).
13. P. Wappner *et al.*, *White pupa: A Ceratitis capitata mutant lacking catecholamines for*
tanning the puparium (1995), vol. 25, pp. 365-373.
14. Y. Rössler, H. Rosenthal, Genetics of the Mediterranean fruit fly (Diptera:
Tephritidae): morphological mutants on chromosome five. *Annals of the*
Entomological Society of America **85**, 525-531 (1992).
15. P. Kerremans, G. Franz, Cytogenetic analysis of chromosome 5 from the
Mediterranean fruit fly, *Ceratitis capitata*. *Chromosoma* **103**, 142-146 (1994).
16. A. Papanicolaou *et al.*, The whole genome sequence of the Mediterranean fruit fly,
Ceratitis capitata (Wiedemann), reveals insights into the biology and adaptive
evolution of a highly invasive pest species. *Genome Biology* **17**, 192 (2016).
17. P. A. Papathanos *et al.*, A perspective on the need and current status of efficient sex
separation methods for mosquito genetic control. *Parasites & Vectors* **11**, 654 (2018).
18. S. B. Sim, S. M. Geib, A chromosome-scale assembly of the *Bactrocera cucurbitae*
genome provides insight to the genetic basis of *white pupae*. *G3 (Bethesda, Md.)* **7**,
1927-1940 (2017).

19.   S. B. Sim, R. Ruiz-Arce, N. B. Barr, S. M. Geib, A new diagnostic resource for
Ceratitis capitata strain identification based on QTL mapping. *G3 (Bethesda, Md.)* **7**,
3637-3647 (2017).
20.   E. M. Zdobnov *et al.*, OrthoDB v9.1: cataloging evolutionary and functional
annotations for animal, fungal, plant, archaeal, bacterial and viral orthologs. *Nucleic*
Acids Res **45**, D744-D749 (2017).
21.   M. San Jose *et al.*, Incongruence between molecules and morphology: A seven-gene
phylogeny of *Dacini* fruit flies paves the way for reclassification (Diptera: Tephritidae).
Mol Phylogenet Evol **121**, 139-149 (2018).
22.   A. Choo, P. Crisp, R. Saint, L. V. O'Keefe, S. W. Baxter, CRISPR/Cas9-mediated
mutagenesis of the white gene in the tephritid pest *Bactrocera tryoni*. *Journal of*
Applied Entomology **142**, 52-58 (2018).
23.   A. Gariou-Papalexiou *et al.*, Polytene chromosomes as tools in the genetic analysis
of the Mediterranean fruit fly, *Ceratitis capitata*. *Genetica* **116**, 59-71 (2002).
24.   C. E. Ogaugwu, M. F. Schetelig, E. A. Wimmer, Transgenic sexing system for
Ceratitis capitata (Diptera: Tephritidae) based on female-specific embryonic lethality.
Insect Biochemistry and Molecular Biology **43**, 1-8 (2013).
25.   A. W. Davis *et al.*, Rescue of hybrid sterility in crosses between *D. melanogaster* and
D. simulans. *Nature* **380**, 157-159 (1996).
26.   L. O. Araripe, H. Montenegro, B. Lemos, D. L. Hartl, Fine-scale genetic mapping of a
hybrid sterility factor between *Drosophila simulans* and *D. mauritiana*: the varied and
elusive functions of "speciation genes". *BMC Evol Biol* **10**, 385 (2010).
27.   N. J. Brideau, D. A. Barbash, Functional conservation of the Drosophila hybrid
incompatibility gene Lhr. *BMC Evol Biol* **11**, 57 (2011).
28.   A. A. Kotov *et al.*, piRNA silencing contributes to interspecies hybrid sterility and
reproductive isolation in *Drosophila melanogaster*. *Nucleic Acids Res* **47**, 4255-4271
(2019).
29.   D. A. Barbash, Ninety years of *Drosophila melanogaster* hybrids. *Genetics* **186**, 1-8
(2010).
30.   D. G. Bedo, A. Zacharopoulou, Inter-tissue variability of polytene chromosome
banding patterns. *Trends Genet* **4**, 90-91 (1988).
31.   E. V. Kriventseva *et al.*, OrthoDB v10: sampling the diversity of animal, plant, fungal,
protist, bacterial and viral genomes for evolutionary and functional annotations of
orthologs. *Nucleic Acids Res* **47**, D807-d811 (2019).
32.   Y. Zhao *et al.*, A major facilitator superfamily protein participates in the reddish brown
pigmentation in *Bombyx mori*. *J Insect Physiol* **58**, 1397-1405 (2012).
33.   The modEncode Consortium *et al.*, Identification of functional elements and
regulatory circuits by Drosophila modENCODE. *Science (New York, N.Y.)* **330**, 1787-
1797 (2010).
34.   T. R. Wright, The genetics of biogenic amine metabolism, sclerotization, and
melanization in *Drosophila melanogaster*. *Adv Genet* **24**, 127-222 (1987).
35.   K. Bourtzis, C. Psachoulia, V. J. Marmaras, Evidence that different integumental
phosphatases exist during development in the Mediterranean fruit fly *Ceratitis*
capitata: possible involvement in pupariation. *Comparative Biochemistry and*
Physiology Part B: Comparative Biochemistry **98**, 411-416 (1991).
36.   A. Meccariello *et al.*, *Maleness-on-the-Y* (*MoY*) orchestrates male sex determination
in major agricultural fruit fly pests. *Science* **365**, 1457-1460 (2019).
37.   A. B. Hall *et al.*, Sex determination. A male-determining factor in the mosquito *Aedes*
aegypti. *Science* **348**, 1268-1270 (2015).
38.   P. Liu *et al.*, *Nix* is a male-determining factor in the Asian tiger mosquito *Aedes*
albopictus. *Insect Biochem Mol Biol* **118**, 103311 (2019).
39.   C. M. Ward, T. H. To, S. M. Pederson, ngsReports: a Bioconductor package for
managing FastQC reports and other NGS related log files. *Bioinformatics* **36**, 2587-
2588 (2020).

40.   F. J. Sedlazeck, P. Rescheneder, A. von Haeseler, NextGenMap: fast and accurate
917   read mapping in highly polymorphic genomes. *Bioinformatics* **29**, 2790-2791 (2013).
918   41.   E. Garrison, G. Marth, Haplotype-based variant detection from short-read
919   sequencing. arXiv:1207.3907.
920   42.   H. Li, A statistical framework for SNP calling, mutation discovery, association
921   mapping and population genetical parameter estimation from sequencing data.
922   *Bioinformatics* **27**, 2987-2993 (2011).
923   43.   X. Zheng *et al.*, SeqArray-a storage-efficient high-performance data format for WGS
924   variant calls. *Bioinformatics* **33**, 2251-2257 (2017).
925   44.   F. A. Simão, R. M. Waterhouse, P. Ioannidis, E. V. Kriventseva, E. M. Zdobnov,
926   BUSCO: assessing genome assembly and annotation completeness with single-copy
927   orthologs. *Bioinformatics* **31**, 3210-3212 (2015).
928   45.   R. M. Waterhouse *et al.*, BUSCO applications from quality assessments to gene
929   prediction and phylogenomics. *Mol Biol Evol* **35**, 543-548 (2018).
930   46.   A. Stamatakis, RAxML version 8: a tool for phylogenetic analysis and post-analysis of
931   large phylogenies. *Bioinformatics* **30**, 1312-1313 (2014).
932   47.   C. Zhang, M. Rabiee, E. Sayyari, S. Mirarab, ASTRAL-III: polynomial time species
933   tree reconstruction from partially resolved gene trees. *BMC Bioinformatics* **19**, 153
934   (2018).
935   48.   S. H. Martin, S. M. Van Belleghem, Exploring evolutionary relationships across the
936   genome using topology weighting. *Genetics* **206**, 429-438 (2017).
937   49.   R. D. Finn, J. Clements, S. R. Eddy, HMMER web server: interactive sequence
938   similarity searching. *Nucleic Acids Res* **39**, W29-37 (2011).
939   50.   H. Li *et al.*, The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
940   2078-2079 (2009).
941   51.   A. V. Zimin *et al.*, The MaSuRCA genome assembler. *Bioinformatics* **29**, 2669-2677
942   (2013).
943   52.   M. Kearse *et al.*, Geneious Basic: an integrated and extendable desktop software
944   platform for the organization and analysis of sequence data. *Bioinformatics* **28**, 1647-
945   1649 (2012).
946   53.   M. R. Green, J. Sambrook, Isolation of High-Molecular-Weight DNA using organic
947   solvents. *Cold Spring Harb Protoc* **2017**, pdb.prot093450 (2017).
948   54.   S. Koren *et al.*, Canu: scalable and accurate long-read assembly via adaptive k-mer
949   weighting and repeat separation. *Genome Res* **27**, 722-736 (2017).
950   55.   S. Huang *et al.*, HaploMerger: reconstructing allelic relationships for polymorphic
951   diploid genome assemblies. *Genome Res* **22**, 1581-1588 (2012).
952   56.   J. Ghurye, M. Pop, S. Koren, D. Bickhart, C. S. Chin, Scaffolding of long read
953   assemblies using long range contact information. *BMC Genomics* **18**, 527 (2017).
954   57.   B. J. Haas *et al.*, De novo transcript sequence reconstruction from RNA-seq using
955   the Trinity platform for reference generation and analysis. *Nat Protoc* **8**, 1494-1512
956   (2013).
957   58.   H. Li, Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics* **34**,
958   3094-3100 (2018).
959   59.   A. Dobin *et al.*, STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15-21
960   (2013).
961   60.   J. T. Robinson *et al.*, Integrative genomics viewer. *Nature biotechnology* **29**, 24-26
962   (2011).
963   61.   A. Zacharopoulou, G. Franz, Genetic and cytogenetic characterization of genetic
964   sexing strains of *Bactrocera dorsalis* and *Bactrocera cucurbitae* (Diptera:
965   Tephritidae). *J Econ Entomol* **106**, 995-1003 (2013).
966   62.   A. Zacharopoulou *et al.*, The genome of the Mediterranean fruit fly *Ceratitis capitata*:
967   localization of molecular markers by *in situ* hybridization to salivary gland polytene
968   chromosomes. *Chromosoma* **101**, 448-455 (1992).
969   63.   P. Mavragani-Tsipidou *et al.*, "Protocols for cytogenetic mapping of arthropod
970   genomes: Tephritid fruit flies of economic importance" in *Protocols for cytogenetic*

mapping of arthropod genomes, I. Sakharov, Ed. (CRC Press, Taylor and Francis
Group, LLC, Florida, USA 2014), pp. 1-62.
64.   A. Zacharopoulou, Polytene chromosome maps in the medfly *Ceratitis capitata*
Genome **33**, 184–197 (1990).
65.   J. P. Concordet, M. Haeussler, CRISPOR: intuitive guide selection for CRISPR/Cas9
genome editing experiments and screens. *Nucleic Acids Res* **46**, W242-W245
(2018).
66.   J. G. Doench *et al.*, Rational design of highly active sgRNAs for CRISPR-Cas9-
mediated gene inactivation. *Nature biotechnology* **32**, 1262-1267 (2014).
67.   R. A. Aumann, M. F. Schetelig, I. Häcker, Highly efficient genome editing by
homology-directed repair using Cas9 protein in *Ceratitis capitata*. *Insect Biochemistry*
and Molecular Biology **101**, 85-93 (2018).
68.   A. Burger *et al.*, Maximizing mutagenesis with solubilized CRISPR-Cas9
ribonucleoprotein complexes. *Development (Cambridge, England)* **143**, 2025-2037
(2016).
69.   G. B. Carvalho, W. W. Ja, S. Benzer, Non-lethal PCR genotyping of single
Drosophila. *Biotechniques* **46**, 312-314 (2009).