SARS-CoV-2 Spike Glycoprotein and ACE2 interaction reveals modulation of viral

entry in different hosts

Manas Ranjan Praharaj[1], Priyanka Garg[1], Raja Ishaq Nabi Khan[2], Shailesh Sharma[1],

Manjit Panigrahi[2], B P Mishra[2], Bina Mishra[2], G Sai kumar[2], Ravi Kumar Gandham[1*],

Raj Kumar Singh[2*], Subeer Majumdar[1*] and Trilochan Mohapatra[3]

[1] National Institute of Animal Biotechnology, Hyderabad, Telangana, India

[2] ICAR-Indian Veterinary Research Institute, Izatnagar, UP, India

[3] Indian Council of Agricultural Research, New Delhi, India

[*] Corresponding authors: Ravi Kumar Gandham: ravigandham@niab.org.in; R.K Singh:

rks_virology@rediffmail.com; Subeer Majumdar: director@niab.org.in

Abstract

Background

SARS-CoV-2 is a viral pathogen causing life threatening disease in human. Interaction

between spike protein of SARS-CoV-2 and ACE2 receptor on the cells is a potential

factor in the infectivity of a host. The interaction of SARS-CoV-2 spike receptor-binding

domain with its receptor - ACE2, in different hosts was evaluated to understand and

predict viral entry. The protein and nucleotide sequences of ACE2 were initially

compared across different species to identify key differences among them. The ACE2

receptor of various species was homology modeled (6LZG as a reference) and its

binding ability to the spike receptor-binding domain of SARS-CoV-2 was assessed.

Initially, the spike binding properties of ACE2 of known infected and uninfected species

was compared with each Order (of animals) as a group. Finally, a logistic regression

model vis-a-vis the spike binding properties of ACE2 (Interaction energy, entropy sidechain and entropy mainchain) was constructed to predict the probability of viral entry in different hosts.

Results

Phylogeny and alignment comparison did not lead to any meaningful conclusion on viral entry in different hosts. Out of several spike binding properties of ACE2, a significant difference between the known infected and uninfected species was observed for five spike binding properties of ACE2. However, these parameters did not specifically categorize the animals into infected or uninfected, for all the Orders (of animals). The logistic regression model constructed revealed that in the mammalian class, most of the species of Carnivores, Artiodactyls, Perissodactyls, Pholidota, and Primates had high probability of viral entry. However, among the primates, baboons have very low probability of viral entry. Among rodents, hamsters were highly probable for viral entry with rats and mice having a very low probability. Rabbits have a medium probability of viral entry. In Birds, ducks have a very low probability, while chickens seemed to have medium probability and turkey showed the highest probability of viral entry.

Conclusions

Most of the species considered in this study showed high probability of viral entry. This study would prompt us to closely follow certain species of animals for determining pathogenic insult by SARS-CoV-2 and for determining their ability to act as a carrier and/or disseminator.

Keywords: SARS-CoV-2; COVID-19; Livestock; ACE2; modeling

Background

Three large-scale disease outbreaks during the past two decades, *viz.*, Severe Acute Respiratory Syndrome (SARS), Middle East Respiratory Syndrome (MERS), and Swine Acute Diarrhea Syndrome (SADS) were caused by three zoonotic coronaviruses. SARS and MERS, which emerged in 2003 and 2012, respectively, caused a worldwide pandemic claiming 774 (8,000 SARS cases) and 866 (2,519 MERS cases) human lives, respectively[1], while SADS devastated livestock production by causing fatal diseases in pigs in 2017. The SARS and MERS viruses had several common factors in having originated from bats in China and being pathogenic to human or livestock[2-4]. Seventeen years after the first highly pathogenic human coronavirus, SARS-COV-2 is devastating the world with 4,907,387 cases and 320,397 deaths (as on May 19, 2020)[5]. This outbreak was first identified in Wuhan City, Hubei Province, China, in December 2019 and notified by WHO on 5[th] January 2020. The disease has since been named as COVID-19 by WHO.

Coronaviruses (CoVs) are an enveloped, crown-like viral particles belonging to the subfamily Orthocoronavirinae in the family Coronaviridae and the order Nidovirales. They harbor a positive-sense, single-strand RNA (+ssRNA) genome of 27–32 kb in size. Two large overlapping polyproteins, ORF1a and ORF1b, that are processed into the viral polymerase (RdRp) and other nonstructural proteins involved in RNA synthesis or host response modulation, cover two thirds of the genome. The rest 1/3 of the genome encodes for four structural proteins (spike (S), envelope (E), membrane (M), and nucleocapsid (N)) and other accessory proteins. The four structural proteins and the ORF1a/ORF1b are relatively consistent among the CoVs, however, number and size of

accessory proteins govern the length of the CoV genome[4]. This genome expansion is

said to have facilitated acquisition of genes that encode accessory proteins, which are

beneficial for CoVs to adapt to a specific host[6, 7]. Next generation sequencing has

increased the detection and identification of new CoV species resulting in expansion of

CoV subfamily. Currently, there are four genera (α-, β-, δ-, and γ-) with thirty-eight

unique species in CoV subfamily (ICTV classification) including the three highly

pathogenic CoVs, *viz.*, SARS-CoV-1, MERS-CoV, SARS-CoV-2 are β-CoVs[8].

Coronaviruses are notoriously promiscuous. Bats host thousands of these types,

without succumbing to illness. The CoVs are known to infect mammals and birds,

including dogs, chickens, cattle, pigs, cats, pangolins, and bats. These viruses have the

potential to leap to new species and in this process mutate along the way to adapt to

their new host(s). COVID -19, global crisis likely started with CoV infected horseshoe

bat in China. The SARS-CoV-2 is spreading around the world in the hunt of entirely new

reservoir hosts for re-infecting people in the future[9]. Recent reports of COVID-19 in a

Pomeranian dog and a German shepherd in Hong Kong[10]; in a domestic cat in

Belgium[11]; in five Malayan tigers and three lions at the Bronx Zoo in New York

City[12] and in minks[13] make it all the more necessary to predict species that could be

the most likely potential reservoir hosts in times to come.

Angiotensin-converting enzyme 2 (ACE2), an enzyme that physiologically

counters RAAS activation functions as a receptor for both the SARS viruses (SARS-

CoV-1 and SARS-CoV-2)[14-16]. The ACE2 human RefSeqGene is 48037 bp in length

with18 exons and is located on chromosome X. ACE2 is found attached to the outer

surface of cells in the lungs, arteries, heart, kidney, and intestines[17, 18]. The potential

factor in the infectivity of a cell is the interaction between SARS viruses and the ACE2 receptor[19, 20]. By comparing the ACE2 sequence, several species that might be infected with SARS-CoV2 have been identified[21]. Recent studies, exposing cells/animals to the SARS-CoV2, revealed humans, horseshoe bats, civets, ferrets, cats and pigs could be infected with the virus and mice, dogs, pigs, chickens, and ducks could not be or poorly infected[16, 22]. Pigs, chickens, fruit bats, and ferrets are being exposed to SARS-CoV2 at Friedrich-Loeffler Institute and initial results suggest that Egyptian fruit bats and ferrets are susceptible, whereas pigs and chickens are not[23]. In this cause of predicting potential hosts, no studies on ACE2 sequence comparison among species along with homology modeling and prediction, to define its interaction with the spike protein of SARS-CoV-2 are available. Therefore, the present study is taken to identify viral entry in potential hosts through sequence comparison, homology modeling and prediction.

Results and Discussion

Recognition of the receptor is an important determinant in identifying the host range and cross-species infection of viruses[24]. It has been established that ACE2 is the cellular receptor of SARS-CoV-2[16]. This study is targeted to predict viral entry in a host, *i.e.,* hosts that can be reservoir hosts (Artiodactyla, Perrisodactyla, Chiroptera, Carnivora, Lagomorpha, Primates, Pholidota, Proboscidea, Testutides, Crocodilia, Acciptriformes and Galliformes) and hosts that can be appropriate small animal laboratory models (Rodentia) of SARS-CoV-2 through sequence comparison and homology modeling of ACE2 and prediction

The protein and DNA sequence lengths of ACE2 varied in different hosts (**Table 1**). Among the sequences that were compared, the longest CDS was found in the Order - Chiroptera (*Myotis braditii* - 811 aa) and the smallest in the Order – Proboscidea (*Loxodonta africana* - 800 aa). *Homo sapiens* ACE2 is taken as a standard to compare all the sequences because of the on-going pandemic nature of COVID-19 and the availability of its 3D structure - 6LZG[25]. The within group mean distance, the parameter indicative of variability of nucleotide sequences within the group was found to be minimum in Perrisodactyla followed by Primates and was maximum among the Galliformes followed by Chiroptera (**Table 2**). This indicates that within the group of primates, all the considered species are prone to be equally infected with SARS-CoV-2 as humans. Further, to establish the probability of SARS-CoV-2 entry into species of other Orders, the distance of all orders from Primates was assessed **(Table 3)**. This distance was found minimum for Perissodactyls followed by Carnivores and maximum for Galliformes followed by Anseriformes. This confirms with the recent reports of Chicken (Galliformes) and ducks (Anseriformes) not being infected with SARS-CoV-2[22], and tigers and lions being infected[12]. To decide a cut-off distance that can establish whether the species can be infected or not, the individual distance of each species from *Homo sapiens* was evaluated **(Supplementary Table 3)**. *Melaegris gallapova* (Turkey) is the species, which had the greatest distance from *Homo sapiens*. Recently, it was reported that SARS-CoV-2 does not infect pigs, chickens, ducks[22] and rats[26]. The minimum distance that corresponds to the species that is already established to be uninfected with the SARS-CoV-2 would be 0.187 of *Rattus norvegicus* (Rat). Considering this distance from *Homo sapiens* as a cut-off, would include all the

carnivores, perissodactyls and few artiodactyls viz. Goat, buffalo, Bison and sheep, to be infected and excludes cattle (Artiodactyla), all the bats (Chiroptera) and birds (Galliformes, Anseriformes and Accipitriformes). Similar distance values were observed on evaluating the protein sequences as well (**Table 2 &Table 3**). These results do not lead to meaningful conclusions on viral entry in different species, thereby, making it inevitable to depend on other parameters like evaluating the spike-interacting domain of ACE2.

The spike interacting domain of the *Homo sapiens* ACE2 protein is defined in the UniProt ID Q9BYF1. The family and domains sections of the UniProt ID Q9BYF1 clearly marks the sequence location of the ACE2 - spike interacting domains as 30 - 41aa, 82 - 84 aa and 353 - 357 aa. The nucleotide sequence alignments at positions that correspond to the spike-binding domain of *Homo sapiens* ACE2 are 90-123 bp; 244-252 bp and 1058-1071 bp. This spike interacting ACE2 domain sequences at the nucleotide level and protein level (**Figure 1 and Figure 2**) were compared and evaluated. The alignment shows that the sequence is well conserved within the Orders, suggesting that the structure defined by the sequence is conserved within the Orders. The maximum variability *w.r.t.* the *Homo sapiens* sequence within these regions was observed for Galliformes, followed by Acciptriformes, Testidunes, Crocodilia and Chiroptera. The protein sequence alignment at 30-41aa, 82-84 aa and 353-357 also showed similar sequence conservation and variability (**Figure 2**). The Codon-based Test of Neutrality to understand the selection pressure on the ACE2 sequence in the process of evolution was done. The analysis showed that there was a significant negative selection between and within orders for the ACE2 sequence indicating that, though, there is a variation at

the nucleotide level, the protein translation had synonymous substitutions dominating over the non-synonymous substitutions. This negative selection indicates that the structure of ACE2 is being conserved through the process of evolution.

The protein sequences that were aligned were further subjected to find the best substitution model for phylogenetic analysis. The best model on the basis of BIC was found to be JTT + G. The phylogenetic analysis clearly classified the sequences of the species into their Orders. All the sequences were clearly grouped into two clusters. The first cluster represented the Mammalian class and the second cluster was represented by two sub- clusters of Avian and Reptilian classes with high bootstrap values (**Figure 3**). Within the mammalian cluster, the artiodactyls were sub-clustered farthest to the primates and the rodents, lagomorphs and carnivores were found clustered close to the primates with reliable bootstrap values. This partially corroborates with the occurrence of SARS-CoV-2 infection in carnivores[22] since rats were found uninfected with SARS-CoV-2[26]. The Chiroptera sub-cluster had a sub-node constituting horseshoe bat (*Rhinolophus ferrumequinum*) and the fruit bats (*Pteropus Alecto and Rousettus aegyptiacus*). The COVID-19 outbreak in Wuhan in Dec 2019 was traced back to have a probable origin from horseshoe bat[16]. The virus strain RaTG13 isolated from this bat was found to have 96.2% sequence similarity with the human SARS-CoV-2. This suggests that the virus probably could enter the fruit-bat as well, since it clustered with horseshoe bat to a common sub-node. These results again leave us with no concrete conclusions on viral entry in various hosts. Therefore, to assess the probability of viral entry in various species, homology modeling of ACE2 along with its interaction with coronavirus spike receptor-binding domain was analyzed for all the 48 hosts.

Homology modeling was done for all the ACE2 sequences based on the X-ray

diffraction structure defined in 6LZG (PDB database). The models constructed were

then studied for their interaction with the spike receptor binding domain defined in the

same ID. It was observed that the modelled interaction of human ACE2 showed four

hydrogen bonds between the ACE2 and Spike receptor binding domain. The hydrogen

bonds between the ACE2 and Spike receptor binding domain varied for different

species (**Supplementary Figure 1**). In FoldX, several parameters (spike binding

properties of ACE2) were estimated for the binding of ACE2 with spike receptor binding

domain. Logistic regression model was constructed on 17 species (known infected or

uninfected) using these parameters. When each parameter was considered individually,

significant difference between the infected and uninfected groups was observed for

Entropy side chain, Van der Waals, Solvation Polar, Solvation Hydrophobic and

Interface Residues (**Supplementary Table 4**). Each of the Order(s) was tested as a

group for their possibility of infection by comparing them with the infected and

uninfected groups for all these significant parameters (**Figure 4, Figure 5 & Figure 6**).

For the parameters - solvation hydrophobic and entropy side chain, artiodactyls were

found significantly ($P<0.05$) different from the uninfected group and not significantly

($P<0.05$) different from the infected group (**Figure 4**). This indicates that the artiodactyls

considered in the study can be infected. The testudines were significantly different from

the infected and not significantly different from the uninfected groups for all the

parameters (**Figure 5**). This suggests that the species considered under testudines may

not be infected. However, analysis for the Order - Chiroptera revealed that this group is

not significantly different from both the infected and uninfected groups (**Figure 6**) for all

the five parameters, leaving no clue about the probability of infection in this group. This suggests that a single parameter at a time, as has been considered in recent reports[21], may not be considered and evaluated for estimating the probability of virus entry. Therefore, all the estimated parameters were considered in logistic regression to find the best possible independent variables that would influence the entry of the SARS-CoV-2.

On evaluating several models, we finally included a model with Interaction energy, entropy side chain and entropy main chain, as independent variables, with an R^2 of 0.807. Hosmer and Lemeshow goodness of fit test showed no significant difference between the model and the observed data ($p > 0.05$) indicating that the model constructed is a good fit. There was also a statistically significant reduction in null deviance on inclusion of these three parameters (**Supplementary Table 5**). The predicted probabilities are given in **Table 4**. Within the Order Artiodactyla, all species except *Sus scrofa* (Pig) had 99% probability of viral (SARS-CoV-2) entry using ACE2 as a receptor. It has been predicted that *Bos indicus* (Indian cattle) and *Bos taurus* (Exotic cattle) can act as intermediate hosts of SARS-CoV-2[27] and that pigs are not susceptible[22]. Also, Camels, which are reported to be infected with SARS-CoV[28] are equally capable of SARS-CoV-2 infection. Among the rodents, hamsters had the highest probability of viral entry[26]. It has been established that SARS-CoV-2 effectively infects hamster[29] and, rats and mice were found less probable[26]. All the Carnivores except *Lontra canadensis* (Otter) in the study had high probability of viral entry. Reports of SARS-CoV2 infection in cats[22], tigers and lions[12] substantiate our estimates obtained in the study. Rabbits had medium probability of viral entry showing

some resemblance to the recent evidence of SARS-CoV-2 replication in rabbit cell lines[30]. In bats, the probability of viral entry was high in family Vespertilionidae. *Rhinolophus ferrumequinum* (horse-shoe bat) and *Phyllostomus discolor* (Pale spear-nosed bat) had lower probability of viral entry. The kidney cell line from the Rhinolophus genus was found infected with SARS-CoV but not with SARS-CoV-2[30]. However, probability of viral entry in chicken and ducks was found to be low. All the primates except baboon were predicted to have ~ 100% probability of viral entry as evident from the devasting nature of the disease in humans. Among the reptiles, both the testudines and crocodilia, showed low probability of viral entry. In the class Aves, *Anas platyrhynchos* (ducks) and *Haliaeetus albicilla* (eagles) showed the lowest probability followed by *Gallus gallus* (chicken). *Aquila chrysaetos chrysaetos* (Golden Eagle) and *Meleagris gallapova* (turkey) showed highest probability of viral entry.

Conclusion

Species considered under different orders, in this study, showed high probability of viral entry. However, viral entry is not the only factor that determines infection in COVID-19 as viral loads were found to be high in asymptomatic patients[31, 32]. The important factors that determine disease/infection(COVID-19) in host(s) are – Host defense potential, underlying health conditions, host behavior and number of contacts, Age, Atmospheric temperature, Population density, Airflow and ventilation and Humidity[33]. This study allows us to take a cue about the probable pathogenic insult that can be caused by SARS-CoV-2 to different species.

Materials and methods

Sequence analysis

In this study, 48 (mammalian, reptilian and avian species) ACE2 complete/partial protein and nucleotide sequences available on NCBI were analyzed (**Table 1**) to understand the possible difference(s) in the ACE2 sequences that may correlate with SARS-CoV-2 viral entry into the cell. Within the mammalian class, Orders - Artiodactyla, Perrisodactyla, Chiroptera, Rodentia, Carnivora, Lagomorpha, Primates, Pholidota and Proboscidea; within the Reptilian class, Orders - Testutides and Crocodile; and within the Avian class, Orders – Acciptriformes, Anseriformes and Galliformes, were considered in the study. These orders were considered keeping in view all the possible reservoir hosts/ laboratory animal models that can possibly be infected with the SARS-CoV-2. The within between group distances were calculated in Mega 6.0[34]. The Codon-based Z test of selection (strict-neutrality (dN=dS)) to evaluate synonymous and non-synonymous substitutions across the ACE2 sequences among the Orders was done.

Phylogenetic analysis

Phylogenetic analysis of the protein sequences was done using MEGA 6.0[34]. Initially, the sequence alignment was done using Clustal W[35]. The aligned sequences were analyzed for the best nucleotide substitution model on the basis of Bayesian information criterion scores using the **JModel**Test software v2.1.7[36]. The tree was constructed by the Neighbor-joining method with the best model obtained using 1000 bootstrap replicates.

Homology modeling

The Structure of novel coronavirus spike receptor-binding domain complexed with its receptor ACE2 which was determined through X-ray diffraction is available at PDB database with ID 6LZG[25]. This available ACE2 model from PDB databse is used for homology modeling using SWISS-MODEL[37]. SWISS-MODEL is a fully automatic homology modeling server for protein structure, which can be accessed through ExPASy web server.

Protein-protein docking

The spike receptor-binding domain of 6LZG was used in docking along with the homology modelled structures of ACE2 proteins of all the hosts, *i.e.*, ACE2 of 48 hosts as a receptor and spike receptor-binding domain of SARS-CoV-2 (from 6LZG) as a ligand for protein-protein docking. GRAMM-X docking server was used for protein-protein docking, which generated a docked complex[38]. Post-docking analysis was carried out using Chimera software[39], which is an extensible program for interactive visualization and analysis of molecular structures for use in structural biology. It provides the user with high quality 3D images, density maps, trajectories of small molecules and biological macromolecules, such as proteins. The homology modelled structure(s) of each species are compared with the human 6LZG to calculate the RMSD (root mean squared deviation). As most the deviation values could not be calculated with 6LZG model, the deviation(s) with respect to different human models 108a and 6M18[40] were calculated. A significant ($P < 0.05$) correlation in the deviation values calculated from 6LZG and 6M18 was observed. As most of the values could be calculated as deviations from 6M18, these values were used for further analysis along with the parameters below.

For the binding of the modelled structure(s) of ACE2 and the spike receptor-

binding domain, using FoldX software[41], several parameters (referred as spike

binding properties of ACE2) – Interaction Energy, Backbone Hydrogen bond, Side chain

Hydrogen bond, Van-der-Waals interaction, Electrostatic interaction, Solvation polar,

Solvation hydrophobic and Entropy sidechain, entropy mainchain, torsional clash,

backbone clash, helix dipole, disulfide, electrostatic kon, Interface Residues, Interface

Residue Clashing and Interface Residues VdW Clashing, were estimated.

Statistical analysis for prediction

Till date, clear-cut information of 17 species that are either infected or uninfected

with SARS-CoV2 is available (**Supplementary table 1**). Initially, for each parameter

(spike binding properties of ACE2), the difference between the infected and uninfected

is tested using both Mann-Whitney non-parametric test was done using GraphPad

Prism 7.00 (GraphPad Software, La Jolla, California, USA). For those parameters that

were significant the difference between Order(s) and the infected/uninfected groups was

established using Mann-Whitney non-parametric test (Note: if a species is included in

the infected/uninfected group, the same is not included in its Order on comparing the

Order(s) with infected/uninfected group) (**Supplementary table 2** for more information).

Later, a Logistic regression model was constructed on all the 18 parameters (17 from

FoldX and RMSD *w.r.t* 6M18) estimated above. With 18 parameters, the minimum

sample size required to derive statistics that represent each parameter, is 1000[42] *(n*

=100 + xi i.e here :- n = 100 + (100 + (50 × 18) = 1000, with a minimum of 50 events per

parameter). The data needed to be extrapolated to at least 1000. This needed us to

take an assumption that the ACE2 structure and sequence is conserved within a

species. For the species - *Homo sapiens*, we compared several ACE2 sequences and

found that all the compared sequences were identical. With this assumption that the

spike binding properties of ACE2 within a species are conserved and due to the

pandemic nature of the disease the data was extrapolated. All the parameters were

included in the glm - logistic regression to construct the best model (based on R^2) for

prediction. The goodness of fit was tested with Hosmer and Lemeshow goodness of fit

test. The reduction in null deviance was tested with Chi-square test.

Table 1. Species considered in this study

Order	Name (common name)	Accession number	Nucleotide length (bp)	Accession number	Amino acid length(aa)
Artiodactyla	*Bos indicus* **(Indian Cattle)**	XM_019956160.1	2436	XP_019811719.1	811
	Bos indicus x Bos taurus **(Indian crossbred Cattle)**	XM_027533926.1	2436	XP_027389727.1	811
	Bos taurus **(Exotic Cattle)**	XM_005228428.4	2436	XP_005228485.1	811
	Bubalus bubalis **(Buffalo)**	XM_006041540.2	2412	XP_006041602.1	803
	Bison bison bison **(American bison)**	XM_010834699.1	1294	XP_010833001.1	431
	Camelus bactrianus **(Double humped Camel)**	XM_010968001.1	2418	XP_010966303.1	805
	Camelus dromedaries **(Single humped camel)**	XM_010993415.2	2418	XP_010991717.1	805
	Capra hircus **(Goat)**	NM_001290107.1	2415	NP_001277036.1	804
	Ovis aries **(Sheep)**	XM_012106267.3	2415	XP_011961657.1	804
	Sus scrofa **(Pig)**	NM_001123070.1	2418	NP_001116542.1	805
Perissodactyla	*Equus asinus* **(Donkey)**	XM_014857647.1	2352	XP_014713133.1	783
	Equus caballus **(Horse)**	XM_001490191.5	2418	XP_001490241.1	805

	Pteropus alecto (**Black fruit bat**)	XM_006911647.1	2418	XP_006911709.1	805
	Rhinolophus ferrumequinum (**Greater horseshoe bat**)	AB297479.1	2418	BAH02663.1	805
	Myotis brandtii (**Brandt's bat**)	XM_014544294.1	2460	XP_014399780.1	819
Chiroptera	*Eptesicus fuscus* (**Big brown bat**)	XM_008154928.2	2436	XP_008153150.1	811
	Desmodus rotundus (**Common vampire bat**)	XM_024569930.1	2415	XP_024425698.1	804
	Phyllostomus discolor (**Pale spear-nosed bat**)	XM_028522516.1	2415	XP_028378317.1	804
	Rousettus aegyptiacus (**Egyptian fruit bat**)	XM_016118926.1	2418	XP_015974412.1	805
Pholidota	*Manis javanica* (**Sunda pangolin**)	XM_017650257.1	2418	XP_017505746.1	805
	Felis catus (**Cat**)	XM_023248796.1	2424	XP_023104564.1	807
	Panthera tigris altaica (**Siberian Tiger**)	XM_007090080.2	2394	XP_007090142.1	797
Carnivora	*Mustela putorius furo* (**Ferret**)	XM_004758885.2	2418	XP_004758942.1	805
	Canis lupus familiaris (**Dog**)	NM_001165260.1	2415	NP_001158732.1	804
	Vulpes vulpes (**Red Fox**)	XM_025986727.1	2415	XP_025842512.1	804
	Lontra canadensis (**North American river otter**)	XM_032880138.1	2418	XP_032736029.1	805
Rodentia	*Mus musculus* (**Mouse**)	NM_027286.4	2418	NP_081562.2	805
	Rattus norvegicus (**Rat**)	NM_001012006.1	2418	NP_001012006.1	805

	Cricetulus griseus **(Hamster)**	XM_027432806.1	2412	XP_027288607.1	805
Lagomorpha	*Oryctolagus cuniculus* **(Rabbit)**	XM_002719845.3	2418	XP_002719891.1	805
	Ochotona princeps **(American pika)**	XM_004597492.2	2427	XP_004597549.2	808
Primates	*Homo sapiens* **(Human)**	NM_001371415.1	2418	NP_001358344.1	805
	Pan troglodytes **(Chimpanzee)**	XM_016942979.1	2418	XP_016798468.1	805
	Papio anubis **(Baboon)**	XM_021933040.1	2418	XP_021788732.1	805
	Macaca nemestrina **(Southern pig-tailed monkey)**	XM_011735203.2	2418	XP_011733505.1	805
	Macaca mulatta **(Rhesus monkey)**	NM_001135696.1	2418	NP_001129168.1	805
	Macaca fascicularis **(Crab eating monkey)**	XM_005593037.2	2418	XP_005593094.1	805
Proboscidea	*Loxodonta Africana* **(African elephant)**	XM_023555192.1	2403	XP_023410960.1	800
Galliformes	*Gallus gallus* **(Chicken)**	XM_416822.5	2427	XP_416822.2	808
	Meleagris gallopavo **(Turkey)**	XM_019612009.2	2586	XP_019467554.1	861
Anseriformes	*Anas platyrhynchos* **(Mallard)**	XM_013094461.3	2418	XP_012949915.2	805
Accipitriformes	*Aquila chrysaetos chrysaetos* **(Golden Eagle)**	XM_029999165.1	2430	XP_029855025.1	809
	Haliaeetus albicilla **(White-tailed eagle)**	XM_009927339.1	1887	XP_009925641.1	629
Crocodilia	*Alligator sinensis* **(Chinese alligator)**	XM_025210843.1	2412	XP_025066628.1	803

	Crocodylus porosus **(Salt water alligator)**	XM_019529281.1	2412	XP_019384826.1	803
Testudines	*Pelodiscus sinensis* **(Chinese softshell turtle)**	XM_006122829.3	2427	XP_006122891.1	808
	Chelonia mydas **(Green sea turtle)**	XM_007070499.1	2436	XP_007070561.1	811
	Chrysemys picta bellii **(Painted turtle)**	XM_024108749.1	2487	XP_023964517.1	828

Table 2. With Mean group distance among the Orders

Order	Within Mean group distance (DNA)	Within Mean group distance (Protein)
Perrisodactyla	0.01	0.02
Primates	0.02	0.03
Accipitriformes	0.03	0.03
Crocodilia	0.04	0.05
Carnivora	0.07	0.10
Testudines	0.07	0.11
Artiodactyla	0.08	0.10
Rodentia	0.10	0.12
Lagomorpha	0.12	0.13
Chiroptera	0.14	0.23
Galliformes	0.21	0.28

Table 3. Between group distance (between Primates and other groups)

Order	Primates (DNA)	Primates (Protein)
Perrisodactyla	0.131	0.164
Carnivora	0.162	0.199
Pholidota	0.163	0.183
Lagomorpha	0.165	0.197
Rodentia	0.179	0.211
Chiroptera	0.181	0.249
Artiodactyla	0.186	0.241
Proboscidea	0.189	0.237

Testudines	0.516	0.573
Crocodilia	0.518	0.565
Accipitriformes	0.562	0.528
Anseriformes	0.594	0.587
Galliformes	0.605	0.653

Table 4. Probability of viral entry in different species

Class	Order	Family	Species (Common name)	Probability of Viral Entry (95% Confidence Interval)
Mammalia	Artiodactyla	Bovidae	*Bos indicus* **(Indian Cattle)**	9.979E-01(9.91E-01 – 9.99E-01)
			Bos taurus **(Exotic Cattle)**	9.978E-01(9.91E-01 – 9.99E-01)
			Bubalus bubalis **(Buffalo)**	9.954E-01(9.92E-01 – 1.00E+00)
			Bison bison bison **(American bison)**	1.00E+00(1.00E+00 – 1.00E+00)
			Bos indicus x *Bos taurus* **(Indian crossbred Cattle)**	9.979E-01(9.92E-01 – 1.00E+00)
		Camilidae	*Camelus bactrianus* **(Double humped Camel)**	9.989E-01(9.93E-01 – 1.00E+00)
			Camelus dromedaries **(Single humped camel)**	9.989E-01(9.93E-01 – 1.00E+00)
		Caprinae	*Capra hircus* **(Goat)**	9.998E-01(9.99E-01 – 1.00E+00)
			Ovis aries **(Sheep)**	9.998E-01(9.99E-01 – 1.00E+00)
		Suidae	*Sus scrofa* **(Pig)**	1.40E-02(2.44E-03 – 7.58E-02)
	Perrisodactyla	Equidae	*Equus asinus* **(Donkey)**	1.00E+00(1.00E+00 – 1.00E+00)

			Equus caballus (Horse)	4.647E-01(4.12E-02 – 9.46E-01)
	Carnivora	Mustelidae	Mustela putorius furo (Ferret)	9.95E-01(9.50E-01 – 1.00E+00)
			Lontra canadensis (North American river otter)	4.971E-07(5.74E-09 – 4.31E-05)
		Felidae	Panthera tigris altaica (Siberian Tiger)	9.57E-01(8.80E-01 – 9.86E-01)
		Canidae	Vulpes Vulpes (Red Fox)	9.889E-01(9.41E-01 – 9.98E-01)
			Canis lupus familiaris (Dog)	1.000E+00(9.99E-01 – 1.00E+00)
		Felidae	Felis catus (Cat)	1.000E+00(1.00E+00 – 1.00E+00)
	Chiroptera	Rhinolophidae	Rhinolophus ferrumequinum (Greater horseshoe bat)	9.269E-04(8.24E-05 – 1.03E-02)
		Phyllostomidae	Desmodus rotundus (Common vampire bat)	9.928E-01(9.47E-01 – 9.99E-01)
			Phyllostomus discolor (Pale spear-nosed bat)	7.237E-04(3.23E-05 – 1.60E-02)
		Vespertilionidae	Eptesicus fuscus (Big brown bat)	1.000E+00(1.00E+00 – 1.00E+00)
			Myotis brandtii (Brandt's bat)	9.998E-01(9.99E-01 – 1.00E+00)
		Pteropodidae	Pteropus Alecto (Black fruit bat)	2.650E-01(9.60E-03 – 9.31E-01)
			Rousettus aegyptiacus (Egyptian fruit bat)	4.83E-01(3.84E-01 – 5.84E-01)
	Rodentia	Cricetidae	Cricetulus griseus (Hamster)	8.92E-01(7.92E-01 – 9.47E-01)

		Muridae	*Mus musculus* **(Mouse)**	2.05E-04(4.56E-06 – 9.12E-03)
			Rattus norvegicus **(Rat)**	1.41E-03(1.91E-04 – 1.03E-02)
	Lagomorpha	Leporidae	*Oryctolagus cuniculus* **(Rabbit)**	6.760E-01(2.73E-01 – 9.20E-01)
		Ochotonidae	*Ochotona princeps* **(American pika)**	1.275E-01(2.94E-02 – 4.13E-01)
	Pholidota	Manidae	*Manis javanica* **(Sunda pangolin)**	1.000E+00(9.99E-01 – 1.00E+00)
	Primates	Hominidae	*Homo sapiens* **(Human)**	1.00E+00(9.98E-01 – 1.00E+00)
		Cercopithecidae	*Macaca fascicularis* **(Crab eating monkey)**	1.00E+00(1.00E+00 – 1.00E+00)
			Macaca mulatta **(Rhesus monkey)**	1.00E+00(9.99E-01 – 1.00E+00)
			Macaca nemestrina **(Southern pig-tailed monkey)**	1.00E+00(1.00E+00 – 1.00E+00)
		Hominidae	*Pan troglodytes* **(Chimpanzee)**	1.00E+00(9.98E-01 - 1.00E+00)
		Cercopithecidae	*Papio Anubis* **(Baboon)**	1.109E-09(4.36E-13 – 2.82E-06)
	Probosidae	Elephantidae	*Loxodonta Africana* **(African elephant)**	9.998E-01(9.98E-01 – 1.00E+00)
Reptiles	Testidunes	Cheloniidae	*Chelonia mydas* **(Green sea turtle)**	9.371E-03(9.11E-05 – 4.95E-01)
		Emydidae	*Chrysemys picta bellii* **(Painted turtle)**	3.781E-09(2.23E-13 – 6.41E-05)
		Trionychidae	*Pelodiscus sinensis* **(Chinese softshell turtle)**	3.851E-04(4.26E-06 – 3.37E-02)

	Crocodilia	Alligatoridae	*Alligator sinensis* **(Chinese alligator)**	6.27E-04(1.64E-06 – 1.94E-01)
		Crocodylidae	*Crocodylus porosus* **(Salt water alligator)**	2.223E-02(1.66E-04 – 7.57E-01)
Aves	Galliformes	Phasianidae	*Gallus gallus* **(Chicken)**	6.58E-01(5.81E-01 – 7.28E-01)
			Meleagris gallapova **(Turkey)**	1.00E+00(1.00E+00 – 1.00E+00)
	Anseriformes	Anatidae	*Anas platyrhynchus* **(Mallard)**	1.84E-10(1.11E-14 – 3.05E-06)
	Accipitriformes	Accipitridae	*Haliaeetus albicilla* **(White-tailed eagle)**	4.168E-01(1.61E-01 – 7.27E-01)
			Aquila chrysaetos chrysaetos **(Golden Eagle)**	9.999E-01(9.99E-01 – 1.00E+00)

Abbreviations

ACE2: Angiotensin-converting enzyme 2

CDS: Coding Sequence

COVID-19: Coronavirus disease 2019

ICTV: International Committee on Taxonomy of Viruses

MERS: Middle East Respiratory Syndrome

PDB: Protein Data Bank

RMSD: Root-mean-square deviation

SADS: Swine Acute Diarrhea Syndrome

SARS-CoV-2: Severe Acute Respiratory Syndrome Coronavirus 2

SARS: Severe Acute Respiratory Syndrome

WHO: World Health Organization

References

1. **COVID-19, MERS & SARS** [https://www.niaid.nih.gov/diseases-conditions/covid-19]
2. Drosten C, Gunther S, Preiser W, van der Werf S, Brodt HR, Becker S, Rabenau H, Panning M, Kolesnikova L, Fouchier RA *et al*: **Identification of a novel coronavirus in patients with severe acute respiratory syndrome**. *The New England journal of medicine* 2003, **348**(20):1967-1976.

3.   Zhou P, Fan H, Lan T, Yang XL, Shi WF, Zhang W, Zhu Y, Zhang YW, Xie QM, Mani S *et al*: **Fatal swine acute diarrhoea syndrome caused by an HKU2-related coronavirus of bat origin**. *Nature* 2018, **556**(7700):255-258.

4.   Fan Y, Zhao K, Shi ZL, Zhou P: **Bat Coronaviruses in China**. *Viruses* 2019, **11**(3).

5.   COVID-19 CORONAVIRUS PANDEMIC [https://www.worldometers.info/coronavirus/]

6.   Subissi L, Posthuma CC, Collet A, Zevenhoven-Dobbe JC, Gorbalenya AE, Decroly E, Snijder EJ, Canard B, Imbert I: **One severe acute respiratory syndrome coronavirus protein complex integrates processive RNA polymerase and exonuclease activities**. *Proceedings of the National Academy of Sciences of the United States of America* 2014, **111**(37):E3900-3909.

7.   Forni D, Cagliani R, Clerici M, Sironi M: **Molecular Evolution of Human Coronavirus Genomes**. *Trends in microbiology* 2017, **25**(1):35-48.

8.   Zaki AM, van Boheemen S, Bestebroer TM, Osterhaus AD, Fouchier RA: **Isolation of a novel coronavirus from a man with pneumonia in Saudi Arabia**. *The New England journal of medicine* 2012, **367**(19):1814-1820.

9.   Scientists hunt for the next potential coronavirus animal host [https://www.nationalgeographic.com/animals/2020/03/coronavirus-animal-reservoir-research/]

10. **Hong Kong: Blood tests confirm that pomerarian caught COVID-19 after its owner** [https://www.theweek.in/news/world/2020/03/26/hong-kong-blood-tests-confirm-that-pomerarian-caught-covid-19-after-its-owner.html]

11. **A cat appears to have caught the coronavirus, but it's complicated** [https://www.sciencenews.org/article/cats-animals-pets-coronavirus-covid19]

12. **Four tigers, three lions test Covid-19 positive at Bronx Zoo** [https://timesofindia.indiatimes.com/world/us/four-tigers-three-lions-test-covid-19-positive-at-bronx-zoo/articleshow/75319387.cms]

13. **Mink found to have coronavirus on two Dutch farms: ministry** [https://in.reuters.com/article/health-coronavirus-netherlands-mink/mink-found-to-have-coronavirus-on-two-dutch-farms-ministry-idINKCN2280K2]

14. Li W, Moore MJ, Vasilieva N, Sui J, Wong SK, Berne MA, Somasundaran M, Sullivan JL, Luzuriaga K, Greenough TC *et al*: **Angiotensin-converting enzyme 2 is a functional receptor for the SARS coronavirus**. *Nature* 2003, **426**(6965):450-454.

15. Hoffmann M, Kleine-Weber H, Schroeder S, Kruger N, Herrler T, Erichsen S, Schiergens TS, Herrler G, Wu NH, Nitsche A *et al*: **SARS-CoV-2 Cell Entry Depends on ACE2 and TMPRSS2 and Is Blocked by a Clinically Proven Protease Inhibitor**. *Cell* 2020, **181**(2):271-280 e278.

16. Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, Si HR, Zhu Y, Li B, Huang CL *et al*: **A pneumonia outbreak associated with a new coronavirus of probable bat origin**. *Nature* 2020, **579**(7798):270-273.

17. Hamming I, Timens W, Bulthuis ML, Lely AT, Navis G, van Goor H: **Tissue distribution of ACE2 protein, the functional receptor for SARS coronavirus. A first step in understanding SARS pathogenesis.** *The Journal of pathology* 2004, **203**(2):631-637.

18. Donoghue M, Hsieh F, Baronas E, Godbout K, Gosselin M, Stagliano N, Donovan M, Woolf B, Robison K, Jeyaseelan R *et al*: **A novel angiotensin-converting enzyme-related carboxypeptidase (ACE2) converts angiotensin I to angiotensin 1-9.** *Circulation research* 2000, **87**(5):E1-9.

19. Li F, Li W, Farzan M, Harrison SC: **Structure of SARS coronavirus spike receptor-binding domain complexed with receptor.** *Science* 2005, **309**(5742):1864-1868.

20. Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh CL, Abiona O, Graham BS, McLellan JS: **Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation.** *Science* 2020, **367**(6483):1260-1263.

21. Qiu Y, Zhao YB, Wang Q, Li JY, Zhou ZJ, Liao CH, Ge XY: **Predicting the angiotensin converting enzyme 2 (ACE2) utilizing capability as the receptor of SARS-CoV-2.** *Microbes and infection* 2020.

22. Shi J, Wen Z, Zhong G, Yang H, Wang C, Huang B, Liu R, He X, Shuai L, Sun Z *et al*: **Susceptibility of ferrets, cats, dogs, and other domesticated animals to SARS-coronavirus 2.** *Science* 2020.

23. **Novel Coronavirus SARS-CoV-2: Fruit bats and ferrets are susceptible, pigs and chickens are not** [https://www.fli.de/en/press/press-releases/press-singleview/novel-coronavirus-sars-cov-2-fruit-bats-and-ferrets-are-susceptible-pigs-and-chickens-are-not/]

24. Li F: **Receptor recognition and cross-species infections of SARS coronavirus.** *Antiviral research* 2013, **100**(1):246-254.

25. Wang Q, Zhang Y, Wu L, Niu S, Song C, Zhang Z, Lu G, Qiao C, Hu Y, Yuen KY *et al*: **Structural and Functional Basis of SARS-CoV-2 Entry by Using Human ACE2.** *Cell* 2020.

26. Zhang H, Penninger JM, Li Y, Zhong N, Slutsky AS: **Angiotensin-converting enzyme 2 (ACE2) as a SARS-CoV-2 receptor: molecular mechanisms and potential therapeutic target.** *Intensive care medicine* 2020, **46**(4):586-590.

27. Luan J, Jin X, Lu Y, Zhang L: **SARS-CoV-2 spike protein favors ACE2 from Bovidae and Cricetidae.** *Journal of medical virology* 2020.

28. Gong SR, Bao LL: **The battle against SARS and MERS coronaviruses: Reservoirs and Animal Models.** *Animal models and experimental medicine* 2018, **1**(2):125-133.

29. Lau SY, Wang P, Mok BW, Zhang AJ, Chu H, Lee AC, Deng S, Chen P, Chan KH, Song W *et al*: **Attenuated SARS-CoV-2 variants with deletions at the S1/S2 junction.** *Emerging microbes & infections* 2020:1-15.

30. Chu H, Chan JF-W, Yuen TT-T, Shuai H, Yuan S, Wang Y, Hu B, Yip CC-Y, Tsang JO-L, Huang X *et al*: **Comparative tropism, replication kinetics, and cell damage profiling of SARS-CoV-2 and SARS-CoV with implications for clinical manifestations,**

transmissibility, and laboratory studies of COVID-19: an observational study. *The Lancet Microbe* 2020.

31. Rabi FA, Al Zoubi MS, Kasasbeh GA, Salameh DM, Al-Nasser AD: **SARS-CoV-2 and Coronavirus Disease 2019: What We Know So Far**. *Pathogens* 2020, **9**(3).

32. Zou L, Ruan F, Huang M, Liang L, Huang H, Hong Z, Yu J, Kang M, Song Y, Xia J *et al*: **SARS-CoV-2 Viral Load in Upper Respiratory Specimens of Infected Patients**. *The New England journal of medicine* 2020, **382**(12):1177-1179.

33. Lakshmi Priyadarsini S, Suresh M: **Factors influencing the epidemiological characteristics of pandemic COVID 19: A TISM approach**. *International Journal of Healthcare Management* 2020:1-10.

34. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S: **MEGA6: Molecular Evolutionary Genetics Analysis version 6.0**. *Molecular biology and evolution* 2013, **30**(12):2725-2729.

35. Thompson JD, Higgins DG, Gibson TJ: **CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice**. *Nucleic acids research* 1994, **22**(22):4673-4680.

36. Darriba D, Taboada GL, Doallo R, Posada D: **jModelTest 2: more models, new heuristics and parallel computing**. *Nature methods* 2012, **9**(8):772.

37. Waterhouse A, Bertoni M, Bienert S, Studer G, Tauriello G, Gumienny R, Heer FT, de Beer TAP, Rempfer C, Bordoli L *et al*: **SWISS-MODEL: homology modelling of protein structures and complexes**. *Nucleic acids research* 2018, **46**(W1):W296-W303.

38. Tovchigrechko A, Vakser IA: **GRAMM-X public web server for protein-protein docking**. *Nucleic acids research* 2006, **34**(Web Server issue):W310-314.

39. Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE: **UCSF Chimera--a visualization system for exploratory research and analysis**. *Journal of computational chemistry* 2004, **25**(13):1605-1612.

40. Yan R, Zhang Y, Li Y, Xia L, Guo Y, Zhou Q: **Structural basis for the recognition of SARS-CoV-2 by full-length human ACE2**. *Science* 2020, **367**(6485):1444-1448.

41. Strokach A, Corbi-Verge C, Kim PM: **Predicting changes in protein stability caused by mutation using sequence-and structure-based methods in a CAGI5 blind challenge**. *Human Mutation* 2019, **40**(9):1414-1423.

42. Bujang MA, Sa'at N, Sidik T, Joo LC: **Sample Size Guidelines for Logistic Regression from Observational Studies with Large Population: Emphasis on the Accuracy Between Statistics and Parameters Based on Real Life Clinical Data**. *Malays J Med Sci* 2018, **25**(4):122-130.

Figure legends

Figure 1. **Nucleotide sequence alignment of the CDS region of ACE2**. The shaded

regions show the spike interacting domains.

Figure 2. Protein sequence alignment of ACE2. The shaded regions show the spike

interacting domains.

Figure 3. **Phylogenetic analysis of ACE2 protein sequences**. The tree was

constructed using neighbor joining method in MEGA 6.0. The bootstrap values

are given at each node.

Figure 4. Scatterplot showing the comparison of Artiodactyls with infected and

uninfected groups for the all five significant parameters (A). Van der Waals

– No Significant difference on comparison of Artiodactyls with infected and

uninfected groups. (B). Entropy side chain - Significant difference on comparison

of Artiodactyls with uninfected group and no significant difference from infected

group. (C). Solvation hydrophobic - Significant difference on comparison of

Artiodactyls with uninfected and no significant difference from infected. (D).

Interface residues - No Significant difference on comparison of Artiodactyls with

infected and uninfected groups. (E). Solvation polar - No Significant difference on

comparison of Artiodactyls with infected and uninfected groups. ** Significance at

$P < 0.01$; * Significance at $P < 0.05$ after Mann-Whitney test on comparing two

groups at a time.

Figure 5. Scatterplot showing the comparison of Testudines with infected and

uninfected groups for the all five significant parameters (A). Van der Waals

– Significant difference on comparison of Testudines with infected and no significant difference from uninfected. (B). Entropy side chain - Significant difference on comparison of Testudines with infected and no significant difference from uninfected. (C). Solvation hydrophobic - Significant difference on comparison of Testudines with infected and no significant difference from uninfected. (D). Interface residues- Significant difference on comparison of Testudines with infected and no significant difference from uninfected. (E). Solvation polar - Significant difference on comparison of Testudines with infected and no significant difference from uninfected. ** Significance at $P < 0.01$; * Significance at $P < 0.05$ after Mann-Whitney test on comparing two groups at a time.

Figure 6. Scatterplot showing the comparison of Chiroptera with infected and uninfected groups for the all five significant parameters (A). Van der Waals – No Significant difference on comparison of Chiroptera with infected and uninfected groups. (B). Entropy side chain - No Significant difference on comparison of Chiroptera with infected and uninfected groups. (C). Solvation hydrophobic - No Significant difference on comparison of Chiroptera with infected and uninfected groups. (D). Interface residues- No Significant difference on comparison of Chiroptera with infected and uninfected groups. (E). Solvation polar - No Significant difference on comparison of Chiroptera with infected and uninfected groups. ** Significance at $P < 0.01$; * Significance at $P < 0.05$ after Mann-Whitney test on comparing two groups at a time.

1 **Supplementary Figure 1.** Representative protein modelled structures showing the

2   interaction between ACE2 of (A) Human (B) Cat (C) Donkey (D) Exotic cattle (E)

3   Chinese alligator & (F) Greater horseshoe bat, and spike receptor binding

4   domain of SARS-CoV-2.

5 **Author's contributions**

13 **Competing interests**

14 The author has declared no competing interests.

15 **Acknowledgments**

16 We are grateful to Director NIAB and Director IVRI for the support.

Shaded region - Spike interacting region : 117bp - 123bp of *Homo sapiens*

```
GTCTTATCAAAGTTCACTTGCTTCTTGGAATTATAACACCAATATTACCGATGAGAATGTCCAAAAGATGAATGAAGCTGGGGCCAAATGGTCTGCCTTTTATGAAGAACAGTCCAAGATTGCCA Majority
    ---+--------+---------+---------+---------+---------+---------+---------+---------+---------+---------+---------+---------+---------+
       130       140       150       160       170       180       190       200       210       220       230       240       250
```

```
117 ..........................................................................CA.A...........................................CG...G.... Bos_indicus.seq
117 ..........................................................................CA.A...........................................CG...G.... Bos_indicus x Bos taurus.seq
117 ..........................................................................CA.A...........................................CG...G.... Bos_taurus.seq
117 ..........................................................................CA.A...........................................CG...G.... Bison bison bison.seq
114 ..........................................................................CA....................C.......................CG...G.... Bubalus_bubalis.seq
117 ..........................................................................CA...........................................CG...G.... Ovis_aries.seq
117 ..........................................................................CA........................................A...CG...G.... Capra_hircus.seq
117 .G.....................C.AT...............A...................T.CA.....................C.............A..TCG........ Sus_scrofa.seq
117 ...........................A...................T.CA........A....................A.......C... Camelus_bactrianus.seq
117 ...........................A...................T.CA........A....................A.......C... Camelus_dromedarius.seq
117 ....C.....................G.....................T.C..............C.......G.......................G....C.... Equus_asinus.seq
117 ....C.....................G.....................T.C..............C.......G.......................G....C.... Equus_caballus.seq
117 A.A.............G...G...................C.....G....A....G..C.ACAAG.G.........A.......................G.C.... Phyllostomus_discolor.seq
117 ..T.........C.........G..........................CA.....................C.......................C..... Pteropus_alecto.seq
117 ....C.G.....G....C...........A.........C.ACAG...G................C.................C....C Myotis_brandtii.seq
117 ....C.G......G.G..........A.....ACAG...G.........A.....................C....C Eptesicus_fuscus.seq
117 ..T........A..........C.............A....G...A.CAG.TG.....A.............AG.A.C.....C.... Desmodus_rotundus.seq
117 ....C...........G.A.......G...............G...G......A.....A.......C... Rhinolophus_ferrumequinum.seq
117 ..T.......G...T.......C.T..........G.A....CA.....C......................C... Rousettus_aegyptiacus.seq
117 ...................C..........TT......CA.......................C... Manis_javanica.seq
117 .................C.....C..A..C.......A...G.C................C.... Felis_catus.seq
 93 .................C.....C..A..C.......A...G.C................C...G Panthera_tigris.seq
114 ...............G...........T....C........A.T.........................C.... Vulpes_vulpes.seq
114 .................T...........C........A.T.........................C.... Canis_lupus_familiaris.seq
117 .....A...............C....A......ATT.....G.................G.....C..CA..... Mustela_putorius_furo.seq
117 .....A...............C....A......ATT..A....G.................G.....C...A..... Lontra canadensis.seq
117 ..............T..T..C....T..A..A...C..........G...G....CA.............T....C....C Mus_musculus.seq
117 ..........C..........C......G..G....C..........C..G....C.....................C...C Rattus_norvegicus.seq
117 ....G.................T..A....C..........G....CT.......T...............T...C..... Cricetulus_griseus.seq
117 ...........G..........G............A.......T..A....................C... Oryctolagus_cuniculus.seq
126 C........G......G...G.C......C...T..A...C...A...C.T..C.A....G.....................C... Ochotona_princeps.seq
117 ..TC.....................T..A.......C....A.T.......A................TAA.G..........CAC....C Homo_sapiens.seq
117 ..TC.....................T..A.......C....A.T......AA.......TAA.........CAC...C Macaca_fascicularis.seq
117 ..TC.......................A.......C....A.T......AA.......TAA.........CAC..C Macaca_mulatta.seq
117 ..TC.....................T..A.......C....A.T......AA.......TAA.........CAC...C Macaca_nemestrina.seq
117 ..TC.....................T..A.......C....A.T.......A................TAA.G........CAC...C Pan_troglodytes.seq
117 ..TC.....................T..A.......C....A.T......AA.......TAA.........GCAC...C Papio_anubis.seq
117 .................G.................C.........T...A....G.......T.......GAG......C..C.... Loxodonta_africana.seq
114 CAGC...G...AC.........C..G......C..C...........C..A..T.G..CA.C.AGG..A....G...G..A..T........G.A..A..........G..GGCC.....G..AC..T. Gallus_gallus.seq
114 CAGC...G...AC.........C..A...G.C..C...........C..A..T.G..CA.C.AGG..G...G..G..C........G.A..A..........G..GGCC.....G..AC..T. Meleagris_gallopavo.seq
114 CAAC...G...AC..C.......C..A..C.........C...T.G...CG.C.AC.......G.C.GT...G...G..G........C........G..GGCC.....G..A.... Anas platyrhynchos.seq
114 CAGC..G.G..C..C..Seq.....C..C.A...C..C.T..C.AGG..A.A...G..GT...G...C.A........C...G........TC..GGCC..G..AC.... Aquila_chrysaetos_chrysaetos.seq
113 CAGC..G.G..C.C.......C.A...C..C.T..C.A..........G..GT...G...C.A.....C.GGC-.---------..C.... Haliaeetus albicilla.seq
102 ..A.....G...........C..A...GCC..C.........C..A...C.A.G..A.........A.AT...G......AG......CA.TA..GCA..T..C.A...G. Crocodylus_porosus.seq
102 ..A.....G.............G...GCC..C.....C...A..A....C.A.G..........A.AT...G......AG......TA..GCA..T..C.A..... Alligator_sinensis.seq
114 ......GC.............C..C...C.........C..C.T...A....C.A.............A..A............A.TA..C......T...GCA......CC.A...... Pelodiscus_sinensis.seq
114 A......GC.........C..C...G.C.........C..C..G.........TC.AG................A..A............A..C...T...GCA........C.A..... Chrysemys picta bellii.seq
114 .........GC.........C..C...G.C..C.......C..C..A.........C.CTG.........A..A............A..C....T.C.GCA.......C.A..... Chelonia_mydas.seq
```

Artiodactyla

Perissodactyla

Chiroptera

Pholidota

Carnivora

Rodentia

Lagomorpha

Primates

Proboscidea

Galliformes

Anseriformes

Accipitriformes

Crocodilia

Testudines

Mammalia

Aves

Reptilia

Shaded region - Spike interacting region : 244bp - 252bp of *Homo sapiens*

```
---MSGSSWLLLSLVAVTAAQSTTEELAKTFLEKFNHEAEDLSYQSSLASWNYNTNITDENVQKMNEAGAKWSAFYEEQSKLAKTYPLEEIQNLTVKRQLQALQQSGSSVLSADKSK--RLNTILNTMST  Majority

         10        20        30        40        50        60        70        80        90       100       110       120       130
```

Sequence	Group
Bos_taurus.pro	Artiodactyla
Bos_indicus x Bos_taurus.pro	
Bos_indicus.pro	
Bison bison bison.pro	
Bubalus_bubalis.pro	
Capra_hircus.pro	
Ovis_aries.pro	
Sus_scrofa.pro	
Camelus_bactrianus.pro	
Camelus_dromedarius.pro	
Equus_caballus.pro	Perissodactyla
Equus_asinus.pro	
Myotis_brandtii.pro	Chiroptera
Pteropus alecto.pro	
Eptesicus_fuscus.pro	
Desmodus_rotundus.pro	
Phyllostomus_discolor.pro	
Rhinolophus_ferrumequinum.pro	
Rousettus_aegyptiacus.pro	
Manis_javanica.pro	Pholidota
Vulpes_vulpes.pro	Carnivora
Felis_catus.pro	
Canis_lupus familiaris.pro	
Panthera_tigris_altaica.pro	
Mustela_putorius_furo.pro	
Lontra_canadensis.pro	
Rattus_norvegicus.pro	Rodentia
Mus_musculus.pro	
Cricetulus griseus.pro	
Ochotona princeps.pro	Lagomorpha
Oryctolagus_cuniculus.pro	
Homo_sapiens.pro	Primates
Pan_troglodytes.pro	
Macaca_nemestrina.pro	
Papio_anubis.pro	
Macaca_fascicularis.pro	
Macaca_mulatta.pro	
Loxodonta africana.pro	Proboscidea
Gallus_gallus.pro	Galliformes
Meleagris_gallopavo.pro	
Anas_platyrhynchos.pro	Anseriformes
Aquila chrysaetos chrysaetos.pro	Accipitriformes
Haliaeetus albicilla.pro	
Alligator_sinensis.pro	Crocodilia
Crocodylus_porosus.pro	
Pelodiscus sinensis.pro	Testudines
Chrysemys picta bellii.pro	
Chelonia_mydas.pro	

Higher-order groups: Mammalia (Artiodactyla through Proboscidea), Aves (Galliformes, Anseriformes, Accipitriformes), Reptilia (Crocodilia, Testudines)