Targeted sequence capture array for phylogenetics and population genomics in the

Salicaceae[1]

Brian J. Sanderson[2,3,6], Stephen P. DiFazio[3], Quentin C. Cronk[4], Tao Ma[5], Matthew S. Olson[2]

[2]Department of Biological Sciences, Texas Tech University, Lubbock, TX 79409-3131 USA

[3]Department of Biology, West Virginia University, Morgantown, WV, 26506 USA

[4]Department of Botany, University of British Columbia, Vancouver, BC, V6T 1Z4 Canada

[5]Key Laboratory of Bio-Resource and Eco-Environment of Ministry of Education, College of

Life Sciences, Sichuan University, Chengdu 610065, People's Republic of China

[6]Author for correspondence: brian@biologicallyrelevant.com

Number of words: 3649

Abstract

•   **Premise of the study**: The family Salicaceae has proved taxonomically challenging,

16   especially in the genus *Salix*, which is speciose and features frequent hybridization and

17   polyploidy. Past efforts to reconstruct the phylogeny with molecular barcodes have failed to

18   resolve the species relationships of many sections of the genus.

19   •   **Methods**: We used the wealth of resequencing data in the family to design sequence capture

20   probes to target regions of 300-1200 base pairs of exonic regions of 972 genes.

21   •   **Results**: We recovered sequence data for nearly all of the targeted genes in three species of

22   *Populus* and three species of *Salix*. We present a species tree, discuss concordance among

23   gene trees, as well as some population genomic summary statistics for these loci.

24   •   **Conclusions**: Our sequence capture array has extremely high capture efficiency within the

25   genera *Populus* and *Salix*, resulting in abundant phylogenetic information. Additionally,

26   these loci show promise for population genomic studies.

Key words: Phylogenetics; *Populus*; Salicaceae; *Salix*; targeted sequence capture

Introduction

Although the cost of whole-genome sequencing has continued to dramatically decrease over the

past decade, the cost and complexity of whole-genome analyses still limit their utility and

accessibility for answering evolutionary questions in novel taxa (Richards, 2018). However,

robust genome assembly is not required for many questions, and several methods have been

developed to reduce the cost and effort required to obtain genomic information in novel species

(McKain et al., 2018). The recent development of targeted sequence capture presents an

affordable method for consistently isolating specific, long, phylogenetically informative regions

in the taxa of interest (Gnirke et al., 2009; Mamanova et al., 2010; Hale et al., 2020). Targeted

sequence capture uses biotinylated RNA baits to target prepared sequencing library fragments.

The baited library fragments can then be pulled out of solution with streptavidin-coated magnetic

beads to selectively enrich the fragments that contain loci of interest, while discarding the

majority of library fragments that do not. This method offers many advantages over other

methods of genome sequence partitioning, such as genome skimming and RAD-seq, because it

does not necessarily depend on a highly polished, annotated reference genome, and because the

same loci can be consistently sequenced at a high depth across individuals without requiring

comprehensive, concurrent sequencing of all individuals (Mamanova et al., 2010; Grover et al.,

2012; Jones and Good, 2016).

In this paper, we report on the design and implementation of a targeted sequence capture

array to collect data for phylogenetic analysis within the Salicaceae, the plant family that

includes poplars and willows. Understanding species relationships within this family, and in

particular the genus *Salix*, has presented challenges to taxonomists as early as Linnaeus, who

noted that "species of this genus are extremely difficult to clarify" (Linnaeus, 1753; Skvortsov,

1999). *Salix* species present challenges to classification due to their wide geographic ranges,

hysteranthous phenology, extensive interspecific hybridization, polyploidy, and the lack of well-

defined morphological characters (Raup, 1959; Skvortsov, 1999; Percy et al., 2014; Wang et al.,

2020). Species of *Salix* exhibit holarctic distributions, and there are several classifications which

differ among continents and are challenging to synthesize due to incomplete overlap of species

(Dickmann and Kuzovkina, 2014). Past efforts to reconstruct the phylogeny of *Salix* using

nuclear AFLP and plastid barcode sequences have resulted in a lack of clearly resolved species

relationships, especially in the subgenus *Vetrix* (Trybush et al., 2008; Percy et al., 2014). A more

recent study using a supermatrix approach with RAD-seq data showed resolution within a subset

of species of the subgenera *Vetrix* and *Chamaetia*, highlighting the potential of genome-scale

markers to resolve this phylogenetically challenging group (Wagner et al., 2018).

The utility of RAD-seq for collecting data for phylogeny, however, is limited by several

issues. First, RAD-seq does not consistently screen homologous regions across species and

across different experiments, which limits its utility for adding species to a phylogeny at a later

time. Second, because RAD-seq assesses diversity in very short segments of the genome,

phylogenetic analysis requires concatenation and supermatix approaches (de Queiroz and

Gatesy, 2007) and does not allow separate exploration of gene and species phylogenies using

super tree methods (Sanderson et al., 1998). Additionally, concatenation approaches are likely to

exacerbate problems associated with maximum-likelihood methods for species with rapid

diversification (Edwards et al., 2007; Edwards, 2009). Targeted sequence capture does not have

these limitations, and thus may be a more appropriate genotyping platform for phylogenetics.

Species of *Populus* and *Salix* have been of great interest for the development of forestry

and biofuel products, resulting in polished reference genomes for *P. trichocarpa*, *P. tremula*, *P.*

euphratica, *S. purpurea* and *S. suchowensis*., as well as shallow resequencing data for many

additional species (Tuskan et al., 2018). Our design strategy leveraged this abundance of existing

genomic information to quantify polymorphism and the distribution of insertion- deletion events

within and among species in order to maximize capture efficiency. Additionally, because we

consistently target regions of exons we are able to characterize the nucleotide-site degeneracy

with these data to quantify population genomic summary statistics. We demonstrate the utility of

this resource for *Populus* and *Salix* species by presenting a fully resolved phylogenetic tree for

six species and an outgroup, and by estimating the distribution of nucleotide diversity within

species for our targeted genes.

Methods

Probe Design

Our goal was to identify regions that could be efficiently captured using RNA bait hybridization

across the entire family Salicaceae. Although we were interested in using these probes for

phylogenetics with both *Populus* and *Salix* species, we focused on maximizing capture efficiency

for the species in *Salix*, because the phylogeny for *Populus* is already much better resolved than

that for *Salix* (Trybush et al., 2008; Wang et al., 2014; Percy et al., 2014; Liu et al., 2017; Wang

et al., 2020). For this reason, the capture baits were designed to target regions in *Salix pupurea*

that also would have high capture efficiency across the Salicaceae. The efficiency of RNA bait

binding, and thus capture efficiency, is reduced as target regions diverge due to sequence

polymorphism (Lemmon and Lemmon, 2013). To improve capture efficiency, we quantified

sequence polymorphism among whole-genome resequencing data from a diverse array of

Populus and *Salix* species (Table S1). The whole-genome short reads of the *Populus* and *Salix*

species were aligned to the *Populus trichocarpa* genome assembly version 3 (Tuskan et al.,

2006) using bwa mem (Li, 2013). We used the *P. trichocarpa* genome as our initial reference

because it was the most polished annotated genome in genus. Variable sites and insertion-

deletion mutations (indels) were identified using samtools mpileup (Li, 2011), and read depth for

the variant calls was quantified using vcftools (Danecek et al., 2011). Custom Python scripts

were used to identify variant and indel frequencies for all exons in the *P. trichocarpa* genome

annotation (scripts available at https://github.com/BrianSanderson/phylo-seq-cap).

Orthologs for our candidate loci in the in the *Salix purpurea* 94006 genome assembly

version 1 (*Salix purpurea* v1.0, DOE-JGI; Carlson et al., 2017; Zhou et al., 2018) were identified

using a list of orthologs between the *P. trichocarpa* and *S. purpurea* prepared using a tree-based

approach by Phytozome (Goodstein et al., 2012). We further screened candidate regions to

exclude high-similarity duplicated regions by accepting only loci with single BLAST (Camacho

et al., 2009) hits against the highly contiguous assembly of *S. purpurea* 94006 version 5 (Zhou et

al., 2020), which is less fragmented than the *S. purpurea* 94006 version 1. Genes from the

Salicoid whole-genome duplication were identified using MCScanX (Wang et al., 2012), using

default parameters but selecting segments for which the average K_S value for paralogous genes

was between 0.2 and 0.8. Genes for which at least 600 base pairs (bp) of exon sequence

contained 2-12% polymorphism and fewer than 2 indels were selected for probe design by Arbor

Biosciences (Ann Arbor, MI, USA). Probes were designed with 50% overlap across the targeted

regions, so that each nucleotide position would potentially be captured by two probes. Finally, to

ensure that loci with high divergence across the family would be captured, we identified targets

with less than 95% identity (based on BLAST results) between *S. purpurea* and *P. trichocarpa*

and designed supplementary probes from orthologs of these genes in the *Idesia polycarpa*

genome.

Library Preparation and Sequence Capture

Libraries for two individuals from each of three species of *Populus* and three species of *Salix*

(Table 1) were prepared using the NEBNext Ultra II DNA Prep Kit (New England Biolabs,

Ipswitch, MA, USA), and quantified using an Agilent Bioanalyzer 2100 DNA 1000 kit (Agilent

Technologies, Santa Clara, CA, USA). Libraries were pooled at equimolar concentrations into

two pools of six prior to probe hybridization following the Arbor Biosciences myBaits protocol v

3.0.1 and Hale et al. (2020). The hybridized samples were subsequently pooled at equimolar

ratios and sequenced at the Texas Tech Center for Biotechnology and Genomics using a MiSeq

with the Micro chemistry and 150 bp paired-end reads (Illumina, Inc., San Diego, CA, USA).

Analysis of Sequence Capture Data

Read data was trimmed for primer sequences and low quality scores using Trimmomatic v. 0.36

(Bolger et al., 2014). The trimmed read data, as well as the whole-genome reads for *I. polycarpa*,

were assembled into gene sequences using the HybPiper pipeline (Johnson et al., 2016). We

estimated the depth of read coverage across all targeted genes using custom Python and R

scripts. The assembled amino acid sequences were aligned with mafft v. 7.310 (Katoh and

Standley, 2013), converted into codon-aligned nucleotide alignments with pal2nal v. 14 (Suyama

et al., 2006), and trimmed for quality and large gaps with trimal v. 1.4.rev15 (Capella-Gutiérrez

et al., 2009).

HybPiper provides warnings for genes that have multiple competing assemblies that are

within 80% of the length of the target region, because the alternate alignments may indicate that

those genes have paralogous copies in the genome. We estimated phylogenetic relationships

using both the full set of gene sequences recovered from our sequence capture data, as well as a

restricted set of putatively single copy genes, based on our *a priori* list of paralogs between *S.*

purpurea and *P. tricocarpa*, as well as the list of paralog warnings from HybPiper.

We estimated gene trees using RAxML v. 8.2.10, specifying a GTRΓ model of sequence

evolution (Stamatakis, 2014). A set of 250 bootstrap replicates was generated for each gene tree.

We used ASTRAL-III to infer the species tree from the RAxML gene trees (Zhang et al., 2018;

Rabiee et al., 2019). Because all nodes are weighted equally during quartet decomposition in

ASTRAL-III, we used sumtrees in the Python package DendroPy v. 4.4.0 to collapse nodes with

less than 33% bootstrap support values prior to species tree estimation (Sukumaran and Holder,

2010). A set of 100 multilocus bootstrap replicates was generated for the species tree. We used

phyparts to determine the extent of congruence among gene trees for each node in the species

tree (Smith et al., 2015). Cladograms representing the gene tree congruence and alternate

topologies were plotted with the scripts phypartspiecharts.py and minority_report.py, written by

Matt Johnson (scripts available at https://github.com/mossmatters/phyloscripts).

Finally, we used custom Python and R scripts to quantify nucleotide diversity at

synonymous and non-synonymous sites between the individuals of the same species, as well as

correlations in values of per-site nucleotide diversity between all species. The scripts described

above as well as the full details of these analyses including are available in Jupyter notebooks at

https://github.com/BrianSanderson/phylo-seq-cap).

Results

Sequence capture efficiency

The final capture kit targets 972 genes covered by 12,951 probes based on the *S. purpurea*

reference, and an additional 7049 (redundant) probes based on the *I. polycarpa* genome that

target genes with the highest divergence between *S. purpurea* and *P. trichocarpa*. This included

an average of 680 $\pm$ 309 (mean $\pm$ sd) probes on each *S. purpurea* chromosome (Table S2), with

an average of 1098 $\pm$ 489 (mean $\pm$ sd) bp of exon sequence per gene. Of the 972 target genes,

593 are putatively single copy based on the list of paralogs between *S. purpurea* and *P.*

trichocarpa, 142 represent pairs of paralogs from the shared Salicoid whole-genome duplication,

and 237 are genes that have known paralogs for which we were not able to design targets in this

kit. We included a total of 1219 genes in the target file used to assemble the capture data, which

includes the 972 targeted genes as well as paralogous copies for which probes were not designed.

Because the issues of paralogy become more complex when we add species other than *S.*

purpurea and *P. trichocarpa*, we advise using the HybPiper warnings of multiple competing

long assemblies to assess paralogy in novel species following guidance here

https://github.com/mossmatters/HybPiper/wiki/Paralogs. The sequence capture kit is available

from Arbor Biosciences (Ref#170424-30 "Salicaceae").

Sequence capture efficiency was high among the libraries. We recovered 805,820 $\pm$

178,482 reads (mean $\pm$ sd) from our *Populus* and *Salix* target capture libraries, of which 86.7 $\pm$

1.15% (mean $\pm$ sd) mapped to the target sequence reference (Table 1). An average of 94.48 $\pm$

1.37% of targeted exon sequences were covered by $\geq$ 10 reads. The average read depth was

44.65 $\pm$ 1.61 for on-target sites, and 14.48 $\pm$ 2.10 for off-target sites (Table S3).

Phylogenetics

The species tree estimated with putatively single copy genes correctly paired all individuals of

the same species and revealed a fully resolved phylogeny for the *Populus* and *Salix* species with

100% multilocus bootstrap support for all nodes (Fig. 1A). The majority of gene trees supported

the topology of the species tree (Fig. 1B), with the exceptions of the bipartition that separates *P.*

balsamifera and *P. tremula*, and the bipartition that separates *S. phlebophylla* from the other

Salix species, which had dominant alternate topologies that were supported by a large number of

gene trees (Figs. S1 and S2). The topology of the species tree estimated with the full set of genes

and known paralogs was nearly identical to the tree estimated with only the putatively single

copy genes. The major difference between these trees was evident in the bipartition separating *P.*

balsamifera and *P. tremula*, where there were a large number of alternative topologies supported

by small numbers of gene trees (the top 3 were supported by 13, 11, and 10 gene trees; Fig. S3).

Population genomics

Patterns of nucleotide diversity, measured as Nei's π (Nei and Li, 1979), varied among species,

with the greatest variation at synonymous sites (Fig. S4; Table S4). *P. tremula* had the highest

average values of π at both synonymous and non-synonymous sites (Fig. 2). The values of π

among species were highly correlated for species within genus and exhibited lower correlations

between genera (Fig. 3).

Discussion

The decreasing cost of obtaining genomic and transcriptomic sequence data holds great promise

for unlocking our understanding of phylogenetic relationships and population genetic patterns

within and among complex taxonomic groups. However, assembling complete genomes is still

not a trivial task, and there exist relatively few polished plant reference genomes onto which

genome skimming data can be mapped. Many methods have been developed to reduce the

sequencing and analytical burdens associated with obtaining genome data. We believe that

targeted sequence capture is one of the most promising contemporary methods of inexpensively

generating genomic information.

209   The efficiency of our targeted sequence capture array was extremely high, which yielded

abundant phylogenetic information for six species of *Populus* and *Salix*. Overall, the phylogeny

was fully resolved and conformed to our general understanding the relationships among the taxa.

One strength of the sequence capture approach is that it provides sufficiently long contiguous

segments of gene sequences to assemble gene trees and can overcome the problems introduced

by concatenation of multiple gene regions with divergent histories (Edwards et al., 2007;

Edwards, 2009). The super tree approach also allowed for the identification of alternative

evolutionary histories that are supported by different regions of the genome, as often occurs

during historical hybridization and introgression (Zhang et al., 2018; Rabiee et al., 2019). Our

species tree identified three alternative gene tree relationships among the three *Populus* species

(Fig. S1). Previous studies have provided evidence of historical introgression among these

species, including a history of chloroplast capture and hybridization between *P. mexicana* and

species in the section *Tacamahaca* (including *P. balsamifera*; Wang et al., 2014; Liu et al., 2017;

Wang et al., 2020). The second most supported alternative topology that we recovered placed *P.*

mexicana and *P. tremula* as sister taxa, a pattern that does not support this hypothesis, likely due

to incomplete lineage sorting (Wang et al., 2020). *P. tremula* likely has a greater long-term

effective population size than *P. balsamifera* (Wang et al., 2016), and so coalescence times may

be shorter on average in *P. balsamifera*. Among the *Salix* species, we identified three alternative

gene tree relationships between the *S. phlebophylla* and *S. exigua* individuals, which may reflect

the histories of rapid speciation and hybridization that have long vexed attempts at phylogenetic

reconstruction in the genus *Salix* (Appendix 5; Trybush et al., 2008; Percy et al., 2014). Both of

these patterns in *Populus* and *Salix* may be better understood once additional taxa are added to

this phylogeny.

We have also shown that this sequence capture design can be applied to address questions

related to population genomics in the Salicaceae. Many of the advantages of targeted sequence

capture over competing methods are of particular relevance for population genomics studies,

including specific knowledge of loci being sequenced, the ability to differentiate among

synonymous, non-synonymous, intronic, and intergenic loci, and the ability to collect data on the

same set of loci across different experiments, either within species or across species, for

comparative studies. In particular, synonymous sites, especially four-fold synonymous sites, are

among the fastest evolving regions of the genome and the sites within genic regions least

influenced by selection (Wright and Andolfatto, 2008), and are thus among the best regions for

estimating patterns of historical demography. Our estimates of nucleotide diversity are similar to

those that have been previously reported for *P. balsamifera* and *P. tremula* using Sanger

sequencing data (Ingvarsson, 2005; Olson et al., 2010) and whole-genome sequencing data

(Wang et al., 2016). The high estimates of diversity in *S. phlebophylla* compared to the other two

Salix species is curious and may result from a history with relatively little migration due to the

absence of glaciation over a large portion of its Beringian distribution (Hultén, 1937).

The current study is based on a small sample size per species (n = 2), and so our ability to

account for population structure or robustly perform population genomic inferences with these

data is limited. Additionally, a potential limitation for using this sequence capture array for

comparative population genomics is that we screened loci for a range of among-species

variability between 2-12%, which excludes loci that exhibit extremely high or low values of

nucleotide diversity. This may bias estimates of nucleotide diversity arising from these probes

toward greater evenness. The ability to identify synonymous sites, which are the closest to

neutral among all classes of sites (Wright and Andolfatto, 2008), should partially address this

bias. Another feature of sequence capture data is the recovery of "off-target" sequences that

result from the fact that the insert size of libraries is larger than the 120 bp bait length, and so

regions upstream and downstream of the target will be sequenced as well. These regions may

include intronic and intergenic regions, as well as exonic sequences that deviate from the

constraints we used for our design. The results we report here only incorporate the "on-target"

sites that we sequenced, but HybPiper implements methods to assemble intronic sequences as

well. However, the potential effects of hitchhiking selection on synonymous site variation will

likely remain apparent.

We also found that it was straightforward to integrate the targeted sequence capture data

with whole-genome sequence data using the HybPiper pipeline by simply including the FASTQ

files from whole-genome reads in the pipeline. This strategy was used to successfully incorporate

whole-genome sequencing data from *Idesia polycarpa*, to act as our outgroup. The proportion of

gene coverage as well as the read depth for the *I. polycarpa* data was similar to the sequence

capture libraries (Table 1).

A whole genome duplication occurred prior to the divergence of *Salix* and *Populus*, and

there are at least 8000 known paralog pairs in the *P. trichocarpa* reference genome (Tuskan et

al., 2006). Genes with paralogous copies in the genome can complicate gene assemblies, because

sequence data from both copies may alternately align to the same target sequence. We used the

list of known paralogs in the *S. purpurea* genome to design the sequence capture array, and it

includes 593 putatively single copy genes, 142 pairs of paralogs, and 237 genes which have

paralogs but for which we were not able to include both paralogs in the kit due our selection

criteria. The target reference file we used to map the sequence capture data thus includes 1219

genes including the single copy and known paralogs from *S. purpurea*. In addition to this,

HybPiper provides warnings for genes that have multiple competing alignments that cover the

majority of the target sequence, which may indicate the presence of multiple paralogous copies

in the genome (Johnson et al., 2016). This will be particularly useful because the list of known

paralogs in *S. purpurea* is likely to differ among species throughout the diversification of

willows. We estimated evolutionary relationships using both the full set of 1219 single copy and

known paralog genes, as well as a limited set of just single copy genes that did not report paralog

warnings. The results from both analyses were nearly the same, but this will likely not be true for

a more complex phylogenetic analysis that includes more than six species and an outgroup. For

those more complex phylogenetic analyses, the ability to compare trees constructed with single

copy genes with those using paralogous copies may provide crucial information for reconciling

evolutionary relationships.

This sequence capture array will provide the community with an excellent resource to

consistently sequence a set of variable regions of the genome for phylogenetic and population

genomic investigations in the Salicaceae. The rate of read mapping and coverage of target genes

was remarkably consistent across both genera, despite the fact that the taxa were selected to

maximize sampling of phylogenetic diversity within each genus. The Salicaceae are important

plants in the northern hemisphere both ecologically and economically and have been the subjects

of numerous population genetics and genomics investigations of speciation, hybridization,

introgression, selection, and historical population size and migration. This resource will allow

phylogenetic and comparative population genomic studies to assess the same loci across different

studies, which will allow us to build a worldwide diversity database and facilitate more precise

comparative research questions. Our results demonstrate that the rate of gene capture is

extremely high, such that it would be unnecessary to filter data and determine appropriate

overlapping genotype thresholds, as is necessary with random genome partitioning methods such

as RAD-seq.

Acknowledgements

We thank M. Johnson for help with data analysis, P. Ingvarsson for help collecting *Populus*

tremula, and J. Martinez, Instituto de Geología, UNAM, Hermosillo, Mexico for help collecting

Populus mexicana. We thank J. Phillips for his work at Phytozome in identifying orthologs

between *P. trichocarpa* and *S. purpurea*. Funding for this project was provided by the US

National Science Foundation (1542509 and 1542599), Genome Canada (168BIO), and the

National Natural Science Foundation of China (31561123001).

Author Contributions

S.P.D. and M.S.O. conceived the study. S.P.D., Q.C.C., T.M., and M.S.O. secured funding to

support the project. B.J.S. and S.P.D. designed the sequence capture array. Q.C.C. and T.M.

provided whole genome resequencing data. B.J.S and M.S.O. prepared and sequenced the DNA

samples, analyzed the data, interpreted the results, and wrote the manuscript. All authors edited

drafts of the manuscript.

Data Accessibility

The raw reads of targeted sequence capture from the six species of *Populus* and *Salix* will be

available on the NCBI sequence read archive upon acceptance of this manuscript. The full details

of the analysis, including all of the custom Python scripts that were used, are summarized in

notebooks available at https://github.com/BrianSanderson/phylo-seq-cap, and upon acceptance

of this manuscript a DOI will be generated from Zenodo to reference the state of the repository at

the time of acceptance.

Literature Cited

BOLGER, A.M., M. LOHSE, and B. USADEL. 2014. Trimmomatic: A flexible trimmer for Illumina

sequence data. *Bioinformatics* 30: 2114–2120.

CAMACHO, C., G. COULOURIS, V. AVAGYAN, N. MA, J. PAPADOPOULOS, K. BEALER, and T.L.

MADDEN. 2009. BLAST+: Architecture and applications. *BMC Bioinformatics* 10: 421–421.

CAPELLA-GUTIÉRREZ, S., J.M. SILLA-MARTÍNEZ, and T. GABALDÓN. 2009. trimAl: A tool for

automated alignment trimming in large-scale phylogenetic analyses. *Bioinformatics* 25: 1972–

1973.

CARLSON, C.H., Y. CHOI, A.P. CHAN, M.J. SERAPIGLIA, C.D. TOWN, and L.B. SMART. 2017.

Dominance and Sexual Dimorphism Pervade the Salix purpurea L. Transcriptome. *Genome*

Biology and Evolution 9: 2377–2394.

DANECEK, P., A. AUTON, G. ABECASIS, C.A. ALBERS, E. BANKS, M.A. DEPRISTO, R.E.

HANDSAKER, ET AL. 2011. The variant call format and VCFtools. *Bioinformatics* 27: 2156–2158.

DE QUEIROZ, A., and J. GATESY. 2007. The supermatrix approach to systematics. *Trends in*

Ecology and Evolution 22: 34–41.

DICKMANN, D.I., and J. KUZOVKINA. 2014. Poplars and Willows of the World, With Emphasis

on Silviculturally Important Species. *In* J. Isebrands, and J. Richardson [eds.], Poplars and

Willows Trees for Society and the Environment, 8–91. The Food and Agriculture Organization

of the United Nations, Rome, Italy.

DOE-JGI. Salix purpurea version 1.

http://phytozome.jgi.doe.gov/pz/portal.html#!info?alias=Org_Spurpurea

EDWARDS, S.V. 2009. Is a new and general theory of molecular systematics emerging? *Evolution*

63: 1–19.

EDWARDS, S.V., L. LIU, and D.K. PEARL. 2007. High-resolution species trees without

concatenation. *Proceedings of the National Academy of Sciences* 104: 5936–5941.

GNIRKE, A., A. MELNIKOV, J.R. MAGUIRE, P. ROGOV, E.M. LEPROUST, W. BROCKMAN, T.J.

FENNELL, ET AL. 2009. Solution hybrid selection with ultra-long oligonucleotides for massively

parallel targeted sequencing. *Nature Biotechnology* 27: 182–189.

GOODSTEIN, D.M., S. SHU, R. HOWSON, R. NEUPANE, R.D. HAYES, J. FAZO, T. MITROS, ET AL.

2012. Phytozome: A comparative platform for green plant genomics. *Nucleic Acids Research* 40:

D1178–D1186.

GROVER, C.E., A. SALMON, and J.F. WENDEL. 2012. Targeted sequence capture as a powerful

tool for evolutionary analysis. *American Journal of Botany* 99: 312–319.

HALE, H., E.M. GARDNER, J. VIRUEL, L. POKORNY, and M.G. JOHNSON. 2020. Strategies for

reducing per-sample costs in target capture sequencing for phylogenomics and population

genomics in plants: Low-cost Hyb-Seq. *Applications in Plant Sciences*e11337.

HULTÉN, E. 1937. Outline of the history of arctic and boreal biota during the Quarternary period :

Their evolution during and after the glacial period as indicated by the equiformal progressive

areas of present plant species. Bokförlags aktiebolaget Thule, Stockholm, Sweden.

INGVARSSON, P.K. 2005. Nucleotide polymorphism and linkage disequilibrium within and

among natural populations of European aspen (Populus tremula L., Salicaceae). *Genetics* 169:

945–953.

JOHNSON, M.G., E.M. GARDNER, Y. LIU, R. MEDINA, B. GOFFINET, A.J. SHAW, N.J.C. ZEREGA,

and N.J. WICKETT. 2016. HybPiper: Extracting Coding Sequence and Introns for Phylogenetics

from High-Throughput Sequencing Reads Using Target Enrichment. *Applications in Plant*

Sciences 4: 1600016–1600016.

JONES, M.R., and J.M. GOOD. 2016. Targeted capture in evolutionary and ecological genomics.

Molecular Ecology 25: 185–202.

KATOH, K., and D.M. STANDLEY. 2013. MAFFT Multiple Sequence Alignment Software

Version 7: Improvements in Performance and Usability. *Molecular Biology and Evolution* 30:

772–780.

LEMMON, E.M., and A.R. LEMMON. 2013. High-Throughput Genomic Data in Systematics and

Phylogenetics. *Annual Review of Ecology, Evolution, and Systematics* 44: 99–121.

LI, H. 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM.

00: 1–3.

LI, H. 2011. A statistical framework for SNP calling, mutation discovery, association mapping

and population genetical parameter estimation from sequencing data. *Bioinformatics* 27: 2987–

2993.

LINNAEUS, C. 1753. Species plantarum. Impensis Laurentii Salvii, Stockholm.

LIU, X., Z. WANG, W. SHAO, Z. YE, and J. ZHANG. 2017. Phylogenetic and Taxonomic Status

Analyses of the Abaso Section from Multiple Nuclear Genes and Plastid Fragments Reveal New

Insights into the North America Origin of Populus (Salicaceae). *Frontiers in Plant Science* 7: 1–

9.

MAMANOVA, L., A.J. COFFEY, C.E. SCOTT, I. KOZAREWA, E.H. TURNER, A. KUMAR, E. HOWARD,

ET AL. 2010. Target-enrichment strategies for next-generation sequencing. *Nature Methods* 7:

111–118.

MCKAIN, M.R., M.G. JOHNSON, S. URIBE-CONVERS, D. EATON, and Y. YANG. 2018. Practical

considerations for plant phylogenomics. *Applications in Plant Sciences* 6: e1038–e1038.

NEI, M., and W.H. LI. 1979. Mathematical model for studying genetic variation in terms of

restriction endonucleases. *Proceedings of the National Academy of Sciences* 76: 5269–5273.

OLSON, M.S., A.L. ROBERTSON, N. TAKEBAYASHI, S. SILIM, W.R. SCHROEDER, and P. TIFFIN.

2010. Nucleotide diversity and linkage disequilibrium in balsam poplar (Populus balsamifera).

New Phytologist 186: 526–536.

PERCY, D.M., G.W. ARGUS, Q.C. CRONK, A.J. FAZEKAS, P.R. KESANAKURTI, K.S. BURGESS,

B.C. HUSBAND, ET AL. 2014. Understanding the spectacular failure of DNA barcoding in willows

(Salix): Does this result from a trans-specific selective sweep? *Molecular Ecology* 23: 4737–

4756.

RABIEE, M., E. SAYYARI, and S. MIRARAB. 2019. Multi-allele species reconstruction using

ASTRAL. *Molecular Phylogenetics and Evolution* 130: 286–296.

RAUP, H.M. 1959. The willows of boreal Western America. *Contributions from the Gray*

Herbarium of Harvard University 185: 3–95.

RICHARDS, S. 2018. Full disclosure: Genome assembly is still hard. *PLOS Biology* 16:

e2005894–e2005894.

SANDERSON, M.J., A. PURVIS, and C. HENZE. 1998. Phylogenetic supertrees: Assembling the

trees of life. *Trends in Ecology and Evolution* 13: 105–109.

SKVORTSOV, A.K. 1999. Willows of Russia and Adjacent Countries. I. N. (. Kadis, G. R. Argus,

and A. G. Zinovjev [eds.], University of Joensuu, Joensuu.

SMITH, S.A., M.J. MOORE, J.W. BROWN, and Y. YANG. 2015. Analysis of phylogenomic datasets

reveals conflict, concordance, and gene duplications with examples from animals and plants.

BMC Evolutionary Biology 15: 150–150.

STAMATAKIS, A. 2014. RAxML version 8: A tool for phylogenetic analysis and post-analysis of

large phylogenies. *Bioinformatics* 30: 1312–1313.

SUKUMARAN, J., and M.T. HOLDER. 2010. DendroPy: A Python library for phylogenetic

computing. *Bioinformatics* 26: 1569–1571.

SUYAMA, M., D. TORRENTS, and P. BORK. 2006. PAL2NAL: Robust conversion of protein

sequence alignments into the corresponding codon alignments. *Nucleic Acids Research* 34:

W609–W612.

TRYBUSH, S., Š. JAHODOVÁ, W. MACALPINE, and A. KARP. 2008. A genetic study of a Salix

germplasm resource reveals new insights into relationships among subgenera, sections, and

species. *Bioenergy Research* 1: 67–79.

TUSKAN, G.A., S.P. DIFAZIO, S. JANSSON, J. BOHLMANN, I. GRIGORIEV, U. HELLSTEN, N.

PUTNAM, ET AL. 2006. The Genome of Black Cottonwood, Populus trichocarpa (Torr. & Gray).

Science 313: 1596–1604.

TUSKAN, G.A., A.T. GROOVER, J. SCHMUTZ, S.P. DIFAZIO, A. MYBURG, D. GRATTAPAGLIA, L.B.

SMART, ET AL. 2018. Hardwood Tree Genomics: Unlocking Woody Plant Biology. *Frontiers in*

Plant Science 9: 1799–1799.

WAGNER, N.D., S. GRAMLICH, and E. HÖRANDL. 2018. RAD sequencing resolved phylogenetic

relationships in European shrub willows (Salix L. Subg. Chamaetia and subg. Vetrix) and

revealed multiple evolution of dwarf shrubs. *Ecology and Evolution* 8: 8243–8255.

WANG, J., N.R. STREET, D.G. SCOFIELD, and P.K. INGVARSSON. 2016. Natural Selection and

Recombination Rate Variation Shape Nucleotide Polymorphism Across the Genomes of Three

Related Populus Species. *Genetics* 202: 1185–1200.

WANG, M., L. ZHANG, Z. ZHANG, M. LI, D. WANG, X. ZHANG, Z. XI, ET AL. 2020.

Phylogenomics of the genus *Populus* reveals extensive interspecific gene flow and balancing

selection. *New Phytologist* 225: 1370–1382.

WANG, Y., H. TANG, J.D. DEBARRY, X. TAN, J. LI, X. WANG, T.-H. LEE, ET AL. 2012. MCScanX:

A toolkit for detection and evolutionary analysis of gene synteny and collinearity. *Nucleic Acids*

Research 40: e49–e49.

WANG, Z., S. DU, S. DAYANANDAN, D. WANG, Y. ZENG, and J. ZHANG. 2014. Phylogeny

reconstruction and hybrid analysis of populus (Salicaceae) based on nucleotide sequences of

multiple single-copy nuclear genes and plastid fragments. *PLoS ONE* 9:.

WRIGHT, S.I., and P. ANDOLFATTO. 2008. The Impact of Natural Selection on the Genome:

Emerging Patterns in Drosophila and Arabidopsis. *Annual Review of Ecology, Evolution, and*

Systematics 39: 193–213.

ZHANG, C., M. RABIEE, E. SAYYARI, and S. MIRARAB. 2018. ASTRAL-III: Polynomial time

species tree reconstruction from partially resolved gene trees. *BMC Bioinformatics* 19: 153–153.

ZHOU, R., D. MACAYA-SANZ, C.H. CARLSON, J. SCHMUTZ, J.W. JENKINS, D. KUDRNA, A.

SHARMA, ET AL. 2020. A willow sex chromosome reveals convergent evolution of complex

palindromic repeats. *Genome Biology* 21: 38–38.

ZHOU, R., D. MACAYA-SANZ, E. RODGERS-MELNICK, C.H. CARLSON, F.E. GOUKER, L.M.

EVANS, J. SCHMUTZ, ET AL. 2018. Characterization of a large sex determination region in Salix

purpurea L. (Salicaceae). *Molecular Genetics and Genomics* 293: 1437–1452.

Tables

Name	Num. Reads	Reads Mapped	Prop. Mapped	Genes Mapped	Genes with 25% Seq	Genes with 50% Seq	Genes with 75% Seq	Genes with 100% Seq
I_polycarpa_WGS-2	223470714	1653494	0.007	971	970	966	944	123
P_balsamifera_MGR-01	614093	523321	0.852	972	971	960	884	122
P_balsamifera_MGR-04	769303	659712	0.858	972	972	965	915	145
P_mexicana_PM3	843032	739728	0.878	972	972	964	917	140
P_mexicana_PM5	880962	768927	0.873	972	972	967	913	142
P_tremula_R01-01	749220	638002	0.852	971	970	960	907	134
P_tremula_R04-01	634625	539805	0.851	971	969	956	876	122
S_exigua_SE002	1139616	998698	0.876	969	969	966	937	229
S_exigua_SE053	843120	741938	0.88	969	969	964	928	195
S_nigra_SG037	1166615	1028635	0.882	971	971	967	932	205
S_nigra_SG051	602649	524993	0.871	971	970	961	903	136
S_phlebophylla_SP15M	753628	651791	0.865	972	972	967	939	204
S_phlebophylla_SP7F	672975	581147	0.864	972	972	967	925	203

Table 1. Coverage summary statistics for sequence capture read data. For each library, values

represent the number of reads in the sequenced library, the number of those reads that mapped to

the reference file for the targeted genes, the proportion of mapped reads, the number of targeted

genes (out of 972) that had read data mapped to them, and the number of genes that had 25%,

50%, 75%, and 100% of the targeted sequences covered with > 10X reads. Footnote: the

I_polycarpa_WGS-2 data is from whole-genome sequencing data, rather than targeted sequence

capture.

Figures

Figure 1. Species trees estimated for the 432 putatively single copy genes that did not have

paralog warnings reported by HybPiper. **A)** Species tree generated by ASTRAL-III for the gene

trees. Node values represent bootstrap support from 100 multilocus bootstrap replicates in

ASTRAL-III. Branch lengths represent coalescent units. **B)** Cladogram showing the congruence

of gene trees for all nodes in the ASTRAL-III species tree. The numbers above each node

represent the number of gene trees that support the displayed bipartition, and numbers below the

node represent the number of gene trees that support all alternate bipartitions. Purple wedges

represent the proportion of gene trees that support the displayed bipartition. Blue wedges

represent the proportion of gene trees that support a single alternative bipartition (see Figs S1 &

S2). Green wedges represent the proportion of gene trees that have multiple conflicting

bipartitions. Yellow wedges represent the proportion of gene trees that have no supported

bipartition. Plotting code and its interpretation were provided by Matt Johnson (for more detail,

see:

https://github.com/mossmatters/MJPythonNotebooks/blob/master/PhyParts_PieCharts.ipynb)

Figure 2. Means and 95% confidence intervals of values of nucleotide diversity (Nei's π) within

each species at synonymous (yellow) and nonsynonymous (purple) sites.

Figure 3. Pairwise correlation of values of Nei's π between all species. Values above the

diagonal represent the correlation of π at synonymous sites, values below the diagonal represent

non-synonymous sites. Black boxes represent within-genus comparisons.

Supporting Information

Supplemental Tables

Table S1. Coverage summary statistics for whole-genome reads used to design sequence capture

array. For each library, values represent the name of the sequenced individual, the number of

reads in the sequenced library, the number of reads that mapped to the *Populus trichocarpa* v3

reference genome, the proportion of reads that mapped to the reference genome, and the mean

and standard deviation of read depth.

Table S2. Distribution of probes across the *Salix purpurea* genome.

Table S3. Summary of read depth at on- and off-target sites. For each library values represent

the 5%, 25%, 50%, 75%, and 99% quantiles of read depth, the maximum number of reads

mapped to a site, and the mean and standard deviation of read depth.

Table S4. Nucleotide diversity expressed as Nei's π for nonsynonymous and synonymous sites.

Supplemental Figures

Figure S1. Alternate bipartitions for the three species of *Populus*, based on gene tree

concordance. The cladogram in all three panels is that of the ASTRAL-III species tree (Figure

1), and the blue color represents the bipartition supported by the indicated number of gene trees

in each panel. **A)** 158 gene trees support the displayed ASTRAL-III species tree topology. **B)**

114 gene trees support a bipartition that places *P. tremula* and *P. mexicana* together. **C)** 94 gene

trees support a bipartition that places *P. balsamifera* and *P. mexicana* together.

Figure S2. Alternate bipartitions for the three species of *Salix*, based on gene tree concordance.

The cladogram in all three panels is that of the ASTRAL-III species tree (Figure 1), and the blue

color represents the bipartition supported by the indicated number of gene trees in each panel. **A)**

327 gene trees support the displayed ASTRAL-III species tree topology. **B)** 44 gene trees and **C)**

39 gene trees place one of the *S. phlebophylla* individuals within *S. exigua*.

Figure S3. Species trees estimated for all genes and known paralogs. **A)** Species tree generated

by ASTRAL-III for the gene trees. Node values represent bootstrap support from 100 multilocus

bootstrap replicates in ASTRAL-III. Branch lengths represent coalescent units. **B)** Cladogram

showing the congruence of gene trees for all nodes in the ASTRAL-III species tree. The numbers

above each node represent the number of gene trees that support the displayed bipartition, and

numbers below the node represent the number of gene trees that support all alternate bipartitions.

Purple wedges represent the proportion of gene trees that support the displayed bipartition. Blue

wedges represent the proportion of gene trees that support a single alternative bipartition (see

Appendices 3 & 4). Green wedges represent the proportion of gene trees that have multiple

conflicting bipartitions. Yellow wedges represent the proportion of gene trees that have no

supported bipartition. Plotting code and its interpretation were provided by Matt Johnson (for

more detail, see:

https://github.com/mossmatters/MJPythonNotebooks/blob/master/PhyParts_PieCharts.ipynb)

Figure S4. Distributions of values of nucleotide diversity (Nei's π) within each species at synonymous (yellow) and nonsynonymous (purple) sites.