Microbial helpers allow cyanobacteria to thrive in ferruginous waters

Nadia Szeinbaum*[1], Yael Toporek[2], Christopher T. Reinhard[1], and Jennifer B. Glass*[1]

[1]School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta, GA

[2]School of Biological Sciences, Georgia Institute of Technology, Atlanta, GA

*Corresponding authors: nadia.szeinbaum@biology.gatech.edu; jennifer.glass@eas.gatech.edu

Running title: Cyanobacterial fitness in early oceans

Acknowledgements. This research was funded by NASA Exobiology grant NNX14AJ87G,

NASA Astrobiology Institute grant 13-13NAI7_2-0027, and a NASA Astrobiology Postdoctoral

Fellowship to NS. We thank Jennifer Thweatt for advice on culturing *Synechococcus* sp. PCC

7002. We thank Amit Reddi for allowing us to use his plate reading spectrometer. We thank

Sommer Durham for technical assistance with the cytometer. We thank Patricia Sanchez-

Baracaldo and Manuel Kleiner for helpful discussions.

Competing Interests: The authors declare no competing interests.

Microbial helpers allow cyanobacteria to thrive in ferruginous waters

Abstract

The Great Oxidation Event (GOE) was a rapid accumulation of oxygen in the atmosphere as a

result of the photosynthetic activity of cyanobacteria. This accumulation reflected the

pervasiveness of O_2 on the planet's surface, indicating that cyanobacteria had become

ecologically successful in Archean oceans. Micromolar concentrations of Fe^{2+} in Archean oceans

would have reacted with hydrogen peroxide, a byproduct of oxygenic photosynthesis, to produce

hydroxyl radicals, which cause cellular damage. Yet cyanobacteria colonized Archean oceans

extensively enough to oxygenate the atmosphere, which likely required protection mechanisms

against the negative impacts of hydroxyl radical production in Fe^{2+}-rich seas. We identify several

factors that could have acted to protect early cyanobacteria from the impacts of hydroxyl radical

production and hypothesize that microbial cooperation may have played an important role in

protecting cyanobacteria from Fe^{2+} toxicity before the GOE. We found that several strains of

facultative anaerobic heterotrophic bacteria (*Shewanella*) with ROS defense mechanisms

increase the fitness of cyanobacteria (*Synechococcus*) in ferruginous waters. *Shewanella* species

with manganese transporters provided the most protection. Our results suggest that a tightly

regulated response to prevent Fe^{2+} toxicity could have been important for the colonization of

ancient ferruginous oceans, particularly in the presence of high manganese concentrations, and

may expand the upper bound for tolerable Fe^{2+} concentrations for cyanobacteria.

Microbial helpers allow cyanobacteria to thrive in ferruginous waters

Introduction

Earth's first biogeochemical cycles were driven by anaerobic microorganisms (Canfield

et al., 2006; Martin et al., 2018). At around 2.3 Ga, the Great Oxidation Event (GOE) resulted in

the initial oxygenation of the atmosphere and surficial biosphere, which ultimately led to the

modern dominance of aerobic organisms on Earth's surface (Bar-On et al., 2018; Luo et al.,

2016). Although biological O_2 production was a prerequisite for the GOE (Haqq-Misra et al.,

2011; Holland, 2002), oxygenic photosynthesis may have emerged in cyanobacteria hundreds of

millions of years prior to the initial accumulation of O_2 in Earth's atmosphere (Cardona et al.,

2019; Lalonde and Konhauser, 2015; Ossa Ossa et al., 2018; Planavsky et al., 2014). The delay

between the emergence of cyanobacterial O_2 production and O_2 accumulation in the atmosphere

may have been modulated by geophysical drivers (Catling et al., 2001; Holland, 2009; Lee et al.,

2016), but may also reflect the time required for metabolic innovations to appear in early

cyanobacteria or for the emergence of ecological linkages with other microbes facilitating the

success of cyanobacteria (Blank and Sanchez-Baracaldo, 2010; Johnston et al., 2009; Lyons et

al., 2014; Ozaki et al., 2019). Understanding how cyanobacteria cooperated with other microbes

to colonize the Earth's surface is thus essential to understand the ecology and tempo of the GOE.

The emergence of oxygenic photosynthesis in cyanobacteria occurred in the Archean

(Chisholm, 2017; Kendall et al., 2010; Konhauser et al., 2011; Lalonde and Konhauser, 2015;

Lyons et al., 2014; Olson et al., 2013; Planavsky et al., 2014; Reinhard et al., 2013b). The

metabolic expansion of cyanobacteria before the GOE may reflect their transition from land to

Fe^{2+}-rich Archean oceans (Herrmann and Gehringer, 2019). This transition would have been

physiologically challenging due to Fe^{2+} toxicity from its reactions with reactive oxygen species

(ROS) produced during photosynthesis (Swanner et al., 2015a). Archean oceans likely contained

tens to hundreds of micromolar Fe^{2+} within the ocean interior (Canfield, 2005; Derry, 2015;

Drever, 1974; Holland, 1973; Song et al., 2017), which would have reacted rapidly in the surface

ocean with O_2 produced from photosynthesis and any hydrogen peroxide (H_2O_2) from

photochemical reactions between O_2 and dissolved organic matter, as well as enzymes like

superoxide dismutase (Hansel and Diaz, 2020; Zinser, 2018b). This Fe^{2+}-driven reaction, known

as the Fenton reaction, produces hydroxyl radicals ($\cdot$OH; **Eq. 1**):

65                       $H_2O_2 + Fe^{2+} \rightarrow Fe^{3+} + OH^- + \cdot OH$ **(Eq. 1)**

Hydroxyl radicals cause cellular damage, especially to DNA (Imlay, 2003; Imlay, 2008).

Toxicity of Fe^{2+} may have prevented cyanobacteria from extensively colonizing Archean oceans

(Swanner et al., 2015a). Yet, cyanobacteria must have colonized vast areas of the ocean in order

to oxygenate the atmosphere. This paradox can be resolved by considering that anti-oxidants

such as elevated dissolved manganese (Mn^{2+}) and ancient Mn-based catalases may have

protected ancient cyanobacteria against ROS toxicity (Fischer et al., 2016).

Here, we test the hypothesis that heterotrophic microbial "helpers" may have protected

cyanobacteria from ROS produced by Fenton chemistry in Archean oceans, thereby increasing

cyanobacterial fitness and enabling their ecological success. Such microbial cooperation is

common among modern cyanobacteria and heterotrophic proteobacteria (Christie-Oleza et al.,

2017; Morris et al., 2011; Morris et al., 2008; Zinser, 2018a), whose intimate relationship is

evidenced by extensive horizontal gene transfer (Ben Said and Or, 2017; Braakman et al., 2017;

Goldenfeld and Woese, 2011). At the time of the GOE, many bacterial lineages, including

Proteobacteria, had already diversified (Battistuzzi et al., 2004; Cavalier-Smith, 2006a; Cavalier-

Smith, 2006b), which would have increased the phenotypic pool available for cooperation.

Including microbial cooperation as an ecological mechanism in models of early Earth's

ecological history might provide a more realistic picture of the ancient interactions that

ultimately led to the GOE.

We explored whether the presence of "helper" heterotrophic proteobacteria leads to

increased fitness of cyanobacteria in ferruginous conditions. For a model cyanobacterium, we

chose *Synechococcus* sp. PCC 7002 (hereafter *Synechococcus*), which was previously shown to

experience Fe^{2+} toxicity at >100 µM Fe^{2+} associated with increased intracellular ROS production

(Swanner et al., 2015a). As potential "helper" bacteria, we chose *Shewanella*, facultative

anaerobic gammaproteobacteria that can survive O_2 intrusions in the presence of high Fe^{2+} using

diverse H_2O_2-scavenging enzymes (Jiang et al., 2014; Mishra and Imlay, 2012; Sekar et al.,

2016). Experimental conditions simulated a pre-GOE illuminated ferruginous surface ocean

overlain by a CO_2- and H_2-rich anoxic atmosphere. We found that several *Shewanella* species

allowed *Synechococcus* to grow in ferruginous conditions that significantly impaired growth of

Synechococcus monocultures. The "helper" *Shewanella* strains all contained the ability to

actively uptake dissolved manganese (Mn^{2+}) via the natural resistance-associated macrophage

protein (NRAMP) family MntH Mn^{2+} transporter, a strategy that has previously been shown to

correlate with ROS survival (Daly et al., 2004). Our results stress the importance of considering

microbial cooperation and alternative ROS strategies, such as manganese protection, in models

of early Earth microbial ecology.

Results

Cyanobacteria growth is impaired in ferruginous conditions and is restored in the

presence of some proteobacteria. We found that *Synechococcus* growth in the presence of

elevated Fe^{2+} improved (to a varying degree) in the presence of all *Shewanella* spp. tested. In

monoculture, *Synechococcus* had similar growth rate and yield at 25 and 500 μM Fe^{2+}, but a

longer lag period at 500 μM Fe^{2+} (~2 days) than at 25 μM Fe^{2+} (~1 day; **Fig. 1A**). At 1000 μM

Fe^{2+}, *Synechococcus* growth was significantly impaired in monoculture, reaching only 10% the

cell density of cultures with 25 and 500 μM Fe^{2+} **(Fig. 1A)**. In contrast, *Synechococcus* growth

was minimally suppressed in the presence of high Fe^{2+} when grown in co-culture with

Shewanella baltica OS-155, although the initial lag period was extended **(Fig. 1B)**. In the

presence of *Shewanella algae* MN-01 **(Fig. 1C)** and *Shewanella loihica* PV-4 **(Fig. 1D)**,

Synechococcus growth was slightly suppressed at 500 and 1000 μM Fe^{2+}. Other than an extended

lag phase, *Shewanella algae* BrY **(Fig. 1E)** and *Shewanella oneidensis* MR-1 **(Fig. 1F)** had

minimal influence on *Synechococcus* growth, compared to the monoculture **(Fig. 1A)**, in all three

Fe^{2+} treatments. Although difficult to quantify due to spectral interference of Fe(III) oxide

particles, *Shewanella* cell numbers declined throughout the experiment (data not shown).

The best proteobacterial helpers are the least H_2O_2 sensitive, and the best H_2O_2

scavengers. We measured growth and H_2O_2 scavenging rates of *Shewanella* spp. in the presence

of varying H_2O_2. *S. baltica* OS-155 was the least sensitive to H_2O_2 **(Fig. 2A)**. *S. algae* MN-01

(Fig. 2B), *S. loihica* PV-4 **(Fig. 2C)**, and *S. algae* BrY **(Fig. 2D)** were moderately sensitive to

H_2O_2. *S. oneidensis* MR-01 was the most sensitive to H_2O_2 **(Fig. 2D)**. Along with being most

H_2O_2 tolerant, *S. algae* MN-01 and *S. baltica* OS-155 had the highest rates of H_2O_2 scavenging

activity, followed by *S. algae* BrY **(Fig. 3)**. *S. loihica* PV-4 and *S. oneidensis* MR-1 had the

lowest H_2O_2 scavenging rates **(Fig. 3)**.

Manganese may protect cyanobacteria from Fe^{2+} toxicity. To test whether Mn^{2+} can

protect cyanobacteria from Fe^{2+} toxicity, we grew *Synechococcus* PCC 7002 (four replicates per

treatment) under anoxic conditions with the addition of 1 mM Fe^{2+} and/or 1 mM Mn^{2+}. Cells

with 1 mM Mn^{2+} grew similarly to the controls **(Fig. 4)**. The A+ medium contained background

concentrations of ~140 μM Fe^{2+} and ~220 μM Mn^{2+}. No growth occurred with 1 mM Fe^{2+}. Red

Fe(III) oxide particles indicated that Fe^{2+} had been oxidized and precipitated, as observed by

Swanner et al. (2015b). Treatments with 1 mM Fe^{2+} and 1 mM Mn^{2+} resembled 1 mM Fe^{2+}

treatments for approximately the first week. Between 4-13 days, two out of four of the Fe^{2+} and

Mn^{2+} treatments grew to maximal OD_{750}. These results show that 1 mM Mn^{2+} is not toxic to

cyanobacteria and may in fact aid in the survival of cyanobacteria Fe^{2+} toxicity, after an

acclimation period. The mechanism that rescued growth of two out of four of the cultures to

grow in the presence of high Fe^{2+} after an extended lag phase remains unknown.

The best proteobacterial helpers encode additional genes for H_2O_2 degradation.

Synechococcus PCC 7002's susceptibility to Fe^{2+} toxicity is consistent with the limited number

of catalase genes in its genome; it encodes cytoplasmic KatG but not periplasmic KatE **(Table**

1). Without KatE to scavenge H_2O_2 in the periplasm, H_2O_2 can react with Fe^{2+} to generate ·OH

intracellularly **(Eq. 1).** Like *Synechococcus* PCC 7002, most marine cyanobacteria are KatE-

negative; a BLAST search of cyanobacterial genomes in NCBI recovered KatE catalase

homologs almost exclusively in freshwater and soil cyanobacteria (**Table S1**).

To identify genes in *Shewanella* that may have helped alleviate Fe^{2+} toxicity to

Synechococcus, we compared the genomes of the *Shewanella* strains in our experiments.

Notably, several *Shewanella* spp. contained catalases predicted to have multiple cellular

locations, as previously observed for other microbial catalases (Hanaoka et al., 2013). Overall,

the genomic inventory of catalase and peroxidase proteins was generally similar between the

more protective and less protective species (**Table 1)**, suggesting (an)other mechanism(s) for

ROS survival. We found 52 proteins in the best helpers (OS-155 and MN-01) that were not

present in the other *Shewanella* strains, including genes for flagella, phenazine biosynthesis, and

transporters **(Table S3)**. Flagella may be involved in the ROS-stress response in eukaryotes

(Hajam et al., 2017), but their connection to ROS protection in bacteria, if any, is unknown.

Phenazines are known to produce oxidative stress (Imlay, 2013), and can also mediate

extracellular redox transfers (Hernandez et al., 2004; Wang and Newman, 2008), but are unlikely

to be responsible for the protective effect because *Synechococcus* PCC 7002 also possesses the

phzF gene for phenazine synthesis.

The high H_2O_2 sensitivity of *S. oneidensis* MR-1, which contains a similar repertoire of

H_2O_2-scavenging enzymes as less H_2O_2-sensitive *Shewanella* spp., is thought to be due to its

inability to actively transport and accumulate intracellular Mn^{2+} (Daly et al., 2004; Jiang et al.,

2014). We surmised that differences in ROS scavenging rates between *Shewanella* strains may

be due to differences in acquisition of Mn^{2+}. We found MntH Mn^{2+} transporters in the genomes

of the three top *Shewanella* helpers: OS-155, MN-01, and PV-4 (**Table 1**). *Shewanella* BrY and

MR-1 lacked characterized Mn^{2+} transporters. *Synechococcus* PCC 7002 contained genes for the

ATP-binding cassette (ABC) family Mn^{2+} transporters MntABCD and SitABCD transporter

instead of MntH.

Discussion

The rise of O_2 and ROS from oxygenic photosynthesis would have severely stressed

strictly anaerobic microbes (Khademian and Imlay, 2020), resulting in what was perhaps Earth's

first mass extinction. Experiments demonstrating that catalase-negative cyanobacteria

(*Synechococcus* PCC 7002) grew poorly in >100 μM Fe^{2+} led to the idea that Fe^+-rich oceans

would have slowed cyanobacterial colonization of the ocean surface and possibly delayed global

oxygenation (Swanner et al., 2015a). Our study confirms the previous finding that

Synechococcus PCC 7002, originally isolated from marine mud, has impaired growth when Fe^{2+}

was 180 μM and higher. We show that this Fe^{2+} toxicity can be alleviated by some strains of

"helper" *Shewanella* spp., with the best protection afforded by *Shewanella* strains possessing the

most varied sets of ROS-defense pathways (e.g. catalases, MntH transporters) and the highest

rates of H_2O_2 degradation. Likely, this protection was afforded by *Shewanella* scavenging H_2O_2

prior to its reaction with Fe^{2+}, thereby decreasing the production of damaging hydroxyl radicals.

Thus, our findings align with previous findings (Brown et al., 2010; Ward et al., 2019)

that cyanobacterial colonization of early oceans would not have been hampered by micromolar

Fe^{2+} concentrations, if Mn^{2+}-transporting and H_2O_2-scavenging genes were present within the

microbial communal gene pool. Early marine cyanobacteria, like modern terrestrial

cyanobacteria, likely had myriad protections against Fe^{2+} and/or may have benefitted from the

presence of co-existing "helper" bacteria to cope with the harmful byproducts produced by their

own metabolism in a ferruginous ocean, which would have later been lost due to genome

streamlining in marine cyanobacteria.

Catalase-based protection. The ubiquity of catalase genes in the genomes of all the

Shewanella strains we studied suggests that catalase accounts for the background protection

provided by all *Shewanella* spp. tested. The enhanced protection provided by *Shewanella* spp.

with similar catalase inventories implies that a mechanism other than catalase was likely at play,

presumably at the level of gene expression. This process may also be related to the centralized

regulation of H_2O_2-related genes in *Shewanella* spp. In *S. oneidensis* MR-1, the transcriptional

regulator OxyR is key for suppression of Fenton chemistry by derepression of KatE and Dps

(Jiang et al., 2014; Wan et al., 2018) whereas H_2O_2-based regulation is performed by multiple

regulators (including PerR) in *Synechococcus* PCC 7002 and other cyanobacteria (Latifi et al.,

2009).

A protective effect of proteobacterial catalase has previously been observed for the

marine cyanobacterium *Prochlorococcus,* which grows in symbiosis with the

gammaproteobacterium *Alteromonas* (Biller et al., 2016; Morris et al., 2011; Morris et al., 2008).

(For more examples of microbe-microbe H_2O_2 protection, see review by Zinser (2018a).) Yet,

unlike those long-lived catalase-based symbioses, the presence of *Shewanella* in our co-cultures

was ultimately transient. A transient cooperative interaction can be a natural consequence of the

dynamics of public goods between microbial groups (Corno et al., 2013; Rodríguez Amor and

Dal Bello, 2019), when the members of the population that provide the extracellular "public

good" are at a fitness disadvantage by providing a costly product that nonproducers use (Cremer

et al., 2019; Özkaya et al., 2017). In our system, *Synechococcus* gained the fitness advantage of

protection from Fe^{2+} toxicity at the expense of *Shewanella*, whose population was eliminated

from the system as *Synechococcus* grew. Indeed, previous attempts to co-culture *Shewanella*

with cyanobacteria with ~15 μM Fe^{2+}, resulted in cyanobacterial dominance, with *Shewanella*'s

growth yield compromised by the presence of *Synechococcus* sp. 7002 even in the presence of

organic carbon (Beliaev et al., 2014).

Thus, our co-culture experiments illustrate that cyanobacteria can benefit from the

presence of "helper" proteobacteria under ferruginous conditions. This protection may have been

one of the ways that cyanobacteria were able to cope with the harmful byproducts produced by

their own metabolism as they incipiently colonized a ferruginous ocean, which would have no

longer been necessary once cyanobacteria increased in numbers and seawater Fe^{2+} concentrations

dropped. The precise levels of dissolved O_2 prevailing on different spatial scales in the surface

ocean prior to the GOE are not fully known. However, there is theoretical evidence to suggest

that dissolved O_2 would have been locally more than sufficient to support aerobic bacterial

respiration (Olson et al., 2013; Reinhard et al., 2013a).

223   *Manganese to the rescue*. One of the genes regulated by OxyR is the Mn^{2+} transporter

MntH, which is used for accumulation of intracellular manganese (Mn^{2+}) as a potent ROS

detoxification method (Anjem et al., 2009; Chen et al., 2008; Kehres et al., 2002). Unlike Fe^{2+},

Mn^{2+} does not undergo Fenton-type reactions. Instead, Mn^{2+} has strong antioxidant properties

(Cheton and Archibald, 1988) and is highly effective at protecting against H_2O_2-induced

oxidative stress through multiple mechanisms (Aguirre and Culotta, 2012; Hansel, 2017;

Horsburgh et al., 2002; Latour, 2015; Papp-Wallace and Maguire, 2006). Mn^{2+}-carbonate and

Mn^{2+}-phosphate complexes can chemically disproportionate H_2O_2 (Archibald and Fridovich,

1982; Barnese et al., 2012; Stadtman et al., 1990). Mn^{2+}-containing catalase, a very ancient

member of the ferritin superfamily, detoxifies H_2O_2 (Klotz and Loewen, 2003; Zamocky et al.,

2008). Under H_2O_2 stress, OxyR facilitates Mn^{2+} replacement of Fe^{2+} in ROS-sensitive enzymes,

preventing their inactivation by Fenton chemistry (Anjem et al., 2009; Smethurst et al., 2020;

Sobota and Imlay, 2011).

236   MntH transporters were found in the most protective *Shewanella* strains, but not in

Synechococcus PCC 7002 (which instead encodes two ABC-type Mn^{2+} transporters) nor in the

less protective *Shewanella* spp. **(Table S1)**. In monocultures, the rescued growth of

Synechococcus PCC 7002 in two out of four of high Fe^{2+} treatments with Mn^{2+} was likely related

to the antioxidant properties of Mn^{2+}, although the details of the protective mechanism, chemical

or enzymatic, await further study. Our findings generally support the hypothesis that elevated

seawater Mn^{2+} in early Earth environments (~5-120 μM; Holland, 1984; Johnson et al., 2016;

Komiya et al., 2008; Liu et al., 2020) played a role in protecting marine cyanobacteria from Fe^{2+}

toxicity (Fischer et al., 2016).

245   *Modern microbial models for ancient physiologies.* The choice of a model

cyanobacterium for physiological experiments applicable to the Precambrian oceans is of great

importance (Hamilton, 2019; Hamilton et al., 2016). Many terrestrial cyanobacteria thrive under

the 10-100 μM Fe^{2+} concentrations predicted for Archean oceans (Brown et al., 2005; Ionescu et

al., 2014; Ward et al., 2019; Ward et al., 2017) and either possess multiple catalases **(Table S1)**

and/or have novel defense mechanisms such as intracellular iron precipitation (Brown et al.,

2010). In contrast, modern marine cyanobacteria (e.g. *Prochlorococcus*) tend to be genetically

streamlined for specific modern oceanographic provinces (Partensky and Garczarek, 2010),

which are extremely Fe^{2+}-poor compared to modern terrestrial and ancient ecosystems.

The closest modern descendants of the ancestral cyanobacteria that evolved into modern

marine plankton cyanobacteria are filamentous non-heterocystous *Synechococcales* (Sánchez-

Baracaldo, 2015; Sánchez☐Baracaldo and Cardona, 2020). KatG was likely present in ancestors

of marine cyanobacteria (Bernroitner et al., 2009; Zamocky et al., 2012), whereas KatE was

likely horizontally transferred from Proteobacteria and Planctomycetes to some cyanobacterial

linages (e.g. Nostocales; Zamocky et al., 2012). *Synechococcus* PCC 7002 lacks Mn^{2+}-catalase,

which is widespread in terrestrial cyanobacteria (Ballal et al., 2020; Banerjee et al., 2012; Bihani

et al., 2016; Chakravarty et al., 2016; Chen et al., 2020) **(Table S1)** and was likely present in

early cyanobacterial lineages (Klotz and Loewen, 2003; Zamocky et al., 2012).

Previous genetic studies of Fe^{2+}-induced oxidative stress have studied cyanobacteria that

cannot cope with high Fe^{2+} and H_2O_2, e.g. *Synechocystis* PCC 6803 (Li et al., 2004; Shcolnick et

al., 2009) in monoculture. In nature, ROS and O_2 cycling are communal processes. Thus, models

that include shared mechanisms of survival are important to consider on the early Earth,

particularly as gene pools were more limited and were in the process of expansion. We advocate

for future studies on more deeply branching cyanobacterial species with additional ROS defense

mechanisms and encourage more explicit incorporation of microbial interactions in large-scale

models of biogeochemical cycling on the ancient Earth.

Materials and Methods

Bacterial strains. *Synechococcus* sp. PCC 7002 was ordered from the Pasteur Culture

collection of Cyanobacteria. *Shewanella oneidensis* MR-1 and *Shewanella algae* BrY were kind

gifts from the lab of Dr. Thomas DiChristina (Georgia Institute of Technology). *Shewanella*

loihica PV-4 was a kind gift from Dr. Jeffrey Gralnick (University of Minnesota).

Experimental setup and growth conditions. *Synechococcus* sp. PCC 7002 was grown in

serum bottles containing modified A+ medium (Stevens Jr. et al., 1973) with 10 g L^{-1} NaCl,

TRIS buffer (pH 7.2), and 10 mM NH_4^+ as the nitrogen source. *Shewanella* spp. were grown

overnight in lysogeny broth (LB, 10 g L^{-1} NaCl, 10 g L^{-1} tryptone, 5 g L^{-1} yeast extract) and

transferred into serum bottles containing modified A+ medium with amino acids (20 mg L^{-1} L-

serine, 20 mg L^{-1} L-arginine, and 20 mg L^{-1} L-glutamic acid), 20 mM lactate as electron donor,

and 20 mM fumarate as electron acceptor. Bottles were flushed with 90% N$_2$/10% CO$_2$ and

opened inside an anoxic chamber (5% CO$_2$/4% H$_2$/91% N$_2$). Cultures were washed with anoxic

A+ medium and combined at optical density of 600 nm (OD$_{600}$) = 0.01. Co-cultures were grown

in triplicate in 10 mL-well tissue culture plates inside the anoxic chamber (5% CO$_2$/4% H$_2$/91%

N$_2$). Cultures were mixed daily by gentle pipetting ~50% of the volume three times in order to

resuspend cells and particulate Fe(III) oxides; if not mixed regularly, PCC 7002 would grow at

the bottom of the well. Light was provided with a fluorescent light in a 12:12 light:dark timer-

controlled cycle. FeCl$_2$ was added at a final concentration of 25, 500, or 1000 μM.

Cyanobacterial quantification by flow cytometry. Cell numbers of *Synechococcus* sp.

PCC 7002 were quantified in an LSR Fortessa flow cytometer using FACSDiva^{TM} (BD

Biosciences, CA). At each time point, 200 μL of culture was loaded into a 96-well plate inside

the anoxic chamber, covered in parafilm to minimize Fe^{2+} oxidation, transported to the

cytometer, and mixed twice in the cytometer. Samples (10 μL) were injected and run at a rate of

0.5 μL s^{-1}. Cyanobacteria were detected by phycocyanin/chlorophyll autofluorescence using blue

(488 nm) and yellow-green lasers (561 nm) measured at 655-684 nm (Hill et al., 2017).

Optimization and calibration of the quantification parameters were achieved using yellow-green

1 μm microspheres (441/485 ex/em; Polysciences, PA). Live cyanobacteria were also quantified

using Syto9 using FITC filters. Events above the thresholds of PerCP and FITC were considered

live cyanobacteria. Propidium iodide could not be used to identify "dead' cyanobacteria, as the

emission spectra overlapped with that of their autofluorescence. Due to spectral overlap with iron

particles, *Shewanella* cells could not be accurately quantified by cytometry.

H$_2$O$_2$ resistance assays. Six *Shewanella* strains were incubated in 96-well plates with

minimal M1 media (Myers and Nealson, 1988) with lactate (10 mM) or acetate (10 mM) as

electron donor under oxic conditions. Growth was monitored periodically (every 1-2 hours) by

OD$_{600}$ in a spectrophotometer with plate-reading capacity (Tecan, Switzerland). Hydrogen

peroxide was added after an initial period of growth for 4 hours, at a final concentration of 10,

25, 50, 100, or 250 μM, after which growth continued to be monitored.

H$_2$O$_2$ scavenging assays. We compared the abilities of the *Shewanella* spp. to remove

H$_2$O$_2$ from their environment in cell suspensions. Strains were seeded in lysogeny broth (LB, 10

g L^{-1} NaCl, 10 g L^{-1} tryptone, 5 g L^{-1} yeast extract, Sigma-Aldrich) at 30ºC with shaking

overnight, harvested by centrifugation at 12,300 x g, washed, and transferred into minimal M1

medium amended with 20 mM lactate at $OD_{600} = 0.02$. Cells were incubated at 30ºC with

shaking until harvesting at mid-log phase ($OD_{600} = 0.15$-0.35), washed twice with minimal

medium, then inoculated at $OD_{600} = 0.05$ into a 24-well plate holding 2 mL minimal M1 medium

amended with 20 mM lactate and various concentrations of H_2O_2 (0-5000 μM). Samples were

collected every 3-5 minutes and analyzed immediately for exogenous H_2O_2 using the resorufin-

horseradish peroxidase colorimetric assay (Zhou et al., 1997). Plates were incubated under oxic

conditions at room temperature with shaking for the duration of the experiment (30-200

min). H_2O_2 disappearance followed an exponential decay **(Eq. 2)**. First-order apparent rate

constants (k) were obtained by plotting the data as shown in **Eq. 3**, where k is the slope of the

graph with $\ln[H_2O_{2(t=0)}/H_2O_{2(t=n)}]$ on the y-axis and time on the x-axis.

$$H_2O_{2(t=n)} = H_2O_{2(t=0)}\ e^{-kt} \qquad \textbf{(Eq. 2)}$$

$$\ln[H_2O_{2(t=0)}/H_2O_{2(t=n)}] = kt \qquad \textbf{(Eq. 3)}$$

Unique proteins. Proteins present in *Shewanella algae* MN-01 and *Shewanella baltica*

OS155 and absent in *Shewanella algae* BrY, *Shewanella oneidensis* MR-1, and *Shewanella*

loihica PV-4 were identified using the Protein Families tool in PATRIC using three protein

family databases: PATRIC cross-genus families (PGfams), PATRIC genus-specific families

(PLfams), and FIGfam.

Synechococcus monoculture experiments. To determine the influence of 1 mM Mn^{2+} on

Synechococcus PCC 7002 growth with and without 1 mM Fe^{2+}, *Synechococcus* was grown in

modified A+ medium containing 82 mM bicarbonate in Hungate tubes with bromobutyl rubber

stoppers containing 95% N_2/5% H_2 headspace with constant shaking at 200 rpm under constant

light. Growth was determined by measurement of optical density at 750 nm (OD_{750}).

Table 1. Locus tags of ROS response proteins in the bacterial species in this study. None
contained Mn-catalase or Ni-superoxide dismutase. "—" indicates no homolog in genome.
OxyR: hydrogen peroxide inducible gene activator; KatG: catalase-peroxidase (clade 1); KatE:
periplasmic catalase (clade 3); AhpC: alkyl hydroperoxide reductase; Dps/MrgA: DNA-binding
ferritin-like protein; BtuE: glutathione peroxidase; SitA and MntA: ABC-type Mn^{2+} transporters;
MntH: NRAMP-type MntH Mn^{2+} transporter. KatG cellular localization based on PSORTb (Yu
et al., 2010): C: cytoplasmic; U: unknown.

Species	Locus prefix	OxyR or PerR	KatG	KatE	AhpC	Dps/ MrgA	BtuE	MntA	SitA	MntH
Synecho- coccus **PCC 7002**	**SYN PCC 7002_A**	1836	2422[C]	—	0558	0031	0117	1734	2501	—
Shewanella baltica **OS155**	**Sbal_**	1181	0875[C]	0894	0849	3285	1384	—	—	0678
Shewanella algae **MN-01**	**AMR44_**	05180	15815[U] 20550[C]	04925	RS11285	RS07 720	08935	—	—	05235
Shewanella loihica **PV-4**	**Shew_**	1035	0709[U]	3190	0792	—	2741	—	—	2965
Shewanella oneidensis **MR-1**	**SO_**	1328	0725[U] 4405[C]	1070	0958	1158	1563 3349	—	—	—
Shewanella algae **BrY**	**BFS86_**	05230	17460[U]	16195	00635	RS11 070	19140	—	—	—

Figure 1. Growth of *Synechococcus* PCC 7002 in mono- or co-culture with *Shewanella* spp. with varying Fe^{2+}. Co-cultures are: A) none, B) *Shewanella baltica* OS-155, C) *Shewanella algae* MN-01, D) *Shewanella loihica* PV-4, E) *Shewanella algae* BrY, F) *Shewanella oneidensis* MR-1. Error bars represent the standard error of the mean (n=3).

350
351 **Figure 2. Growth of *Shewanella* spp. with varying H$_2$O$_2$.** A) *Shewanella baltica* OS-155, B)
352 *Shewanella algae* MN-01, C) *Shewanella loihica* PV-4, D) *Shewanella algae* BrY, E)
353 *Shewanella oneidensis* MR-1. Error bars represent the standard error of the mean (n=3 for all
354 except *S. baltica* OS-155, n=2). H$_2$O$_2$ was added at the four-hour timepoint.

355
356 **Figure 3. H_2O_2 peroxide scavenging capacity of *Shewanella* spp. shown as the first order**
357 **rate constant plotted versus initial H_2O_2 concentration.** No change in H_2O_2 was observed in
358 the abiotic control.

Figure 4. Growth of *Synechococcus* sp. PCC 7002 in monoculture with 1 mM Fe^{2+} and/or 1 mM Mn^{2+}. Controls had background levels of ~140 μM Fe^{2+} and ~220 μM Mn^{2+}. The growth curve for the 1 mM Fe^{2+} + 1 mM Mn^{2+} treatment is shown for the two replicates (out of four) that grew. Photos were taken on day 13.

369   **Supplemental tables**

370   **Table S1. Cyanobacterial genomes containing MnKat, KatE, or KatG**. Appended as separate
371   spreadsheet.
372
373   **Table S2. Proteins that are present in *Shewanella algae* MN-01 and *Shewanella baltica***
374   **OS155 and are not present in *Shewanella oneidensis* MR-1, *Shewanella algae* BrY, and**
375   *Shewanella loihica* **PV-4.** Appended as separate spreadsheet.
376

References

Aguirre, J. D., & Culotta, V. C. (2012). Battles with iron: manganese in oxidative stress protection. Journal of Biological Chemistry, 287, 13541-13548.

Anjem, A., Varghese, S., & Imlay, J. A. (2009). Manganese import is a key element of the OxyR response to hydrogen peroxide in *Escherichia coli*. Molecular Microbiology, 72, 844-858.

Archibald, F. S., & Fridovich, I. (1982). The scavenging of superoxide radical by manganous complexes: in vitro. Archives of Biochemistry and Biophysics, 214, 452-463.

Ballal, A., Chakravarty, D., Bihani, S. C., & Banerjee, M. (2020). Gazing into the remarkable world of non-heme catalases through the window of the cyanobacterial Mn-catalase 'KatB'. Free Radical Biology and Medicine, 160, 480-487.

Banerjee, M., Ballal, A., & Apte, S. K. (2012). Mn☐catalase (Alr0998) protects the photosynthetic, nitrogen☐fixing cyanobacterium Anabaena PCC7120 from oxidative stress. Environmental Microbiology, 14, 2891-2900.

Bar-On, Y. M., Phillips, R., & Milo, R. (2018). The biomass distribution on Earth. Proceedings of the National Academy of Sciences, 115, 6506-6511.

Barnese, K., Gralla, E. B., Valentine, J. S., & Cabelli, D. E. (2012). Biologically relevant mechanism for catalytic superoxide removal by simple manganese compounds. Proceedings of the National Academy of Sciences, 109, 6892-6897.

Battistuzzi, F. U., Feijao, A., & Hedges, S. B. (2004). A genomic timescale of prokaryote evolution: insights into the origin of methanogenesis, phototrophy, and the colonization of land. BMC Evolutionary Biology, 4.

Beliaev, A. S., Romine, M. F., Serres, M., Bernstein, H. C., Linggi, B. E., Markillie, L. M., Isern, N. G., Chrisler, W. B., Kucek, L. A., Hill, E. A., Pinchuk, G. E., Bryant, D. A., Wiley, H. S., Fredrickson, J. K., & Konopka, A. (2014). Inference of interactions in cyanobacterial–heterotrophic co-cultures via transcriptome sequencing. The ISME Journal, 8, 2243-2255.

Ben Said, S., & Or, D. (2017). Synthetic microbial ecology: engineering habitats for modular consortia. Frontiers in Microbiology, 8, 1125.

Bernroitner, M., Zamocky, M., Furtmüller, P. G., Peschek, G. A., & Obinger, C. (2009). Occurrence, phylogeny, structure, and function of catalases and peroxidases in cyanobacteria. Journal of Experimental Botany, 60, 423-440.

Bihani, S. C., Chakravarty, D., & Ballal, A. (2016). KatB, a cyanobacterial Mn-catalase with unique active site configuration: Implications for enzyme function. Free Radical Biology and Medicine, 93, 118-129.

Biller, S. J., Coe, A., & Chisholm, S. W. (2016). Torn apart and reunited: impact of a heterotroph on the transcriptome of *Prochlorococcus*. The ISME Journal, 10, 2831-2843.

Blank, C. E., & Sanchez-Baracaldo, P. (2010). Timing of morphological and ecological innovations in the cyanobacteria - a key to understanding the rise in atmospheric oxygen. Geobiology, 8, 1-23.

Braakman, R., Follows, M. J., & Chisholm, S. W. (2017). Metabolic evolution and the self-organization of ecosystems. Proceedings of the National Academy of Sciences, 114, E3091-E3100.

Brown, I. I., Bryant, D. A., Casamatta, D., Thomas-Keprta, K. L., Sarkisova, S. A., Shen, G., Graham, J. E., Boyd, E. S., Peters, J. W., & Garrison, D. H. (2010). Polyphasic characterization of a thermotolerant siderophilic filamentous cyanobacterium that produces intracellular iron deposits. Applied and Environmental Microbiology, 76, 6664-6672.

Brown, I. I., Mummey, D., & Cooksey, K. E. (2005). A novel cyanobacterium exhibiting an elevated tolerance for iron. FEMS Microbiology Ecology, 52, 307-314.

Canfield, D. E. (2005). The early history of atmospheric oxygen: homage to Robert M. Garrels. Annual Review of Earth and Planetary Sciences, 33, 1-36.

Canfield, D. E., Rosing, M. T., & Bjerrum, C. (2006). Early anaerobic metabolisms. Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences, 361, 1819-1836.

Cardona, T., Sánchez-Baracaldo, P., Rutherford, A. W., & Larkum, A. W. (2019). Early Archean origin
of Photosystem II. Geobiology, 17, 127-150.
Catling, D. C., Zahnle, K. J., & Mckay, C. P. (2001). Biogenic methane, hydrogen escape, and the
Iirreversible oxidation of early Earth. Science, 293, 839-843.
Cavalier-Smith, T. (2006a). Cell evolution and Earth history: stasis and revolution. Philosophical
Transactions of the Royal Society of London. Series B, Biological Sciences, 361, 969-1006.
Cavalier-Smith, T. (2006b). Rooting the tree of life by transition analyses. Biology Direct, 1, 19.
Chakravarty, D., Banerjee, M., Bihani, S. C., & Ballal, A. (2016). A salt-inducible Mn-catalase (KatB)
protects cyanobacterium from oxidative stress. Plant Physiology, 170, 761-773.
Chen, H., Xu, G., Zhao, Y., Tian, B., Lu, H., Yu, X., Xu, Z., Ying, N., Hu, S., & Hua, Y. (2008). A novel
OxyR sensor and regulator of hydrogen peroxide stress with one cysteine residue in *Deinococcus*
radiodurans. PLoS ONE, 3, e1602.
Chen, M.-Y., Teng, W.-K., Zhao, L., Hu, C.-X., Zhou, Y.-K., Han, B.-P., Song, L.-R., & Shu, W.-S.
(2020). Comparative genomics reveals insights into cyanobacterial evolution and habitat
adaptation. The ISME Journal, 1-17.
Cheton, P. L., & Archibald, F. S. (1988). Manganese complexes and the generation and scavenging of
hydroxyl free radicals. Free Radical Biology and Medicine, 5, 325-333.
Chisholm, S. W. (2017). Prochlorococcus. Current Biology, 27, R447-R448.
Christie-Oleza, J. A., Sousoni, D., Lloyd, M., Armengaud, J., & Scanlan, D. J. (2017). Nutrient recycling
facilitates long-term stability of marine microbial phototroph–heterotroph interactions. Nature
Microbiology, 2, 17100.
Corno, G., Villiger, J., & Pernthaler, J. (2013). Coaggregation in a microbial predator–prey system affects
competition and trophic transfer efficiency. Ecology, 94, 870-881.
Cremer, J., Melbinger, A., Wienand, K., Henriquez, T., Jung, H., & Frey, E. (2019). Cooperation in
microbial populations: theory and experimental model systems. Journal of Molecular Biology,
431, 4599-4644.
Daly, M. J., Gaidamakova, E. K., Matrosova, V., Vasilenko, A., Zhai, M., Venkateswaran, A., Hess, M.,
Omelchenko, M., Kostandarithes, H. M., & Makarova, K. (2004). Accumulation of Mn (II) in
Deinococcus radiodurans facilitates gamma-radiation resistance. Science, 306, 1025-1028.
Derry, L. A. (2015). Causes and consequences of mid-Proterozoic anoxia. Geophysical Research Letters,
42, 8538-8546.
Drever, J. (1974). Geochemical model for the origin of Precambrian banded iron formations. GSA
Bulletin, 85, 1099-1106.
Fischer, W. W., Hemp, J., & Valentine, J. S. (2016). How did life survive Earth's great oxygenation?
Current Opinion in Chemical Biology, 31, 166-178.
Goldenfeld, N., & Woese, C. (2011). Life is physics: Evolution as a collective phenomenond far from
equilibrium. Annual Review of Condensed Matter Physics, 2, 375-399.
Hajam, I. A., Dar, P. A., Shahnawaz, I., Jaume, J. C., & Lee, J. H. (2017). Bacterial flagellin—a potent
immunomodulatory agent. Experimental & Molecular Medicine, 49, e373-e373.
Hamilton, T. L. (2019). The trouble with oxygen: The ecophysiology of extant phototrophs and
implications for the evolution of oxygenic photosynthesis. Free Radical Biology and Medicine,
140, 233-249.
Hamilton, T. L., Bryant, D. A., & Macalady, J. L. (2016). The role of biology in planetary evolution:
cyanobacterial primary production in low-oxygen Proterozoic oceans. Environmental
Microbiology, 18, 325-340.
Hanaoka, Y., Takebe, F., Nodasaka, Y., Hara, I., Matsuyama, H., & Yumoto, I. (2013). Growth-
dependent catalase localization in *Exiguobacterium oxidotolerans* T-2-2T reflected by catalase
activity of cells. PLoS ONE, 8, e76862.
Hansel, C. M. (2017). Manganese in marine microbiology. In: Advances in Microbial Physiology.
Elsevier, pp. 37-83.

Hansel, C. M., & Diaz, J. M. (2020). Production of extracellular reactive oxygen species by marine biota.
Annual Review of Marine Science, 13.
Haqq-Misra, J., Kasting, J. F., & Lee, S. (2011). Availability of O2 and H2O2 on pre-photosynthetic
Earth. Astrobiology, 11, 293-302.
Hernandez, M. E., Kappler, A., & Newman, D. K. (2004). Phenazines and Other Redox-Active
Antibiotics Promote Microbial Mineral Reduction. Applied and Environmental Microbiology, 70,
921-928.
Herrmann, A. J., & Gehringer, M. M. (2019). An investigation into the effects of increasing salinity on
photosynthesis in freshwater unicellular cyanobacteria during the late Archaean. Geobiology, 17,
343-359.
Hill, E. A., Chrisler, W. B., Beliaev, A. S., & Bernstein, H. C. (2017). A flexible microbial co-culture
platform for simultaneous utilization of methane and carbon dioxide from gas feedstocks.
Bioresource Technology, 228, 250-256.
Holland, H. D. (1973). The Oceans; A Possible Source of Iron in Iron-Formations. Economic Geology,
68, 1169-1172.
Holland, H. D. (1984). The chemical evolution of the atmosphere and oceans, Princeton University Press.
Holland, H. D. (2002). Volcanic gases, black smokers, and the great oxidation event. Geochimica et
Cosmochimica Acta, 66, 3811-3826.
Holland, H. D. (2009). Why the atmosphere became oxygenated: A proposal. Geochimica et
Cosmochimica Acta, 73, 5241-5255.
Horsburgh, M. J., Wharton, S. J., Karavolos, M., & Foster, S. J. (2002). Manganese: elemental defence
for a life with oxygen. Trends in Microbiology, 10, 496-501.
Imlay, J. A. (2003). Pathways of oxidative damage. Annual Reviews in Microbiology, 57, 395-418.
Imlay, J. A. (2008). Cellular defenses against superoxide and hydrogen peroxide. Annual Review of
Biochemistry, 77, 755-776.
Imlay, J. A. (2013). The molecular mechanisms and physiological consequences of oxidative stress:
lessons from a model bacterium. Nature Reviews Microbiology, 11, 443-454.
Ionescu, D., Buchmann, B., Heim, C., Häusler, S., De Beer, D., & Polerecky, L. (2014). Oxygenic
photosynthesis as a protection mechanism for cyanobacteria against iron-encrustation in
environments with high Fe2+ concentrations. Frontiers in Microbiology, 5, 459.
Jiang, Y., Dong, Y., Luo, Q., Li, N., Wu, G., & Gao, H. (2014). Protection from oxidative stress relies
mainly on derepression of OxyR-dependent KatB and Dps in *Shewanella oneidensis*. Journal of
Bacteriology, 196, 445-458.
Johnson, J. E., Webb, S. M., Ma, C., & Fischer, W. W. (2016). Manganese mineralogy and diagenesis in
the sedimentary rock record. Geochimica et Cosmochimica Acta, 173, 210-231.
Johnston, D. T., Wolfe-Simon, F., Pearson, A., & Knoll, A. H. (2009). Anoxygenic photosynthesis
modulated Proterozoic oxygen and sustained Earth's middle age. Proceedings of the National
Academy of Sciences, 106, 16925-16929.
Kehres, D. G., Janakiraman, A., Slauch, J. M., & Maguire, M. E. (2002). Regulation of *Salmonella*
enterica serovar Typhimurium *mntH* transcription by H2O2, Fe2+, and Mn2+. Journal of
Bacteriology, 184, 3151-3158.
Kendall, B., Reinhard, C. T., Lyons, T. W., Kaufman, A. J., Poulton, S. W., & Anbar, A. D. (2010).
Pervasive oxygenation along late Archaean ocean margins. Nature Geoscience, 3, 647–652.
Khademian, M., & Imlay, J. A. (2020). How microbes evolved to tolerate oxygen. Trends in
Microbiology.
Klotz, M. G., & Loewen, P. C. (2003). The molecular evolution of catalatic hydroperoxidases: evidence
for multiple lateral transfer of genes between prokaryota and from bacteria into eukaryota.
Molecular Biology and Evolution, 20, 1098-1112.
Komiya, T., Hirata, T., Kitajima, K., Yamamoto, S., Shibuya, T., Sawaki, Y., Ishikawa, T., Shu, D., Li,
Y., & Han, J. (2008). Evolution of the composition of seawater through geologic time, and its
influence on the evolution of life. Gondwana Research, 14, 159-174.

Konhauser, K. O., Lalonde, S. V., Planavsky, N. J., Pecoits, E., Lyons, T. W., Mojzsis, S. J., Rouxel, O.
J., Barley, M. E., Rosìere, C., Fralick, P. W., Kump, L. R., & Bekker, A. (2011). Aerobic
bacterial pyrite oxidation and acid rock drainage during the Great Oxidation Event. Nature, 478,
369-373.
Lalonde, S. V., & Konhauser, K. O. (2015). Benthic perspective on Earth's oldest evidence for oxygenic
photosynthesis. Proceedings of the National Academy of Sciences, 112, 995-1000.
Latifi, A., Ruiz, M., & Zhang, C.-C. (2009). Oxidative stress in cyanobacteria. FEMS Microbiology
Reviews, 33, 258-278.
Latour, J.-M. (2015). Manganese, the stress reliever. Metallomics, 7, 25-28.
Lee, Yeung, L. Y., Mckenzie, N. R., Yokoyama, Y., Ozaki, K., & Lenardic, A. (2016). Two-step rise of
atmospheric oxygen linked to the growth of continents. Nature Geoscience, 9, 417.
Li, H., Singh, A. K., Mcintyre, L. M., & Sherman, L. A. (2004). Differential gene expression in response
to hydrogen peroxide and the putative PerR regulon of *Synechocystis* sp. strain PCC 6803.
Journal of Bacteriology, 186, 3331-3345.
Liu, W., Hao, J., Elzinga, E. J., Piotrowiak, P., Nanda, V., Yee, N., & Falkowski, P. G. (2020). Anoxic
photogeochemical oxidation of manganese carbonate yields manganese oxide. Proceedings of the
National Academy of Sciences, 117, 22698-22704.
Luo, G., Ono, S., Beukes, N. J., Wang, D. T., Xie, S., & Summons, R. E. (2016). Rapid oxygenation of
Earth's atmosphere 2.33 billion years ago. Science Advances, 2, e1600134.
Lyons, T. W., Reinhard, C. T., & Planavsky, N. J. (2014). The rise of oxygen in Earth's early ocean and
atmosphere. Nature, 506, 307-315.
Martin, W. F., Bryant, D. A., & Beatty, J. T. (2018). A physiological perspective on the origin and
evolution of photosynthesis. FEMS Microbiology Reviews, 42, 205-231.
Mishra, S., & Imlay, J. (2012). Why do bacteria use so many enzymes to scavenge hydrogen peroxide?
Archives of Biochemistry and Biophysics, 525, 145-160.
Morris, J. J., Johnson, Z. I., Szul, M. J., Keller, M., & Zinser, E. R. (2011). Dependence of the
cyanobacterium *Prochlorococcus* on hydrogen peroxide scavenging microbes for growth at the
ocean's surface. PLoS ONE, 6, e16805.
Morris, J. J., Kirkegaard, R., Szul, M. J., Johnson, Z. I., & Zinser, E. R. (2008). Facilitation of robust
growth of *Prochlorococcus* colonies and dilute liquid cultures by "helper" heterotrophic bacteria.
Applied and Environmental Microbiology, 74, 4530-4534.
Myers, C. R., & Nealson, K. H. (1988). Bacterial manganese reduction and growth with manganese oxide
as the sole electron acceptor. Science, 240, 1319-1321.
Olson, S. L., Kump, L. R., & Kasting, J. F. (2013). Quantifying the areal extent and dissolved oxygen
concentrations of Archean oxygen oases. Chemical Geology, 362, 35-43.
Ossa Ossa, F., Hofmann, A., Wille, M., Spangenberg, J. E., Bekker, A., Poulton, S. W., Eickmann, B., &
Schoenberg, R. (2018). Aerobic iron and manganese cycling in a redox-stratified Mesoarchean
epicontinental sea. Earth and Planetary Science Letters, 500, 28-40.
Ozaki, K., Thompson, K. J., Simister, R. L., Crowe, S. A., & Reinhard, C. T. (2019). Anoxygenic
photosynthesis and the delayed oxygenation of Earth's atmosphere. Nature Communications, 10,
3026.
Özkaya, Ö., Xavier, K. B., Dionisio, F., & Balbontín, R. (2017). Maintenance of microbial cooperation
mediated by public goods in single- and multiple-trait scenarios. Journal of Bacteriology, 199,
e00297-00217.
Papp-Wallace, K. M., & Maguire, M. E. (2006). Manganese transport and the role of manganese in
virulence. Annual Reviews in Microbiology, 60, 187-209.
Partensky, F., & Garczarek, L. (2010). *Prochlorococcus*: advantages and limits of minimalism. Annual
Review of Marine Science, 2, 305-331.
Planavsky, N. J., Asael, D., Hofmann, A., Reinhard, C. T., Lalonde, S. V., Knudsen, A., Wang, X., Ossa
Ossa, F., Pecoits, E., Smith, A. J. B., Beukes, N. J., Bekker, A., Johnson, T. M., Konhauser, K.

O., Lyons, T. W., & Rouxel, O. J. (2014). Evidence for oxygenic photosynthesis half a billion
years before the Great Oxidation Event. Nature Geoscience, 7, 283-286.
Reinhard, C. T., Lalonde, S. V., & Lyons, T. W. (2013a). Oxidative sulfide dissolution on the early Earth.
Chemical Geology, 362, 44-55.
Reinhard, C. T., Planavsky, N. J., & Lyons, T. W. (2013b). Long-term sedimentary recycling of rare
sulphur isotope anomalies. Nature, 497, 100-103.
Rodríguez Amor, D., & Dal Bello, M. (2019). Bottom-up approaches to synthetic cooperation in
microbial communities. Life, 9, 22.
Sánchez-Baracaldo, P. (2015). Origin of marine planktonic cyanobacteria. Scientific Reports, 5, 17418.
Sánchez□Baracaldo, P., & Cardona, T. (2020). On the origin of oxygenic photosynthesis and
Cyanobacteria. New Phytologist, 225, 1440-1446.
Sekar, R., Shin, H. D., & Dichristina, T. J. (2016). Direct conversion of cellulose and hemicellulose to
fermentable sugars by a microbially-driven Fenton reaction. Bioresource Technology, 218, 1133-
1139.
Shcolnick, S., Summerfield, T. C., Reytman, L., Sherman, L. A., & Keren, N. (2009). The mechanism of
iron homeostasis in the unicellular cyanobacterium *Synechocystis* sp. PCC 6803 and its
relationship to oxidative stress. Plant Physiology, 150, 2045-2056.
Smethurst, D. G., Kovalev, N., Mckenzie, E. R., Pestov, D. G., & Shcherbik, N. (2020). Iron-mediated
degradation of ribosomes under oxidative stress is attenuated by manganese. Journal of
Biological Chemistry, 295, 17200-17214.
Sobota, J. M., & Imlay, J. A. (2011). Iron enzyme ribulose-5-phosphate 3-epimerase in Escherichia coli is
rapidly damaged by hydrogen peroxide but can be protected by manganese. Proceedings of the
National Academy of Sciences, 108, 5402-5407.
Song, H., Jiang, G., Poulton, S. W., Wignall, P. B., Tong, J., Song, H., An, Z., Chu, D., Tian, L., She, Z.,
& Wang, C. (2017). The onset of widespread marine red beds and the evolution of ferruginous
oceans. Nature Communications, 8, 399.
Stadtman, E., Berlett, B., & Chock, P. (1990). Manganese-dependent disproportionation of hydrogen
peroxide in bicarbonate buffer. Proceedings of the National Academy of Sciences, 87, 384-388.
Stevens Jr., S. E., Patterson, C. O. P., & Myers, J. (1973). The production of hydrogen peroxide by blue-
green algae: a survey. Journal of Phycology, 9, 427-430.
Swanner, E. D., Mloszewska, A. M., Cirpka, O. A., Schoenberg, R., Konhauser, K. O., & Kappler, A.
(2015a). Modulation of oxygen production in Archaean oceans by episodes of Fe(II) toxicity.
Nature Geoscience, 8, 126-130.
Swanner, E. D., Wu, W., Hao, L., Wüstner, M. L., Obst, M., Moran, D. M., Mcilvin, M. R., Saito, M. A.,
& Kappler, A. (2015b). Physiology, Fe(II) oxidation, and Fe mineral formation by a marine
planktonic cyanobacterium grown under ferruginous conditions. Frontiers in Earth Science, 3.
Wan, F., Kong, L., & Gao, H. (2018). Defining the binding determinants of Shewanella oneidensis OxyR:
implications for the link between the contracted OxyR regulon and adaptation. Journal of
Biological Chemistry, 293, 4085-4096.
Wang, Y., & Newman, D. K. (2008). Redox reactions of phenazine antibiotics with ferric (hydr)oxides
and molecular oxygen. Environmental Science & Technology, 42, 2380-2386.
Ward, L. M., Idei, A., Nakagawa, M., Ueno, Y., Fischer, W. W., & Mcglynn, S. E. (2019). Geochemical
and metagenomic characterization of Jinata Onsen, a Proterozoic-analog hot spring, reveals novel
microbial diversity including iron-tolerant phototrophs and thermophilic lithotrophs. Microbes
and Environments, ME19017.
Ward, L. M., Idei, A., Terajima, S., Kakegawa, T., Fischer, W. W., & Mcglynn, S. E. (2017). Microbial
diversity and iron oxidation at Okuoku□hachikurou Onsen, a Japanese hot spring analog of
Precambrian iron formations. Geobiology, 15, 817-835.
Yu, N. Y., Wagner, J. R., Laird, M. R., Melli, G., Rey, S., Lo, R., Dao, P., Sahinalp, S. C., Ester, M., &
Foster, L. J. (2010). PSORTb 3.0: improved protein subcellular localization prediction with

refined localization subcategories and predictive capabilities for all prokaryotes. Bioinformatics,
26, 1608-1615.
Zamocky, M., Furtmuller, P. G., & Obinger, C. (2008). Evolution of catalases from bacteria to humans.
Antioxidants & Redox Signaling, 10, 1527-1548.
Zamocky, M., Gasselhuber, B., Furtmuller, P. G., & Obinger, C. (2012). Molecular evolution of hydrogen
peroxide degrading enzymes. Archives of Biochemistry and Biophysics, 525, 131-144.
Zhou, M., Diwu, Z., Panchuk-Voloshina, N., & Haugland, R. P. (1997). A stable nonfluorescent
derivative of resorufin for the fluorometric determination of trace hydrogen peroxide:
applications in detecting the activity of phagocyte NADPH oxidase and other oxidases.
Analytical Biochemistry, 253, 162-168.
Zinser, E. R. (2018a). Cross-protection from hydrogen peroxide by helper microbes: the impacts on the
cyanobacterium *Prochlorococcus* and other beneficiaries in marine communities. Environmental
Microbiology Reports, 10, 399-411.
Zinser, E. R. (2018b). The microbial contribution to reactive oxygen species dynamics in marine
ecosystems. Environmental Microbiology Reports, 10, 412-427.

643
644