Microbial helpers allow cyanobacteria to thrive in ferruginous waters

Nadia Szeinbaum*[1], Yael Toporek[2], Christopher T. Reinhard[1], Jennifer B. Glass[1]

[1]School of Earth and Atmospheric Sciences, Georgia Institute of Technology, Atlanta,

GA

[2]School of Biological Sciences, Georgia Institute of Technology, Atlanta, GA

*corresponding author: nadia.szeinbaum@biology.gatech.edu

Running title: Metal-reducing bacteria improve cyanobacterial fitness in early oceans

Abstract

The Great Oxidation Event (GOE) was a rapid accumulation of oxygen (O_2) in the atmosphere

as a result of the photosynthetic activity of cyanobacteria. This accumulation reflected the

pervasiveness of O_2 on the planet's surface, indicating that cyanobacteria had become

ecologically successful. The GOE must have required the emergence of several metabolic

innovations, in particular those used for protection from reactive O_2 species (ROS), including

superoxides (O·) and hydroperoxides (H_2O_2), which are byproducts of oxygenic photosynthesis.

In Archean oceans, near-millimolar levels of Fe^{2+} would have reacted with H_2O_2, exacerbating

oxidative damage further. It is thought that cyanobacteria must have not endured the ROS

toxicity of Fe^{2+}-rich oceans. Yet, cyanobacteria colonized Archean oceans extensively enough

to oxygenate the atmosphere. In modern oceans, catalase-lacking cyanobacteria can be

protected by catalase-producing heterotrophs. Here we show experimentally that facultative

metal-reducing helper heterotrophs increase the fitness of *Synechococcus* in high Fe^{2+}

concentrations, expanding the upper bound for tolerable Fe^{2+} concentrations for cyanobacteria

in the ocean. These results suggest that helper bacteria could have been important for the

colonization of ancient ferruginous oceans and imply that analyses considering multiple

evolutionary lineages may reveal metabolic strategies that were shared by co-evolving microbial

populations.

Introduction

Earth's first biogeochemical cycles were driven by anaerobic microorganisms (Martin, Bryant, & Beatty, 2018; Weiss et al., 2016). At around 2.3 billion years ago (Ga), the Great Oxidation Event (GOE) resulted in the initial oxygenation of the atmosphere and surficial biosphere, which ultimately led to the modern dominance of aerobic organisms on Earth's surface (Bar-On, Phillips, & Milo, 2018; Luo et al., 2016). Although biological O_2 production by cyanobacteria was a prerequisite for the GOE (Haqq-Misra, Kasting, & Lee, 2011; Holland, 2002; Lee et al., 2016), oxygenic photosynthesis may have emerged hundreds of millions of years prior to the GOE, during the early Archean (Cardona, Sánchez-Baracaldo, Rutherford, & Larkum, 2019; Lalonde & Konhauser, 2015; Ossa Ossa et al., 2018; Planavsky et al., 2014). The delay between the emergence of biologically-produced O_2 and its accumulation in the atmosphere may have been modulated by geophysical drivers (Catling, Zahnle, & McKay, 2001; Holland, 2009; Lee et al., 2016), but may also reflect the time required for the emergence of several metabolic innovations and the ecological success of cyanobacteria (Blank & Sanchez-Baracaldo, 2010; Johnston, Wolfe-Simon, Pearson, & Knoll, 2009; Lyons, Reinhard, & Planavsky, 2014; Ozaki, Thompson, Simister, Crowe, & Reinhard, 2019). Understanding how cyanobacteria adapted to changing environmental conditions to colonize most of the Earth's surface is essential to understand the microbial ecology of the GOE.

Although the sunlit Archean oceans would have provided cyanobacteria with a large surface area for water oxidation compared to land (Chisholm, 2017; Lyons et al., 2014), high Fe^{2+} concentrations would have created a significant challenge due to its reaction with reactive O_2 species (ROS). Oxygenic photosynthesis produces ROS, namely superoxides (O·) and hydroperoxides (H_2O_2) (Mishra & Imlay, 2012). Hydrogen peroxide can penetrate cellular membranes more easily than O· and subsequently inflict widespread damage inside the cell (Klotz & Loewen, 2003). In Archean oceans, elevated levels of Fe^{2+} (Song et al., 2017) would

have reacted with H_2O_2 to produce highly toxic hydroxyl radicals, further exacerbating oxidative

damage and delaying cyanobacterial colonization of the oceans (Mishra & Imlay, 2012;

Shcolnick, Summerfield, Reytman, Sherman, & Keren, 2009; Swanner et al., 2015). If

cyanobacteria were not equipped to resist ROS from oxygenic photosynthesis, what allowed

them to colonize the ocean and ultimately oxygenate the atmosphere?

Cooperation between different microbial organisms may explain the apparent

evolutionary conundrum of the absence of an essential trait in a population (Morris, Johnson,

Szul, Keller, & Zinser, 2011; Morris, Lenski, & Zinser, 2012). Cooperation occurs when

members of a community produce metabolically costly molecules ("public goods") that become

available to neighboring members (Özkaya, Xavier, Dionisio, & Balbontín, 2017). Cooperation is

common among cyanobacteria and proteobacteria (Christie-Oleza, Sousoni, Lloyd, Armengaud,

& Scanlan, 2017; Zinser, 2018), whose intimate relationship is also evidenced by extensive

horizontal gene transfer between the two lineages (Ben Said & Or, 2017; Braakman, Follows, &

Chisholm, 2017; Cantine & Fournier, 2017; Cavalier-Smith, 2006a, 2006b; Christie-Oleza et al.,

2017; Goldenfeld & Woese, 2011). Cooperation by "helper" organisms providing ROS protection

may have also allowed early cyanobacteria to overcome ROS toxicity in Fe^{2+}-rich oceans (**Fig.**

1A). At the time of the GOE, several potential helper lineages had already diversified

(Battistuzzi, Feijao, & Hedges, 2004), including proteobacteria (Cavalier-Smith, 2006a, 2006b),

which would have provided many opportunities for cooperation.

Aerobic microbes use a variety of hydroperoxidases to inactivate H_2O_2 by reduction or

dismutation to water (Mishra & Imlay, 2012; Zamocky, Furtmuller, & Obinger, 2008), the most

widespread of which are catalases. Catalases first evolved in bacteria independently as heme

and non-heme catalases, possibly before ~3.5 Ga (Klotz & Loewen, 2003; Zamocky et al.,

2008). Among the heme catalases, monofunctional KatEs originated as clade 2 KatEs and later

formed clade 1 and clade 3 (Klotz & Loewen, 2003). Modern proteobacteria can enhance the

growth of cyanobacteria by providing ROS protection by secreting catalase (Morris et al., 2011;

Zinser, 2018). Extracellular catalases have mostly been found to be of the clade 3 KatE type

(Hanaoka et al., 2013; Naclerio, Baccigalupi, Caruso, De Felice, & Ricca, 1995; Pezzoni,

Pizarro, & Costa, 2014). We hypothesize that a similar microbial cooperation, in which a "helper"

organism consumed O_2 or scavenged hydroperoxides, protected cyanobacteria from ROS

toxicity in early, Fe^{2+}-rich oceans (**Fig. 1A**). Our hypothesis predicts that cyanobacteria in co-

culture with proteobacteria should show increased fitness under ferruginous conditions

compared to a cyanobacterial monoculture, and that proteobacteria with higher hydroperoxidase

activity will have a more protective effect than those with lower activity.

Synthetic communities are a useful method to study ecological and geochemical

dynamics of individual species within a community because the initial conditions can be

precisely controlled (Widder et al., 2016). Synthetic communities are particularly useful for

studies of phototrophic-driven consortia in simulated pre-GOE conditions that do not exist on

Earth today (illuminated, eukaryotic-free, Fe^{2+}-rich, and anoxic). To test our hypothesis, we used

synthetic consortia of model bacterial species consisting of one KatE-lacking cyanobacterium

Synechococcus sp. PCC7002 (hereafter *Synechococcus*) and one of six different KatE-

containing gammaproteobacterial helpers (metal-reducing *Shewanella*). *Synechococcus* sp.

PCC7002 has been used as a model cyanobacteria to understand modern and past

phototrophic metabolisms (Alexander S. Beliaev et al., 2014; Bryant & Frigaard, 2006; Swanner

et al., 2015). *Shewanella* spp. are facultative anaerobes that can survive O_2 intrusions in the

presence of high Fe^{2+} concentrations (Sekar, Shin, & DiChristina, 2016) and contain variable

numbers and types of KatEs. Co-cultures were grown under parameters simulating a pre-GOE,

Fe^{2+}-rich shallow continental margin overlain by a CO_2- and H_2-rich anoxic atmosphere (**Fig. 1**).

The relative fitness of cyanobacterial cultures was compared with and without the inclusion of

gammaproteobacterial helpers. We found that some *Shewanella* species allowed

Synechococcus to grow in Fe^{2+}-rich conditions that precluded growth of *Synechococcus*

monocultures. Our findings stress the importance of considering microbial cooperation and

complementary genetic repertoires in models of early Earth ecology leading to the GOE.

Results

Impaired growth of KatE-lacking cyanobacteria in high Fe^{2+} is restored in the presence of

KatE-containing proteobacteria. *Synechococcus* was grown with 25 µM (low Fe^{2+}), 500 µM

Fe^{2+}, or 1000 µM Fe^{2+} (high iron) under an anoxic atmosphere**.** *Synechococcus* reached highest

cell densities (~2.7×10^7 cells mL^{-1}) at 25 µM Fe^{2+} **(Fig. 2)**. At 500 µM Fe^{2+}, *Synechococcus*

showed an increased lag period (2 days, **Fig. S1B**) and similar growth rates **(Fig. S2)**. Growth

was significantly impaired at 1,000 µM Fe^{2+} (~20% of the yields under low-Fe^{2+} conditions**; Fig.**

2). When *Synechococcus* was grown in co-culture with different *Shewanella* strains at 1000 µM

Fe^{2+}, growth yields increased 30-100% compared to yields from co-cultures with low Fe^{2+} (**Fig.**

2). At low levels of Fe^{2+}, the presence of *Shewanella* generally hindered *Synechococcus* growth,

as evidenced by a longer lag phase of 2-4 days in co-culture with 25 µM Fe^{2+} compared to 0.5

days in monoculture (**Fig. S1A),** and 3-4 days in co-culture with 500 µM Fe^{2+} compared to 2

days in monoculture (**Fig. S1B**). Among the strains tested, *Shewanella algae* MN-01 and

Shewanella baltica OS155 had the most positive effect on the growth rates and yields of

Synechococcus. Shewanella oneidensis MR-1 had a lesser positive effect. By the end of all of

the incubations, *Synechococcus* was the only strain remaining in the cultures (**Fig. S3**). Iron

levels remained close to initial concentrations only in co-cultures and decreased by the second

day in *Synechococcus* monocultures (**Fig. S4**).

Proteobacteria with higher catalase activity are better cyanobacterial helpers. Higher

growth yields and stable Fe^{2+} levels in co-cultures with high Fe^{2+} suggested that the mechanism

for protection could be due to O_2 or H_2O_2 scavenging. In order to test whether *Shewanella*

partners protected cyanobacteria by scavenging O_2 or H_2O_2, we measured the maximum O_2

respiration rate and H_2O_2 scavenging rate in cell suspensions of the most and least "helpful"

helper strains. *S. algae* MN-01 and *Shewanella* sp. BrY had higher O_2 respiration rates than *S.*

baltica and *S. oneidensis* MR-1 (**Fig. 3A**). *S. algae* MN-01 showed the highest H_2O_2 scavenging

capacity, followed by *S. baltica*, while *Shewanella* sp. BrY and *S. oneidensis* MR-1 showed

similar lower rates (25-40% of the other two strains; **Fig. 3B**). These differences were also

reflected in the different growth rates of *Shewanella* strains upon exposure to increasing

concentrations of H_2O_2 (**Fig. S5**). No differences were detected at the amino acid level that

would explain these differences (**Fig. S6**). Final growth yields in *Synechococcus* co-cultures

were proportional to the H_2O_2 scavenging activity of each *Shewanella* strain (**Fig. 3C**).

Distribution of KatE catalases in Cyanobacteria suggests link with cell morphology and

salinity. The well-known extracellular role of KatE on scavenging extracellular H_2O_2 (Naclerio et

al., 1995; Shin, Choi, & Cho, 2008), the presence of KatE in metal-reducing strains, and the link

between higher rates of extracellular H_2O_2 scavenging and growth of cyanobacteria in pre-GOE

conditions is suggestive of an important ecological role of KatE in catalase-based interactions,

which are thought to be prevalent in the environment (Zinser, 2018). KatE was absent in our

model species, *Synechococcus* spp. PCC7002. Out of 2,886 cyanobacterial genomes, only 135

(~5%) contained a KatE homolog. Of those with assigned taxonomy, the vast majority were

Nostocales from freshwater or soils (**Table S1**). A small minority were Synechococcales,

Chroococcales, or Oscillatoriales from marine or high-salt environments.

We created a phylogeny of bacterial KatEs to determine the evolutionary history of this

gene in cyanobacteria. Almost all of the cyanobacterial clade 3 KatEs grouped together and

branched off of gammaproteobacterial KatEs (**Fig. 4**). Multiple phylogenetic models confirmed

this result (not shown). The tree topology of KatE is incongruent with the previously established

history of bacteria, where Cyanobacteria are ancestral to Proteobacteria (Gupta, 2000; Marin,

Battistuzzi, Brown, & Hedges, 2017), suggesting later horizontal gene transfer of *katE* genes to

cyanobacteria.

Among the gammaproteobacterial KatE genes, KatEs formed three distinct groups. The

first group branched off earlier than the rest of the gammaproteobacterial KatEs and grouped

more closely to KatEs from Flavobacteriia, Deinococci, and Clostridia. This group contained

KatEs experimentally verified to be extracellular from *Bacillus subtilis*, *P. aeruginosa* PAO1,

Exiguobacterium oxidotolerans, and *Alteromonas macleodii* (Hanaoka et al., 2013; Morris et al.,

2011; Naclerio et al., 1995; Pezzoni et al., 2014). The second group contained other

Alteromonadales, including iron-reducing *Shewanella* and *Ferrimonas* species. The third group

contained the majority of the sequences analyzed and also grouped together with

proteobacterial KatEs of the alpha, beta, and delta classes. Unlike the fragmented distribution of

KatEs among Gammaproteobacteria, the rest of the KatEs from other taxonomic groups

clustered together.

Discussion

The fitness of cyanobacteria increases in the presence of helper proteobacteria under

pre-GOE conditions. In our experiments, axenic growth of *Synechococcus* was impaired at

high Fe^{2+} concentrations but was reestablished in the presence of *Shewanella*. These results

provide empirical evidence that ecological interactions between members of different species

with phenotypes relevant to the onset of the GOE can provide means to overcome Archean

levels of Fe^{2+}-induced ROS stress. Our findings underscore the need to consider ecological

interactions between members of different species in the reconstruction of our microbial past

and Earth's biogeochemical cycles.

Reconstructions of the evolutionary history of oxygenic phototrophs have revealed a

variety of functional and morphological innovations that likely contributed to the GOE (Blank &

Sanchez-Baracaldo, 2010; Chrismas, Barker, Anesio, & Sánchez-Baracaldo, 2016; Di Cesare et

al., 2018; Hohmann-Marriott & Blankenship, 2011; Sanchez-Baracaldo, Bianchini, Di Cesare,

Callieri, & Chrismas, 2019). Expanding analyses to include organisms other than cyanobacterial

lineages, but important to their ecological success, can provide new explanations for the

interpretation of the geochemical record. For instance, lack of cyanobacteria in modern

environments containing above ~80 μM Fe^{2+} and lack of cyanobacterial growth in laboratory

culture above ~200 μM Fe^{2+} have contributed to the idea that Fe^{2+}-rich oceans would have

slowed down the colonization of the ocean surface due to toxicity, possibly delaying global

oxygenation (Swanner et al., 2015). However, based on our data, catalase-producing, iron-

reducing bacteria could have augmented the ability of cyanobacteria to respond to Fe^{2+} toxicity

while keeping Fe^{2+} levels constant, presumably due to a combination of ROS scavenging and

O_2 respiration. Including facultative metal-reducing bacteria in models would attribute the delay

in oxygenation to cryptic oxygen cycling (Garcia-Robledo et al., 2017) and reduced levels of

Fe^{2+}-catalyzed ROS, conditions that would have allowed cyanobacteria to grow in a largely

anoxic ocean during the GOE's "lag phase".

Under our experimental conditions, the beneficial role of *Shewanella* to *Synechococcus*

was ultimately transient. This may be a natural consequence of the dynamics of public goods in

cooperative interactions between microbial groups (Corno, Villiger, & Pernthaler, 2013;

Rodríguez Amor & Dal Bello, 2019), when the members of the population that provide the

extracellular "public good" are at a fitness disadvantage by providing a costly product that

nonproducers use and benefit from (Cremer et al., 2019; Özkaya et al., 2017). In our system,

Synechococcus gained the fitness advantage of ROS protection at the expense of *Shewanella*,

whose population was eliminated from the system possibly due to limiting organic carbon (A. S.

Beliaev et al., 2014), as *Synechococcus* grew in numbers.

The transient partnership between *Synechococcus* and *Shewanella* in our cultures may

reflect that the conditions in which cyanobacteria and proteobacteria grow once overlapped

(illuminated, anoxic, and Fe^{2+}-rich), but might have changed as Earth became oxidizing enough

to reduce the pool of free Fe^{2+}, leading to spatial separation of these populations. This scenario

is consistent with the different ecological niches that *Shewanella* and *Synechococcus* occupy,

and the different environments in which they are found. While *Synechococcus* dominates the

open ocean surface today (Flombaum et al., 2013), *Shewanella* species have been isolated

mostly from dark and organic-rich environments such as sediments or inside the gut of animals

(Hau & Gralnick, 2007; Konstantinidis et al., 2009). An association with ROS-scavenging

proteobacteria would have been beneficial to cyanobacteria only during the initial stages of the

evolution of oxygenic photosynthesis and colonization of the ocean (**Fig. 5**). It is possible that a

close interaction could have been sustained in Archean oceans as long as both populations

were maintained near our initial concentrations (10^5 cells mL^{-1}).

Differences in fitness effects can be explained by catalase activity and ROS survival

capacity of helper partners. Although the six *Shewanella* strains we examined have similar

ecological niches, we noted substantial differences in their effect on *Synechococcus'* growth.

Strains that scavenged H_2O_2 faster resulted in higher cyanobacterial yields. Genomic

sequences of catalase genes, however, did not explain the phenotypic difference observed.

These differences may be explained by higher catalase expression levels in strains that are

more resistant to ROS stress (Wang et al., 2009). These results support the idea that genomic

information alone may not be enough to understand the interplay between microbial community

members and their ecological role (M. Lee et al., 2017).

Microbes are thought to have developed resistance to ROS after the GOE, which

allowed them to colonize oxic environments (Hamilton, Bryant, & Macalady, 2016). Yet genomic

and biochemical evidence suggest that catalases could have evolved earlier, in the Archean

(Klotz & Loewen, 2003; Olson et al., 2017). In an Archean ferruginous environment, Fe^{2+}-driven

ROS production could have imposed a selection pressure strong enough for the evolution of

catalases, much earlier than the oxygenation of the atmosphere and the ecological expansion of

aerobic life. The selection pressure for the evolution of catalases could have been stronger in

microbes other than oxygenic cyanobacteria. In fact, the evolution of catalases is thought to

have evolved initially in proteobacteria (Klotz & Loewen, 2003). The evolutionary pressure to

evolve an ROS scavenging system would have been strong in iron-cycling organisms. Among

the metal-reducing bacteria, Gammaproteobacteria are the most widespread facultative

aerobes. Due to the high reactivity of Fe^{2+} with hydroperoxides, the iron-rich, multi-heme

cytochromes involved in extracellular iron respiration in proteobacteria (Shi et al., 2016) could

become targets of extracellular H_2O_2 toxicity. Modern facultative anaerobic metal-reducing

bacteria, often found at oxic/anoxic transition zones, may experience strong selection pressure

to retain outer membrane or extracellular H_2O_2 scavenging enzymes. The association between

catalases and Fe^{2+}-enhanced ROS toxicity is consistent with genomic correlation between the

presence of catalase in *Shewanella* species and their ability to reduce metals (**Fig. S7**); for

example, *S. denitrificans* is the only *Shewanella* species that lacks *katE* genes and is also the

only one unable that reduce metals, due to its lack of heme-rich extracellular electron conduits.

The distribution pattern of clade 3 KatEs in cyanobacteria suggests that extracellular

KatEs providing ROS protection may be phylogenetically distinct and adapted to marine

conditions. In the modern open ocean, the majority of the nitrogen fixers are filamentous non-

heterocyst-forming   *Trichodesmium*,   heterocyst-forming   eukaryotic   symbionts

Richelia, *Calothrix,* and relatives, or unicellular *Crocosphaera* and *Cyanothece,* yet none of the

sequenced *Trichodesmium*, *Richelia,* and *Crocosphaera* genomes, and only 9 out of the 66

Calothrix genomes, contain KatE (**Table S2**). *Trichodesmium* is usually found in association

with *A. macleodii,* which provides protection from ROS (M. Lee et al., 2017). *Trichodesmium*

epibionts also include Piscirickettsiaceae (Gradoville, Crump, Letelier, Church, & White, 2017)

whose   KatEs   grouped   in   the   second   gammaproteobacterial   clade   together   with

Alteromonadales and separate from most other Gammaproteobacteria (**Fig. 4**). Similarly, *A.*

macleodii provides catalase as a "public good" to marine *Prochlorococcus* (Morris et al., 2011)

and their KatEs grouped in the first cluster. Alongside nutrient scarcity (Christie-Oleza et al.,

2017; Gradoville et al., 2017), protection from toxic byproducts such as ROS toxicity can thus

also be an important cooperative ecological mechanism between cyanobacteria and non-

phototrophic helpers in modern marine environments (Zinser, 2018), or oceans on early Earth.

The *Prochlorococcus-Alteromonas* interaction via catalase has been used as a case

study to explain reductive evolution (Morris et al., 2012). The history of KatE, however, might

have had a different evolutionary path. The modern pattern of low KatE incidence in

cyanobacteria can be explained by clade 3 KatE originating in proteobacteria and later

transferred to a few cyanobacterial lineages (Klotz & Loewen, 2003; Zamocky et al., 2008)

supporting a model in which a mutualistic association between cyanobacteria and

proteobacteria based on ROS protection originated in the Archean under anoxic conditions

(Olson et al., 2017).

This study expands the upper bound for tolerable Fe^{2+} concentrations for cyanobacteria

in oceans. For around two billion years, the anoxic conditions of early Earth sustained abundant

soluble Fe^{2+} in Archean seawater (Anbar et al., 2007; Hazen & Ferry, 2010; Holland, 2006;

Johnson, Beard, & Roden, 2008; Reinhard et al., 2017). Fe^{2+} buffered by carbonates (siderite,

$FeCO_3$) or phosphates (vivianite, $Fe_3(PO_4)_2 \cdot 8H_2O$) would have been present at up to ~100 uM

to ~1 mM or more (Derry, 2015; Drever, 1974; Holland, 1973). Under these conditions,

cyanobacteria might have not survived alone (Swanner et al., 2015). Recent metagenomic

analyses suggest that oxygenic photosynthesis preceded aerobic respiration in cyanobacteria,

which was acquired later by horizontal gene transfer (Cavalier-Smith, 2006b; Soo, Hemp, Parks,

Fischer, & Hugenholtz, 2017), pointing to a close association with other microbial lineages that

had already developed ROS resistance and respiration capacity, such as proteobacteria. Based

on our study, the presence of other microbial lineages would have allowed cyanobacteria to

grow in Fe^{2+} concentrations close to the most recent estimates of Archean seawater levels

(Derry, 2015).

Our results suggest that while the metabolic repertoire of different microbial lineages

often provides us with patterns that reveal how life navigated alternative environmental

landscapes, metabolic strategies can be shared among closely interacting microbial populations

and thus critical evolutionary information may be stored in separate microbial members of a

community. Therefore, analyses focused on only one evolutionary lineage may obscure

metabolic strategies that were shared by co-evolving microbial populations. Considering

microbial interactions in the establishment of novel biochemical pathways can open new

windows through which to understand how the physiology of microbes evolved.

Materials and Methods

Strains used in this study. *Synechococcus spp.* PCC7002 was obtained from the Paris

Culture Collection. *S. putrefaciens* strain 200, *S. oneidensis* MR-1, *Shewanella spp.* strain BrY

were kind gifts from the lab of Dr. Thomas DiChristina (Georgia Institute of Technology). *S.*

denitrificans and *S. loihica* were a kind gift from Dr. Jeffrey Gralnick (University of Minnesota).

All strains were grown in lysogeny broth (LB, 10 g L^{-1} NaCl, 10 g L^{-1} tryptone, 5 g L^{-1} yeast

extract, Sigma-Aldrich) or a modified A+ medium (Stevens Jr., Patterson, & Myers, 1973) with

NH_4^+ (10 mM) instead of NO_3^- as carbon source and no iron(III), and additional amino acids L-

serine, L-arginine and L-glutamic acid (20 mg L^{-1}) when used to culture *Shewanella* strains. The

use of LB or modified A+ is specified in each one of the methods, where appropriate.

GOE-mimicking experimental setup and co-culture growth conditions. *Synechococcus*

PCC 7002 was grown on modified A+ medium. To eliminate O_2 as much as possible at the start

of the co-culture experiment, *Synechococcus PCC7002* sp. PCC 7002 was incubated in serum

bottles flushed for 10 min with $N_2:CO_2$ 90:10 daily, and once before transferring to the anaerobic

chamber. In parallel, *Shewanella* strains were transferred from an overnight culture in LB into

anoxic bottles with minimal medium (flushed once with $N_2:CO_2$ 90:10) with lactate as electron

donor and fumarate as electron acceptor. After overnight growth, all bottles were opened inside

the anoxic chamber ($H_2:CO_2:N_2$ 4:5:91), and mixed at appropriate concentrations based on their

OD600 to achieve a 1:1 ratio of OD 0.01 each. Co-cultures were grown in 10 mL in triplicate in

6-well tissue culture plates and homogenized daily by gentle pipetting ~50% of the volume up

and down. Light was provided with a fluorescent light in a 12:12 light:dark timer-controlled cycle.

$FeCl_2$ was added at 25, 500 or 1000 µM.

Analytical methods during co-culture. Cell numbers were quantified in a LSR Fortessa flow

cytometer in a high-throughput manner using the software FACSDiva™ (BD Biosciences, CA) to

set the parameters and monitor each quantification. At each time point, 200 µL of sample were

loaded into a 96-well plate inside the anoxic chamber, covered in parafilm to minimize iron

oxidation, and then transported to the cytometer. Each sample consisted of 10 µL taken for

injection after mixing twice (100 µL mixing volume) and run at a rate of 0.5 µL s^{-1}. Cyanobacteria

were detected by phycocyanin/chlorophyll autofluorescence using blue (488 nm) and yellow-

green lasers (561nm) measured at 655-684 (Hill, Chrisler, Beliaev, & Bernstein, 2017).

Optimization and calibration of the quantification parameters were achieved using yellow-green

1 µm microspheres (441/485 ex/em) (Polysciences, PA). Live cyanobacteria were also

quantified using Syto9 using FITC filters. Events above the thresholds of PerCP and FITC were

considered live cyanobacteria. Propidium iodide could not be used to identify "dead'

cyanobacteria, as the emission spectra overlapped with that of their autofluorescence.

Shewanella cells could potentially overlap with iron particles in the cytometer, except for their

complete absence in the sample. Iron was determined colorimetrically via the ferrozine method

(Stookey, 1970), sampling inside the chamber into 0.5 M HCl-filled wells of a 96-well plate and

using a spectrophotometer with plate-reading capacity for quantification (Tecan, Switzerland).

Oxygen was monitored using 3 mm tip diameter fiber-optic sensors (Robust Oxygen Sensors;

Pyroscience, Germany).

Hydroperoxide resistance assays. Six *Shewanella* strains were incubated in 96-well plates

with minimal media and lactate (10 mM) or acetate (10 mM) as electron donor under aerobic

conditions. Growth was monitored continuously by OD at 600 nm in a spectrophotometer with

plate-reading capacity (Tecan, Switzerland). Hydrogen peroxide was added after an initial

period of growth for 4 hours, at a final concentration of 10, 25, 50, 100, or 250 µM.

Whole-cell extracellular hydroperoxide scavenging assays. To compare the abilities of the

selected *Shewanella* strains to remove exogenous H_2O_2 from their surrounding environment,

strains were first activated from the -80ºC freezer stock by incubating them at 30ºC overnight in

lysogeny broth (LB, 10 g L^{-1} NaCl, 10 g L^{-1} tryptone, 5 g L^{-1} yeast extract, Sigma-Aldrich).

Activated cultures were streaked onto LB plates (LB in 1.5% w/v agar, Sigma-Aldrich). Cultures

were harvested by centrifugation at 3000 rpm, washed and transferred into minimal medium

with lactate at an OD_{600} = 0.02. Cells were incubated at 30ºC until harvesting at mid-log phase

(OD_{600} = 0.15-0.35), washed twice with minimal medium, then inoculated at OD_{600} = 0.05 into

minimal medium amended with various concentrations of exogenous H_2O_2 (3-5000 µM).

Samples were collected and analyzed for residual H_2O_2 every 3-5 minutes using the resorufin-

horseradish peroxidase colorimetric assay. H_2O_2 disappearance followed an exponential decay

[$H_2O_{2(t=n)}$ = $H_2O_{2(t=0)}$ x e^{-kt}] and thus constants were calculated from the estimated curves.

Whole-cell O_2 consumption rates. Each *Shewanella* strain was grown in LB from an overnight

tube of the same media until reaching early exponential growth (~0.3). Cells were then washed

once and resuspended in LB at an OD at 600nm of 0.1 and flushed with air for 10 minutes.

Immediately after, 15 mL of cells were transferred to a screw-top vial with contactless optical

sensors for O_2 to monitor O_2 consumption (TOVIAL20; Pyroscience, Germany). Flushed media

was used as the abiotic control.

Catalase phylogeny. All KatE sequences used in this study were downloaded from the

PATRIC database searching by taxonomic group. All KatE sequences available were chosen,

but only sequences with more than 300 amino acids were kept for analysis. The cyanobacterial

sequences types were identified by conserved domain analysis in NCBI. Phylogenies were

constructed using MEGA X (Kumar, Stecher, Li, Knyaz, & Tamura, 2018). Initial tree(s) for the

heuristic search were obtained automatically by applying Neighbor-Join and BioNJ algorithms to

a matrix of pairwise distances estimated using a JTT model, and then selecting the topology

with superior log likelihood value. A discrete Gamma distribution was used to model

evolutionary rate differences among sites (5 categories (+*G*, parameter = 1.0309). The rate

variation model allowed for some sites to be evolutionarily invariable ([+*I*], 8.13% sites). This

analysis involved 115 amino acid sequences. All positions containing gaps and missing data

were eliminated (complete deletion option). There was a total of 309 positions in the final

dataset.

KatE amino acid sequence comparison. KatE amino acid sequences from strains *Shewanella*

algae MN-01, *Shewanella* spp. BrY, *Shewanella oneidensis* MR-1, and *S. baltica* OS155 were

aligned with MUSCLE in MEGAX (Kumar et al., 2018). Conserved residues in the alignment

were color coded using BOXSHADE 3.2.3 (http://sourceforge.net/projects/boxshade/).

Conserved sequences of the proteins' heme-binding, NADH-binding and tetramer-binding motifs

as determined by NCBI's Conserved Domain search were manually inspected for residue

differences.

Acknowledgements. This research was funded by NASA Exobiology grant NNX14AJ87G and

a NASA Astrobiology Postdoctoral Fellowship to NS. We thank Jennifer Thweatt for advice on

culturing *Synechococcus* sp. PCC7002. We thank Amit Reddi for sharing equipment with us.

We thank Sommer Durham for technical assistance with the cytometer. We thank Patricia

Sanchez-Baracaldo and Manuel Kleiner for helpful discussions.

Competing Interests: The authors declare no competing interests.

Figure 1. Experimental setup of synthetic communities in pre-GOE environments. A)
Conceptual model of potential interactions of metal-reducing proteobacteria and cyanobacteria
in Fe^{2+}-rich, anoxic liquid cultures; B) qualitative growth differences in representative culture
plates of monocultured S*ynechococcus* or co-cultured with *Shewanella;* C) experimental setup
mimicking pre-GOE shallow water environments recreated by incubating synthetic communities
in small-volume tissue culture wells inside an anoxic chamber.

393 **Figure 2. Growth yields of *Synechococcus* in mono- or co-culture in pre-GOE conditions.**
394 This bar graph shows the maximum *Synechococcus* growth yields alone or in co-culture with
395 *Shewanella* with 25 µM (light gray), 500 µM (gray) or 1000 µM (dark gray) Fe^{2+}. Yields
396 correspond to the highest number of cells observed during the growth period. Error bars
397 represent the standard deviation of triplicate samples.

Figure 3. Oxygen and hydrogen peroxide scavenging capacity of _Shewanella_ spp.
Maximum (A) H_2O_2 scavenging and (B) O_2 utilization rates of _S. oneidensis_ MR-1 (MR-1),
Shewanella sp. BrY (BrY), _S. baltica_ OS155 (Baltica), and _Shewanella algae_ MN-01 (MN-01).
(C) Correlation between the H_2O_2 scavenging rate of each _Shewanella_ strain and the growth
yield of _Synechococcus_ in co-culture when incubated with that strain. Error bars represent
standard errors of triplicate measurements. The abiotic control was set up identically but not
inoculated.

Figure 4. Maximum likelihood phylogenetic inference of clade 3 KatE proteins. All KatE
sequences from all sequenced genomes of Cyanobacteria were used to construct a maximum
likelihood phylogeny. The tree is drawn to scale, with branch lengths measured in the number of
substitutions per site. This analysis involved 115 amino acid sequences. Clade 2 KatEs were
used as the outgroup.

Figure 5. A Proterozoic paleo-eco-evolutionary scenario for Cyanobacteria and
Proteobacteria. This schematic is a conceptual model for a potential ecological association
between cyanobacteria and proteobacteria coexisting in the Archean ocean. The RSS
detoxifying machinery emerging in euxinic margins could have evolved into ROS scavenging
proteins, aiding phototrophs. In modern environments, previously coexisting Cyanobacteria and
Proteobacteria are now found in separate niches (Cavalier-Smith, 2006b; Hau & Gralnick, 2007;
Klotz & Loewen, 2003; Lyons et al., 2014; Olson et al., 2017; Zamocky et al., 2008).

References

Anbar, A. D., Duan, Y., Lyons, T. W., Arnold, G. L., Kendall, B., Creaser, R. A., . . . Buick, R. (2007). A Whiff of Oxygen Before the Great Oxidation Event? *Science, 317*(5846), 1903-1906. doi:10.1126/science.1140325

Bar-On, Y. M., Phillips, R., & Milo, R. (2018). The biomass distribution on Earth. *Proceedings of the National Academy of Sciences, 115*(25), 6506-6511. doi:10.1073/pnas.1711842115

Battistuzzi, F. U., Feijao, A., & Hedges, S. B. (2004). A genomic timescale of prokaryote evolution: insights into the origin of methanogenesis, phototrophy, and the colonization of land. *Bmc Evolutionary Biology, 4*. doi:Artn 44

10.1186/1471-2148-4-44

Beliaev, A. S., Romine, M. F., Serres, M., Bernstein, H. C., Linggi, B. E., Markillie, L. M., . . . Konopka, A. (2014). Inference of interactions in cyanobacterial–heterotrophic co-cultures via transcriptome sequencing. *The Isme Journal, 8*, 2243. doi:10.1038/ismej.2014.69

https://www.nature.com/articles/ismej201469#supplementary-information

Beliaev, A. S., Romine, M. F., Serres, M., Bernstein, H. C., Linggi, B. E., Markillie, L. M., . . . Konopka, A. (2014). Inference of interactions in cyanobacterial-heterotrophic co-cultures via transcriptome sequencing. *ISME J, 8*(11), 2243-2255. doi:10.1038/ismej.2014.69

Ben Said, S., & Or, D. (2017). Synthetic Microbial Ecology: Engineering Habitats for Modular Consortia. *Front Microbiol, 8*, 1125. doi:10.3389/fmicb.2017.01125

Blank, C. E., & Sanchez-Baracaldo, P. (2010). Timing of morphological and ecological innovations in the cyanobacteria - a key to understanding the rise in atmospheric oxygen. *Geobiology, 8*(1), 1-23. doi:10.1111/j.1472-4669.2009.00220.x

Braakman, R., Follows, M. J., & Chisholm, S. W. (2017). Metabolic evolution and the self-organization of ecosystems. *Proc Natl Acad Sci U S A, 114*(15), E3091-E3100. doi:10.1073/pnas.1619573114

Bryant, D. A., & Frigaard, N. U. (2006). Prokaryotic photosynthesis and phototrophy illuminated. *Trends Microbiol, 14*(11), 488-496. doi:10.1016/j.tim.2006.09.001

Cantine, M. D., & Fournier, G. P. (2017). Environmental Adaptation from the Origin of Life to the Last Universal Common Ancestor. *Orig Life Evol Biosph*. doi:10.1007/s11084-017-9542-5

462   Cardona, T., Sánchez-Baracaldo, P., Rutherford, A. W., & Larkum, A. W. (2019). Early Archean origin of
463   Photosystem II. *Geobiology, 17*(2), 127-150. doi:10.1111/gbi.12322

464
465   Catling, D. C., Zahnle, K. J., & McKay, C. P. (2001). Biogenic Methane, Hydrogen Escape, and the
466   Irreversible Oxidation of Early Earth. *Science, 293*(5531), 839-843. doi:10.1126/science.1061976

467
468   Cavalier-Smith, T. (2006a). Cell evolution and Earth history: stasis and revolution. *Philos Trans R Soc Lond
469   B Biol Sci, 361*(1470), 969-1006. doi:10.1098/rstb.2006.1842

470
471   Cavalier-Smith, T. (2006b). Rooting the tree of life by transition analyses. *Biol Direct, 1*, 19.
472   doi:10.1186/1745-6150-1-19

473
474   Chisholm, S. W. (2017). Prochlorococcus. *Curr Biol, 27*(11), R447-R448. doi:10.1016/j.cub.2017.02.043

475
476   Chrismas, N. A. M., Barker, G., Anesio, A. M., & Sánchez-Baracaldo, P. (2016). Genomic mechanisms for
477   cold tolerance and production of exopolysaccharides in the Arctic cyanobacterium
478   Phormidesmis priestleyi BC1401. *BMC Genomics, 17*(1), 533. doi:10.1186/s12864-016-2846-4

479
480   Christie-Oleza, J. A., Sousoni, D., Lloyd, M., Armengaud, J., & Scanlan, D. J. (2017). Nutrient recycling
481   facilitates long-term stability of marine microbial phototroph–heterotroph interactions. *2*,
482   17100. doi:10.1038/nmicrobiol.2017.100

483
484   Corno, G., Villiger, J., & Pernthaler, J. (2013). Coaggregation in a microbial predator–prey system affects
485   competition and trophic transfer efficiency. *Ecology, 94*(4), 870-881. doi:10.1890/12-1652.1

486
487   Cremer, J., Melbinger, A., Wienand, K., Henriquez, T., Jung, H., & Frey, E. (2019). Cooperation in
488   Microbial Populations: Theory and Experimental Model Systems. *Journal of Molecular Biology,
489   431*(23), 4599-4644. doi:https://doi.org/10.1016/j.jmb.2019.09.023

490
491   Derry, L. A. (2015). Causes and consequences of mid-Proterozoic anoxia. *Geophysical Research Letters,
492   42*(20), 8538-8546. doi:10.1002/2015gl065333

493
494   Di Cesare, A., Cabello-Yeves, P. J., Chrismas, N. A. M., Sanchez-Baracaldo, P., Salcher, M. M., & Callieri, C.
495   (2018). Genome analysis of the freshwater planktonic Vulcanococcus limneticus sp. nov. reveals
496   horizontal transfer of nitrogenase operon and alternative pathways of nitrogen utilization. *BMC
497   Genomics, 19*(1), 259. doi:10.1186/s12864-018-4648-3

498
Drever, J. (1974). Geochemical Model for the Origin of Precambrian Banded Iron Formations. *GSA
Bulletin, 85*(7), 1099-1106. doi:10.1130/0016-7606(1974)85<1099:Gmftoo>2.0.Co;2

Flombaum, P., Gallegos, J. L., Gordillo, R. A., Rincon, J., Zabala, L. L., Jiao, N., . . . Martiny, A. C. (2013).
Present and future global distributions of the marine Cyanobacteria Prochlorococcus and
Synechococcus. *Proc Natl Acad Sci U S A, 110*(24), 9824-9829. doi:10.1073/pnas.1307701110

Garcia-Robledo, E., Padilla, C. C., Aldunate, M., Stewart, F. J., Ulloa, O., Paulmier, A., . . . Revsbech, N. P.
(2017). Cryptic oxygen cycling in anoxic marine zones. *Proc Natl Acad Sci U S A, 114*(31), 8319-
8324. doi:10.1073/pnas.1619844114

Goldenfeld, N., & Woese, C. (2011). Life is Physics: Evolution as a Collective Phenomenon Far From
Equilibrium. *Annual Review of Condensed Matter Physics, 2*(1), 375-399. doi:10.1146/annurev-
conmatphys-062910-140509

Gradoville, M. R., Crump, B. C., Letelier, R. M., Church, M. J., & White, A. E. (2017). Microbiome of
Trichodesmium Colonies from the North Pacific Subtropical Gyre. *Frontiers in Microbiology,
8*(1122). doi:10.3389/fmicb.2017.01122

Gupta, R. S. (2000). The Natural Evolutionary Relationships among Prokaryotes. *Critical Reviews in
Microbiology, 26*(2), 111-131. doi:10.1080/10408410091154219

Hamilton, T. L., Bryant, D. A., & Macalady, J. L. (2016). The role of biology in planetary evolution:
cyanobacterial primary production in low-oxygen Proterozoic oceans. *Environ Microbiol, 18*(2),
325-340. doi:10.1111/1462-2920.13118

Hanaoka, Y., Takebe, F., Nodasaka, Y., Hara, I., Matsuyama, H., & Yumoto, I. (2013). Growth-Dependent
Catalase Localization in Exiguobacterium oxidotolerans T-2-2T Reflected by Catalase Activity of
Cells. *PLoS One, 8*(10), e76862. doi:10.1371/journal.pone.0076862

Haqq-Misra, J., Kasting, J. F., & Lee, S. (2011). Availability of O2 and H2O2 on Pre-Photosynthetic Earth.
Astrobiology, 11(4), 293-302. doi:10.1089/ast.2010.0572

Hau, H. H., & Gralnick, J. A. (2007). Ecology and biotechnology of the genus Shewanella. *Annual Review
of Microbiology, 61*, 237-258. doi:10.1146/annurev.micro.61.080706.093257

Hazen, R. M., & Ferry, J. M. (2010). Mineral Evolution: Mineralogy in the Fourth Dimension. *Elements,
6*(1), 9-12. doi:10.2113/gselements.6.1.9

Hill, E. A., Chrisler, W. B., Beliaev, A. S., & Bernstein, H. C. (2017). A flexible microbial co-culture platform
for simultaneous utilization of methane and carbon dioxide from gas feedstocks. *Bioresource
Technology, 228*, 250-256. doi:https://doi.org/10.1016/j.biortech.2016.12.111

Hohmann-Marriott, M. F., & Blankenship, R. E. (2011). Evolution of photosynthesis. *Annu Rev Plant Biol,*
62, 515-548. doi:10.1146/annurev-arplant-042110-103811

544
545   Holland, H. D. (1973). The Oceans; A Possible Source of Iron in Iron-Formations. *Economic Geology,*
546   *68*(7), 1169-1172. doi:10.2113/gsecongeo.68.7.1169

547
548   Holland, H. D. (2002). Volcanic gases, black smokers, and the great oxidation event. *Geochimica Et*
549   *Cosmochimica Acta, 66*(21), 3811-3826. doi:https://doi.org/10.1016/S0016-7037(02)00950-X

550
551   Holland, H. D. (2006). The oxygenation of the atmosphere and oceans. *Philosophical transactions of the*
552   *Royal Society of London. Series B, Biological sciences, 361*(1470), 903-915.
553   doi:10.1098/rstb.2006.1838

554
555   Holland, H. D. (2009). Why the atmosphere became oxygenated: A proposal. *Geochimica Et*
556   *Cosmochimica Acta, 73*(18), 5241-5255. doi:https://doi.org/10.1016/j.gca.2009.05.070

557
558   Johnson, C. M., Beard, B. L., & Roden, E. E. (2008). The Iron Isotope Fingerprints of Redox and
559   Biogeochemical Cycling in Modern and Ancient Earth. *Annual Review of Earth and Planetary*
560   *Sciences, 36*(1), 457-493. doi:10.1146/annurev.earth.36.031207.124139

561
562   Johnston, D. T., Wolfe-Simon, F., Pearson, A., & Knoll, A. H. (2009). Anoxygenic photosynthesis
563   modulated Proterozoic oxygen and sustained Earth's middle age. *Proc Natl Acad Sci USA,*
564   *106*(40), 16925-16929. doi:10.1073/pnas.0909248106

565
566   Klotz, M. G., & Loewen, P. C. (2003). The molecular evolution of catalatic hydroperoxidases: evidence for
567   multiple lateral transfer of genes between prokaryota and from bacteria into eukaryota. *Mol*
568   *Biol Evol, 20*(7), 1098-1112. doi:10.1093/molbev/msg129

569
570   Konstantinidis, K. T., Serres, M. H., Romine, M. F., Rodrigues, J. L. M., Auchtung, J., Mccue, L. A., . . .
571   Tiedje, J. M. (2009). Comparative systems biology across an evolutionary gradient within the
572   Shewanella genus. *Proceedings of the National Academy of Sciences of the United States of*
573   *America, 106*(37), 15909-15914. doi:10.1073/pnas.0902000106

574
575   Kumar, S., Stecher, G., Li, M., Knyaz, C., & Tamura, K. (2018). MEGA X: Molecular Evolutionary Genetics
576   Analysis across Computing Platforms. *Mol Biol Evol, 35*(6), 1547-1549.
577   doi:10.1093/molbev/msy096

578
579   Lalonde, S. V., & Konhauser, K. O. (2015). Benthic perspective on Earth's oldest evidence for oxygenic
580   photosynthesis. *Proceedings of the National Academy of Sciences, 112*(4), 995-1000.
581   doi:10.1073/pnas.1415718112

583   Lee, Yeung, L. Y., McKenzie, N. R., Yokoyama, Y., Ozaki, K., & Lenardic, A. (2016). Two-step rise of
584   atmospheric oxygen linked to the growth of continents. *Nature Geoscience, 9*, 417.
585   doi:10.1038/ngeo2707

586
587   Lee, M., Walworth, N. G., McParland, E. L., Fu, F.-X., Mincer, T. J., Levine, N. M., . . . Webb, E. A. (2017).
588   The Trichodesmium consortium: conserved heterotrophic co-occurrence and genomic
589   signatures of potential interactions. *The Isme Journal, 11*(8), 1813-1824.
590   doi:10.1038/ismej.2017.49

591
592   Luo, G., Ono, S., Beukes, N. J., Wang, D. T., Xie, S., & Summons, R. E. (2016). Rapid oxygenation of Earth's
593   atmosphere 2.33 billion years ago. *Science Advances, 2*(5), e1600134.
594   doi:10.1126/sciadv.1600134

595
596   Lyons, T. W., Reinhard, C. T., & Planavsky, N. J. (2014). The rise of oxygen in Earth's early ocean and
597   atmosphere. *Nature, 506*(7488), 307-315. doi:10.1038/nature13068

598
599   Marin, J., Battistuzzi, F. U., Brown, A. C., & Hedges, S. B. (2017). The Timetree of Prokaryotes: New
600   Insights into Their Evolution and Speciation. *Mol Biol Evol, 34*(2), 437-446.
601   doi:10.1093/molbev/msw245

602
603   Martin, W. F., Bryant, D. A., & Beatty, J. T. (2018). A physiological perspective on the origin and evolution
604   of photosynthesis. *FEMS microbiology reviews, 42*(2), 205-231. doi:10.1093/femsre/fux056

605
606   Mishra, S., & Imlay, J. (2012). Why do bacteria use so many enzymes to scavenge hydrogen peroxide?
607   *Archives of Biochemistry and Biophysics, 525*(2), 145-160. doi:10.1016/j.abb.2012.04.014

608
609   Morris, J. J., Johnson, Z. I., Szul, M. J., Keller, M., & Zinser, E. R. (2011). Dependence of the
610   cyanobacterium Prochlorococcus on hydrogen peroxide scavenging microbes for growth at the
611   ocean's surface. *PLoS One, 6*(2), e16805. doi:10.1371/journal.pone.0016805

612
613   Morris, J. J., Lenski, R. E., & Zinser, E. R. (2012). The Black Queen Hypothesis: evolution of dependencies
614   through adaptive gene loss. *MBio, 3*(2). doi:10.1128/mBio.00036-12

615
616   Naclerio, G., Baccigalupi, L., Caruso, C., De Felice, M., & Ricca, E. (1995). Bacillus subtilis Vegetative
617   Catalase Is an Extracellular Enzyme. *Appl Environ Microbiol, 61*(12), 4471-4473.

618
619   Olson, K. R., Gao, Y., DeLeon, E. R., Arif, M., Arif, F., Arora, N., & Straub, K. D. (2017). Catalase as a
620   sulfide-sulfur oxido-reductase: An ancient (and modern?) regulator of reactive sulfur species
621   (RSS). *Redox Biology, 12*(Supplement C), 325-339.
622   doi:https://doi.org/10.1016/j.redox.2017.02.021

Ossa Ossa, F., Hofmann, A., Wille, M., Spangenberg, J. E., Bekker, A., Poulton, S. W., . . . Schoenberg, R.
(2018). Aerobic iron and manganese cycling in a redox-stratified Mesoarchean epicontinental
sea. *Earth and Planetary Science Letters, 500*, 28-40.
doi:https://doi.org/10.1016/j.epsl.2018.07.044

Ozaki, K., Thompson, K. J., Simister, R. L., Crowe, S. A., & Reinhard, C. T. (2019). Anoxygenic
photosynthesis and the delayed oxygenation of Earth's atmosphere. *Nature Communications,
10*(1), 3026. doi:10.1038/s41467-019-10872-z

Özkaya, Ö., Xavier, K. B., Dionisio, F., & Balbontín, R. (2017). Maintenance of Microbial Cooperation
Mediated by Public Goods in Single- and Multiple-Trait Scenarios. *Journal of Bacteriology,
199*(22), e00297-00217. doi:10.1128/jb.00297-17

Pezzoni, M., Pizarro, R. A., & Costa, C. S. (2014). Protective role of extracellular catalase (KatA) against
UVA radiation in Pseudomonas aeruginosa biofilms. *J Photochem Photobiol B, 131*, 53-64.
doi:10.1016/j.jphotobiol.2014.01.005

Planavsky, N. J., Asael, D., Hofmann, A., Reinhard, C. T., Lalonde, S. V., Knudsen, A., . . . Rouxel, O. J.
(2014). Evidence for oxygenic photosynthesis half a billion years before the Great Oxidation
Event. *Nature Geoscience, 7*(4), 283-286. doi:10.1038/ngeo2122

Reinhard, C. T., Planavsky, N. J., Gill, B. C., Ozaki, K., Robbins, L. J., Lyons, T. W., . . . Konhauser, K. O.
(2017). Evolution of the global phosphorus cycle. *Nature, 541*(7637), 386-389.
doi:10.1038/nature20772

Rodríguez Amor, D., & Dal Bello, M. (2019). Bottom-Up Approaches to Synthetic Cooperation in
Microbial Communities. *Life (Basel, Switzerland), 9*(1), 22. doi:10.3390/life9010022

Sanchez-Baracaldo, P., Bianchini, G., Di Cesare, A., Callieri, C., & Chrismas, N. A. M. (2019). Insights Into
the Evolution of Picocyanobacteria and Phycoerythrin Genes (mpeBA and cpeBA). *Front
Microbiol, 10*, 45. doi:10.3389/fmicb.2019.00045

Sekar, R., Shin, H. D., & DiChristina, T. J. (2016). Direct conversion of cellulose and hemicellulose to
fermentable sugars by a microbially-driven Fenton reaction. *Bioresour Technol, 218*, 1133-1139.
doi:10.1016/j.biortech.2016.07.087

Shcolnick, S., Summerfield, T. C., Reytman, L., Sherman, L. A., & Keren, N. (2009). The Mechanism of Iron
Homeostasis in the Unicellular Cyanobacterium Synechocystis sp. PCC 6803 and Its Relationship
to Oxidative Stress. *Plant Physiology, 150*(4), 2045-2056. doi:10.1104/pp.109.141853

Shi, L., Dong, H., Reguera, G., Beyenal, H., Lu, A., Liu, J., . . . Fredrickson, J. K. (2016). Extracellular
electron transfer mechanisms between microorganisms and minerals. *Nat Rev Microbiol, 14*(10),
651-662. doi:10.1038/nrmicro.2016.93

Shin, D.-H., Choi, Y.-S., & Cho, Y.-H. (2008). Unusual Properties of Catalase A (KatA) of
Pseudomonas aeruginosa PA14 Are Associated with Its Biofilm Peroxide Resistance.
Journal of Bacteriology, 190(8), 2663-2670. doi:10.1128/jb.01580-07

Song, H., Jiang, G., Poulton, S. W., Wignall, P. B., Tong, J., Song, H., . . . Wang, C. (2017). The onset of
widespread marine red beds and the evolution of ferruginous oceans. *Nature Communications,*
8(1), 399. doi:10.1038/s41467-017-00502-x

Soo, R. M., Hemp, J., Parks, D. H., Fischer, W. W., & Hugenholtz, P. (2017). On the origins of oxygenic
photosynthesis and aerobic respiration in Cyanobacteria. *Science, 355*(6332), 1436-1440.
doi:10.1126/science.aal3794

Stevens Jr., S. E., Patterson, C. O. P., & Myers, J. (1973). THE PRODUCTION OF HYDROGEN PEROXIDE BY
BLUE-GREEN ALGAE: A SURVEY1. *Journal of Phycology, 9*(4), 427-430. doi:10.1111/j.1529-
8817.1973.tb04116.x

Stookey, L. L. (1970). Ferrozine---a new spectrophotometric reagent for iron. *Analytical Chemistry, 42*(7),
779-781. doi:10.1021/ac60289a016

Swanner, E. D., Mloszewska, A. M., Cirpka, O. A., Schoenberg, R., Konhauser, K. O., & Kappler, A. (2015).
Modulation of oxygen production in Archaean oceans by episodes of Fe(II) toxicity. *Nature*
Geosci, 8(2), 126-130. doi:10.1038/ngeo2327

Wang, S., Deng, K., Zaremba, S., Deng, X., Lin, C., Wang, Q., . . . Zhang, W. (2009). Transcriptomic
Response of *Escherichia coli* O157:H7 to Oxidative Stress. *Applied and*
Environmental Microbiology, 75(19), 6110-6123. doi:10.1128/aem.00914-09

Weiss, M. C., Sousa, F. L., Mrnjavac, N., Neukirchen, S., Roettger, M., Nelson-Sathi, S., & Martin, W. F.
(2016). The physiology and habitat of the last universal common ancestor. *Nat Microbiol, 1*(9),
16116. doi:10.1038/nmicrobiol.2016.116

Widder, S., Allen, R. J., Pfeiffer, T., Curtis, T. P., Wiuf, C., Sloan, W. T., . . . Soyer, O. S. (2016). Challenges
in microbial ecology: building predictive understanding of community function and dynamics.
The Isme Journal, 10(11), 2557-2568. doi:10.1038/ismej.2016.45

Zamocky, M., Furtmuller, P. G., & Obinger, C. (2008). Evolution of catalases from bacteria to humans.
Antioxid Redox Signal, 10(9), 1527-1548. doi:10.1089/ars.2008.2046

Zinser, E. R. (2018). Cross-protection from hydrogen peroxide by helper microbes: the impacts on the

cyanobacterium Prochlorococcus and other beneficiaries in marine communities. *Environ*

Microbiol Rep, 10(4), 399-411. doi:10.1111/1758-2229.12625

710