Alterations in odor hedonics in the Tg6799 Alzheimer's disease mouse model and the influence of sex

Elizabeth R. Roberts[1#], Amanda M. Dossat[1#], María del Mar Cortijo[1], Patrik Brundin[2], Daniel W. Wesson[1*]

[1]Department of Pharmacology & Therapeutics, University of Florida, 1200 Newell Dr., Gainesville, FL, 32610

[2]Van Andel Institute, 333 Bostwick Ave NE, Grand Rapids, MI 49503

#equal contributing first authors

*corresponding author: danielwesson@ufl.edu

RUNNING HEAD: Tg6799 mice and odor hedonics.

Acknowledgements: This work was supported by National Institutes of Health grants: NIDCD R01DC016519 (to D.W. and P.B.) and NIDCD R01DC014443 to D.W. (the latter including a supplement from the NIA to support research into Alzheimer's disease to P.B. and D.W.) and NIDA R01DA049545 and R01DA049449 (to D.W.).

Abstract

Olfactory impairments, including deficits in odor detection, discrimination, recognition, and changes in odor hedonics are reported in the early stages of Alzheimer's disease (AD). Rodent models of AD display deficits in odor learning, detection, and discrimination – recapitulating the clinical condition. However, the impact of familial AD genetic mutations on odor hedonics is unknown. We tested 2-, 4-, and 6-months old 5XFAD (Tg6799) mice in the five-port odor multiple-choice task designed to assay a variety of odor-guided behaviors, including odor preferences/hedonics. We found that 5XFAD mice investigated odors longer than controls, an effect that was driven by 6-months old mice. Interestingly, this effect was carried by females in the 5XFAD group, who investigated odors longer than age-matched males. Upon examining behavior directed towards individual odors to test for aberrant odor preferences, we uncovered that 5XFAD females at several ages displayed heightened preferences towards some of the odors, indicating aberrant hedonics. We observed no impairments in the ability to engage in the task in 5XFAD mice. Taken together, 5XFAD mice, particularly 5XFAD females, display prolonged odor investigation behavior and enhanced preferences to certain odors. The data provide insight into hedonic alterations which may occur in AD mouse models, and how these are influenced by biological sex.

Keywords: 5XFAD, olfaction, sex difference, valence, behavior

Introduction

Alzheimer's disease (AD) is a neurodegenerative disorder characterized by cognitive, motor, and sensory dysfunction. Olfactory dysfunction is a common feature of the AD prodrome, with patients displaying deficits in olfactory detection, discrimination, and recognition (Devanand et al., 2008; Doty, 2017; Claire Murphy, 2019; Rey et al., 2018). The emotional responses to odors, known as hedonics, are also disturbed in persons with AD (Joussain et al., 2015). While alterations in odor hedonics have received less attention in AD research than manifestations in perceptual impairments, this is a significant issue since odor preferences influence food selection and ingestive behaviors (C Murphy, 2008) which, if aberrant, might impact disease progression, and at a minimum, quality of life (Gopinath et al., 2012; Claire Murphy, 2019).

Mouse models of AD may provide insight into the origins of aberrant odor preferences in the disease. These mouse models have already yielded clues into some mechanisms of olfactory dysfunction and how these deficits might relate to disease progression. For instance, the Tg2576 mouse model of AD, which over-expresses two familial mutations in the amyloid precursor protein gene (*APP*), displays impairments in short-term odor learning and odor discrimination (D W Wesson et al., 2010). These deficits are associated with alterations in neural activity in the olfactory bulb and piriform cortex – two important regions of the olfactory system (Daniel W Wesson et al., 2011). Transient over-expression of mutant *APP* in solely the olfactory sensory neurons in the nasal epithelium is sufficient to impair olfactory perception (Cheng et al., 2011). While not well understood, it is possible that products of the *APP* gene, and/or other genes implicated in AD, impact olfactory system physiology and thus perturb perception (Morales-Corraliza et al., 2012; D W Wesson et al., 2010; Daniel W Wesson et al., 2011; Xu et al., 2015; Yoo et al., 2017). Whether animal models which express these pathologies have alterations in their odor hedonics is unknown.

We sought to investigate odor preferences in an AD mouse model. To address this, we used the 5XFAD (Tg6799) model which overexpresses five familial AD mutations and thus is considered an accelerated AD model (Oakley et al., 2006). The 5XFAD mouse model displays progressive AD pathology throughout the brain, starting at just two months

of age (Oakley et al., 2006), which differentially burdens males and females (Bundy et al., 2019). This includes amyloid-β burden in some key olfactory structures, such as the olfactory bulb and piriform cortex (Oakley et al., 2006; Wang et al., 2017; Xiao et al., 2015). Further, this mouse model shows impairments in synaptic plasticity and cognition (Devi & Ohno, 2016; Kimura & Ohno, 2009; Oakley et al., 2006). Importantly, 5XFAD mice appear to exhibit normal olfactory detection abilities, as assayed in a sensitive psychophysical assay (Roddick et al., 2016). However, the possibility that despite intact sensitivity for odors, these mice may have aberrant tendencies to investigate odors and possibly altered hedonics to select odors, has not yet been investigated. In both humans and mice, the duration of time spent investigating odors reflects the attractiveness or preference of an odor reflective of its hedonics (Baum & Keverne, 2002; Mandairon et al., 2009). Therefore, we monitored the duration of time 5XFAD mice spent investigating odors in a five-port odor multiple-choice task, which is a sensitive and semi-automated assay of odor preferences (Jagetia et al., 2018). We did so grouping across all odors to test for general tendencies to investigate odors, and separately, within individual odors to explore possible alterations in hedonics. Our results show that 5XFAD mice, particularly 5XFAD females, display prolonged odor investigation behavior and enhanced preferences to certain odors.

Materials & Methods

Subjects

We used ninety-six 5XFAD/B6SJL-Tg6799 (5XFAD) mice and age-matched nontransgenic (NTg) littermates (2-, 4-, and 6-months of age; 48 males and 48 females). Breeder stock were obtained from The Jackson Laboratory (stock #006554; Bar Harbor, ME) and their offspring were raised and maintained at the University of Florida. Animals were genotyped using PCR of tail clip DNA upon weaning, and genotypes were confirmed with a second round of genotyping after completion of behavioral testing. All experiments were performed in accordance with the guidelines of the National Institutes of Health and were approved by the University of Florida Institutional Animal Care and Use Committee. From weaning onward, all animals were group-housed (2-5 per cage), placed on a

reversed 12:12h light cycle, and allowed *ad libitum* access to food and water. Cohorts
ranging from four to 16 mice were pseudo-randomly assigned to maximize diversity
across sex, age, and genotype. All behavioral testing occurred during the light phase
(0900:1800 hr).

5-port odor preference apparatus

We used the five-port odor multiple-choice behavioral apparatus as we previously
described in detail (Jagetia et al., 2018). Briefly, this apparatus allows animals to freely
investigate different 3D-printed odor ports and provides automated, millisecond-precise
monitoring of nose-poking behavior towards odors by means of infrared photobeams. In
this apparatus, animals gain access to the odor by nose-poking into a port with nose-
pokes detected by an 880 nm infrared beam positioned at the entrance of the odor port.
The odor ports were designed to scavenge odor out of the port, thereby requiring the
animal to poke to gain access to the odor (Jagetia et al., 2018). On the testing day, 20 µL
of liquid odor was dispensed into a flange cap (MOCAP, Park Hills, MO) and inserted into
the bottom of a port. The onset, duration, and location (port identity) of nose-pokes were
recorded by a microcontroller (Arduino; www.arduino.cc), digitized, and stored on a
computer for subsequent analysis.

Five-port odor multiple-choice preference testing

On acclimation and testing days alike, the mice were brought into the testing room where
they were allowed to acclimate in their home cages prior to being placed in the testing
chamber. The mice were allowed to explore the five-port chamber for increasing durations
over three consecutive days (10, 20, and 30 minutes, respectively). During acclimation,
clean plastic flange caps were placed in the bottom of the odor ports. Within two days
following acclimation, 30 minute testing sessions were conducted over five consecutive
days. The same five odors were presented every day, with each odor being placed in a
different port each day to minimize the influence of spatial location on odor investigation

(Jagetia et al., 2018). The apparatus was cleaned with 70% ethanol before onset of

testing and after each mouse was removed from the chamber.

Odors

Odors were purchased at their highest available purity (>97%) and diluted to 1 Pa in light

mineral oil (Sigma Aldrich; St. Louis, MO) prior to placing in the flange caps / odor ports

and included L-carvone, decanol, geraniol, guaiacol, and thioglycolic acid (Sigma Aldrich;

St. Louis, MO). Odors in their stock concentration and diluted concentration were each

stored under nitrogen to maintain purity. The panel of odors selected was based upon our

prior work (Jagetia et al., 2018), wherein we found that [at least in a different background

strain of mice] these odors elicited the widest assortment of odor preferences, with

decanol and thioglycolic acid being investigated the least and most, respectively.

Data Analysis

Individual animals' behavior was evaluated by the duration of nose-pokes (*viz.*, odor

investigation), nose-poke number, and inter-poke interval. Statistical analyses were

performed in Prism 8.0 (GraphPad Software; San Diego, CA), included mixed-effects

models with Geisser-Greenhouse corrections and two-way analysis of variance

(ANOVA), and were followed by Sidak's post hoc tests where indicated. Two-tailed

unpaired t-tests were used to determine effects of genotype within each sex and age

group. Any values that were > 2 standard deviations beyond the group mean were

classified as outliers and removed from the data set. Across variables there was a range

of 0-3 outliers per group, represented similarly across all genotypes, ages, and sexes.

Over testing days, mice robustly habituate to odors in this task (Jagetia et al., 2018). In

the data set from the present study we found that all groups of mice habituated over the

five testing days so that by the 5^{th} day very little odor investigation occurred. Indeed, there

was an effect of mice to display shorter odor investigation durations on day 5 vs. day 1

(NTg: $t(85)=7.523$, $p<0.0001$; 5XFAD: $t(85)=10.76$, $p<0.0001$). Therefore, in order to

investigate possible changes in odor investigation and odor preferences between

genotypes (as well as sex and age effects), we restricted our analyses to the behavior

from day 1 of testing since this allows for examination of animals' response to odors when

they are highest.

Results

5XFAD mice investigate odors longer than controls.

We first analyzed the duration of odor investigation (cumulative, throughout the entire

session) with data from all odors grouped, prior to determining if there was unique

preference towards individual odors in later sections of the results. Examination of odor

investigation durations revealed an effect of genotype (t(42)=2.98, p=0.005, collapsing all

ages and sexes; **Figure 1A**) wherein 5XFAD mice exhibited longer odor investigation

durations vs. NTg mice. We then separated the data in order to investigate the

contributions of age upon the effect for 5XFAD mice to investigate odors longer. 6-months

old 5XFAD mice engaged in odor investigation longer than age-matched NTg

(t(13)=2.279, p=0.040; **Figures 1B – 1D**). There was no effect of genotype in 2-months

and 4-months old mice (p>0.05). These results indicate that 5XFAD mutations and aging

interact to influence odor investigation behavior.

Females investigate odors longer than males.

Due to established sex differences in AD (Nebel et al., 2018), and in pathological

accumulation in this mouse model (Bundy et al., 2019), we sought to determine if sex

influenced the above outcomes. Females exhibited longer odor investigation durations

vs. males (t(92)=4.302, p<0.0001, collapsing ages and genotypes; **Figure 2A**). As above,

we next separated the data to investigate the contributions of age upon the sex difference

in odor investigation time. We found that 4- and 6-months old female mice exhibited

longer odor investigation durations vs. age-matched males (t(29)=2.099, p=0.045;

t(28)=5.314, p<0.0001, respectively, **Figure 2C & 2D**), with no sex difference at 2-months

of age (p>0.05). These data show that females exhibit longer odor investigation durations
vs. males, an effect carried by the older age groups.

5XFAD genotype impacts preferences towards individual odors.

In all of the above analyses, odor investigation durations were collapsed across all odors
as an assay for the tendencies of mice to investigate odors. However, collapsing data
across all stimuli from a panel of molecularly diverse odors does not indicate changes
towards a given odor which is essential in order to identify changes in odor preferences.
Therefore, we next looked for possible influences of 5XFAD genotype on preferences
towards specific odors.

As expected based upon the results in **Figure 1A**, this analysis revealed a main
effect of genotype to influence duration of odor investigation across all age groups ($F_{(1, 456)}$=20.07, p<0.0001, **Figure 3A**). 5XFAD mice exhibited a stronger preference for
carvone and decanol vs. NTg control (t(456)=3.419, p=0.003; t(456)=3.233, p=0.007,
respectively) with no effect for the other odors (p>0.05 for each).

When separating the data by age, we observed a main effect of genotype to
influence duration of odor investigation in 2-months and 6-months mice ($F_{(1,146)}$=4.92,
p=0.028; $F_{(1,147)}$=12.54, p<0.001, respectively) but not in 4-months mice (p>0.05).
However, post hoc tests failed to uncover 'attraction' to an individual odor when testing
within age groups for effects of genotype (p>0.05 for each).

Sex and genotype impact preferences towards individual odors.

While we found an effect of 5XFAD genotype on preferences towards odors, it is possible
an effect was influenced by the pronounced sex difference wherein females exhibit longer
odor investigation (as in **Figure 2**). Therefore, we next investigated if sex of the 5XFAD
mice influences odor preferences.

There was an effect of sex to influence preferences for all odors (carvone: $F_{(1, 89)}$=21.49, p<0.0001; decanol: $F_{(1, 89)}$=10.02, p=0.002; geraniol: $F_{(1, 89)}$=14.89, p<0.001;

guaiacol: $F_{(1, 90)}=14.91$, $p<0.001$; thioglycolic acid: $F_{(1, 88)}=7.895$, $p=0.006$; **Figure 4)**. NTg

females exhibited greater preferences vs. NTg males for some odors (carvone:

$t(89)=2.454$, $p=0.032$; guaiacol: $t(90)=2.496$, $p=0.029$; $p>0.05$ for decanol, geraniol, and

thioglycolic acid). Further, 5XFAD females exhibited greater preferences for most odors

vs. 5XFAD males (carvone: $t(89)=4.094$, $p<0.001$; decanol: $t(89)=3.323$, $p=0.003$;

geraniol: $t(89)=4.017$, $p=0.002$; guaiacol: $t(90)=2.961$, $p=0.008$; $p>0.05$ for thioglycolic

acid). These data are in-line with our above findings from total odor investigation duration

(**Figure 2**), where females investigated odors longer than males. The above analyses

within individual odors indicates that sex influences odor preferences, as is well

established in other mouse lines/strains (*e.g.,* (Baum & Keverne, 2002)).

5XFAD genotype also influenced preferences to some of the odors tested

(carvone: $F_{(1, 89)}=14.79$, $p<0.001$; decanol: $F_{(1, 89)}=12.3$, $p<0.001$). Post hoc analysis

revealed that the effect of genotype was carried by 5XFAD females, which displayed

greater preferences vs. NTg females for carvone ($t(89)=3.539$, $p=0.001$) and decanol

($t(89)=3.526$, $p=0.001$; $p>0.05$ for geraniol, guaiacol, and thioglycolic acid). Overall, these

data indicate that females within each age group exhibit more preference towards most

odors vs. males. Further, in some cases, the enhanced preferences displayed by females

is influenced by 5XFAD genotype.

As a final analysis into preference towards individual odors, we sought to test

whether there was also an influence of age. Having uncovered that preference towards

two of the odors in the panel was impacted by genotype and sex (**Figure 4A & 4B**), this

resulted in the need of comparing possible effects of age not just within odor types, but

also within genotypes (**Figure 5**).

This analysis revealed that within the 2-months females, there was an effect of

genotype on preference for geraniol ($t(13)=2.177$, $p=0.049$) with no effect on any other

odor ($p>0.05$ for all). In 6-months females, 5XFAD genotype influenced preference for

decanol and guaiacol ($t(13)=2.745$, $p=0.017$; $t(14)=2.29$, $p=0.038$, respectively). There

was no effect of genotype in 4-months females ($p>0.05$). Within males there was an effect

of genotype wherein 5XFAD 4-months mice more greatly preferred carvone vs NTg

($t(14)=3.135$, $p=0.007$; $p>0.05$ for all other odors). There was no effect of genotype in 2-

months and 6-months age groups (p>0.05 for all). These data indicate that females
largely drive effects of 5XFAD mutations on odor investigation behaviors, particularly in
the 6-months age group. Taken together, 5XFAD mice, especially females, show aberrant
odor preferences.

No effect of 5XFAD genotype on measures of general task engagement.

Finally, we performed two analyses on the above data as simple 'controls' to indicate the
mice, regardless of group, were capable of engaging in the task. More greatly aged
5XFAD mice are known to have impaired motor function (O'Leary et al., 2018, 2020) and
changes in the ability of mice to nose-poke would confound use of this measure as a
variable whereby to assay odor investigation and preferences. We first examined inter-
poke interval, which is indicative of an animal's ability to coordinate motor movements
from one odor port to another. There was neither an effect of genotype nor sex to
influence inter-poke intervals in both 2- and 4-month old mice (p>0.05 each; **Figs 6A &**
6B). In 6-month old mice there is an effect of sex ($F_{(1, 26)}$=5.307, p=0.03; **Figure 6C**), but
not genotype (p>0.05), to shorten inter-poke interval.

An additional index of general task engagement is the total number of nose-pokes
animals execute within a session. There was neither an effect of genotype nor sex to
influence nose-poke count in both 2- and 4-months old mice (p>0.05 each; **Figure 6D &**
6E). 6-months old mice exhibited a significant effect of sex ($F_{(1, 28)}$=5.212, p=0.03; **Figure**
6F) but not genotype (p>0.05), to increase nose-poke count. Reconciling these outcomes,
these significant differences however do not support impairments in the task. Indeed, a
greater number of pokes in 6-months old female 5XFAD mice (**Figure 6C**), together with
their reduced inter-poke intervals (**Figure 6F**), point towards increased task engagement.
Taken together, data from both analyses suggest the differences in odor investigation
and preferences observed in the present study are not due to deficits in task engagement.

Discussion

We used the 5XFAD mouse AD model in order to define odor investigation behavior using
the five-port odor multiple-choice task. Our long-term goal is to understand how odor

hedonics and possibly other affective symptoms of AD result mechanistically. To this end,

we assessed if odor preferences, reflected by the time sampling individual odors, are

perturbed in the 5XFAD mouse AD model. We found, paradoxical to what is known in

persons with AD to display reduced hedonic ratings of odors (Joussain et al., 2015), that

5XFAD mice sample certain odors longer, reflective of increased hedonics.

No gross behavioral impairments are detected in 5XFAD mice in the 5-port odor multiple

choice task.

As an important starting point, our results indicate that 5XFAD mice do not display overt

impairments in their ability to engage in the task we used (**Figure 6**). This suggests that

5XFAD mice, at the ages we tested, have intact motor function at least as it is required to

coordinate nose-pokes in this task. A robust literature suggests that there are significant

motor impairments in these mice occurring at much later ages (O'Leary et al., 2018,

2020). If anything, female mice in our most aged group (6-months) showed changes in

task engagement measures suggesting *increased* engagement in the task (engaged

more frequently and promptly between pokes; **Figure 6**), yet this was not influenced by

genotype.

We also uncovered that 2-6-months old 5XFAD mice of both sexes are capable of

engaging with odors (**Figure 1**). This latter outcome suggests the 5XFAD mice do not

have grossly impaired abilities to detect and discriminate among odors since if they would

simply generalize across all odors, one would predict to display similar investigation

regardless of the odor. In contrast, our analyses among the individual odors (**Figures 4-**

5) support that mice can differentiate among the odors we used. These results at their

most basic level are consistent with a report from one group that measured odor detection

thresholds using sophisticated olfactometry in 5XFAD mice (Roddick et al., 2016). Using

the go/no-go olfactometer as an assay, Roddick and colleagues quantified odor detection

thresholds and concluded that odor sensitivity is not dramatically impaired in 6-months

old 5XFAD mice (Roddick et al., 2016). A study by a separate group used the buried food

pellet test to perform testing of odor-influenced food motivation behavior (Xiao et al.,

2015). The work by Xiao indicated possible deficits in food-finding behavior at three

months of age which may suggest deficits in olfaction. However, this test is difficult to use
as a specific metric of olfaction since it relies upon the ability for animals to coordinate
foraging behavior which is fundamentally regulated by states of motivation. Overall, the
results of the present study together with especially that of Roddick et al., (2016) highlight
that the most basic aspects of olfaction are largely preserved in 5XFAD mice. This
outcome is perhaps surprising given that the brains of 6-months old 5XFAD mice are
burdened with extensive AD pathologies (Devi & Ohno, 2016; Oakley et al., 2006),
including within the olfactory bulbs themselves (Oakley et al., 2006; Wang et al., 2017;
Xiao et al., 2015).

Alterations in odor investigation behavior and odor preferences in 5XFAD mice.

Collapsing across all odors, our results demonstrate that 5XFAD mice, all ages and both
sexes included, investigate odors longer than NTg controls (**Figure 1**). This was due to
longer investigation behavior specifically in the 6-month old animals, but not the 2- and
4-months mice. We used these data pooled across all odors as a simple metric of
generalized odor investigation behavior. Importantly, generalized odor investigation
behavior may not be used to understand hedonics necessarily, since this may also be
resultant from enhanced arousal and/or novelty behavior. Nevertheless, it is informative
to consider the tendency for 5XFAD mice, especially 6-months old female 5XFAD mice,
to display enhanced propensities to investigate odors.

By analyzing the data within individual odors grouped by sex and age, we
uncovered several examples supporting our conclusion that 5XFAD mice display altered
hedonics to certain odors. Two of the five odors employed (carvone and decanol) often
entailed aberrant investigation responses in 5XFAD mice, but did so depending on age
and sex. Notably, we anticipate the uncovered changes in odor hedonics of the 5XFAD
mice may be more prevalent than that found herein. Our use of a simple panel of five
odors was selected to provide a spectrum of odor/chemical types to thereby yield a first-
pass screen of olfactory preferences in 5XFAD mice. Due the extensive numbers of
animals tested in this study, and testing labor this study required, we did not also include
additional panels of odors to further assay olfactory abilities. Certainly, investigating the

behavior of 5XFAD mice in the 5-port task with differing panels of odors is an important future goal to provide a complete assessment of the magnitude of odor preference alterations. One reason for this is that the differences in preferences we found were not highly overt like one would expect to see if a mouse was allowed to investigate conspecific urine or a food source odor versus that of a predator. Instead, the preferences were subtle and suggest modest, although functionally significant at a behavioral level, changes in affective brain systems supporting odor preferences. In the future it will be important to understand what brain systems might be impacted in these animals which results in altered preferences to select odors.

Perplexingly, while in persons with AD, hedonic ratings towards odors are notably reduced (Joussain et al., 2015), we found that 5XFAD mice show opposite responses when engaging with odors during altered hedonic states. Indeed, our results show that in some cases, 5XFAD mice investigate odors longer, and specific odors longer, than control mice (**Figures 4 & 5**). While increases in odor investigation are known to be reliable measures of odor preferences in both humans and mice (Mandairon et al., 2009), why 5XFAD mice would display increased hedonic responses to odors is unclear. It is possible that an AD pathologically-burdened mouse model displays hedonic responses to odors in fundamentally different manners than a human.

Sex influences on odor investigation and preferences.

Olfaction is influenced by sex and gonadal hormones (Dillon et al., 2013; Doty & Cameron, 2009; Good et al., 1976; Kass et al., 2017; Sorwell et al., 2008; D W Wesson et al., 2006). This also holds true to odor preferences (Baum & Keverne, 2002; Beach, 1974). In line with this, we found that females exhibited longer durations of odor investigation compared to males specifically at 4- and 6-months of age (**Figure 2**). We also found that females across all ages and in both genotypes exhibited higher preference for most of the odors as compared to males (**Figure 4**). AD differentially impacts males and females (Nebel et al., 2018), and this is somewhat recapitulated in neuronal pathology in the 5XFAD mouse model wherein females exhibit more AD-associated pathology compared to males (Bundy et al., 2019). Indeed, the longer odor investigation

duration of the 5XFAD mice was driven primarily by the behavior of the females.
Consistent with this, male and female 5XFAD mice display different odor learning curves,
yet comparable odor detection abilities (Roddick et al., 2016). Taken together, these
results emphasize that it is important to consider sex as a biological variable in AD model
research, especially given the established sex differences in human AD presentation,
development, and progression (Nebel et al., 2018).

Conclusions.

Our results show that young adult 5XFAD mice investigate odors longer than NTg
controls. This effect was not simply due to genotype. Indeed, prolonged odor investigation
in 5XFAD mice was influenced significantly by both age and sex with females
investigating odors longer than males regardless of genotype. These results provide
some evidence for altered odor investigation behaviors and further, odor preferences, in
an AD mouse model, the latter of which was surprisingly increased among 5XFAD mice
towards some odors. Future work employing larger odor sets are needed to best define
the nature of these changes in preferences, as are studies to explore their mechanistic
causes.

Competing interests: P.B. has received commercial support as a consultant from Axial Biotherapeutics, CuraSen, Fujifilm-Cellular Dynamics International, IOS Press Partners, LifeSci Capital LLC, Lundbeck A/S and Living Cell Technologies LTD. He has received commercial support for grants/research from Lundbeck A/S and Roche. He has ownership interests in Acousort AB and Axial Biotherapeutics and is on the steering committee of the NILO-PD trial. The authors declare no additional competing financial interest.

References

Baum, M. J., & Keverne, E. B. (2002). Sex difference in attraction thresholds for volatile odors from male and estrous female mouse urine. *Hormones and Behavior*, *41*(2), 213–219.

Beach, F. A. (1974). Effects of gonadal hormones on urinary behavior in dogs. *Physiology and Behavior*, *12*(6), 1005–1013.

Bundy, J. L., Vied, C., Badger, C., & Nowakowski, R. S. (2019). Sex-biased hippocampal pathology in the 5XFAD mouse model of Alzheimer's disease: A multi-omic analysis. *The Journal of Comparative Neurology*, *527*(2), 462–475. https://doi.org/10.1002/cne.24551

Cheng, N., Cai, H., & Belluscio, L. (2011). In Vivo Olfactory Model of APP-Induced Neurodegeneration Reveals a Reversible Cell-Autonomous Function. *The Journal of Neuroscience*, *31*(39), 13699–13704. https://doi.org/10.1523/jneurosci.1714-11.2011

Devanand, D. P., Liu, X., Tabert, M. H., Pradhaban, G., Cuasay, K., Bell, K., de Leon, M. J., Doty, R. L., Stern, Y., & Pelton, G. H. (2008). Combining Early Markers Strongly Predicts Conversion from Mild Cognitive Impairment to Alzheimer's Disease. *Biological Psychiatry*, *64*(10), 871–879. http://www.sciencedirect.com/science/article/B6T4S-4T8TW52-1/2/166fa8cd739a0242ecc9096da4f9e303

Devi, L., & Ohno, M. (2016). Cognitive benefits of memantine in Alzheimer's 5XFAD model mice decline during advanced disease stages. *Pharmacology Biochemistry and Behavior*, *144*, 60–66. https://doi.org/https://doi.org/10.1016/j.pbb.2016.03.002

Dillon, T. S., Fox, L. C., Han, C., & Linster, C. (2013). 17β-estradiol enhances memory duration in the main olfactory bulb in CD-1 mice. *Behavioral Neuroscience*, *127*(6), 923–931. https://doi.org/10.1037/a0034839

Doty, R. L. (2017). Olfactory dysfunction in neurodegenerative diseases: is there a common pathological substrate? *The Lancet Neurology*, *16*(6), 478–488.

https://doi.org/https://doi.org/10.1016/S1474-4422(17)30123-0

Doty, R. L., & Cameron, E. L. (2009). Sex differences and reproductive hormone
influences on human odor perception. *Physiology & Behavior*, *97*(2), 213–228.
https://doi.org/10.1016/j.physbeh.2009.02.032

Good, P. R., Geary, N., & Engen, T. (1976). The effect of estrogen on odor detection.
Chemical Senses and Flavor, *2*, 45–50.

Gopinath, B., Anstey, K. J., Kifley, A., & Mitchell, P. (2012). Olfactory impairment is
associated with functional disability and reduced independence among older
adults. *Maturitas*, *72*(1), 50–55. https://doi.org/10.1016/j.maturitas.2012.01.009

Jagetia, S., Milton, A. J., Stetzik, L. A., Liu, S., Pai, K., Arakawa, K., Mandairon, N., &
Wesson, D. W. (2018). Inter- and intra-mouse variability in odor preferences
revealed in an olfactory multiple-choice test. In *Behavioral Neuroscience* (Vol. 132,
Issue 2, pp. 88–98). American Psychological Association.
https://doi.org/10.1037/bne0000233

Joussain, P., Bessy, M., Fournel, A., Ferdenzi, C., Rouby, C., Delphin-Combe, F.,
Krolak-Salmon, P., & Bensafi, M. (2015). Altered Affective Evaluations of Smells in
Alzheimer's Disease. *Journal of Alzheimer's Disease*, *49*(2), 433–441.

Kass, M. D., Czarnecki, L. A., Moberly, A. H., & McGann, J. P. (2017). Differences in
peripheral sensory input to the olfactory bulb between male and female mice.
Scientific Reports, *7*(1), 45851. https://doi.org/10.1038/srep45851

Kimura, R., & Ohno, M. (2009). Impairments in remote memory stabilization precede
hippocampal synaptic and cognitive failures in 5XFAD Alzheimer mouse model.
Neurobiology of Disease, *33*(2), 229–235.
https://doi.org/https://doi.org/10.1016/j.nbd.2008.10.006

Mandairon, N., Poncelet, J., Bensafi, M., & Didier, A. (2009). Humans and Mice Express
Similar Olfactory Preferences. *PLoS ONE*, *4*(1), e4209.
https://doi.org/10.1371/journal.pone.0004209

Morales-Corraliza, J., Schmidt, S. D., Mazzella, M. J., Berger, J. D., Wilson, D. A.,

Wesson, D. W., Jucker, M., Levy, E., Nixon, R. A., & Mathews, P. M. (2012).

Immunization targeting a minor plaque constituent clears β-amyloid and rescues

behavioral deficits in an Alzheimer's disease mouse model. *Neurobiology of Aging*,

in press(0). https://doi.org/10.1016/j.neurobiolaging.2012.04.007

Murphy, C. (2008). The chemical senses and nutrition in older adults. *Journal of*

Nutrition For the Elderly, *27*(3–4), 247–265.

Murphy, Claire. (2019). Olfactory and other sensory impairments in Alzheimer disease.

Nature Reviews Neurology, *15*(1), 11–24. https://doi.org/10.1038/s41582-018-

0097-5

Nebel, R. A., Aggarwal, N. T., Barnes, L. L., Gallagher, A., Goldstein, J. M., Kantarci, K.,

Mallampalli, M. P., Mormino, E. C., Scott, L., Yu, W. H., Maki, P. M., & Mielke, M.

M. (2018). Understanding the impact of sex and gender in Alzheimer's disease: A

call to action. *Alzheimer's & Dementia : The Journal of the Alzheimer's Association*,

14(9), 1171–1183. https://doi.org/10.1016/j.jalz.2018.04.008

O'Leary, T., Mantolino, H., Stover, K., & Brown, R. (2020). Age-related deterioration of

motor function in male and female 5xFAD mice from 3 to 16 months of age. *Genes*

Brain and Behavior, *19*(3), e12538. https://doi.org/10.1111/gbb.12538

O'Leary, T., Robertson, A., Chipman, P. H., Rafuse, V. F., & Brown, R. E. (2018). Motor

function deficits in the 12 month-old female 5xFAD mouse model of Alzheimer's

disease. *Behavioural Brain Research*, *337*, 256–263.

https://doi.org/10.1016/j.bbr.2017.09.009

Oakley, H., Cole, S. L., Logan, S., Maus, E., Shao, P., Craft, J., Guillozet-Bongaarts, A.,

Ohno, M., Disterhoft, J., Van Eldik, L., Berry, R., & Vassar, R. (2006). Intraneuronal

beta-Amyloid Aggregates, Neurodegeneration, and Neuron Loss in Transgenic

Mice with Five Familial Alzheimer's Disease Mutations: Potential Factors in Amyloid

Plaque Formation. *Journal of Neuroscience*, *26*(40), 10129–10140.

https://doi.org/10.1523/jneurosci.1202-06.2006

Rey, N. L., Wesson, D. W., & Brundin, P. (2018). The olfactory bulb as the entry site for

prion-like propagation in neurodegenerative diseases. *Neurobiology of Disease*, *109*, 226–248. https://doi.org/http://dx.doi.org/10.1016/j.nbd.2016.12.013

Roddick, K. M., Roberts, A. D., Schellinck, H. M., & Brown, R. E. (2016). Sex and Genotype Differences in Odor Detection in the 3×Tg-AD and 5XFAD Mouse Models of Alzheimer's Disease at 6 Months of Age. *Chemical Senses*, *41*(5), 433–440. https://doi.org/10.1093/chemse/bjw018

Sorwell, K. G., Wesson, D. W., & Baum, M. J. (2008). Sexually dimorphic enhancement by estradiol of male urinary odor detection thresholds in mice. *Behavioral Neuroscience*, *122*(4), 788–793. https://doi.org/2008-09788-007 [pii] 10.1037/0735-7044.122.4.788

Wang, Y., Wu, Z., Bai, Y.-T., Wu, G.-Y., & Chen, G. (2017). Gad67 haploinsufficiency reduces amyloid pathology and rescues olfactory memory deficits in a mouse model of Alzheimer's disease. *Molecular Neurodegeneration*, *12*(1), 73. https://doi.org/10.1186/s13024-017-0213-9

Wesson, D W, Keller, M., Douhard, Q., Baum, M. J., & Bakker, J. (2006). Enhanced urinary odor discrimination in female aromatase knockout (ArKO) mice. *Hormones and Behavior*, *49*(5), 580–586.

Wesson, D W, Levy, E., Nixon, R. A., & Wilson, D. A. (2010). Olfactory Dysfunction Correlates with β-Amyloid Plaque Burden in an Alzheimer's Disease Mouse Model. *The Journal of Neuroscience*, *30*(2), 505–514.

Wesson, Daniel W, Borkowski, A. H., Landreth, G. E., Nixon, R. A., Levy, E., & Wilson, D. A. (2011). Sensory Network Dysfunction, Behavioral Impairments, and Their Reversibility in an Alzheimer's beta-Amyloidosis Mouse Model. *The Journal of Neuroscience*, *31*(44), 15962–15971. https://doi.org/10.1523/jneurosci.2085-11.2011

Xiao, N.-A., Zhang, J., Zhou, M., Wei, Z., Wu, X.-L., Dai, X.-M., Zhu, Y.-G., & Chen, X.-C. (2015). Reduction of Glucose Metabolism in Olfactory Bulb is an Earlier Alzheimer's Disease-related Biomarker in 5XFAD Mice. *Chinese Medical Journal*,

128(16), 2220–2227. https://doi.org/10.4103/0366-6999.162507

Xu, W., Fitzgerald, S., Nixon, R. A., Levy, E., & Wilson, D. A. (2015). Early hyperactivity
in lateral entorhinal cortex is associated with elevated levels of AβPP metabolites in
the Tg2576 mouse model of Alzheimer's disease. *Experimental Neurology*, *264*,
82–91. https://doi.org/https://doi.org/10.1016/j.expneurol.2014.12.008

Yoo, S.-J., Lee, J.-H., Kim, S. Y., Son, G., Kim, J. Y., Cho, B., Yu, S.-W., Chang, K.-A.,
Suh, Y.-H., & Moon, C. (2017). Differential spatial expression of peripheral olfactory
neuron-derived BACE1 induces olfactory impairment by region-specific
accumulation of β-amyloid oligomer. *Cell Death Disease*, *8*, e2977.
https://doi.org/10.1038/cddis.2017.349

Figures

Figure 1. 5XFAD mice investigate odors longer than controls. Cumulative odor investigation duration in **A,** all mice (n=44-48/group), and in **B,** 2-months, **C,** 4-months, and **E,** 6-months old mice (n=15-16/group). *p<0.05, **p<0.01 5XFAD Tg vs. NTg.

Figure 2. Females investigate odors longer than males. A, Cumulative odor investigation duration in males and females (n=46-48/group), and in **B,** 2-months, **C,** 4-months, and **D,** 6-months old mice with genotypes collapsed (n=14-16/group). *p<0.05, ****p<0.0001 vs. males.

594 **Figure 3. 5XFAD genotype impacts preferences displayed towards individual**

595 **odors.** Duration of odor investigation in **A,** all mice (n=45-48/group), and in **B,** 2-months,

596 **C,** 4-months, and **D,** 6-months old mice with sexes collapsed (n=15-16/group). *p<0.05,

597 ***p<0.001, ****p<0.0001 vs. NTg.

612 **Figure 4. Sex and genotype impact preferences displayed towards individual**

613 **odors.** Odor investigation duration for **A,** carvone, **B,** decanol, **C,** geraniol, **D,** guaiacol,

614 and **E,** thioglycolic acid (n=22-24/group). *$p<.05$, **$p<.01$, ***$p<.001$ vs. genotype-

615 matched males. #$p<.05$ vs. NTg females.

629 **Figure 5. Sex, age, and genotype impact preferences displayed towards individual**

630 **odors.** Odor investigation duration for females at 2-months (**Ai-Av**), 4-months (**Bi-Bv**),

631 and 6-months (**Ci-Cv**) and males at 2-months (**Di-Dv**), 4-months (**Ei-Ev**) and 6-months

632 (**Fi-Fv**) for carvone, decanol, geraniol, guaiacol, and thioglycolic (n=7-8/group). *p<.05,

633 **p<.01 vs. same-sex NTg counterpart.

Figure 6. No effect of 5XFAD genotype on measures of general task engagement.
Inter-poke interval in **A,** 2-months (n=7-8/group), **B,** 4-months (n=7-8/group), and **C,** 6-months old mice (n=8/group). Nose-poke count in **D,** 2-months (n=8/group), **E,** 4-months (n=8/group), and **F,** 6-months old mice (n=8/group). *p<.05 vs. males.