Competitive state of actions during planning predicts sequence execution accuracy.

Myrto Mantziara[1,2], Tsvetoslav Ivanov[1], George Houghton[1], and Katja Kornysheva[1,2]*

[1] School of Psychology, Bangor University, Bangor, Wales LL57 2AS, UK
[2] Bangor Imaging Unit, Bangor University, Bangor, Wales LL57 2AS, UK

*Correspondence: Dr Katja Kornysheva at e.kornysheva@bangor.ac.uk**,** School of Psychology**,** Bangor University**,** Wales, LL57 2AS**,** United Kingdom

Word count: Abstract: 240; Introduction: 786; Discussion: 2222; Methods: 3627; 39 pages, 5 figures, 0 tables, 3 supplementary figures.

Author contributions: M.M. and K.K. conceived the experiments; M.M. and T.I. collected the data; M.M. and K.K. designed the analysis; G.H. and K.K. designed the computational model; M.M., T.I. and K.K. performed the analysis; M.M., T.I., G.H. and K.K. wrote the paper. All authors contributed to editing of the manuscript.

Acknowledgements: The authors wish to thank Tom Hartley, Ken Valyear and Simon Watt for helpful comments on the study.

Disclosures: The authors declare no conflicts of interest.

Abstract

Humans can learn and retrieve novel skilled movement sequences from

memory, yet the content and structure of sequence planning are not well understood.

Previous computational and neurophysiological work suggests that actions in a

sequence are planned as parallel graded activations and selected for output through

competition (competitive queuing; CQ). However, the relevance of CQ during planning

to sequence fluency and accuracy, as opposed to sequence timing, is unclear. To

resolve this question, we assessed the competitive state of constituent actions

behaviourally during sequence preparation. In three separate multi-session

experiments, 55 healthy participants were trained to retrieve and produce 4-finger

sequences with particular timing from long-term memory. In addition to sequence

production, we evaluated reaction time (RT) and error rate increase to constituent

action probes at several points during the preparation period. Our results demonstrate

that longer preparation time produces a steeper CQ activation and selection gradient

between adjacent sequence elements, whilst no effect was found for sequence speed

or temporal structure. Further, participants with a steeper CQ gradient tended to

produce correct sequences faster and with a higher temporal accuracy. In a

computational model, we hypothesize that the CQ gradient during planning is driven

by the width of acquired positional tuning of each sequential item, independently of

timing. Our results suggest that competitive activation during sequence planning is

established gradually during sequence planning and predicts sequence fluency and

accuracy, rather than the speed or temporal structure of the motor sequence.

Keywords:

motor sequence; preparation; reaction time; finger accuracy; competitive

queuing

Introduction

Producing a variety of movement sequences from memory fluently is an essential capacity of primates, in particular humans. It enables a skilled and flexible interaction with the world for a range of everyday activities - from tool-use, speech and gestural communication, to sports and music. Key to fluent sequence production is sequence planning before the initiation of the first movement[1,2], with longer preparation time benefitting sequence execution, i.e. reducing initiation time after a 'Go' cue and improving accuracy[3]. However, the underlying nature and content of sequence planning is still debated[4].

Computational models of sequence control, such as competitive queuing (CQ) models, suggest that preparatory activity reactivates sequence segments *concurrently* by means of a parallel activation gradient in the parallel planning layer[5]. Here the neural activation pattern for each sequence segment is weighted according to its temporal position in the sequence[6,7]. A rich literature indirectly supporting CQ in sequence control stems from observations of serial recall including transposition of neighbouring items and items occupying the same position in different chunks[6,8,9], and excitability of forthcoming items during sequence production[10]. Moreover, the CQ account has also been substantiated directly at the neurophysiological level in the context of well-trained finger sequences[11,12], saccades[13] and drawing geometrical shapes[14]. Importantly, these results have demonstrated that the neural gradient during planning is relevant to subsequent execution. In particular, response separation in the competitive gradient during sequence planning is predictive of sequence production accuracy[11,14]. Together, these data suggest that skilled sequence production involves the concurrent planning of several movements in advance before sequence initiation to achieve fluent performance.

While neural CQ during planning has been shown to predict subsequent production, it remains unclear which properties of the sequence this preparatory pattern encapsulates – the accuracy of the sequence (fluency of initiation and production quality), or the temporal structure of the sequence (speed and temporal grouping). Some CQ models assume the presence of a temporal context layer and that the activity gradients are learned by associations of the latter to each sequence element in the parallel planning layer, e.g. through Hebbian learning[7]. The form of activity in the context layer can be as simple as a decaying start signal[15], a combination

of start and end signals[5,16] or a sequence of overlapping states[7,17]. Although primarily

encoding serial order of sequence items, models utilizing overlapping states can

implement effects of temporal grouping or sequence rhythm[7]. Therefore, it is possible

that the competitive activation of actions during sequence planning encodes the

temporal structure of the upcoming sequence.

In order to investigate the nature of sequence preparation and its relation to

subsequent performance, we have developed a behavioural paradigm to capture the

preparatory state of each item during planning of a well-learned sequence. Following

training, participants prepared a motor sequence from memory following an abstract

visual stimulus associated with a particular sequence of finger presses performed with

a particular temporal structure and speed. In half of the trials during the test phase,

the 'Go' cue was replaced by a finger press cue probing presses occurring at different

positions of the sequence. We used reaction time (RT) and finger press accuracy to

these 'probes' to compute as measures of the relative activation of planned actions

during sequence planning.

We hypothesized that if competitive queuing primarily reflected the accuracy of

the sequence plan, we would on average observe an enhancement of the CQ gradient

with longer preparation time, as well as a correlation of the gradient with measures of

sequence fluency and skill, specifically more rapid sequence initiation of correct

sequences after the 'Go' cue, more accurate timing and fewer finger press errors. If,

however, the gradient reflected the temporal structure – the speed and temporal

grouping of the sequence, we should see that it is less pronounced for sequences

twice as fast (speed manipulation), and shortened vs lengthened inter-press-intervals

(IPI; temporal structure manipulation), because the actions are closer together in time.

We find that the relative level of activation of probed actions at the end of the

planning period accords with their intended serial position. Contrary to the timing

hypothesis, we found no reliable association with speed or temporal structure of the

sequences. In contrast, we report that the corresponding CQ gradient is enhanced

with longer preparation time, and is correlated with faster initiation of correct

sequences and better temporal accuracy. Our data suggests that the competitive

queuing gradient during planning primarily encodes the intended order of actions and

the accuracy of a sequence plan, and not its overall speed or temporal structure.

Based on this data, we propose a computational model that explains how the width of

purely positional tuning could act on the relative activation state of actions during

sequence planning independently of timing to enable accurate and fluent sequence

performance.

Preparation
       500/1000/1500 ms

Figure 1 | Experimental conditions. a. Participants were trained to produce 4-element finger sequences following a *Go* cue from memory. Each finger sequence or timing corresponded to a unique abstract visual *Sequence* cue presented up to 1500 ms before the Go cue (Preparation period). Experiment 1 cued the production of sequences with two different finger press orders. Here we manipulated the duration of the Preparation period (500, 1000 or 1500 ms). Experiments 2 and 3 had a fixed preparation duration of 1500 ms, but *Sequence* cues prompted the production of sequences with a different temporal structure (slow, fast and irregular). Participants received visual feedback in each trial on the accuracy of the press order and their timing, and received points based on press accuracy, temporal accuracy and initiation time (cf. Methods section). **b**. In all experiments, we introduced *Probe* trials, in which following the preparation period the *Go* cue was replaced with a *Probe* cue prompting a particular finger digit to be pressed, corresponding to each sequence position or control (thumb, which did not feature in any sequence production). This condition was used to obtain the reaction time (RT) and error rate for each position at the end of the preparation period. They received points for accurate production and fast RTs.

Results

Finger press accuracy in sequences produced from memory was matched across conditions

Across three experiments, participants were trained for two days to associate two or three abstract visual cues with a particular four-element finger sequence performed with a particular temporal structure (Timing: slow, fast or irregular) following a brief preparation period (Delay between *Sequence* and *Go* cue onsets: short / 500ms, intermediate / 1000 ms and long / 1500 ms). In the test phase on the third day, they produced the respective sequences entirely from memory (Figure 1, all panels).

The finger error rate in sequence production from memory was higher in Experiment 1 than in Experiments 2 and 3. This is likely due to Experiment 1 involving the production of two different finger sequences produced with the same timing, and Experiments 2 and 3 involving the production of one finger sequence with different timings. The mean occurrence of finger errors, as indicated by either incorrect finger order or incomplete sequences, ranged from 0% to 26.6% in the short ($M = 5.6\%$, $SD = 6.9$), 0% to 21.9% in the intermediate ($M = 5.7\%$ $SD = 6$), and from 0% to 15.6% in the long Delay condition ($M = 4.6\%$, $SD = 4.5$) in Experiment 1. In Experiment 2, finger error rate varied between 0% and 5.5% at the slow timing ($M = 2.2\%$, $SD = 2.1$), between 0% and 8.5% at the fast timing ($M = 2.2\%$, $SD = 2.5$), and between 0% and 7% at the irregular timing ($M = 2.3\%$, $SD = 2.7$). Error performance in Experiment 3 showed a rate between 0% and 13.3% at the slow timing ($M = 2.6\%$, $SD = 3.4$), between 0% and 7.5% at the fast timing ($M = 2.8\%$, $SD = 2.5$), and between 0% and 9.2% at the irregular timing ($M = 2.4\%$, $SD = 2.6$).

Neither Delay (Experiment 1, $F(2, 36) = .993$, $p = .451$, $\eta p^2 = .052$) between the *Sequence* and the *Go* cue, nor the sequence Timing condition affected finger press accuracy during sequence production (Experiment 2, $F(2, 34) = .006$, $p = .994$, $\eta p^2 = .000$; Experiment 3, $F(1.458, 24.787) = .249$, $p = .711$, $\eta p^2 = .014$, Greenhouse-Geisser corrected, $\chi_2 (2) = 7.436$, $p = .024$, Figure 2c). This means that participants learned and prepared the finger order of all target sequences with the same finger accuracy, regardless of the preparation time or the temporal structure of the planned sequence.

Participants produced sequences from memory with correct relative

timing

When producing the sequences from memory during the test phase,

participants had a general tendency to produce faster versions of the finger press

sequences (Figure 2a), similar to the effects found in previous work[18]. The produced

sequence duration was shorter than the target duration by 28.4% (SD = 10.7%), 38.6%

(SD = 20.6%) and 46.2% (SD = 17.4%) in Experiment 1, 2 and 3, respectively (Figure

2a). However, the goal of our experimental design was to train participants to either

retain or to modulate the *relative* timing across conditions according to the target

relative IPIs, respectively (Figure 2b). Importantly, the majority of participants

produced the sequences with the correct relative timing across conditions – on

average the same temporal structure (slow) across preparation durations in

Experiment1, and three different temporal structures (slow, fast, irregular) in

Experiments 2 and 3 (Figure 2b).

Figure 2 | Sequence production. a. Individual participants' raster plots show the timing of single button presses for each correct sequence trial produced from memory after the *Go* cue (t = 0) following training (target timing superimposed, grey lines). The colour code of the button presses corresponds to the

press position in Figures 1 and 5. Within each condition, trials are ordered from most accurate to least accurate with regard to target onset (colour coding for conditions, cf. side bars in first participant panels, respectively). **b.** On average IPI production followed the target IPI structure, with a slow, twice as fast and an irregular sequence. IPIs were normalized across trials relative to slow isochronous condition. Preparation duration did not modulate IPI production of slow sequences, in contrast to the timing conditions. Error bars represent standard errors. **c.** Sequence initiation RT (Go cue to first press latency) decreased with preparation time (foreperiod effect). Relative temporal error was elevated for the irregular sequence in both experiments 2 and 3, in which it occurred. There was no effect of any of the conditions on finger press error rate, defined as proportion of incorrect trials. | * P ≤ 0.05 | ** P ≤ 0.01 | *** P ≤ 0.001.

Despite a largely overlapping sequence timing across preparation durations in

Experiment 1 (Figure 2b, left), we found a small, but significant IPI × Delay interaction

(3×3 repeated measures ANOVA: $F_{(4, 72)} = 2.528$, $p = .048$, $\eta p^2 = .123$) explained by

a modulation of 9 ms. Post-hoc comparisons (Bonferroni-corrected for nine tests)

revealed a significant shortening of the 1_{st} interval in the short ($p = .002$) and in the

intermediate delay ($p = .002$) compared to the long delay conditions. Additionally, the

3_{rd} interval was larger in the long delay than at the intermediate delay ($p = .004$). This

shows that there was a tendency to slightly compress the 1_{st} interval with shorter

preparation time and slightly expand the 3_{rd} interval with longer preparation time.

However, the size of this temporal modulation in Experiment 1 was 97% smaller than

the temporal structure modulation induced in Experiments 2 and 3. Any potential

sequence timing effect on CQ activation of actions during preparation should thus be

vastly augmented in Experiments 2 and 3.

As expected, Experiment 2 showed a significant IPI × Timing interaction (3×3

repeated measures ANOVA: $F_{(1.260, 21.417)} = 59.485$, $p < .001$, $\eta p^2 = .778$,

Greenhouse-Geisser corrected, $\chi_2 (9) = 97.832$, $p < .001$). The pairwise comparisons

(Bonferroni-corrected for nine tests) of the produced intervals confirmed that the

participants modulated their relative interval production according to the trained target

interval structure. In accordance with the target sequence, in the slow timing condition

the 1_{st} interval was significantly longer than in the fast ($p < .001$) and the irregular

timing conditions ($p < .001$), while there was no difference between the fast and

irregular conditions ($p = 1.000$) for the latter. The 2_{nd} interval length increased slightly,

yet proportionally for both the slow and the fast conditions, retaining the significant

difference ($p < .001$) and doubled in length for irregular relative to the slow timing

condition ($p < .001$). Finally, the 3_{rd} interval exhibited a very similar profile to the 1_{st}

interval (slow *vs* fast, $p < .001$; slow *vs* irregular, $p < .001$), but showed a slightly lower

percent interval for the fast compared to the irregular conditions ($p = .027$). Overall,

the IPI production data shows that the fast sequence was on average half as long as

the slow sequence, and the irregular timing condition changed the relative interval

structure from regular slow to an irregular short–long–short pattern of the same

sequence duration. Experiment 3 replicated the findings of Experiment 2 showing a

significant IPI × Timing interaction (3×3 repeated measures ANOVA: $F_{(1.558, 26.485)}$

= 17.369, $p < .001$, $\eta p^2 = .505$, Greenhouse-Geisser corrected, $\chi^2_{(9)} = 61.311$, p

$< .001$). Again, post-hoc pairwise comparisons (Bonferroni-corrected for nine tests)

confirmed that the 1_{st} interval of the slow condition was longer than that of the fast

($p = .001$) and irregular ($p = .003$) conditions, while no difference was found between

fast and irregular conditions ($p = 1.000$). The 2_{nd} interval was significantly longer in the

slow compared to the fast condition ($p = .001$), but shorter compared to the irregular

condition ($p = .005$). Similarly, the fast condition was half as long in the irregular

condition ($p < .001$). The 3_{rd} interval was twice as long for the slow relative to the fast

condition ($p < .001$), but failed to show a significant shortening for the irregular relative

to the regular slow sequence conditions ($p = 1.000$), and there was only a marginally

significant difference between fast and irregular conditions ($p = .096$). Overall, our

findings demonstrate that, on average, participants retrieved and produced the finger

sequences form memory with distinct temporal structures according to the relative

timing of slow, fast and irregular target intervals.

Longer preparation duration speeds up sequence initiation

The time to initiate a correct action sequence after the *Go* cue can be taken as

a marker of the state of action planning after the preparatory delay[3,19,20]. We found a

significant difference in mean initiation RT with Delay (Experiment 1, one-way

repeated measures ANOVA: $F_{(1.382, 24.877)} = 52.809$, $p < .001$, $\eta p^2 = .746$,

Greenhouse-Geisser corrected, $\chi^2_{(2)} = 10.074$, $p = .006$) (Figure 2c, left). Pairwise

comparisons (Bonferroni-corrected for three tests) confirmed that initiation time for the

intermediate and long delay conditions was significantly shorter than following the

short delay (intermediate *vs* short delay, $p < .001$; long *vs* short delay, $p < .001$).

Similarly, sequence initiation following a long delay performed at significantly faster

mean RT as compared to the intermediate delay period ($p = .005$). Notably, this effect

is also in line with the classic foreperiod effect identified for single actions showing

faster RTs for longer foreperiod durations [21,22] suggesting that temporal expectation of

the *Go* cue may also contribute to faster sequence initiation in addition to the state of

sequence planning.

In contrast to the effect of preparation time, the planned temporal structure of

the sequence did not consistently affect sequence initiation RT (Figure 2c, left). There

was no main effect of sequence Timing in Experiment 2 (one-way repeated measures

ANOVA: $F(1.407, 23.917) = 1.700$, $p = .207$, $\eta p^2 = .091$, Greenhouse-Geisser

corrected, $\chi_2(2) = 8.759$, $p = .013$), but a main effect of Timing in Experiment 3 (one-

way repeated measures ANOVA: $F(1.294, 21.993) = 11.590$, $p = .001$, $\eta p^2 = .405$,

Greenhouse-Geisser corrected, $\chi_2(2) = 12.632$, $p = .002$). Specifically, as explained

by pairwise comparisons (Bonferroni-corrected for three tests), participants in

Experiment 3 were slower at initiating a slow regular sequence when compared to a

fast regular sequence ($p = .006$) and an irregular sequence ($p = .010$) whilst there was

no difference in initiation RT between the fast regular and the irregular conditions (p

$= .118$).

Sequences involving irregular inter-press-intervals were produced with

less accurate timing

Next, we aimed to establish whether preparation time and the temporal interval

structure of sequences modulated the observed relative timing accuracy across

presses (Figure 2c, middle). In Experiment 1, the mean relative temporal error did not

differ across the three delay conditions (one-way repeated measures ANOVA: $F(2, 36)$

$= .105$, $p = .901$, $\eta p^2 = .006$). This indicates that time to prepare the sequence did not

affect the degree of relative temporal accuracy. Here, sequence accuracy in conditions

with a shorter preparation time might have been compensated by slower initiation RT

(cf. above). In Experiment 2, there was a significant effect of Timing (one-way repeated

measures ANOVA: $F(2, 34) = 28.226$, $p < .001$, $\eta p^2 = .624$). Pairwise comparisons

(Bonferroni-corrected for three tests) revealed that participants performed at a higher

relative temporal accuracy when producing a slow regular sequence compared to an

irregular ($p < .001$) and a fast regular compared to an irregular sequence ($p < .001$),

while there was no difference between the slow and fast regular conditions ($p = 1.000$).

Experiment 3 replicated the main effect of Timing (one-way repeated measures

ANOVA: $F(1.454, 24.723) = 7.060$, $p = .007$, $\eta p^2 = .293$, Greenhouse-Geisser

corrected, $\chi_2(2) = 7.527$, $p = .023$). In line with the findings of Experiment 2, there

were smaller temporal errors in the slow sequence condition compared to the irregular

condition (p = .049) and in the fast compared to the irregular sequence (p =. 008).
There was no significant difference in temporal performance between the two regular
conditions (p = 1.000). These results suggest that the production of sequences which
consist of several different IPIs (irregular sequence) as opposed to only one interval
length (isochronous/regular sequence) is associated with decreases in temporal
accuracy of the sequence.

Action probes show graded activation of sequence elements at the end of
preparation and are modulated by preparation duration, not sequence timing

In half of the trials in the test phase on Day 3, instead of the *Go* cue that
prompted the production of the prepared finger sequence, participants encountered a
visual *Probe* cue which instructed them to respond with the corresponding finger press
as quickly and accurately as possible (Figure 1b). This allowed us to obtain two
behavioural measures related to the competitive state of each constituent press at the
end of the preparation period – the relative action activation and the probability of
correct action selection – for each action in the sequence, respectively. Specifically,
lower RT would suggest higher activation of the correctly selected action and lower
press error rate a higher probability of selection, and thus availability of the associated
action.

Reaction times. To evaluate the relative activation state of the probed actions
associated with different sequence positions (Figure 3a), we normalized the RT of
each probe position relative to the RT of the first position of the prepared sequence
(RT increase in % relative to 1_{st} position; cf. Supplementary Figure 1a for raw RT
values). In Experiment 1, results showed a main effect of Position (F (1,615, 29.062)
= 45.958, $p <$ .001, $\eta p^2 =$ .719, Greenhouse-Geisser corrected, $\chi 2$ (5) = 22.621, p
$<$ .001). Planned contrasts to detect differences between adjacent positions revealed
a significant RT increase with each position compared to its preceding one (2_{nd} *vs* 1_{st}
position, F (1, 18) = 63.360, $p <$ .001, $\eta p^2 =$ .779; 3_{rd} *vs* 2_{nd} position, F (1, 18) = 54.534,
$p <$ .001, $\eta p^2 =$ .752; 4_{th} *vs* 3_{rd} position, F (1, 18) = 24.900, $p <$ .001, $\eta p^2 =$ .580). In line
with our hypothesis, these differences indicate a graded activation of actions according
to their serial position in the sequence, with the first action being the most activated.
We also found a significant Position × Delay interaction (F (6, 108) = 2.980, $p =$ .010,
$\eta p^2 =$ .142). Planned contrasts using the long preparation duration as the reference
condition for the Delay factor, showed a significantly greater increase of the 2_{nd}

position relative to the 1st position at the long *vs* the short delay (F (1, 18) = 7.349, p

= .014, ηp^2 = .290). These results suggest that a longer preparation time prior to

sequence execution boosted the competitive activation of actions.

Figure 3 | Competitive state of actions during sequence preparation. *Probe* trials prompting the production of an action associated with the 1st-4th press position of the prepared sequence or a control action not present in any sequence (Experiment 3). **a.** Reaction time (RT) gradually increased for later sequence positions relative to the first position and became more pronounced with longer preparation duration, with responses to actions in later sequence positions becoming slower on average, when participants had more time to prepare the sequence (Experiment 1). No significant changes to RT increase were observed between conditions in which participants prepared sequences performed with different timing (Experiment 2 and 3). The RT increase was most pronounced for the action not featuring in the planned sequence (control action was a thumb press; Experiment 3), cf. raw RT graphs in Supplementary Figure 1. **b.** Press error rate also increased gradually for later sequence positions relative to the first position, with the exception of the last position, thus approaching an inverted U-shape. The press error gradient became more pronounced with longer preparation duration, i.e. responses to probes associated with later positions became less accurate when participants had more time to prepare the cued sequence. **c.** This pronounced effect on error increase was driven by trials where the response to action probes was short (lower RT quartile). When participants slowed down

their response allowing more time for deliberation and correction (upper RT quartile), the characteristic error increase was absent or less pronounced. Error bars represent standard errors.

In Experiment 2, we replicated the main effect of Position ($F_{(2.230, 37.904)}$ =

25.131, $p < .001$, $\eta p^2 = .596$, Greenhouse-Geisser corrected, $\chi 2_{(5)} = 15.333$, p

$= .009$). Planned contrasts showed significant differences when contrasting all pairs of

adjacent probe positions replicating the findings of Experiment 1. Specifically, RT

increase of the 2nd position was larger compared to the 1st ($F_{(1, 17)} = 48.072$, $p < .001$,

$\eta p^2 = .739$). Similarly, there was a significantly greater RT increase for the 3rd position

vs the 2nd ($F_{(1, 17)} = 32.040$, $p = .001$, $\eta p^2 = .653$) and the 4th vs the 3rd ($F_{(1, 17)} =$

28.873, $p < .001$, $\eta p^2 = .629$). Crucially, there was no interaction between Position and

Timing ($F_{(2.430, 41.318)} = 2.823$, $p = .061$, $\eta p^2 = .142$, Greenhouse-Geisser

corrected, $\chi 2_{(20)} = 59.308$, $p < .001$). This suggests that preparing a sequence with a

different temporal structure did not impact the competitively cued activations at the

end of preparation.

Experiment 3 once more replicated a main effect of Position ($F_{(3, 51)} =$

29.852, $p < .001$, $\eta p^2 = .637$). Planned contrasts, similarly, revealed a significantly

greater RT increase for the 2nd, 3rd and 4th positions over their preceding 1st, 2nd and

3rd positions, respectively (2nd vs 1st, $F_{(1, 17)} = 61.485$, $p < .001$, $\eta p^2 = .783$; 3rd vs

2nd, $F_{(1, 17)} = 69.762$, $p < .001$, $\eta p^2 = .804$; $F_{(1, 17)} = 14.180$, $p = .002$, $\eta p^2 = .455$).

In addition, the finger press which did not feature in any of the planned sequences

(control action: thumb) showed a further RT increase relative to the last (4th position)

item of the sequence (paired samples t-test: $t_{(17)} = 3.062$, $p = .007$, $d = .840$, two-

tailed). Timing did not interact with Position ($F_{(3.743, 63.632)} = 1.089$, $p = .367$,

$\eta p^2 = .060$, Greenhouse-Geisser corrected, $\chi 2_{(20)} = 36.727$, $p = .014$). This result

replicates Experiment 2. Thus, CQ during sequence preparation was not dependent

upon the speed or temporal structure of the planned sequences, and suggests that

fine grained competitive activation gradient of constituent actions of a sequence is

activated above the level of an unrelated effector.

Error rate. To evaluate the relative probability of selection of the probed actions

associated with different sequence positions (Figure 3b), in each experiment the finger

error rate for each probe action was calculated and normalized to that of the first

position (Error rate increase in % relative to 1st position; cf. Supplementary Figure 1b

for raw error rate values). Equally, we assessed the same factors, predicting an

ascending error rate increase by position. Experiment 1 showed a main effect of

Position (4 × 3 repeated measures ANOVA: $F(1,948, 35.073) =$

18.017, $p < .001$, $\eta p^2 = .500$, Greenhouse-Geisser corrected, $\chi_2(5) = 13.595$, $p = .019$).

Planned contrasts revealed a significantly increased error rate for the 2_{nd} position

compared to the 1_{st} position ($F(1, 18) = 29.675$, $p < .001$, $\eta p^2 = .622$) and for the 3_{rd}

position compared to the 2_{nd} position ($F(1, 18) = 25.937$, $p < .001$, $\eta p^2 = .590$), whilst

the last, 4_{th} position showed a lower error increase than the 3_{rd} position ($F(1, 18) =$

5.092, $p = .037$, $\eta p^2 = .220$). This error rate pattern during preparation shows an

inverted U-shape, similar to serial position curves during production[23] and suggests

the presence of a ranked probability across sequence positions for action selection.

The Position × Delay interaction ($F(4.137, 74.466) = 3.813$, $p = .007$, $\eta p^2 = .175$,

Greenhouse-Geisser corrected, $\chi_2(20) = 34.036$, $p = .028$) was driven by a significant

increase of the 2_{nd} position compared to 1_{st} position at the long vs the short delay (F

$(1, 18) = 10.877$, $p = .004$, $\eta p^2 = .377$) as revealed by planned contrasts. No other

pairs showed a significant difference. In accordance with the RT results, these findings

suggest that accuracy of probe elements during sequence planning is modulated by

preparation, in that a longer preparation time is associated with more pronounced error

increases for the 2_{nd} position when compared to a short preparation period, suggesting

less availability for selection of actions in later positions the more sequence planning

advances.

Experiment 2 showed a significant main effect of Position ($F(3, 51) =$

14.397, $p < .001$, $\eta p^2 = .459$). Planned contrasts revealed that this effect was driven

by a significant error rate increase for the 2_{nd} position vs the 1_{st} position ($F(1, 17) =$

24.070, $p < .001$, $\eta p^2 = .586$). The 3_{rd} position performed at a greater error increase

than the 2_{nd} position ($F(1, 17) = 15.510$, $p = .001$, $\eta p^2 = .477$), whilst the 4_{th} position

was not significantly different from the 3_{rd} position ($F(1, 17) = 1.284$, $p = .273$,

$\eta p^2 = .070$). These results replicate the CQ effect of serial actions during preparation,

found in Experiment 1, with a graded increase in error rates for later elements up to

the 3_{rd} position. We did not find a significant Position × Timing interaction ($F(6, 102)$

$= 1.583$, $p = .160$, $\eta p^2 = .085$).

Similarly in Experiment 3, there was a main effect of Position ($F(3, 51) =$

13.725, $p < .001$, $\eta p^2 = .447$), with the 2_{nd} position showing a greater error increase

than the 1_{st} position ($F(1, 17) = 29.074$, $p < .001$, $\eta p^2 = .631$), and the 3_{rd} position

performing with more errors than the 2_{nd} position (F (1, 17) = 17.903, p = .001,

ηp^2 = .513). Error rates of the 4_{th} position did not differ from the 3_{rd} position (F (1, 17)

= 3.791, p = .068, ηp^2 = .182). Our prediction that the control action would not be part

of this queuing pattern, implying a much weaker probability to be selected for

execution, was refuted by a non-significant difference from the 4_{th} position (paired

samples t-test: t (17) = -.323, p = .751, d = .111, two-tailed). As in Experiment 2,

Position did not interact with Timing (F (3.803, 64.654) = 1.869, p = .130, ηp^2 = .099,

Greenhouse-Geisser corrected, χ_2 (20) = 42.899, p = .002). Together, this indicates

that the competitive error rate for probed actions during preparation was not modulated

by the speed or temporal structure of the planned sequence.

Whilst there was no significant interaction of Timing and Position for action

probes during preparation (neither for RT, nor for error rate), we observed a non-

significant, but consistent flattening of the CQ curve for the temporally irregular

sequence across Experiments 2 and 3. However, this patterns of results cannot be

attributed to changes in temporal grouping of actions per se, but may be driven by

accuracy: The irregularly (non-isochronously) timed sequence was characterized by a

highly significant increase in relative temporal error when compared to both the slow

and fast regularly (isochronously) timed sequences (Figure 2b, Experiments 2 and 3)

due to the increased temporal complexity. This lends support to the alternative

hypothesis, namely that the precision of the sequence plan is driving the CQ state of

actions during the preparation period.

A steeper CQ error gradient is bound to fast responses

Next, we sought to determine whether the characteristic error rate gradients were the

result of automatic responses, or deliberated action selection after the *Probe* cue. To

test this hypothesis, we assessed the error rate gradient for fast *vs* slow responses

following the *Probe* cue. We extracted the relative error rate increases for action

probes in the first and third RT quartiles for each experiment (Figure 3c). Only for the

fast responses, we found a main effect of Position (Experiment 1, F (1.758, 31.650) =

19.731, p < .001, ηp^2 = .523, Greenhouse-Geisser corrected, χ_2 (5) = 17.279, p = .004;

Experiment 2, F (2.033, 34.559) = 16.325, p < .001, ηp^2 = .490, Greenhouse-Geisser

corrected, χ_2 (5) = 19.928, p = .001; Experiment 3, F (3, 51) =

12.749, p < .001, ηp^2 = .429). Planned contrasts for adjacent positions confirmed a

graded increase in finger errors up to the 3_{rd} position (Experiment 1, 2_{nd} position *vs* 1_{st}

position, F (1, 18) = 26.954, $p < .001$, $\eta p^2 = .600$; 3rd position vs 2nd position, F (1, 18) = 35.745, $p < .001$, $\eta p^2 = .665$; 4th position vs 3rd position, F (1, 18) = 2.347, $p = .143$, $\eta p^2 = .115$; Experiment 2, 2nd position vs 1st position, F (1, 17) = 27.138, $p < .001$, $\eta p^2 = .615$; 3rd position vs 2nd position, F (1, 17) = 15.222, $p = .001$, $\eta p^2 = .472$; 4th position vs 3rd position, F (1, 17) = 4.982, $p = .039$, $\eta p^2 = .227$; Experiment 3, 2nd position vs 1st position, F (1, 17) = 21.580, $p < .001$, $\eta p^2 = .559$; 3rd position vs 2nd position, F (1, 17) = 13.888, $p = .002$, $\eta p^2 = .450$; 4th position vs 3rd position, F (1, 17) = 1.845, $p = .192$, $\eta p^2 = .098$). We also found a significant Position × Delay interaction (Experiment 1, F (6, 108) = 4.003, $p = .001$, $\eta p^2 = .182$), driven by a significant increase of the 2nd position 1st position at the long vs the short delay (F (1, 18) = 18.132, $p < .001$, $\eta p^2 = .502$) and at the long vs the intermediate delay (F (1, 18) = 10.370, $p = .005$, $\eta p^2 = .366$). Error rate increases did not change by sequence position number in the slow responses (Experiment 1, F (3, 54) = .313, $p = .816$, $\eta p^2 = .017$; Experiment 2, F (3, 51) = .552, $p = .649$, $\eta p^2 = .031$; Experiment 3, F (3, 51) = 1.672, $p = .185$, $\eta p^2 = .090$), accompanied by an absent Position × Delay interaction (Experiment 1, F (3.559, 64.058) = 1.302, $p = .280$, $\eta p^2 = .067$, Greenhouse-Geisser corrected, χ_2 (20) = 37.340, $p = .012$). The control action was not different from the 4th position in either RT pole (fast RTs, t (17) = 1.654, $p = .117$, $d = .473$, two-tailed; slow RTs, t (17) = .203, $p = .842$, $d = .050$, two-tailed). These results suggest that the CQ gradient at the end of a preparation period of 500 to 1500 ms was driven by automatic responses rather than by cognitive action selection and replanning, and constitute a readout for the state of actions during sequence planning.

Preparatory CQ gradient correlates with temporal accuracy and initiation speed

Neurally derived CQ of sequence actions during planning predicts the participants' subsequent performance accuracy as shown previously[11]. In line with this finding, we found that more time to prepare a sequence is associated with a more pronounced competitive RT and error rate increase for action probes. To test the association directly, we predicted that a more pronounced (steeper) CQ gradient of RTs and error rates would correlate with a better performance in sequence production, specifically with faster correct sequence initiation, and less temporal and finger press errors. Correlation analyses were performed on group data obtained from trials in the slow timing condition and long preparation duration (1500 ms) present across all three

experiments (N = 55). The magnitude of the CQ gradient during preparation was

calculated based on RT and error rate increase data in *Probe* trials (difference

between adjacent positions; Figure 4a and 4b). Results showed that participants with

a more pronounced CQ based on relative RT and error rate increase initiated correct

sequences faster (CQ RT increase: r = -.393, p = .002, one-tailed; CQ error increase:

r = -.539, p < .001, one-tailed). Higher RT based CQ also predicted smaller relative

temporal error (r = -.345, p = .005, one-tailed). However, in contrast to the correlations

with the neural measure of CQ reported in a previous study[11], neither CQ RT increase,

nor CQ error increase showed negative correlations with finger error (CQ RT increase:

r = .083, p = .273, one-tailed; CQ error increase: r = .118, p = .196, one-tailed). Also,

contrary to the CQ RT increase, a more pronounced CQ error increase of probe

actions at the end of preparation was not associated with reduced temporal error

during execution (r = -.051, p = .356, one-tailed). Thus, CQ error increase may be a

less sensitive predictor for temporal accuracy than CQ RT increase.

Figure 4 | Correlation between overall CQ measures during preparation and subsequent production. Average relative RT (**a**) and press error increase (**b**) between adjacent positions (1st to 2nd, 2nd to 3rd, 3rd to 4th) obtained through probe trials was taken as a proxy for CQ of actions during preparation. Larger CQ during preparation was associated with faster initiation speed of correct sequences, and smaller relative temporal errors. Larger CQ was not associated with reduced finger error rate (proportion of trials with wrong finger order, finger order repetitions, or missing presses), as predicted based on neural CQ findings (Kornysheva et al. 2019). All correlations are one-tailed, in line with one-sided predictions.

In sum, consistent with previous neurophysiological findings[11], our behavioural

results show that during sequence preparation of sequence from memory participants

establish a competitive activation and selection gradient of constituent actions

according to their serial order. This competitive gradient expands with longer

preparation durations and is more pronounced in participants with faster sequence

initiation and more precise interval timing.

Discussion

Sequence planning is central to skilled action control, however its content and

organisation is poorly understood[2,4]. Neurophysiological findings in humans have

demonstrated that a trained action sequence is pre-planned by establishing a

competitive activation gradient of action patterns according to their serial position, and

that the quality of this neural pattern during planning predicts subsequent

performance[11–14]. Here we have established a behavioural measure of the preparatory

action activation gradient and demonstrate that it reflects the skill (sequence

production accuracy and fluency of initiation), rather than the temporal structure

(sequence production speed and temporal interval pattern) of the planned sequence.

Both the time to respond and the probability of making a finger press mistake

increased progressively when participants responded to action cues during

preparation that were associated with later *vs* earlier positions in the respective

sequence. The non-linear increase was particularly pronounced for the first three out

of four planned actions in the sequence. This response gradient demonstrates that the

relative availability of each planned action in the sequence decreases with serial

position, as predicted by competitive queuing (CQ) models[6,7,24,25] and previous

neurophysiological findings[11,14].

The preparatory action activation gradient markedly contrasts with mechanisms

for non-sequential action planning involving multiple actions: A cued set of possible

actions triggers equal activity increase in cortical populations tuned to the respective

actions, and the preparatory competition is only resolved once an action cue specifies

the target action[26]. In contrast, sequence preparation establishes a fine-tuned gradient

of action activations, with the latter switching flexibly depending on the retrieved

sequence. Notably, actions that were part of the planned sequence were activated

above the level of a control action which was not part of the retrieved sequence (Figure

3a, right). This suggests that all constituent actions were concurrently activated above

a passive baseline, albeit to a different degree depending on their position in the

planned sequence.

Our study provides a behavioural measure of the competitive state of

constituent actions during sequence planning. This is complementary to previous

behavioural work which revealed CQ of actions during production, such as accuracy

and RT curves obtained from sequence execution[27,28], or on-the-fly action planning

following sequence initiation, assessed behaviourally[29] and through measures of

cortico-spinal excitability[10]. Gilbert and colleagues have employed a paradigm at the

interface between sequence preparation and production to characterize the CQ

profiles the respective sequential actions – silent rehearsal[30]. Here participants were

asked to listen to sequences of spoken digits and silently rehearse the items during a

retention interval. They received explicit instructions to rehearse the sequence at the

same pace as active production. After an unpredictable delay, a tone prompted the

report of an item being rehearsed at that moment and revealed graded overlapping

probabilities of neighbouring items, suggesting potential CQ during internal rehearsal.

In contrast to the latter study, our paradigm did not allow for active rehearsal during

preparation: First, our participants retrieved the sequence entirely from memory

without a sensory instruction period which might have facilitated active entrainment

with the sequence prior to planning. Second, the period for sequence retrieval and

planning was comparatively brief (ranging from 500 to 1500 ms after *Sequence* cue

onset) and not sufficient to cycle through the full sequence at the rate participants

employed for active production. In addition, if the observed CQ gradient were

somehow driven by silent rehearsal at the target rate, it would have been more

pronounced for the fast sequences, as more of the planned sequence could fit into the

preparation phase. However, there was no significant difference between relative

activation curves for fast and slow sequences.

Whilst active motor rehearsal at scale during the short preparation phase is

unlikely, an alternative serial mechanism underlying the different levels of action

activation may be mediated by rapid sequence replay. The latter has been observed

in the hippocampus during navigation tasks[31] and perceptual sequence encoding[32],

as well as in the motor cortex in the context of motor sequence learning tasks[33]. Replay

has been shown to involve fast sweeps through the neural patterns associated with

the sequence during wakeful rest and planning (preplay)[31,34–36], and is characterized

by a multifold temporal sequence compression[32,33,37,38]. How replay could translate

into a parallel activation of serial items described here is uncertain. One possibility is

that serial sweeps during motor sequence preparation involve fast repeated replay

fragments[37,39] of different length during planning, starting with the first elements – e.g.

1st-2nd-3rd, 1st-2nd, 1st, 1st-2nd-3rd-4th, 1st-2nd etc. This would produce an overall bias

towards the activation of earlier rather than later parts of the planned sequence, which

may be translated into a cumulative ramping activity for each constituent action by a

separate neuronal mechanism during the preparation period[26,40]. Future analysis of

the 'sequenceness'[32,33] of the corresponding neural patterns during preparation should

shed light on the presence of preplay and its hypothesized relationship to parallel CQ

of actions[11].

The CQ gradient was established after a brief retrieval and preparation period,

and revealed through faster rather than slower responses to probes (Figure 3c). This

suggests that out behavioural measure of CQ during sequence planning reflects a

rapid and automatic process involved in the execution of well-trained motor sequences

from memory, and is not a result of slow deliberation or higher-level decision making.

Contrary to a prominent account of motor control of skill learning[41,42], this data implies

that discrete motor sequence production incorporates automatic planning

mechanisms which are associated with fluent and accurate execution of sequential

actions.

Remarkably, longer preparation times reinforced the competitive activation

gradient making responses to action probes for later sequence positions even slower

and more inaccurate relative to those for earlier positions. Whilst counterintuitive in the

context of single action performance gains from longer foreperiod durations[21], the

gradient expansion with time suggests a dynamic refinement of the plan for sequence

production during the retrieval and preparation phase. This refinement involves the

graded suppression of later actions in the sequence, making them less available for

production, even more so with time. Crucially, the gradient increase with preparation

duration was not accompanied by substantial expansion or compression of sequence

production. Instead, the CQ gradient was associated with a faster initiation of correctly

performed sequences whilst retaining the same level of press and timing accuracy,

suggesting increased sequence fluency. This demonstrates that the action activation

gradient established during planning reflects the preparedness for correct and fluent

production, rather than the planned temporal structure of the sequence.

551   Furthermore, participants who had a more pronounced competitive activation

during planning exhibited both faster initiation times and a more accurate temporal

execution of the sequence after the "Go" cue, particularly when looking at the RT

based CQ gradient. These findings strengthen the interpretation that an ordered

competitive activation of actions during planning preempts subsequent fluency and

temporal accuracy of the sequence[11]. Yet, we did not replicate the association of the

planning gradient with finger error probability found in the latter study. This may be due

to a smaller pool of timing and finger order sequences that the participants had to learn

relative to the previous paradigm, and the presence of only one finger order (but

different sequence timing) in Experiments 2 and 3. This likely facilitated finger

accuracy to reach ceiling levels in a substantial number of participants. Future

experiments should resolve an association with finger accuracy through the inclusion

of a larger pool of trained sequences to provoke more frequent finger errors.

Alternatively, reaching or drawing tasks would allow to make the spatial in addition to

temporal feature of the sequential behaviour continuous and capture fine-grained

spatial errors at overall high accuracy levels of sequence production.

567   In contrast, doubling the speed of sequence production did not change the

relative activation between sequential actions at the end of the preparation period.

This suggests invariance of the gradient in the competitive planning layer across

sequences produced at different time scales. This transfer across speed profiles is in

line with the presence of flexible motor timing and temporal scaling in dynamic

neuronal populations[43,44], and a separate neural process controlling the speed of an

action or action sequence during execution, e.g. through the strength of an external

input to the network involved in the generation of timed behavior.[43] Preparing a

sequence of the same length with an irregular compared to isochronous interval

structure was associated with a tendency for a dampened CQ gradient during

sequence planning. However, this non-significant trend on CQ is unlikely to be the

effect of temporal grouping, as the irregular interval sequence was characterized by a

significant increase in temporal interval production error (Figure 2c, middle panel).

Instead, we hypothesize that longer preparation time (above 1500 ms) would have

benefitted the participants and enhanced the relative activation gradient in line with

Experiment 1 in order to form a more accurate plan for the complex sequencing of two
different (non-isochronous) rather than just one constituent IPI (isochronous).
Our results show that CQ of actions during sequence planning reflects the
overall action order and temporal accuracy of the sequence, but not its temporal
structure – neither its speed, nor its temporal grouping. This dissociation is
counterintuitive, however, we propose that temporal accuracy can be dissociated from
timing in CQ models. In our model (Figure 5 and Methods), we assume that positional
associations of the items in the sequence (positional context and parallel planning
layer) are determined by the respective sequence cue, and the corresponding start-
state of the cued sequence becomes gradually activated. Crucially, we show that
changing the width of the receptive field for each position (Figure 5a) affects the
activation gradient of action items during sequence planning (Figure 5b). Specifically,
our model demonstrates that narrowing this positional tuning will cause a steeper
relative activation gradient at the end of sequence preparation, with actions in later
positions being progressively less activated. Conversely, wider tuning, would broaden
the excitation from the positional context to parallel planning layer and lead to smaller
relative activation differences between actions at the end of the planning period.
Notably, while the positional tuning in CQ models is hypothesized to be acquired
through exposure (Hebbian learning)[45], we assume that it is dynamically established
throughout the preparation period (cf. Methods section). Thus, the width of the
positional tuning of individual actions in the parallel planning layer may underly the
accuracy of actions, independently of the overall speed and temporal structure of
sequences.

Figure 5 | Competitive queuing (CQ) model and the role of positional tuning in sequence preparation. a. The Parallel planning and Competitive choice layers of the CQ model contain nodes representing possible sequence items, such as finger presses A, B, C and D. When learning a sequence, connections are formed from sequentially activated nodes in the Positional context layer to item nodes in the Parallel planning layer as each is activated in turn. Crucially, the current model incorporates a positional tuning of the nodes. The receptive field of this positional tuning has a tuning (variance) parameter σ, controlling the model's sensitivity to positional differences. This tuning curve may be acquired though training (variability of instructive stimulus and training exposure), as well as reflect an intrinsic variability of each participant (sensory or motor variability). **b.** The tuning width of the receptive field determines, inversely, the spread of the competitive activations of corresponding action items following the *Sequence* cue, with a narrow or sharper tuning producing more pronounced CQ during preparation. Time steps represent linear arbitrary values.

Although our empirical data and CQ model do not support the integration of the

timing signal with action order before sequence execution, they do not exclude the

presence of a separate preparation process for the speed and timing of the respective

sequence, which may take place concurrently or at different time points during

preparation[46–48]. In previous work, we proposed a drift-diffusion based model which

contains input from separate modules that activate action order and timing.[49] This

model was based on behavioural sequence learning data demonstrating that

sequence timing is encoded independently of the action order, but requires

multiplicative, rather than additive integration with each action. This enables previously

learnt sequence timing to be transferred to new sequences, but only after the action

order has been acquired, reconciling previous experimental findings[50–54]. Most

recently Zeid and Bullock proposed how such plans might be generated in the context

of CQ models[51], proposing that two CQ modules could operate in parallel - one

controlling the item order and the other controlling the sequence of inter-onset-
intervals that define a rhythmic pattern, including separate parallel planning and
competitive choice layers. While this model is in line with neurophysiological and
imaging evidence for a separate control of timing for sequence generation[50,55–59],
empirical data for a dedicated CQ process for temporal intervals is lacking.

Conclusions

In sum, our findings indicate that the graded relative activation state during a
brief period of retrieval and planning reflects the subsequent action order and
correlates with the individual's sequence fluency and accuracy. It appears to be
invariant to the exact timing of the sequence, but is instead bound to the precision of
the positional tuning. In contrast to neurophysiological approaches involving advanced
neural pattern analysis[11,14], a simple behavioural paradigm could provide a
straightforward and cost-effective proxy to assess the state of action preparation
across trials in individual participants. This behavioural readout may help advance our
understanding of the neural processes associated with disorders affecting the fluent
production motor sequences, such as stuttering, dyspraxia, and task-dependent
dystonia[60–64].

Methods

Participants

Data were collected from a total of 55 right-handed University students
(Experiment 1: N = 19, 11 females; M = 24.2 years, SD = 4.1; Experiment 2: N=18, 11
females; M = 24.2 years, SD = 4.5; Experiment 3: N = 18, 9 females; M = 20.8 years,
SD = 2.4). Four additional participants were tested, but excluded from analysis based
on their sequence production error rate (cf. Participant exclusion criteria). They were
hypothesis-naive and had no previous exposure in performing a similar experimental
task. All participants had normal or corrected-to-normal vision and reported no history
of neurological or psychiatric disorders or hearing problems. Handedness was
evaluated through the online Handedness Questionnaire
(http://www.brainmapping.org/shared/Edinburgh.php) adapted from the Edinburgh
Handedness Inventory [65] (Experiment 1, M = 88.4, SD = 9.4; Experiment 2, M = 90.6,
SD = 9.7; Experiment 3, M = 90, SD = 11.8). All participants provided written informed
consent before participation and were debriefed after completing the study. They were

compensated either monetarily or with course credits at the end of the experiment. All

procedures were approved by the Bangor University School of Psychology Research

Ethics Committee (Ethics Review Board Approval Code 2017-16100-A14320).

Participant exclusion criteria

Mean finger and temporal interval error rate during sequence production in the

test phase (Day 3) above three standard deviations of the group mean performance

was considered as outlier performance, in each experiment separately. This was to

ensure that participants reached a comparable skill level in sequence performance

from memory and to have sufficient number of trials for RT analysis per participant,

which included correct trials only. We set this blindly to the individual *Probe* trial

performance to ensure that data exclusion was independent of the data analysed to

to test our main hypotheses. This resulted in the exclusion of data from one participant

in Experiment 1 who showed 53.1% finger error in the short delay, 54.7% in the

intermediate delay and 53.9% in the long preparation duration conditions. Two

participants' data sets were removed from Experiment 2, one with 25% finger error in

the slow timing and 18.8% in the irregular timing conditions, and the other with 44.5%

in the fast timing conditions. The data of one participant was excluded from Experiment

3 due to 12.5% finger error in the fast timing condition. No outlier performance was

found for temporal interval production in any condition of each experiment according

to the above criteria. Overall, the data of 19 participants were analyzed for Experiment

1, 18 participants for Experiment 2, and 18 participants for Experiment 3.

Apparatus

For all three experiments participants were seated in a quiet room in front of a

19-inch LCD monitor (LG Flatron L1953HR, 1280 x 1024 pixels, refresh rate 60Hz),

wearing headphones for noise isolation. All instructions about when each block began,

visual stimuli and feedback were controlled by Cogent 2000 (v1.29)

(http://www.vislab.ucl.ac.uk/cogent.php) through a custom-written MATLAB program

(v9.2 R2017a, The MathWorks, Inc., Natick, Massachusetts, United States) and

projected on to the LCD screen with inter-stimulus-intervals calculated in refresh rates

to ensure precise stimulus timing. In Experiments 1 and 2, a customized foam channel

was attached to the outer-half surface of the table to stabilize the cable of a Pyka 5-

button fiber optic response device (Current Designs). A thin anti-slip black mat was

placed underneath the response device to prevent sliding during the task. The

response device was positioned horizontally and adjusted accordingly for each

participant to ensure good control over the target buttons as well as arm and wrist

comfort. Participants were instructed to place the right index, middle, ring and little

fingers on the respective target buttons of the device. Experiment 3 used an identical

experimental set-up with the exception that responses were recorded using a

computer keyboard and participants were instructed to place their right thumb in

addition to the rest of the right-hand fingers on the designated keyboard keys. For

hand stabilization and comfort their wrist was positioned on a wrist rest.

Behavioural task and design

In Experiments 1 and 2, the task involved the recording of sequential and single

button presses produced with the four fingers (index, middle, ring and little) of the right

hand on a response device while performing a visually cued motor learning task

adapted from Kornysheva et al.[11]. Experiment 3 additionally required single presses

with the thumb. Participants were trained to associate a visual cue (an abstract fractal

shape, henceforth *Sequence* cue) with a specific a four-element finger sequence

produced with a specific timing. In all experiments, the paradigm employed two main

trial types: sequence and single-press (*Probe*) trials. *Sequence* trials were further

divided into visually instructed and memory-guided trials. Instructed trials involved the

presentation of four visual digit cues (index, middle, ring and little) at specified intervals

comprising a unique target sequence. These were only used during training in the first

two days, and during two refresher blocks on the third day. The test phase on the third

day only involved sequence production without visual guidance (memory-guided trials,

Supplementary Figure 3). *Probe* trials involved the production of only one visual digit

cue (*Probe* cue) corresponding to one of the serial positions in the target sequence

(Figure 1b).

Experiment 1. All participants were trained in producing two four-element target

sequences comprising two different finger order types (F1, F2) with isochronous

temporal intervals of 800 ms between presses (T1). Two additional finger order types

(F3, F4) of the same temporal sequence (T1) served as practice sequences to impose

familiarization with the task. Four additional finger order types (F5, F6, F7, F8) with

isochronous intervals of 800 ms (T1) were used to evaluate sequence-specific learning

in a visually cued task alongside the target sequences, immediately before and after

the training phase. The data from this control task is not presented here, as the current

work evaluates the preparation and performance during trials involving production
from memory. As a result, the experiment employed a total of eight unique *Sequence*
cues associated with eight finger sequences. The sequences were randomly
generated offline through a custom-written MATLAB code for each participant.
Specifically, the sequence generation process produced sequences for each
participant randomly excluding sequences with ascending and descending digit triplets.
The trained sequences started with different digits.

All trial types started with a *Sequence* cue. The *Sequence* cue had a fixed
duration of 400ms followed by a fixation cross, the latency of which varied depending
on the delay period from *Sequence* cue onset to *Go* cue. The resultant short (500 ms),
intermediate (1000 ms), and long (1500 ms) delay periods following the *Sequence* cue
comprised the three preparation duration conditions employed in the task. After the
delay period, a black hand stimulus appeared as the *Go* cue. In an instructed trial, the
Go cue was presented on a grey background for 2400 ms, guiding the participants
throughout the execution of the sequence by sequentially displaying a small white
circle on the digits of the hand stimulus. This acted as a visual digit cue appearing
sequentially on each of the four digits, with the time intervals between the digit cues
forming the target temporal structure of the sequence (T1) and defining its duration of
2400 ms. To achieve finger and temporal accuracy during training, participants were
asked to press the correct target buttons in synchrony with the digit cues until the
completion of the sequence, with the aim to progress towards synchronization with the
target timing. As the first digit cue of a sequence appeared at the same time as the *Go*
cue, immediate initiation of the sequence was emphasized in the instructions.

In memory-guided trials, a green rectangle was used as a background for the
Go cue, remaining on the screen for 2400 ms. Memory-guided trials featured the *Go*
cue without the appearance of digit cues, requiring participants to produce the
upcoming target sequence from memory. In these trials, participants were instructed
to initiate the sequence as quickly as possible and produce the sequence according
to its target finger and temporal structure (i.e. F1T1, F2T1).

In probe trials, the *Go* cue was displayed for 1000 ms on a grey background
with a digit cue presented on one digit (*Probe* cue), prompting a single press of the
corresponding target button. Here, the instructions were to respond to the *Probe* cue

as fast and accurately as possible. Participants were encouraged to avoid premature
responses (before the *Go* cue) in all trial types.

Following the *Go* cue, a fixation cross (1000 ms) and, subsequently, feedback
(1000 ms) were presented on the screen. The duration of a sequence trial including
feedback was 5.4 s, while a probe trial had a duration of 4 s. The inter-trial-interval (ITI)
was fixed at 800 ms. The experiment consisted of two 90min long training (Days 1 and
2) and a test (Day 3) sessions taking place over three consecutive days. Day 1
commenced with a practice block which involved two instructed and two memory-
guided *Sequence* trials for each of the target finger sequences as well as two random
probe trials, all randomly combined with the three delays. Over the three days,
participants serially underwent a pre-training (2 blocks), a training (36 blocks), a post-
training (2 blocks) and a test phase (2 refresher training blocks + 16 test blocks)
completing a total of 58 blocks. Participants were naïve as to the structure of the
gradual transition from the training through to the testphase and which block type they
were administered. The pre- and post-training blocks consisted of 24 instructed trials
each; each block was 2.48 min long and contained randomized mixed repetitions of
the two target and four control sequences matched equally with the delay conditions.
The training phase was organized in three stages: 12 blocks of 288 instructed and 72
probe trials (stage A, 80% instructed and 20% probe trials in each block), 12 blocks of
144 instructed, 144 from memory and 72 probe trials (stage B, 40% for each sequence
type and 20% probe trials in each block), and 12 blocks of 288 memory-guided and
72 probe trials (stage C, 80% memory and 20% probe trials in each block). A training
block (3 min long) of either stage consisted of 30 trials. On each block there was a
stable 20% occurrence of probe trials (6 in each block) comprising a total of 216 probes
throughout the training blocks. Distribution of probe trials in this phase was determined
by the minimum number of trials possible, namely 24 (2 sequences × 3 delays × 4
probe digits), and the block repeats. Eventually each probe digit occurred 18 times in
each training stage. All 40 blocks were evenly assigned to the study sessions such
that from Day 1 through the end of Day 2 participants had completed the training and
the post-training synchronization task. The testphase (Day 3) started with two
refresher training blocks of mixed type and immediately progressed to 16 blocks of 48
trials each, in which 24 memory-guided and 24 probe trials were randomly presented.
Duration of a test block was 4.4 min. The two trained sequences used in the memory-

guided trials were matched to the three delay conditions with each combination being
repeated four times within the block. This gave a total of 128 memory-guided trials per
delay condition, across blocks. In probe trials, each probe digit was combined with the
three delay conditions resulting in 32 trials per digit per delay condition. The testphase
had a total of 768 trials (384 memory-guided sequences and 384 probes). Overall, the
participants underwent 2004 trials excluding the practice trials.

Experiment 2. Procedures for Experiment 2 were identical to Experiment 1
except that the preparation period was fixed at 1500 ms and participants were trained
in associating three unique *Sequence* cues with one finger sequence (F1) to be
performed with three target temporal structures (T1, T2, T3) or IPIs: slow (T1, 800-
800-800 ms), fast (T2, 400-400-400 ms) and irregular (T3, 400-1600-400 ms), forming
respective target sequence durations of 2400 ms, 1200 ms, and 2400ms. The trial
followed the same structure as in Experiment 1, but the *Go* cue remained on the
screen for 3000 ms in a sequence trial and for 1000 in a probe trial. This was followed
by a fixation cross (1000 ms) and feedback (1000 ms) with a varying ITI of 500, 900
and 1300 ms. As a result, a sequence trial was 6.5 min long and a probe trial 4.5 min
long. The participants underwent the same structure of training and testsessions as in
Experiment 1. Similarly, we conducted a synchronization task, in a pre-post design.
Here, three additional control timing conditions were used to test for temporal transfer
(trained timing conditions) with the target timing patterns combined with a different
finger sequence. Overall, in this experiment participants were exposed to 15 unique
temporally structured sequences associated with their respective *Sequence* cue and
completed 2016 trials over 58 blocks.

Experiment 3. The training/testprocedures, trial structure, and the pre/post-
training synchronization task in Experiment 3 were identical to those of Experiment 2,
except that probe trials would additionally cue the thumb. This served as a control
condition to obtain reaction times and error rates for unplanned responses as thumb
presses were not part of any learnt finger sequence. Across each training stage, there
were 60 probe trials, while the testphase (30 blocks × 26 trials) contained 360 memory-
guided *Sequence* trials (120 trials per timing condition), 360 probe trials (30 trials per
digit per timing condition), and 60 thumb probe trials (20 trials per timing condition).
Overall, participants completed 1990 trials over 72 blocks, excluding the practice block.

Feedback. In all experiments, a points system was designed to reward fast
initiation and accurate performance, and avoid any drift in the motor production from
memory. After each sequence trial, feedback was presented on the screen for 1000
ms in the form of points (0-10) based on three performance criteria: reaction time (RT)
to assess sequence initiation, percentage of deviation from the target temporal
intervals of the sequence, and finger press accuracy. Points gained from the RT
component of the sequence, i.e. response from *Go* cue to the first press, were defined
by tolerance RT windows of 0-200, 200-360, 360-480, 480-560, 560-600 ms resulting
in 5, 4, 3, 2 and 1 points, respectively. For late (> 600) responses, 0 points were given.
A schematic feedback provided information on both finger accuracy and temporal
sequence accuracy performance. An '*x*' or a '-' symbol was shown for every correct or
incorrect press, respectively. Temporal errors were calculated after each trial as
deviations of press from target timing in percent of the target interval to account for the
scalar variability of timing[66,67]. Thresholds for mean absolute percentage deviation
across all correct presses were set at 10, 20, 30, 40 and 50 percent assigning 5, 4, 3,
2 and 1 points, respectively. Timing interval deviation > 50% resulted in 0 points. If a
press was performed too early the respective symbol was displayed below the midline,
while for a late response it was displayed above. This applied only to the second, third
and fourth presses of the sequence, whilst the first symbol reflecting the first press
was always positioned on the midline, representing the starting point of the sequence.
Deviation from target onset (presented or assumed) rather than interval timing
encouraged participants to synchronize with the visually cued sequences during
training, however, may have contributed to a tendency to compress the overall
sequence length during trials produced from memory.
Participants were instructed to adjust their performance by keeping the crosses
as close to the midline as possible. If at least one incorrect press or an incorrect
number of presses was recorded (< 4 or > 4), 0 points were given on that trial. The
points on each trial were displayed above the schematic feedback, and were the sum
of the RT, interval deviation and finger accuracy points. The feedback following a probe
trial displayed only points (0-5) gained based on RT and finger press accuracy utilizing
the same tolerance windows as described above for assessing sequence initiation RT.
In the case of an incorrect press or incorrect number of presses (< 1 or > 1), 0 points
were given regardless of the RT length. To incentivize the participants to gain as many

points as possible on each trial, we offered an extra monetary reward (10£) to those

two with the highest total points.

Data analysis

Data analyses were performed using custom written code in Matlab (v9.2

R2017a, The MathWorks, Inc., Natick, Massachusetts, United States) and SPSS

version 22.0 (IBM Corp., Armonk, N.Y., USA). Median reaction time (RT; correct trials

only) and mean error rates for each *Probe* position and condition were calculated

relative to the 1st position in each participant and condition (RT and error rate increase

in %).

Repeated measures ANOVAs were undertaken for RT and error rate in *Probe*

trials, and for inter-press-intervals (IPI), temporal error, finger error, and sequence

initiation RT in *Sequence* trials produced from memory. Planned contrast analyses for

the main and interaction terms of interest in each ANOVA model involved user-defined

orthogonal contrasts. To evaluate the RT and error rate increase for the control action

(Experiment 3), we used two-tailed paired-samples t-tests (control *vs* 4th position).

Mean relative increase between adjacent positions (1st to 2nd, 2nd to 3rd, 3rd

to 4th) for RT (CQ RT increase) and press error rate (CQ error increase) in *Probe* trials

were taken as a measure for the strength of the competitive action gradient during

preparation. Using the group data (N = 55), we conducted six planned one-tailed

Pearson correlations between the CQ strength derived from RT and error increase,

respectively, and each of the sequence production measures.

CQ model of sequence preparation

To our knowledge, no CQ model has previously been applied to response

preparation. While models differ somewhat with respect to how sequence position is

represented, they all require some form of "start state", which has stronger links to

items or responses that should occur earlier in the sequence[5,8,68,69]. In the

implementation, we use the Start-End CQ model[16], but we expect that other CQ

models with distinct start states would behave similarly if the same assumptions

regarding preparation were added to them.

The model makes the following assumptions: Each learned sequence is

hierarchically organized, with a "sequence node (or nodes)" linking to, and activating,

the responses which make up the sequence. The sequence node activates the

position codes (associations) of the items in the sequence. Following the *Sequence*

cue, the start-state of the cued sequence becomes gradually activated. The start-state

of the intended sequence produces an activation gradient over the planned responses

based on its strength of association with them. The additions to the published CQ

model of Houghton (2018)[16] consist of: (i) the gradual increase of the start state

activation, (Equation 1); (ii) the damping of response activations prior to the *Go* cue

(Equation 3); and inhibition of the competitive response selection process, also prior

to the *Go* cue. The model's activation of planned responses during preparation is

shown in Figure 5.

Sequence node activation. Following presentation of the *Sequence* cue, the

associated sequence node s_j becomes gradually activated. We implement a simple

linear "ramp",

$$a_j(t) = b_j + ct$$

where t is (discrete) time (since cue presentation), b_j is the baseline activation,

and c is the rate of increase. The activation of the start-state S_j retrieved by the learned

sequence s_j follows this activation,

$$S_j(t) = a_j(t)S_j \qquad (1)$$

S_j (without a time index) is the asymptotic (i.e., stored) value for the sequence,

set here to unity. The effect of this is simply that the start-state gradually increases its

activation following the cue.

Input to response nodes. Activation spreads from the start-state $S_j(t)$ to finger

responses via its positional associations W_j (a matrix) with the response tokens[16]. The

input from sequence s_j to its associated actions is given by,

$$Q(t) = g\big(S_j(t)W_j\big) \qquad (2)$$

Here $Q(t)$ (a matrix) represents the state of the queue of response tokens. W_j

encodes the positional weights from the sequence level to the responses in s_j, and the

product $S_j(t)W_j$ computes the differences between the start state of the sequence

$(S_j(t))$ and the position codes W_j of the sequence items (represented using the phase

code of Houghton, 2018).

Finally, the function g represents a Gaussian receptive field, or positional tuning

curve, applied element-wise to the signals coming from the matrix W_j. The receptive

field is sensitive to the difference between the state of the start-signal S_j and the

position codes in W_j, peaking when they are identical (see Houghton, 2018, Equation

1). The receptive field has a tuning (variance) parameter σ, controlling the model's

sensitivity to positional differences (Figure 5a).

Finger response activation. Each finger response in the sequence s_j becomes

gradually active during preparation, due to the increasing activation of the sequence's

start state (Equation 1), sending an increasing degree of activation to the responses

in the sequence, modulated by the similarity of the response's position code to the

start state (Equation 2). For a finger response $F_{k,j}$ (i.e., *k*th finger in sequence *j*), its

activation during preparation is, in discrete time form,

930
$$F_{k,j}(t+1) = 1 - F_{k,j}(t) \times \left(\delta F_{k,j}(t) + Q_k(t) \right) \qquad (3)$$

In the second term on the right, δ controls the decay rate (of the current

activation level), and $Q_k(t)$ is the input to finger F_k given by Equation 2. Note that if the

latter equals 0, then response activation spontaneously decays due to the decay term,

$\delta < 1$.

The first term on the right, $1 - F_{k,j}(t)$, acts as "damping" factor; it becomes

smaller as the response activation increases towards a ceiling of 1. This prevents

activations growing without bound as the preparation interval increases (Figure 5b). It

is proposed that on detection of the *Go* signal, this damping term ceases to act,

permitting activations to rapidly increase, and initiating the competitive response

selection process intrinsic to CQ models.

References

1.   Rosenbaum, D. A. Motor programming. *Massachsetts Inst. Technol. Cambridge,*

MA. (1984).

2.   Lashley, K. S. The problem of serial order in behavior. in *Cerebral mechanisms*
in behavior 112–131 (Wiley, 1951).

3.   Ariani, G. & Diedrichsen, J. Sequence learning is driven by improvements in
motor planning. *J. Neurophysiol.* (2019). doi:10.1152/jn.00041.2019

4.   Remington, E. D., Egger, S. W., Narain, D., Wang, J. & Jazayeri, M. A Dynamical
Systems Perspective on Flexible Motor Timing. *Trends Cogn. Sci.* **22**, 938–952
(2018).

5.   Houghton, G. The problem of serial order: A neural network model of sequence
learning and recall. in *Current Research in Natural Language Generation* (1990).

6.   Hartley, T. & Houghton, G. A linguistically constrained model of short-term
memory for nonwords. *J. Mem. Lang.* **35**, 1–31 (1996).

7.   Burgess, N. & Hitch, G. J. Memory for serial order: A network model of the
phonological loop and its timing. *Psychol. Rev.* **106**, 551–581 (1999).

8.   Henson, R. N. A. Short-Term Memory for Serial Order: The Start-End Model.
Cogn. Psychol. **36**, 73–137 (1998).

9.   Glasspool, D. W. & Houghton, G. Serial order and consonant-vowel structure in
a graphemic output buffer model. *Brain Lang.* **94**, 304–330 (2005).

10.   Behmer, L. P. *et al.* Parallel regulation of past, present, and future actions during
sequencing. *J. Exp. Psychol. Hum. Percept. Perform.* **44**, 1147–1152 (2018).

11.   Kornysheva, K. *et al.* Neural Competitive Queuing of Ordinal Structure Underlies
Skilled Sequential Action. *Neuron* **101**, 1166-1180.e3 (2019).

12.   Pinet, S., Dell, G. S. & Alario, F. X. Tracking keystroke sequences at the cortical
level reveals the dynamics of serial order production. *J. Cogn. Neurosci.* (2019).
doi:10.1162/jocn_a_01401

13.   Basu, D. & Murthy, A. Parallel programming of saccades in the macaque frontal
eye field: Are sequential motor plans coactivated? *J. Neurophysiol.* (2020).
doi:10.1152/jn.00545.2018

14.   Averbeck, B. B., Chafee, M. V., Crowe, D. A. & Georgopoulos, A. P. Parallel
processing of serial movements in prefrontal cortex. *Proc. Natl. Acad. Sci.* **99**,
13172–13177 (2002).

15.   Page, M. P. A. & Norris, D. The Primacy Model: A New Model of Immediate Serial
Recall. *Psychol. Rev.* **105**, 761–781 (1998).

16.   Houghton, G. Action and perception in literacy: A common-code for spelling and

reading. *Psychol. Rev.* (2018). doi:10.1037/rev0000084

17. Burgess, N. & Hitch, G. J. A revised model of short-term memory and long-term

learning of verbal sequences. *J. Mem. Lang.* **55**, 627–652 (2006).

18. Friedman, J. & Korman, M. Offline optimization of the relative timing of

movements in a sequence is blocked by retroactive behavioral interference.

Front. Hum. Neurosci. **10**, 1–15 (2016).

19. Klapp, S. T. Reaction time analysis of central motor control. in *Advances in Motor*

Learning and Control (1996).

20. Sternberg, S., Monsell, S., Knoll, R. L. & Wright, C. E. The Latency and Duration

of Rapid Movement Sequences: Comparisons of Speech and Typewriting. in

Information Processing in Motor Control and Learning (1978).

doi:10.1016/b978-0-12-665960-3.50011-6

21. Niemi, P. & Naatanen, R. The foreperiod and simple reaction time. *Psychol. Bull.*

89, 133–162 (1981).

22. Vallesi, A., Shallice, T. & Walsh, V. Role of the prefrontal cortex in the foreperiod

effect: TMS evidence for dual mechanisms in temporal preparation. *Cereb.*

Cortex (2007). doi:10.1093/cercor/bhj163

23. Hurlstone, M. J. Functional similarities and differences between the coding of

positional information in verbal and spatial short-term order memory. *Memory*

27, 147–162 (2019).

24. Bullock, D. Adaptive neural models of queuing and timing in fluent action. *Trends*

Cogn. Sci. **8**, 426–433 (2004).

25. Hartley, T., Hurlstone, M. J. & Hitch, G. J. Effects of rhythm on memory for

spoken sequences: A model and tests of its stimulus-driven mechanism. *Cogn.*

Psychol. **87**, 135–178 (2016).

26. Cisek, P. & Kalaska, J. F. Neural correlates of reaching decisions in dorsal

premotor cortex: specification of multiple direction choices and final selection of

action. *Neuron* **45**, 801–814 (2005).

27. Verwey, W. B. & Abrahamse, E. L. Distinct modes of executing movement

sequences: reacting, associating, and chunking. *Acta Psychol. (Amst).* **140**,

274–82 (2012).

28. Rhodes, B. J., Bullock, D., Verwey, W. B., Averbeck, B. B. & Page, M. P. A.

Learning and production of movement sequences: Behavioral,

neurophysiological, and modeling perspectives. *Hum. Mov. Sci.* **23**, 699–746
(2004).

29.   Behmer, L. P. & Crump, M. J. C. The dynamic range of response set activation
during action sequencing. *J. Exp. Psychol. Hum. Percept. Perform.* (2017).
doi:10.1037/xhp0000335

30.   Gilbert, R. A., Hitch, G. J. & Hartley, T. Temporal precision and the capacity of
auditory–verbal short-term memory. *Q. J. Exp. Psychol.* (2017).
doi:10.1080/17470218.2016.1239749

31.   Ólafsdóttir, H. F., Bush, D. & Barry, C. The Role of Hippocampal Replay in
Memory and Planning. *Current Biology* (2018). doi:10.1016/j.cub.2017.10.073

32.   Liu, Y., Dolan, R. J., Kurth-Nelson, Z. & Behrens, T. E. J. Human Replay
Spontaneously Reorganizes Experience. *Cell* **178**, 640-652.e14 (2019).

33.   Eichenlaub, J. B. *et al.* Replay of Learned Neural Firing Sequences during Rest
in Human Motor Cortex. *Cell Rep.* **31**, 107581 (2020).

34.   Jafarpour, A., Fuentemilla, L., Horner, A. J., Penny, W. & Duzel, E. Replay of
Very Early Encoding Representations during Recollection. *J. Neurosci.* **34**, 242–
248 (2014).

35.   Drieu, C. & Zugaro, M. Hippocampal sequences during exploration:
Mechanisms and functions. *Front. Cell. Neurosci.* **13**, 1–22 (2019).

36.   Dragoi, G. & Tonegawa, S. Preplay of future place cell sequences by
hippocampal cellular assemblies. *Nature* (2011). doi:10.1038/nature09633

37.   Michelmann, S., Staresina, B. P., Bowman, H. & Hanslmayr, S. Speed of time-
compressed forward replay flexibly changes in human episodic memory. *Nat.*
Hum. Behav. **3**, 143–154 (2019).

38.   Kurth-Nelson, Z., Economides, M., Dolan, R. J. & Dayan, P. Fast Sequences of
Non-spatial State Representations in Humans. *Neuron* (2016).
doi:10.1016/j.neuron.2016.05.028

39.   Davidson, T. J., Kloosterman, F. & Wilson, M. A. Hippocampal Replay of
Extended Experience. *Neuron* (2009). doi:10.1016/j.neuron.2009.07.027

40.   Li, N., Daie, K., Svoboda, K. & Druckmann, S. Robust neuronal dynamics in
premotor cortex during motor planning. *Nature* **532**, 459–464 (2016).

41.   Wong, A. L. & Krakauer, J. W. Why Are Sequence Representations in Primary
Motor Cortex So Elusive? *Neuron* (2019). doi:10.1016/j.neuron.2019.09.011

42.   Krakauer, J. W. & Mazzoni, P. Human sensorimotor learning: adaptation, skill,
and beyond. *Curr. Opin. Neurobiol.* **21**, 636–644 (2011).

43.   Wang, J., Narain, D., Hosseini, E. A. & Jazayeri, M. Flexible timing by temporal
scaling of cortical responses. *Nat. Neurosci.* (2018). doi:10.1038/s41593-017-
0028-6

44.   Goudar, V. & Buonomano, D. V. Encoding sensory and motor patterns as time-
invariant trajectories in recurrent neural networks. *Elife* (2018).
doi:10.7554/eLife.31134

45.   Burgess, N. & Hitch, G. Computational models of working memory: Putting long-
term memory into context. *Trends Cogn. Sci.* **9**, 535–541 (2005).

46.   Bortoletto, M., Cook, A. & Cunnington, R. Motor timing and the preparation for
sequential actions. *Brain Cogn.* (2011). doi:10.1016/j.bandc.2010.11.016

47.   Bortoletto, M. & Cunnington, R. Motor timing and motor sequencing contribute
differently to the preparation for voluntary movement. *Neuroimage* (2010).
doi:10.1016/j.neuroimage.2009.11.048

48.   Maslovat, D., Chua, R., Klapp, S. T. & Franks, I. M. Preparation of timing
structure involves two independent sub-processes. *Psychological Research* **82**,
981–996 (2018).

49.   Kornysheva, K., Sierk, A. & Diedrichsen, J. Interaction of temporal and ordinal
representations in movement sequences. *J. Neurophysiol.* **109**, 1416–24 (2013).

50.   Kornysheva, K. & Diedrichsen, J. Human premotor areas parse sequences into
their spatial and temporal features. *Elife* **3**, e03043 (2014).

51.   Zeid, O. & Bullock, D. Moving in time: Simulating how neural circuits enable
rhythmic enactment of planned sequences. *Neural Networks* (2019).
doi:10.1016/j.neunet.2019.08.006

52.   Ullén, F., Bengtsson, S. L. & Ullen, F. Independent Processing of the Temporal
and Ordinal Structure of Movement Sequences. *J. Neurophysiol.* **90**, 3725–3735
(2003).

53.   Shin, J. C. & Ivry, R. B. Spatial and Temporal Sequence Learning in Patients
with Parkinson's Disease or Cerebellar Lesions. *J. Cogn. Neurosci.* **15**, 1232–
1243 (2003).

54.   O'Reilly, J. X., McCarthy, K. J., Capizzi, M. & Nobre, A. C. Acquisition of the
temporal and ordinal structure of movement sequences in incidental learning. *J.*

Neurophysiol. **99**, 2731–2735 (2008).

55.   Bengtsson, S. L., Ehrsson, H. H., Forssberg, H. & Ullén, F. Dissociating brain
regions controlling the temporal and ordinal structure of learned movement
sequences. *Eur. J. Neurosci.* **19**, 2591–2602 (2004).

56.   Bengtsson, S. L., Ehrsson, H. H., Forssberg, H. & Ullén, F. Effector-independent
voluntary timing: behavioural and neuroimaging evidence. *Eur. J. Neurosci.* **22**,
3255–3265 (2005).

57.   Friston, K. & Buzsáki, G. The Functional Anatomy of Time: What and When in
the Brain. *Trends Cogn. Sci.* **20**, 500–511 (2016).

58.   Crowe, D. A., Zarco, W., Bartolo, R. & Merchant, H. Dynamic representation of
the temporal and sequential structure of rhythmic movements in the primate
medial premotor cortex. *J. Neurosci.* **34**, 11972–83 (2014).

59.   Merchant, H., Perez, O., Zarco, W. & Gamez, J. Interval Tuning in the Primate
Medial Premotor Cortex as a General Timing Mechanism. *J. Neurosci.* **33**,
9082–9096 (2013).

60.   Miller, N. The assessment of limb dyspraxia. *Clin. Rehabil.* **2**, 177–181 (1988).

61.   Sadnicka, A., Kornysheva, K., Rothwell, J. C. & Edwards, M. J. A unifying motor
control framework for task-specific dystonia. *Nature Reviews Neurology* **14**,
116–124 (2018).

62.   Howell, P. A model of serial order problems in fluent, stuttered and agrammatic
speech. *Hum. Mov. Sci.* (2007). doi:10.1016/j.humov.2007.07.004

63.   Craig-McQuaide, A., Akram, H., Zrinzo, L. & Tripoliti, E. A review of brain
circuitries involved in stuttering. *Front. Hum. Neurosci.* **8**, 1–20 (2014).

64.   Ingham, R. J., Ingham, J. C., Euler, H. A. & Neumann, K. Stuttering treatment
and brain research in adults: A still unfolding relationship. *Journal of Fluency*
Disorders (2018). doi:10.1016/j.jfludis.2017.02.003

65.   Oldfield, R. C. The assessment and analysis of handedness: the Edinburgh
inventory. *Neuropsychologia* **9**, 97–113 (1971).

66.   Rakitin, B. C. *et al.* Scalar expectancy theory and peak-interval timing in humans.
J. Exp. Psychol. Anim. Behav. Process. **24**, 15–33 (1998).

67.   Jazayeri, M. & Shadlen, M. N. Temporal context calibrates interval timing. *Nat.*
Neurosci. **13**, 1020–1026 (2010).

68.   Brown, G. D. A., Hulme, C. & Preece, T. Oscillator-Based Memory for Serial

Order. *Psychol. Rev.* **107**, 127–181 (2000).

69.   Burgess, N. & Hitch, G. J. Toward a network model of the articulatory loop. *J.*

Mem. Lang. (1992). doi:10.1016/0749-596X(92)90022-P

1114