The effect of plant-derived carbon on rhizospheric microbial communities decomposing different litter types and nutrient return

Running title: Plant effects on nutrient cycling in the rhizosphere

Somak Chowdhury[1], Markus Lange[1], Ashish A Malik[2] Timothy Goodall[3], Jianbei Huang[1], Robert I Griffiths[4] and Gerd Gleixner[1]

Affiliations –

[1] Department of Biogeochemical Processes, Max Planck Institute for Biogeochemistry, 07745, Jena, Germany

[2] School of Biological Sciences, University of Aberdeen, Aberdeen, UK

[3] Centre for Ecology and Hydrology, Wallingford, Oxfordshire, UK

[4] Centre for Ecology and Hydrology, Bangor, UK

Corresponding author: Gerd Gleixner

Institutional Affiliation: Department of Biogeochemical Processes,

Max Planck Institute for Biogeochemistry,

07745, Jena, Germany

Email: gerd.gleixner@bgc-jena.mpg.de

Phone no: +493641576172

Competing Interests statement: The authors declare that they have no conflict of interest.

Keywords: AMF; Pulse labeling; ^{13}C; ^{15}N; ingrowth core; PLFA; 16S rRNA Gene; ITS;

Abstract

Interactions between plants and microorganisms strongly affect ecosystem functioning as processes of plant productivity, litter decomposition and nutrient cycling are controlled by both organisms. Though two-sided interactions between plants and microorganisms and between microorganisms and litter decomposition are areas of major scientific research, our understanding of the three-sided interactions of plant-derived carbon flow into the soil microbial community and their follow-on effects on ecosystem processes like litter decomposition and plant nutrient acquisition remains limited. Therefore, we performed a microcosm experiment with two plant communities differing in their association with arbuscular mycorrhizal fungi (AMF). By applying a $^{13}CO_2$ pulse label to the plant communities and adding various ^{15}N labelled litter types to ingrowth cores, we simultaneously traced the flow of plant-derived carbon into soil microbial communities and the return of mineralized nitrogen back to the plant communities. We observed that net ^{13}C assimilation by the rhizospheric microbial communities and community composition depended on both, plant-AMF association and the litter type being decomposed. AMF-associated plant communities invested less ^{13}C into the litter decomposing soil microbial community but gained similar amounts of ^{15}N from litter decomposition compared to non-AMF plant communities. This effect was driven by significantly lower net ^{13}C assimilation per ^{15}N gain of saprotrophic fungi and bacterial, and significantly higher one of AM fungi compared to NM plant communities. Additionally the net ^{13}C assimilation of the rhizospheric microbial community from AMF-associated plant communities depended on litter type. Lowest ^{13}C assimilation was found for microbial and plant litter decomposition and highest for inorganic nitrogen. Large parts of the soil microbial community showed most interestingly redundant functions and different parts of the microbial community were dependent on the

50   plant-microorganism-litter context activated by plant-derived carbon in order to decompose

51   specific litter types. Therefore, our results suggest that ecosystem processes like litter

52   decomposition can only be fully understood if the whole plant-microorganism-litter context is

53   investigated. Moreover our results give first hints that the plant AMF association might help

54   to get less carbon costly access to nutrients locked in bacterial and plant necromass.

Introduction

Plant growth is mostly limited by the availability of nutrients, such as nitrogen (N) and phosphorous (P) in soil (Veresoglou et al., 2012). These nutrients are not readily available to plants as they are bound in soil organic matter and the mineral matrix. Consequently plants invest substantial parts of their assimilated carbon into soil via root exudates (sugars, amino acids and organic acids) and root growth associated cell debris, which stimulates microbial decomposition of soil organic matter and releases nutrients for plant and microbial reuse (Jacoby et al., 2017; Kuzyakov and Xu, 2013). However, immediate uptake of the released nutrients by plants is not always possible (Dijkstra et al., 2013) as the soil microbial communities themselves have a high nutrient demand and assimilate nutrients more efficiently than plants (Levin and Angert, 2015; Zhu et al., 2017). Since nutrient cycles in soil are almost entirely controlled by microorganisms, variations in plant exudation patterns and associations of plants and microorganism can have a tremendous impact on the rates of nutrient cycling (Craine et al., 2007; Veresoglou et al., 2012).

To adequately benefit from the microbial nutrient release some plants rely solely on their root systems while many others associate with specific soil microorganisms that colonize roots (Bonfante and Genre, 2010; Nuccio et al., 2013). A well-known association of this kind occurs between plants and arbuscular mycorrhizal fungi (AMF) (Davison et al., 2015). Though AMF do not decompose organic matter themselves (Tisserant et al., 2013), they colonize other saprophytic organisms that mineralize soil organic matter and compete with them for the nutrients at similar uptake rates (Bunn et al., 2019; Cheng et al., 2012; Thirkell et al., 2016). Plant support this process by altering their root exudate composition (Bonfante and Genre, 2010) and use the hyphal network to increase the soil volume that is available for nutrient uptake (Bitterlich et al., 2018). In contrast, non-mycorrhizal plants only release labile carbon to the microbial communities around their roots and lack the advantages of extended

soil volume accessibility and nutrient uptake efficiency (Karlowsky et al., 2018). As a result,

they only benefit indirectly from decomposition related nutrient release. However, it remains

unclear whether there are trade-offs for plants that depend on the type of belowground

microbial associations and to what extent presence or absence of such microbial associations

can affect decomposition related microbial processes.

For plants, the AMF-associated (AM) or non-mycorrhizal (NM) nutrient uptake strategies

involve export of different amounts of carbon into microbial communities. AM plants are

suggested to export more carbon to microbial communities than NM plants (Drigo et al.,

2010; Scheffknecht et al., 2007) as the former supports an additional hyphal network

(Cavagnaro et al., 2012). In comparison to NM plants this hyphal network supports the AM

plants in multiple ways such as offering its roots greater soil coverage (Sanders and Tinker,

1973), access to immobile nutrient pools (Cavagnaro et al., 2012) and access to distant water

pockets (16, 21). Hence, the overall plant carbon allocation into the root systems and the

carbon export to associated microbial processes vary depending on the nutrient uptake

strategy, which was successfully demonstrated using mesh guarded ingrowth cores (Nuccio et

al., 2013; Throckmorton et al., 2012). However, it remains unclear if carbon intensive specific

mutualistic associations like with AMF also result in higher carbon costs per nutrient return

and if the net plant carbon uptake by decomposing microorganisms is directly related to the

plant nutrient uptake.

The microbial decomposition process itself is well understood. Microbial growth and

activity in soil is primarily limited by the availability of labile carbon which is either released

via plant exudation or decomposition of plant litter and soil organic matter (Hütsch et al.,

2002). Both quantity of labile carbon inputs (de Graaff et al., 2010) and alteration of

biochemical profile of root exudates by AMF infection (Shi et al., 2016; Zhalnina et al., 2018)

can alter activity and composition of microbial communities associated with the rhizosphere

and litter decomposition (Cheng et al., 2012; Nottingham et al., 2013; Nuccio et al., 2013).

Similarly, interactions between litter quality and decomposer community are also well known and suggest that the amount and form of nitrogen play an important role in the extent of litter decomposition and nitrogen uptake by the plants (Schimel and Bennett, 2004). So far it was considered that bacterial members of microbial community favor nitrogen rich substrates whereas fungal members favor carbon rich lignocellulosic substrates during decomposition (Güsewell and Gessner, 2009; Meier et al., 2015; Strickland et al., 2009). Recent studies, however, have demonstrated that many bacteria also decompose lignocellulose-rich substrates (López-Mondéjar et al., 2016) and that the soil microbial community is in general characterized by a high functional redundancy (Louca et al., 2018).

Soil receives nitrogen containing substances either via carbon rich plant residues, nitrogen rich microbial necromass or inorganic nitrogen from deposition and fertilizers (Gleixner, 2013; Liang et al., 2017; Thuille et al., 2015). However, these nitrogen inputs differ in their biochemical properties (litter type) and carbon to nitrogen ratios (litter quality), thus confounding the effects that plant communities have on the decomposer communities in soil. Under natural conditions, the decomposer community is shaped by litter characteristics and their physiological boundaries, which are defined by their energetic and nutritional demands (Thomson et al., 2013). Hence, by exposing rhizospheric microbial communities to litter of identical quality but different types, a better understanding of the mechanisms by which plants affect the decomposer community can be obtained.

In summary, large uncertainties still exist regarding the mechanistic benefits of plant-microbial associations with respect to litter decomposition and nutrient return. Therefore, we assembled two plant communities that differed in their ability to establish an AMF-based mycorrhizal association (AM or NM) and amended the soil inside the root ingrowth cores with various ^{15}N labeled litter types (plant root litter, microbial necromass and inorganic N) of similar quality (C/N ratio) and traced the net plant derived carbon flow into the microbial community (PLFA) using a $^{13}CO_2$ pulse label experiment and investigated the associated

microbial community structure (DNA and RNA). We first hypothesized that AM associations result in greater plant carbon export to the microbial community (including AMF) for nitrogen mineralization. We also hypothesized that carbon cost per nutrient gain is higher in AM association than in NM association, since AM associations require a considerable amount of plant carbon for maintaining the hyphal network. We finally hypothesize that the developed soil microbial community depends in the first order on the plant community (AM vs NM) and its carbon export, and in the second order on the litter type independent from its quality.

Materials and Methods

Microcosm Experiment

We set up microcosms (19 x 12.5 x 9.5 cm) and established mixed temperate plant communities of two types namely AMF dependent plant community (*Linum ussitatissimum, Linum perenne, Festuca rubra* and *Tagetes patula*) (n = 72 pots) termed as the AM plant community and non-mycorrhizal community (*Dianthus caryophyllus*, *Matthiola longipetala sub-species bicornis, Dianthus barbaratus* and *Carex arenaria*) (n = 36 pots) termed as NM plant community (Brundrett, 2015b, 2015a). The soil used for the experiment was collected from The Jena Experiment (Roscher et al., 2004) located in Jena, Germany in March 2015. After removing the root mat in upper 2 cm, the top 5 cm of soil was collected, sieved (<2 mm) and all remaining roots were removed. The soil was stored at 4 °C until the establishment of the microcosms

Each microcosm unit had a central PVC ingrowth core (diameter = 5 cm) with two tiers of 8 windows (4 x 1.5 cm) each guarded by a mesh (Figure 1). The mesh with a 1000 micron pore size allowed entry of fine roots and hyphae and the one with 30 micron pore size allowed entry of AMF hyphae only. The microcosms and the ingrowth cores were filled with soil and pre-equilibrated for 2 weeks in the greenhouse. Plant saplings that had developed the

first true leaves were planted with equal evenness into the pots excluding the ingrowth core. The microcosms were randomly distributed in the greenhouse and incubated for 4 months until the plants were fully developed. Irrigation was provided using an ebb and flood system for 5 min daily and lighting for the plants (Son- T Agro 430 W HPS bulbs, primary light range = 520–610 nm, Philips Lighting Company, Somerset, NJ, USA) was provided daily for 12 h.

For litter amendment to the fully developed microcosms, the central ingrowth core was removed. The soil was sieved (<2mm) to remove fine roots. 7.5, 0.8 & 0.012 g of ^{15}N labelled plant root litter, microbial necromass and inorganic nitrogen were respectively added to 200 g of sieved soil (Figure 1). An additional amount of 1.28 and 1.58 g of xanthan gum (Sigma-Aldrich, Inc) was added to the microbial necromass and inorganic nitrogen treatment respectively to keep the carbon and nitrogen inputs across all amendments (avg. C:N = 46 ± 1.1 s.e.m.) similar. Overall, the amendments were kept small, the total carbon and nitrogen content of the ingrowth core soil was increased by ca. 0.15 % and ca. 0.01 %, respectively. The amended soil was mixed, refilled into the ingrowth core and replaced into the microcosms (Figure 1).

2 weeks after the amendment, which allowed roots and hyphae to regrow into the ingrowth core, the microcosms were exposed to a $^{13}CO_2$ pulse in an air tight plexiglass chamber (Malik et al., 2015). For each litter amendment, all pots (n=27) were subjected to a $^{13}CO_2$ pulse together in a well-lit (PAR intensity >500 µmol photons m^{-2}s^{-1}) chamber (volume = 0.15 m^3). The chamber air was flushed with CO_2 free synthetic air until the CO_2 concentration fell below 150 ppm. Enriched $^{13}CO_2$ (99 atom %) was introduced into the chamber using a mass flow controller (model- MF1-20sccm-N2, MKS Instruments Inc.) at a flow rate of 3.33 x 10^{-7} m^{3}s^{-1} for 10 min and circulated using fans. The CO_2 concentration was monitored online by a cavity ringdown spectrometer (G2101-I, Picarro Inc., Santa Clara, CA, USA) throughout the labeling period of 1h. An additional maintenance stream of $^{13}CO_2$ was

184   introduced halfway at a flow rate of 1.67×10^{-7} m^3s^{-1} for 5 min that kept the chamber CO_2

185   concentration between 350 ppm and 410 ppm.

186   At the end of the labelling period the pots were removed from the chamber and

returned back to the greenhouse and destructively harvested at 0 h, 4 h, 24 h and 240 h from

the pulse. Soil from the ingrowth compartments were sieved (<2 mm) and divided into three

sub-samples and stored at -80 °C for molecular analysis within 1 h of harvest, dried at 105 °C

for 48 h and stored at room temperature for elemental analysis. The above and below ground

plant tissues were collected at the last destructive harvest (240 h) and dried at 60 °C for 48 h

and subsequently stored for future analysis. Control pots for each litter amendment (n=9)

were harvested prior to the labeling in order to avoid cross contamination from ^{13}C labeled

air.

196   Production of labeled litter

197   To follow nutrients released by decomposition to the plant tissues, ^{15}N labelled

organic matter of three types namely plant root litter, microbial necromass and synthetic

substrate were used in this study. Plant root litter was harvested from perennial grass *Holcus*

lanatus grown in a pot (57x 37 x 20 cm) containing sand in a plant growth chamber that was

lit daily for 12 h at day temperatures of 23 °C and at 63 % rel. humidity. The plants were

watered 2-3 times a week and fertilized weekly with a balanced N free Hoagland's solution

amended with 0.40 atom% $K^{15}NO_3$ as the sole nitrogen source. At the end of 4 weeks the

plants were harvested and roots were cleaned with distilled water, dried and shredded to

remove effects of particle size.

206   Inoculum for producing microbial necromass was produced by slow centrifugation of

6 g of Jena Experiment soil in 20 ml of distilled water at 650 g for 15 min. The supernatant

was inoculated into M9 minimal medium amended with 20% glucose and contained $K^{15}NO_3$

(0.40 atom%) as only nitrogen source. The culture grew for 7 days at 19 °C on a horizontal

shaker. The entire cell mass was harvested by centrifugation (@ 3500 g for 15 mins) and re-

suspended in 10 ml of sterile distilled water and sterilized by autoclaving. The sterile pellet

was freeze dried for further use.

The inorganic nitrogen amendments were generated by mixing KNO_3 and $K^{15}NO_3$

such that the mixture had an isotopic signature of 0.40 atom % and used directly for the

amendments.

Elemental and isotopic analysis of plant samples

The shoot samples that were harvested at the end of experiment were subjected to

carbon and nitrogen elemental and isotope analysis. Briefly, oven dried shoot and root

biomass was ball-milled to a fine powder and weighed into tin capsules. Separate samples

were prepared for bulk ^{13}C and bulk ^{15}N analysis. The samples were analyzed using an

elemental analyzer EA1100 (CE, Mainland) coupled via a conflow III interface to a DELTA+

(Finnigan MAT) isotope ratio mass spectrometer (Malik et al., 2015).

Biomarker extraction, quantification and compound specific isotope analysis

To assess the ^{13}C flow into the microbial community, fatty acids from neutral lipids

(NL) and phospholipids (PL) were extracted from ingrowth core soil sampled at 0 h and 240

h. All biological replicates of each litter-rhizosphere combination that were collected at the

same time point were pooled and homogenized to create a composite sample from which

always 5g aliquots of soil were used for lipid extraction. We applied a modified Bligh and

Dyer (1959) lipid extraction protocol using pressurized solvent extraction, as described earlier

Karlowsky et al., (2018). The concentration and compound specific ^{13}C incorporated relative

to an unlabeled sample for bacterial PLFA markers (14:0, 15:0, 16:0, 17:0, 18:0, 14:0i, 15:0i,

15:0a, 16:0i, 17:0i, 17:0a, 16:0(10Me), 17:0(10Me), 18:0(10Me), 17:0cy, 19:0cy, 15:1,

16:1ω7, 16:1ω5, 16:1, 17:1, 18:1ω7 and 18:1ω9) and fungal PLFA (18:2ω6,9) was

determined using a gas chromatograph coupled with a flame ionization detector (FID) and

isotope ratio mass spectrometer (IRMS) respectively (Karlowsky et al., 2018). Additionally,

the NLFA biomarker 16:1ω5 was used as the marker for AMF.

The amount of ^{13}C incorporated was determined from samples collected after the pulse

minus the isotope content of unlabeled soil collected before the pulse (Karlowsky et al.,

2018). The amounts of saprotrophic fungal markers (PLFA - 18:2ω6,9) were converted into

biomass C values by applying conversion factor 11.8 nmol = 1 mg C (Klamer, 2004), for

bacterial biomarkers by applying conversion factor 363.3 nmol = 1mg C (Frostegård and

Bååth, 1996) and for 16:1ω5 NLFA by applying conversion factor 1.047 nmol = 1μg C

(Williams et al., 2017). The amount of ^{13}C in the specific biomarker was determined from the

C pool size of the biomarker corrected for soil weight and moisture (Karlowsky et al., 2018).

The total amount of ^{13}C in the bacterial and fungal biomarkers was used as a proxy for

microbial biomass that incorporated freshly fixed plant C. Net ^{13}C assimilated by soil

microbial community and its parts (saprophytic fungi, AMF and free-living bacteria) for

every unit of ^{15}N gained by the plant was used as an estimate of net plant C allocation for

nutrient gain to specific microbial groups.

DNA and RNA based microbial community profiling

To ascertain the microbial community composition, 1.09 ± 0.01 g (n = 27) of soil was

used to co-extract total RNA and DNA using the MoBio RNA Powersoil kit. The samples

were bead-beaten for one cycle at 5.5 m/s for 30 s with a FastPrep instrument (MP

Biomedicals Germany GmbH). DNA was extracted using a MoBio RNA Powersoil DNA

elution accessory kit. Traces of DNA that was co-extracted in the final RNA elution was

removed using a MoBio DNase Max kit. The purified DNA and RNA concentrations were

measured using Qubit 4 fluorometer (Thermo Fisher Scientific) using the broad range assay

kits for DNA and RNA. Additionally, the purity of the samples was checked using a nanodrop

spectrometer 2000c (Thermo Fisher Scientific) and all samples were found to have A260/280

and A260/230 ratios of 2 suggesting no protein or organic compound contamination. The

integrity of the RNA samples was visually verified using 0.8 % agarose gel electrophoresis.

Presence of sharp well-defined ribosomal RNA bands and smear in between and above these

bands was considered as an indicator of high integrity of the RNA. 200 ng of total RNA

extracted was converted into cDNA using Superscript™ VILO kit (Thermo Fisher Scientific)

using random hexamers to ensure uniform conversion of all the RNA fragments. The DNA

and cDNA samples were further used for amplifying the 16S rRNA gene and ITS regions

respectively. Briefly, for the 16S rRNA gene sequencing, amplicon libraries were prepared

with V3-V5 primers (Haas et al., 2011) using PCR conditions described by Kozich et al.,

2013. ITS2 gene libraries were constructed by amplifying ITS region 2 using the fITS7

(forward) and ITS4 (reverse) primers (Ihrmark et al., 2012) with previously described PCR

conditions (Gweon et al., 2015). Both sets of primers also consisted of an appropriate

Illumina adapter, an 8-nt index sequence, a 10-nt pad sequence, a 2-nt linker (Kozich et al.,

2013). The amplicons were sequenced using Illumina MiSeq, v3 chemistry, 600 cycles.

Processing of sequencing data

For the bacterial community 16S rRNA gene sequencing data, the forward and reverse

reads obtained were merged using PEAR (v 0.9.6) with default parameters (Zhang et al.,

2014). The bacterial community composition was analyzed using the QIIME (1.8.0) pipeline

(Caporaso et al., 2010). Representative OTU's were picked using UCLUST using de novo

method at 97% similarity level and were assigned taxonomy using RDP classifier method

with the SILVA reference data set (release 132 for QIIME) (Edgar, 2010; Wang et al., 2007).

Total number of sequencing reads classified within the OTU table for the samples varied

between 4000 to 21982 reads. In order to make inter-sample comparisons OTU table obtained

was rarefied to 4000 sequences per sample using QIIME command single_rarefraction.py

(Caporaso et al., 2010).

Since fungal spores are ubiquitous, the ITS2 regions that were transcribed into RNA

were used to sample transcriptionally active fungi (Anderson and Parkin, 2007). The

amplified ITS region from total cDNA samples were used for sequencing. The amplified ITS

sequencing data was analyzed using the PIPITS pipeline using the default parameters (Gweon

et al., 2015). Briefly, the paired end sequences were merged using VSEARCH after which the

ITS2 region was extracted using ITSx after which they are matched to a trained UNITE

database at 97% similarity using RDP classifier to obtain an OTU table (Bengtsson-Palme et

al., 2013; Rognes et al., 2016). The OTUs detected were then assigned to fungal guilds using

FUNguild assignment app (http://www.stbates.org/guilds/app.php) (Nguyen et al., 2016).

The bacterial and fungal amplicon sequencing data and the associated metadata is

publicly available through the sequence read archive project number: SRP132171.

Statistical data analysis

One-way ANOVA was used to determine statistical differences amongst rhizosphere

types (AM vs NM) for root:shoot ratio and net microbial ^{13}C assimilation into total microbial

biomass. Two way ANOVA was applied to identify statistically significant differences due to

rhizosphere and litter types for ^{15}N assimilated by plant shoot and plant carbon allocation per

unit ^{15}N gain. All two-way ANOVA models that showed significant interactions of

rhizosphere-litter types were subject to Tukey's post-hoc test to identify differences amongst

litter types nested within rhizosphere types. Subsampled OTU tables that were binned to

different taxonomic groups at the phylum and class levels (bacteria) whereas fungal OTUs

were binned into guilds to compare effect of AM vs NM plant communities and litter

amendments on microbial (bacterial and fungal) community composition. The effects of litter

and rhizosphere type on microbial groups was separately determined by comparison of the F

values generated by a two-way ANOVA of their relative abundance. Differences between

microbial (bacterial and fungal) community composition affected by the experimental

treatment were determined using non-parametric PERMANOVA ("adonis" function in the

vegan package 2.5-2) (Oksanen et al., 2018). The dissimilarity based upon microbial

community composition due to litter and rhizosphere treatments were visualized as an NMDS

ordination of Bray-Curtis dissimilarity coefficients on a Hellinger transformed OTU table so

as to obtain the least amount of stress in the ordination (Ramette, 2007). All analysis was

performed using R software (v 3.4.3) with visualization being done with ggplot2 package

(Wickham, 2011).

Results

Carbon allocation and Nitrogen return

The shoot biomass of the AM plant community was smaller (4.66 ± 0.17 g, n = 18)

compared to NM the plant community (7.42 ± 0.44 g, n = 9) but both had similar root

biomass ($F_{1,25} = 0.32$; $p = 0.582$), which resulted in a higher root to shoot ratio for the AM

plant community. The AM plant community also allocated more carbon into their roots,

which was reflected in the root to shoot ratio of ^{13}C incorporated into the AM plant

community compared to the NM plant community ($F_{1,16} = 10.11$, $p = 0.005$; Figure 2A).

Unexpectedly, the net ^{13}C assimilation of the rhizospheric microbial communities from the

AM plant community was significantly lower than that of the NM plant community ($F_{1,16} =$

21.75, $p < 0.001$, Figure 2B).

The AM plant community gained 93.1 ± 4.7 µg ^{15}N from litter decomposition, which was not

significantly different from the NM plant community that gained 82.3 ± 10.7 µg ^{15}N (($F_{1,16} =$

0.36, $p = 0.558$, Figure 3A). Neither the different litter types (AM: $F_{2,15} = 0.31$, $p = 0.743$;

NM: $F_{2,6} = 0.99$, $p = 0.424$) nor the mesh sizes of the ingrowth cores (%N - $F_{1,16} = 1.19$, $p =$

0.290; ^{15}N - $F_{1,16}$ = 0.25, p = 0.623) had a significant influence on the ^{15}N return into both

plant communities. Consequently, the AM plant community invested 66% less ^{13}C into the

microbial community per ^{15}N gain than the NM plant community ($F_{1,16}$ = 14.92, p = 0.002;

Figure 3B). Moreover, the AM plant community allocated significantly more ^{13}C per ^{15}N

gained to AMF biomass ($F_{1,16}$ = 30.53, p < 0.001; Figure 4A) whereas NM plant communities

allocated significantly more ^{13}C per ^{15}N gained into saprophytic fungi ($F_{1,16}$ = 15.47, p =

0.001; Figure 4B) and bacteria ($F_{1,16}$ =4.78, p=0.044). Unexpectedly the amount of ^{13}C

allocated per ^{15}N gained depended only in the AM plant community on the type of

decomposed litter (litter – $F_{2,15}$ = 7.13, p = 0.009, Figure 4). Specifically, in the AM

rhizosphere, colonization of inorganic N amended soil resulted in significantly more plant-

derived ^{13}C to be allocated to bacteria and saprotrophic fungi compared to microbial

necromass and plant litter amended soil (Figure 4B, C). Such litter-based differences were

missing in the NM rhizosphere.

Microbial community composition

Most individual microbial groups were responsive to either rhizosphere type (NM or

AM) or litter type (root litter or microbial necromass or inorganic N) (Figure 5). Litter type

strongly affected bacterial OTUs binned under *Bacteroidetes*, *Gemmamonadetes*, *Nitrospira*

and *Deltaproteobacteria* whereas amongst fungal guilds saprotrophic and ectomycorrhizal

guilds responded to litter types (Figure 5A). The bacterial groups affected by rhizosphere type

belonged to the *Acidobacteria*, *Proteobacteria* and *Betaproteobacteria* whereas amongst the

fungal guilds AMF responded the most to rhizosphere type (Figure 5B). Such responsiveness

of specific microbial groups to rhizosphere and litter type significantly affected the microbial

community composition (Figure 6 & Table 1). Expectedly, the fungal communities varied

strongly along the expected lines between both plant communities (Figure 6A, Table 1). This

was mainly explained by the higher abundance of AMF (Figure 6C). Litter amendments also significantly affected the fungal community (Table 1). The bacterial community was also influenced by both rhizosphere and litter type (Figure 6D & 6E; Table 1). The rhizosphere of the NM plant community was dominated by phylum *Acidobacteria* whereas for AM plant communities were dominated by Proteobacterial classes (esp. *Betaproteobacteria*) (Figure 6F). Specifically, root litter amendments also resulted in dominance of phylum *Proteobacteria* whereas inorganic N amendments resulted in increased relative abundance of phylum *Bacteroidetes* (Figure 6F). OTUs grouped under *Deltaproteobacteria* were in general responsive to all litter amendments made in this study (Figure 6F).

Discussion

In this study we investigated how plant communities, differing in their association with AMF, affected the carbon allocation to the microbial communities and the return of mineralized nitrogen from different litter types. By combining a dual stable isotope labelling with a sequencing analyses and compound specific isotope analysis of the soil microbial community we found that AMF associated (AM) plant communities did not gain significantly more nitrogen from litter decomposition than non-mycorrhizal (NM) plant communities. However, AM plant communities had to invest less carbon into the soil microbial communities, resulting in a significantly lower carbon per nitrogen cost than NM plant communities (Figure 7). Even though AM plant communities foster a carbon demanding fungal network, they benefit strongly from the AMF-targeted access to the decomposition sites where plant-derived carbon is used to stimulate selected microorganisms for litter decomposition. These findings are in line with previous studies showing the importance of the AMF hyphal network on the associated soil bacterial community (Drigo et al., 2010; Vestergård et al., 2008). The close interaction of AMF and bacteria potentially increases decomposition efficiency and makes in the long run more nutrients available to the plant

communities. NM plant communities likely miss the targeted growth of AMF hyphae and

relay on untargeted saprophytic fungi and bacteria for decomposition, which increases the

overall carbon per nitrogen cost.

Furthermore, our results also showed that the net assimilation of plant carbon by the

rhizosphere microbial community depended on the biochemical characteristics of the litter

being decomposed. This suggests strongly that litter chemistry (litter type) is as important as

litter quality in the decomposition process and can modulate the extent of plant-derived

carbon required for decomposition (Jones et al., 2004). AMF-associated rhizospheric

microbial communities generally assimilated less plant carbon than the microbial

communities with NM plant communities. This suggests AMF allocated less labile C to the

free-living soil microbial community in order to take up mineralized N from decomposition.

The different net assimilation of the free-living microbial community from AM and NM plant

communities also suggests that the decomposition of different litter types strongly depended

on the association with AMF. This confirms that AMF indeed act as target oriented root

extensions that more effectively feed plant-derived carbon to the free-living microbial

community. Moreover, in response to the litter biochemistry, and thus the quantity of plant

available nutrients, plants and AMF likely control the amounts of exudates released. In turn,

the significant response of specific microbial groups to either the litter type and rhizosphere

type as well as the shift in the microbial community composition suggests that in addition to

the litter nutrient content (Stempfhuber et al., 2016) the interplay of litter type with plant

carbon inputs shapes microbial communities in the rhizosphere. Hence the composition of the

microbial decomposer community is adjusted to plant exudation and type of the litter to be

decomposed (Figure 7), making plant communities an important part of the decomposition

process (McLaren and Turkington, 2011). Expectedly, the fungal community primarily

differed between AM or NM plant communities, emphasizing the fungal role in

decomposition and nutrient gathering (Bukovská et al., 2018; Meier et al., 2015). However, in

the presence of AMF, we observed lower plant carbon assimilation by the free living

microbial community (Mueller et al., 2014), which was dominated by the bacterial

community. This suggests that primary decomposers in the AM rhizosphere were bacteria that

were fed and adapted to litter types by the AMF (Herman et al., 2012; Nuccio et al., 2013).

Most interestingly our results indicate for the first time that AM plant communities strongly

benefit from the association with AMF during microbial litter decomposition. AMF

assimilated more ^{13}C from plants but invested less carbon into the bacterial community for

decomposition. This might indicated that AMF, which have no capability to decomposed

lignocellulose structures of plant cell walls themselves (Tisserant et al., 2013), might be able

to decomposed peptidoglycan structures of microbial cell walls. This idea would be supported

by the dramatic up-regulation of genes encoding for enzymes containing a peptidoglycan

binding LysM domain and glycan hydrolases like lysozyme, glucosamidase and alpha

amylase that would be necessary for microbial cell wall decomposition (Tisserant et al., 2013;

Gude et al., 2012). If we consider that large parts of soil organic matter is microbial

necromass (Gleixner, 2013) the association of plants with AMF gain a new overall

importance for ecosystem functioning. However, additional experiments are needed to

confirm the new role of plant-AMF associations for microbial cell wall decomposition.

Conclusion
428

Our results suggest that three way interactions between the plant community, the

microbial community and the decomposing litter type have fundamental importance for

ecosystem functioning. Especially nutrient recycling in the plant-microorganism-soil

continuum is shaped by these interactions leading to a plant-driven and context-dependent

stimulation of decomposers. Their decomposition capabilities are likely highly redundant as

biochemically identical litter is decomposed by different parts of the microbial community. In

consequence, litter decomposition and nutrient recycling can only be fully understood if the

whole continuum is investigated. Moreover, our results demonstrate for the first time that

plant nutrient gain via symbiotic AMF is less carbon intensive and likely more targeted then

untargeted exudation od NM plant communities. This might explain why a large part of the

plant community relay on the AM nutrient uptake strategy.

Acknowledgements

SC received funding from long term DAAD scholarship to carry out the research. ML is funded by the German Research Foundation (DFG; FOR 456, FOR 1451 – "The Jena Experiment") and by the "Zwillenberg-Tietz Stiftung". We acknowledge help from Agnes Fastnacht with greenhouse resources and monitoring of the experiment. Special thanks to Karl Kübler for construction and deployment of the pulse labelling setup in the greenhouse. We acknowledge Heike Geilmann and Steffen Ruehlow for help with stable isotope measurements, and Maria Foerster for helping with fatty acid analysis We also thank Erika Kothe, Ruchira Mukherji, Elisa Catao and Huei Ying Gan for helpful comments and discussions.

Author contributions

SC, GG and AM designed the study. SC and JH performed the experiment. SC performed the extractions and stable isotope analyses. TG and RG completed the downstream processing of nucleic acids, sequencing and supervised bioinformatics analysis. SC and ML completed the statistical analysis. SC, ML and GG conceptualized the work and wrote the manuscript and all authors contributed to revisions.

Conflict of Interest

The authors declare no conflict of interest

References:

Anderson, I. C., and Parkin, P. I. (2007). Detection of active soil fungi by RT-PCR

amplification of precursor rRNA molecules. *J. Microbiol. Methods* 68, 248–253.

doi:10.1016/j.mimet.2006.08.005.

Bengtsson-Palme, J., Ryberg, M., Hartmann, M., Branco, S., Wang, Z., Godhe, A., et al.

(2013). Improved software detection and extraction of ITS1 and ITS2 from ribosomal

ITS sequences of fungi and other eukaryotes for analysis of environmental sequencing

data. *Methods Ecol. Evol.* 4, 914–919. doi:10.1111/2041-210X.12073.

Bitterlich, M., Franken, P., and Graefe, J. (2018). Arbuscular Mycorrhiza Improves Substrate

Hydraulic Conductivity in the Plant Available Moisture Range Under Root Growth

Exclusion. *Front. Plant Sci.* 9, 301. doi:10.3389/fpls.2018.00301.

Bligh, E. G., and Dyer, W. J. (1959). A rapid method of total lipid extraction and purification.

Can. J. Biochem. Physiol. 37, 911–917. doi:10.1139/y59-099.

Bonfante, P., and Genre, A. (2010). Mechanisms underlying beneficial plant-fungus

interactions in mycorrhizal symbiosis. *Nat. Commun.* 1, 48. doi:10.1038/ncomms1046.

Bowles, T. M., Jackson, L. E., and Cavagnaro, T. R. (2018). Mycorrhizal fungi enhance plant

nutrient acquisition and modulate nitrogen loss with variable water regimes. *Glob.*

Chang. Biol. 24, e171–e182. doi:10.1111/gcb.13884.

Brundrett, M. . (2015a). Arbuscular Mycorrhizas. In: Mycorrhizal Associations: The Web

Resource. Version 2.0. 24.04.2015.‹mycorrhizas.info›. Available at:

http://mycorrhizas.info/vam.html.

Brundrett, M. . (2015b). Mycorrhizal Associations: The Web Resource. Version 2.0.

24.04.2015. ‹mycorrhizas.info›. Available at: http://mycorrhizas.info/nmplants.html.

Bukovská, P., Bonkowski, M., Konvalinková, T., Beskid, O., Hujslová, M., Püschel, D., et al.

(2018). Utilization of organic nitrogen by arbuscular mycorrhizal fungi—is there a

specific role for protists and ammonia oxidizers? *Mycorrhiza* 28, 269–283.

doi:10.1007/s00572-018-0825-0.

Bunn, R. A., Simpson, D. T., Bullington, L. S., Lekberg, Y., and Janos, D. P. (2019).

Revisiting the 'direct mineral cycling' hypothesis: arbuscular mycorrhizal fungi colonize

leaf litter, but why? *ISME J.*, 1. doi:10.1038/s41396-019-0403-2.

Caporaso, J. G., Kuczynski, J., Stombaugh, J., Bittinger, K., Bushman, F. D., Costello, E. K.,

et al. (2010). QIIME allows analysis of high-throughput community sequencing data.

Nat. Methods 7, 335–336. doi:10.1038/nmeth.f.303.

Cavagnaro, T. R., Barrios-Masias, F. H., and Jackson, L. E. (2012). Arbuscular mycorrhizas

and their role in plant growth, nitrogen interception and soil gas efflux in an organic

production system. *Plant Soil* 353, 181–194. doi:10.1007/s11104-011-1021-6.

Cheng, L., Booker, F. L., Tu, C., Burkey, K. O., Zhou, L., Shew, H. D., et al. (2012).

Arbuscular Mycorrhizal Fungi Increase Organic Carbon Decomposition Under Elevated

CO_2. *Science* 337, 1084–1087. doi:10.1126/science.1224304.

Craine, J. M., Morrow, C., and Fierer, N. (2007). Microbial nitrogen limitation increases

decomposition. *Ecology* 88, 2105–2113. doi:10.1890/06-1847.1.

Davison, J., Moora, M., Opik, M., Adholeya, A., Ainsaar, L., Ba, A., et al. (2015). Global

assessment of arbuscular mycorrhizal fungus diversity reveals very low endemism.

Science 349, 970–973. doi:10.1126/science.aab1161.

de Graaff, M.-A., Classen, A. T., Castro, H. F., and Schadt, C. W. (2010). Labile soil carbon

inputs mediate the soil microbial community composition and plant residue

decomposition rates. *New Phytol.* 188, 1055–1064. doi:10.1111/j.1469-

8137.2010.03427.x.

Dijkstra, F. A., Carrillo, Y., Pendall, E., and Morgan, J. A. (2013). Rhizosphere priming: a

nutrient perspective. *Front. Microbiol.* 4, 216. doi:10.3389/fmicb.2013.00216.

Drigo, B., Pijl, A. S., Duyts, H., Kielak, A. M., Gamper, H. A., Houtekamer, M. J., et al.

(2010). Shifting carbon flow from roots into associated microbial communities in

response to elevated atmospheric CO_2. *Proc. Natl. Acad. Sci. U. S. A.* 107, 10938–42.

doi:10.1073/pnas.0912421107.

Edgar, R. C. (2010). Search and clustering orders of magnitude faster than BLAST.

Bioinformatics 26, 2460–2461. doi:10.1093/bioinformatics/btq461.

Frostegård, A., and Bååth, E. (1996). The use of phospholipid fatty acid analysis to estimate

bacterial and fungal biomass in soil. *Biol. Fertil. Soils* 22, 59–65.

doi:10.1007/s003740050076.

Gleixner, G. (2013). Soil organic matter dynamics: a biological perspective derived from the

use of compound-specific isotopes studies. *Ecol. Res.* 28, 683–695. doi:10.1007/s11284-

012-1022-9.

Gude, A., Kandeler, E. and Gleixner, G. (2012). Input related microbial carbon dynamic of

soil organic matter in particle size fractions. *Soil Biol. Biochem.* 47, 209-219.

Güsewell, S., and Gessner, M. O. (2009). N : P ratios influence litter decomposition and

colonization by fungi and bacteria in microcosms. *Funct. Ecol.* 23, 211–219.

doi:10.1111/j.1365-2435.2008.01478.x.

Gweon, H. S., Oliver, A., Taylor, J., Booth, T., Gibbs, M., Read, D. S., et al. (2015). PIPITS:

An automated pipeline for analyses of fungal internal transcribed spacer sequences from

the Illumina sequencing platform. *Methods Ecol. Evol.* 6, 973–980. doi:10.1111/2041-

210X.12399.

Haas, B. J., Gevers, D., Earl, A. M., Feldgarden, M., Ward, D. V, Giannoukos, G., et al.

(2011). Chimeric 16S rRNA sequence formation and detection in Sanger and 454-

pyrosequenced PCR amplicons. *Genome Res.* 21, 494–504. doi:10.1101/gr.112730.110.

Herman, D. J., Firestone, M. K., Nuccio, E., and Hodge, A. (2012). Interactions between an

arbuscular mycorrhizal fungus and a soil microbial community mediating litter

decomposition. *FEMS Microbiol. Ecol.* 80, 236–247. doi:10.1111/j.1574-

6941.2011.01292.x.

Hütsch, B. W., Augustin, J., and Merbach, W. (2002). Plant rhizodeposition — an important

source for carbon turnover in soils. *J. Plant Nutr. Soil Sci.* 165, 397. doi:10.1002/1522-

2624(200208)165:4<397::AID-JPLN397>3.0.CO;2-C.

Ihrmark, K., Bödeker, I. T. M., Cruz-Martinez, K., Friberg, H., Kubartova, A., Schenck, J., et

al. (2012). New primers to amplify the fungal ITS2 region - evaluation by 454-

sequencing of artificial and natural communities. *FEMS Microbiol. Ecol.* 82, 666–677.

doi:10.1111/j.1574-6941.2012.01437.x.

Jacoby, R., Peukert, M., Succurro, A., Koprivova, A., and Kopriva, S. (2017). The Role of

Soil Microorganisms in Plant Mineral Nutrition-Current Knowledge and Future

Directions. *Front. Plant Sci.* 8, 1617. doi:10.3389/fpls.2017.01617.

Jones, D. L., Hodge, A., and Kuzyakov, Y. (2004). Plant and mycorrhizal regulation of

rhizodeposition. *New Phytol.* 163, 459–480. doi:10.1111/j.1469-8137.2004.01130.x.

Karlowsky, S., Augusti, A., Ingrisch, J., Hasibeder, R., Lange, M., Lavorel, S., et al. (2018).

Land use in mountain grasslands alters drought response and recovery of carbon

allocation and plant-microbial interactions. *J. Ecol.* 106, 1230–1243. doi:10.1111/1365-

2745.12910.

Klamer, M. (2004). Estimation of conversion factors for fungal biomass determination in

compost using ergosterol and PLFA 18:2ω6,9. *Soil Biol. Biochem.* 36, 57–65.

doi:10.1016/j.soilbio.2003.08.019.

Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K., and Schloss, P. D. (2013).

Development of a dual-index sequencing strategy and curation pipeline for analyzing

amplicon sequence data on the miseq illumina sequencing platform. *Appl. Environ.*

Microbiol. 79, 5112–5120. doi:10.1128/AEM.01043-13.

Kuzyakov, Y., and Xu, X. (2013). Competition between roots and microorganisms for

nitrogen: mechanisms and ecological relevance. *New Phytol.* 198, 656–69.

doi:10.1111/nph.12235.

Levin, P. A., and Angert, E. R. (2015). Small but mighty: Cell size and bacteria. *Cold Spring*

Harb. Perspect. Biol. 7, 1–11. doi:10.1101/cshperspect.a019216.

Liang, C., Schimel, J. P., and Jastrow, J. D. (2017). The importance of anabolism in microbial

control over soil carbon storage. *Nat. Microbiol.* 2, 17105.

doi:10.1038/nmicrobiol.2017.105.

López-Mondéjar, R., Zühlke, D., Becher, D., Riedel, K., and Baldrian, P. (2016). Cellulose

and hemicellulose decomposition by forest soil bacteria proceeds by the action of

structurally variable enzymatic systems. *Sci. Rep.* 6, 25279. doi:10.1038/srep25279.

Louca, S., Polz, M. F., Mazel, F., Albright, M. B. N., Huber, J. A., O'Connor, M. I., et al.

(2018). Function and functional redundancy in microbial systems. *Nat. Ecol. Evol.* 2,

936–943. doi:10.1038/s41559-018-0519-1.

Malik, A. A., Dannert, H., Griffiths, R. I., Thomson, B. C., and Gleixner, G. (2015).

Rhizosphere bacterial carbon turnover is higher in nucleic acids than membrane lipids:

implications for understanding soil carbon cycling. *Front. Microbiol.* 6, 268.

doi:10.3389/fmicb.2015.00268.

McLaren, J. R., and Turkington, R. (2011). Plant identity influences decomposition through

more than one mechanism. *PLoS One* 6, e23702. doi:10.1371/journal.pone.0023702.

Meier, I. C., Pritchard, S. G., Brzostek, E. R., McCormack, M. L., and Phillips, R. P. (2015).

The rhizosphere and hyphosphere differ in their impacts on carbon and nitrogen cycling

in forests exposed to elevated CO_2. *New Phytol.* 205, 1164–74. doi:10.1111/nph.13122.

Mueller, R. C., Balasch, M. M., and Kuske, C. R. (2014). Contrasting soil fungal community

responses to experimental nitrogen addition using the large subunit rRNA taxonomic

marker and cellobiohydrolase I functional marker. *Mol. Ecol.* 23, 4406–17.

doi:10.1111/mec.12858.

Nguyen, N. H., Song, Z., Bates, S. T., Branco, S., Tedersoo, L., Menke, J., et al. (2016).

FUNGuild: An open annotation tool for parsing fungal community datasets by ecological

guild. *Fungal Ecol.* 20, 241–248. doi:10.1016/J.FUNECO.2015.06.006.

Nottingham, A. T., Turner, B. L., Winter, K., Chamberlain, P. M., Stott, A., and Tanner, E. V.

J. (2013). Root and arbuscular mycorrhizal mycelial interactions with soil

microorganisms in lowland tropical forest. *FEMS Microbiol. Ecol.* 85, 37–50.

doi:10.1111/1574-6941.12096.

Nuccio, E. E., Hodge, A., Pett-Ridge, J., Herman, D. J., Weber, P. K., and Firestone, M. K.

(2013). An arbuscular mycorrhizal fungus significantly modifies the soil bacterial

community and nitrogen cycling during litter decomposition. *Environ. Microbiol.* 15,

1870–1881. doi:10.1111/1462-2920.12081.

Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., et al. (2018).

vegan: Community Ecology Package. Available at: https://cran.r-

project.org/package=vegan.

Ramette, A. (2007). Multivariate analyses in microbial ecology. *FEMS Microbiol. Ecol.* 62,

142–60. doi:10.1111/j.1574-6941.2007.00375.x.

Rognes, T., Flouri, T., Nichols, B., Quince, C., and Mahé, F. (2016). VSEARCH: a versatile

open source tool for metagenomics. *PeerJ* 4, e2584. doi:10.7717/peerj.2584.

Roscher, C., Schumacher, J., Baade, J., Wilcke, W., Gleixner, G., Weisser, W. W., et al.

(2004). The role of biodiversity for element cycling and trophic interactions: an

experimental approach in a grassland community. *Basic Appl. Ecol.* 5, 107–121.

doi:10.1078/1439-1791-00216.

Sanders, F. E., and Tinker, P. B. (1973). Phosphate flow into mycorrhizal roots. *Pestic. Sci.* 4,

385–395. doi:10.1002/ps.2780040316.

Scheffknecht, S., St-Arnaud, M., Khaosaad, T., Steinkellner, S., and Vierheilig, H. (2007). An

altered root exudation pattern through mycorrhization affecting microconidia

germination of the highly specialized tomato pathogen Fusarium oxysporum f. sp.

lycopersici (Fol) is not tomato specific but also occurs in Fol nonhost plants. *Can. J.*

Bot. 85, 347–352. doi:10.1139/B07-015.

Schimel, J. P., and Bennett, J. (2004). Nitrogen mineralization: Challenges of a changing

paradigm. *Ecology* 85, 591–602. doi:10.1890/03-8002.

Shi, S., Nuccio, E. E., Shi, Z. J., He, Z., Zhou, J., and Firestone, M. K. (2016). The

interconnected rhizosphere: High network complexity dominates rhizosphere

assemblages. *Ecol. Lett.* 19, 926–936. doi:10.1111/ele.12630.

Stempfhuber, B., Richter-Heitmann, T., Regan, K. M., Kölbl, A., Wüst, P. K., Marhan, S., et

al. (2016). Spatial interaction of archaeal ammonia-oxidizers and nitrite-oxidizing

bacteria in an unfertilized grassland soil. *Front. Microbiol.* 6, 1567.

doi:10.3389/fmicb.2015.01567.

Strickland, M. S., Osburn, E., Lauber, C., Fierer, N., and Bradford, M. A. (2009). Litter

quality is in the eye of the beholder: initial decomposition rates as a function of inoculum

characteristics. *Funct. Ecol.* 23, 627–636. doi:10.1111/j.1365-2435.2008.01515.x.

Thirkell, T. J., Cameron, D. D., and Hodge, A. (2016). Resolving the "nitrogen paradox" of

arbuscular mycorrhizas: fertilization with organic matter brings considerable benefits for

plant nutrition and growth. *Plant. Cell Environ.* 39, 1683–90. doi:10.1111/pce.12667.

Thomson, B. C., Ostle, N. J., McNamara, N. P., Oakley, S., Whiteley, A. S., Bailey, M. J., et

al. (2013). Plant soil interactions alter carbon cycling in an upland grassland soil. *Front.*

Microbiol. 4, 253. doi:10.3389/fmicb.2013.00253.

Throckmorton, H. M., Bird, J. a., Dane, L., Firestone, M. K., and Horwath, W. R. (2012). The

source of microbial C has little impact on soil organic matter stabilisation in forest

ecosystems. *Ecol. Lett.* 15, 1257–1265. doi:10.1111/j.1461-0248.2012.01848.x.

Thuille, A., Laufer, J., Höhl, C., and Gleixner, G. (2015). Carbon quality affects the nitrogen

partitioning between plants and soil microorganisms. *Soil Biol. Biochem.* 81, 266–274.

doi:10.1016/J.SOILBIO.2014.11.024.

Tisserant, E., Malbreil, M., Kuo, A., Kohler, A., Symeonidi, A., Balestrini, R., et al. (2013).

Genome of an arbuscular mycorrhizal fungus provides insight into the oldest plant

symbiosis. *Proc. Natl. Acad. Sci.* 110, 20117–20122. (Tisserant et al., 2013).

Veresoglou, S. D., Chen, B., and Rillig, M. C. (2012). Arbuscular mycorrhiza and soil

nitrogen cycling. *Soil Biol. Biochem.* 46, 53–62. doi:10.1016/J.SOILBIO.2011.11.018.

Vestergård, M., Henry, F., Rangel-Castro, J. I., Michelsen, A., Prosser, J. I., and Christensen,

S. (2008). Rhizosphere bacterial community composition responds to arbuscular

mycorrhiza, but not to reductions in microbial activity induced by foliar cutting. *FEMS*

Microbiol. Ecol. 64, 78–89. doi:10.1111/j.1574-6941.2008.00447.x.

Wang, Q., Garrity, G. M., Tiedje, J. M., and Cole, J. R. (2007). Naive Bayesian classifier for

rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl. Environ.*

Microbiol. 73, 5261–7. doi:10.1128/AEM.00062-07.

Wickham, H. (2011). ggplot2. *Wiley Interdiscip. Rev. Comput. Stat.* 3, 180–185.

doi:10.1002/wics.147.

Williams, A., Manoharan, L., Rosenstock, N. P., Olsson, P. A., and Hedlund, K. (2017).

Long-term agricultural fertilization alters arbuscular mycorrhizal fungal community

composition and barley (*Hordeum vulgare*) mycorrhizal carbon and phosphorus

exchange. *New Phytol.* 213, 874–885. doi:10.1111/nph.14196.

Zhalnina, K., Louie, K. B., Hao, Z., Mansoori, N., da Rocha, U. N., Shi, S., et al. (2018).

Dynamic root exudate chemistry and microbial substrate preferences drive patterns in

rhizosphere microbial community assembly. *Nat. Microbiol.* 3, 470–480.

doi:10.1038/s41564-018-0129-3.

Zhang, J., Kobert, K., Flouri, T., and Stamatakis, A. (2014). PEAR: a fast and accurate

Illumina Paired-End reAd mergeR. *Bioinformatics* 30, 614–20.

doi:10.1093/bioinformatics/btt593.

Zhu, Q., Riley, W. J., and Tang, J. (2017). A new theory of plant-microbe nutrient

competition resolves inconsistencies between observations and model predictions. *Ecol.*

Appl. 27, 875–886. doi:10.1002/eap.1490.

Figures and Tables Legends

 Figure 1: Experimental setup of the study.

Figure 2: Photosynthetically fixed carbon assimilated by plants and associated microbial

community. A: Comparison of ratio of root to shoot ^{13}C content between AM (n = 9) and

NM (n = 9) the plant community types. Height of the bar represents the mean and error bars

represent standard error of mean. Different letters on bars designate significance differences.

B: Mean ^{13}C (ng) excess in soil microbial biomass of each plant community types. Colors

represent plant community types where Dark green=AM plant community and Orange= NM

plant community. Bars sharing the same letter are not significantly different.

Figure 3: Carbon investment into microbial community and return of decomposition released nitrogen. A: ^{15}N content (µg) of shoot biomass compared between AM (n = 9) and NM (n = 9) plant community (AM – Dark green and NM - Orange). Horizontal bars represent means for each group and error bars represent standard error of mean (n=9 per rhizosphere). Groups sharing the same letters are not significantly different. **B:** Comparison of carbon investment between AM (Dark green) and NM (Orange) plant communities represented as µg of freshly fixed carbon (^{13}C) incorporated into total microbial biomass per µg ^{15}N gained by plants with NM and AM plant community (n = 9 per community).

Figure 4: ^{13}C investment for ^{15}N gain by plants partitioned based on microbial groups

Partioning of plant ^{13}C investment into microbial biomass ($^{13}C_{mic}$) per ^{15}N gained by into key

microbial groups like AMF (A), free-living fungi (B), and free-living bacteria (C) in

rhizospheres of AM and NM plant communities amended with the three litter types (Plant

Root Litter, Microbial Necromass and Inorganic N). Significant differences between the

rhizosphere types are depicted with asterisks when $p < 0.05$ (*) , $p < 0.01$ (**) and $p<0.001$

(***) . Different letters within the bar is shows significant differences between litter types

amended in that rhizosphere.

Figure 5: Effects of rhizosphere type and litter type on specific microbial groups.

Comparison of effect size (F values from two-way ANOVA) for relative abundance of

individual microbial groups in response to experimental treatments (Litter (A) and

Rhizopshere (B) types). Filled bars represent significant effect of a treatment on the relative

abundance of the microbial group. Relative abundances of bacterial and fungal groups can be

found in supplementary Figures S1, S2 and S3. Microbial group abbreviations – Nitro

(*Nitrospira*), Gemma (*Gemmamonadetes*), Actino (*Actinobacteria*), Chlor (*Chloroflexi*),

DeltaP (*Deltaproteobacteria*), BetaP (*Betaproteobacteria*), Proteo (*Proteobacteria*), AlphaP

(*Alphaproteobacteria*), GammaP (*Gammaproteobacteria*), Acido (*Acidobacteria*), Verruco

(*Verrucomicrobia*), Bac (*Bacteroidetes*).

Figure 6: Variation in microbial bacterial and fungal community composition

Non-metric multidimensional scaling (NMDS) plots displaying Bray-Curtis distances

amongst microbial community composition under different combinations of rhizosphere types

(panel A&D) and litter types (panel B&E). Colors represent rhizosphere types where Dark

green= AM plant community and Orange= NM plant communtiy. Symbols represent litter

type being used with plant root litter (circle), Microbial necromass (square) and Inorganic N

(triangle). Microbial group abbreviations – Nitro (*Nitrospira*), Gemma (*Gemmamonadetes*),

Actino (*Actinobacteria*), Chlor (*Chloroflexi*), DeltaP (*Deltaproteobacteria*), BetaP

(*Betaproteobacteria*), Proteo (*Proteobacteria*), AlphaP (*Alphaproteobacteria*), GammaP

(*Gammaproteobacteria*), Acido (*Acidobacteria*), Verruco (*Verrucomicrobia*), Bac

(*Bacteroidetes*). Stress values for fungal (A, B &C) and bacterial (D, E &F) plots are 0.05 and

0.06 respectively.

Figure 7 – Conceptual model showing key differences between plants associated with AMF

and not associated with AMF. Arrows on each side represent the C and N flow differences

arising due to presence or absence of AMF. The microbial community influenced by plant

fresh C input and litter inputs are represented as sectors. Microbial groups that are specified in

the key are responsive to either litter or plant inputs.

Table 1: Permanova analysis of bacterial and fungal communities of the microcosms

Microbial Community	Treatment	Df	F.value	R^2	p-value
Bacterial	**Rhizosphere Type**	**2**	**3.68**	**0.19**	**0.01**
	Litter type	**2**	**3.83**	**0.2**	**0.01**
Fungal	**Rhizosphere Type**	**2**	**16.22**	**0.57**	**0.00**
	Litter type	**2**	**3.40**	**0.12**	**0.05**

Rows in bold face represent significant effects. Interaction terms (Rhizosphere Type x Litter type) were not significant.