Original article

Reassembly of soil fungal communities under reforestation and herbivore exclusion

Shinichi Tatsumi[,2,]*, Shunsuke Matsuoka[3,]*, Saori Fujii[4], Kobayashi Makoto[5], Takashi Osono[6], Forest Isbell[7], Akira S. Mori[8]

* These authors contributed equally to this study.

Department of Biological Sciences, University of Toronto Scarborough, Toronto, ON, Canada

Hokkaido Research Center, Forest and Forestry Products Research Institute, Hokkaido, Japan

Graduate School of Simulation Studies, University of Hyogo, Hyogo, Japan

Forest and Forestry Products Research Institute, Ibaraki, Japan

Field Science Center for Northern Biosphere, Hokkaido University, Hokkaido, Japan

Department of Environmental Systems Science, Doshisha University, Kyoto, Japan

Department of Ecology, Evolution and Behavior, University of Minnesota, St Paul, MN, USA

Graduate School of Environment and Information Sciences, Yokohama National University, Kanagawa, Japan

Corresponding author: S. Tatsumi

Department of Biological Sciences, University of Toronto Scarborough, 1265 Military Trail, Toronto, ON M1C 1A4, Canada

Email: community.ecologist@gmail.com, Tel: +1-416-208-5130

Abstract

Soil fungi can help improve ecosystem restoration, yet our understanding of how fungi reassemble

in degraded land is limited. Here, we studied fungal community structure using DNA

metabarcoding in reforested sites following agricultural abandonment and overgrazing. We used a

natural experiment in which reforestation with different numbers of tree species and deer exclusion

have been applied for multiple decades. We found that local fungal richness (alpha diversity) was

1.9 to 2.9 times greater in reforested stands than in natural forests and total fungal richness (gamma

diversity) was 1.3 to 1.9 times greater. These results were regardless of the number of tree species

planted in the reforested stands. Conversely, reforested stands had a homogenized community

structure with relatively lower degrees of compositional dissimilarity among sites within each

stand (beta diversity). These findings were attributable to lower environmental heterogeneity,

stronger dispersal limitation, and a comparatively shorter time since the onset of community

assembly in reforested stands. Deer exclosures had no detectable effect on fungal community

structure. Overall, the agricultural legacy in fungal community structure appears to have persisted

for decades, even under proactive restoration of aboveground vegetation. Direct human

intervention belowground may therefore be necessary for the recovery of soil biota once

homogenized, which may facilitate ecosystem restoration.

Keywords: Biodiversity, Biotic homogenization, Community assembly, DNA metabarcoding,

Ecosystem restoration, Soil microbes

Introduction

Given the extent of the human-induced damage to the biosphere, ecosystem restoration has become

increasingly relevant in today's world (Hobbs and Harris, 2001). While restoration ecology has

typically focused on aboveground vegetation (Brudvig 2011; Perring et al., 2015; Young 2000),

less attention has been paid to belowground biota (Kardol and Wardle 2010). The diversity and

composition of soil organisms, particularly fungi, can be largely altered in degraded land due to

the decreased availability of organic substrates and their symbionts (Kardol and Wardle, 2010).

Understanding how soil fungi respond to restoration is of ecosystem-level significance, because

they can regulate key belowground functions and create feedbacks that support the aboveground

recovery (Young et al. 2005).

Recently, there are increasing social concerns on reforestation using mixed-species

plantations over monocultures (Verheyen et al., 2016; Tatsumi 2020). Plant diversity has often

shown to have positive effects on soil fungal diversity in grassland experiments (Milcu et al., 2013;

Scherber et al., 2010). In forests, however, experimental tests on the relationships between tree and

fungal diversity remain scarce (Weißbecker et al., 2018). Particularly, compared with local fungal

richness (alpha diversity), we still know little about the spatial variation in fungal composition

(beta diversity), as influenced by tree-species mixing. Assessing both alpha and beta diversity can

inform ecosystem recovery at local scales and larger scales that are relevant to land management

and policy making (e.g., forest stands, landscapes).

Human-induced increases in herbivore density can drive ecosystem degradation and

hamper vegetative recovery (Opperman and Merenlender, 2000). The changes in vegetation caused

by herbivory are known to subsequently impact soil organisms, including fungi (Kardol et al.,

2014). Thus far, such indirect effects have typically been tested by sole manipulations of herbivores

(Bardgett and Wardle, 2003; Kardol et al., 2014). It is also possible, however, for the impacts of

herbivores to be mediated by plant diversity. Specifically, the rates of herbivory can be reduced in

plant mixtures compared to monocultures because of associational protection among neighbouring

species (Cook-Patton et al., 2014). There has yet been little experiment on whether herbivores and

plant diversity have individual or interactive effects on soil fungi.

In this study, we quantified the effects of aboveground forest restoration on soil fungal

communities. As a study system, we selected a restoration site in northern Japan. We used a natural

experiment in which the following treatments were applied in a fully-crossed design: 'tree planting

with different numbers of species' and 'deer exclusion' (Fujii et al., 2017; Mori et al., 2016). Our

specific objectives were to test (1) whether the community structure of soil fungi — namely, their

alpha and beta diversity — differ among monocultures and mixtures as well as nearby grasslands

and natural forests, and (ii) whether tree diversity and deer herbivory have individual or interactive

effects on soil fungi. Addressing these questions provides a step towards ecosystem restoration

grounded on above- and belowground linkages.

Methods

Study site

We conducted a natural experiment in lowland coastal areas of the Shiretoko National Park,

northern Japan (44°08'–11' N, 145°03'–08' E, elevation 140–220 m). The area has been designated

a World Heritage Site by the United Nations Educational, Scientific and Cultural Organization

(UNESCO) on account of it being home to one of the most species rich, northern temperate

ecosystems in the world (http://whc.unesco.org/en/list/1193). The mean monthly temperature

ranges from −6.1°C in February to 20.8°C in August. The mean annual precipitation is 1149 mm.

According to the soil classification system of Japan (Obara et al., 2011) and the Japan soil

inventory (https://soil-inventory.dc.affrc.go.jp/), the soil type in our study sites are low-humic

allophanic Andosols, which corresponds to Typic Hapludands and Hydric Hapludands in the

USDA soil taxonomy (Soil Survey Staff, 2010) (Supplementary Fig. S1).

Approximately 90% of the park's terrestrial area is covered by pristine natural vegetation,

most of which is composed of mixed conifer–broadleaf forests. Parts of the remaining area had

been used for agriculture from the early twentieth century until the Government of Japan ordered

the settlers to abandon the land, a process that was completed by the late 1960s. Since then, a

number of reforestation initiatives has been conducted in the area to restore the arable land to

mixed conifer–broadleaf forests (>861 ha; the Shiretoko National Trust Movement Area). Such

activities included reforestation with different numbers of tree species and the establishment of

fences to prevent overgrazing and browsing by sika deer (*Cervus nippon yesoensis*). Deer density

has increased rapidly from the late 1980s to the late 1990s in the park, with a current density of

6.1–13.6 individuals km^{-2}. Currently, the landscape is composed of mosaics of multiple vegetation

types, including monoculture and mixture stands, grasslands, and natural forests.

Study design and sampling

The design and sampling methods are explained in detail in Mori et al. (2016) and Fujii et al.

(2017). Although we used the same study setting and a partly overlapping dataset with Mori et al.

(2016) and Fujii et al. (2017), this study completely differs from them. Specifically, while the

previous studies investigated the effects of fungal richness on ecosystem functioning, we instead

focused on the fungal community structure under different restoration treatments.

We used a natural experiment with a 4 × 2 factorial design — namely, four habitat types

and the inside/outside of deer exclosures. The four habitat types were (i) monoculture stands

reforested with *Larix kaempferi*, (ii) mixture stands reforested with *Abies sachalinensis*, *Picea*

glehnii, and *Betula ermanii*, (iii) grasslands dominated by a dwarf bamboo species (*Sasa cernua*)

as a negative control group (i.e., the initial state of restoration), and (iv) mixed conifer–broadleaf

natural forests dominated by *A. sachalinensis*, *Quercus crispula*, and *Kalopanax septemlobus* as a

positive control group (i.e., the reference state of restoration).

The deer-exclusion and control sites were established adjacent to each other (i.e., inside

and outside of deer fences) in each habitat type. Each site was ~1 ha in size. All the sites were

distributed within an area of 2 km × 5 km (Supplementary Fig. S1). Given the restricted number

of reforested stands and deer fences in the region, our experiment consisted of eight sites with one

replication for each factorial combination. Reforested stands and deer fences were respectively

established >30 years and ca. 10 years prior to our field sample collection (Fujii et al., 2017). The

present vegetation inside and outside deer exclosures show significant structural differences (Fujii

et al., 2017; Nishizawa et al., 2016) (Supplementary Fig. S2).

The collection of soil samples and chemical analyses were conducted as previously

described (Fujii et al., 2017; Mori et al., 2016). Briefly, in May 2013 we established three 10 m ×

10 m plots in each of the eight factorial combinations. In each plot, we randomly selected three

points and collected topsoil from a 0–5 cm depth at each point (i.e., totalling 72 soil samples). Soil

samples were transported from the field on ice and kept at −20°C until further analysis. Soil water

content, pH, total carbon (C) and nitrogen (N) content, and inorganic N (ammonium and nitrate)

content were measured to characterize soil properties. Total C and N content were measured using

an organic elemental analyzer (Macro Coder JM1000CN, J-Science Lab Co., Ltd., Kyoto, Japan).

Ammonium and nitrate were extracted from soil using a 2-M KCL solution and then measured

with an auto-analyzer (AACS-4, BL-TEC Co., Ltd., Osaka, Japan).

Molecular analyses and bioinformatics

Molecular analysis and bioinformatics were conducted as previously described (Fujii et al., 2017;
Mori et al., 2016) and are explained in detail in Matsuoka et al. (2016b). Briefly, total DNA was
extracted from each of the 72 soil samples (0.25 g sample^{-1}) using the Soil DNA Isolation kit
(Norgen Biotek Corp., Thorold, ON, Canada). A semi-nested polymerase chain reaction (PCR)
was then performed to amplify the nuclear internal transcribed spacer 1 region. The pooled
amplicons were sequenced with a GS Junior sequencer (454 Life Sciences, Branford, CT, USA).

The reads were clustered with a cut-off sequence similarity of 97% (Osono, 2014) using
the Minimus genome assembler (Sommer et al., 2007). Consensus sequences were used as
molecular operational taxonomic units (OTUs). A total of 389 OTUs were obtained. For each OTU,
taxonomic identification was conducted using the QCauto method implemented in Claident
(Tanabe and Toju, 2013). Hereafter, we refer to 'OTUs' as 'species' for simplicity, bearing in mind
that OTUs defined by a fixed sequence similarity do not necessarily represent species in a
biological sense. The functional group of each species was determined based on the FUNGuild
database (Nguyen et al., 2016) and an intensive literature review. See Supplementary Materials for
details of molecular analyses and bioinformatics. Raw sequence data files are available at the DNA
Data Bank of Japan (DRA003024).

Community structure analyses

We defined fungal alpha diversity as the number of species within a community (i.e., a soil sample)
and beta diversity as the extent of community dissimilarity within a treatment combination. We
tested the effects of habitat types, deer fences, and their interactions on fungal alpha diversity using
two-way analysis of variance (ANOVA) and Tukey's HSD test with 'plot' as a random effect.
Variation among samples in sequencing depths (i.e., read counts), which can bias alpha diversity
estimations, was standardized by rarefying read numbers. We used two rarefaction methods,
namely sample size-based and coverage-based rarefactions (Chao and Jost, 2012), to confirm the
robustness of results. In addition to habitat types, deer fences, and their interactions, we also
included soil properties (pH, total C, total N, C:N ratio, inorganic N, and water content) as
explanatory variables in separate ANOVA and linear regression analyses to account for the
potential confounding effects these variables could have on alpha diversity. The effects of habitat
types, deer fences, and their interactions on soil properties were tested by two-way ANOVA and

Tukey's HSD tests.

To detect the types of rank-frequency distribution of the species in each treatment

combination, four models — preemption, log-normal, Zipf, and null models (Wilson, 1991) —

were fitted to the distribution of occurrence by species rank (i.e., the number samples in which

species occurred vs. species rank). The best model was selected based on the Akaike information

criterion (AIC).

The effects of habitat types, deer fences, and their interactions on fungal community

composition were tested using two-way permutational multivariate analysis of variance

(PERMANOVA). The compositional dissimilarity between each treatment pair was tested using

pairwise PERMANOVA where *P*-values were adjusted using the Bonferroni correction method.

We compared beta diversity among the treatment combinations using the homogeneity of

multivariate dispersions test (Anderson, 2006). The communities were ordinated using nonmetric

multidimensional scaling (NMDS). The effects of soil properties on fungal community

composition were tested by fitting their vectors onto the NMDS ordination. For all the above

community dissimilarity analyses, we used two dissimilarity measurements — the Jaccard

(Jaccard, 1912) and Raup–Crick indices (Raup and Crick, 1979) — to confirm the robustness of

results. We used the indices based on presence/absence information in order to minimize the

possible influence of read-count biases resulting from interspecific variation in the number of

ribosomal DNA tandem repeats and from PCR processes (Toju, 2015). The habitat preferences of

each fungal species and functional group were tested based on the association between their

occurrence patterns and treatment combinations (De Cáceres et al., 2010). All statistical analyses

were implemented in R 3.5.1 (R Core Team, 2018). The R packages used are listed in

Supplementary Table S1.

169   Results

Fungal alpha diversity was 1.9 to 2.9 times greater in grasslands, monocultures, and mixtures than

in natural forests (Fig. 1). It was confirmed that the sample size- and coverage-based rarefactions,

as well as the case without rarefaction, yielded qualitatively consistent results; therefore, the

results of the last two are provided in Supplementary Fig. S3. The effect of deer exclosures on

fungal alpha diversity was not significant (Figs. 1, S3). The interactions between habitat types and

deer exclosures also had no detectable effect on alpha diversity (Figs. 1, S3). Soil properties (i.e.,

Figure 1. Effects of habitat type and deer fence on rarefied fungal operational taxonomic unit (OTU) richness (i.e., alpha diversity). Results from two-way ANOVA is shown in the upper right; *** $P < 0.001$; n.s. $P \geq 0.05$. Different letters indicate significant differences ($P < 0.05$) among treatments (Tukey's test). Error bars indicate standard errors.

pH, total C, total N, C:N ratio, inorganic N, and water content) varied among the treatment combinations (Fig. S4). Habitat type was the most significant predictor of fungal alpha diversity even when we included the soil properties as explanatory variables in ANOVA (Table S2). Regression analyses which controlled for the effect of soil pH (which was found to be a significant variable in ANOVA) also showed that alpha diversity was lower in natural forests than in grasslands (Table S3; Fig. S5).

The frequency of occurrence against species rank decreased less steeply in grasslands, monocultures, and mixtures than in natural forests (Fig. 2). Based on AIC, the preemption model was selected as the best model for habitats other than natural forests. The Zipf model was selected for natural forests. Gamma diversity (i.e., the total number of species present in each treatment combination) was 1.3 to 1.9 times greater in grasslands, monocultures, and mixtures (>130 species) than in natural forests (≤100 species) (Fig. 2).

Fungal community composition differed significantly among the habitat types (Fig. 3a, b). We confirmed that the Jaccard and Raup–Crick indices qualitatively yielded the same results; therefore, results from the Raup–Crick index are provided in Supplementary Fig. S6. The effect of deer exclosures on community composition was not significant (Figs. 3a, S6a). The interaction

Figure 2. Relationships between occurrence frequencies of fungal operational taxonomic units (OTUs) and their rank. The y axes indicate the number of times each OTU occurred out of the nine replications in each treatment. The OTUs are arranged in decreasing order of occurrence frequency. Note that the maximum number of rank (on the x axis) in each panel equals the total number of OTUs observed in each treatment (i.e., gamma diversity). The numbers next to the model names indicate the Akaike information criterion (AIC). Curves show the selected model (shown in boldface), based on AIC, which was fitted to the distribution.

between reforestation types and deer exclosures also had no significant effect on community composition. The soil properties were significantly correlated to community composition (Figs. 3a, S6a). Increases in inorganic N and water content were mainly associated with shifts in fungal composition among habitat types (i.e., arrows roughly paralleled shift direction), whereas soil pH and C:N ratio were associated with community dissimilarity within each habitat type (Figs. 3a, S6a). Fungal beta diversity was significantly higher in natural forests compared to the other three habitat types (Figs. 3c, S6c).

Most fungal species and functional groups occurred more frequently in certain habitat types (Fig. 4). For example, ectomycorrhizal fungi, ericoid mycorrhizal fungi, and other symbionts that coexist with Ericaceae species occurred more frequently in monocultures and mixtures than in the other two habitat types. Conversely, the majority of fungal species and functional groups

Figure 3. Dissimilarity of fungal communities within and among treatments. (a) Ordination of communities based on nonmetric multidimensional scaling (NMDS) and the effects of treatments (habitat type and deer fence) on community composition tested by two-way permutational multivariate analysis of variance (PERMANOVA). Community dissimilarity was measured using the Jaccard index. Arrows show the associations of soil properties with community composition. (b) Community dissimilarity between pairs of vegetation types. (c) Among-vegetation differences in the size of within-vegetation community dissimilarity (i.e., beta diversity) tested by the permutation test of homogeneity of multivariate dispersion (PERMDISP).

occurred at approximately the same frequency both inside and outside deer fences (Fig. 4). Exceptions were *Mortierella* sp. 1 and coprophilous fungi that more often occurred outside than inside fenced exclosures.

Discussion

Effects of tree diversity and herbivores on soil fungi

Tree planting and herbivore exclusion are among the globally conducted approaches in terrestrial restoration (Kardol and Wardle, 2010; Verheyen et al., 2016). In this study, we tested the effects of these restoration practices on soil fungal communities using a natural experiment. Most notably, we found that local fungal richness (alpha diversity) was 1.9 to 2.9 times greater in grasslands and monoculture and mixture stands than in natural forests (Fig. 1). This result was contrary to the often reported, positive association between aboveground and belowground diversity (Peay et al.,

Figure 4. Differences in occurrence frequency of fungal operational taxonomic units (OTUs) and functional groups among restoration treatments. OTUs and functional groups with adequate sample sizes (OUTs occurring at >25 sampling points out of 72 points and functional groups having >3 OTUs) are presented. The transverse lines in boxes indicate treatments in which a given species or a functional group occurred significantly more often than in other treatments.

2013; Prober et al., 2015). We also found that fungal beta diversity and rank-frequency distribution in reforested stands were more similar to those of grasslands compared to natural forests, regardless of the number of tree species planted (Figs. 2, 3). The total fungal richness (gamma diversity) was 1.3 to 1.9 time greater in grasslands and reforested stands than in natural forests (Fig. 2). The reforested stands shared more indicator species and functional groups with grasslands than with natural forests (Fig. 4). These results stand in contrast with the fact that aboveground

vegetation in our restoration sites is steadily recovering (Nishizawa et al. 2016, Fujii et al. 2017;
Fig. S2). Previous studies on naturally regenerating forests showed that historical effects of past
land-use activity (e.g., farming, logging) on soil fungal communities can last for decades (Bachelot
et al., 2016; Hartmann et al., 2012). Our results further suggest that, even after decades of proactive
aboveground restoration, the way soil fungi assemble in reforested stands still differs from how
communities are structured in natural forests.

Deer exclusion had no detectable effect on soil fungal communities at the species level
(Figs. 1, 2, 3a). Nevertheless, at the level of functional groups, coprophilous fungi occurred more
frequently outside fenced exclosures than inside (Fig. 4), which is attributable to the input of deer
faeces. Coprophilous fungi were also found most often in grasslands (Fig. 4), coinciding with the
habitat preference of sika deer for grasslands over forests (Yabe, 1995). Such deer-induced changes
in the functional composition of fungi may, in turn, affect ecosystem nutrient cycling (Kardol et
al., 2014). In fact, a faeces decomposition experiment conducted at our study area (Yabe, 1995)
revealed more rapid faeces decomposition (and thus nutrient release) in grasslands than forests.
This indicates that, in our grassland sites, a positive feedback exists between faeces production,
fungal activity, and the growth of plants on which deer feed (van der Wal et al., 2004). Conversely,
we found no evidence to support the interactive effects of deer and tree diversity on soil fungi
(Figs. 1, 3a). A previous study (Cook-Patton et al., 2014) found a diversity-derived reduction in
seedling herbivory by white-tailed deer (*Odocoileus virginianus*) owing to the associational
protection of palatable species by unpalatable species. We did not observe such associational
effects presumably because sika deer unselectively feed on a wide variety of plants including
unpalatable species when its population density is high (Takahashi and Kaji, 2001). Overall, the
results indicate that aboveground plant–herbivore interactions in our experimental site have limited
impact on the species diversity of soil fungi, yet have potential to affect their functional
composition and ecosystem functioning.

Fungal community assembly
Understanding the ecological processes underlying diversity patterns can provide a critical step
towards the development of theory-driven restoration (Mori et al., 2017). Here, based on a
conceptual synthesis in community ecology (Vellend, 2010), we discuss three potential assembly
processes by which the lower fungal alpha diversity and higher beta diversity in natural forests

than reforested stands (Figs. 1, 3a, 3c) can arise — that is, ecological selection across space and
time, and dispersal limitation.

286   The first possibility is the selection under different degrees of spatial environmental
heterogeneity. Increasing environmental heterogeneity can increase beta diversity (i.e., the
dissimilarity among local communities within each treatment combination) through the selection
of different species at different sites (Mouquet and Loreau, 2003). The high fungal beta diversity
we found in natural forests (Fig. 3a, c) is attributable to the fact that natural forests often have
higher habitat heterogeneity compared to grasslands and plantations (Mori, 2011). The greater
variation in water content and inorganic N of natural forests compared to grassland and reforested
stands (Fig. 3a) also coincides with previous findings that soil properties are often homogenized
in ex-arable land (Bachelot et al., 2016; Fraterrigo et al., 2005). Moreover, it is often expected that
environmental heterogeneity increases alpha diversity (i.e., local species richness) via source-sink
effects (Pulliam, 1988). Nonetheless, a theory by Kadmon and Allouche (2007) suggests that
environmental heterogeneity can reduce the available habitat size of each species and, thus, could
conversely decrease overall species richness. According to this theory, the low alpha diversity in
natural forests (Fig. 1) could have derived from the high environmental heterogeneity that brought
some species to such low abundances that they go locally extinct.

301   The second possibility is the time dependency in community response to environmental
conditions. Constrained rates of mortality and reproduction prevent species from going extinct
immediately, even when their population growth rates are negative. The higher alpha and lower
beta diversity in reforested stands compared to natural forests (Figs. 1, 3a, 3c) can be interpreted
simply that the process of local species selection has not yet been completed in reforested stands,
regardless of environmental variation *per se*. Moreover, the long duration of time in natural forests
could have allowed the effect of species' arrival order to be amplified and thereby the communities
to diverge (i.e., priority effects; Fukami, 2015). In fact, a multi-year monitoring study of fungal
communities (Matsuoka et al. 2016a) found that compositional similarities among communities
were largely explained by the closeness in the time of survey. This suggests that the elapsed time
after the onset of community assembly can have a major control on fungal diversity patterns
observed in the field. Furthermore, species occurrence patterns in grasslands and reforested stands
exhibited the preemption (geometric) distribution (Fig. 2). This type of distribution is often found
in the early stage of succession for various taxa, including fungi (Visser, 1995), plants (Whittaker,

1965), and soil invertebrates (Caruso and Migliorini, 2006). We note that caution is needed in

comparing our study to previous studies because we used rank-frequency distributions instead of

rank-abundance distributions due to the methodological concerns in high-throughput DNA data

analyses (Toju, 2015). Nonetheless, we do offer a potential interpretation that fungal species in our

reforested sites are still in the course of ecological selection even after decades of tree planting.

The third possibility is dispersal limitation. Despite the small size and immense number

of propagules, dispersal limitation is now acknowledged as a crucial determinant of fungal

community structure (Peay et al., 2010). Fungal diversity patterns in our experiment (Figs. 1, 3a,

3c) can be explained by dispersal limitation at two conceptual scales: external dispersal from

outside the set of local communities (as assumed in the mainland-island model) and internal

dispersal among the local communities (as described in the metacommunity model) (Fukami,

2015). In the former scenario, the low beta diversity of grasslands and reforested stands (Fig. 3a,

3c) can be explained by limitations in the external dispersal of habitat specialists, which may have

led the communities to become composed of a same suite of generalists, irrespective of local

habitat conditions (Vellend et al., 2007). Indeed, a study that compared fungal community structure

in primary forests and artificial pastures (Mueller et al., 2016) reported an increase in dominance

of generalist fungi in pastures. Under the latter metacommunity scenario, maximal alpha and beta

diversity is predicted to occur at intermediate and low levels of internal dispersal, respectively

(Mouquet and Loreau 2003). This indicates that the low alpha and high beta diversity of natural

forests (Figs. 1, 3a, 3c) resulted from limited within-treatment dispersal. This view is further

supported by the presence of competition-dispersal tradeoffs among fungal species (Peay et al.,

2007; Smith et al., 2018); that is, only species with high competitive but low dispersal abilities

were able to subsist in natural forests.

Future challenges

We found clear differences in fungal community structure among restoration treatments, but

uncertainty still persists regarding causal relationships among some variables. Specially, given the

lack of site-level replication for each treatment in our experiment, we cannot explicitly separate

the effects of human activities (i.e., restoration treatments and past agricultural practices) from the

among-site environmental variation that might have existed from before (e.g., soil types). For

example, we found that the C:N ratio was relatively low in the natural forests (Fig. S3), similarly

to what was found for fungal richness (Fig. 1); this similarity, however, can be due to either human
activities or the original environment of the sites. Future studies with multiple site replications are
thus required in order to draw conclusions about causalities among the variables (e.g., potential
reductions in species richness caused by increased N availability; Cline et al., 2018).

Nevertheless, some of our results did, in fact, indicate that the fungal community structure
was driven by the human activities in the study sites. Most notably, adding the measured soil
properties as explanatory variables to the statistical models did not alter our result that fungal
richness differs among habitat types (Table S2). Additional analyses showed that natural forests
had the lowest fungal richness even when we controlled for the soil pH which negatively affected
the richness (Table S3; Fig. S5). The fact that all the study sites have the same soil type (low-humic
allophanic Andosols; Fig. S1) further supports the possibility that the soil environment was
homogeneous across the sites prior to the treatments and land-use changes. Moving forward,
accumulations of additional field data will allow us to explicitly disentangle the relationships
among site conditions, human activities, and fungal diversity.

Implications for restoration

In this study, we investigated soil fungal communities in a restoration landscape and found that
aboveground-oriented restoration treatments (i.e., tree planting and herbivore exclusion) do not
necessarily translate into the recovery of fungal diversity. Our results suggest that in order to
enhance the recovery of soil fungi, direct intervention to the soil, in conjunction with the
application of vegetative treatments, may be necessary. For example, supplying organic substrates
(e.g., deadwoods) to the soil surface can help create additional habitats for soil fungi (Mäkipää et
al., 2017). Especially in restored sites with homogenized community structure like ours (Fig. 3a,
3c), creating mosaics of habitat patches by supplemental substrates could increase environmental
heterogeneity and thus beta diversity. Another commonly applied approach in soil restoration is
fungal inoculation, although caution is needed because fungal inoculation can occasionally cause
negative impacts on the ecosystem (Janoušková et al., 2013). In fact, in our study site, natural
forests had the lowest fungal alpha diversity (Fig. 1), indicating that simply adding multiple species
to degraded land may not necessarily shift the community structure to that of natural forests. Rather,
it might be effective to selectively inoculate a small number of species that could otherwise not
reach the sites, considering the limited colonization of habitat specialists in our restored sites (as

indicated by low beta diversity; Fig. 3a, 3c). Furthermore, inoculating fungal species in different
order at different locations can increase beta diversity via priority effect. We believe that such
direct treatments to the soil and belowground biota will allow us to better enhance the recovery of
degraded ecosystems.

Acknowledgements
We would like to thank the Shiretoko Nature Foundation for providing logistical support for our
field work, and the members of the Shiretoko Biodiversity Evaluation Project for assistance in the
field and laboratory work. We are also grateful to Marc W. Cadotte for thoughtful feedbacks, Jia
Pu for her support in statistical analysis, Keiichi Okada for his advice on fungal guilds, and Bill
Liu for English language editing. S.T. was supported by the Grant-in-Aids for JSPS Fellows PD
(No. 15J10614) and the Young Scientists B (No. 16K18715) from the Japan Society for the
Promotion of Science (JSPS). S.M. was supported by the Grant-in-Aids for Young Scientists B
(No. 17K15199) from the JSPS. A.S.M. was supported by the Mitsui & Co., Ltd. Environment
Fund and the Pro Natura Foundation Japan.

References
Anderson, M.J., 2006. Distance-based tests for homogeneity of multivariate dispersions.
Biometrics 62, 245–253. doi:10.1111/j.1541-0420.2005.00440.x
Bachelot, B., Uriarte, M., Zimmerman, J.K., Thompson, J., Leff, J.W., Asiah, A., Koshner, J.,
McGuire, K., 2016. Long-lasting effects of land use history on soil fungal communities in
second-growth tropical rain forests. Ecol. Appl. 26, 1881–1895. doi:10.1890/15-1397.1
Bardgett, R.D., Wardle, D.A., 2003. Herbivore-mediated linkages between aboveground and
belowground communities. Ecology 84, 2258–2268. doi:10.1890/02-0274
Brudvig, L.A., 2011. The restoration of biodiversity: Where has research been and where does it
need to go? Am. J. Bot. 98, 549–558. doi:10.3732/ajb.1000285
Caruso, T., Migliorini, M., 2006. A new formulation of the geometric series with applications to
oribatid (Acari, Oribatida) species assemblages from human-disturbed Mediterranean areas.
Ecol. Modell. 195, 402–406. doi:10.1016/j.ecolmodel.2005.11.042
Chao, A., Jost, L., 2012. Coverage-based rarefaction and extrapolation: Standardizing samples by
completeness rather than size. Ecology 93, 2533–2547. doi:10.1890/11-1952.1

Chase, J.M., 2003. Community assembly: When should history matter? Oecologia 136, 489–498.
doi:10.1007/s00442-003-1311-7

Cline, L.C., Huggins, J.A., Hobbie, S.E., Kennedy, P.G., 2018. Organic nitrogen addition
suppresses fungal richness and alters community composition in temperate forest soils. Soil
Biol. Biochem. 125, 222–230. doi:10.1016/j.soilbio.2018.07.008

Cook-Patton, S.C., LaForgia, M., Parker, J.D., 2014. Positive interactions between herbivores and
plant diversity shape forest regeneration. Proc. R. Soc. B Biol. Sci. 281, 20140261.
doi:10.1098/rspb.2014.0261

De Cáceres, M., Legendre, P., Moretti, M., 2010. Improving indicator species analysis by
combining groups of sites. Oikos 119, 1674–1684. doi:10.1111/j.1600-0706.2010.18334.x

Fraterrigo, J.M., Turner, M.G., Pearson, S.M., Dixon, P., 2005. Effects of past land use on spatial
heterogeneity of soil nutrients in southern Appalachian forests. Ecol. Monogr. 75, 215–230.
doi:10.1890/03-0475

Fujii, S., Mori, A.S., Koide, D., Makoto, K., Matsuoka, S., Osono, T., Isbell, F., 2017.
Disentangling relationships between plant diversity and decomposition processes under forest
restoration. J. Appl. Ecol. 54, 80–90. doi:10.1111/1365-2664.12733

Fukami, T., 2015. Historical contingency in community assembly: Integrating niches, species
pools, and priority effects. Annu. Rev. Ecol. Evol. Syst. 46, 1–23. doi:10.1146/annurev-
ecolsys-110411-160340

Hartmann, M., Howes, C.G., VanInsberghe, D., Yu, H., Bachar, D., Christen, R., Henrik Nilsson,
R., Hallam, S.J., Mohn, W.W., 2012. Significant and persistent impact of timber harvesting
on soil microbial communities in Northern coniferous forests. ISME J. 6, 2199–2218.
doi:10.1038/ismej.2012.84

Hobbs, R.J., Harris, J. a, 2001. Restoration ecology: Repairing the Earth's ecosystems in the new
millennium. Restor. Ecol. 9, 239–246. doi:10.1046/j.1526-100x.2001.009002239.x

Jaccard, P., 1912. The distribution of the flora in the alpine zone. New Phytol. 11, 37–50.
doi:10.1111/j.1469-8137.1912.tb05611.x

Janoušková, M., Krak, K., Wagg, C., Štorchová, H., Caklová, P., Vosátka, M., 2013. Effects of
inoculum additions in the presence of a preestablished arbuscular mycorrhizal fungal
community. Appl. Environ. Microbiol. 79, 6507–6515. doi:10.1128/AEM.02135-13

Kadmon, R., Allouche, O., 2007. Integrating the effects of area, isolation, and habitat heterogeneity

on species diversity: A unification of island biogeography and niche theory. Am. Nat. 170,

443–54. doi:10.1086/519853

Kardol, P., Dickie, I.A., St. John, M.G., Husheer, S.W., Bonner, K.I., Bellingham, P.J., Wardle,

D.A., 2014. Soil-mediated effects of invasive ungulates on native tree seedlings. J. Ecol. 102,

622–631. doi:10.1111/1365-2745.12234

Kardol, P., Wardle, D.A., 2010. How understanding aboveground-belowground linkages can assist

restoration ecology. Trends Ecol. Evol. 25, 670–679. doi:10.1016/j.tree.2010.09.001

Mäkipää, R., Rajala, T., Schigel, D., Rinne, K.T., Pennanen, T., Abrego, N., Ovaskainen, O., 2017.

Interactions between soil- and dead wood-inhabiting fungal communities during the decay of

Norway spruce logs. ISME J. 11, 1964–1974. doi:10.1038/ismej.2017.57

Matsuoka, S., Kawaguchi, E., Osono, T., 2016. Temporal distance decay of similarity of

ectomycorrhizal fungal community composition in a subtropical evergreen forest in Japan.

FEMS Microbiol. Ecol. 92, fiw061. doi:10.1093/femsec/fiw061

Milcu, A., Allan, E., Roscher, C., Jenkins, T., Meyer, S.T., Flynn, D., Bessler, H., Buscot, F., Engels,

C., Gubsch, M., König, S., Lipowsky, A., Loranger, J., Renker, C., Scherber, C., Schmid, B.,

Thébault, E., Wubet, T., Weisser, W.W., Scheu, S., Eisenhauer, N., 2013. Functionally and

phylogenetically diverse plant communities key to soil biota. Ecology 94, 1878–1885.

doi:10.1890/12-1936.1

Mori, A.S., 2011. Ecosystem management based on natural disturbances: Hierarchical context and

non-equilibrium paradigm. J. Appl. Ecol. 48, 280–292. doi:10.1111/j.1365-

2664.2010.01956.x

Mori, A.S., Isbell, F., Fujii, S., Makoto, K., Matsuoka, S., Osono, T., 2016. Low multifunctional

redundancy of soil fungal diversity at multiple scales. Ecol. Lett. 19, 249–259.

doi:10.1111/ele.12560

Mori, A.S., Lertzman, K.P., Gustafsson, L., 2017. Biodiversity and ecosystem services in forest

ecosystems: A research agenda for applied forest ecology. J. Appl. Ecol. 54, 12–27.

doi:10.1111/1365-2664.12669

Mouquet, N., Loreau, M., 2003. Community patterns in source-sink metacommunities. Am. Nat.

162, 544–557. doi:10.1086/378857

Mueller, R.C., Rodrigues, J.L.M., Nüsslein, K., Bohannan, B.J.M., 2016. Land use change in the

Amazon rain forest favours generalist fungi. Funct. Ecol. 30, 1845–1853. doi:10.1111/1365-

2435.12651

Nguyen, N.H., Song, Z., Bates, S.T., Branco, S., Tedersoo, L., Menke, J., Schilling, J.S., Kennedy, P.G., 2016. FUNGuild: An open annotation tool for parsing fungal community datasets by ecological guild. Fungal Ecol. 20, 241–248. doi:10.1016/j.funeco.2015.06.006

Nishizawa, K., Tatsumi, S., Kitagawa, R., Mori, A.S., 2016. Deer herbivory affects the functional diversity of forest floor plants via changes in competition-mediated assembly rules. Ecol. Res. 31, 569–578. doi:10.1007/s11284-016-1367-6

Obara, H., Ohkura, T., Tanaka, Y., Kohyama, K., Maejima, Y., Hamazaki, T., 2011. Comprehensive soil classification system of Japan first approximation. Bull. Natl. Inst. Agro-Environ. Sci. 29, 1–73.

Opperman, J.J., Merenlender, A.M., 2000. Deer herbivory as an ecological constraint to restoration of degraded riparian corridors. Restor. Ecol. 8, 41–47. doi:10.1046/j.1526-100x.2000.80006.x

Osono, T., 2014. Metagenomic approach yields insights into fungal diversity and functioning, in: Sota, T., Kagata, H., Ando, Y., Utsumi, S., Osono, T. (Eds.), Species diversity and community structure. Springer, pp. 1–23. doi:10.1007/978-4-431-54261-2_1

Peay, K.G., Baraloto, C., Fine, P.V., 2013. Strong coupling of plant and fungal community structure across western Amazonian rainforests. ISME J. 7, 1852–1861. doi:10.1038/ismej.2013.66

Peay, K.G., Bruns, T.D., Kennedy, P.G., Bergemann, S.E., Garbelotto, M., 2007. A strong species-area relationship for eukaryotic soil microbes: Island size matters for ectomycorrhizal fungi. Ecol. Lett. 10, 470–480. doi:10.1111/j.1461-0248.2007.01035.x

Peay, K.G., Garbelotto, M., Bruns, T.D., 2010. Evidence of dispersal limitation in soil microorganisms: Isolation reduces species richness on mycorrhizal tree islands. Ecology 91, 3631–3640. doi:10.1890/09-2237.1

Perring, M.P., Standish, R.J., Price, J.N., Craig, M.D., Erickson, T.E., Ruthrof, K.X., Whiteley, A.S., Valentine, L.E., Hobbs, R.J., 2015. Advances in restoration ecology: Rising to the challenges of the coming decades. Ecosphere 6, art131. doi:10.1890/ES15-00121.1

Prober, S.M., Leff, J.W., Bates, S.T., Borer, E.T., Firn, J., Harpole, W.S., Lind, E.M., Seabloom, E.W., Adler, P.B., Bakker, J.D., Cleland, E.E., Decrappeo, N.M., Delorenze, E., Hagenah, N., Hautier, Y., Hofmockel, K.S., Kirkman, K.P., Knops, J.M.H., La Pierre, K.J., Macdougall, A.S., Mcculley, R.L., Mitchell, C.E., Risch, A.C., Schuetz, M., Stevens, C.J., Williams, R.J.,

Fierer, N., 2015. Plant diversity predicts beta but not alpha diversity of soil microbes across grasslands worldwide. Ecol. Lett. 18, 85‑95. doi:10.1111/ele.12381

Pulliam, H.R., 1988. Sources, sinks, and population regulation. Am. Nat. 132, 652–661. doi:10.1086/284880

R Core Team, 2018. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.

Raup, D.M., Crick, R.E., 1979. Measurement of faunal similarity in paleontology. J. Paleontol. 53, 1213–1227. doi:10.2307/1304099

Scherber, C., Eisenhauer, N., Weisser, W.W., Schmid, B., Voigt, W., Fischer, M., Schulze, E.-D., Roscher, C., Weigelt, A., Allan, E., Beßler, H., Bonkowski, M., Buchmann, N., Buscot, F., Clement, L.W., Ebeling, A., Engels, C., Halle, S., Kertscher, I., Klein, A.-M., Koller, R., König, S., Kowalski, E., Kummer, V., Kuu, A., Lange, M., Lauterbach, D., Middelhoff, C., Migunova, V.D., Milcu, A., Müller, R., Partsch, S., Petermann, J.S., Renker, C., Rottstock, T., Sabais, A., Scheu, S., Schumacher, J., Temperton, V.M., Tscharntke, T., 2010. Bottom-up effects of plant diversity on multitrophic interactions in a biodiversity experiment. Nature 468, 553–556. doi:10.1038/nature09492

Smith, G.R., Steidinger, B.S., Bruns, T.D., Peay, K.G., 2018. Competition–colonization tradeoffs structure fungal diversity. ISME J. 12, 1758–1767. doi:10.1038/s41396-018-0086-0

Soil Survey Staff, 2010. Keys to soil taxonomy: Eleventh edition. USDA, NRCS, pp. 338, Washington, DC

Sommer, D.D., Delcher, A.L., Salzberg, S.L., Pop, M., 2007. Minimus: A fast, lightweight genome assembler. BMC Bioinformatics 8, 64. doi:10.1186/1471-2105-8-64

Takahashi, H., Kaji, K., 2001. Fallen leaves and unpalatable plants as alternative foods for sika deer under food limitation. Ecol. Res. 16, 257–262. doi:10.1046/j.1440-1703.2001.00391.x

Tanabe, A.S., Toju, H., 2013. Two new computational methods for universal DNA barcoding: A benchmark using barcode sequences of bacteria, archaea, animals, fungi, and land plants. PLoS One 8, e76910. doi:10.1371/journal.pone.0076910

Tatsumi, S., 2020. Tree diversity effects on forest productivity increase through time because of spatial partitioning. For. Ecosyst. 24, 7. doi:10.1186/s40663-020-00238-z

Toju, H., 2015. High-throughput DNA barcoding for ecological network studies. Popul. Ecol. 57, 37–51. doi:10.1007/s10144-014-0472-z

van der Wal, R., Bardgett, R.D., Harrison, K. a, Stien, A., 2004. Vertebrate herbivores and
ecosystem control: Cascading effects of faeces on tundra ecosystems. Ecography 27, 242–
252. doi:10.1111/j.0906-7590.2004.03688.x

Vellend, M., 2010. Conceptual synthesis in community ecology. Q. Rev. Biol. 85, 183–206.
doi:10.1086/652373

Vellend, M., Verheyen, K., Flinn, K.M., Jacquemyn, H., Kolb, A., Van Calster, H., Peterken, G.,
Graae, B.J., Bellemare, J., Honnay, O., Brunet, J., Wulf, M., Gerhardt, F., Hermy, M., 2007.
Homogenization of forest plant communities and weakening of species-environment
relationships via agricultural land use. J. Ecol. 95, 565–573. doi:10.1111/j.1365-
2745.2007.01233.x

Verheyen, K., Vanhellemont, M., Auge, H., Baeten, L., Baraloto, C., Barsoum, N., Bilodeau-
Gauthier, S., Bruelheide, H., Castagneyrol, B., Godbold, D., Haase, J., Hector, A., Jactel, H.,
Koricheva, J., Loreau, M., Mereu, S., Messier, C., Muys, B., Nolet, P., Paquette, A., Parker,
J., Perring, M., Ponette, Q., Potvin, C., Reich, P., Smith, A., Weih, M., Scherer-Lorenzen, M.,
2016. Contributions of a global network of tree diversity experiments to sustainable forest
plantations. Ambio 45, 29–41. doi:10.1007/s13280-015-0685-1

Visser, S., 1995. Ectomycorrhizal fungal succession in jack pine stands following wildfire. New
Phytol. 129, 389–401. doi:10.1111/j.1469-8137.1995.tb04309.x

Weißbecker, C., Wubet, T., Lentendu, G., Kühn, P., Scholten, T., Bruelheide, H., Buscot, F., 2018.
Experimental evidence of functional group-dependent effects of tree diversity on soil fungi
in subtropical forests. Front. Microbiol. 9, 1–16. doi:10.3389/fmicb.2018.02312

Whittaker, R.H., 1965. Dominance and diversity in land plant communities: Numerical relations
of species express the importance of competition in community function and evolution.
Science. 147, 250–260. doi:10.1126/science.147.3655.250

Wilson, J.B., 1991. Methods for fitting dominance/diversity curves. J. Veg. Sci. 2, 35–46.
doi:10.2307/3235896

Yabe, T., 1995. A fundamental study on habitat management for wildlife: Habitat use of sika deer
and a change in the vegetation on Shiretoko peninsula, Hokkaido. Res. Bull. Hokkaido Univ.
For. 52, 115–180.

Young, T.P., 2000. Restoration ecology and conservation biology. Biol. Conserv. 92, 73–83.
doi:10.1016/S0006-3207(99)00057-9

Young, T.P., Petersen, D.A., Clary, J.J., 2005. The ecology of restoration: Historical links,

emerging issues and unexplored realms. Ecol. Lett. 8, 662–673. doi:10.1111/j.1461-

0248.2005.00764.x