**Title:** Microniche sampling of the microbiome in the Starlet Sea Anemone, *Nematostella*

*vectensis*, reveals a compartment-specific dominance of Spirochetes

**Authors:** A. M. Bonacolta[1], M. T. Connelly[1], S. Rosales[2,3], J. del Campo[1], N. Traylor-

Knowles[1,*]

[1]: Rosenstiel School of Marine and Atmospheric Science, Department of Marine Biology and

Ecology, University of Miami, 4600 Rickenbacker Causeway, Florida 33149, USA

[2]: Atlantic Oceanographic and Meteorological Laboratory, National Oceanographic and

Atmospheric Administration, Miami, Florida, USA

[3]: Cooperative Institute for Marine and Atmospheric Studies, University of Miami, Miami,

Florida, USA

*: Corresponding author: Nikki Traylor-Knowles, ngt14@miami.edu

**Abstract**

Microniche sampling can reveal highly specialized bacterial associations within the holobiont and also facilitate identification of core microbial symbionts that would otherwise be overlooked by bulk sampling methods. We characterized compartment-specific associations present within the model cnidarian *Nematostella vectensis* by dividing its morphology into three distinct body regions. This sampling design allowed us to uncover a compartment-specific dominance of spirochetes within the capitulum of *N. vectensis*. Bacteria from the Spirochaetaceae family made up 65.8% of the community in the capitulum, while only representing 1.2% and 0.1% of the communities in the mesenteries and physa, respectively. The capitulum community also exhibited significantly lower alpha-diversity compared to the other two compartments. A phylogenetic analysis of the predominate spirochete sequence recovered from *N. vectensis* showed a close relation to spirochetes previously recovered from wild *N. vectensis*. These sequences clustered closer to the recently described *Oceanispirochaeta*, rather than *Spirochaeta perfilievii*, supporting them as members of this genus. This suggests a consistent and potentially important association between *N. vectensis* and spirochetes from the Spirochaetales family.

**Introduction**

The discovery of microniche specific microbial communities has become important for our understanding of microbial interactions and diversity (Ainsworth, Thurber and Gates 2010; Bourne, Morrow and Webster 2016). Microbial niches have been identified in many different organisms from an ecosystem scale, all the way down to the cellular level (Aagaard *et al.* 2014; Foster *et al.* 2017; Ricci *et al.* 2019). For example, in the Giant Clam, *Tridacna maxima*,

different tissues harbor distinct microbial communities, with the gills harboring a high

prevalence of *Endozoicomonas* bacteria. Compartmental evaluation of *T. maxima* helped uncover

*Endozoicomonas*'s role in nitrogen-cycling within the holobiont (Rossbach 2019). The

microniches within animal's is proposed to strongly influence the composition of the microbial

community (Vaishnava *et al.* 2008; Hooper, Littman and Macpherson 2012). Consequently, the

specific composition of each microniche can provide clues as to the community's functional

contribution to host-physiology (Ainsworth, Thurber and Gates 2010).

In the Phylum Cnidaria (sea anemones, corals etc.) many species are known to associate

with different symbiotic microbial partners, including bacteria, and these relationships are known

to be important for the health of the cnidarian host (Rädecker *et al.* 2015; Bourne, Morrow and

Webster 2016; Robbins *et al.* 2019). Previous studies have identified that whole organism

sampling methods used for microbiome analysis can skew results (Sweet, Croquer and Bythell

2011). An advantage of separating an organism into microniches is that it can facilitate the

identification of core microbial symbionts in different compartments (Apprill, Weber and

Santoro 2016). Microorganisms inhabiting specific compartments within a cnidarian host, such

as the epithelium and gastrodermal tissue, likely represent highly specialized symbiotic

associations (Bourne, Morrow and Webster 2016). However, in the model sea anemone,

*Nematostella vectensis* (Phylum: Cnidaria), the presence of microniches of microbial

communities has not previously been identified.

*N. vectensis* is an estuarine sea anemone found in salt marshes along both the Atlantic

and Pacific coastlines of North America (Har 2009). They are diploblastic (i.e., only two

embryonic cell layers) and possess a single oral opening that is surrounded by 12 to 16 tentacles

(Renfer *et al.* 2010). They have a clearly defined pharynx and mesenteries that run along the

oral-aboral axis and a physa at the aboral end used for digging (Layden *et al.* 2010). *N. vectensis*

has rapidly gained popularity as a model organism because it is easy to rear in the lab, simple to

manipulate, and has a wide range of available genomic resources (Putnam *et al.* 2007; Reitzel,

Ryan and Tarrant 2012).

Microbiome analyses through 16S rRNA high-throughput sequencing have increased our

understanding of the microbial community diversity within many organisms including *N.*

*vectensis.* In cnidarians, the microbiome is crucial for nutrient cycling, defense against diseases,

thermoregulation, and the ability to rapidly adapt to environmental stressors (Rosenberg *et al.*

2007; Mortzfeld *et al.* 2016). A previous study conducted comprehensive 16S rRNA gene

sequencing, gene expression analysis, and whole genome sequencing of associated bacteria from

*N. vectensis* (Har *et al.* 2015). This work identified 11 bacterial phyla that were commonly

associated in both wild caught and lab reared *N. vectensis*. These included Cytophaga-

Flexibacter-Bacteroides, Chloroflexi, Cyanobacteria, Deferribacteres, Firmicutes, OD1,

Planctomycetes, Proteobacteria, Spirochetes, Tenericutes and Verrucomicrobia (Har *et al.* 2015).

The microbiome of *N. vectensis* can also be dynamic, with diel lighting and developmental stage

heavily influence microbial community composition (Mortzfeld *et al.* 2016; Leach, Carrier and

Reitzel 2019). However, the potential role of these bacteria within *N. vectensis* is still not fully

understood. The current information on *N. vectensis* suggests that 1) the *N. vectensis* microbiome

can be a highly dynamic and complex community, 2) there are external cues that may affect the

microbiome community, and 3) the specific contribution of certain bacterial taxa may be

overshadowed by bulk sampling methods.

In this study, we distinguished the microbial microniches present within the *N. vectensis*

by using 16S rRNA amplicon sequencing of the three body compartments: the capitulum

(pharynx + tentacles), mesenteries, and physa (Fig. 1). We found that the capitulum possesses

significantly different bacterial community abundances compared with the mesenteries and

physa and that spirochetes are the dominant bacteria phylum present in the capitulum.

Spirochetes were previously identified in *N. vectensis*; however, their compartment specific

nature was not understood. This work better clarifies our understanding of bacteria community

diversity in cnidarians and sheds light on the importance of compartmental sampling in

microbiome studies.

**Methods**

*Nematostella Vectensis Culture and Care*

The individuals used for this study are general cultured animals that have been housed at

the Rosenstiel School of Marine and Atmospheric Science (RSMAS) Cnidarian Immunity Lab

since 2016. All animals were kept in the dark in covered, glass dishes filled with non-circulating

11 parts per thousand (ppt), 2-micron filtered seawater (FSW) at room temperature (~23.8˚C).

Two months prior to the beginning of this study, 30 *N. vectensis* adults were placed into a single

glass dish and fed mussels 3 days a week, and *Artemia* 5 days a week for 2 months in order to

increase their size. Water changes were conducted an hour after each mussel feeding. One week

before beginning the trisection, 12 of the largest individuals were transferred individually into a

12-well plate with sterilized FSW and starved for seven days to minimize microbial carryover

from food.

*Trisection*

The 12 adult *N. vectensis* were anesthetized using 7% $MgCl_2$ solution and then

aseptically trisected. The trisection resulted in the isolation of three tissue compartments from

each anemone: (1) capitulum (pharynx and tentacles), (2) mesenteries, and (3) physa (Fig. 1) The

scalpel and tweezers were both flame and ethanol sterilized between tissue divisions. Each cut

was made as cleanly as possible to limit any cross-contamination of tissue compartments. The

capitulum was removed first, followed by the mesenteries, then the physa. The cut for the

capitulum and mesentery boundary was established as just below the visible pharynx. Some

physa tissue was found in the mesentery compartment as the mesenteries were not directly

removed from the body cavity. However, the physa compartment did not contain any mesentery

material. Each sample was then stored in a 2 mL centrifuge tube with RNALater™ (Sigma

Aldrich, St. Louis, MO) and placed in a -80°C freezer.

*16S rRNA PCR and Sequence Library Preparation*

Total DNA was extracted from 36 samples using the Qiagen™ DNeasy Powersoil Kit.

The concentration and quality of the DNA extracts were assessed using a Nanodrop™

spectrophotometer (Thermo Fisher Scientific Inc., Waltham, MA). DNA was then stored in a

-80°C freezer until library preparation. High-throughput 16s rRNA amplicon sequencing

methods were adapted from the protocol developed by the Earth Microbiome Project (Thompson

*et al.* 2017). In short, the V4 region of the 16S rRNA gene (515F-806R primers) was amplified

using polymerase chain reaction (PCR) on 18 tissue samples (Apprill *et al.* 2015). A negative

control was also amplified to account for extraction and PCR contamination. A total of 50 µL of

PCR reaction mixtures were prepared according to the following recipe: 23 µL PCR-grade water,

20 µL PCR master mix (2x), 1 µL 806R primer (10 µM), 1 µL barcoded 515F primer (10µM),

and 5 µL template DNA. PCR was performed with the following program: 1 x 3 min at 94°C, 35

x (45 s at 94°C, 60 s at 50°C, 90 s at 72°C), 1 x 10 min at 72°C. Amplified DNA was stored at -

20°C for one week. The PCR product was run on a 1.5% agarose gel to check for correct size

amplification. Following the gel, the 16S rRNA V4 amplicon was cleaned using AMPure XP

Beads™ (Beckman Coulter Inc., Brea, CA). DNA concentrations were quantified using the

Qubit dsDNA high sensitivity (HS) kit™ (Thermo Fisher Scientific Inc., Waltham, MA) and

samples were diluted to 4 nM using PCR-grade water. Then 5 µL of each sample was pooled

into a single 1.5 mL tube. The pooled DNA library was sent for sequencing on an Illumina

MiSeq™ PE300 sequencer at the Center for Genome Technology (CGT) at the University of

Miami's Miller School of Medicine for quality control and sequencing. Raw Illumina paired-end

reads have been deposited in the NCBI Sequence Read Archive (SRA) database under

BioProject PRJNA610635.

*Post-Sequencing Processing and Statistical Analysis*

Post-sequencing processing was conducted using the Quantitative Insights in Microbial

Ecology 2 (QIIME2) v2019.1 pipeline (Bolyen *et al.* 2019). Demultiplexed paired-end reads

were merged using DADA2 v1.10.0 and processed to yield amplicon sequence variants (ASVs)

(Callahan *et al.* 2016). Within the DADA2 pipeline, forward sequences were truncated at 200 bp

and reverse sequences were truncated at 180 bp. Taxonomy was assigned using the SILVA 132

99% classifier (https://data.qiime2.org/2019.10/common/silva-132-99-nb-classifier.qza) (Quast

*et al.* 2013). Mitochondria and chloroplast sequences were filtered from the feature table. The

Phyloseq R package v1.28.0 was used for data handling (McMurdie and Holmes 2013).

Normalization across samples was conducted using a variance stabilizing transformation (VST)

on the data using DESeq2 v1.24.0 (Love, Huber and Anders 2014). To account for abundance

and evenness of microbial taxa within the VST dataset, the Shannon-Weiner Diversity Index and

Simpson's Diversity Index were calculated for each compartment region using Phyloseq.

DESeq2 v1.24.0 and the QIIME2 add-on, Analysis of Composition of Microbiomes (ANCOM),

were used for microbial differential abundance analysis against tissue-compartments and

differences were considered significant if they had a p-value < 0.05.  Random forest classifier

modeling was performed using randomForest in order to determine the main bacterial taxa

driving tissue compartment composition (Liaw and Wiener 2002). For beta-diversity

measurements, both Bray-Curtis and Aitchison (Euclidean distances of center-log ratio

transformed counts) distance matrices were assessed as both are commonly used in microbial

community studies. For Bray-Curtis distances, the VST-transformed data was used. For

Aitchison distances, the R package Microbiome v1.6.0 was used to perform a center-log ratio

transformation on the raw count data (Lahti *et al.* 2019). Vegan v2.5.6 was used to calculate

Bray-Curtis and Euclidean distances and the *Adonis* function was used to perform

PERMANOVA analyses in order to statistically assess dissimilarity in microbial community

composition between tissue compartments (Oksanen *et al.* 2013). Ampvis2 v2.5.7 was used to

generate a heat-map of the most abundant bacterial families according to tissue compartment

(Andersen *et al.* 2018). All associated code can be found on Github:

https://github.com/Abonacolta/Nvec_Compartmentalization.

*Mapping N. vectensis ASVs to a phylogenetic tree of Spirochetes*

Spirochaetae phylum 16s rRNA sequences were retrieved from NCBI for representatives

from every major lineage in FASTA format. Using BLASTn, closely clustering (97%) sequences

to the two spirochete ASVs recovered in this study were added to the FASTA file along with a

Planctomycete sequence (Accession: X56305) as an outgroup, giving a total of 64 sequences.

These sequences were aligned using MAFFT v7.450 (auto-setting) with the L-INS-i strategy

(Katoh and Standley 2013). The alignment was trimmed to remove columns with gaps in more

than 70% of the sequences or with a similarity score below 0.001 using TrimAl v1.2 (Capella-

Gutiérrez, Silla-Martínez and Gabaldón 2009). A maximum-likelihood tree was constructed with

RAxML v7.3.0 using the GTRCAT substitution model and 1000 sub-samplings to generate a

best-fit tree (Stamatakis 2014). The final tree was visualized and refined in FigTree v 1.4.4

(Drummond *et al.* 2012).

**Results**

*Spirochetes dominate the capitulum of the Nematostella vectensis microbiome*

The microbial composition of the capitulum (n= 6) is distinct from the mesenteries (n= 6)

and physa (n= 6; Fig. 1). Spirochetes dominated the capitulum, while Proteobacteria dominated

the mesenteries and physa (Fig. 1). In the mesenteries a larger prevalence of

Gammaproteobacteria was found (Fig. S2). The physa of *N. vectensis* was dominated by the

classes Gammaproteobacteria, Alphaproteobacteria, and Bacteroidetes (Fig. S2). Of the

Gammaproteobacteria which dominated the mesenteries and physa compartments,

Oceanospirillales and Pseudomonadales were the primary orders, with Pseudomonadales being

more abundant than Oceanospirillales (Fig. 2). The relative abundance of the top nine most

abundant bacterial families revealed a capitulum-specific dominance of Spirochaetaceae bacteria

(~66%), with a high abundance of Pseudomonadaceae in the mesenteries and physa (~36% and

~38%, respectively) (Fig. 3). At the family-level, Pseudomonadaceae, Rhodobacteraceae,

Alteromonadaceae, and Saccharospirillaceae bacteria made up the majority of amplicon

sequence variants (ASVs) in the physa (Fig. S3). The mesenteries showed similar bacterial

community composition as the physa, with the exclusion of Alteromonadaceae (only appearing

in >5% abundance in just one individual) (Fig. S3). Overall the microbiomes were consistent

between replicates, however, one individual harbored a different compartmental microbiome

from the others in this study (Fig. S1).

Comparing the differential abundance of taxa across tissue compartments using a random

forest classifier showed that the phyla Lentisphaerae, Proteobacteria, and Spirochetes were the

strongest predictors of tissue compartments. According to differential abundance analysis,

Spirochetes and Fusobacteria were significantly more abundant in the capitulum compared to the

other compartments (DESeq2; p-adj=0.0000 and 0.0003, respectively). Lentisphaerae,

Proteobacteria, and Chloroflexi were more abundant in the mesenteries compared to the

capitulum (DESeq2; p-adj= 0.0000, 0.0058, and 0.0009, respectively). According to differential

abundance analysis using ANCOM, Spirochetes and Bacteroidetes were the most differentially

abundant phyla among tissue compartments (ANCOM; W = 16 & 14, respectively).

*Alpha-diversity metrics differed significantly between the capitulum and other compartments*

Alpha-diversity metrics showed a significant difference in the biodiversity and species

richness present in different tissue compartments. Species richness among the capitulum and the

mesenteries and physa were significantly different (ANOVA; p = 0.023, p = 0.033, respectively;

Fig. 4). The Simpson's Index between the capitulum and the mesenteries and physa was also

significantly different (ANOVA; p = 0.039, p = 0.022, respectively; Fig. 4). However, the

Shannon Diversity Index did not show a significant difference in alpha-diversity among body

regions (Fig. 4). Overall, the capitulum of *N. vectensis* showed lower species richness compared

to the mesenteries and physa, which had very similar species richness across individuals. A

similar pattern was identified in the Simpson's Index, with the capitulum showing a lower

species diversity, but with a higher variation between individuals.

*Beta diversity metrics inconclusive regarding the significance between tissue compartments*

Both Bray-Curtis Dissimilarity and Aitchison distance were used to assess the differences

in the microbial communities between tissue compartments (Fig. 5). A multivariate homogeneity

of groups dispersions analysis confirmed that an ANOVA test could appropriately be performed

on both datasets (Koleff, Gaston and Lennon 2003). Bacterial communities between tissue

compartments showed significant dissimilarity (PERMANOVA; permutations = 9999, p =

0.0315) according to the Bray-Curtis Dissimilarity Index. However, the same PERMANOVA

test on the Aitchison distances found insignificant dissimilarity between tissue compartments

(PERMANOVA; permutations = 9999, p = 0.3294).

*Phylogenetic analysis of the Spirochete ASVs*

The majority of spirochetes found in the capitulum was from a single ASV (Spirochete 1)

within the Spirochaetales family. This ASV along with the other spirochete ASV (Spirochete 2)

recovered in this study were mapped to a maximum-likelihood tree of representative

Spirochaetae sequences. Strong bootstrap support (93% bootstrap) placed Spirochete 1 as closely

related to four spirochete ASVs recovered from wild *N. vectensis* from Har *et al.* 2015 and to a

spirochete ASV recovered from the cold-water coral, *Lophelia pertusa* (Fig. 6). Together, these

sequences formed a strongly supported (90%) monophyletic clade with the two representative

species of the recently described *Oceanispirocheata* genus. Spirochete 2, along with several

closely clustering wastewater ASVs, showed strong relation (100% bootstrap) to *Turneriella*

*parva*. Interestingly spirochetes associated with a diverse group of corals, including *Corallium*

*rubrum, Orbicella faveolata, and Lophelia pertusa,* did form a well-supported (100% bootstrap)

monophyletic clade within the *Leptospira* genus.

**Discussion**

In this study, spirochetes (predominantly from 1 ASV) within the Spirochaetaceae family

made up over ~65% of the bacterial community within the capitulum of *N. vectensis*, but only

~1% of the community in the mesenteries and physa (Figs. 1 & 3). Although spirochetes are

known to be one of the most abundant phyla within the microbiome of *N. vectensis* (Mortzfeld *et*

*al.* 2016), their compartment-specific distribution within the microbiome was masked by whole

organism sampling and the high prevalence of Proteobacteria. The differences in dominant taxa

between the capitulum and the other two tissue compartments in *N. vectensis* highlight the need

to use compartmentalization to better understand the roles of the microbiome within the

holobiont- be that in metabolism, defense, or nutrient cycling. This finding is similar to another

microbiome compartmentalization study of the Crown-of-thorns Starfish, *Acanthaster planci*,

which found a high relative abundance (43-64%) of spirochetes (predominantly from 1 OTU)

within body wall samples. Prior to this Alphaproteobacteria was thought to be the most abundant

phylum of bacteria across the sea stars (Høj *et al.* 2018).

Spirochetes are gram-negative, organo-chemoheterotrophs known to attach to each other

and to other live cells, commonly as epibionts (Wier *et al.* 2010). They have been identified as

symbionts in bivalves, corals, oligochaetes, and termites (as epibionts on intestinal protists)

where they perform reductive acetogenesis and nitrogen fixation (Margulis, Nault and Sieburth

1991; Lilburn, Schmidt and Breznak 1999; Campbell and Cary 2001; Ohkuma *et al.* 2015). In

cnidarians, spirochetes are the predominant bacteria of the red coral, *Corallium rubrum*, and the

octocoral, *Lobophytum pauciflorum,* where they are hypothesized to play a significant role in

nitrogen fixation and even microbial community structuring through antibiotic secretion (Van De

Water *et al.* 2016; Wessels *et al.* 2017). They have also been identified in *Hydra vulgaris,*

predominately in transparent vacuoles at the apical end of epitheliomuscular cells (Hufnagel and

Myhal 1977). The predominance of spirochetes as symbionts in some cnidarians suggests a

potential conserved benefit to the Cnidaria hosts. While we do not know why there is a

consistent, and highly specific compartment abundance of spirochetes in the capitulum of *N.*

*vectensis,* we hypothesize that it may play a significant role in the host physiology. In the future,

testing the capabilities of spirochetes to perform nitrogen fixation in the capitulum of

*Nematostella* may inform our understanding of these potential symbionts.

In order to better understand the phylogenetic placement of the spirochetes within *N.*

*vectensis,* we conducted a phylogenetic analysis of our ASVs with representative sequences from

every major lineage within Spirochaetae. The predominant spirochete ASV (Spirochete 1)

recovered from the capitulum of *N. vectensis* is part of a strongly supported monophyletic clade

of spirochete sequences recovered from wild *N. vectensis* supporting this spirochete as a

consistent associate of the anemone (Fig. 6). The potential role of the spirochetes within the

capitulum of *N. vectensis* should be investigated further as this relationship appears to be

conserved, regardless of environment. These *N. vectensis* ASVs also showed a strong relation to

a spirochete recovered from the cold-water coral, *Lophelia pertusa*. Together, these sequences

clustered closer to the recently described *Oceanispirochaeta*, rather than *Spirochaeta perfilievii*,

supporting them as members of this genus. The *Oceanispirochaeta* consists solely of two

representatives, *O. litoralis* and *O. sediminicola*, which are gram-negative, moderately

halophilic, and obligately anaerobic (Subhash and Lee 2017). Spirochete 2, along with various

wastewater associated spirochetes, formed a monophyletic clade with *Turneriella parva* (Fig. 6).

*Turneriella parva* is a gram-negative, obligately aerobic spirochete with morphological

similarities to *Leptospira* (Stackebrandt *et al.* 2013). Interestingly, a clade of cnidarian-

associated spirochetes clustered closely within the *Leptospira* genus (Fig. 6). Despite the broad

geographical ranges of these coral species (*O. faveolata* is a Caribbean species, *C. rubrum* is a

Mediterranean species, & *L. pertusa* is a deep sea species), they all share an association with

*Leptospira*. The potential role of these cnidarian associated *Leptospira* species should be

investigated further.

Other abundant phyla of the *N. vectensis* microbiome revealed in this study include

Proteobacteria, Actinobacteria, Firmicutes, and Tenericutes (Fig 1). In previous studies on the

microbiome of lab-raised *N. vectensis*, Proteobacteria was found to be the most abundant

bacterial phylum, with Firmicutes, Planctomycetes, Tenericutes, Verrucomicrobia, and

Spirochetes having lower, but still significant, abundances (Har *et al.* 2015; Mortzfeld *et al.*

2016; Leach, Carrier and Reitzel 2019). In particular, Gammaproteobacteria have been shown to

play an important role in microbial community structuring in juvenile *N. vectensis* (Domin *et al.*

2018). Additionally, fluorescent *in situ* hybridization revealed that Proteobacteria and

Pseudomonas occur predominantly in aggregates along the walls of the mesenteries (Har 2009).

We found a similar pattern of compartmentalization in our study, where only the physa and

mesentery tissue compartments had a high abundance of Proteobacteria (Orders: Pseudomonas

and Oceanospirillales). These two compartments also showed significantly higher alpha-diversity

compared to the capitulum, most likely due to the dominance of the singular Spirochete ASV in

the capitulum. One of our beta-diversity analyses also supported our hypothesis that microbial

communities differed significantly between tissue compartments, with the capitulum harboring a

notably distinct microbial community from the mesenteries and physa.

Interestingly, one study found that as *N. vectensis* develops, Betaproteobacteria,

Actinobacteria, and Bacteroidetes decrease, while Spirochetes become one of the most abundant

phyla in the adult organism (Mortzfeld *et al.* 2016). The development and growth of the

capitulum during this time span may explain the increase in abundance of spirochetes in the adult

organism compared to its earlier developmental stages. Spirochetes may play an active role in

the capitulum development of *N. vectensis* or may preferentially thrive in this tissue

compartment over other bacteria. Their ability to establish dominance over the Proteobacteria

present throughout the rest of *N. vectensis* implies a competitive advantage on the part of the

Spirochete.

In this study we characterized the microbiome of three distinct body compartments within

*N. vectensis*. A capitulum-specific dominance of spirochetes was uncovered highlighting the

possible importance of this bacteria in *N. vectensis* physiology. Previous bulk-sampling

techniques could not untangle compartment-specific bacterial associations, which should be

taken into consideration in future cnidarian studies. Specifically, the role of spirochetes within

the capitulum of *N. vectensis* should be further considered in subsequent studies. Overall, our

findings further establish spirochetes as consistent bacteria associates within *N. vectensis* and

suggests that the functional role of this bacteria in the capitulum may be important for the

anemone.

**Acknowledgements**

346       Authors would like to thank members of the Cnidarian Immunity Lab for comments and

348    suggestions as this project was developing. AMB: was supported by the RSMAS Small

349    Undergraduate Research Grant Experience (SURGE) Award. NTK, JDC, and MTC: were

350    supported by startup funds from University of Miami, Rosenstiel School of Marine and

351    Atmospheric Sciences.

**References**

Aagaard K, Ma J, Antony KM *et al.* The Placenta Harbors a Unique Microbiome. *Sci Transl*

*Med* 2014;**6**, DOI: 10.1126/scitranslmed.3009864.

Ainsworth TD, Thurber RV, Gates RD. The future of coral reefs: a microbial perspective. *Trends*

*Ecol Evol* 2010;**25**:233–40.

Apprill A, Mcnally S, Parsons R *et al.* Minor revision to V4 region SSU rRNA 806R gene

primer greatly increases detection of SAR11 bacterioplankton. *Aquat Microb Ecol*

2015;**75**:129–37.

Apprill A, Weber LG, Santoro AE. Distinguishing between Microbial Habitats Unravels

Ecological Complexity in Coral Microbiomes. *mSystems* 2016;**1**:e00143-16.

Bolyen E, Rideout JR, Dillon MR *et al.* Reproducible, interactive, scalable and extensible

microbiome data science using QIIME 2. *Nat Biotechnol* 2019;**37**:852–7.

Bourne DG, Morrow KM, Webster NS. Insights into the Coral Microbiome: Underpinning the

Health and Resilience of Reef Ecosystems. *Annu Rev Microbiol* 2016;**70**:317–40.

Callahan BJ, Mcmurdie PJ, Rosen MJ *et al.* DADA2: High resolution sample inference from

Illumina amplicon data. *Nat Methods* 2016;**13**:581–3.

Campbell BJ, Cary SC. Characterization of a Novel Spirochete Associated with the

Hydrothermal Vent Polychaete Annelid, Alvinella pompejana. *Appl Environ Microbiol*

2001;**67**:110–7.

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: A tool for automated alignment

trimming in large-scale phylogenetic analyses. *Bioinformatics* 2009;**25**:1972–3.

Domin H, Zurita-Gutiérrez YH, Scotti M *et al.* Predicted bacterial interactions affect in vivo

microbial colonization dynamics in Nematostella. *Front Microbiol* 2018;**9**:1–12.

Drummond AJ, Suchard MA, Xie D *et al.* Bayesian phylogenetics with BEAUti and the BEAST

1.7. *Mol Biol Evol* 2012;**29**:1969–73.

Foster KR, Schluter J, Coyte KZ *et al.* The evolution of the host microbiome as an ecosystem on

a leash. *Nature* 2017;**548**:43–51.

Har JY. Introducing the starlet sea anemone Nematostella vectensis as a model for investigating.

2009.

Har JY, Helbig T, Lim JH *et al.* Microbial diversity and activity in the Nematostella vectensis

holobiont: Insights from 16S rRNA gene sequencing, isolate genomes, and a pilot-scale

survey of gene expression. *Front Microbiol* 2015;**6**, DOI: 10.3389/fmicb.2015.00818.

Høj L, Levy N, Baillie B *et al.* Crown-of-Thorns Sea Star Acanthaster cf. solaris Has Tissue-

Characteristic Microbiomes with Potential Roles in Health and Reproduction. *Appl Environ*

*Microbiol* 2018;**84**:1–18.

Hooper L V., Littman DR, Macpherson AJ. Interactions between the microbiota and the immune

system. *Science (80- )* 2012;**336**:1268–73.

Hufnagel LA, Myhal ML. Observations on a Spirochaete Symbiotic in Hydra. *Trans Am Microsc*

*Soc* 1977;**96**:406–11.

Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: Improvements

in performance and usability. *Mol Biol Evol* 2013;**30**:772–80.

Koleff P, Gaston KJ, Lennon JJ. Measuring beta diversity for presence-absence data. *J Anim

Ecol* 2003;**72**:367–82.

Layden MJ, Meyer NP, Pang K *et al.* Expression and phylogenetic analysis of the zic gene

family in the evolution and development of metazoans. *Evodevo* 2010;**1**:1–16.

Leach WB, Carrier TJ, Reitzel AM. Diel patterning in the bacterial community associated with

the sea anemone Nematostella vectensis. *Ecol Evol* 2019;**9**:9935–47.

Liaw A, Wiener M. Classification and Regression by randomForest. *R News* 2002;**2**:18–22.

Lilburn TG, Schmidt TM, Breznak JA. Phylogenetic diversity of termite gut spirochaetes.

*Environ Microbiol* 1999;**1**:331–45.

Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq

data with DESeq2. *Genome Biol* 2014;**15**:1–21.

Mandal S, Van Treuren W, White RA *et al.* Analysis of composition of microbiomes: a novel

method for studying microbial composition. *Microb Ecol Heal Dis* 2015;**26**, DOI:

10.3402/mehd.v26.27663.

Margulis L, Nault L, Sieburth JM. Cristispira from oyster styles: complex morphology of large

symbiotic spirochetes. *Symbiosis* 1991;**11**:1–17.

McMurdie PJ, Holmes S. Phyloseq: An R Package for Reproducible Interactive Analysis and

Graphics of Microbiome Census Data. *PLoS One* 2013;**8**, DOI:

10.1371/journal.pone.0061217.

Mortzfeld BM, Urbanski S, Reitzel AM *et al.* Response of bacterial colonization in Nematostella

vectensis to development, environment and biogeography. *Environ Microbiol*

2016;**18**:1764–81.

Ohkuma M, Noda S, Hattori S *et al.* Acetogenesis from H2 plus CO2and nitrogen fixation by an

endosymbiotic spirochete of a termite-gut cellulolytic protist. *Proc Natl Acad Sci U S A*

2015;**112**:10224–30.

Price MN, Dehal PS, Arkin AP. FastTree 2 - Approximately maximum-likelihood trees for large

alignments. *PLoS One* 2010;**5**, DOI: 10.1371/journal.pone.0009490.

Putnam NH, Srivastava M, Hellsten U *et al.* Sea anemone genome reveals ancestral eumetazoan

gene repertoire and genomic organization. *Science (80- )* 2007;**317**:86–94.

Quast C, Pruesse E, Yilmaz P *et al.* The SILVA ribosomal RNA gene database project:

Improved data processing and web-based tools. *Nucleic Acids Res* 2013;**41**:590–6.

Rädecker N, Pogoreutz C, Voolstra CR *et al.* Nitrogen cycling in corals: The key to

understanding holobiont functioning? *Trends Microbiol* 2015;**23**:490–7.

Reitzel AM, Ryan JF, Tarrant AM. Establishing a model organism: A report from the first annual

Nematostella meeting. *BioEssays* 2012;**34**:158–61.

Renfer E, Amon-Hassenzahl A, Steinmetz PRH *et al.* A muscle-specific transgenic reporter line

of the sea anemone, Nematostella vectensis. *Proc Natl Acad Sci U S A* 2010;**107**:104–8.

Ricci F, Rossetto Marcelino V, Blackall LL *et al.* Beneath the surface: Community assembly and

functions of the coral skeleton microbiome. *Microbiome* 2019;**7**:1–10.

Robbins SJ, Singleton CM, Chan CX *et al.* A genomic view of the reef-building coral Porites

lutea and its microbial symbionts. *Nat Microbiol* 2019:1–11.

Rosenberg E, Koren O, Reshef L *et al.* The role of microorganisms in coral health , disease and

evolution. *Nat Rev Microbiol* 2007;**5**:355–62.

Rossbach S. Tissue-Specific Microbiomes of the Red Sea Giant Clam Tridacna maxima

Highlight Differential Abundance of Endozoicomonadaceae. 2019;**10**, DOI:

10.3389/fmicb.2019.02661.

Stackebrandt E, Chertkov O, Lapidus A *et al.* Genome sequence of the free-living aerobic

spirochete Turneriella parva type strain (HT), and emendation of the species Turneriella

parva. *Stand Genomic Sci* 2013;**8**:228–38.

Stamatakis A. RAxML version 8: A tool for phylogenetic analysis and post-analysis of large

phylogenies. *Bioinformatics* 2014;**30**:1312–3.

Subhash Y, Lee SS. Description of oceanispirochaeta sediminicola gen. Nov., sp. nov., an

obligately anaerobic bacterium isolated from coastal marine sediments, and reclassification

of spirochaeta litoralis as oceanispirochaeta litoralis comb. nov. *Int J Syst Evol Microbiol*

2017;**67**:3403–9.

Sweet MJ, Croquer A, Bythell JC. Bacterial assemblages differ between compartments within

the coral holobiont. *Coral Reefs* 2011;**30**:39–52.

Thompson LR, Sanders JG, McDonald D *et al.* A communal catalogue reveals Earth's multiscale

microbial diversity. *Nature* 2017;**551**:457–63.

Vaishnava S, Behrendt CL, Ismail AS *et al.* Paneth cells directly sense gut commensals and

maintain homeostasis at the intestinal host-microbial interface. *Proc Natl Acad Sci U S A*

2008;**105**:20858–63.

Van De Water JAJM, Melkonian R, Junca H *et al.* Spirochaetes dominate the microbial

community associated with the red coral Corallium rubrum on a broad geographic scale. *Sci*

*Rep* 2016;**6**:1–7.

Wessels W, Sprungala S, Watson SA *et al.* The microbiome of the octocoral Lobophytum

pauciflorum: Minor differences between sexes and resilience to short-term stress. *FEMS*

*Microbiol Ecol* 2017;**93**:1–13.

Wier AM, Sacchi L, Dolan MF *et al.* Spirochete Attachment Ultrastructure: Implications for the

Origin and Evolution of Cilia. *Biol Bull* 2010;**218**:25–35.

**Fig 1. Compartmentalization of the microbiome in *N. vectensis*.** The microbiome of *N.*

*vectensis* was divided into three tissue compartments for this study: the capitulum, mesenteries,

and physa. The capitulum had a bacterial community dominated by Spirochetes. The mesenteries

and physa had a bacterial community dominated by Gammaproteobacteria and

Alphaproteobacteria, with very little presence of Spirochetes.

**Fig 2. The relative abundance of bacterial orders recovered from *N. vectensis* in individuals**

**and tissue compartments.** Only taxonomic orders with a relative abundance above 2% were

incorporated. Spirochaetales dominated the bacterial communities in the majority of capitulum

samples (Individual 1 being an outlier). The mesentery and physa body regions showed more

variety in their order composition with Pseudomonadales, Alteromonadales, and

Oceanospirillales making up the majority of most samples.

**Fig 3. Heat-map of the percent read abundance for top bacterial families found in *N.***

***vectensis* separated by tissue compartment.** Bacteria of the family Pseudomonadaceae were

found to be the most prevalent in the mesenteries and physa tissue of *N. vectensis* (37.9% and

36.3% abundance, respectively). In contrast, Pseudomonadaceae only had a 7.3% abundance in

capitulum tissue. Likewise, Spirochaetaceae were found to make up 65.8% of the community

within the capitulum tissue of *N. vectensis*, while only being 1.2% and 0.1% abundant in the

mesentery and physa tissues, respectively. Overall, the capitulum tissue appears to have a very

distinct community from the mesenteries and physa.

**Fig 4. Bacterial community Alpha diversity measures between tissue regions.** Species

richness between the capitulum, mesenteries, and physa was significantly different (ANOVA;

p=0.023,  p=0.033, respectively). The Simpson Alpha Diversity Index between the capitulum,

mesenteries and physa was significantly different (ANOVA; p=0.039,  p=0.022, respectively).

The Shannon Diversity Index did not show a significant difference in the bacterial communities

between body regions.

**Fig 5. Beta diversity measures between tissue regions of *N. vectensis*.** (A) Bray-Curtis PCoA

of *N. vectensis* bacterial communities after a variance stabilizing transformation. Capitulum

samples clustered tightly, while physa and mesentery communities clustered closely together.

Individual 1 appeared to be an outlier. (B) Aitchison PCA (Euclidean distance of center-log ratio

transformed sample counts) of *N. vectensis* bacterial communities. Again, most capitulum

samples clustered closely together, however they were also clustered with mesenteries and physa

samples.

**Fig 6. Best-fit maximum likelihood tree based on 16S rRNA sequences depicting the**

**phylogenetic relationship of the spirochetes recovered from the capitulum of *Nematostella***

***vectensis*.** This tree was constructed using RAxML v7.3.0 with 1000 sub samplings. Bootstrap

values (> 50%) are shown. Nodes with 100% bootstrap support are indicated by open circles.

GenBank accession numbers are listed before each sequence's name. The host/habitat of

uncultured species is in parentheses. Spirochaetales associated with cnidarians are in bold.

*Gemmata obscuriglobus* was used as an outgroup to root the tree. The *N. vectensis* spirochetes

from this study are indicated by an asterisk. The predominant spirochete ASV from this study

(Spirochete 1) forms a strongly supported (100% bootstrap) monophyletic clade with spirochete

sequences recovered from wild *N. vectensis*.  Spirochete 2 clusters within the *Turneriella* genus.

**Fig 1. Compartmentalization of the microbiome in *N. vectensis*.** The microbiome of *N. vectensis* was divided into three tissue compartments for this study: the capitulum, mesenteries, and physa. The capitulum had a bacterial community dominated by Spirochetes. The mesenteries and physa had a bacterial community dominated by Gammaproteobacteria and Alphaproteobacteria, with very little presence of Spirochetes.

338x218mm (300 x 300 DPI)

**Fig 2. The relative abundance of bacterial orders recovered from *N. vectensis* in individuals and tissue compartments.** Only taxonomic orders with a relative abundance above 2% were incorporated. Spirochaetales dominated the bacterial communities in the majority of capitulum samples (Individual 1 being an outlier). The mesentery and physa body regions showed more variety in their order composition with Pseudomonadales, Alteromonadales, and Oceanospirillales making up the majority of most samples.

740x529mm (72 x 72 DPI)

**Fig 3. Heat-map of the percent read abundance for top bacterial families found in *N. vectensis* separated by tissue compartment.** Bacteria of the family Pseudomonadaceae were found to be the most prevalent in the mesenteries and physa tissue of *N. vectensis* (37.9% and 36.3% abundance, respectively). In contrast, Pseudomonadaceae only had a 7.3% abundance in capitulum tissue. Likewise, Spirochaetaceae were found to make up 65.8% of the community within the capitulum tissue of *N. vectensis*, while only being 1.2% and 0.1% abundant in the mesentery and physa tissues, respectively. Overall, the capitulum tissue appears to have a very distinct community from the mesenteries and physa.

846x529mm (72 x 72 DPI)

**Fig 4. Bacterial community Alpha diversity measures between tissue regions.** Species richness between the capitulum, mesenteries, and physa was significantly different (ANOVA; p=0.023, p=0.033, respectively). The Simpson Alpha Diversity Index between the capitulum, mesenteries and physa was significantly different (ANOVA; p=0.039, p=0.022, respectively). The Shannon Diversity Index did not show a significant difference in the bacterial communities between body regions.

740x529mm (72 x 72 DPI)

**Fig 5. Beta diversity measures between tissue regions of *N. vectensis*.** (A) Bray-Curtis PCoA of *N. vectensis* bacterial communities after a variance stabilizing transformation. Capitulum samples clustered tightly, while physa and mesentery communities clustered closely together. Individual 1 appeared to be an outlier. (B) Aitchison PCA (Euclidean distance of center-log ratio transformed sample counts) of *N. vectensis* bacterial communities. Again, most capitulum samples clustered closely together, however they were also clustered with mesenteries and physa samples.

355x127mm (300 x 300 DPI)

**Fig 6. Best-fit maximum likelihood tree based on 16S rRNA sequences depicting the phylogenetic relationship of the spirochetes recovered from the capitulum of *Nematostella vectensis*.** This tree was constructed using RAxML v7.3.0 with 1000 sub samplings. Bootstrap values (> 50%) are shown. Nodes with 100% bootstrap support are indicated by open circles. GenBank accession numbers are listed before each sequence's name. The host/habitat of uncultured species is in parentheses. Spirochaetales associated with cnidarians are in bold. *Gemmata obscuriglobus* was used as an outgroup to root the tree. The *N. vectensis* spirochetes from this study are indicated by an asterisk. The predominant spirochete ASV from this study (Spirochete 1) forms a strongly supported (100% bootstrap) monophyletic clade with spirochete sequences recovered from wild *N. vectensis*.  Spirochete 2 clusters within the *Turneriella* genus.

294x243mm (300 x 300 DPI)

**Title:** Microniche sampling of the microbiome in the Starlet Sea Anemone, *Nematostella*

*vectensis*, reveals a compartment-specific dominance of Spirochetes

**Authors:** A. M. Bonacolta, M. T. Connelly, S. Rosales, J. del Campo, N. Traylor-Knowles

**Supplemental Material**

**Fig S1. The relative abundance of bacterial phyla recovered from *N. vectensis* in**

**individuals and tissue compartments** Only taxonomic phyla with a relative abundance above

2% were incorporated. Spirochetes dominated the bacterial communities in the majority of

capitulum samples (Individual 1 again being an outlier). Proteobacteria dominated the bacterial

communities of mesentery and physa samples.

**Fig S2. The relative abundance of bacterial classes recovered from *N. vectensis* in**

**individuals and tissue compartments** Only taxonomic classes with a relative abundance above

2% were incorporated. Spirochaetia dominated the bacterial communities in the majority of

capitulum samples (Individual 1 again being an outlier). Gammaproteobacteria dominated the

bacterial communities of mesentery and physa samples.

**Fig S3. The relative abundance of bacterial families recovered from *N. vectensis* in**

**individuals and tissue compartments.** Only taxonomic families with a relative abundance

above 5% were incorporated. Spirocheataceae dominated the bacterial communities in the

majority of capitulum samples (Individual 1 again being an outlier). The mesentery and physa

body regions showed more variety in their bacterial family composition with

Pseudomonadaceae, Saccharospirillaceae, Colwelliaceae, and Alteromonadaceae making up the

majority of most samples.