Title: The Starlet Sea Anemone, *Nematostella vectensis,* possesses body region-specific bacterial

associations with spirochetes dominating the capitulum

Authors: A. M. Bonacolta[1], M. T. Connelly[1], S. Rosales[2,3], J. del Campo[1], N. Traylor-

Knowles[1,*]

[1]: Rosenstiel School of Marine and Atmospheric Science, Department of Marine Biology and

Ecology, University of Miami, 4600 Rickenbacker Causeway, Florida 33149, USA

[2]: Atlantic Oceanographic and Meteorological Laboratory, National Oceanographic and

Atmospheric Administration, Miami, Florida, USA

[3]: Cooperative Institute for Marine and Atmospheric Studies, University of Miami, Miami,

Florida, USA

*: Corresponding author: Nikki Traylor-Knowles, ngt14@miami.edu

Abstract

Sampling of different body regions can reveal highly specialized bacterial associations

within the holobiont and facilitate identification of core microbial symbionts that would

otherwise be overlooked by bulk sampling methods. Here we characterized compartment-

specific associations present within the model cnidarian *Nematostella vectensis* by dividing its

morphology into three distinct body regions. This sampling design allowed us to uncover a

capitulum-specific dominance of spirochetes within *N. vectensis*. Bacteria from the family

Spirochaetaceae made up 66% of the community in the capitulum, while only representing 1.2%

and 0.1% of the communities in the mesenteries and physa, respectively. A phylogenetic analysis

of the predominant spirochete sequence recovered from *N. vectensis* showed a close relation to

spirochetes previously recovered from wild *N. vectensis*. These sequences clustered closer to the

recently described genus *Oceanispirochaeta*, rather than *Spirochaeta perfilievii*, supporting them

as members of this clade. This suggests a consistent and potentially important association

between *N. vectensis* and spirochetes from the order Spirochaetales.

Introduction

The discovery of microniche specific microbial communities has become important for

our understanding of microbial interactions and diversity (Ainsworth, Thurber and Gates 2010;

Bourne, Morrow and Webster 2016). Microbial niches have been identified in many different

organisms from an ecosystem scale, all the way down to the cellular level (Aagaard *et al.* 2014;

Foster *et al.* 2017; Ricci *et al.* 2019). For example, in the Giant Clam, *Tridacna maxima*,

different tissues harbor distinct microbial communities, with the gills harboring a high

prevalence of *Endozoicomonas* bacteria. Compartmental evaluation of *T. maxima* helped uncover

Endozoicomonas's role in nitrogen-cycling within the holobiont (Rossbach 2019). The

microniches within animal tissues are proposed to strongly influence the composition of the

microbial community (Vaishnava *et al.* 2008; Hooper, Littman and Macpherson 2012).

Consequently, the specific composition of each microniche can provide clues as to the

community's functional contribution to host-physiology (Ainsworth, Thurber and Gates 2010).

In the Phylum Cnidaria (sea anemones, corals etc.) many species are known to associate

with different symbiotic microbial partners, including bacteria, and these relationships are known

to be important for the health of the cnidarian host (Rädecker *et al.* 2015; Bourne, Morrow and

Webster 2016; Robbins *et al.* 2019). Previous studies have identified that whole organism

sampling methods used for microbiome analysis can skew results by preferentially sampling

certain compartments or by combining them and drawing conclusions based on the

agglomeration (Sweet, Croquer and Bythell 2011). An advantage of compartmental sampling of

an organism's body is that it can facilitate the identification of core microbial symbionts in

different body compartments, which likely represent highly specialized symbiotic associations

(Apprill, Weber and Santoro 2016; Bourne, Morrow and Webster 2016). However, in the model

sea anemone, *Nematostella vectensis*, the presence of body region-specific associations of

microbial communities has not previously been identified.

N. vectensis is an estuarine sea anemone found in salt marshes along both the Atlantic

and Pacific coastlines of North America (Har 2009). They possess only two embryonic cell

layers (i.e. diploblastic) and possess a single oral opening that is surrounded by 12 to 16 tentacles

(Renfer *et al.* 2010). They have a clearly defined pharynx and mesenteries that run along the

oral-aboral axis and a physa at the aboral end used for digging (Layden *et al.* 2010). *N. vectensis*

has rapidly gained popularity as a model organism because it is easy to rear in the lab, simple to

manipulate, and has a wide range of available genomic resources (Putnam *et al.* 2007; Reitzel,

Ryan and Tarrant 2012).

72   *N. vectensis* microbiome 16S rRNA amplicon high-throughput sequencing has increased

our understanding of the microbial community diversity which is crucial for understanding

nutrient cycling, defense against diseases, thermoregulation, and the ability to rapidly adapt to

environmental stressors (Rosenberg *et al.* 2007; Mortzfeld *et al.* 2016). Previous studies on wild

and lab raised *N. vectensis* using 16S rRNA gene sequencing, gene expression analysis, and

whole genome sequencing found that they primarily associate with the bacteria phylum :

Bacteroides, Chloroflexi, Cyanobacteria, Firmicutes, Proteobacteria, and Spirochetes (Har *et al.*

2015a). Other studies have also found that the *N. vectensis* microbiome can be dynamic with diel

lighting and developmental stage heavily influencing the microbial community composition

(Mortzfeld *et al.* 2016; Domin et al. 2018; Leach, Carrier and Reitzel 2019). The current

information on *N. vectensis* suggests that the microbiome is a highly dynamic and complex

community that responds to environmental cues. However, these patterns were observed in the

whole organism, which may obscure important niche specific bacteria compartments within *N.*

vectensis.

86   In this study, we distinguished body region-specific bacterial associations present within

N. vectensis by using 16S rRNA gene amplicon sequencing of the three body compartments: the

capitulum (pharynx + tentacles), mesenteries, and physa (Fig. 1). We found that the capitulum

possesses significantly different bacterial community abundances compared with the mesenteries

and physa and that spirochetes are the dominant bacterial phylum present in the capitulum.

Spirochetes were previously identified in *N. vectensis*; however, their compartment specific

nature was unknown. The work presented better clarifies our understanding of bacteria

community diversity in *N. vectensis* and sheds light on the importance of compartmental

sampling in microbiome studies.

Methods

Nematostella vectensis Culture and Care

The individuals used for this study are lab cultured animals that have been housed at the

Rosenstiel School of Marine and Atmospheric Science (RSMAS) Cnidarian Immunity Lab since

2016. All animals were kept in the dark within covered, glass dishes filled with non-circulating

11 parts per thousand (ppt), 2-micron filtered seawater (FSW) at room temperature (~23.8˚C).

Two months prior to the beginning of this study, *N. vectensis* adults were placed into a single

glass dish and fed mussels three days a week, and *Artemia* five days a week for two months in

order to increase their size. Water changes were conducted an hour after each mussel feeding.

One week before beginning the trisection (see section below), 12 of the largest individuals were

transferred individually into a 12-well plate with sterilized FSW and starved for seven days to

minimize microbial carryover from food. Previous studies have also used a starving period of

two days to eliminate digested food debris (Har *et al.* 2015b; Leach, Carrier and Reitzel 2019)

Trisection

The 12 adult *N. vectensis* were anesthetized using 7% $MgCl_2$ solution and then

aseptically trisected. The trisection resulted in the isolation of three tissue compartments from

each anemone: (1) capitulum (pharynx and tentacles), (2) mesenteries, and (3) physa (Fig. 1).

The scalpel and tweezers were both flame and ethanol sterilized between tissue divisions. Each

cut was made as cleanly as possible to limit any cross-contamination of tissue compartments.

The capitulum was removed first, followed by the mesenteries, then the physa. The cut for the

capitulum and mesentery boundary was established as just below the visible pharynx. Some

physa tissue was found in the mesentery compartment as the mesenteries were not directly

removed from the body cavity. However, the physa compartment did not contain any mesentery

material. Each sample was then stored in a 2 mL centrifuge tube with RNALater™ (Sigma

Aldrich, St. Louis, MO) and placed in a -80°C freezer. In total there were 18 samples (n=6

capitulum, n=6 mesenteries, and n=6 physa) that were further processed.

16S rRNA PCR and Sequence Library Preparation

Total DNA was extracted from 18 samples using the Qiagen™ DNeasy Powersoil Kit.

The concentration and quality of the DNA extracts were assessed using a Nanodrop™

spectrophotometer (Thermo Fisher Scientific Inc., Waltham, MA). DNA was then stored in a

-80°C freezer until library preparation. High-throughput 16S rRNA amplicon sequencing

methods were adapted from the protocol developed by the Earth Microbiome Project (Thompson

et al. 2017). In short, the V4 region of the 16S rRNA gene (515F-806R primers) was amplified

using polymerase chain reaction (PCR) on 18 tissue samples (Apprill *et al.* 2015). A negative

non-template control was also amplified to account for extraction and PCR contamination. A

total of 50 µL of PCR reaction mixtures were prepared according to the following recipe: 23 µL

PCR-grade water, 20 µL PCR master mix (2x), 1 µL 806R primer (10 µM), 1 µL barcoded 515F

primer (10µM), and 5 µL template DNA. PCR was performed with the following program: 1 x 3

min at 94°C, 35 x (45 s at 94°C, 60 s at 50°C, 90 s at 72°C), 1 x 10 min at 72°C. Amplified DNA

was stored at -20°C for one week. The PCR product was run on a 1.5% agarose gel to check for

correct amplification size. Following the gel, PCR products were cleaned using AMPure XP

Beads™ (Beckman Coulter Inc., Brea, CA). DNA concentrations were quantified using the

Qubit dsDNA high sensitivity (HS) kit™ (Thermo Fisher Scientific Inc., Waltham, MA) and

samples were diluted to 4 nM using PCR-grade water. Then 5 µL of each sample was pooled

into a single 1.5 mL tube. The pooled DNA library was sent for sequencing on an Illumina

MiSeq™ PE300 sequencer at the Center for Genome Technology (CGT) at the University of

Miami's Miller School of Medicine for quality control and sequencing. Raw Illumina paired-end

reads have been deposited in the NCBI Sequence Read Archive (SRA) database under

BioProject PRJNA610635.

Post-Sequencing Processing and Statistical Analysis

Post-sequencing processing was conducted using the Quantitative Insights in Microbial

Ecology 2 (QIIME2) v2019.1 pipeline (Bolyen *et al.* 2019). Demultiplexed paired-end reads

were merged using DADA2 v1.10.0 and processed to yield amplicon sequence variants (ASVs)

(Callahan *et al.* 2016). Within the DADA2 pipeline, forward sequences were truncated at 200 bp

and reverse sequences were truncated at 180 bp. Taxonomy was assigned using the SILVA 132

99% classifier (https://data.qiime2.org/2019.10/common/silva-132-99-nb-classifier.qza) (Quast

et al. 2013). Mitochondria and chloroplast sequences were filtered from the feature table. The

Phyloseq R package v1.28.0 was used for data handling (McMurdie and Holmes 2013). The R

package decontam v1.4.0 was used to eliminate microbial contaminants by removing all

sequences that were more prevalent in the negative control than in the real samples (Davis *et al.*

2018). Normalization across samples was conducted using a variance stabilizing transformation

(VST) on the data using DESeq2 v1.24.0 (Love, Huber and Anders 2014). To account for

abundance and evenness of microbial taxa within the VST dataset, the Shannon-Weiner

Diversity Index and Simpson's Diversity Index were calculated for each compartment region

using Phyloseq. DESeq2 v1.24.0 and the QIIME2 add-on, Analysis of Composition of

Microbiomes (ANCOM), were used for microbial differential abundance analysis against tissue-

compartments and differences were considered significant if they had a p-value < 0.05. Random

forest classifier modeling was performed using randomForest in order to determine the main

bacterial taxa driving tissue compartment composition (Liaw and Wiener 2002). For beta-

diversity measurements, both Bray-Curtis and Aitchison (Euclidean distances of center-log ratio

transformed counts) distance matrices were assessed as both are commonly used in microbial

community studies. For Bray-Curtis distances, the VST-transformed data was used. For

Aitchison distances, the R package Microbiome v1.6.0 was used to perform a center-log ratio

transformation on the raw count data (Lahti *et al.* 2019). Vegan v2.5.6 was used to calculate

Bray-Curtis and Euclidean distances and the *Adonis* function was used to perform

PERMANOVA analyses in order to statistically assess dissimilarity in microbial community

composition between tissue compartments (Oksanen *et al.* 2013). Ampvis2 v2.5.7 was used to

generate a heat-map of the most abundant bacterial families according to tissue compartment

(Andersen *et al.* 2018). All associated code can be found on GitHub:

https://github.com/Abonacolta/Nvec_Compartmentalization.

Mapping N. vectensis ASVs to a phylogenetic tree of Spirochetes

The 16S rRNA gene sequences from the phylum Spirochaetae were retrieved from NCBI

for representatives from every major spirochete lineage in FASTA format. Using BLASTn,

closely clustering (97%) sequences to the two spirochete ASVs recovered in this study were

added to the FASTA file along with a Planctomycete sequence (Accession: X56305) as an

outgroup, giving a total of 64 sequences. These sequences were aligned using MAFFT v7.450

(auto-setting) with the L-INS-i strategy (Katoh and Standley 2013). The alignment was trimmed

to remove columns with gaps in more than 70% of the sequences or with a similarity score below

0.001 using TrimAl v1.2 (Capella-Gutiérrez, Silla-Martínez and Gabaldón 2009). A maximum-

likelihood tree was constructed with RAxML v7.3.0 using the GTRCAT substitution model and

1000 sub-samplings to generate a best-fit tree (Stamatakis 2014). The final tree was visualized

and refined in FigTree v 1.4.4 (Drummond *et al.* 2012).

Results and Discussion

The body regions of N. vectensis exhibited distinct bacterial associations

We had an average of 113,758 reads per sample and there was a total of 620 ASVs with

five ASVs removed as they were likely contaminants. The microbial composition of the

capitulum (n= 6) is distinct from the mesenteries (n= 6) and physa (n= 6; Fig. 1). With the

mesenteries showing a similar bacterial community composition as the physa. Overall the

microbiomes were consistent between replicates, however, one individual harbored a different

compartmental microbiome from the others in this study (Figs. 3, S1, S2, & S3). Spirochetes

(predominantly from 1 ASV) from the family Spirochaetaceae made up ~66% of the bacterial

community within the capitulum of *N. vectensis*, but only 1.2% and 0.1% of the community in

the mesenteries and physa, respectively (Fig. 2). This finding is similar to the microbiome of the

Crown-of-thorns Starfish, *Acanthaster planci*, which found a high relative abundance (43-64%)

of spirochetes (predominantly from 1 ASV) only within the body wall compartment. Prior to this

Alphaproteobacteria was thought to be the most abundant bacterial phylum across the sea stars

(Høj *et al.* 2018).

In the mesenteries a larger prevalence of Gammaproteobacteria was found (Fig. 1 & S2),

which may play an important role in microbial community structuring in juvenile *N. vectensis*

(Domin *et al.* 2018). The physa of *N. vectensis* was dominated by the classes

Gammaproteobacteria, Alphaproteobacteria, and Bacteroidetes (Fig. S2). Similarly, previous

studies on the microbiome of lab-raised *N. vectensis* found that Proteobacteria (including the

classes Gammaproteobacteria and Alphaproteobacteria) was the most abundant bacterial phylum,

with Firmicutes, Planctomycetes, Tenericutes, Verrucomicrobia, and Spirochetes having lower,

but still significant, abundances (Har *et al.* 2015a; Mortzfeld *et al.* 2016; Leach, Carrier and

Reitzel 2019). In addition to the physa, the mesenteries were also dominated by the class

Gammaproteobacteria, among them Oceanospirillales and Pseudomonadales were the primary

orders, with Pseudomonadales being more abundant than Oceanospirillales (Fig. S3). Among

bacterial families, Pseudomonadaceae had the highest relative abundance in both the mesenteries

and physa (~36.5% and ~38.1%, respectively) (Fig. 2). This result is in line with a fluorescent *in*

situ hybridization study that revealed that Proteobacteria and Pseudomonas occur predominantly

in aggregates along the walls of the mesenteries (Har 2009).

Interestingly, one study found that as *N. vectensis* develops, Betaproteobacteria,

Actinobacteria, and Bacteroidetes decrease, while Spirochetes become one of the most abundant

phyla in the adult organism (Mortzfeld *et al.* 2016). The development and growth of the

capitulum during this time span may explain the high relative abundance of spirochetes that we

saw in the capitulum. Spirochetes may play an active role in the capitulum development of *N.*

vectensis and their ability to establish dominance over the Proteobacteria present throughout the

rest of *N. vectensis* implies a competitive advantage on the part of the Spirochete. Of note is that

Spirochetes are not found in high abundance in artemia or mussel, the two food sources for the

Nematostella used in this study (Quiroz *et al.* 2015; Weingarten, Atkinson and Jackson 2019).

Spirochetes have been identified as symbionts in bivalves, corals, oligochaetes, and

termites (as epibionts on intestinal protists) where they perform reductive acetogenesis and

nitrogen fixation (Margulis, Nault and Sieburth 1991; Lilburn, Schmidt and Breznak 1999;

Campbell and Cary 2001; Ohkuma *et al.* 2015). In cnidarians, spirochetes have been identified in

a wide range of cnidarians including red coral, *Corallium rubrum*, *Hydra vulgaris* and the

octocoral, *Lobophytum pauciflorum (*Fig. 4; Van De Water *et al.* 2016; Wessels *et al.* 2017;

Hufnagel and Myhal 1977). In *L. pauciflorum* spirochetes are hypothesized to play a role in

nitrogen fixation and microbial community (Van De Water *et al.* 2016; Wessels *et al.* 2017)

however, in other cnidarians, their function is not well understood. The predominance of

spirochetes as symbionts in multiple cnidarians suggests a potential conserved benefit to the

cnidarian hosts. In the future, testing the role of spirochetes in nitrogen fixation in the capitulum

of *N. vectensis* may inform our understanding of these potential symbionts.

Differential microbial community structure between body regions

The taxonomic make-up of the microbiome of *N. vectensis* varied according to body

region sampled. Comparing the differential abundance of taxa across tissue compartments using

a random forest classifier showed that the phyla Lentisphaerae, Proteobacteria, and Spirochetes

were the strongest predictors of tissue compartments. According to differential abundance

analysis, Spirochetes and Fusobacteria were significantly more abundant in the capitulum

compared to the other compartments (DESeq2; p-adj=0.0000 and 0.0003, respectively).

Lentisphaerae, Proteobacteria, and Chloroflexi were more abundant in the mesenteries compared

to the capitulum (DESeq2; p-adj= 0.0000, 0.0058, and 0.0009, respectively). According to

differential abundance analysis using ANCOM, Spirochetes and Bacteroidetes were the most

differentially abundant phyla among tissue compartments (ANCOM; W = 16 & 14,

respectively). Spirochetes was the only taxon to be considered significantly differentially

abundant or as a strong predictor of body region in *N. vectensis* by all programs, emphasizing the

significance of this association (Fig. S4). The differences in dominant taxa between the

capitulum and the other two tissue compartments in *N. vectensis* highlight the need to use

compartmentalization to better understand the roles of the microbiome within the holobiont - be

that in metabolism, defense, or nutrient cycling.

The body compartments of *N. vectensis* sampled in this study exhibited distinct microbial

community structures. Alpha-diversity metrics showed a significant difference in the biodiversity

and species richness present in different tissue compartments. Species richness among the

capitulum and the mesenteries and physa were significantly different (ANOVA; p = 0.023, p =

0.033, respectively; Fig. S5). The Simpson's Index between the capitulum and the mesenteries

and physa was also significantly different (ANOVA; p = 0.039, p = 0.022, respectively; Fig. S5).

However, the Shannon Diversity Index did not show a significant difference in alpha-diversity

among body regions (Fig. S5). Overall, the capitulum of *N. vectensis* showed lower species

richness compared to the mesenteries and physa, which had very similar species richness across

individuals. A similar pattern was identified in the Simpson's Index, with the capitulum showing

a lower species diversity, but with a higher variation between individuals. The lower species

richness in the capitulum was likely driven by the dominance of the one single Spirochete ASV.

Both Bray-Curtis Dissimilarity and Aitchison distance were used to assess the differences in the

microbial communities between tissue compartments (Fig. 3). A multivariate homogeneity of

groups dispersions analysis confirmed that an ANOVA test could appropriately be performed on

both datasets (Koleff, Gaston and Lennon 2003). Bacterial communities between tissue

compartments showed significant dissimilarity (PERMANOVA; permutations = 9999, p =

0.0315) according to the Bray-Curtis Dissimilarity Index. However, the same PERMANOVA

test on the Aitchison distances found insignificant dissimilarity between tissue compartments

(PERMANOVA; permutations = 9999, p = 0.3294).

Phylogenetic analysis of N. vectensis Spirochetes

The majority of spirochetes found in the capitulum was from a single ASV (Spirochete 1)

within the order Spirochaetales. In order to better understand the phylogenetic placement of the

spirochetes within *N. vectensis,* we conducted a phylogenetic analysis of this ASV along with the

other spirochete ASV (Spirochete 2) recovered in this study. Strong bootstrap support (93%

bootstrap) placed Spirochete 1 as closely related to four spirochete ASVs recovered from wild *N.*

vectensis from Har *et al.* 2015 supporting this spirochete as a consistent associate of the anemone

(Fig. 4). The potential role of the spirochetes within the capitulum of *N. vectensis* should be

investigated further as this relationship appears to be conserved, regardless of environment.

These *N. vectensis* ASVs also showed a strong relation to a spirochete recovered from the cold-

water coral, *Lophelia pertusa*. Together, these sequences formed a strongly supported (90%)

monophyletic clade with the two representative species of the recently described genus

Oceanispirocheata. The genus *Oceanispirochaeta* consists solely of two representatives, *O.*

litoralis and *O. sediminicola*, which are Gram-negative, moderately halophilic, and obligately

anaerobic (Subhash and Lee 2017).

Spirochete 2, along with several closely clustering wastewater ASVs, showed strong

relation (100% bootstrap) to *Turneriella parva*, which is a Gram-negative, obligately aerobic

spirochete with morphological similarities to *Leptospira* (Stackebrandt *et al.* 2013). Spirochetes

associated with a diverse group of corals, including *Corallium rubrum, Orbicella faveolata, and*

Lophelia pertusa, did form a well-supported (100% bootstrap) monophyletic clade within the

genus *Leptospira*, despite the broad geographical ranges of these coral species (*O. faveolata* is a

Caribbean species*, C. rubrum* is a Mediterranean species, & *L. pertusa* is a deep sea species)

indicating a potentially conserved association of this bacteria with Cnidarians.

Conclusion

In this study we characterized the microbiome of three distinct body compartments within

N. vectensis. A capitulum-specific dominance of spirochetes was uncovered highlighting the

possible importance of this bacteria in *N. vectensis* physiology. Previous bulk-sampling

techniques could not untangle compartment-specific bacterial associations, which should be

taken into consideration in future cnidarian studies. Specifically, the role of spirochetes within

the capitulum of *N. vectensis* should be further considered in subsequent studies. Overall, our

findings further establish spirochetes as consistent bacteria associates within *N. vectensis* and

suggests that the functional role of this bacteria in the capitulum may be important for the

anemone.

Acknowledgements

318   Authors would like to thank members of the Cnidarian Immunity Lab for comments and

319   suggestions as this project was developing. AMB: was supported by the RSMAS Small

320   Undergraduate Research Grant Experience (SURGE) Award. NTK was supported by startup

321   funds from University of Miami, Rosenstiel School of Marine and Atmospheric Sciences and by

322   NSF Award #: 1951826

References

Aagaard K, Ma J, Antony KM *et al.* The Placenta Harbors a Unique Microbiome. *Sci Transl*

Med 2014;**6**, DOI: 10.1126/scitranslmed.3009864.

Ainsworth TD, Thurber RV, Gates RD. The future of coral reefs: a microbial perspective. *Trends*

Ecol Evol 2010;**25**:233–40.

Apprill A, Mcnally S, Parsons R *et al.* Minor revision to V4 region SSU rRNA 806R gene

primer greatly increases detection of SAR11 bacterioplankton. *Aquat Microb Ecol*

2015;**75**:129–37.

Apprill A, Weber LG, Santoro AE. Distinguishing between Microbial Habitats Unravels

Ecological Complexity in Coral Microbiomes. *mSystems* 2016;**1**:e00143-16.

Bolyen E, Rideout JR, Dillon MR *et al.* Reproducible, interactive, scalable and extensible

microbiome data science using QIIME 2. *Nat Biotechnol* 2019;**37**:852–7.

Bourne DG, Morrow KM, Webster NS. Insights into the Coral Microbiome: Underpinning the

Health and Resilience of Reef Ecosystems. *Annu Rev Microbiol* 2016;**70**:317–40.

Callahan BJ, Mcmurdie PJ, Rosen MJ *et al.* DADA2: High resolution sample inference from

Illumina amplicon data. *Nat Methods* 2016;**13**:581–3.

Campbell BJ, Cary SC. Characterization of a Novel Spirochete Associated with the

Hydrothermal Vent Polychaete Annelid, Alvinella pompejana. *Appl Environ Microbiol*

2001;**67**:110–7.

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: A tool for automated alignment

trimming in large-scale phylogenetic analyses. *Bioinformatics* 2009;**25**:1972–3.

Davis NM, Proctor DiM, Holmes SP *et al.* Simple statistical identification and removal of

contaminant sequences in marker-gene and metagenomics data. *Microbiome* 2018;**6**:1–14.

Domin H, Zurita-Gutiérrez YH, Scotti M *et al.* Predicted bacterial interactions affect in vivo

microbial colonization dynamics in Nematostella. *Front Microbiol* 2018;**9**:1–12.

Drummond AJ, Suchard MA, Xie D *et al.* Bayesian phylogenetics with BEAUti and the BEAST

1.7. *Mol Biol Evol* 2012;**29**:1969–73.

Foster KR, Schluter J, Coyte KZ *et al.* The evolution of the host microbiome as an ecosystem on

a leash. *Nature* 2017;**548**:43–51.

Har JY. Introducing the starlet sea anemone Nematostella vectensis as a model for investigating.

2009.

Har JY, Helbig T, Lim JH *et al.* Microbial diversity and activity in the Nematostella vectensis

holobiont: Insights from 16S rRNA gene sequencing, isolate genomes, and a pilot-scale

survey of gene expression. *Front Microbiol* 2015a;**6**, DOI: 10.3389/fmicb.2015.00818.

Har JY, Helbig T, Lim JH *et al.* Microbial diversity and activity in the Nematostella vectensis

holobiont: Insights from 16S rRNA gene sequencing, isolate genomes, and a pilot-scale

survey of gene expression. *Front Microbiol* 2015b;**6**, DOI: 10.3389/fmicb.2015.00818.

Høj L, Levy N, Baillie B *et al.* Crown-of-Thorns Sea Star Acanthaster cf. solaris Has Tissue-

Characteristic Microbiomes with Potential Roles in Health and Reproduction. *Appl Environ*

Microbiol 2018;**84**:1–18.

Hooper L V., Littman DR, Macpherson AJ. Interactions between the microbiota and the immune

system. *Science (80-)* 2012;**336**:1268–73.

Hufnagel LA, Myhal ML. Observations on a Spirochaete Symbiotic in Hydra. *Trans Am Microsc*

Soc 1977;**96**:406–11.

Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: Improvements

in performance and usability. *Mol Biol Evol* 2013;**30**:772–80.

Koleff P, Gaston KJ, Lennon JJ. Measuring beta diversity for presence-absence data. *J Anim*

Ecol 2003;**72**:367–82.

Layden MJ, Meyer NP, Pang K *et al.* Expression and phylogenetic analysis of the zic gene

family in the evolution and development of metazoans. *Evodevo* 2010;**1**:1–16.

Leach WB, Carrier TJ, Reitzel AM. Diel patterning in the bacterial community associated with

the sea anemone Nematostella vectensis. *Ecol Evol* 2019;**9**:9935–47.

Liaw A, Wiener M. Classification and Regression by randomForest. *R News* 2002;**2**:18–22.

Lilburn TG, Schmidt TM, Breznak JA. Phylogenetic diversity of termite gut spirochaetes.

Environ Microbiol 1999;**1**:331–45.

Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq

data with DESeq2. *Genome Biol* 2014;**15**:1–21.

Margulis L, Nault L, Sieburth JM. Cristispira from oyster styles: complex morphology of large

symbiotic spirochetes. *Symbiosis* 1991;**11**:1–17.

McMurdie PJ, Holmes S. Phyloseq: An R Package for Reproducible Interactive Analysis and

Graphics of Microbiome Census Data. *PLoS One* 2013;**8**, DOI:

10.1371/journal.pone.0061217.

Mortzfeld BM, Urbanski S, Reitzel AM *et al.* Response of bacterial colonization in Nematostella

vectensis to development, environment and biogeography. *Environ Microbiol*

2016;**18**:1764–81.

Ohkuma M, Noda S, Hattori S *et al.* Acetogenesis from H2 plus CO2and nitrogen fixation by an

endosymbiotic spirochete of a termite-gut cellulolytic protist. *Proc Natl Acad Sci U S A*

2015;**112**:10224–30.

Putnam NH, Srivastava M, Hellsten U *et al.* Sea anemone genome reveals ancestral eumetazoan

gene repertoire and genomic organization. *Science (80-)* 2007;**317**:86–94.

Quast C, Pruesse E, Yilmaz P *et al.* The SILVA ribosomal RNA gene database project:

Improved data processing and web-based tools. *Nucleic Acids Res* 2013;**41**:590–6.

Quiroz M, Triadó-Margarit X, Casamayor EO *et al.* Comparison of Artemia–bacteria

associations in brines, laboratory cultures and the gut environment: a study based on

Chilean hypersaline environments. *Extremophiles* 2015;**19**:135–47.

Rädecker N, Pogoreutz C, Voolstra CR *et al.* Nitrogen cycling in corals: The key to

understanding holobiont functioning? *Trends Microbiol* 2015;**23**:490–7.

Reitzel AM, Ryan JF, Tarrant AM. Establishing a model organism: A report from the first annual

Nematostella meeting. *BioEssays* 2012;**34**:158–61.

Renfer E, Amon-Hassenzahl A, Steinmetz PRH *et al.* A muscle-specific transgenic reporter line

of the sea anemone, Nematostella vectensis. *Proc Natl Acad Sci U S A* 2010;**107**:104–8.

Ricci F, Rossetto Marcelino V, Blackall LL *et al.* Beneath the surface: Community assembly and

functions of the coral skeleton microbiome. *Microbiome* 2019;**7**:1–10.

Robbins SJ, Singleton CM, Chan CX *et al.* A genomic view of the reef-building coral Porites

lutea and its microbial symbionts. *Nat Microbiol* 2019:1–11.

Rosenberg E, Koren O, Reshef L *et al.* The role of microorganisms in coral health , disease and

evolution. *Nat Rev Microbiol* 2007;**5**:355–62.

Rossbach S. Tissue-Specific Microbiomes of the Red Sea Giant Clam Tridacna maxima

Highlight Differential Abundance of Endozoicomonadaceae. 2019;**10**, DOI:

10.3389/fmicb.2019.02661.

Stackebrandt E, Chertkov O, Lapidus A *et al.* Genome sequence of the free-living aerobic

spirochete Turneriella parva type strain (HT), and emendation of the species Turneriella

parva. *Stand Genomic Sci* 2013;**8**:228–38.

Stamatakis A. RAxML version 8: A tool for phylogenetic analysis and post-analysis of large

phylogenies. *Bioinformatics* 2014;**30**:1312–3.

Subhash Y, Lee SS. Description of oceanispirochaeta sediminicola gen. Nov., sp. nov., an

obligately anaerobic bacterium isolated from coastal marine sediments, and reclassification

of spirochaeta litoralis as oceanispirochaeta litoralis comb. nov. *Int J Syst Evol Microbiol*

2017;**67**:3403–9.

Sweet MJ, Croquer A, Bythell JC. Bacterial assemblages differ between compartments within

the coral holobiont. *Coral Reefs* 2011;**30**:39–52.

Thompson LR, Sanders JG, McDonald D *et al.* A communal catalogue reveals Earth's multiscale

microbial diversity. *Nature* 2017;**551**:457–63.

Vaishnava S, Behrendt CL, Ismail AS *et al.* Paneth cells directly sense gut commensals and

maintain homeostasis at the intestinal host-microbial interface. *Proc Natl Acad Sci U S A*

2008;**105**:20858–63.

Van De Water JAJM, Melkonian R, Junca H *et al.* Spirochaetes dominate the microbial

community associated with the red coral Corallium rubrum on a broad geographic scale. *Sci*

Rep 2016;**6**:1–7.

Weingarten EA, Atkinson CL, Jackson CR. The gut microbiome of freshwater Unionidae

mussels is determined by host species and is selectively retained from filtered seston. *PLoS*

One 2019;**14**:1–17.

Wessels W, Sprungala S, Watson SA *et al.* The microbiome of the octocoral Lobophytum

pauciflorum: Minor differences between sexes and resilience to short-term stress. *FEMS*

Microbiol Ecol 2017;**93**:1–13.

Wier AM, Sacchi L, Dolan MF *et al.* Spirochete Attachment Ultrastructure: Implications for the

Origin and Evolution of Cilia. *Biol Bull* 2010;**218**:25–35.

Fig 1. Compartmentalization of the microbiome in *N. vectensis*. The microbiome of *N.*

vectensis was divided into three tissue compartments for this study: the capitulum, mesenteries,

and physa. The capitulum had a bacterial community dominated by Spirochetes. The mesenteries

and physa had a bacterial community dominated by Gammaproteobacteria and

Alphaproteobacteria, with very little presence of Spirochetes.

Fig 2. Heat-map of the percent read abundance for top bacterial families found in *N.*

***vectensis* separated by tissue compartment.** Bacteria of the family Pseudomonadaceae were

found to be the most prevalent in the mesenteries and physa tissue of *N. vectensis* (38.1% and

36.5% abundance, respectively). In contrast, Pseudomonadaceae only had a 7.4% abundance in

capitulum tissue. Likewise, Spirochaetaceae were found to make up 66% of the community

within the capitulum tissue of *N. vectensis*, while only being 1.2% and 0.1% abundant in the

mesentery and physa tissues, respectively. Overall, the capitulum tissue appears to have a very

distinct community from the mesenteries and physa.

Fig 3. Beta diversity measures between tissue regions of *N. vectensis*. (A) Bray-Curtis PCoA

of *N. vectensis* bacterial communities after a variance stabilizing transformation. Capitulum

samples clustered tightly, while physa and mesentery communities clustered closely together.

(B) Aitchison PCA (Euclidean distance of center-log ratio transformed sample counts) of *N.*

vectensis bacterial communities. Again, most capitulum samples clustered closely together,

however they were also clustered with mesenteries and physa samples.

Fig 4. Best-fit maximum likelihood tree based on 16S rRNA sequences depicting the

phylogenetic relationship of the spirochetes recovered from the capitulum of *Nematostella*

vectensis. This tree was constructed using RAxML v7.3.0 with 1000 sub samplings. Bootstrap

values (> 50%) are shown. Nodes with 100% bootstrap support are indicated by open circles.

GenBank accession numbers are listed before each sequence's name. The host/habitat of

uncultured species is in parentheses. Spirochetes from this study are in bold. Shading

corresponds to major clades within the phylum Spirochaetae. *Gemmata obscuriglobus* was used

as an outgroup to root the tree. The predominant spirochete ASV from this study (Spirochete 1)

forms a strongly supported (100% bootstrap) monophyletic clade with spirochete sequences

recovered from wild *N. vectensis*. Spirochete 2 clusters within the genus *Turneriella*.

Compartmentalization of the microbiome in *N. vectensis*. The microbiome of *N. vectensis* was divided into three tissue compartments for this study: the capitulum, mesenteries, and physa. The capitulum had a bacterial community dominated by Spirochetes. The mesenteries and physa had a bacterial community dominated by Gammaproteobacteria and Alphaproteobacteria, with very little presence of Spirochetes.

338x218mm (300 x 300 DPI)

Beta diversity measures between tissue regions of *N. vectensis*. (A) Bray-Curtis PCoA of *N. vectensis* bacterial communities after a variance stabilizing transformation. Capitulum samples clustered tightly, while physa and mesentery communities clustered closely together. (B) Aitchison PCA (Euclidean distance of center-log ratio transformed sample counts) of *N. vectensis* bacterial communities. Again, most capitulum samples clustered closely together, however they were also clustered with mesenteries and physa samples.

355x127mm (300 x 300 DPI)

Best-fit maximum likelihood tree based on 16S rRNA sequences depicting the phylogenetic relationship of the spirochetes recovered from the capitulum of *Nematostella vectensis*. This tree was constructed using RAxML v7.3.0 with 1000 sub samplings. Bootstrap values (> 50%) are shown. Nodes with 100% bootstrap support are indicated by open circles. GenBank accession numbers are listed before each sequence's name. The host/habitat of uncultured species is in parentheses. Spirochetes from this study are in bold. Shading corresponds to major clades within the phylum Spirochaetae. *Gemmata obscuriglobus* was used as an outgroup to root the tree. The predominant spirochete ASV from this study (Spirochete 1) forms a strongly supported (100% bootstrap) monophyletic clade with spirochete sequences recovered from wild *N. vectensis*. Spirochete 2 clusters within the genus *Turneriella*.

Title: The Starlet Sea Anemone, *Nematostella vectensis,* possesses body region-specific bacterial

associations with spirochetes dominating the capitulum

Authors: A. M. Bonacolta, M. T. Connelly, S. Rosales, J. del Campo, N. Traylor-Knowles

Supplemental Material

Fig S1. The relative abundance of bacterial phyla recovered from *N. vectensis* in

individuals and tissue compartments Only taxonomic phyla with a relative abundance above

2% were incorporated. Spirochetes dominated the bacterial communities in the majority of

capitulum samples (Individual 1 again being an outlier). Proteobacteria dominated the bacterial

communities of mesentery and physa samples.

Fig S2. The relative abundance of bacterial classes recovered from *N. vectensis* in

individuals and tissue compartments Only taxonomic classes with a relative abundance above

2% were incorporated. Spirochaetia dominated the bacterial communities in the majority of

capitulum samples (Individual 1 again being an outlier). Gammaproteobacteria dominated the

bacterial communities of mesentery and physa samples.

Fig S3. The relative abundance of bacterial orders recovered from *N. vectensis* in

individuals and tissue compartments. Only taxonomic orders with a relative abundance above

2% were incorporated. The order Spirochaetales dominated the bacterial communities in the

majority of capitulum samples (Individual 1 being an outlier). The mesentery and physa body

regions showed more variety in their composition with the orders Pseudomonadales,

Alteromonadales, and Oceanospirillales making up the majority of most samples.

34   **Fig S4. Venn diagram of significant taxa by program.** Taxa with significant differential

35   abundance between body regions were determined by ANCOM and DESeq2. Taxa which were

36   shown to be the strongest predictor of body compartment by randomForest are also listed.

37   Spirochetes was the only taxon to be considered significantly differentially abundant or as a

38   strong predictor of body region in *N. vectensis* by all programs.

Fig S5. Bacterial community Alpha diversity measures between tissue regions. Species

richness between the capitulum, mesenteries, and physa was significantly different (ANOVA;

p=0.023, p=0.033, respectively). The Simpson Alpha Diversity Index between the capitulum,

mesenteries and physa was significantly different (ANOVA; p=0.039, p=0.022, respectively).

The Shannon Diversity Index did not show a significant difference in the bacterial communities

between body regions.

529x529mm (72 x 72 DPI)

740x529mm (72 x 72 DPI)

338x190mm (300 x 300 DPI)